

TWO RAMSEY PROBLEMS IN BLOWUPS OF GRAPHS

ANTÓNIO GIRÃO AND ROBERT HANCOCK

ABSTRACT. Given graphs G and H , we say $G \xrightarrow{r} H$ if every r -colouring of the edges of G contains a monochromatic copy of H . Let $H[t]$ denote the t -blowup of H . The blowup Ramsey number $B(G \xrightarrow{r} H; t)$ is the minimum n such that $G[n] \xrightarrow{r} H[t]$. Fox, Luo and Wigderson refined an upper bound of Souza, showing that, given G , H and r such that $G \xrightarrow{r} H$, there exist constants $a = a(G, H, r)$ and $b = b(H, r)$ such that for all $t \in \mathbb{N}$, $B(G \xrightarrow{r} H; t) \leq ab^t$. They conjectured that there exist some graphs H for which the constant a depending on G is necessary. We prove this conjecture by showing that the statement is true in the case of H being 3-chromatically connected, which in particular includes triangles. On the other hand, perhaps surprisingly, we show that for forests F , the function $B(G \xrightarrow{r} F; t)$ is independent of G .

Second, we show that for any $r, t \in \mathbb{N}$, any sufficiently large r -edge coloured complete graph on n vertices with $\Omega(n^{2-1/t})$ edges in each colour contains a member from a certain finite family \mathcal{F}_t^r of r -edge coloured complete graphs. This answers a conjecture of Bowen, Hansberg, Montejano and Můyesser.

1. BLOWUP RAMSEY NUMBERS

We write $G \xrightarrow{r} H$ if every r -colouring of the edges of G contains a monochromatic copy of H . Given a graph H and $t \in \mathbb{N}$, the t -blowup of H , denoted $H[t]$, is the graph obtained from H by replacing each vertex of H by an independent set of size t , and replacing every edge of H by a complete bipartite graph $K_{t,t}$ between the corresponding parts. We call a copy of $H[t]$ in $G[n]$ *canonical* if it is the t -blowup of a copy of H in G . Suppose G, H are graphs and $r \in \mathbb{N}$ such that $G \xrightarrow{r} H$. Then for $t \in \mathbb{N}$, the *blowup Ramsey number* $B(G \xrightarrow{r} H; t)$ is the minimum n such that every r -colouring of the edges of $G[n]$ contains a monochromatic canonical copy of $H[t]$.

Blowup Ramsey numbers were introduced by Souza in [13], as a natural generalisation of bipartite Ramsey numbers. At the same time, he proved the following upper bound on them.

Theorem 1.1 (Souza [13]). *Let G, H be graphs and $r \geq 2$ an integer such that $G \xrightarrow{r} H$. Then there is a constant $c = c(G, H, r)$ such that for every $t \in \mathbb{N}$,*

$$B(G \xrightarrow{r} H; t) \leq c^t.$$

Souza also proved that the exponential-type bound is necessary, finding a lower bound with an exponential constant which does not depend on G . This result led him to conjecture that in general the dependence on G is not necessary, that is, for any H , the dependence on G in the upper bound can be removed from the function c .

AG: Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK. E-mail: girao@maths.ox.ac.uk. Research supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC-2181/1 - 390900948 (the 14 Heidelberg STRUCTURES Cluster of Excellence) and by EPSRC grant EP/V007327/1.

RH: Institut für Informatik, Heidelberg University, Im Neuenheimer Feld 205, 69120, Heidelberg, Germany. E-mail: hancock@informatik.uni-heidelberg.de. Research supported by a Humboldt Research Fellowship.

Shortly after, Fox, Luo and Wigderson proved that the exponential constant does not depend on G , however, the upper bound still has some dependence on G . In particular, they proved the following.

Theorem 1.2 (Fox–Luo–Wigderson [7]). *Let G, H be graphs and $r \geq 2$ an integer such that $G \xrightarrow{r} H$. There exist constants $a = a(G, H, r)$ and $b = b(H, r)$ such that for every $t \in \mathbb{N}$,*

$$B(G \xrightarrow{r} H; t) \leq ab^t.$$

However, Fox, Luo and Wigderson conjectured that the dependence on G is necessary for some graphs H , contrary to the conjecture of Souza.

Conjecture 1.3 (Fox–Luo–Wigderson [7]). *There exists a graph H and integers $r, t \geq 2$ for which the following holds. There exist graphs G_1, G_2, \dots such that $G_i \xrightarrow{r} H$ for all i and $\sup_i B(G_i \xrightarrow{r} H; t) = \infty$.*

In particular Fox, Luo and Wigderson conjectured that this holds with H a triangle and $r = t = 2$. Our first main result is to confirm this conjecture.

Theorem 1.4. *For every integer $s \geq 2$, there exists a graph G such that $G \xrightarrow{2} K_3$ but $G[s] \not\xrightarrow{2} K_3[2]$.*

In fact we can prove the result for a much larger class of graphs. A graph is *3-chromatically connected* if $|V(G)| \geq 3$ and for all $V \subseteq V(G)$ such that $G[V]$ is bipartite, the graph $G - V$ is connected. (In particular, note that a triangle is 3-chromatically connected.) Our theorem extends to all graphs which are 3-chromatically connected. We also note that it makes no difference in the proof to replace 2 by any $r \geq 2$.

Theorem 1.5. *Let H be a 3-chromatically connected graph and let $r \geq 2$. For every integer $s \geq 2$, there exists a graph G such that $G \xrightarrow{r} H$ but $G[s] \not\xrightarrow{r} H[2]$.*

We prove this result in Section 1.1. Our theorem relies on a sparse Ramsey theorem of Nešetřil and Rödl (see Theorem 1.7), which in particular, requires the graph in question to be 3-chromatically connected. We are therefore as of yet unable to extend our result to graphs which are not 3-chromatically connected. It would be very interesting to determine precisely which graphs H require some dependence on G in their blowup Ramsey numbers, and which do not.

We call G *r -Ramsey-minimal* for H , if $G \xrightarrow{r} H$, and for any proper subgraph G' of G , we have $G' \not\xrightarrow{r} H$. A graph H is *r -Ramsey-finite* if its class of r -Ramsey-minimal graphs is finite. Souza proved that r -Ramsey-finite graphs do not require the dependence on G .

While any graph containing a cycle is not r -Ramsey-finite (as observed by Souza), there are some graphs, for example a star with an odd number of edges, which are 2-Ramsey-finite [3]. However, Fox, Luo and Wigderson showed that P_3 , the path with two edges, is not r -Ramsey-finite, yet it also does not have any dependence on G . We are also able to show that the same phenomenon holds for any tree T .

Theorem 1.6. *Let G be a graph, T a tree and $r \geq 2$ an integer such that $G \xrightarrow{r} T$. Then for every $t \in \mathbb{N}$, there exists a function $f = f(T, r, t)$ such that*

$$B(G \xrightarrow{r} T; t) \leq f.$$

We prove this result in Section 1.2.

1.1. **Proof of Theorems 1.4 and 1.5.** Given two graphs H and G , we define $H(G)$, the *hypergraph of copies of H in G* , to be a $|E(H)|$ -uniform hypergraph with vertex set $E(G)$, and edge set consisting of each set of edges of G which correspond to a copy of H in G .

We define the *girth* of a hypergraph H , to be the number of edges in a shortest cycle of H and denote it by $g(H)$. In particular, note that if two edges share at least two vertices, then these two edges form a cycle of length 2. Note therefore that a hypergraph with girth at least three is what is often referred to in literature as a *linear* hypergraph, and all its cycles are *loose* cycles.

The graph G which we use in the proof will be obtained using the following sparse Ramsey theorem of Nešetřil and Rödl.

Theorem 1.7 (Nešetřil–Rödl, Theorem 4.2 [12]). *Let $r, s \in \mathbb{N}$ and let H be a 3-chromatically connected graph. Then there exists a graph G with the following properties:*

- (1) $G \xrightarrow{r} H$;
- (2) $g(H(G)) > s$.

The only property of a 3-chromatically connected graph we require in our proof is the existence of the above G , together with having minimum degree at least 2 (which is required only at stage 1, step (i)(1) of the proof).

So for the rest of the proof, to make it clearer and easier to read, we will take $r = 2$ and $H = K_3$, noting that it easily extends to general $r \geq 2$ and H which are 3-chromatically connected.

Fix $s \geq 2$ and choose G according to Theorem 1.7 so that $g(H(G)) > 2s + 2$. Note that we can choose G so that it is (2-)Ramsey-minimal, which we recall means that for any edge e , we have $G - e \not\xrightarrow{2} K_3$. In order to complete the proof, it suffices to find a red/blue colouring of $G[s]$ which does not contain any monochromatic canonical copies of $K_3[2]$.

We first give an outline of the proof. The idea is to step by step recolour edges of $G[s]$ so that in each step all monochromatic copies of $K_3[2]$ created by the previous step are destroyed, while in each edge blowup, less edges are recoloured compared to the previous step. In stage 1, we will look at colourings of G . Since G is Ramsey-minimal, we can select an edge $e \in E(G)$, for which there exists a red/blue-colouring of $E(G - e)$ such that there are no monochromatic triangles. Now, we extend the red/blue-colouring of $E(G - e)$ to a red/blue-colouring of $E(G)$ by colouring e red. Since $G \xrightarrow{2} K_3$, there must exist some red triangles; note that all of these red triangles contain e , and further there are no blue triangles. Further note that these red triangles intersect at e and are otherwise edge-disjoint from each other, since $H(G)$ contains no cycle of length 2. Therefore we can select precisely one edge per red triangle to recolour blue. We obtain a colouring which no longer contains red triangles, although it does contain new blue triangles, each of which must contain one of the recoloured edges. Note that again since $H(G)$ does not contain short cycles, the set of blue triangles each share at most one edge with any other triangle (precisely the edge which was just recoloured from red to blue) and are otherwise entirely edge-disjoint from all other monochromatic triangles created previously. We do this recolouring step s times. In stage 2, we will transfer these edge colourings of G to $G[s]$ so that in each step, whenever we recolour edges of G , we recolour proportionally less of the edges in the blowup of these edges of G compared to the previous step, while still destroying all monochromatic copies of $K_3[2]$ created during the previous step. Eventually we get to a step where we do not have to recolour anything at all in order to destroy all monochromatic copies of $K_3[2]$.

We now make this recolouring procedure precise.

Stage 1: the colourings in G .

Step 0: Fix an edge $e \in E(G)$ and let c_0 be a red/blue colouring of $E(G)$ such that there are no monochromatic triangles in $E(G - e)$ and e is red. Further, fix a vertex v incident to e , and set $E_0 = \{e\}$. Let T_0 be the set of triangles in G which are red in c_0 .

Note that all triangles in T_0 contain e , and further except for e , these triangles are edge-disjoint. This follows from $g(H(G)) > 2s + 2$.

We now repeat the following step s times, for each $i = 1, \dots, s$.

Step i :

- (1) For each $T \in T_{i-1}$, change the colour of the edge of T which is not in E_{i-1} and is incident to v .
- (2) Call the new colouring c_i ;
- (3) Call the set of edges which were just recoloured E_i .
- (4) Let T_i be the set of monochromatic triangles in c_i .

Note that each time we have the following. In c_i , there are only monochromatic triangles of one colour; whichever one did not occur in the previous colouring c_{i-1} . Every edge in E_i contains the fixed vertex v . All triangles in T_i contain exactly one edge from E_i and all of these triangles contain the vertex v .

Crucially, all triangles in $\cup_{j=0}^i T_j$ are edge-disjoint from each other, except for the following cases:

- A triangle $A_i \in T_i$ shares precisely one edge e with one triangle $A_{i-1} \in T_{i-1}$, where $e \in E_i$;
- A triangle $A_i \in T_i$ may share precisely one edge e with other triangles in T_i ; if so, then $e \in E_i$.

This edge-disjointness follows from $g(H(G)) > 2s + 2$; suppose for a contradiction that there exists triangles $A_i \in T_i$ and $B_j \in T_j$, $j \leq i$, which violate the above. If they share two edges then this immediately contradicts $g(H(G)) > 2s + 2$ since they would form a cycle of length 2 in $H(G)$. Otherwise, by design, for each $0 \leq k < i$ we can find triangles $A_k \in T_k$ such that $A_k \cap A_{k+1} = e_{k+1} \in E_{k+1}$, and for each $0 \leq k < j$ we can find triangles $B_k \in T_k$ such that $B_k \cap B_{k+1} = f_{k+1} \in E_{k+1}$. Let ℓ be the maximum index for which $A_\ell = B_\ell$. If ℓ exists then the triangles $A_\ell, A_{\ell+1}, \dots, A_i, B_j, B_{j-1}, \dots, B_{\ell+1}, A_\ell$ form a cycle in $H(G)$. If not, then the triangles $A_0, \dots, A_i, B_j, B_{j-1}, \dots, B_0, A_0$ form a cycle in $H(G)$. In both cases we have a cycle of length at most $2s + 2$, a contradiction to $g(H(G)) > 2s + 2$.

Stage 2: transferring the colourings to $G[s]$.

Step 0: First we transfer the colouring c_0 of $E(G)$ to a colouring c'_0 of $E(G[s])$ in the obvious way; that is, colour every edge in the $K_{s,s}$ blowup of uw by $c_0(uw)$ for all edges $uw \in E(G)$.

For the fixed vertex v , label the vertices in its blowup by v_1, \dots, v_s . We now repeat the following step s times, for each $i = 1, \dots, s$.

Step i :

- (1) For each $uw \in E_i$ (recall v is fixed and $c_{i-1}(uw) \neq c_i(uw)$), recolour edges from $\{v_{i+1}, \dots, v_s\}$ to all vertices in the blowup of u . (Do not change the colour of edges involving $\{v_1, \dots, v_i\}$.)
- (2) Call the new colouring c'_i .

See Figure 1 for an example of step 3.

We now aim to prove that c'_s is a colouring of $G[s]$ which contains no monochromatic canonical copies of $K_3[2]$. This will follow quickly from the following claim.




FIGURE 1. An example of step 3 of stage 2. The triangle uvw is the monochromatic $T \in T_2$, with $wv \in E_2$ and $vw \in E_3$. Originally (on the left) all edges between vertices in the blowups of v and w are red. In step 3, we recolour all edges between $\{v_4, \dots, v_k\}$ and vertices in the blowup of w from red to blue (on the right). It is not too hard to see that there are no monochromatic canonical copies of $K_3[2]$ with parts in u, v, w . Note we can see evidence of the previous recolourings too; in step 2, all edges between $\{v_3, \dots, v_k\}$ and vertices in the blowup of u were coloured red; in step 1, all edges between $\{v_2, \dots, v_k\}$ and vertices in the blowup of y were coloured blue.

Claim 1.8. *In step i , for $i \in [s]$, all monochromatic canonical copies of $K_3[2]$ contained in $\{T[s] : T \in T_{i-1}\}$ are destroyed.*

Indeed, recall that in step i of stage 1, all monochromatic copies of K_3 are collected in the set T_i . Therefore, the only possible monochromatic copies of $K_3[2]$ must contain edges which were recoloured in step s of stage 2. However, no such edges exist since in fact no edges were recoloured in step s of stage 2. It remains to prove the claim.

Proof of Claim 1.8. Since all $T \in T_{i-1}$ contain v , we must use 2 vertices from $\{v_1, \dots, v_s\}$ in any monochromatic $K_3[2]$. Since all edges involving $\{v_{i+1}, \dots, v_s\}$ are recoloured, all monochromatic copies of $K_3[2]$ using these vertices are destroyed, and so we must use 2 vertices from $\{v_1, \dots, v_i\}$. In step 1, we are done, since there are not enough vertices.

Now consider any step i with $i \geq 2$. By construction, all triangles T_{i-1} are of the form uvw , with $wv \in E_i$ and $uv \in E_{i-1}$. Since $uv \in E_{i-1}$, in step $i-1$, the colour of the edges $\{v_1, \dots, v_{i-1}\}$ to vertices in the blowup of u were not changed, and hence are a different colour to the edges from $\{v_1, \dots, v_i\}$ to vertices in the blowup of w . Therefore v_i is the only vertex with the same colour of edges to vertices in the blowups of u and w , and hence we cannot find a monochromatic $K_3[2]$ using the vertices u, v and w . \square

1.2. Proof of Theorem 1.6. We will prove a stronger result, Theorem 1.10, from which our main result Theorem 1.6 will immediately follow. First we need to introduce some notation. Let G be a graph. We say c is a *partial 2-colouring* of G if there exists a subgraph $G' \subset G$ such that $c : E(G') \rightarrow \{\text{red, blue}\}$.

Given a tree T , a graph G and a partial 2-colouring c of G , we say that $H \subset G$ is a *possible red (blue) copy* of T if H is isomorphic to T and all edges of H coloured by c are red (blue). (Note that not all edges of

H need to be coloured.) We say H is a *possible monochromatic copy* of T if H is either a possible red or possible blue copy of T .

Let G be a graph, as usual we denote by $G[m]$ the m -blowup of G and more generally, given a function $m : V(G) \rightarrow \mathbb{N}$, we denote by $G[\{m(x)\}_{x \in V(G)}]$ the blowup of G where each vertex x is blown up by $m(x)$ vertices. For every $x \in V(G)$, we denote by \bar{x} the blowup of x in $G[\{m(x)\}_{x \in V(G)}]$. We write (\bar{x}, \bar{y}) to denote the complete bipartite graph between \bar{x} and \bar{y} in $G[\{m(x)\}_{x \in V(G)}]$.

Let G be a graph, c a partial 2-colouring of G and let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a non-decreasing function such that $\lim_{n \rightarrow \infty} f(n) = \infty$. (We introduce this function entirely so that we get a cleaner argument which avoids tracing all of the constants.) We say a colouring $c' : E(G[\{m(x)\}_{x \in V(G)}]) \rightarrow \{\text{red}, \text{blue}\}$ is *f -coherent with c* if for every edge $e = (x, y) \in E(G)$ if $c(e)$ is red (blue) then the complete bipartite graph (\bar{x}, \bar{y}) does not contain a blue (red) $K_{f(m(x)), f(m(y))}$ in c' .

We now prove a technical lemma.

Lemma 1.9. *Let T be a tree and $T' \subset T$ a subtree and $xy \in E(T) \setminus E(T')$ where $x \in V(T')$. Let G be a graph where $G \xrightarrow{2} T$ and let c be a partial colouring of G . Let $z \in V(G)$ and let $A = \{T'_1, \dots, T'_k\}$ be a set of distinct copies of T' such that z plays the role of x in all of these copies of T' , and for every $i \in [k]$ there is a possible monochromatic copy of T , say T_i , such that $T'_i \subset T_i$. If k is sufficiently large then there exists $i < j$ such that $T'_i \cup (T_j \setminus T'_j)$, T_i and T_j are each the same colour possible monochromatic copies of T .*

Proof. First we may pass to a subset $A' = \{T'_1, \dots, T'_{k/2}\}$ of A of at least half the size where without loss of generality we may assume all copies of T'_i lie in a possible red copy of T . If $T_j \setminus T'_j$ is vertex disjoint from T'_1 for some $1 < j \leq k/2$, then we are done. Therefore, we may assume $T_j \setminus T'_j$ has non-empty intersection with T'_1 for all $1 < j \leq k/2$. Let x_1 be the vertex of T'_1 which appears most often (at least $\frac{k}{2v(T')}$ times) in the intersection of T'_1 with $T_j \setminus T'_j$, $1 < j \leq k/2$. We may pass to a subset of A' , say $A_1 = \{T'_2, \dots, T'_{k_1}\}$ such that every $T_\ell \setminus T'_\ell$ contains the vertex x_1 . Now if $T_j \setminus T'_j$ is vertex disjoint from T'_2 for some $2 < j \leq k_1$, then we are done. Hence, again, we may pass to a subset of A_1 , say $A_2 = \{T'_3, \dots, T'_{k_2}\}$ where every $T_\ell \setminus T'_\ell$ contains the vertex x_2 , which was the vertex of T'_2 appearing most often in the intersection of T'_2 with $T_j \setminus T'_j$, $2 < j \leq k_1$. (Note that $x_2 \neq x_1$ since $x_1 \in T_2 \setminus T'_2$.) We keep doing this process, forming further subsets A_3, A_4, \dots , and finding further vertices x_3, x_4, \dots which lie in each $T_\ell \setminus T'_\ell$. If k is sufficiently large (in particular $k \geq 2v(T')^{v(T)-v(T')}$ suffices) then this process must stop, and we have the conclusion. \square

We are now ready to prove the following result which easily implies Theorem 1.6. It turns out to be more convenient to allow each vertex to be blown up a different number of times.

Theorem 1.10. *For every tree T , any subtree $T' \subset T$ there exists a non-decreasing function f such that $\lim_{n \rightarrow \infty} f(n) = \infty$ and the following holds. Let $G \xrightarrow{2} T$ and c be a partial 2-colouring of G . Then, for any 2-colouring c' of $E(G[\{m(x)\}_{x \in V(G)}])$ which is f -coherent with c there is a $H'[\{f(m(x))\}_{x \in V(H')}]$ monochromatic in red (or blue) and $H, H' \subset G$ such that $H' \subset H$ and H' is a copy of T' and H is a possible red (or blue) copy of T in G .*

Note that Theorem 1.6 easily follows by applying this theorem with $T' = T$, letting c be empty and by choosing m in hindsight so that for all $x \in V(G)$, $m(x) = m$ and $f(m) \geq t$.

Proof. First fix the larger tree T . We prove by induction on $|E(T')|$. For the base case T' is a single edge. Let f be a non-decreasing function f such that $\lim_{n \rightarrow \infty} f(n) = \infty$ and which grows sufficiently slowly so

that after an application of Ramsey's Theorem to $K_{m(x),m(y)}$ there exists a monochromatic $K_{f(m(x)),f(m(y))}$. Let c, c' be as in the statement. Since $G \xrightarrow{2} T$ there exists a possible monochromatic copy of T in G . If it contains a red (blue) edge xy , then since c' is f -coherent with c , then by Ramsey's Theorem (\bar{x}, \bar{y}) contains a red (blue) $T'[\{f(m(x)), f(m(y))\}]$ as required. If not, then T does not have any colours and is a possible red copy and a possible blue copy. Then applying Ramsey's Theorem to any edge of T , one can find a red or blue $T'[\{f(m(x)), f(m(y))\}]$ (it does not matter which colour), as required.

Now suppose that the result holds for T'' and we wish to prove the result for T' , where $T'' = T' \setminus y$ and y is a leaf of T' . Suppose y is adjacent to x in T' . We need to show that the function f exists. By induction with T, T'' there is a function g satisfying the properties in the statement. Let $f = g^k$ i.e. the composition of g k times, where k is the number outputted by Lemma 1.9 using T and T'' .

By induction we may find $H''[\{g(m(x))\}_{x \in V(H'')}]$ monochromatic in c' (note that c' is g -coherent because $f < g$) where $H'' \subset H$ and H is a possible red (blue) copy of T and H'' is a copy of T'' . Let $\{H_1, \dots, H_t\}$ be all possible red (blue) copies of T which contain H'' and for each $i \in [t]$, let $y_i \in H_i$ be the vertex which corresponds to the vertex y . Let $H''(x)$ be the subset of \bar{x} of size $g(m(x))$ which lies in the monochromatic $H''[\{g(m(x))\}_{x \in V(H'')}]$ in c' . If we can find a red (blue) copy of $K_{g(m(x)),g(m(y_i))}$ between $H''(x)$ and a subset of size $g(m(y_i))$ in \bar{y}_i (for some i) we are done; this red (blue) complete bipartite subgraph of $(H''(x), \bar{y}_i)$ together with the monochromatic $H''[\{g(m(x))\}_{x \in V(H'')}]$ forms a monochromatic $H'[\{f(m(x))\}_{x \in V(H')}]$ where $H'' \subset H' \subset H_i = H$, H is a possible red (blue) copy of T and H' is a copy of T' , as required.

So suppose this is not the case. In particular this means that no edge xy_i was coloured red (blue) in c ; as otherwise by applying Ramsey's Theorem to $(H''(x), \bar{y}_i)$ we would have been able to find a red (blue) $K_{g(m(x)),g(m(y_i))}$ in $(H''(x), \bar{y}_i)$. These edges also cannot be blue (red) in c since they are supposed to form part of a red (blue) possible copy of T .

Hence, we may shrink \bar{x} such that only the vertices in $H''(x)$ remain and we denote the new blowup of G by G_1^* . We also update c by colouring all edges xy_i blue (red) and denote it by c_1 . Clearly, the colouring c' of G_1^* is still g -coherent with c_1 , by construction. Also note that now H'' cannot be a subgraph of a possible monochromatic copy of H . Further, without replacing \bar{x} by $H''(x)$, it may not be true that c' is g -coherent with c_1 , since there may be a large blue (red) bipartite graph between \bar{x} and \bar{y}_i for some i . We only know that no such bipartite graph exists between $H''(x)$ and \bar{y}_i . Finally, we define a new function $m_1 : V(G) \rightarrow \mathbb{N}$ by defining $m_1(x) = g(m(x))$ and $m_1(y) = m(y)$ for all $y \in V(G) \setminus \{x\}$.

Now, we keep applying induction, each time shrinking the blowup of one vertex of G , and adding new colours to c and updating the function m . (For simplicity we keep the same names G, c and m .) Note that after shrinking a blowup of a vertex \bar{x} for the i -th time, it goes from size $g^{i-1}(m(x))$ to $g^i(m(x))$.

The essential fact is that each blowup of a vertex will not be shrunk more than k times because of Lemma 1.9. To see this, observe the following: suppose that $\{H_1'', H_2'', \dots, H_k''\}$ are each copies of T'' found by induction each using the same vertex x . Let H_i be the possible monochromatic copy of T which H_i'' lies in at the point when H_i'' was first found by induction. Note that later after c is updated H_i is no longer a monochromatic copy. However, crucially, in the initial colouring c , H_i is a possible monochromatic copy which H_i'' lies in, since by removing colours of edges, we do not destroy any possible monochromatic copies of T . Therefore we may apply Lemma 1.9 with $z = x$, $T_i' = H_i''$, $T_i = H_i$ and the original colouring c . Since k is large enough, the lemma outputs $i < j$ such that $H_i'' \cup (H_j \setminus H_j'')$, H_i and H_j are each possible red (blue) copies of H . Let y be the vertex of $H_j \setminus H_j''$ which is connected to x . Since the process did not stop after finding H_i'' , we must therefore have updated c by colouring xy blue (red). Now when we apply induction and

find H_j'' , since H_j contains xy , H_j must be a possible blue (red) copy of T , a contradiction, since we know it should be a possible red (blue) copy.

Since each blowup of a vertex is shrunk at most k times and each application of induction results in the shrinking of one blowup of a vertex, this process must stop and therefore we must have found the desired monochromatic blowup of T' . \square

2. UNAVOIDABLE COLOUR PATTERNS

Ramsey's Theorem states that for any $r, t \in \mathbb{N}$ and n sufficiently large, any r -colouring of the edges of K_n contains a monochromatic copy of a clique on t vertices. In other words, Ramsey's Theorem states that a monochromatic colouring is an unavoidable colour pattern of a clique. It is therefore natural to ask whether there are other colour patterns of a clique which are unavoidable. Of course, the answer is negative since the colouring of the large clique might itself be monochromatic. One may then restrict the attention to a proper subset of the collection of all r -edge colourings of a clique and ask whether within that class there are other colour patterns which are unavoidable.

First consider the case of $r = 2$. We say that a 2-edge coloured K_{2t} is $(2, t)$ -unavoidable if one of the colours induces either a clique on t vertices or two vertex disjoint cliques on t vertices. Bollobás asked whether for every $\varepsilon > 0$ and $t \in \mathbb{N}$, there is $n(\varepsilon, t)$ such that for every $n \geq n(\varepsilon, t)$, every 2-coloured clique on n vertices, where each colour appears in at least $\varepsilon \binom{n}{2}$ edges must necessarily contain a $(2, t)$ -unavoidable coloured K_{2t} . This was confirmed by Cutler and Montágh [6]. Observe that the colour patterns which are defined as $(2, t)$ -unavoidable are the only ones which one could hope to find here, since the given colouring of K_n might itself be of this form.

Later, Fox and Sudakov [8] proved $n(\varepsilon, t) = \varepsilon^{-O(t)}$, which is tight up to the implied constant. Instead of imposing both colours to appear in a positive proportion of the edges, we could ask the following.

Problem 2.1. *Let $t \in \mathbb{N}$ and $n_0(t)$ be a sufficiently large integer. Given $n \geq n_0$, how many edges $m(t, n)$ of each colour must exist in a 2-edge colouring of K_n to guarantee the existence of a $(2, t)$ -unavoidable pattern?*

Note that the Cutler and Montágh result implies that $m(t, n) = o(n^2)$, for fixed t as $n \rightarrow \infty$. Quite recently, Caro, Hansberg and Montejano [4] proved that for every $t \in \mathbb{N}$, there exists a $\delta = \delta(t)$ such that $m(t, n) = O(n^{2-\delta})$, following which the first author and Narayanan [10] showed that $\delta = 1/t$, by proving that for every $t \in \mathbb{N}$, there exists a constant $C(t)$ such that $m(t, n) \leq C(t)n^{2-1/t}$. It is not hard to see that this bound is tight up to the constant $C(t)$, conditional on a famous conjecture of Kővári-Sós-Turán [11], which asserts that $\text{ex}(n, K_{a,b}) = \Omega(n^{2-1/a})$ for all $b \geq a \geq 2$.

Very recently, Caro, Hansberg and Montejano have obtained various generalisations, as well as cases of unconditional sharpness, of this result, see [5] for the precise results. Here, we instead consider the generalisation of the above problem to r colours. The concept of a $(2, t)$ -unavoidable colour pattern was first generalised to r colours in [2]. Let G be a vertex and edge coloured complete graph with r colours. Moreover, suppose that all r colours are present and G is minimal with this property, i.e. no proper induced subgraph of G spans all r colours. Then, we say G is an r -minimal graph. The t -blowup of a vertex and edge coloured graph G on k vertices is obtained in the following natural way:

- Replace each coloured vertex of G by a clique of order t with every edge within it that colour;
- Replace each coloured edge of G by a complete bipartite graph $K_{t,t}$ with every edge that colour.

Finally, given $t \geq 2$, we call an edge-coloured clique G' (r, t) -unavoidable if G' is a t -blowup of *some* r -minimal graph. We denote by \mathcal{F}_t^r the set of all (r, t) -unavoidable graphs. Note that putting $r = 2$ recovers the definition of $(2, t)$ -unavoidable from earlier, and also that one can show G being r -minimal implies G has at most $2r$ vertices, and so \mathcal{F}_t^r is a finite set (see the argument in [2]).

Bowen, Lamaison and Müyesser [2] proved the generalisation of the result of Cutler and Montágh to r colours. Not long after, Bowen, Hansberg, Montejano and Müyesser [1] attempted to generalise the result of the first author and Narayanan to r colours, showing that for any $r, t \in \mathbb{N}$, any r -edge colouring of a sufficiently large graph on n vertices with $\Omega(n^{2-1/tr^r})$ edges in each colour contains a member of \mathcal{F}_t^r . They conjectured that $\Omega(n^{2-1/t})$ edges in each colour is sufficient. Our main result of this section is to confirm this conjecture.

Theorem 2.2. *For every $r, t \in \mathbb{N}$, there exists $C(r, t) > 0$ such that any r -edge colouring of a sufficiently large complete graph on n vertices with the property that any colour appears in at least $C(r, t)n^{2-1/t}$ edges must contain an (r, t) -unavoidable graph.*

2.1. Proof of Theorem 2.2. We require the following version of the dependent random choice lemma.

Lemma 2.3 ([9]). *For all $K, t \in \mathbb{N}$, there exists a constant C such that any graph with at least $Cn^{2-1/t}$ edges contains a set S of K vertices in which each subset $X \subseteq S$ with t vertices has a common neighbourhood of size at least K . Moreover the neighbourhoods may be chosen so that they are pairwise disjoint from each other and from S .*

We will also repeatedly apply Ramsey's Theorem; since we do not attempt to optimise the bound on the constant C which we will obtain, the following statement suffices for our proof. We write it exactly as we will apply it.

Lemma 2.4. *For all $r, s, t \in \mathbb{N}$ with $s \leq t$, there exists s', t' with $s' \leq t'$ such that however one r -colours the edges of two cliques $K_{s'}$ and $K_{t'}$ and the edges between them, one can find a monochromatic $K_s \subseteq K_{s'}$ and monochromatic $K_t \subseteq K_{t'}$ with all edges between K_s and K_t monochromatic (where we may have different colours for each of the different monochromatic parts).*

The idea of the proof is as follows. We initially apply the dependent random choice lemma to find large sets A_i , $i \in [r]$, such that every subset T of size t of each A_i forms a large monochromatic bipartite graph in colour i with some set $C_i(T)$. The aim will be to find subsets $D_i \subseteq A_i$, which together with $C_i(D_i)$ form a graph which will then easily be shown to contain an (r, t) -unavoidable colour pattern. First we apply Lemma 2.4 to the pairs (A_i, A_j) so that we have large monochromatic graphs between each pair. Now we get to the crucial step in our proof which results in the improvement on the bound given by Bowen et al; we apply Lemma 2.4 between A_j and *all* sets $C_i(T)$ with $i < j$. Only then do we fix a t -set D_j within A_j . We fix the sets D_j , $j = r, r-1, \dots, 1$ one at a time, each time applying Lemma 2.4 to $C_j(D_j)$ and A_i , $i < j$, before going on to fix D_{j-1} . Finally, we apply Lemma 2.4 to each pair $(C_i(D_i), C_j(D_j))$.

The large number of applications of Lemma 2.4 results in the A_i 's needing to be extremely large, and also increasing in size, i.e. $|A_i| = m_i$, with $1/m_r \ll \dots \ll 1/m_1$. However, crucially, we can skip the step required by Bowen et al. to obtain monochromatic subgraphs between the D_j 's and $C_i(T)$'s, which is the step which led to the unwanted r^r factor.

Note that by $a \ll b$ we mean that there exists a non-decreasing function $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that the statement holds for all a, b satisfying $a \leq f(b)$.

Proof of Theorem 2.2. Fix beforehand positive constants $1/m_r \ll \dots \ll 1/m_2 \ll 1/m_1 \ll 1/r, 1/t$ and let $1/C \ll 1/\ell \ll 1/r, 1/t$. Suppose that G is an r -edge coloured complete graph on n vertices such that every colour appears in at least $Cn^{2-1/t}$ edges. We wish to show that G contains an (r, t) -unavoidable graph.

First, using Lemma 2.3, we may find r pairwise disjoint sets A_1, \dots, A_r such that

- (i) For every $i \in [r]$, $|A_i| := m_i$,
- (ii) For every $i \in [r]$ and every subset $T \subset A_i$ of size t , there is a set $C_i(T)$ such that $(T, C_i(T))$ forms a complete bipartite graph in colour i , $|C_i(T)| = \ell$, $C_i(T)$ is vertex disjoint from $\bigcup_{i=1}^r A_i$, and the $C_i(T)$ are all pairwise disjoint from each other.

Now we proceed to the main stage of the argument, where we iteratively apply Lemma 2.4 to find subsets of the A_i 's and subsets of the $C_i(T)$'s which satisfy certain properties. For ease of notation, we will keep calling these sets A_i and $C_i(T)$. We also note that in every step the relations between the sizes of the new sets are maintained; that is, we make the m_i and ℓ sufficiently large so that the relations are maintained after each application of Lemma 2.4.

- (1) For each $i, j \in [r]$ with $i < j$, apply Lemma 2.4 to the pair (A_i, A_j) .
- (2) For each $j \in \{2, \dots, r\}$, for each $i \in [j-1]$, and for each subset T of size t of A_i , apply Lemma 2.4 to the pair $(A_j, C_i(T))$.
- (3) For j from r to 1 (start at r and decrease by 1 in the iteration), fix $D_j \subseteq A_j$ of size t , let $F_j := C_j(D_j)$, and for each $i \in [j-1]$, apply Lemma 2.4 to the pair (A_i, F_j) .
- (4) For each $i, j \in [r]$ with $i < j$, apply Lemma 2.4 to the pair (F_i, F_j) .

As mentioned, after each application of Lemma 2.4 we rename everything while maintaining the relations between the sizes of the sets; for example in (1), when we apply Lemma 2.4 to the pair (A_i, A_j) , what happens is the following: Ramsey's Theorem guarantees graphs $A'_i \subset A_i$, $A'_j \subset A_j$ such that there is a monochromatic complete bipartite graph between A'_i and A'_j , and also A'_i and A'_j are each monochromatic complete graphs. We have $|A'_i| = m'_i$, $|A'_j| = m'_j$ where we have

$$1/m_r \ll \dots \ll 1/m_{j+1} \ll 1/m'_j \ll 1/m_{j-1} \ll \dots \ll 1/m_{i+1} \ll 1/m'_i \ll 1/m_{i-1} \ll \dots \ll 1/m_1.$$

Then we relabel so that $A_i := A'_i$, $A_j := A'_j$, $m_i := m'_i$ and $m_j := m'_j$. We do similar relabellings for (2), (3) and (4).

Following steps (1) and (2), in addition to properties (i) and (ii) above, we have the following:

- (iii) For every $i, j \in [r]$ with $i < j$, (A_i, A_j) forms a complete monochromatic bipartite graph.
- (iv) For every $i, j \in [r]$ with $i < j$ and every subset T of size t of A_i , $(A_j, C_i(T))$ forms a complete monochromatic bipartite graph.

Now we examine the result of step (3). By construction, for each $j \in [r]$, we have

- (v) By (1), D_j is a subset of A_j of size t which is monochromatic.
- (vi) By (ii), (D_j, F_j) forms a complete monochromatic (in colour j) bipartite graph.
- (vii) By (iv), $(D_j, C_i(T))$ forms a monochromatic complete bipartite graph for every $i < j$ and every $T \subset A_i$ of size t . In particular, D_j will form a monochromatic complete bipartite graph with F_i , once D_i and hence F_i is chosen.
- (viii) By (3), for each $k > j$, (A_j, F_k) and therefore (D_j, F_k) forms a monochromatic complete bipartite graph.

Finally we examine the result of step (4). We have

- (ix) For all $i \in [r]$, F_i is monochromatic.
- (x) For every $i, j \in [r]$ with $i < j$, (F_i, F_j) forms a complete monochromatic bipartite graph.

The overall result is that the subgraph H of G induced by $\{D_i, F_i : i \in [r]\}$ forms a t -blowup of a vertex and edge coloured clique J on $2r$ vertices where each of the r colours appears at least once. Therefore by deleting vertices from J if necessary, we can obtain an r -minimal graph. Hence G contains as a subgraph a member of \mathcal{F}_t^r , as required. \square

3. CONCLUDING REMARKS

We have shown that for any 3-chromatically connected graph H , and any $s \in \mathbb{N}$, there are graphs G for which $G \xrightarrow{r} H$ and yet $G[s] \not\xrightarrow{r} H[2]$. On the other hand, we showed that for forests there is no dependence on the ground graph G .

The first natural problem is to classify all finite graphs for which their blowup Ramsey number does not depend on the ground graph. We conjecture that forests are the only such graphs.

Conjecture 3.1. *Let H be a graph containing a cycle. Then, there is a t such that for every s , there is a graph G where $G \xrightarrow{r} H$ but $G[s] \not\xrightarrow{r} H[t]$.*

It would already be interesting to prove the conjecture for cycles C_k with $k \geq 4$ (which are not 3-chromatically connected).

Regarding Theorem 1.6 it is natural to ask whether the function $f(T, 2, t)$ is exponential in t , as the result of Fox, Luo and Wigderson suggests. A slightly more careful analysis of our argument shows that we may take $f(T, 2, t) = 2^{2^{O(t)}}$ but we suspect single exponential might be the truth.

Problem 3.2. *Is $f(T, 2, t) = 2^{O(t)}$?*

REFERENCES

- [1] M. Bowen, A. Hansberg, A. Montejano, and A. Müyesser. Colored unavoidable patterns and balanceable graphs. *arXiv preprint arXiv:1912.06302*, 2019. [9](#)
- [2] M. Bowen, A. Lamaison, and A. Müyesser. Finding unavoidable colorful patterns in multicolored graphs. *Electron. J. Combin.*, 27(4), 2020. # P4.4. [8](#), [9](#)
- [3] S. A. Burr, P. Erdős, R. J. Faudree, C. C. Rousseau, and R. H. Schelp. Ramsey-minimal graphs for star-forests. *Discrete Math.*, 33(3):227–237, 1981. [2](#)
- [4] Y. Caro, A. Hansberg, and A. Montejano. Unavoidable chromatic patterns in 2-colorings of the complete graph. *J. Graph Theory*, 97(1):123–147, 2021. [8](#)
- [5] Y. Caro, A. Hansberg, and A. Montejano. The evolution of unavoidable bi-chromatic patterns and extremal cases of balanceability. *arXiv preprint arXiv:2204.04269*, 2022. [8](#)
- [6] J. Cutler and B. Montágh. Unavoidable subgraphs of colored graphs. *Discrete Math.*, 308:4396–4413, 2008. [8](#)
- [7] J. Fox, S. Luo, and Y. Wigderson. Extremal and Ramsey results on graph blowups. *J. Comb.*, 12(1):1–15, 2021. [2](#)
- [8] J. Fox and B. Sudakov. Unavoidable patterns. *J. Combin. Theory Ser. A*, 115:1561–1569, 2008. [8](#)
- [9] J. Fox and B. Sudakov. Dependent random choice. *Random Structures Algorithms*, 38:68–99, 2011. [9](#)
- [10] A. Girão and B. Narayanan. Turán theorems for unavoidable patterns. In *Math. Proc. Cambridge Philos. Soc.*, pages 1–20. Cambridge University Press, 2021. [8](#)

- [11] T. Kővári, V. T. Sós, and P. Turán. On a problem of K. Zarankiewicz. *Colloquium Math.*, 3:50–57, 1954. [8](#)
- [12] J. Nešetřil and V. Rödl. Sparse Ramsey graphs. *Combinatorica*, 4(1):71–78, 1984. [3](#)
- [13] V. Souza. Blowup Ramsey numbers. *Eur. J. Comb.*, 92:103238, 2021. [1](#)