

# Griffiths inequalities for the $O(N)$ -spin model

Benjamin Lees\*

University of Bristol

## Abstract

We prove Griffiths inequalities for the  $O(N)$ -spin model with inhomogeneous coupling constants and external magnetic field for any  $N \geq 2$ .

## 1 Introduction

Correlation inequalities for models in statistical mechanics were first presented by Griffiths [10] for the Ising model. These *Griffiths' inequalities* were extended by Ginibre [9] to a more general framework that also includes quantum spin systems. This framework includes the classical XY (or  $O(2)$ -spin) model [9, 11] and the spin- $\frac{1}{2}$  XY model [4]. A Griffiths inequality was also proven for the classical Heisenberg (or  $O(3)$ -spin) ferromagnet in the case of homogeneous rotations and for four-component spin systems such as the  $|\phi|^4$  lattice euclidean field [8, 11]. We direct the reader to [5] for an overview of results in these cases. Griffiths inequalities are useful tools for proofs of the existence of infinite volume limit of correlation functions and monotonicity of spontaneous magnetisation and also allowing comparisons of aspects, such as the critical temperature, of models with different spin and/or spatial dimension. The extension of these inequalities to the  $O(N)$ -spin model and to higher spin quantum XY models or quantum Heisenberg models has been an important problem.

The representation of spin models in terms of random geometric objects has enjoyed increasing interest in recent years and has led to new results and new proofs for several important systems. The idea finds its origins in the work of Symanzik [16] and Brydges, Fröhlich and Spencer [6] for the random walk expansion of spin systems. Later, Aizenman [1] developed techniques for the *random current* representation of the Ising model, including the famous and powerful *switching lemma* for this model. There are analogous representations for quantum models, such as the representation of the *Heisenberg ferromagnet* in terms of sequences of random transpositions which were used by Tóth to bound the pressure of this model, or the representation of the *Heisenberg anti-ferromagnet* as a type of *random cluster* model by Aizenman and Nachtergaele [2]. It was later shown by Ueltschi [18] that these two representations can be combined into a *random loop* model that reduces to the previous models for a particular choice of the model's parameters. This representation has enjoyed significant interest in the literature.

In this article we will make use of a representation of the  $O(N)$ -spin model in terms of *random paths* (walks and loops) that was introduced in [12] and is closely related to the model introduced in [3]. The difference with previous representations is that realisations can be defined purely in terms of objects local to individual edges and vertices. This distinction has allowed the use important tools such as reflection positivity [12, 15, 17], a proof of exponential decay of transverse correlations in the presence of an external magnetic field [13] that goes beyond results proven using the well-known Lee-Yang method, and a new proof of the BKT phase transition for the  $O(2)$ -spin model in two dimensions [14].

---

\*Heilbronn Institute for Mathematical Research and School of Mathematics, University of Bristol. Email: *benjamin.lees@bristol.ac.uk*

We prove a Griffiths inequality for the  $O(N)$ -spin model for any  $N \geq 2$  on an arbitrary simple graph with inhomogeneous ferromagnetic coupling constants that may vary in different spin-directions. To the author's knowledge, this is the first proof of a Griffiths inequality for the  $O(N)$ -spin model with  $N \geq 4$  and the first proof for the case of inhomogeneous rotations with  $N \geq 3$ . We are also able to treat the case of an inhomogeneous external field. This is achieved by using the connection of this system with the random path model. By presenting the random path model in a slightly different way to the presentation in [12] we obtain a Griffiths inequality by using the proof of the switching lemma for the random current model presented in [1] as a template. Unfortunately, where the proof for random currents identifies a bijection between two sets, we are only able to identify an injection from the first set into the set of interest. It would be extremely useful to identify a natural subset of the set of interest which is in bijection with the first set and hence obtain a "true" switching lemma for  $O(N)$ -spins.

## 1.1 Model and main result

Here we define the  $O(N)$ -spin model on an arbitrary graph with inhomogeneous coupling constants. In Section 4 we introduce the model with an external field. Consider a finite simple graph  $G = (V, E)$  and, for  $N \in \mathbb{N}_{>0}$ , define the configuration space  $\Omega_{G,N} := (\mathbb{S}^{N-1})^V$ , where  $\mathbb{S}^{N-1} \subset \mathbb{R}^N$  is the  $N-1$  dimensional unit sphere. For  $\varphi \in \mathbb{S}^{N-1}$  we denote by  $\varphi^i \in \mathbb{R}$  the  $i^{\text{th}}$  component of  $\varphi$ . For a collection of non-negative constants  $\mathbf{J} = (J_e^1, \dots, J_e^N)_{e \in E}$  we introduce the hamiltonian function acting on  $\varphi = (\varphi_x)_{x \in V} \in \Omega_{G,N}$ ,

$$H_{G,N,\mathbf{J}}^{\text{free}}(\varphi) = - \sum_{\{x,y\} \in E} \sum_{i=1}^N J_{\{x,y\}}^i \varphi_x^i \varphi_y^i. \quad (1.1)$$

We define the expectation operator  $\langle \cdot \rangle_{G,N,\mathbf{J}}^{\text{free}}$  acting on  $f : \Omega_{G,N} \rightarrow \mathbb{R}$  by

$$\langle f \rangle_{G,N,\mathbf{J}}^{\text{free}} = \frac{1}{Z_{G,N,\mathbf{J}}^{\text{free}}} \int_{\Omega_{G,N}} d\varphi f(\varphi) e^{-H_{G,N,\mathbf{J}}^{\text{free}}(\varphi)}, \quad (1.2)$$

where  $d\varphi = \prod_{x \in V} d\varphi_x$  is a product measure with  $d\varphi_x$  the uniform measure on  $\mathbb{S}^{N-1}$  and  $Z_{G,N,\mathbf{J}}^{\text{free}}$  is a normalising constant that ensures  $\langle 1 \rangle_{G,N,\beta} = 1$ . Note that this definition includes the case of *free* and *periodic* boundary conditions (for example a torus  $\mathbb{Z}^d/L\mathbb{Z}^d$  with nearest neighbour edges). We may also consider other boundary conditions such as the "+" boundary condition which we define here in analogy with the Ising model.

Suppose that  $G$  is a subgraph of some larger (possibly infinite) graph  $\mathcal{G}$ . Let  $\partial V \subset V$  be the set of vertices that share an edge with some vertex of  $\mathcal{G} \setminus G$  and let  $\partial E$  be this set of edges. We also denote by  $\partial^{\text{ext}} V \subset V^c$  the set of vertices in  $V^c$  that share an edge with a vertex in  $V$  (the exterior vertex boundary). We define the hamiltonian with *+ boundary conditions* as

$$H_{G,N,\mathbf{J}}^+(\varphi) = - \sum_{\{x,y\} \in E} \sum_{i=1}^N J_{\{x,y\}}^i \varphi_x^i \varphi_y^i - \sum_{\{x,y\} \in \partial E: x \in V} J_{\{x,y\}}^1 \varphi_x^1. \quad (1.3)$$

We interpret this as having all spins outside of  $V$  fixed as  $(1, 0, \dots, 0)$ , i.e. fully aligned along the first spin-direction, so that  $\sum_{i=1}^N J_{\{x,y\}}^i \varphi_x^i \varphi_y^i = J_{\{x,y\}}^1 \varphi_x^1$  for edges  $\{x,y\} \in \partial E$  with  $x \in V, y \in V^c$ . We have analogous expectations to the case of free or periodic boundary. For  $f : \Omega_{G,N} \rightarrow \mathbb{R}$

$$\langle f \rangle_{G,N,\mathbf{J}}^+ = \frac{1}{Z_{G,N,\mathbf{J}}^+} \int_{\Omega_{G,N}} d\varphi f(\varphi) e^{-H_{G,N,\mathbf{J}}^+(\varphi)}. \quad (1.4)$$

Our main result concerns correlations between the components of spins. For any  $x \in V$ , define the random variable  $S_x : \Omega_{G,N} \mapsto \mathbb{S}^{N-1}$  representing the spin at  $x$  as,  $S_x(\varphi) := \varphi_x$ , moreover we represent its components as  $S_x = (S_x^1, \dots, S_x^N)$ . So, for example,  $S_x^1(\varphi) = \varphi_x^1$ .

**Theorem 1.1.** *Let  $G = (V, E)$  be a finite simple graph and  $N \in \mathbb{N}_{>1}$ . Assume that  $J_e^i \geq 0$  for every  $e \in E$ ,  $i \in [N]$ . For  $A, B \subset V$  and  $\eta \in \{\text{free}, +\}$  we have*

$$\left\langle \prod_{x \in A} S_x^1 \prod_{y \in A} S_y^1 \right\rangle_{G, N, \mathbf{J}}^\eta \geq \left\langle \prod_{x \in A} S_x^1 \right\rangle_{G, N, \mathbf{J}}^\eta \left\langle \prod_{y \in B} S_y^1 \right\rangle_{G, N, \mathbf{J}}^\eta.$$

This holds, for example, on  $\mathbb{Z}^d$  with finite range (not necessarily translation invariant) coupling constants. The following monotonicity with respect to coupling constants is a straightforward corollary of Theorem 1.1.

**Corollary 1.2.** *Let  $G = (V, E)$  be a finite simple graph and  $N \in \mathbb{N}_{>1}$ . Assume that  $J_e^i \geq 0$  for every  $e \in E$ ,  $i \in [N]$  and  $\eta \in \{\text{free}, +\}$ . For  $A \subset V$  and  $e \in E$  we have*

$$\frac{\partial}{\partial J_e^i} \left\langle \prod_{x \in A} S_x^1 \right\rangle_{G, N, \mathbf{J}}^\eta \geq 0.$$

**Remark 1.3.** *We could think of the free boundary case as simply the case of  $+$  boundary where  $J_e^1 = 0$  for every  $e \in \partial E$ , we encourage the reader to adopt this point of view if it is helpful. We can also prove Theorem 1.1 for more general boundary conditions, however this is very similar to the case of an inhomogeneous external magnetic field, which we present in Section 4. The random path model used for the proof has a corresponding extension to this case, as explained in [13]. Because we already need to make slight modifications to the proof for the cases  $\eta = \text{free}$  or  $\eta = +$  and the cases  $A \cap B = \emptyset$  or  $A \cap B \neq \emptyset$ , we will describe the  $O(N)$ -spin model with external field, the corresponding changes to the random path model, and the differences in the proof in Section 4. In this way we avoid the need for the extra modifications to be present throughout the article and can focus on the case of no external field, which already contains all the important ideas of the proof.*

## 1.2 Proof method

To prove Theorem 1.1 we will use the connection between the  $O(N)$ -spin model and the random path model [12]. We will present the random path model in a slightly different way that is convenient for proofs. Both presentations are equivalent and hence reduce to the random current representation of the Ising model for  $N = 1$ . With this in mind, we will attempt to recreate the proof of the switching lemma for random currents and find that, at the point of the proof of the switching lemma where a bijection is identified, we are only able to identify an injection. This leads to the inequality in Lemma 3.1 from which Theorem 1.1 follows easily.

## Notation

$\mathbb{N}$	$\{0, 1, \dots\}$
$\mathbb{N}_{>0}$	$\{1, 2, \dots\}$
$2\mathbb{N}$	the set of even integers in $\mathbb{N}$
$2\mathbb{N} + 1$	the set of odd integers in $\mathbb{N}$
$G = (V, E)$	an undirected, simple, finite graph
$A\Delta B$	the symmetric difference of set $A$ and $B$
$N \in \mathbb{N}_{>0}$	the number of colours or spin components
$[N]$	$\{1, \dots, N\}$
$\mathcal{M}_G^N$	the set of $(m^1, \dots, m^N) \in \mathbb{N}^{E \times [N]}$
$\mathbf{m}$	an element $(m^1, \dots, m^N)$ of $\mathcal{M}_G^N$
$\mathcal{P}_G(\mathbf{m})$	the set of pairings of configurations for $\mathbf{m} = (m^1, \dots, m^N)$
$n_x^i(\mathbf{m})$	the local time of $i$ -objects at $x$
$n_x(\mathbf{m})$	$\sum_{i=1}^N n_x^i(\mathbf{m})$
$m_z^j$	$\sum_{e \ni z} m_e^j$
$Z_{G,N,\mathbf{J},U}(A)$	the total weight of configurations such that $m_z^j$ is odd iff $j = 1$ and $z \in A$
$\mathbb{G}_{G,N,\mathbf{J},U}(A)$	the correlation function for $A \subset V$ in the random path model

## 2 The Random Path Model

In this section we introduce the Random Path Model (RPM) in two equivalent ways. The first way was introduced in [12] and was shown to include many interesting models for specific choices of its parameters. The second way differs only slightly and will be more convenient for proofs. In both cases we will introduce the model for the case of free or periodic boundary conditions and then comment on the very minor difference needed to make the connection with  $+$  boundary conditions.

The RPM differs from previous representations of  $O(N)$ -spins in terms of loops and walks, such as the one introduced by Symanzik [16] and studied by Brydges, Fröhlich and Spencer [6], due to the presence of *colourings* of our basic objects into  $N$  colours and *pairings*.

The first description of the Random Path Model introduced in [12] has colourings and pairings “built-in”. The purpose of introducing this representation is to allow us to take advantage of the proven connections between this model and the  $O(N)$ -spin model that we require. The reader is encouraged to consult [12] or [13] for an alternative introduction of the model.

The case  $N = 1$  reduces to the random current representation of the Ising model where pairings are not present (as we will see below, the pairings are perfectly cancelled by the *vertex weight function* in this case).

### 2.1 The Random Path Model with Colourings and Pairings

Let  $G = (V, E)$  be a finite simple graph and  $N \in \mathbb{N}_{>0}$  be the *number of colours*. RPM configurations will correspond to undirected walks and loops, collectively referred to as *paths*, taking colours in  $[N] := \{1, \dots, N\}$ .

First, define  $\mathcal{M}_G := \mathbb{N}^E$ . An element  $m = (m_e)_{e \in E} \in \mathcal{M}_G$  will be thought of as a *collection of links* on  $E$  with  $m_e$  referred to as the *number of links* on  $e \in E$ . We say a link is incident to  $x \in V$  if it is on an edge incident to  $x$ .

For  $m \in \mathcal{M}_G$ , a *colouring*  $c = (c_e)_{e \in E}$  is a collection of functions, one for each  $e \in E$ , with  $c_e : \{1, \dots, m_e\} \mapsto [N]$  interpreted as an assignment of an integer (colour) in  $[N]$  to each link. More precisely, if we use  $(e, p)$  to represent the  $p^{\text{th}}$  link on  $e$  then  $c((e, p)) \in [N]$  is the colour of the  $p^{\text{th}}$  link on  $e \in \mathcal{E}$ . A link with colour  $i \in [N]$  is called an  $i$ -link and we denote by  $m_e^i = m_e^i(m, c)$  the number




Figure 2.1: Left: an example of a collection of links on a graph consisting of a  $3 \times 3$  box with nearest neighbour edges, Centre: an example of a collection of links and a colouring of the links into two colours (red and blue). Right: an example of a RPM configuration consisting of a collection of links, a colouring, and a pairing. Pairs are denoted by lines joining links to form continuous paths through vertices. This example consists of 4 red walks, 1 red loop and 2 blue loops.

of  $i$ -links on  $e$  for a pair  $(m, c)$  of a collection of links and a colouring of those links. We denote by  $\mathcal{C}_G(m)$  the set of all possible colourings  $c = (c_e)_{e \in E}$  for  $m$ .

Given a collection of links  $m \in \mathcal{M}_G$ , and a colouring  $c \in \mathcal{C}_G(m)$ , we say  $\pi = (\pi_x)_{x \in V}$  is a *pairing* of  $(m, c)$  if, for each  $x \in V$ ,  $\pi_x(m, c)$  is a partition of the links incident to  $x$  into sets with at most two links each such that if two links are in the same set of the partition, then they have the same colour. If a set contains two links then we say those links are *paired at  $x$*  and think of them being joined (or *wired*) together at  $x$  to form part of a continuous path passing through  $x$ . If a set contains only one link then we say this link is *unpaired at  $x$* . A link can be paired to at most two other links, one at each end point of its edge. We denote by  $\mathcal{P}_G(m, c)$  the set of all pairings for  $m \in \mathcal{M}_G$  and  $c \in \mathcal{C}_G(m)$ .

We remark here that, in general, there are many possible colourings of  $m \in \mathcal{M}_G$  and given a colouring  $c \in \mathcal{C}_G(m)$  there are many possible pairings of  $(m, c)$ , in particular the “empty pairing” that leaves all links unpaired at every vertex.

A configuration for the random path model is a triple  $w = (m, c, \pi)$  such that  $m \in \mathcal{M}_G$ ,  $c \in \mathcal{C}_G(m)$ , and  $\pi \in \mathcal{P}_G(m, c)$ . Let  $\mathcal{W}_G$  be the set of all random path configurations on  $G$ . As we can see from the example in Figure 2.1, any  $w \in \mathcal{W}_G$  can be viewed as a collection of closed or open *paths*. A *path* is a maximal set of links such that each link in the path is paired to at least one other link in the path. It follows that all links in a path are the same colour. For each path, either every link in the path is paired at both end points, in which case we call the path *closed*, or there are two link end points that are not paired in which case we call the path *open*. We will often refer to closed paths as *loops* and open paths as *walks*, we also refer to the vertices at which an open path has unpaired links as its *end points*. We note that an open path may consist of a single link, unpaired at both end points, or start and end at the same vertex  $z \in V$ , in which case  $z$  is counted as both end points, this case is distinct from the case when the unpaired links at  $z$  are paired to form a loop passing through  $z$ . We refer to paths/loops/walks whose links have colour  $i \in [N]$  as  $i$ -paths/loops/walks, respectively. By a slight abuse of notation, we will also view  $m : \mathcal{W}_G \mapsto \mathcal{M}_G$  as a function such that, for  $w' = (m', c', \pi')$ ,  $m(w') = m'$ .

Let  $u_x^i(w)$  be the number of  $i$ -links incident to  $x$  which are unpaired at  $x$  (this is also the number of walk end-points at  $x$ ). Let  $v_x^i(w)$  be the number of pairs of  $i$ -links incident to  $x$ . Note that  $2v_x^i(w) + u_x^i(w)$  is the number of  $i$ -links incident to  $x$  in  $w$ .

Moreover, let

$$n_x^i(w) := v_x^i(w) + u_x^i(w) \tag{2.1}$$

be the *local time of  $i$ -paths at  $x$* . Unpaired end-points of links incident to  $x$  and pairs of paired links incident to  $x$  both contribute +1 to the local time. In words,  $n_x^i(w)$  is the number of times an  $i$ -path

visits  $x$ , where a visit means that a path (open or closed) passes through  $x$  or an open path ends at  $x$ . We also define

$$n_x(w) := \sum_{i=1}^N n_x^i(w) \quad (2.2)$$

to be the local time of paths of all colours at  $x$ .

We are interested in a particular subset of  $\mathcal{W}_G$  which is relevant for understanding the correlations of the  $O(N)$ -spin model. Consider the set of configuration  $\mathcal{W}'_G \subset \mathcal{W}_G$  such that the only unpaired links have colour 1. Hence, any configuration in  $\mathcal{W}'_G$  consists of open paths (walks) of colour 1, and closed paths (loops) of any colour.

We may also consider a model with boundary conditions. For  $\mathcal{G} \supset G$  we define random path configurations on  $G \cup \partial E$  as above except that we do not specify pairings on the end points of  $\partial E$  that lie in  $V^c$ . Intuitively, we think of random path configurations where paths are allowed to leave  $G$  at the boundary, but we do not track where they go after this. We denote by  $\mathcal{W}'_{G \cup \partial E}$  the set of all such configurations where only 1-links may be unpaired on  $V$  and no links are paired on  $V^c$ . In the definitions, to make the connection with  $+$  boundary conditions for the  $O(N)$ -spin model, we will make the restriction that only 1-paths may leave  $G$ .

Now we are ready to define our measure on configurations.

**Definition 2.1.** Let  $\mathcal{W}'_G$  be the set of configurations  $w \in \mathcal{W}_G$  such that  $u_x^2(w) = \dots = u_x^N(w) = 0$  for every  $x \in V$ . Given  $N \in \mathbb{N}_{>0}$  and non-negative coupling constants  $\mathbf{J} = (J_e^1, \dots, J_e^N)_{e \in E}$  define the measure  $\tilde{\mu}_{G,N,\mathbf{J},U}^{\text{free}}$  on  $\mathcal{W}'_G$  by

$$\tilde{\mu}_{G,N,\beta,U}^{\text{free}}(w) := \prod_{e \in E} \frac{\prod_{i=1}^N (J_e^i)^{m_e^i}}{m_e!} \prod_{x \in V} U_x(w), \quad w \in \mathcal{W}'_G, \quad (2.3)$$

where  $U_x(w)$  is a vertex weight function depending only on the links incident to  $x$  and their pairing. In the case where  $G \subset \mathcal{G}$ , we also define a measure on configurations on  $G \cup \partial E$  that is related to the case of  $+$  boundary conditions

$$\tilde{\mu}_{G,N,\beta,U}^+(w) := \prod_{e \in E \cup \partial E} \frac{\prod_{i=1}^N (J_e^i)^{m_e^i}}{m_e!} \prod_{x \in V} U_x(w) \prod_{e \in \partial E} \prod_{i=2}^N \mathbb{1}_{\{m_e^i=0\}}, \quad w \in \mathcal{W}'_{G \cup \partial E}. \quad (2.4)$$

The difference with  $\tilde{\mu}_{G,N,\beta,U}^{\text{free}}$  is that we allow 1-links on edges of  $\partial E$  (and hence for 1-paths to leave  $G$ ). However we do not allow any links of colour  $2, \dots, N$  on  $\partial E$ .

The central quantities of interest are *correlation functions*. We define this quantity for the first description of the random path model. Below we will define the analogous quantity for the second description of the random path model.

**Definition 2.2.** For  $A \subset V$ , define  $\tilde{\mathcal{S}}(A)$  to be the set of configurations  $w$  in  $\mathcal{W}'_G$  for the free or periodic boundary case or in  $\mathcal{W}'_{G \cup \partial E}$  for the  $+$  boundary case such that  $u_z^1(w) = 1$  for every  $z \in A$  and  $u_z^1(w) = 0$  for every  $z \in V \setminus A$ .

Further, for  $B \subset V$  define  $\tilde{\mathcal{S}}(A; B)$  to be the set of configurations  $w$  in  $\mathcal{W}'_G$  for the free or periodic boundary case or in  $\mathcal{W}'_{G \cup \partial E}$  for the  $+$  boundary case such that  $u_z^1(w) = 1$  for every  $z \in A \Delta B$ ,  $u_z^1(w) = 2$  for every  $z \in A \cap B$ , and  $u_z^1(w) = 0$  for every  $z \in V \setminus A \cup B$ .

For  $\eta \in \{\text{free}, +\}$ , we define  $\tilde{Z}_{G,N,\mathbf{J},U}^\eta(A) = \tilde{\mu}_{G,N,\mathbf{J},U}^\eta(\tilde{\mathcal{S}}(A))$ ,  $\tilde{Z}_{G,N,\mathbf{J},U}^\eta(A; B) = \tilde{\mu}_{G,N,\mathbf{J},U}^\eta(\tilde{\mathcal{S}}(A; B))$  and  $\tilde{Z}_{G,N,\mathbf{J},U}^{\text{loop},\eta} = \tilde{\mu}_{G,N,\mathbf{J},U}^\eta(\tilde{\mathcal{S}}(\emptyset))$ . Finally, we define the correlation functions by

$$\mathbb{G}_{G,N,\mathbf{J},U}^\eta(A) := \frac{\tilde{Z}_{G,N,\mathbf{J},U}^\eta(A)}{\tilde{Z}_{G,N,\mathbf{J},U}^{\text{loop},\eta}}, \quad \mathbb{G}_{G,N,\mathbf{J},U}^\eta(A; B) := \frac{\tilde{Z}_{G,N,\mathbf{J},U}^\eta(A; B)}{\tilde{Z}_{G,N,\mathbf{J},U}^{\text{loop},\eta}}.$$

We call the cases where  $|A| = 2$  two-point functions. Note that if  $\eta = \text{free}$  and  $|A|$  is odd then  $\tilde{S}(A) = \emptyset$ . If  $A \cap B = \emptyset$  then  $\mathbb{G}_{G,N,\mathbf{J},U}^\eta(A; B) = \mathbb{G}_{G,N,\mathbf{J},U}^\eta(A \cup B)$ .

For  $A \subset V$ ,  $\tilde{Z}_{G,N,\mathbf{J},U}^\eta(A)$  corresponds to the total measure of configurations with coloured loops and 1-paths with end points at vertices of  $A$ , such that each  $z \in A$  has precisely one 1-walk end point. In particular, the case  $A = \{x, y\}$  with  $x \neq y$  and  $\eta = \text{free}$  is the measure of configurations with coloured loops and a single 1-walk with end points  $x$  and  $y$ . This quantity is relevant for the correlation  $\langle \prod_{x \in A} S_x^1 \rangle_{G,N,\mathbf{J}}^\eta$ . For  $A, B \subset V$ ,  $\tilde{Z}_{G,N,\mathbf{J},U}^\eta(A; B)$  corresponds to the total measure of configurations with coloured loops and 1-paths with a single 1-walk end point at vertices of  $A \Delta B$ , and two 1-walk end points at vertices of  $A \cap B$ . In the case  $A = B = \{x\}$  this reduces to  $\mu_{G,N,\mathbf{J},U}(\mathcal{R}_x)$  as defined in Definition 2.2 of [12]. This quantity is relevant for the correlation  $\langle \prod_{x \in A} S_x^1 \prod_{y \in B} S_y^1 \rangle_{G,N,\mathbf{J}}^\eta$  when  $A \cap B \neq \emptyset$ .

The next proposition connects the correlation functions defined above to the spin correlations of the  $O(N)$ -spin model. For  $\langle \prod_{x \in A} S_x^1 \rangle_{G,N,\mathbf{J}}^\eta$ , the case of homogeneous coupling constants  $J_e^i = \beta \geq 0$  for all  $e \in E$ ,  $i \in [N]$  and free or periodic boundary conditions, the proof can be found in [12] for the case  $|A| = 2$  and in [13] for the general case, even with an external field. The extension to inhomogeneous coupling and + boundary condition is straightforward. For  $\langle \prod_{x \in A} S_x^1 \prod_{y \in B} S_y^1 \rangle_{G,N,\mathbf{J}}^\eta$ , this is a straightforward extension of the previous case by using the case  $\langle (S_x^1)^2 \rangle_{G,N,\mathbf{J}}^\eta$  dealt with in [12].

**Proposition 2.3** (Lees-Taggi 2020). *Let  $G = (V, E)$  be a finite, simple graph. Let  $N \in \mathbb{N}_{>0}$  and  $\mathbf{J} = (J_e^1, \dots, J_e^N)_{e \in E}$  be non-negative coupling constants. Let the vertex weight function introduced in Definition 2.2 be given by  $U_x^{(N)}(w) := \mathcal{U}^{(N)}(n_x(w))$  with*

$$\mathcal{U}^{(N)}(r) := \frac{\Gamma(\frac{N}{2})}{2r\Gamma(r + \frac{N}{2})}, \quad r \in \mathbb{N}. \quad (2.5)$$

For  $A, B \subset V$  and  $\eta \in \{\text{free}, +\}$  we have that,

$$\mathbb{G}_{G,N,\mathbf{J},U^{(N)}}^\eta(A) = \left\langle \prod_{x \in A} S_x^1 \right\rangle_{G,N,\mathbf{J}}^\eta, \quad \mathbb{G}_{G,N,\mathbf{J},U^{(N)}}^\eta(A; B) \leq \left\langle \prod_{x \in A} S_x^1 \prod_{y \in B} S_y^1 \right\rangle_{G,N,\mathbf{J}}^\eta. \quad (2.6)$$

The inequality for  $\mathbb{G}_{G,N,\mathbf{J},U^{(N)}}^\eta(A; B)$  is due to the possible presence of “degenerate paths” with 0 edges and hence two end-points on the same vertex of  $A \cap B$  when expanding the right side as in [12, 13].

**Remark 2.4.** *In order to obtain a Griffiths inequality when  $A \cap B \neq \emptyset$  we do not need to worry about the degenerate paths mentioned above as we prove the inequality  $\mathbb{G}_{G,N,\mathbf{J},U^{(N)}}^\eta(A) \mathbb{G}_{G,N,\mathbf{J},U^{(N)}}^\eta(B) \leq \mathbb{G}_{G,N,\mathbf{J},U^{(N)}}^\eta(A; B)$ , which is sufficient.*

In the case  $N = 1$  we find that

$$\mathcal{U}^{(1)}(r) = 1/(2r - 1)!!$$

where  $(2r - 1)!!$  is the number of ways to pair  $2r$  objects, this cancels out the contribution from pairings and leads to the random current representation of the Ising model.

## 2.2 The Random Path Model with pre-coloured links

We now introduce a different description of the RPM, there are many similarities with the previous description. We will focus on the case relevant for  $\langle \prod_{x \in A} S_x^1 \rangle_{G,N,\mathbf{J}}^\eta$  and then in Remark 2.8 comment on the changes needed to make the connection with  $\langle \prod_{x \in A} S_x^1 \prod_{y \in B} S_y^1 \rangle_{G,N,\mathbf{J}}^\eta$ . The changes are conceptually simple but result in cumbersome extra notation which, in the author’s opinion, detracts too much from clarity.

Let  $G = (V, E)$  be a finite simple graph and  $N \in \mathbb{N}_{>0}$  be the *number of colours*. We introduce “pre-coloured” link configurations by defining  $\mathcal{M}_G^N := \mathbb{N}^{E \times [N]}$ . An element  $\mathbf{m} = (m^1, \dots, m^N) \in \mathcal{M}_G^N$  is a tuple of  $N$  *collections of links* on  $E$  with  $m^i = (m_e^i)_{e \in E}$  the *collection of  $i$ -links* and  $m_e^i$  the *number of  $i$ -links* on  $e \in E$ . As above, we say a link is incident to  $z \in V$  if it is on an edge incident to  $z$ . We also define

$$m_e = \sum_{i=1}^N m_e^i. \quad (2.7)$$

Given a collection  $\mathbf{m} = (m^1, \dots, m^N) \in \mathcal{M}_G^N$  we again define a pairing  $\pi = (\pi_z)_{z \in V}$  to be a collection of functions such that  $\pi_z(\mathbf{m})$  is a partition of the links incident to  $z$  into sets of at most two links each such that if two links are in the same set, then they belong to the same  $m^i$ . As above we say that two links are *paired* at  $z \in V$  if they are in the same set of  $\pi_z(\mathbf{m})$  and links are *unpaired* at  $z$  if they are in a set by themselves in  $\pi_z(\mathbf{m})$ . Denote by  $\mathcal{P}_G(\mathbf{m})$  the set of all pairings of  $\mathbf{m} \in \mathcal{M}_G^N$ . As above, a pair  $(\mathbf{m}, \pi)$  where  $\mathbf{m} \in \mathcal{M}_G^N$  and  $\pi \in \mathcal{P}_G(\mathbf{m})$  has a natural interpretation as a collection of open and closed paths taking colours in  $[N]$ .

Configurations in this description are simply the tuples  $\mathbf{m} = (m^1, \dots, m^N) \in \mathcal{M}_G^N$ , they do not have a specific pairing “built in” and so to make the connection between correlations for the two descriptions a sum over pairings will appear in the measure of the model. For this reason we need to immediately restrict the types of pairings we will sum over in order to recover correlations of the  $O(N)$ -spin model. Define  $P_G(\mathbf{m}) \subset \mathcal{P}_G(\mathbf{m})$  to be the set of pairings such that at most one link of each colour is unpaired at each  $z \in V$ . We see from this that for a pairing in  $P_G(\mathbf{m})$  there is an unpaired  $i$ -link at  $z \in V$  if and only if  $\sum_{e \ni z} m_e^i \in 2\mathbb{N} + 1$ . It hence makes sense to define, for  $z \in V$ ,  $i \in [N]$ ,

$$m_z^i := \sum_{e \ni z} m_e^i, \quad m_z = \sum_{i=1}^N m_z^i. \quad (2.8)$$

Pairings in  $P_G(\mathbf{m})$  are *maximal* in the sense that they have as many paired links as possible at each vertex. Because we will only consider pairings in  $P_G(\mathbf{m})$  we can make an alternate definition of *local time* which is consistent with the definition above in this case. For  $z \in V$  and  $i \in [N]$  we define

$$n_z^i(\mathbf{m}) := \lceil \frac{1}{2} m_z^i \rceil, \quad n_z(\mathbf{m}) = \sum_{i=1}^N n_z^i(\mathbf{m}). \quad (2.9)$$

When it is not ambiguous, we will write  $n_z^i = n_z^i(\mathbf{m})$  and  $n_z = n_z(\mathbf{m})$ . As above, we are interested in a particular subset of configurations (in this case - a subset of  $\mathcal{M}_G^N$ ). We consider the set  $\mathcal{M}_G^N(1) \subset \mathcal{M}_G^N$  of configurations  $\mathbf{m}$  such that  $m_z^i \in 2\mathbb{N}$  for every  $i \in \{2, \dots, N\}$  and every  $z \in V$ . Configurations  $\mathbf{m} \in \mathcal{M}_G^N(1)$  can only have an odd number of incident  $i$ -links at a vertex if  $i = 1$  and hence when pairing these configurations with an element of  $P_G(\mathbf{m})$  the only walks present (if any) will be 1-walks.

As above, we may also consider the model with boundary conditions. For  $\mathcal{G} \supset G$  we define random path configurations on  $G \cup \partial E$  as above, allowing links on edges of  $\partial V$ . In this case we do not specify pairings on the end points of  $\partial E$  that lie in  $V^c$  so that  $\mathcal{P}_{G \cup \partial E}(\mathbf{m})$  is only the set of pairings of links incident to each  $z \in V$ . We denote by  $\mathcal{M}_{G \cup \partial E}^N(1)$  the set of all such configurations where  $m_z^i \in 2\mathbb{N}$  for every  $i \in \{2, \dots, N\}$  and every  $z \in V$ . In the definitions, to make the connection with  $+$  boundary conditions for the  $O(N)$ -spin model, we make the restriction that only 1-links may be present on  $\partial E$ .

We are now ready to define our measure on these configurations.

**Definition 2.5.** *Consider the set  $\mathcal{M}_G^N(1) \subset \mathcal{M}_G^N$  of link configurations such that  $m_z^i \in 2\mathbb{N}$  for every  $i \in \{2, \dots, N\}$  and every  $z \in V$ . Given  $N \in \mathbb{N}_{>0}$  and non-negative coupling constant  $\mathbf{J} = (J_e^1, \dots, J_e^N)_{e \in E}$*



define the measure  $\mu_{G,N,\mathbf{J},U}^{\text{free}}$  on  $\mathcal{M}_G^N(1)$  by

$$\mu_{G,N,\mathbf{J},U}^{\text{free}}(\mathbf{m}) = \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} U_z(\mathbf{m}) |P_G(\mathbf{m})|, \quad \mathbf{m} \in \mathcal{M}_G^N(1),$$

where  $U_z(\mathbf{m})$  is a vertex weight function depending only on the links incident to  $z$ . In the case where  $G \subset \mathcal{G}$ , we also define a measure on configurations on  $G \cup \partial E$

$$\mu_{G,N,\mathbf{J},U}^+(\mathbf{m}) := \prod_{e \in E \cup \partial E} \frac{\prod_{i=1}^N (J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} U_z(\mathbf{m}) |P_{G \cup \partial E}(\mathbf{m})| \prod_{e \in \partial E} \prod_{i=2}^N \mathbb{1}_{\{m_e^i=0\}}, \quad \mathbf{m} \in \mathcal{M}_{G \cup \partial E}^N(1), \quad (2.10)$$

the difference with  $\mu_{G,N,\mathbf{J},U}^{\text{free}}$  is that we allow 1-links on edges of  $\partial E$ . However we do not allow any links of colour  $2, \dots, N$  on  $\partial E$ .

In order to make the connection with the previous presentation of the random path model, and therefore with  $O(N)$ -spins, we introduce correlations for this measure in an analogous way to Definition 2.2. We will use the same notation to denote correlations as the correlations are indeed equal to those defined in 2.2, this is proven in Proposition 2.7.

**Definition 2.6.** For  $A \subset V$ , define  $\mathcal{S}(A)$  to be the set of configurations  $\mathbf{m} = (m^1, \dots, m^N)$  in  $\mathcal{M}_G^N(1)$  for the free or periodic boundary case or in  $\mathcal{M}_{G \cup \partial E}^N(1)$  for the + boundary case such that, for  $z \in V$ ,  $m_z^1 \in 2\mathbb{N} + 1$  if and only if  $z \in A$ . For  $\eta \in \{\text{free}, +\}$ , we define  $Z_{G,N,\mathbf{J},U}^\eta(A) = \mu_{G,N,\mathbf{J},U}^\eta(\mathcal{S}(A))$  and  $Z_{G,N,\mathbf{J},U}^{\text{loop},\eta} = \mu_{G,N,\mathbf{J},U}^\eta(\mathcal{S}(\emptyset))$ . Finally, the correlation functions are given by

$$\mathbb{G}_{G,N,\mathbf{J},U}^\eta(A) := \frac{Z_{G,N,\mathbf{J},U}^\eta(A)}{Z_{G,N,\mathbf{J},U}^{\text{loop},\eta}}.$$

The next proposition justifies this re-use of notation for correlations.

**Proposition 2.7.** Let  $N \in \mathbb{N}_{>0}$ ,  $\mathbf{J}$  such that  $J_e^i \geq 0$  for all  $e \in E$  and  $i \in [N]$ , and vertex weight functions  $U_x^{(N)}$  be fixed as in Proposition 2.3. For any  $A \subset V$  and  $\eta \in \{\text{free}, +\}$  we have that

$$\tilde{Z}_{G,N,\mathbf{J},U^{(N)}}^\eta(A) = Z_{G,N,\mathbf{J},U^{(N)}}^\eta(A).$$

*Proof.* The proof is a simple calculation. We present the proof for free or periodic boundary conditions, given this the adaptation to + boundary conditions is trivial, we simply insert indicators that no  $i$ -links are present on  $\partial E$  for  $i = 2, \dots, N$ .

$$\begin{aligned} \tilde{Z}_{G,N,\mathbf{J},U^{(N)}}^{\text{free}}(A) &= \sum_{(m,c,\pi) \in \tilde{\mathcal{S}}(A)} \prod_{e \in E} \frac{\prod_{i=1}^N (J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} U_z^{(N)}(m, c, \pi) \\ &= \sum_{\mathbf{m} \in \mathcal{M}_G} \sum_{c \in \mathcal{C}_G(\mathbf{m})} \sum_{\pi \in \mathcal{P}_G(\mathbf{m}, c)} \mathbb{1}_{(m,c,\pi) \in \tilde{\mathcal{S}}(A)} \prod_{e \in E} \frac{\prod_{i=1}^N (J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} U_z^{(N)}(m, c, \pi) \\ &= \sum_{(m^1, \dots, m^N) \in \mathcal{S}(A)} \sum_{\mathbf{m} \in \mathcal{M}_G} \sum_{c \in \mathcal{C}_G(\mathbf{m})} \left( \prod_{e \in E} \prod_{i=1}^N \mathbb{1}_{m_e^i(\mathbf{m}, c) = m_e^i} \right) \sum_{\pi \in \mathcal{P}_G(\mathbf{m})} \prod_{e \in E} \frac{\prod_{i=1}^N (J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} U_z^{(N)}(\mathbf{m}) \\ &= \sum_{(m^1, \dots, m^N) \in \mathcal{S}(A)} \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} U_z^{(N)}(\mathbf{m}) |P_G(\mathbf{m})| \end{aligned} \quad (2.11)$$

where the third inequality used that the definitions of  $n_z$  coincide on  $\tilde{\mathcal{S}}(A)$  and  $\mathcal{S}(A)$  and the last equality used that there are  $\binom{m_e}{m_e^1, \dots, m_e^N}$  ways to colour  $m_e$  objects so that there are  $m_e^i$  objects of colour  $i$ , for  $i = 1, \dots, N$ . The last expression is  $Z_{G,N,\mathbf{J},U^{(N)}}^{\text{free}}(A)$ , as desired.  $\square$

**Remark 2.8.** For the case  $\langle \prod_{x \in A} S_x^1 \prod_{y \in B} S_y^1 \rangle_{G,N,\mathbf{J}}^\eta$  with  $A \cap B \neq \emptyset$  we consider  $\mathbf{m}$  such that  $m_z^1 \in 2\mathbb{N} + 1$  for  $z \in A \Delta B$  and  $m_z^1 \in 2\mathbb{N}$  otherwise. We also consider pairings such that precisely one 1-link is unpaired at each  $z \in A \Delta B$  and two 1-links are unpaired at each  $z \in A \cap B$ .  $n_z(\mathbf{m})$  is increased by one for  $z \in A \cap B$  as the two links incident to  $z$  that correspond to 1-walk end points in some fixed pairing now contribute one each to the local time. No other links are unpaired. Defining  $Z_{G,N,\mathbf{J},U^{(N)}}^\eta(A; B)$  with these changes we can show  $\tilde{Z}_{G,N,\mathbf{J},U^{(N)}}^\eta(A; B) = Z_{G,N,\mathbf{J},U^{(N)}}^\eta(A; B)$  with an identical proof to above.

### 3 Proof of Theorem 1.1

During the proof we will consider,  $G, N, \mathbf{J}$  and  $U^{(N)}$  fixed and denote  $\mu_{G,N,\mathbf{J},U^{(N)}}^\eta$  by  $\mu_N$ . The proof of Theorem 1.1 follows easily from Lemma 3.1 as explained immediately below the lemma. Before proceeding to the statement and proof of the lemma, we introduce some simple concepts that are necessary. To avoid repeated duplication of expressions and definitions, throughout this section we will stick to the notation for free or periodic boundary conditions unless the required change for + boundary is not immediate. We will also consider the case  $A \cap B = \emptyset$  and comment on required changes where necessary.

For  $\mathbf{m}, \bar{\mathbf{m}} \in \mathcal{M}_G^N$  we define the sum of configurations  $\mathbf{m} + \bar{\mathbf{m}} = (m^1 + \bar{m}^1, \dots, m^N + \bar{m}^N)$  where  $m^i + \bar{m}^i = (m_e^i + \bar{m}_e^i)_{e \in E}$ . In other words,  $\mathbf{m} + \bar{\mathbf{m}}$  is the configuration with  $m_e^i + \bar{m}_e^i$   $i$ -links on  $e$ . We say that  $\bar{\mathbf{m}} \leq \mathbf{m}$  if  $\bar{m}_e^i \leq m_e^i$  for every  $e \in E$  and  $i \in [N]$ . When  $\bar{\mathbf{m}} \leq \mathbf{m}$  we can define their difference by  $\mathbf{m} - \bar{\mathbf{m}} = (m^1 - \bar{m}^1, \dots, m^N - \bar{m}^N)$  where  $m^i - \bar{m}^i = (m_e^i - \bar{m}_e^i)_{e \in E}$ . In the sequel we will make use of the fact that, for  $A, B \subset V$  with  $A \cap B = \emptyset$ , if we take  $\mathbf{m} \in \mathcal{S}(A)$  and  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  then  $n_z(\mathbf{m}) + n_z(\bar{\mathbf{m}}) = n_z(\mathbf{m} + \bar{\mathbf{m}})$ .

We may also interpret  $\mathbf{m} \in \mathcal{M}_G^N$  as a collection of multigraphs  $(m^1, \dots, m^N)$  each with vertex set  $V$  and the  $i^{\text{th}}$  multigraph having  $m_{\{x,y\}}^i$  edges between  $x, y \in V$  labelled  $1, \dots, m_{\{x,y\}}^i$ . This interpretation was used in [14]. Consider a collection of multigraphs  $\bar{\mathbf{m}} = (\bar{m}^1, \dots, \bar{m}^N)$  such that  $\bar{m}^i$  is a subgraph of  $m^i$  for each  $i \in [N]$  (possibly  $\bar{\mathbf{m}} = \mathbf{m}$ ). We can define pairings of the edges of these multigraphs in a completely analogous way to pairings for elements of  $\mathcal{M}_G^N$  and by a small abuse of notation denote by  $P_G(\bar{\mathbf{m}})$  the set of pairings of the multigraph  $\bar{\mathbf{m}}$  that leave at most one edge of each multigraph in the collection unpaired at each  $z \in V$ . For  $A \subset V$  we commit another slight abuse of notation and say that a collection of multigraphs  $\mathbf{m} \in \mathcal{S}(A)$  if  $m_z^i \in 2\mathbb{N} + 1$  for  $i = 1$  and  $z \in A$  and  $m_z^i \in 2\mathbb{N}$  otherwise.

For  $A \subset V$ , define  $\mathcal{F}_A \subset \mathcal{M}_G^N(1)$  to be the set of  $\mathbf{m} \in \mathcal{M}_G^N(1)$  such that each connected component of  $\mathbf{m}$  (when considered as a multigraph) contains an even number (possibly 0) of vertices from  $A$ . Further, for the case with boundary, define  $\mathcal{F}_A^+ \subset \mathcal{M}_{G \cup \partial E}^N(1)$  to be the set of  $\mathbf{m} \in \mathcal{M}_{G \cup \partial E}^N(1)$  such that each connected component of  $\mathbf{m}$  (when considered as a multigraph) either contains an even number (possibly 0) of vertices from  $A$  or is connected to  $\partial E$ . For  $A, B \subset V$  with  $A \cap B = \emptyset$  and  $\mathbf{m} \in \mathcal{S}(A \cup B)$  we define  $P_{A,B}(\mathbf{m}) \subset P_G(\mathbf{m})$  to be those pairings  $\pi$  of  $\mathbf{m}$  for which the collection of coloured paths defined by the pair  $(\mathbf{m}, \pi)$  is such no walk has an end point in  $A$  and an end point in  $B$ .

We have the following lemma.

**Lemma 3.1.** Let  $A, B \subset V$  with  $A \cap B = \emptyset$  and let vertex weight functions  $U_x^{(N)}$  be fixed as in Proposition 2.3 and  $\eta = \text{free}$ . For any  $N \in \mathbb{N}_{>1}$ , non-negative couple constants  $\mathbf{J} = (J_e^1, \dots, J_e^N)_{e \in E}$

and  $F : \mathcal{M}_G^N \rightarrow \mathbb{R}_{\geq 0}$  we have

$$\sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \leq \sum_{\substack{\mathbf{m} \in \mathcal{S}(A \cup B) \\ \bar{\mathbf{m}} \in \mathcal{S}(\emptyset)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \mathbb{1}_{\mathcal{F}_B}(\mathbf{m} + \bar{\mathbf{m}}) \frac{P_{A,B}(\mathbf{m})}{P_G(\mathbf{m})}.$$

The inequality holds for  $\eta = +$  if we replace  $\mathcal{F}_B$  on the right side by  $\mathcal{F}_B^+$ .

For  $A \cap B \neq \emptyset$  we replace  $\mathcal{S}(A \cup B)$  with  $\mathcal{S}(A \Delta B)$  and require that pairings for  $\mathbf{m}$  on the right side have two unpaired 1-links on each  $z \in A \cap B$ . Vertices in  $A \Delta B$  count twice in the definition of  $\mathcal{F}_B^{(+)}$ .

Theorem 1.1 follows from Lemma 3.1 and Proposition 2.3 when we set  $F \equiv 1$ , bound the indicator and ratio on the right side of the inequality by 1 and divide both sides by  $\mu_N(\mathcal{S}(\emptyset))^2$ .

**Remark 3.2.** *The proof of Lemma 3.1 is slightly different depending on whether  $N$  is even or odd. In both cases we use the proof of the switching lemma for  $N = 1$  as a template. By adding configurations in  $\mathcal{S}(A)$  and  $\mathcal{S}(B)$  we obtain an element  $\mathbf{m} \in \mathcal{S}(A \cup B)$ , when summing over these configuration we have a sum over  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  such that  $\bar{\mathbf{m}} \leq \mathbf{m}$  which we can give a geometric interpretation. The proof for the case  $A \cap B \neq \emptyset$  is essentially identical in both cases once we note the differences mentioned in the statement of the lemma and the requirement in the definition of  $\mathcal{S}(A \Delta B)$  that two 1-links are unpaired at each  $z \in A \cap B$ .*

Unlike the case of  $N = 1$  we are not able to give a bijection to the corresponding geometric object with  $\bar{\mathbf{m}} \in \mathcal{S}(\emptyset)$ , only an injection into the corresponding objects with the additional restriction that pairings are in  $P_{A,B}(\mathbf{m})$ . It would be extremely interesting, and likely very useful, to identify a bijection with some specific set of geometric objects, and therefore replace the inequality in Lemma 3.1 with some appropriate equality.

For the proof, we will make use of the following ordering on the pairs of a pairing at each  $z \in V$ .

Given a pairing of  $\mathbf{m} \in \mathcal{M}_G^N$  we may order the pairs at each  $z \in V$  as follows. For pairs  $p_1, p_2$  consisting of one or two edges each,  $p_1 < p_2$  if  $p_1$  contains edges of  $m^i$  and  $p_2$  contains edges of  $m^j$  for  $i < j$ . If  $p_1$  and  $p_2$  both contain edges of  $m^i$  then  $p_1 < p_2$  if the the lowest label of edges in  $p_1 \cup p_2$  belongs to an edge in  $p_1$  with arbitrary tie breaking decided by some ordering on  $E$  (or  $E \cup \partial E$ ) if the lowest label of each pairing is the same. Given a pairing of  $\mathbf{m}$  considered as a multigraph we may order the pairs at each  $z \in V$  in the same way.

We are now ready to prove Lemma 3.1, we begin with the proof for  $N$  even.

### 3.1 Proof of Lemma 3.1 for $N$ even

First we note that, for  $N$  even, we have that

$$\mathcal{U}^{(N)}(r) = \frac{\Gamma(\frac{N}{2})}{2^r (r + \frac{N-2}{2})!}$$

and hence for  $k, r \in \mathbb{N}$  with  $r \leq k$

$$\mathcal{U}^{(N)}(k-r) \mathcal{U}^{(N)}(r) = \frac{\Gamma(\frac{N}{2})^2}{2^k (k+N-2)!} \binom{k+N-2}{r + \frac{N-2}{2}}. \quad (3.1)$$

Let  $A, B \subset V$  such that  $A \cap B = \emptyset$ . A calculation similar to that in [7, Lemma 4.3] gives

$$\begin{aligned}
& \sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \\
&= \sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i + \bar{m}_e^i}}{m_e^i! \bar{m}_e^i!} \prod_{z \in V} U_z^{(N)}(\mathbf{m}) U_z^{(N)}(\bar{\mathbf{m}}) |P_G(\mathbf{m})| |P_G(\bar{\mathbf{m}})| \\
&= \sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i + \bar{m}_e^i}}{(m_e^i + \bar{m}_e^i)! (\bar{m}_e^i)^{-1}} \prod_{z \in V} U_z^{(N)}(\mathbf{m}) U_z^{(N)}(\bar{\mathbf{m}}) |P_G(\mathbf{m})| |P_G(\bar{\mathbf{m}})| \\
&= \sum_{\mathbf{m} \in \mathcal{S}(A \cup B)} F(\mathbf{m}) \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} \frac{\Gamma(\frac{N}{2})^2}{2^{n_z(\mathbf{m})} (n_z(\mathbf{m}) + N - 2)!} \\
&\quad \sum_{\substack{\bar{\mathbf{m}} \in \mathcal{S}(B) \\ \bar{\mathbf{m}} \leq \mathbf{m}}} \prod_{e \in E} \prod_{i=1}^N \binom{m_e^i}{\bar{m}_e^i} \prod_{z \in V} \binom{n_z(\mathbf{m}) + N - 2}{n_z(\bar{\mathbf{m}}) + \frac{N-2}{2}} |P_G(\mathbf{m} - \bar{\mathbf{m}})| |P_G(\bar{\mathbf{m}})|,
\end{aligned} \tag{3.2}$$

where the last equality used (3.1). We note that the set  $\{\bar{\mathbf{m}} \in \mathcal{S}(B) : \bar{\mathbf{m}} \leq \mathbf{m}\}$  is non-empty (and therefore the corresponding sum non-zero) if and only if  $\mathbf{m} \in \mathcal{F}_B$  when  $\eta = \text{free}$ , or  $\mathbf{m} \in \mathcal{F}_B^+$  when  $\eta = +$ . Indeed, for the case  $\eta = \text{free}$  there can only be an odd number of  $i$ -links incident to every vertex of  $B$  for  $\bar{\mathbf{m}} \leq \mathbf{m}$  if vertices of  $B$  can be paired off in some way within each connected component of  $\mathbf{m}$ . This corresponds to having a walk between these pairs. For  $\eta = +$  the vertices of  $B$  do not have to be paired off with each other as long as they can be connected to the boundary.

Consider a subgraph  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  of the multigraph defined by  $\mathbf{m} \in \mathcal{S}(A \cup B)$  and pairings  $\bar{\pi}$  of  $\bar{\mathbf{m}}$  and  $\pi$  of  $\mathbf{m} \setminus \bar{\mathbf{m}}$ . Now replace each vertex  $z \in V$  with the  $n_z(\mathbf{m}) + N - 2$  vertices  $\{z_1, \dots, z_{n_z(\mathbf{m}) + N - 2}\}$ . At each  $z \in V$ , select a subset,  $V_z(\bar{\mathbf{m}})$ , of these vertices of size  $n_z(\bar{\mathbf{m}}) + \frac{N-2}{2}$ . The sum over  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  above is then the number of choices of such a subgraph  $\bar{\mathbf{m}}$ , a pairing of  $\bar{\mathbf{m}}$  and  $\mathbf{m} \setminus \bar{\mathbf{m}}$  and a set,  $V_z(\bar{\mathbf{m}})$ , of size  $n_z(\bar{\mathbf{m}}) + \frac{N-2}{2}$  at each  $z \in V$ . Encode these choices in a tuple  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$ .

We will define an injection from the set of tuples  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$  with  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  into the set of tuples  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$  with  $\bar{\mathbf{m}} \in \mathcal{S}(\emptyset)$  by using the ordering on pairs defined above.

**Remark 3.3.** *Informally speaking, we want to identify the walks defined by  $\bar{\mathbf{m}}$  and  $\bar{\pi}$  (these walks have end points on  $B$  and in the case  $\eta = +$  may also leave  $G$ ) and some corresponding subset of  $V_z(\bar{\mathbf{m}})$  whose size is the number of visits of these walks to  $z$ . The mapping will then move these walks and vertices from  $\bar{\mathbf{m}}, \bar{\pi}$  and  $(V_z(\bar{\mathbf{m}}))_{z \in V}$  to  $\mathbf{m} \setminus \bar{\mathbf{m}}, \pi$ , and  $(V_z(\bar{\mathbf{m}})^c)_{z \in V}$ . To do this, we need a deterministic choice to decide which vertices in  $V_z(\bar{\mathbf{m}})$  “belong” to the walks. This deterministic choice is provided by the ordering on pairs defined above Lemma 3.1.*

For a tuple  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$  with  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  we consider the paths defined by  $\bar{\mathbf{m}}$  and  $\bar{\pi}$  and attach the  $j^{\text{th}}$  pair of  $\bar{\pi}_z$  to the  $j^{\text{th}}$  vertex in  $V_z(\bar{\mathbf{m}})$ . Similarly, we attach the  $j^{\text{th}}$  pair of  $\pi_z$  to the  $j^{\text{th}}$  vertex in  $V_z(\bar{\mathbf{m}})^c$ . This results in a graph with  $m_e$  edges for each  $e \in E$  and  $n_z(\mathbf{m}) + N - 2$  vertices for each  $z \in V$  such that each connected component of the graph is either an isolated vertex ( $N - 2$  isolated vertices for every  $z \in V$ ) or a path defined by  $\bar{\mathbf{m}}$  and  $\bar{\pi}$  or by  $\mathbf{m} \setminus \bar{\mathbf{m}}$  and  $\pi$  with the vertices of these connected components determined by  $V_z(\bar{\mathbf{m}})$  and the ordering on pairs of  $\bar{\pi}, \pi$  at each  $z \in V$ .

Consider the connected components of this graph corresponding to the walks with end points on  $B$ . Let  $\bar{\mathbf{m}}^B \subset \bar{\mathbf{m}}$  be the corresponding set of edges, let  $\bar{\pi}^B$  respectively  $\bar{\pi}^{\text{loops}}$  be the pairing of  $\bar{\mathbf{m}}^B$  respectively  $\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B$  coming from the restriction of the partition defined by  $\bar{\pi}$  to  $\bar{\mathbf{m}}^B$  respectively  $\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B$ , and let  $V_z(\bar{\mathbf{m}}^B) \subset V_z(\bar{\mathbf{m}})$  be the set of vertices attached to pairs of  $\bar{\pi}^B$ . Note that identifying

$\bar{\mathbf{m}}^B$  requires knowledge of  $\bar{\pi}$  and that the pair  $(\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B, \bar{\pi}^{loops})$  can be interpreted as a collection of coloured loops. We define our injection by mapping

$$(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V}) \mapsto (\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B, \bar{\pi}^{loop}, \pi \sqcup \bar{\pi}^B, (V_z(\bar{\mathbf{m}}) \setminus V_z(\bar{\mathbf{m}}^B))_{z \in V}) \quad (3.3)$$

where  $\pi \sqcup \bar{\pi}^B$  is the pairing of  $(\mathbf{m} \setminus \bar{\mathbf{m}}) \cup \bar{\mathbf{m}}^B$  that pairs the edges of  $\mathbf{m} \setminus \bar{\mathbf{m}}$  according to  $\pi$  and the edges of  $\bar{\mathbf{m}}^B$  according to  $\bar{\pi}^B$ . The image of this map consists of a tuple of an element  $\tilde{\mathbf{m}} = \bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B \in \mathcal{S}(\emptyset)$ , a pairing of  $\tilde{\mathbf{m}}$ , a pairing of  $\mathbf{m} \setminus \tilde{\mathbf{m}}$  and a collection of sets  $(V_z(\tilde{\mathbf{m}}))_{z \in V}$  such that  $|V_z(\tilde{\mathbf{m}})| = n_z(\tilde{\mathbf{m}}) + \frac{N-2}{2}$ . This can again be interpreted as a graph whose connected components are isolated vertices and paths defined by the triple  $\tilde{\mathbf{m}}, \bar{\pi}^{loop}$  and  $(V_z(\tilde{\mathbf{m}}))_{z \in V}$  or the triple  $\mathbf{m} \setminus \tilde{\mathbf{m}}, \pi \sqcup \bar{\pi}^B, (V_z(\tilde{\mathbf{m}})^c)_{z \in V}$  where pairs are attached to vertices according to the fixed ordering inherited from the map. Described somewhat informally, the mapping takes the graph obtained from  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$  and moves the walks with end points in  $B$  from the first triple to the second.

The map is an injection. Indeed, the tuple  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$  together with the ordering on pairings defines uniquely the graph described below Remark 3.3 whose connected components are isolate vertices and paths labelled by either the triple  $\bar{\mathbf{m}}, \bar{\pi}, (V_z(\bar{\mathbf{m}}))_{z \in V}$  or the triple  $\mathbf{m} \setminus \bar{\mathbf{m}}, \pi, (V_z(\bar{\mathbf{m}})^c)_{z \in V}$ . The path structure is unchanged by the mapping, it simply changes the label of the walks with end points on  $B$  from the first triple to the second. Hence we can change the walks with end points in  $B$  back to the first triple and recover the original ctuple. It is not clear which tuples with  $\bar{\mathbf{m}} \in \mathcal{S}(\emptyset)$  can be given such an inherited ordering in a consistent way, hence we do not know if there is a natural bijection into an appropriate set of these tuples.

We note that every tuple in the image of this map involved pairings such that no walk has an end point on  $A$  and an end point on  $B$ . In other words these pairings are elements of  $P_G(\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B)$  and  $P_{A,B}((\mathbf{m} \setminus \bar{\mathbf{m}}) \cup \bar{\mathbf{m}}^B)$ , respectively. Putting this together we have that

$$\begin{aligned} & \sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \\ & \leq \sum_{\mathbf{m} \in \mathcal{S}(A \cup B)} F(\mathbf{m}) \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} \frac{\Gamma(\frac{N}{2})^2}{2^{n_z(\mathbf{m})} (n_z(\mathbf{m}) + N - 2)!} \mathbb{1}_{\mathcal{F}_B}(\mathbf{m}) \\ & \quad \sum_{\substack{\bar{\mathbf{m}} \in \mathcal{S}(\emptyset) \\ \bar{\mathbf{m}} \leq \mathbf{m}}} \prod_{e \in E} \prod_{i=1}^N \binom{m_e^i}{\bar{m}_e^i} \prod_{z \in V} \binom{n_z(\mathbf{m}) + N - 2}{n_z(\bar{\mathbf{m}}) + \frac{N-2}{2}} |P_{A,B}(\mathbf{m} - \bar{\mathbf{m}})| |P_G(\bar{\mathbf{m}})|. \end{aligned} \quad (3.4)$$

Now by tracing the first calculation backwards we obtain Lemma 3.1 for  $N$  even.

### 3.2 Proof of Lemma 3.1 for $N \geq 3$ odd

The proof of  $N$  odd differs slightly from that of  $N$  even due to the presence of a half-integer in the gamma functions in the definition of  $U_z^{(N)}$ . Other than this difference, and hence the need for a slightly different geometric interpretation of the sum over  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  to the one for  $N$  even, the proofs are very similar. We include the details here for completeness.

For  $N \geq 3$  odd, a simple calculation shows that

$$\mathcal{U}^{(N)}(r) = \frac{2^{\frac{N-2}{2}} \Gamma(\frac{N}{2})}{(2r + N - 2)!!}$$

where  $(2k - 1)!! = (2k - 1)(2k - 3) \cdots 3 = (2k)! / 2^k k!$  is the double factorial. Hence, for  $k, r \in \mathbb{N}$  with  $r \leq k$  another simple calculation gives

$$\mathcal{U}^{(N)}(k - r) \mathcal{U}^{(N)}(r) = \frac{2^{N-1} \Gamma(\frac{N}{2})^2}{(2k + 2N - 3)!! (k + N - 1)!} \binom{2k + 2N - 2}{2r + N - 1} \left(r + \frac{N-1}{2}\right)! \left(k - r + \frac{N-1}{2}\right)! \quad (3.5)$$

Using this identity the same calculation as in (3.2) gives

$$\begin{aligned}
& \sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \\
&= \sum_{\mathbf{m} \in \mathcal{S}(A \cup B)} F(\mathbf{m}) \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} \frac{2^{N-1} \Gamma(\frac{N}{2})^2}{(2n_z(\mathbf{m}) + 2N - 3)!! (n_z(\mathbf{m}) + N - 1)!} \\
& \sum_{\substack{\bar{\mathbf{m}} \in \mathcal{S}(B) \\ \bar{\mathbf{m}} \leq \mathbf{m}}} \prod_{e \in E} \prod_{i=1}^N \binom{m_e^i}{\bar{m}_e^i} \prod_{z \in V} \binom{2n_z(\mathbf{m}) + 2N - 2}{2n_z(\bar{\mathbf{m}}) + N - 1} (n_z(\bar{\mathbf{m}}) + \frac{N-1}{2})! (n_z(\mathbf{m} - \bar{\mathbf{m}}) + \frac{N-1}{2})! |P_G(\mathbf{m} - \bar{\mathbf{m}})| |P_G(\bar{\mathbf{m}})|.
\end{aligned} \tag{3.6}$$

The set  $\{\bar{\mathbf{m}} \in \mathcal{S}(B) : \bar{\mathbf{m}} \leq \mathbf{m}\}$  is non-empty (and therefore the corresponding sum non-zero) if and only if  $\mathbf{m} \in \mathcal{F}_B$  when  $\eta = \text{free}$ , or  $\mathbf{m} \in \mathcal{F}_B^+$  when  $\eta = +$ . Indeed, for the case  $\eta = \text{free}$  there can only be an odd number of  $i$ -links incident to every vertex of  $B$  for  $\bar{\mathbf{m}} \leq \mathbf{m}$  if vertices of  $B$  can be paired off in some way within each connected component of  $\mathbf{m}$ . This corresponds to having a walk between these pairs. For  $\eta = +$  the vertices of  $B$  do not have to be paired off with each other as long as they can be connected to the boundary. Consider a subgraph  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  of the multigraph defined by  $\mathbf{m} \in \mathcal{S}(A \cup B)$  and pairings  $\bar{\pi}$  of  $\bar{\mathbf{m}}$  and  $\pi$  of  $\mathbf{m} \setminus \bar{\mathbf{m}}$ . Now replace each vertex  $z \in V$  with the  $2n_z(\mathbf{m}) + 2N - 2$  vertices  $\{z_1, \dots, z_{2n_z(\mathbf{m}) + 2N - 2}\}$ . At each  $z \in V$ , select a subset  $V_z(\bar{\mathbf{m}})$  of these vertices of size  $2n_z(\bar{\mathbf{m}}) + N - 1$  and then select an ordering of the first  $n_z(\bar{\mathbf{m}}) + \frac{N-1}{2}$  in this set and the first  $n_z(\mathbf{m}) - n_z(\bar{\mathbf{m}}) + \frac{N-2}{2}$  vertices in the complementary set  $V_z(\bar{\mathbf{m}})^c$ . The sum over  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  above is then the number of choices of such a subgraph  $\bar{\mathbf{m}}$ , a pairing of  $\bar{\mathbf{m}}$  and  $\mathbf{m} \setminus \bar{\mathbf{m}}$ , a set of size  $2n_z(\bar{\mathbf{m}}) + N - 1$  at each  $z \in B$  which we will denote by  $V_z(\bar{\mathbf{m}})$ , an ordering  $\bar{p}_z$  of the first half of  $V_z(\bar{\mathbf{m}})$  and an ordering  $p_z$  of the first half of  $V_z(\bar{\mathbf{m}})^c$ . Encode these choices in a tuple  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V}, (\bar{p}_z)_{z \in V}, (p_z)_{z \in V})$ .

Note that we have interpreted the term  $(n_z(\bar{\mathbf{m}}) + \frac{N-1}{2})!$  as the number of ways to order the first half of  $V_z(\bar{\mathbf{m}})$ , but we could also replace this with an ordering of any other deterministically chosen subset of  $V_z(\bar{\mathbf{m}})$  that has size  $n_z(\bar{\mathbf{m}}) + \frac{N-1}{2}$  and have a valid interpretation of the term  $(n_z(\bar{\mathbf{m}}) + \frac{N-1}{2})!$ . This will be helpful below.

We will define an injection from the set of tuples  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V}, (\bar{p}_z)_{z \in V}, (p_z)_{z \in V})$  with  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  into the set of tuples  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V}, (\bar{p}_z)_{z \in V}, (p_z)_{z \in V})$  with  $\bar{\mathbf{m}} \in \mathcal{S}(\emptyset)$  by extending the orderings  $\bar{p}_z$  and  $p_z$  to all of  $V_z(\bar{\mathbf{m}})$  and  $V_z(\bar{\mathbf{m}})^c$ , respectively, and then using the ordering on pairs defined above. First we describe how we extend the orderings. Given an ordering of the first half of a set  $V_z(\bar{\mathbf{m}})$  of size  $2n_z(\bar{\mathbf{m}}) + N - 1$  we extend the ordering to all vertices by repetition as follows, if the  $k^{\text{th}}$  vertex in  $V_z(\bar{\mathbf{m}})$ ,  $k \leq n_z(\bar{\mathbf{m}}) + \frac{N-1}{2}$ , is in position  $\ell$  of the ordering then the  $(k + n_z(\bar{\mathbf{m}}) + \frac{N-1}{2})^{\text{th}}$  vertex of  $V_z(\bar{\mathbf{m}})$  is placed in position  $\ell + n_z(\bar{\mathbf{m}}) + \frac{N-1}{2}$ . We extend the ordering of the first  $n_z(\mathbf{m}) - n_z(\bar{\mathbf{m}}) + \frac{N-2}{2}$  vertices of  $V_z(\bar{\mathbf{m}})^c$  to all of  $V_z(\bar{\mathbf{m}})^c$  in the same way. Now, for a tuple  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V}, (\bar{p}_z)_{z \in V}, (p_z)_{z \in V})$  with  $\bar{\mathbf{m}} \in \mathcal{S}(B)$  we consider the paths defined by  $\bar{\mathbf{m}}$  and  $\bar{\pi}$  and attach the  $j^{\text{th}}$  pair of  $\bar{\pi}_z$  to the  $j^{\text{th}}$  vertex in the ordering of  $V_z(\bar{\mathbf{m}})$ . Similarly, we attach the  $j^{\text{th}}$  pair of  $\pi_z$  to the  $j^{\text{th}}$  vertex in the ordering of  $V_z(\bar{\mathbf{m}})^c$ . This results in a graph with  $m_e$  edges for each  $e \in E$  and  $2n_z(\mathbf{m}) + 2N - 2$  vertices for each  $z \in V$  such that each connected component of the graph is either an isolated vertex ( $n_z(\mathbf{m}) + 2N - 2$  isolated vertices for every  $z \in V$ ) or a path defined by  $\bar{\mathbf{m}}, \bar{\pi}$  and  $(\bar{p}_z)_{z \in V}$  or by  $\mathbf{m} \setminus \bar{\mathbf{m}}, \pi$  and  $(p_z)_{z \in V}$ .

Informally speaking, the mapping we define below works almost identically to the mapping described in Remark 3.3. The difference with the case of  $N$  even is that for  $N$  even we always took the ‘‘natural’’ ordering on  $V_z(\bar{\mathbf{m}})$  and  $V_z(\bar{\mathbf{m}})^c$  coming from the index of the vertices but in this case the ordering comes from  $\bar{p}$  and  $p$ . There are also extra vertices that will be moved by referring to the extended ordering of vertices.

Consider the connected components of this graph corresponding to the walks with end points on  $B$ .

Let  $\bar{\mathbf{m}}^B \subset \bar{\mathbf{m}}$  be the corresponding set of edges, let  $\bar{\pi}^B$  respectively  $\bar{\pi}^{loop}$  be the pairing of  $\bar{\mathbf{m}}^B$  respectively  $\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B$  coming from the restriction of the partition defined by  $\bar{\pi}$  to  $\bar{\mathbf{m}}^B$  respectively  $\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B$ , and let  $V_z(\bar{\mathbf{m}}^B) \subset V_z(\bar{\mathbf{m}})$  be the set of vertices that are attached to pairs of  $\bar{\pi}^B$  together with the set of isolated vertices that are  $n_z(\bar{\mathbf{m}}) + \frac{N-1}{2}$  places higher in the ordering than a vertex that is attached to a pair of  $\bar{\pi}^B$ . Finally we consider the orderings  $\bar{p}_z$  and  $p_z$ ,  $z \in V$ . Let  $\bar{p}_z^{loop}$  be the ordering on  $V_z(\bar{\mathbf{m}}) \setminus V_z(\bar{\mathbf{m}}^B)$  induced by  $\bar{p}_z$  (i.e.  $\bar{p}_z^{loop}$  maintains the same relative order of the the remaining vertices in  $V_z(\bar{\mathbf{m}}) \setminus V_z(\bar{\mathbf{m}}^B)$ ). Let  $p_z \sqcup \bar{p}_z^B$  be the ordering on  $V_z(\bar{\mathbf{m}})^c \cup V_z(\bar{\mathbf{m}})^B$  induced by the ordering of  $V_z(\bar{\mathbf{m}})^c$  coming from  $p_z$  and the ordering of  $V_z(\bar{\mathbf{m}})^B$  coming from  $\bar{p}$ . In other words, reorder  $V_z(\bar{\mathbf{m}})^c$  according to  $p_z$  and  $V_z(\bar{\mathbf{m}})$  according to  $\bar{p}_z$ , now take the union  $V_z(\bar{\mathbf{m}})^c \cup V_z(\bar{\mathbf{m}})^B$  and define the ordering obtained as  $p_z \sqcup \bar{p}_z^B$ .

We note than  $|V_z(\bar{\mathbf{m}}^B)| = 2n_z(\bar{\mathbf{m}}^B)$  (half the vertices are attach to walks with end points on  $B$  and half are associated isolated vertices that are  $n_z(\bar{\mathbf{m}}) + \frac{N-1}{2}$  places higher in the repeated ordering). Also note that identifying  $\bar{\mathbf{m}}^B$  requires knowledge of  $\bar{\pi}$  and that the pair  $(\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B, \bar{\pi}^{loop})$  can be interpreted as a collection of coloured loops. We define our injection by the mapping

$$(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V}, (\bar{p}_z)_{z \in V}, (p_z)_{z \in V}) \mapsto (\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B, \bar{\pi}^{loop}, \pi \sqcup \bar{\pi}^B, (V_z(\bar{\mathbf{m}}) \setminus V_z(\bar{\mathbf{m}}^B))_{z \in V}, (\bar{p}_z^{loop})_{z \in V}, (p_z \sqcup \bar{p}_z^B)_{z \in V}). \quad (3.7)$$

The image of this map consists of a tuple of an element  $\tilde{\mathbf{m}} = \bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B \in \mathcal{S}(\emptyset)$ , a pairing of  $\tilde{\mathbf{m}}$ , a pairing of  $\mathbf{m} \setminus \tilde{\mathbf{m}}$ , a collection of sets  $(V_z(\tilde{\mathbf{m}}))_{z \in V}$  such that  $|V_z(\tilde{\mathbf{m}})| = 2n_z(\tilde{\mathbf{m}}) + N - 1$ , an ordering of the first  $n_z(\tilde{\mathbf{m}}) + \frac{N-1}{2}$  vertices of  $V_z(\tilde{\mathbf{m}})$  for each  $z \in V$  and an ordering of a set of  $n_z(\mathbf{m} \setminus \tilde{\mathbf{m}}) + \frac{N-1}{2}$  vertices of  $V_z(\tilde{\mathbf{m}})^c$  for each  $z \in V$  such that this set is determined by the previous elements of the tuple. This can again be interpreted as a graph whose connected components are isolated vertices and paths defined by the triple  $\tilde{\mathbf{m}}, \bar{\pi}^{loop}$  and  $(V_z(\tilde{\mathbf{m}}))_{z \in V}$  or the triple  $\mathbf{m} \setminus \tilde{\mathbf{m}}, \pi \sqcup \bar{\pi}^B, (V_z(\tilde{\mathbf{m}})^c)_{z \in V}$  where pairs are attached to vertices according to a fixed ordering. Described somewhat informally, the mapping takes the graph obtained from  $(\bar{\mathbf{m}}, \bar{\pi}, \pi, (V_z(\bar{\mathbf{m}}))_{z \in V})$  and moves the walks with end points in  $B$  from the first triple to the second. This determines the set of  $n_z(\mathbf{m} \setminus \tilde{\mathbf{m}}) + \frac{N-1}{2}$  vertices of  $V_z(\tilde{\mathbf{m}})^c$  that are given a specific ordering.

As for the case of  $N$  even, this map is an injection. Indeed, the path structure is unchanged and the ordering on pairs is determined by the graph interpretation so that we can move the walks with end points in  $B$  and their corresponding isolated vertices back to the first triple and recover the original choice of vertices for each  $z \in V$  and their ordering.

We note that every tuple in the image of this map involved pairings such that no walk has an end point on  $A$  and an end point on  $B$ . In other words these pairings are elements of  $P_G(\bar{\mathbf{m}} \setminus \bar{\mathbf{m}}^B)$  and  $P_{A,B}((\mathbf{m} \setminus \tilde{\mathbf{m}}) \cup \bar{\mathbf{m}}^B)$ , respectively. Putting this together we have that

$$\begin{aligned} & \sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \\ & \leq \sum_{\mathbf{m} \in \mathcal{S}(A \cup B)} F(\mathbf{m}) \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} \frac{2^{N-1} \Gamma(\frac{N}{2})^2}{(2n_z(\mathbf{m}) + 2N - 3)!! (n_z(\mathbf{m}) + N - 1)!} \mathbb{1}_{\mathcal{F}_B}(\mathbf{m}) \\ & \sum_{\substack{\bar{\mathbf{m}} \in \mathcal{S}(\emptyset) \\ \bar{\mathbf{m}} \leq \mathbf{m}}} \prod_{e \in E} \prod_{i=1}^N \binom{m_e^i}{\bar{m}_e^i} \prod_{z \in V} \binom{2n_z(\mathbf{m}) + 2N - 2}{2n_z(\bar{\mathbf{m}}) + N - 1} (n_z(\bar{\mathbf{m}}) + \frac{N-1}{2})! (n_z(\mathbf{m} - \bar{\mathbf{m}}) + \frac{N-1}{2})! |P_{A,B}(\mathbf{m} - \bar{\mathbf{m}})| |P_G(\bar{\mathbf{m}})|. \end{aligned} \quad (3.8)$$

Now by tracing the first calculation backwards we obtain Lemma 3.1 for  $N$  odd.

## 4 Extension to the case of external magnetic field

In this section we introduce the  $O(N)$ -spin model with external field and the corresponding RPM. The setting is as in Section 1.1 except in addition we introduce an *inhomogeneous external field*  $\mathbf{h} = (h_x^1, \dots, h_x^N)_{x \in V}$  with  $h_x^i \geq 0$  for each  $x \in V, i \in [N]$ . For  $\eta \in \{\text{free}, +\}$  and  $\varphi \in \Omega_{G,N}$  we define

$$H_{G,N,\mathbf{J},\mathbf{h}}^\eta(\varphi) = H_{G,N,\mathbf{J}}^\eta(\varphi) - \sum_{x \in V} \sum_{i=1}^N h_x^i \varphi_x^i. \quad (4.1)$$

This corresponds to having an external magnetic field on  $V$  whose magnitude and direction on  $x \in V$  are given by  $|(h_x^1, \dots, h_x^N)|$  and  $(h_x^1, \dots, h_x^N)/|(h_x^1, \dots, h_x^N)|$ , respectively. Note that the magnetic field term looks similar to the boundary term for  $\eta = +$ . Indeed, we can think of this term as a boundary term coming from edges connected to *ghost vertices*. We use this interpretation when introducing the modification of the random path model. This means that the extension to this case can naturally be arrived at from the case with boundary. Nevertheless we make the effort to present the necessary changes. The changes required to prove Theorems 1.1 and 4.1 for a general boundary condition (other than  $+$ ) are virtually identical to the changes required for an external magnetic field.

Expectations are defined analogously to Section 1.1. For  $f : \Omega_{G,N} \rightarrow \mathbb{R}$

$$\langle f \rangle_{G,N,\mathbf{J},\mathbf{h}}^\eta = \frac{1}{Z_{G,N,\mathbf{J},\mathbf{h}}^\eta} \int_{\Omega_{G,N}} d\varphi f(\varphi) e^{-H_{G,N,\mathbf{J},\mathbf{h}}^\eta(\varphi)} \quad (4.2)$$

where  $Z_{G,N,\mathbf{J},\mathbf{h}}^\eta$  is a normalising constant. We also have the analogous theorem and corollary to the case of no external field. For  $x \in V$ , define  $S_x : \Omega_{G,N} \mapsto \mathbb{S}^{N-1}$  by  $S_x(\varphi) = \varphi_x$  and the components of  $S_x = (S_x^1, \dots, S_x^N)$ .

**Theorem 4.1.** *Let  $G = (V, E)$  be a finite simple graph and  $N \in \mathbb{N}_{>1}$ . Assume that  $J_e^i \geq 0$  for every  $e \in E, i \in [N]$  and  $h_x^i \geq 0$  for every  $x \in V, i \in [N]$ . For  $A, B \subset V$  and  $\eta \in \{\text{free}, +\}$  we have*

$$\left\langle \prod_{x \in A} S_x^1 \prod_{y \in A} S_y^1 \right\rangle_{G,N,\mathbf{J},\mathbf{h}}^\eta \geq \left\langle \prod_{x \in A} S_x^1 \right\rangle_{G,N,\mathbf{J},\mathbf{h}}^\eta \left\langle \prod_{y \in B} S_y^1 \right\rangle_{G,N,\mathbf{J},\mathbf{h}}^\eta.$$

**Corollary 4.2.** *Let  $G = (V, E)$  be a finite simple graph and  $N \in \mathbb{N}_{>1}$ . Assume that  $J_e^i \geq 0$  for every  $e \in E, i \in [N]$  and  $h_x^i \geq 0$  for every  $x \in V, i \in [N]$ . Let  $\eta \in \{\text{free}, +\}$ . For  $A \subset V$  and  $e \in E$  we have*

$$\frac{\partial}{\partial J_e^i} \left\langle \prod_{x \in A} S_x^1 \right\rangle_{G,N,\mathbf{J},\mathbf{h}}^\eta \geq 0.$$

For the proof we will stick to the case  $A \cap B = \emptyset$ , the modifications for  $A \cap B \neq \emptyset$  are completely analogous to the modifications for the case of no external field.

In order to prove this result, we use a slight modification of the random path model. As in [13] we do this by introducing *ghost vertices*. We introduce  $N$  ghost vertices, one for each colour. The case where  $h_x^i = 0$  for  $i \in \{1, \dots, N-1\}$  for every  $x \in V$  corresponds to having one ghost vertex as considered in [13], the extension to multiple ghost vertices is straightforward. Let  $g_1, \dots, g_N$  be vertices not in  $G$  (or in the case with boundary, not in  $\mathcal{G}$ ). Further define edge set  $\tilde{E}_g = \{\{x, g_1\}, \dots, \{x, g_N\} : x \in V\}$ . We now define a modified graph  $G_g = (V_g, E_g)$  where  $V_g = V \cup \{g_1, \dots, g_N\}$  and  $E_g = E \cup \tilde{E}_g$ .  $G_g$  consists of the original graph  $G$  together with  $N$  ghost vertices and an edge from  $x$  to each ghost vertex.

Now consider the random path model on  $G_g$  ( $G_g \cup \partial E$  for the case with boundary).  $\mathcal{M}_{G_g}^N(1) \subset \mathcal{M}_{G_g}^N = \mathbb{N}^{E_g \times [N]}$  ( $\mathcal{M}_{G_g \cup \partial E}^N(1)$ ) consists of tuples  $\mathbf{m} = (m^1, \dots, m^N)$  of  $N$  collections of links on  $E_g$  ( $E_g \cup \partial E$ )



such that  $m_z^i \in 2\mathbb{N}$  for every  $i \in \{2, \dots, N\}$  and  $z \in V$  and  $m_{\{z, g_j\}}^i = 0$  whenever  $i \neq j$  for every  $z \in V$  (i.e. no  $i$ -links incident to the  $j^{\text{th}}$  ghost vertex for  $i \neq j$ ). Note that we place no restriction on the parity of  $m_{g_i}^i$ . For  $\mathbf{m} \in \mathcal{M}_{G_g}^N(1)$  ( $\mathbf{m} \in \mathcal{M}_{G_g \cup \partial E}^N(1)$ ) let  $\mathcal{P}_{G_g}$  ( $\mathcal{P}_{G_g \cup \partial E}$ ) be the set of pairings of links incident to each  $z \in V$ . Links incident to  $g_j$  for  $j \in [N]$  (and  $z \in \partial^{\text{ext}}V$ ) are not paired. Let  $P_{G_g}(\mathbf{m})$  ( $P_{G_g \cup \partial E}(\mathbf{m})$ ) be the set of pairings such that at most one link of each colour is unpaired at each  $z \in V$ . Recall that when considering  $\langle \prod_{x \in A} S_x^1 \prod_{y \in A} S_y^1 \rangle_{G, N, \mathbf{J}}^\eta$  this set of pairings is relevant when  $A \cap B = \emptyset$ , when  $A \cap B \neq \emptyset$  we consider pairings where one 1-link is unpaired at each  $z \in A \Delta B$  and two 1-links are unpaired at each  $z \in A \cap B$ .

We see that a tuple  $(\mathbf{m}, \pi)$  with  $\mathbf{m} \in \mathcal{M}_{G_g}^N(1)$  ( $\mathbf{m} \in \mathcal{M}_{G_g \cup \partial E}^N(1)$ ) and  $\pi \in P_{G_g}(\mathbf{m})$  ( $\mathcal{P}_{G_g \cup \partial E}$ ) can be interpreted as a collection of loops with colours in  $[N]$ , 1-paths with end points on  $V$ , and  $i$ -paths with both end points on  $g_i$  for  $i \in [N]$  (in addition these tuples may also have paths leaving  $G_g$  for the case  $G_g \cup \partial E$ ).

We now define our measure on these new configurations. For  $\eta = \text{free}$  the measure of  $\mathbf{m} \in \mathcal{M}_{G_g}^N(1)$  is

$$\mu_{G_g, N, \mathbf{J}, \mathbf{h}, U}^{\text{free}}(\mathbf{m}) = \prod_{e \in E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} \prod_{i=1}^N \frac{(h_z^i)^{m_{\{z, g_i\}}}}{m_{\{z, g_i\}}!} \prod_{z \in V} U_z(\mathbf{m}) |P_{G_g}(\mathbf{m})|. \quad (4.3)$$

For  $\eta = +$  the measure of  $\mathbf{m} \in \mathcal{M}_{G_g \cup \partial E}^N(1)$  is

$$\mu_{G_g, N, \mathbf{J}, \mathbf{h}, U}^+(\mathbf{m}) = \prod_{e \in E \cup \partial E} \prod_{i=1}^N \frac{(J_e^i)^{m_e^i}}{m_e^i!} \prod_{z \in V} \prod_{i=1}^N \frac{(h_z^i)^{m_{\{z, g_i\}}}}{m_{\{z, g_i\}}!} \prod_{z \in V} U_z(\mathbf{m}) |P_{G_g}(\mathbf{m})| \prod_{e \in \partial E} \prod_{i=1}^N \mathbb{1}_{\{m_e^i=0\}}. \quad (4.4)$$

For  $A \subset V$  we define  $\mathcal{S}(A)$  to be the set of  $\mathbf{m} \in \mathcal{M}_{G_g}^N(1)$  ( $\mathbf{m} \in \mathcal{M}_{G_g \cup \partial E}^N(1)$ ) such that, for  $z \in V$ ,  $m_z^1 \in 2\mathbb{N} + 1$  if and only if  $z \in A$ . For  $\eta \in \{\text{free}, +\}$  define  $Z_{G, N, \mathbf{J}, \mathbf{h}, U}^\eta(A) = \mu_{G_g, N, \mathbf{J}, \mathbf{h}, U}^\eta(\mathcal{S}(A))$  and  $Z_{G, N, \mathbf{J}, \mathbf{h}, U}^{\text{loop}, \eta} = \mu_{G_g, N, \mathbf{J}, \mathbf{h}, U}^\eta(\mathcal{S}(\emptyset))$ . Of course, in the case with ghost vertices configurations in  $\mathcal{S}(\emptyset)$  contain not only coloured loops, but also coloured walks ending at the matching ghost vertex (and paths leaving the boundary) but we still use the superscript *loop* to be consistent with the notation above. Correlations are given by

$$\mathbb{G}_{G, N, \mathbf{J}, \mathbf{h}, U}^\eta(A) := \frac{Z_{G, N, \mathbf{J}, \mathbf{h}, U}^\eta(A)}{Z_{G, N, \mathbf{J}, \mathbf{h}, U}^{\text{loop}, \eta}}. \quad (4.5)$$

These correlations are equal to the correlations presented in [13] for the first description of the RPM in Section 2.1. The proof that the description above leads to the same correlation functions is identical to the proof of Proposition 2.7. The proof of the following proposition can be found in [13] for the case of homogeneous couplings and external field. As explained in [13, Section 5.2], the extension to the inhomogeneous case is straightforward.

**Proposition 4.3.** *Let  $N \in \mathbb{N}_{>0}$ ,  $\mathbf{J}$  such that  $J_e^i \geq 0$  for all  $e \in E$  and  $i \in [N]$ ,  $\mathbf{h}$  such that  $h_x^i \geq 0$  for all  $x \in V$  and  $i \in [N]$ , and vertex weight functions  $U_x^{(N)}$  as in Proposition 2.3. For any  $A \subset V$  and  $\eta \in \{\text{free}, +\}$  we have that*

$$\mathbb{G}_{G, N, \mathbf{J}, \mathbf{h}, U}^\eta(A) = \left\langle \prod_{x \in A} S_x^1 \right\rangle_{G, N, \mathbf{J}, \mathbf{h}}^\eta.$$

Now that we have this proposition the proof of Theorem 4.1 can proceed as in the case of no external field. Indeed, we can think of  $\tilde{E}_g$  as being boundary edges of  $G \subset \mathcal{G} = G_g$ .

We can define the sum and difference of configurations, and the partial ordering as in Section 3, we can also interpret configurations as collections of multigraphs and define  $\mathcal{F}_A \subset \mathcal{M}_{G_g}^N(1)$  ( $\mathcal{F}_A^+ \subset$

$\mathcal{M}_{G_{g \cup \partial E}}^N(1)$  as in Section 3 with the difference that connected components must contain an even number of vertices from  $A$  or be connected to  $g_1$  (or  $\partial E$ ). Indeed, the ghost vertices act as boundary vertices, as mentioned above, so 1-walks may leave  $G$  by going to  $g_1$ . We also define  $P_{A,B}(\mathbf{m})$  to be pairings such that no 1-walk has an end point on  $A$  and an end point on  $B$ .

Theorem 4.1 is a consequence of the following lemma, in the same way that Theorem 1.1 follows from Lemma 3.1.

**Lemma 4.4.** *Let  $A, B \subset V$  with  $A \cap B = \emptyset$  and let vertex weight functions  $U_x^{(N)}$  be fixed as in Proposition 2.3 and  $\eta = \text{free}$ . For any  $N \in \mathbb{N}_{>1}$ , non-negative couple constants  $\mathbf{J} = (J_e^1, \dots, J_e^N)_{e \in E}$ , non-negative external field  $\mathbf{h} = (h_x^1, \dots, h_x^N)_{x \in V}$ , and  $F : \mathcal{M}_G^N \rightarrow \mathbb{R}_{\geq 0}$  we have*

$$\sum_{\substack{\mathbf{m} \in \mathcal{S}(A) \\ \bar{\mathbf{m}} \in \mathcal{S}(B)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \leq \sum_{\substack{\mathbf{m} \in \mathcal{S}(A \cup B) \\ \bar{\mathbf{m}} \in \mathcal{S}(\emptyset)}} F(\mathbf{m} + \bar{\mathbf{m}}) \mu_N(\mathbf{m}) \mu_N(\bar{\mathbf{m}}) \mathbb{1}_{\mathcal{F}_B}(\mathbf{m} + \bar{\mathbf{m}}) \frac{P_{A,B}(\mathbf{m})}{P_G(\mathbf{m})}.$$

The inequality holds for  $\eta = +$  if we replace  $\mathcal{F}_B$  on the right side by  $\mathcal{F}_B^+$ .

The proof of this lemma is identical to the proof of Lemma 3.1 and gives us Theorem 4.1.

## Acknowledgements

It is a pleasure to thank Jess Jay for valuable discussions regarding the proofs and Lorenzo Taggi for helpful comments on a draft of this article.

## References

- [1] M. Aizenman: Geometric analysis of  $\varphi^4$  fields and Ising models. *Comm. Math. Phys.*, Vol. 86(1), pp.1-48 (1982)
- [2] M. Aizenman and B. Nachtergaele: Geometric aspects of quantum spin states. *Comm. Math. Phys.*, Vol. 164, pp.17-63 (1994)
- [3] C. Benassi and D. Ueltschi: Loop correlations in random wire models. *Comm. Math. Phys.*, Vol. 374, pp.525-547 (2020)
- [4] C. Benassi, B. Lees, and D. Ueltschi: Correlation inequalities for the quantum XY model. *J. stat. Phys.*, Vol. 164, pp.1157-1166 (2016)
- [5] C. Benassi, B. Lees, and D. Ueltschi: Correlation Inequalities for classical and quantum XY models In: Advances in Quantum mechanics. *Springer INdAM Series*, Vol. 18, pp.15-31 (2017)
- [6] D. Brydges, J. Fröhlich, and T. Spencer: The random walk representation of classical spin systems and correlation inequalities. *Comm. Math. Phys.*, Vol. 83(1), pp.123-150 (1982)
- [7] H. Duminil-Copin: Lectures on the Ising and Potts models on the hypercubic lattice. In: Random graphs, phase transitions, and the Gaussian free field. *Springer Proc. Math. Stat*, Vol. 304, pp.35-161 (2020)
- [8] F. Dunlop: Correlation inequalities for multicomponent rotors. *Comm. Math. Phys.*, Vol. 49, pp.247-256 (1976)
- [9] J. Ginibre: General formulation of Griffiths' inequalities. *Comm. Math. Phys.*, Vol 16(4), pp.310-328 (1970)
- [10] R. B. Griffiths: Correlations in Ising ferromagnets. I *J. Math. Phys.*, Vol. 8, pp.478-483 (1967)

- [11] H. Kunz, C.E. Pfister, P.A. Vuillermot: Correlation inequalities for some classical spin vector models. *Phys. Lett. A* Vol. 54(6), pp.428-430 (1975)
- [12] B. Lees and L. Taggi: Site monotonicity and uniform positivity for interacting random walks and the Spin  $O(N)$  model with arbitrary  $N$ . *Comm. Math. Phys.*, Vol. 376(1), pp.487-520 (2020)
- [13] B. Lees and L. Taggi: Exponential decay of transverse correlations for  $O(N)$  spin systems and related models. *Prob. Th. Rel. Fiel.*, Vol. 180, pp.1099-1133 (2021)
- [14] D. van Engelenburg and M. Lis: An elementary proof of phase transition in the planar XY model. *Arxiv preprint ArXiv:2110.09465*
- [15] A. Quitmann and L. Taggi: Macroscopic loops in the Bose gas, Spin  $O(N)$  and related models. *Arxiv preprint ArXiv:2201.04047*
- [16] K. Symanzik: Euclidean quantum field theory. I. Equations for a scalar model. *J. Math. Phys*, Vol. 7, pp.510-525 (1966)
- [17] L. Taggi: Uniformly Positive Correlations in the Dimer Model and Macroscopic Interacting Self-Avoiding Walk in  $\mathbb{Z}^d$ ,  $d \geq 3$ . *Comm. Pure and App. Math.*, Vol. 75(6), pp.1183-1236 (2021)
- [18] D. Ueltschi: Random loop representations for quantum spin systems. *J. Math. Phys*, Vol. 54, 083301 (2013)