

# Entanglement polygon inequalities for pure states in qudit systems

Xian Shi\*

College of Information Science and Technology, Beijing University of Chemical Technology, Beijing 100029, China  
(Dated: May 27, 2022)

Entanglement is one of the important resources in quantum tasks. Recently, Yang *et al.* [arXiv:2205.08801] proposed an entanglement polygon inequalities (EPI) in terms of some entanglement measures for  $n$ -qudit pure states. Here we continue to consider the entanglement polygon inequalities. Specifically, we show that the EPI is valid for  $n$ -qudit pure states in terms of geometrical entanglement measure (GEM), then we study the residual entanglement in terms of GEM for pure states in three-qubit systems. At last, we present counterexamples showing that the EPI is invalid for higher dimensional systems in terms of negativity, we also present a class of states beyond qubits satisfy the EPI in terms of negativity.

PACS numbers: 03.65.Ud, 03.67.Mn

## I. INTRODUCTION

Quantum entanglement is an essential feature of quantum mechanics. It plays an important role in quantum information and quantum computation theory [1], such as superdense coding [2], teleportation [3], and the speedup of quantum algorithms [4].

One of the important properties of multipartite entanglement is that entanglement cannot be freely shared. For a tripartite entangled state  $\rho_{ABC}$ , there exists six different bipartite entanglements  $E_{A|B}, E_{A|C}, E_{B|C}, E_{A|BC}, E_{B|AC}$  and  $E_{C|AB}$ . Here  $E$  is an entanglement measure for bipartite systems,  $E_{A|BC}$  can be seen as a *one to group* entanglement, and  $E_{A|B}$  can be seen as a *one to one* entanglement. In 2000, Coffman, Kundu and Wootters showed a famous inequality for a three qubit system  $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_C$  [5],

$$E_{A|BC} \geq E_{A|B} + E_{A|C}, \quad (1)$$

This inequality means that the entanglement between a singled out qubit to a group of qubits is bounded by the sum of the entanglement between the singled out qubit to each qubit in the group. Later, this inequality is generalized to multiqubit systems in terms of the squashed entanglement measure [6], the squared entanglement of formation [7], the squared Tsallis- $q$  entanglement measure [8], and the squared Renyi entanglement measure [9]. As (1) is not valid for multipartite higher dimensional systems in terms of generic bipartite entanglement measures, other types of monogamy of entanglement were presented [10–12].

Recently, entanglement polygon inequality (EPI) that is on the *one to group* entanglement was proposed for multipartite entangled pure states  $\otimes_i \mathcal{H}_i$  [13, 14],

$$E_{j|\bar{j}} \leq \sum_{k \neq j} E_{k|\bar{k}}. \quad (2)$$

This inequality was proved valid for pure states in multi-qubit systems in terms of bipartite entanglement measures [13] and in arbitrary dimensional systems in terms of the  $q$ -concurrence [18] when  $q \geq 2$  and the unified- $(q, s)$  entangled measure when  $q \geq 1, s \geq 0$  [14]. However, whether the EPI is valid for pure states in higher dimensional systems in terms of concurrence or negativity is unknown.

This paper is organized as follows. In Sec. II, we present the preliminary knowledge needed here, In Sec. III, we present our main results, first we present that the EPI is valid in terms of GEM for pure states in arbitrary dimensional systems, we also consider the entanglement indicators based the EPI. At last, We present a class of pure states that doesnot satisfy the EPI in terms of negativity. Moreover, we present a class of pure states in higher dimensional systems satisfy the EPI in terms of negativity, we also present that the EPI is valid in terms of concurrence for pure states in arbitrary dimensional systems. In Sec. IV, we end with a summary.

## II. PRELIMINARIES

An  $n$ -partite pure state  $|\psi\rangle_{A_1 A_2 \dots A_n}$  is full product if it can be written as

$$|\psi\rangle_{A_1 A_2 \dots A_n} = |\phi_1\rangle_{A_1} |\phi_2\rangle_{A_2} \dots |\phi_n\rangle_{A_n}, \quad (3)$$

otherwise, it is entangled. A multipartite pure state is called genuinely entangled if

$$|\psi\rangle_{A_1 A_2 \dots A_n} \neq |\phi\rangle_S |\zeta\rangle_{\bar{S}}, \quad (4)$$

for any partite  $S|\bar{S}$ , here  $S$  is a subset of  $\mathbf{A} = \{A_1, A_2, \dots, A_n\}$ , and  $\bar{S} = \mathbf{A} - S$ .

Next we recall some entanglement measures for a bipartite state  $\rho_{AB}$ . A bipartite pure state  $|\psi\rangle_{AB}$  can be always written as

$$|\psi\rangle_{AB} = \sum_{i=0}^{d-1} \sqrt{\lambda_i} |i\rangle_A |i\rangle_B,$$

\*shixian01@gmail.com

where  $\lambda_i \geq \lambda_{i+1} \geq 0$ ,  $i = 0, 1, \dots, d-2$ , and  $\sum_i \lambda_i = 1$ . The geometric entanglement measure for a pure state  $|\phi\rangle_{AB} = \sum_i \sqrt{\lambda_i} |ii\rangle$  is defined as

$$G(|\phi\rangle) := 1 - \max_{|\psi\rangle=|a_1\rangle|a_2\rangle} \langle \psi | \rho | \psi \rangle \quad (5)$$

The geometric entanglement measure for a mixed state  $\rho_{AB}$  is generally defined as [15],

$$G(\rho_{AB}) = \min \sum_i p_i G(|\phi_i\rangle), \quad (6)$$

where the minimum takes over all the decompositions of  $\rho_{AB}$ .

Negativity is an important entanglement measure for bipartite systems. It is defined as follows for a pure state  $|\phi\rangle_{AB}$  [16],

$$N(|\phi\rangle_{AB}) = \frac{\|\psi\| \langle \psi | T_A | \psi \rangle - 1}{2}, \quad (7)$$

here  $\langle ij | \rho_{AB}^{T_A} | kl \rangle = \langle kj | \rho_{AB} | il \rangle$ .

Concurrence is the other entanglement measure for bipartite mixed states  $\rho_{AB}$ . The concurrence of a pure state  $|\psi\rangle_{AB}$  is defined as

$$C(|\psi\rangle_{AB}) = \sqrt{2(1 - \text{Tr} \rho_A^2)}. \quad (8)$$

Here  $\rho_A = \text{Tr}_B |\psi\rangle_{AB} \langle \psi|$ . For a mixed state  $\rho_{AB}$ , it is defined as

$$C(\rho_{AB}) = \min_{\{p_i, |\phi_i\rangle_{AB}\}} \sum_i p_i C(|\phi_i\rangle_{AB}), \quad (9)$$

where the minimum takes over all the decompositions of  $\rho_{AB} = \sum_i p_i |\phi_i\rangle_{AB} \langle \phi_i|$  with  $p_i \geq 0$  and  $\sum p_i = 1$ .

For a two-qubit mixed state  $\rho_{AB}$ , Wootters derived an analytical formula [17]:

$$C_{AB} = \max\{\sqrt{\mu_1} - \sqrt{\mu_2} - \sqrt{\mu_3} - \sqrt{\mu_4}, 0\}, \quad (10)$$

where  $\mu_1, \mu_2, \mu_3$ , and  $\mu_4$ , are the eigenvalues of the matrix  $\rho_{AB}(\sigma_y \otimes \sigma_y) \rho_{AB}^* (\sigma_y \otimes \sigma_y)$  with nonincreasing order.

Recently, a parametrized entanglement measure, the  $q$ -concurrence, was proposed in [18]. The  $q$ -concurrence for a pure state  $|\psi\rangle_{AB}$  is defined as

$$C_q(|\psi\rangle_{AB}) = 1 - \text{Tr} \rho_A^q. \quad (11)$$

In [14], the authors showed that the  $q$ -concurrence satisfies the EPI for pure states when  $q \geq 2$ .

**Lemma 1** [14] *For any  $n$ -qudit pure entangled state  $|\psi\rangle$  in  $\otimes_i \mathcal{H}_i$ . When  $q \geq 2$ , the following inequality holds,*

$$C_q^{j|\bar{j}} \leq \sum_{k \neq j, \forall k} C_q^{k|\bar{k}}(|\psi\rangle), \quad (12)$$

### III. MAIN RESULTS

In this section, we first recall the EPI proposed in [13, 14], then we present that the geometric entanglement measure (GEM) also satisfies the EPI for three qudit pure states, and we also present an application based on the EPI in terms of GEM. At last, we present a class of counterexamples which shows that negativity is invalid for three qudit pure states, we also present a class of examples that satisfies the EPI in terms of negativity.

#### A. EPI IN TERMS OF GEOMETRIC ENTANGLEMENT MEASURE

Assume  $|\psi\rangle_{AB} = \sum_{i=0}^{d-1} \sqrt{\lambda_i} |ii\rangle$  is a pure state with  $\lambda_i \geq \lambda_{i+1} \geq 0$ ,  $i = 0, 1, \dots, d-2$ , and  $\sum_i \lambda_i = 1$ . By the formula (5), we have

$$G(|\psi\rangle) = 1 - \lambda_0, \quad (13)$$

Next we recall the Schatten  $p$ -norm of a bounded operator  $M$  on the Hilber space  $\mathcal{H}$  with finite dimensions,

$$\begin{aligned} \|M\|_p &:= [\text{Tr}(|M|^p)]^{\frac{1}{p}}, \\ &= \left( \sum_{i=0}^{d-1} s_i^p(M) \right)^{\frac{1}{p}}. \end{aligned} \quad (14)$$

here  $s_i(M)$  are the sigular values of  $M$  with decreasing order, *i.e.* the eigenvalues of  $\sqrt{M^\dagger M}$ . And when  $p \rightarrow \infty$ ,  $\|M\|_p \rightarrow s_0(M)$ . Next we recall the following lemma.

**Lemma 2** [19] *For any bipartite state  $\rho$  on a Hilbert space  $\mathcal{H}_1 \otimes \mathcal{H}_2$ , the inequality*

$$1 + \|\rho\|_q \geq \|\text{Tr}_1 \rho\|_q + \|\text{Tr}_2 \rho\|_1 \quad (15)$$

*holds for  $q > 1$ .*

Clearly, the inequality (15) can be written as

$$\sum_{i=1}^2 1 - \|\text{Tr}_i \rho\|_q \geq 1 - \|\rho\|_q, \quad \text{when } q > 1. \quad (16)$$

Then we have the following result.

**Theorem 3** *Assume  $|\psi\rangle_{A_1 A_2 \dots A_n}$  is a pure state in the following system  $\otimes_{i=1}^n \mathcal{H}_i$ , then we have the following inequality,*

$$G(|\psi\rangle_{P_i | \bar{P}_i}) \leq \sum_{j \neq i} G(|\psi\rangle_{P_j | \bar{P}_j}), \quad (17)$$

*here  $\{P_1, P_2, \dots, P_k\}$  is a partition of the set  $\{A_1, A_2, \dots, A_n\}$  with  $k \leq n$ .*

**Proof.** As

$$G(|\psi\rangle_{AB}) = 1 - s_0(\rho_A) = \lim_{p \rightarrow \infty} [1 - \|\rho_A\|_p], \quad (18)$$

then we have

$$\begin{aligned} G(|\psi\rangle_{P_i|\overline{P_i}}) &= \lim_{p \rightarrow \infty} [1 - \|\rho_{P_i}\|_p] \\ &= \lim_{p \rightarrow \infty} [1 - \|\rho_{\cup_{j \neq i} P_j}\|_p] \\ &\leq \lim_{p \rightarrow \infty} \sum_{j \neq i} [1 - \|\rho_{P_j}\|_q] \\ &= \sum_{j \neq i} G(|\psi\rangle_{P_j|\overline{P_j}}). \end{aligned} \quad (19)$$

Here the first equality is due to (18), the first inequality is due to the inequality (16), and the last inequality is due to the equality (18).  $\square$

Next we propose a generalized EPI for  $n$ -qudit pure states in terms of the GEM.

**Lemma 4** When  $a, b, c \in (0, 1]$ , and  $a + b \geq c$ , then

$$a^\alpha + b^\alpha \geq c^\alpha,$$

when  $\alpha \in (0, 1]$ .

**Proof.** Here we can always assume  $a \geq b$ , and when  $0 < \alpha \leq 1$ ,  $(\frac{a}{b} + 1)^\alpha - (\frac{a}{b})^\alpha \leq 1$ , then we have

$$a^\alpha + b^\alpha \geq (a + b)^\alpha \geq c^\alpha. \quad \square$$

**Corollary 5** Assume  $|\psi\rangle_{A_1 A_2 \dots A_n}$  is a pure state in the following system  $\otimes_{i=1}^n \mathcal{H}_i$ , then we have the following inequality,

$$G(|\psi\rangle_{P_i|\overline{P_i}})^\alpha \leq \sum_{j \neq i} G(|\psi\rangle_{P_j|\overline{P_j}})^\alpha, \quad (20)$$

here  $\{P_1, P_2, \dots, P_k\}$  is a partition of the set  $\{A_1, A_2, \dots, A_n\}$  with  $k \leq n$ , and  $\alpha \in (0, 1]$ .

The above corollary is due to the Theorem 3 and Lemma 4. Then we present an example for the above corollary.

**Example 6**

$$|\psi\rangle_{ABC} = \frac{3}{5}|102\rangle + \frac{2\sqrt{2}}{5}|200\rangle + \frac{2}{5}|010\rangle + \frac{\sqrt{2}}{5}|020\rangle + \frac{\sqrt{2}}{5}|001\rangle,$$

through computation, we have

$$G(|\psi\rangle_{A|BC}) = \frac{9}{25}, G(|\psi\rangle_{B|AC}) = \frac{19}{25}, G(|\psi\rangle_{C|AB}) = \frac{14}{25}, \quad (21)$$

then the inequality (20) can be written as

$$\begin{aligned} g &= G(|\psi\rangle_{A|BC})^\alpha + G(|\psi\rangle_{C|BA})^\alpha - G(|\psi\rangle_{B|AC})^\alpha \\ &= \left(\frac{9}{25}\right)^\alpha + \left(\frac{14}{25}\right)^\alpha - \left(\frac{19}{25}\right)^\alpha, \end{aligned} \quad (22)$$

we plot (22) in Fig. 1, from the figure, we see that the inequality (22) is bigger than 0 when  $\alpha \in (0, 1)$ .




FIG. 1: In this figure, we present the inequality (22).

## B. APPLICATIONS OF THE EPI IN TERMS OF GEM

Assume  $|\psi\rangle_{A_1 A_2 \dots A_n}$  is a pure state in  $\otimes_i \mathcal{H}_i$ , then we consider an entanglement indicator defined as [14, 20],

$$\begin{aligned} \delta_\alpha^G(|\psi\rangle) &= \min_i \tau_\alpha^i(|\psi\rangle), \\ \tau_\alpha^i(|\psi\rangle) &= \sum_{j \neq i} G^\alpha(|\psi\rangle_{A_j|\overline{A_j}}) - G^\alpha(|\psi\rangle_{A_i|\overline{A_i}}). \end{aligned} \quad (23)$$

By the Corollary 5, we have that  $\delta_\alpha^G \geq 0$ ,  $\alpha \in (0, 1)$ . When considering  $|\psi\rangle$  is a three-qubit pure state, we have the following theorem.

**Theorem 7** Assume  $|\psi\rangle_{A_1 A_2 A_3}$  is a three-qubit pure state, then  $\delta_\alpha^G = 0$ ,  $\forall \alpha \in (0, 1)$  if and only if  $|\psi\rangle_{A_1 A_2 A_3}$  is a biseparable state.

**Proof.**  $\Rightarrow$ : When  $\alpha \in (0, 0.5)$ , as  $2\alpha \in (0, 1)$ , we have

$$\begin{aligned} &G^{2\alpha}(|\psi\rangle_{i|\overline{i}}) - G^{2\alpha}(|\psi\rangle_{j|\overline{j}}) - G^{2\alpha}(|\psi\rangle_{k|\overline{k}}) \\ &= (G^\alpha(|\psi\rangle_{j|\overline{j}}) + G^\alpha(|\psi\rangle_{k|\overline{k}}))^2 - G^{2\alpha}(|\psi\rangle_{j|\overline{j}}) - G^{2\alpha}(|\psi\rangle_{k|\overline{k}}) \\ &= 2G^\alpha(|\psi\rangle_{j|\overline{j}})G^\alpha(|\psi\rangle_{k|\overline{k}}) = 0. \end{aligned}$$

Here  $j \neq k \neq i$ . And From the above, we have  $G(|\psi\rangle_{j|\overline{j}}) = 0$  or  $G(|\psi\rangle_{k|\overline{k}}) = 0$ .

A three-qubit pure state  $|\psi\rangle_{A_1 A_2 A_3}$  can be written in the generalized Schmidt decomposition [21]:

$$|\psi\rangle = l_0|000\rangle + l_1 e^{i\theta}|100\rangle + l_2|101\rangle + l_3|110\rangle + l_4|111\rangle, \quad (24)$$

where  $\theta \in [0, \pi)$ ,  $l_i \geq 0$  ( $i = 0, 1, 2, 3, 4$ ), and  $\sum_{i=0}^4 l_i^2 = 1$ . From simple computation, we have the Schmidt coefficients of  $|\psi\rangle$  in terms of  $A|BC$  is

$$Sch(|\psi\rangle_{A|BC}) = (l_0^2, 1 - l_0^2),$$

the Schmidt coefficients of  $|\psi\rangle$  in terms of  $B|AC$  is

$$\text{Sch}(|\psi\rangle_{B|AC}) = \left( \frac{1 + \sqrt{1 - 4\Delta_0}}{2}, \frac{1 - \sqrt{1 - 4\Delta_0}}{2} \right),$$

here  $\Delta_0 = l_0^2 l_3^2 + l_0^2 l_4^2 + l_1^2 l_4^2 + l_2^2 l_3^2 - 2l_1 l_2 l_3 l_4 \cos \theta$ , the Schmidt coefficients of  $|\psi\rangle$  in terms of  $C|AB$  is

$$\text{Sch}(|\psi\rangle_{C|AB}) = \left( \frac{1 + \sqrt{1 - 4\Delta_1}}{2}, \frac{1 - \sqrt{1 - 4\Delta_1}}{2} \right),$$

here  $\Delta_1 = l_0^2 l_2^2 + l_0^2 l_4^2 + l_1^2 l_4^2 + l_2^2 l_3^2 - 2l_1 l_2 l_3 l_4 \cos \theta$ .

When  $G(|\psi\rangle_{A|BC}) = 0$ ,  $l_0 = 0$  or  $l_0 = 1$ , then  $|\psi\rangle_{ABC}$  is biseparable. When  $G(|\psi\rangle_{B|AC}) = 0$ , that is,  $\Delta_0 = 0$ , as  $\Delta_0 = l_0^2(l_3^2 + l_4^2) + (l_1 l_4 - l_2 l_3)^2 + 4l_1 l_2 l_3 l_4 \sin^2 \frac{\theta}{2} = 0$ , then  $l_3 = l_4 = 0$ , or  $l_0 = 0$ , then  $|\psi\rangle_{ABC}$  is biseparable. The case  $G(|\psi\rangle_{C|AB}) = 0$  is similar to  $G(|\psi\rangle_{B|AC}) = 0$ .

---

### Example 8

$$|\psi\rangle_{ABC} = \frac{|000\rangle + |101\rangle + |202\rangle + |310\rangle + |411\rangle + |512\rangle + |620\rangle + |721\rangle + |822\rangle}{3}, \quad (25)$$

According to the definition of Negativity (7), we have that  $N(|\psi\rangle_{A|BC}) = 4$ ,  $N(|\psi\rangle_{B|AC}) = 1$ ,  $N(|\psi\rangle_{C|AB}) = 1$ . Clearly,  $N(|\psi\rangle_{A|BC}) \geq N(|\psi\rangle_{B|AC}) + N(|\psi\rangle_{C|AB})$ , that is, the EPI is invalid for  $|\psi\rangle_{ABC}$  in terms of negativity.

**Remark 9** In the above example, we have that  $\rho_{BC} = \frac{1}{9}I_3 \otimes I_3$ , that is,  $|\psi\rangle_{ABC}$  is a purification of a product state  $\rho_{BC}$ .

Next we present a class of states which doesnot satisfy the EPI in terms of negativity.

**Theorem 10** Assume  $\rho$  and  $\sigma$  are two states on the system  $\mathcal{H}_d$ ,  $\rho = \sum_i a_i |i\rangle\langle i|$ ,  $\sigma = \sum_j b_j |j\rangle\langle j|$ ,  $\text{rank}(\rho), \text{rank}(\sigma) > 1$ , and  $\{|i\rangle\}$  and  $\{|j\rangle\}$  are the orthonormal bases of  $\mathcal{H}_d$ . Let  $|\psi\rangle_{ABC} = \sum_{ij} \sqrt{a_i b_j} |ij\rangle_{AB} |ij\rangle_C$  be a purification of  $\rho_A \otimes \sigma_B$ , then we have

$$N(|\psi\rangle_{C|AB}) \geq N(|\psi\rangle_{A|BC}) + N(|\psi\rangle_{BAC}). \quad (26)$$

**Proof.** Assume  $|\phi\rangle_{AB} = \sum_i \sqrt{\lambda_i} |ii\rangle$ , then

$$N(|\phi\rangle_{AB}) = \frac{(\sum_i \sqrt{\lambda_i})^2 - 1}{2}, \quad (27)$$

Due to (27), we have

$$\begin{aligned} & N(|\psi\rangle_{C|AB}) - N(|\psi\rangle_{A|BC}) - N(|\psi\rangle_{BAC}) \\ &= \frac{1}{2} [(\sum_{ij} \sqrt{a_i b_j})^2 - 1 - (\sum_i \sqrt{a_i})^2 + 1 - (\sum_j \sqrt{b_j})^2 + 1] \\ &= \frac{1}{2} [(1 - (\sum_i \sqrt{a_i})^2)(1 - (\sum_j \sqrt{b_j})^2)]. \end{aligned} \quad (28)$$

As  $\sum_i a_i = \sum_j b_j = 1$ , and  $\text{rank} \rho, \text{rank} \sigma > 1$ , then (28)  $> 0$ . Then we finish the proof.  $\square$

$\Leftarrow$ : Assume  $|\psi\rangle = |\phi\rangle_i |\varphi\rangle_{jk}$ ,  $i \neq j \neq k$ , then  $G(|\psi\rangle_{i|jk}) = 0$ ,  $G(|\psi\rangle_{j|ki}) = G(|\psi\rangle_{k|ji})$ , then  $\delta_\alpha^G = 0$ ,  $\forall \alpha \in (0, 1)$ .  $\square$

### C. EPI IN TERM OF NEGATIVITY AND CONCURRENCE

In this subsection, we present a class of counterexamples that does not satisfy the EPI in terms of negativity and concurrence, which answers the problem proposed in [14]. We also propose a class of multipartite pure states satisfying the EPI in terms of negativity.

First we present an example.

At last, we present a class of pure states, the generalized W class (GW) states, satisfy the EPI in terms of negativity. These states were first studied on the problem of the monogamy relations in terms of concurrence for higher dimensional systems [22]. Recently, the general monogamy relations of this class of states attracts much attention of the revelent researchers [23, 24].

Now let us recall the definition of the GW states  $|W_n^d\rangle$ ,

$$|W_n^d\rangle_{A_1 \dots A_n} = \sum_{i=1}^d (a_{1i} |i0 \dots 0\rangle + \dots + a_{ni} |00 \dots i\rangle), \quad (29)$$

where we assume  $\sum_{i=1}^d \sum_{j=1}^n |a_{ji}|^2 = 1$ . Next we present a lemma and then we show the main result of this subsection.

**Lemma 11** [22] Assume  $|\psi\rangle_{AB_1 B_2 \dots B_{n-1}}$  is a GW state, then for an arbitrary partition  $\{P_1, P_2, \dots, P_m\}$  of the set  $S = \{A, B_1 B_2, \dots, B_{n-1}\}$ , the state  $|\psi\rangle_{P_1 P_2 \dots P_m}$  is also a GW state, here we assume  $P_i \cap P_j = \emptyset$  ( $i \neq j$ ) and  $\cup_i P_i = S, m \leq n$ .

**Theorem 12** Assume  $|\psi\rangle_{A_1 \dots A_n}$  is a GW state, and here we denote  $\{P_1, P_2, P_3\}$  is a partition of the set  $\{A_1, A_2, \dots, A_n\}$ ,  $n \geq 3$ , then we have

$$N(|\psi\rangle_{P_1|P_2 P_3}) \leq N(|\psi\rangle_{P_2|P_1 P_3}) + N(|\psi\rangle_{P_3|P_2 P_1}). \quad (30)$$

**Proof.** Due to the above lemma, we have that  $|\psi\rangle_{P_1 P_2 P_3}$  is also a GW state, that is,  $|\psi\rangle_{P_1 P_2 P_3}$  can be

written as

$$|\psi\rangle_{P_1 P_2 P_3} = \sum_{i=1}^d (a_{1i}|i00\rangle + a_{2i}|0i0\rangle + a_{3i}|00i\rangle), \quad (31)$$

through computation, we have that

$$\begin{aligned} N(|\psi\rangle_{P_1|P_2 P_3}) &= \sqrt{\sum_i |a_{1i}|^2} \times \sqrt{\sum_i (|a_{2i}|^2 + |a_{3i}|^2)}, \\ N(|\psi\rangle_{P_2|P_1 P_3}) &= \sqrt{\sum_i |a_{2i}|^2} \times \sqrt{\sum_i (|a_{1i}|^2 + |a_{3i}|^2)}, \\ N(|\psi\rangle_{P_3|P_1 P_2}) &= \sqrt{\sum_i |a_{3i}|^2} \times \sqrt{\sum_i (|a_{2i}|^2 + |a_{1i}|^2)}, \end{aligned} \quad (32)$$

let  $a = \sum_i |a_{1i}|^2$ ,  $b = \sum_i |a_{2i}|^2$ ,  $c = \sum_i |a_{3i}|^2$ , as

$$\begin{aligned} \sqrt{a(b+c)} &\leq \sqrt{ab+ac+2bc+2\sqrt{bc(a+c)(a+b)}}, \\ &= \sqrt{[\sqrt{b(a+c)} + \sqrt{c(a+b)}]^2}. \end{aligned}$$

Then we finish the proof.  $\square$

Similar to the proof of Theorem 12 and Corollary 5, we can generalize the Theorem 12 to the following.

**Corollary 13** Assume  $|\psi\rangle_{A_1 \dots A_n}$  is a GW state, and here we denote  $\{P_1, P_2, \dots, P_k\}$  is a partition of the set  $\{A_1, A_2, \dots, A_n\}$ ,  $n \geq 3$ , then we have the following inequality,

$$N(|\psi\rangle_{P_i|\bar{P}_i})^\alpha \leq \sum_{j \neq i} N(|\psi\rangle_{P_j|\bar{P}_j})^\alpha, \quad (33)$$

here  $\{P_1, P_2, \dots, P_k\}$  is a partition of the set  $\{A_1, A_2, \dots, A_n\}$  with  $k \leq n$ , and  $\alpha \in (0, 1]$ .

Then we present an example to show the meaning of Theorem (12).

**Example 14** Here we consider a tripartite pure state,

$$|\psi\rangle_{ABCD} = 0.3|0001\rangle + 0.4|0020\rangle + 0.5|0100\rangle + \sqrt{0.5}|1000\rangle,$$

here we take  $P_1 = \{A\}$ ,  $P_2 = \{BC\}$ ,  $P_3 = \{D\}$ , then

$$\begin{aligned} N(|\psi\rangle_{P_1|P_2 P_3}) &= 0.5, N(|\psi\rangle_{P_2|P_1 P_3}) = \sqrt{0.2419}, \\ N(|\psi\rangle_{P_3|P_1 P_2}) &= \sqrt{0.0819}, \end{aligned}$$

then the inequality (33) can be written as

$$\begin{aligned} h &= N(|\psi\rangle_{P_2|P_1 P_3})^\alpha + N(|\psi\rangle_{P_3|P_1 P_2})^\alpha - N(|\psi\rangle_{P_1|P_2 P_3})^\alpha \\ &= 0.2419^{\frac{\alpha}{2}} + 0.0819^{\frac{\alpha}{2}} - 0.5^\alpha, \end{aligned} \quad (34)$$

we plot (34) in Fig. 2, from the figure, we see that the inequality (22) is bigger than 0 when  $\alpha \in (0, 1)$ .




FIG. 2: In this figure, we present the inequality (34).

At last, we present that the EPI is valid in terms of concurrence for pure states in arbitrary dimensional systems.

Comparing the (11) with (8), we have when  $|\psi\rangle_{AB}$  a bipartite pure state,  $C(|\psi\rangle_{AB}) = \sqrt{2C_2(|\psi\rangle_{AB})}$ . Then combing the Lemma 1 and similar proof of Corollary 5, we have the following result,

**Theorem 15** For any  $n$ -qudit pure entangled state  $|\psi\rangle$  in  $\otimes_i \mathcal{H}_i$ , we have

$$C_{j|\bar{j}} \leq \sum_{k \neq j, \forall k} C_{k|\bar{k}}(|\psi\rangle), \quad (35)$$

#### IV. CONCLUSION

In this article, we have investigated the EPI for pure states in multipartite systems. First we present the EPI is valid in terms of GEM for pure states in arbitrary dimensional systems. Based on the inequalities, we have presented the entanglement indicators in terms of GEM. At last, we have presented a class of pure states that does not satisfy the EPI in terms of negativity, we have also presented a class of pure states satisfying the EPI in terms of negativity. Moreover, we have presented the EPI is valid in terms of concurrence for pure states in arbitrary dimensional systems. Due to the importance of the study of the higher-dimensional multipartite entanglement systems, our results can provide a reference for future work on the study of multiparty quantum entanglement.

- 
- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, “Quantum entanglement,” *Reviews of modern physics*, vol. 81, no. 2, p. 865, 2009.
- [2] C. H. Bennett and S. J. Wiesner, “Communication via one-and two-particle operators on einstein-podolsky-rosen states,” *Physical review letters*, vol. 69, no. 20, p. 2881, 1992.
- [3] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, “Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels,” *Physical review letters*, vol. 70, no. 13, p. 1895, 1993.
- [4] Y. Shimoni, D. Shapira, and O. Biham, “Entangled quantum states generated by shor’s factoring algorithm,” *Physical Review A*, vol. 72, no. 6, p. 062308, 2005.
- [5] V. Coffman, J. Kundu, and W. K. Wootters, “Distributed entanglement,” *Physical Review A*, vol. 61, no. 5, p. 052306, 2000.
- [6] M. Christandl and A. Winter, ““squashed entanglement”: an additive entanglement measure,” *Journal of mathematical physics*, vol. 45, no. 3, pp. 829–840, 2004.
- [7] Y.-K. Bai, Y.-F. Xu, and Z. Wang, “General monogamy relation for the entanglement of formation in multiqubit systems,” *Physical review letters*, vol. 113, no. 10, p. 100503, 2014.
- [8] Y. Luo, T. Tian, L.-H. Shao, and Y. Li, “General monogamy of tsallis q-entropy entanglement in multi-qubit systems,” *Physical Review A*, vol. 93, no. 6, p. 062340, 2016.
- [9] W. Song, Y.-K. Bai, M. Yang, M. Yang, and Z.-L. Cao, “General monogamy relation of multiqubit systems in terms of aared rényi- $\alpha$  entanglement,” *Physical Review A*, vol. 93, no. 2, p. 022306, 2016.
- [10] T. R. De Oliveira, M. F. Cornelio, and F. F. Fanchini, “Monogamy of entanglement of formation,” *Physical Review A*, vol. 89, no. 3, p. 034303, 2014.
- [11] G. Gour and Y. Guo, “Monogamy of entanglement without inequalities,” *Quantum*, vol. 2, p. 81, 2018.
- [12] X. Shi, L. Chen, and M. Hu, “Multilinear monogamy relations for multiqubit states,” *Physical Review A*, vol. 104, no. 1, p. 012426, 2021.
- [13] X.-F. Qian, M. A. Alonso, and J. H. Eberly, “Entanglement polygon inequality in qubit systems,” *New Journal of Physics*, vol. 20, no. 6, p. 063012, 2018.
- [14] X. Yang, Y.-H. Yang, and M.-X. Luo, “Entanglement polygon inequality in qudit systems,” *arXiv preprint arXiv:2205.08801*, 2022.
- [15] T.-C. Wei and P. M. Goldbart, “Geometric measure of entanglement and applications to bipartite and multipartite quantum states,” *Physical Review A*, vol. 68, no. 4, p. 042307, 2003.
- [16] G. Vidal and R. F. Werner, “Computable measure of entanglement,” *Physical Review A*, vol. 65, no. 3, p. 032314, 2002.
- [17] W. K. Wootters, “Entanglement of formation of an arbitrary state of two qubits,” *Physical Review Letters*, vol. 80, no. 10, p. 2245, 1998.
- [18] X. Yang, M.-X. Luo, Y.-H. Yang, and S.-M. Fei, “Parametrized entanglement monotone,” *Physical Review A*, vol. 103, no. 5, p. 052423, 2021.
- [19] K. M. Audenaert, “Subadditivity of q-entropies for  $q \geq 1$ ,” *Journal of Mathematical Physics*, vol. 48, no. 8, p. 083507, 2007.
- [20] X.-N. Zhu and S.-M. Fei, “Entanglement monogamy relations of qubit systems,” *Physical Review A*, vol. 90, no. 2, p. 024304, 2014.
- [21] A. Acin, A. Andrianov, L. Costa, E. Jane, J. Latorre, and R. Tarrach, “Generalized schmidt decomposition and classification of three-quantum-bit states,” *Physical Review Letters*, vol. 85, no. 7, p. 1560, 2000.
- [22] J. San Kim and B. C. Sanders, “Generalized w-class state and its monogamy relation,” *Journal of Physics A: Mathematical and Theoretical*, vol. 41, no. 49, p. 495301, 2008.
- [23] X. Shi and L. Chen, “Monogamy relations for the generalized w-class states beyond qubits,” *Physical Review A*, vol. 101, no. 3, p. 032344, 2020.
- [24] L.-M. Lai, S.-M. Fei, and Z.-X. Wang, “Tighter monogamy and polygamy relations for a superposition of the generalized w-class state and vacuum,” *Journal of Physics A: Mathematical and Theoretical*, vol. 54, no. 42, p. 425301, 2021.