

# RATIONAL POINTS ON $x^3 + x^2y^2 + y^3 = k$

XIAOAN LANG AND JEREMY ROUSE

ABSTRACT. We study the problem of determining, given an integer  $k$ , the rational solutions to  $C_k : x^3z + x^2y^2 + y^3z = kz^4$ . For  $k \neq 0$ , the curve  $C_k$  has genus 3 and there are maps from  $C_k$  to three elliptic curves  $E_{1,k}$ ,  $E_{2,k}$ ,  $E_{3,k}$ . We explicitly determine the rational points on  $C_k$  under the assumption that one of these elliptic curves has rank zero.

## 1. INTRODUCTION AND STATEMENT OF RESULTS

A fundamental problem in number theory is to determine the rational solutions to an equation or system of equations with rational coefficients. In 1983, Gerd Faltings proved that if  $C$  is a curve of genus at least 2 defined over a number field  $K$ , then  $C(K)$  is finite. However, the proof is not constructive, and it is still not known if there is an algorithm that will determine the finite set  $C(K)$  for any curve  $C$  of genus at least 2 and any number field  $K$ .

A variety of techniques exist that in some cases allow one to determine the set  $C(K)$ . The most basic is that if  $C$  does not have local points at some completion of  $K$ , then  $C(K) = \emptyset$ . The Chabauty-Coleman method provides a way to determine  $C(K)$  under the assumption that the Mordell-Weil rank of the Jacobian is less than the genus. There are also a handful of other techniques that work in more restricted situations (including maps to lower genus curves, étale descent, and elliptic curve Chabauty). In addition, there is the relatively new Chabauty-Kim method.

In this paper, we consider the family of genus 3 curves  $C_k : x^3z + x^2y^2 + y^3z = kz^4$  over  $\mathbb{Q}$  under the restriction that  $k \in \mathbb{Z}$ . For each  $k$ , this curve has two obvious rational points:  $(1 : 0 : 0)$  and  $(0 : 1 : 0)$ . The map which interchanges  $x$  and  $y$  is an automorphism of  $C_k$  over  $\mathbb{Q}$ , and generically the automorphism group of  $C_k/\overline{\mathbb{Q}}$  is isomorphic to  $S_3$ . As a consequence, the curve admits three independent maps to

---

2010 *Mathematics Subject Classification.* Primary 11G05; Secondary 11G30, 14H45.

elliptic curves. In particular, define

$$\begin{aligned} E_{1,k} &: y^2 + 3xy = x^3 + k \\ E_{2,k} &: y^2 = x^3 + 4kx^2 + 16k^2 \\ E_{3,k} &: y^2 = x^3 - 27x^2 - 1728k. \end{aligned}$$

The corresponding morphisms are given by  $\phi_1 : C_k \rightarrow E_{1,k}$ ,  $\phi_2 : C_k \rightarrow E_{2,k}$  and  $\phi_3 : C_k \rightarrow E_{3,k}$  with

$$\begin{aligned} \phi_1(x : y : z) &= (-xz - yz : xy : z^2) \\ \phi_2(x : y : z) &= (-4x^3z - 4xyz^2, -8x^4 + 16x^3z + 8y^3z - 12kz^4 : z^4) \\ \phi_3(x : y : z) &= ((x - y)z(16x^2y^2 + 12(x^3 + x^2y + xy^2 + y^3)z + 36(x^2 - xy + y^2)z^2) : \\ &\quad 72x^4yz + 108x^4z^2 + 64x^3y^3 + 72x^3y^2z + 108x^3z^3 + 72x^2y^3z + 216x^2y^2z^2 + 72xy^4z + 108y^4z^2 + 108y^3z^3 : \\ &\quad (x - y)^3z^3). \end{aligned}$$

For more detail about how these maps were found, see Section 3. In the event that one of these three curves (say  $E_{i,k}$ ) has rank zero, we can determine  $C_k(\mathbb{Q})$  by computing  $E_{i,k}(\mathbb{Q})$  followed by  $C_k(\mathbb{Q}) = \phi_i^{-1}(E_{i,k}(\mathbb{Q}))$ . The original motivation for considering this family of curves came from a question on MathOverflow from Bogdan Grechuk. The question was whether there are integral solutions to  $x^3 + x^2y^2 + y^3 = 7$ . The answer is no since the map  $\phi_1 : C_k \rightarrow E_{1,k}$  sends integral points to integral points. The integral points on  $E_{1,7}$  can be found using Magma and one can then verify that none of the preimages on  $C_7$  of these integral points are themselves integral.

The main result of this paper is that under the assumption that one  $E_{i,k}$  has rank zero, the only rational points are the obvious ones.

**Theorem 1.** *Let  $k \neq 0$  be an integer and suppose that at least one of  $E_{1,k}$ ,  $E_{2,k}$  and  $E_{3,k}$  has rank zero over  $\mathbb{Q}$ . Then*

- If  $k \neq 2a^3 + a^4$  for some integer  $a$ , then  $C_k(\mathbb{Q}) = \{(1 : 0 : 0), (0 : 1 : 0)\}$ .
- If  $k = 2a^3 + a^4$  for some integer  $a \notin \{-1, 3\}$ , then  $C_k(\mathbb{Q}) = \{(1 : 0 : 0), (0 : 1 : 0), (a : a : 1)\}$ .
- If  $k = -1 = 2 \cdot (-1)^3 + (-1)^4$ , then  $C_k(\mathbb{Q}) = \{(1 : 0 : 0), (0 : 1 : 0), (-1 : 0 : 1), (0 : -1 : 1), (-1 : -1 : 1)\}$ .
- If  $k = 135 = 2 \cdot 3^3 + 3^4$ , then  $C_k(\mathbb{Q}) = \{(1 : 0 : 0), (0 : 1 : 0), (3 : 3 : 1), (-6 : 3 : 1), (3 : -6 : 1)\}$ .

**Remark.** *We are not able to extend our main result to handle the case that  $k \in \mathbb{Q}$ . Changing this hypothesis would require provably finding all of the rational points on the genus 2 curve  $y^2 = x^6 + 6x^5 + 39x^4 + 52x^3 + 39x^2 + 6x + 1$ .*

The classical method of Chabauty and Coleman applies when the rank of the Jacobian is less than the genus. If this condition applies to  $C_k$ , then one of the  $E_{i,k}$  must have rank zero and the main theorem applies. The method of quadratic Chabauty, a

special case of the Chabauty-Kim method, proves in an effective way that  $C(\mathbb{Q})$  is finite in other cases. For more detail, see [2] and [3]. In particular, by Lemma 3.2 of [2], this method will work provided  $\text{rank Jac}(C)(\mathbb{Q}) < g + \rho - 1$ , where  $\rho$  is the Picard number of  $\text{Jac}(C)/\mathbb{Q}$ . Since  $\text{Jac}(C_k)$  is the product of three elliptic curves,  $\rho \geq 3$ , and quadratic Chabauty should be able to provably determine  $C_k(\mathbb{Q})$  provided that the sum of the ranks of  $E_{1,k}$ ,  $E_{2,k}$  and  $E_{3,k}$  is  $\leq 4$ . Nonetheless, we do not pursue this method since its application to genus 3 curves is still quite difficult (see [1]).

**Acknowledgements.** *The authors used Magma [5] V2.26-11 for computations.*

## 2. BACKGROUND

For an elliptic curve  $E/\mathbb{Q}$  in the form  $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$ , the set  $E(\mathbb{Q})$  has the structure of an abelian group. More precisely, if  $P$  and  $Q$  are points in  $E(\mathbb{Q})$ , let  $R = (x, y)$  be the third intersection point of the line through  $P$  and  $Q$  with the curve, and define  $P + Q = (x, -y - a_1x - a_3)$ . The Mordell-Weil theorem states that  $E(\mathbb{Q})$  is a finitely generated abelian group, and Mazur [9] showed that the torsion subgroup of  $E(\mathbb{Q})$ , denoted  $E(\mathbb{Q})_{\text{tors}}$ , must be cyclic of order 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 12, or isomorphic to  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2k\mathbb{Z}$  for  $k \in \{1, 2, 3, 4\}$ .

For an elliptic curve  $E : y^2 = x^3 + Ax + B$  in short Weierstarss form, we define as usual  $j(E) = \frac{6912A^3}{4A^3 + 27B^2}$ . We have that two elliptic curves  $E_1/\mathbb{Q}$  and  $E_2/\mathbb{Q}$  are isomorphic over  $\overline{\mathbb{Q}}$  if and only if  $j(E_1) = j(E_2)$ . If  $E : y^2 = x^3 + Ax + B$  is an elliptic curve, the quadratic twist  $E_D$  of  $E$  is given by  $E_D : y^2 = x^3 + AD^2x + BD^3$ .

To determine the torsion subgroup of  $E_{1,k}$ ,  $E_{2,k}$  and  $E_{3,k}$ , we use well-known parametrizations of families of elliptic curves with points of various orders. (See for example page 214 of [7].) In addition, we use that if  $E/\mathbb{Q}$  has a point  $P$  of order 7, then  $\{rP : 0 \leq r < 7\}$  is a Galois-stable subgroup of order 7 and this implies that  $E$  has a cyclic 7-isogeny. The 7-isogeny condition in the lemma below is from [14].

**Lemma 2.** *Let  $E/\mathbb{Q}$  be an elliptic curve with  $j(E) \neq 0, 1728$ .*

- *We have that  $E/\mathbb{Q}$  has a rational point of order 2 if and only if there is a rational number  $t$  so that  $j(E) = \frac{t^3}{t+16}$ .*
- *If  $E/\mathbb{Q}$  has a rational point of order 3, then there is a rational number  $t$  so that  $j(E) = \frac{t^3(t+24)}{t-3}$ . Conversely, if  $j(E)$  has this form, then some quadratic twist of  $E$  has a rational point of order 3.*
- *If  $E/\mathbb{Q}$  has a rational point of order 4, then there is a rational number  $t$  so that  $j(E) = \frac{-256(t^2-t+1/16)^3}{t^4(t-1/16)}$ .*

- If  $E/\mathbb{Q}$  has a rational point of order 5, then there is a rational number  $t$  so that  $j(E) = \frac{-(t^4+12t^3+14t^2-12t+1)^3}{t^5(t^2+11t-1)}$ . Conversely, if  $j(E)$  has this form, then some quadratic twist of  $E$  has a rational point of order 5.
- If  $E/\mathbb{Q}$  has a rational point of order 7, then it necessarily has a cyclic 7-isogeny and hence there is a rational number  $t$  so that  $j(E) = \frac{(t^2+5t+1)^3(t^2+13t+49)}{t}$ .

We will use the above lemma to determine the integer values of  $k$  for which  $j(E_{1,k}) = j(E_{3,k})$  have the above forms. This leads to the equation of a curve (involving the parameters  $t$  and  $k$ ). Frequently, this curve has genus 0 and is isomorphic to  $\mathbb{P}^1$ . This leads to a parametrization  $k = f(x)$ ,  $t = g(x)$ , and we need to know the rational values of  $x$  for which  $f(x) \in \mathbb{Z}$ . We handle this question using [12, Lemma 3', p. 72].

**Lemma 3.** *Let  $\phi(X)$  and  $\psi(X)$  be polynomials with integer coefficients and no common (complex) roots. Let  $d$  be the maximum of the degrees of  $\phi$  and  $\psi$ . There is a number  $R$ , depending on  $\phi$  and  $\psi$ , so that for all rational numbers  $\frac{m}{n}$ ,*

$$\gcd\left(n^d \phi\left(\frac{m}{n}\right), n^d \psi\left(\frac{m}{n}\right)\right)$$

*divides  $R$ . Moreover, if  $a_0$  is the leading coefficient of  $\phi$ ,  $F(X)$  and  $G(X)$  are polynomials with rational coefficients so that  $F(X)\phi(X) + G(X)\psi(X) = 1$  and  $A$  is the least common multiples of the denominators of the coefficients of  $F$  and  $G$ , then we can take  $R = Aa_0^{\deg \phi + \deg \psi}$ .*

In some cases, the above lemma reduces the problem of determining integer values of a rational function to solving a Thue equation: an equation of the form  $F(x, y) = m$ , where  $F(x, y)$  is a homogeneous polynomial of degree  $\geq 3$ . Based on the work of Bilu and Hanrot [4], Thue equations can be solved effectively, and we use the Magma [5] computer algebra system to solve the ones that arise.

To complete our analysis, it will be necessary to study the case where there is a rational point  $P \in C_k(\mathbb{Q})$  whose image on  $E_{3,k}$  has order 2. Pairs  $(P, k)$  with this property are parametrized by an elliptic curve  $E$  which turns out to have positive rank. It is necessary to know the rational points in  $E(\mathbb{Q})$  for which the corresponding value of  $k$  is an integer. To handle these points, we use Corollary IX.3.2.2 of [11]. We summarize this result below.

Given a finite set  $S$  of primes, let  $\mathbb{Z}_S$  denote the set of rational numbers  $\frac{a}{b}$  with  $\gcd(a, b) = 1$  so that if  $p$  is a prime divisor of  $b$ , then  $p \in S$ . This ring is integrally closed in  $\mathbb{Q}$ : if  $q \in \mathbb{Q}$  is the root of a monic polynomial with coefficients in  $\mathbb{Z}_S$ , then  $q \in \mathbb{Z}_S$ . A special case of Corollary IX.3.2.2 shows that if  $f : E \rightarrow \mathbb{P}^1$  is a non-constant function that has at least one pole in  $E(\mathbb{Q})$ , then there is a finite set  $S$  of primes with the property that  $\{P \in E(\mathbb{Q}) : f(P) \in \mathbb{Z}_S\} \subseteq \{P \in E(\mathbb{Q}) : x(P) \in \mathbb{Z}_S\}$ , where  $x$  denotes the  $x$ -coordinate of a Weierstrass equation for  $E$ . The latter set

$\{P \in E(\mathbb{Q}) : x(P) \in \mathbb{Z}_S\}$  is known to be finite (by Corollary IX.3.2.1 of [11]) and this set can be found effectively using Magma's command `SIntegralPoints`. This Magma command in turn relies on work of Stroeker and Tzanakis [13] which in turn relies on explicit bounds on linear forms in elliptic logarithms due to S. David [6].

### 3. MAPS FROM $C_k$ TO ELLIPTIC CURVES

For  $k \notin \{0, -\frac{27}{16}\}$ , the curve  $C_k : x^3z + x^2y^2 + y^3z = kz^4$  is a smooth plane quartic and hence has genus 3. The Jacobian  $J(C_k)$  is three-dimensional abelian variety. As a consequence of the maps given in the introduction, this Jacobian decomposes (up to isogeny) as the product of three elliptic curves. We wish to explain briefly how these maps were found.

First, the affine equation for  $C_k$  is symmetric in  $x$  and  $y$ . So letting  $s = x + y$  and  $t = xy$ , the equation of  $C_k$  becomes  $s^3 - 3st + t^2 = k$ . Setting  $X = -s$  and  $Y = t$  the equation becomes  $Y^2 + 3XY = X^3 + k$  and the map  $(x, y) \mapsto (s, t)$  is a degree 2 map.

In addition, the curve  $C_k$  has an automorphism of order 3 defined by  $\phi(x : y : z) = (\zeta_3x : \zeta_3^{-1}y : z)$  where  $\zeta_3 = e^{2\pi i/3}$ . The quotient of  $C_k$  by this subgroup of  $\text{Aut}(C_k)$  generated by  $\phi$  is a genus 1 curve which is isomorphic to  $E_{2,k}$ .

Since  $J(C_k)$  has maps to at least two non-isomorphic elliptic curves, Poincaré's complete reducibility theorem (see [10, Proposition 12.1]) implies that  $J(C_k)$  is isogenous to the product of three elliptic curves. To find the third elliptic curve, we choose several small values of  $k$  and compute the numerator of the zeta function of  $C_k/\mathbb{F}_p$  for several lots of small primes  $p$  using the algorithm of Tuitman (see [15] and [16]) and implemented in Magma. We observe that for  $p \equiv 1 \pmod{3}$ , the numerator of the zeta function has a repeated quadratic factor, while for  $p \equiv 2 \pmod{3}$  it usually does not have a repeated quadratic factor. This suggests that  $J(C_k)/\mathbb{F}_p$  has a repeated factor when  $p \equiv 1 \pmod{3}$  and suggests the remaining elliptic curve is the quadratic twist by  $-3$  of either  $E_{1,k}$  or  $E_{2,k}$ . Checks for a few values of  $k$  indicate that the new elliptic curve is the quadratic twist of  $E_{1,k}$  by  $-3$ .

We proceed to search for a map from  $C_k$  to  $E_{3,k}$ . We choose two small positive integers  $a$  and  $b$  and let  $k = a^3 + a^2b^2 + b^3$ . This forces the point  $(a : b : 1)$  to be on  $C_k$ . For values of  $k$  that have this form, we test if  $E_{3,k}$  has rank 1 and trivial torsion subgroup. If it does, we search for points of small height on  $E_{3,k}$  in the hope that the simplest point in  $E_{3,k}(\mathbb{Q})$  is the image of  $(a : b : 1)$  under the map  $C_k \rightarrow E_{3,k}$ . This leads easily to the map given in the introduction. Once the map is found, it is straightforward to verify that it is correct, and that  $\phi_3 : C_k \rightarrow E_{3,k}$  has degree 6.

## 4. PROOF OF THEOREM 1

Suppose that  $(x : y : 1)$  is a rational point on  $C_k$ . It follows that  $\phi_1(x : y : 1) \in E_{1,k}(\mathbb{Q})$ ,  $\phi_2(x : y : 1) \in E_{2,k}(\mathbb{Q})$ , and  $\phi_3(x : y : 1) \in E_{3,k}(\mathbb{Q})$ . Either all three points have infinite order, or one or more of points are torsion points. The proof of Theorem 1 follows from classifying the values of  $k$  for which these curves have torsion points, together with a finer analysis of rational points of order 2 on  $E_{1,k}$  and  $E_{3,k}$ .

First, we note that  $E_{2,k}(\mathbb{Q})$  has positive rank unless  $k = -1$  or  $k = -2$ . In particular,  $E_{2,k} : y^2 = x^3 + 4x^2 + 16k^2$  and so  $P = (0, 4k)$  is a rational point on  $E_{2,k}$ . Computing the coordinates of  $2P, 3P, \dots, 12P$  as rational functions in  $k$  we find that  $P$  can have finite order if and only if  $k = -1$  or  $k = -2$ . When  $k = -1$ ,  $P$  has order 5 and when  $k = -2$ ,  $P$  has order 6. It suffices then to determine when  $\phi_1(x : y : 1)$  and  $\phi_3(x : y : 1)$  have finite order.

If either of these points has finite order, its order must be in  $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12\}$ . However, we will show that there are only finitely many integers  $k$  for which  $E_{1,k}$  or  $E_{3,k}$  has a rational point of order 3, 4, 5 or 7, and this will suffice.

**Case 1:** Either  $\phi_1(x : y : 1)$  or  $\phi_3(x : y : 1)$  has order 1.

The point of order 1 should be the point  $(0 : 1 : 0)$  in projective space. For  $E_{1,k}$ , the corresponding morphism is  $\phi_1(x : y : 1) = (-x - y : xy : 1)$ . This cannot equal  $(0 : 1 : 0)$ . So only  $E_{3,k}$  will be in consideration.

For  $E_{3,k}$ , the corresponding morphism is

$$\begin{aligned} \phi_3(x : y : 1) = & (16x^2y^2 + 12(x^3 + x^2y + xy^2 + y^3) + 36(x^2 - xy + y^2))(x - y) : \\ & 72x^4y + 108x^4 + 64x^3y^3 + 72x^3y^2 + 108x^3 + 72x^2y^3 + 216x^2y^2 + \\ & 72xy^4 + 108y^4 + 108y^3 : \\ & (x - y)^3 \end{aligned}$$

This equals  $(0 : 1 : 0)$  if and only if  $x = y$ . This implies that  $(x : y : 1)$  on  $C_k$  satisfies  $2x^3 + x^4 = k$ .

**Case 2:** Either  $\phi_1(x : y : 1)$  or  $\phi_3(x : y : 1)$  has order 2.

We will show that this only occurs for  $k \in \{-17, -1, 135, 368\}$ . If  $k \in \{-17, 368\}$ , then  $E_{1,k}$ ,  $E_{2,k}$  and  $E_{3,k}$  all have positive rank.

If  $k = -1$ , we find that  $E_{3,k}(\mathbb{Q}) \cong \mathbb{Z}/2\mathbb{Z}$  and the two points are  $(0 : 1 : 0)$  and  $(24 : 0 : 1)$ . We compute all rational points on  $C_{-1}$  which are preimages of these two points under  $\phi_3$ . For  $(0 : 1 : 0)$  we find  $(-1 : -1 : 1)$ ,  $(0 : 1 : 0)$  and  $(1 : 0 : 0)$ . For

$(24 : 0 : 1)$  we find  $(-1 : 0 : 1)$  and  $(0 : -1 : 1)$ . These are all the rational points on  $C_{-1}$ .

If  $k = 135$ , we also have  $E_{3,k}(\mathbb{Q}) \cong \mathbb{Z}/2\mathbb{Z}$  and the rational points are  $(0 : 1 : 0)$  and  $(72 : 0 : 1)$ . The rational points on  $C_{135}$  that are preimages of  $(0 : 1 : 0)$  are  $(3 : 3 : 1)$ ,  $(0 : 1 : 0)$ ,  $(1 : 0 : 0)$  and the rational points on  $C_{135}$  that are preimages of  $(72 : 0 : 1)$  are  $(-6 : 3 : 1)$  and  $(3 : -6 : 1)$ .

First consider the case of  $E_{1,k}$ . To easily identify points of order 2, we rewrite the model as  $E'_{1,k} : Y^2 = X^3 - \frac{27}{16}X + (k + \frac{23}{32})$ . The map  $g : E_{1,k} \rightarrow E'_{1,k}$  is given by  $g(x, y) = (x + \frac{3}{4}, y + \frac{3}{2}x)$ . Now we want to determine the values of  $k \in \mathbb{Z}$  for which there is some  $P \in C_k(\mathbb{Q})$  so that the  $y$ -coordinate of  $g \circ \phi_1(P) = 0$ .

If  $P = (x : y : 1)$  we obtain the equation  $xy = \frac{3}{2}(x + y)$  and we also have  $k = x^3 + x^2y^2 + y^3 \in \mathbb{Z}$ . We can express  $x$  in terms of  $y$ , and find  $x = \frac{3y}{2y-3}$ . The assumption that  $k \in \mathbb{Z}$  implies  $\left(\frac{3y}{2y-3}\right)^3 + \left(\frac{3y}{2y-3}\right)^2 y^2 + y^3$  is an integer. Write  $y = \frac{a}{b}$  where  $a, b \in \mathbb{Z}$  with  $\gcd(a, b) = 1$  and  $b > 0$ . Thus  $\frac{27a^3b^3 + 9a^4b(2a-3b) + a^3(2a-3b)^3}{b^3(2a-3b)^3}$  is an integer. So  $b \mid 27a^3b^3 + 9a^4b(2a-3b) + a^3(2a-3b)^3$  which implies  $b \mid a^3(2a-3b)^3$ . Since  $\gcd(a, b) = 1$ , this implies that  $b \mid 8$  and so  $b \in \{1, 2, 4, 8\}$ .

If  $b = 1$ , then  $2a - 3 \mid 27a^3$  and this implies  $2a - 3 \mid 8 \cdot 27a^3$ . Since  $2a - 3 \mid (2a)^3 - 3^3$  and thus  $2a - 3 \mid 27((2a)^3 - 3^3)$ . It follows that  $2a - 3 \mid 729$ . There are only finitely many  $a$  with this property and for these values of  $a$  we find  $k \in \{-17, 0, 135, 368\}$ .

If  $b > 1$ , then  $b$  is even and  $2a - 3b \mid (2a)^3 - (3b)^3$  implies that  $2a - 3b \mid (\frac{b}{2})^3(2a)^3 - (\frac{b}{2})^3(3b)^3$ . Also, the assumption that  $k \in \mathbb{Z}$  implies that  $2a - 3b \mid 27a^3b^3$ . Since  $2a - 3b \mid 27(\frac{b}{2})^3(2a)^3 - 27(\frac{b}{2})^3(3b)^3$ , we get that  $2a - 3b \mid \frac{729}{8}b^3$ . Since  $b \in \{2, 4, 8\}$  there are only finitely many possibilities for  $a$ . We find these using Magma and again determine that  $k \in \{-17, 0, 135, 368\}$ .

Now consider the case of  $E_{3,k}$ . If  $\phi_3 : C_k \rightarrow E_{3,k}$  is the morphism given above and the image of  $P = (x : y : 1)$  on  $E_{3,k}$  has order 2, then the numerator of the  $y$ -coordinate of  $\phi_3(x : y : 1)$  must be zero. This equation defines a curve

$$C : 72x^4yz + 108x^4z^2 + 64x^3y^3 + 72x^3y^2z + 108x^3z^3 \\ + 72x^2y^3z + 216x^2y^2z^2 + 72xy^4z + 108y^4z^2 + 108y^3z^3 = 0$$

of genus 2. This curve  $C$  is isomorphic to  $y^2 = x^6 + 6x^5 + 39x^4 + 52x^3 + 39x^2 + 6x + 1$  and Magma finds that the rank of the Jacobian of  $C$  is 2. For this reason, the method of Chabauty-Coleman cannot provably find all the rational points on  $C$ . The map  $(x : y : 1) \mapsto (y : x : 1)$  is an automorphism of  $C$ , and we proceed by finding a map from  $C$  to an elliptic curve and finding the points on this elliptic curve that correspond to integral values of  $k$ .

Since  $C$  is symmetric in  $x$  and  $y$ , let  $s = x + y$  and  $t = xy$ . We can rewrite the equation for  $C$  in terms of  $s$  and  $t$ , and this map  $(x : y : 1) \mapsto (s : t : 1)$  sends rational points to rational points. The image of this map is the genus 1 curve

$$E_2 : 64t^3 + 72(s^3t - 2st^2) + 108(s^2 - 2t)^2 + 108(s^3 - 3st) = 0.$$

We are able to find a rational point on  $E_2$  and transform this equation into the Weierstrass form  $E : y^2 = x^3 - 12x + 20$ . We may also express the map  $k : C \rightarrow \mathbb{P}^1$  given by  $k(x : y : 1) = x^3 + x^2y^2 + y^3 = s^3 - 3st + t^2$ . To use Corollary IX.3.2.2 of [11] it will be convenient to produce a map  $\tilde{k} : E \rightarrow \mathbb{P}^1$  so that the composition  $C \rightarrow E_2 \rightarrow E \rightarrow \mathbb{P}^1$  is the same as  $k$ . After this map is found, we pre-compose it with a translation to ensure that  $\tilde{k}$  has a pole at  $(0 : 1 : 0)$ . This pre-composition does not affect the set of points of the form  $(k : 1)$  in the image of  $\tilde{k}(E(\mathbb{Q}))$  with  $k \in \mathbb{Z}$ .

Now we need to find all  $P$  that  $P \in E(\mathbb{Q})$  so  $\tilde{k}(P) \in \mathbb{Z}$ . We follow the proof of Corollary IX.3.2.2 and get that if  $\tilde{k}(P) \in \mathbb{Z}$ ,  $x(P) \in \mathbb{Z}_{\{2\}}$ . To show this, we have  $E : y^2 = x^3 - 12x + 20$  and write

$$\tilde{k}(x : y : 1) = \frac{\phi(x) + \psi(x)y}{\eta(x)}.$$

We find

$$\begin{aligned} \phi(x) &= \frac{15}{16}x^7 - 4x^6 + \frac{15}{2}x^5 - \frac{51}{2}x^4 + 113x^3 - 246x^2 + 240x - 64 \\ \psi(x) &= \frac{1}{8}x^6 + \frac{3}{2}x^5 - \frac{39}{4}x^4 + 25x^3 - 33x^2 + 24x \\ \eta(x) &= x^6 - 12x^5 + 60x^4 - 160x^3 + 240x^2 - 192x + 64. \end{aligned}$$

We compute  $(\tilde{k}\eta(x) - \phi(x))^2 = (\psi(x)y)^2 = \psi(x)^2(x^3 - 12x + 20)$ . Writing this out as a polynomial in  $x$  with coefficients in  $\mathbb{Q}[\tilde{k}]$ , we see that the highest power of  $x$  comes from  $\psi(x)^2x^3$ . In particular,  $x$  is the root of a monic polynomial in  $\mathbb{Q}[\tilde{k}]$ . Then by eliminating the denominator of every coefficient, we can get the relation:

$$-4x^{15} + p_1 + \tilde{k}(-480x^{13} + p_2) + \tilde{k}^2(256x^{12} + p_3) = 0$$

where  $p_1(x)$ ,  $p_2(x)$ , and  $p_3(x)$  are some polynomials of degree smaller than 15, 13, and 12 respectively.

If  $(x : y : 1)$  is a point on  $E$  for which  $\tilde{k}(x : y : 1) \in \mathbb{Z}$ , it follows that  $\tilde{k}(x : y : 1) \in \mathbb{Z}_{\{2\}}$ . The equation above shows that  $x$  is integral over  $\mathbb{Z}_{\{2\}}$  and therefore  $x \in \mathbb{Z}_{\{2\}}$ .

Using Magma we find that the points  $(x, y) \in E(\mathbb{Q})$  with  $x \in \mathbb{Z}_{\{2\}}$  are  $(-4, \pm 2)$ ,  $(-2, \pm 6)$ ,  $(\frac{1}{4}, \pm \frac{33}{8})$ ,  $(1, \pm 3)$ ,  $(2, \pm 2)$ ,  $(4, \pm 6)$ ,  $(10, \pm 30)$ ,  $(22, \pm 102)$ , and  $(89, \pm 839)$ . From this it follows that  $\tilde{k}(x, y) = -1$  or  $135$ .



**Remark.** *There are infinitely many integers  $k$  for which  $E_{1,k}$  and  $E_{3,k}$  have rational points of order 2. This is the reason for considering the more delicate question of when there is a rational point  $P \in C_k(\mathbb{Q})$  whose image on  $E_{1,k}$  or  $E_{3,k}$  has order 2.*

**Case 3:**  $E_{1,k}(\mathbb{Q})$  or  $E_{3,k}(\mathbb{Q})$  contains a rational point of order 3.

We will show that this occurs only for  $k = -72$  and  $k = -2$ . If  $k = -72$ , then  $E_{3,k}(\mathbb{Q}) = \{(0 : 1 : 0)\}$  and pulling back this point we find that  $C_k(\mathbb{Q}) = \{(0 : 1 : 0), (1 : 0 : 0)\}$ . If  $k = -2$ , we find that  $E_{1,k}(\mathbb{Q}) = \{(0 : 1 : 0)\}$  and pulling back this point we find that  $C_k(\mathbb{Q}) = \{(0 : 1 : 0), (1 : 0 : 0)\}$ .

If  $E_{1,k}$  or  $E_{3,k}$  has a rational point of order 3, then there is a rational number  $t$  so that

$$j(E_{1,k}) = j(E_{3,k}) = \frac{-3^7}{16k^2 + 27k} = \frac{t^3(t + 24)}{t - 3}.$$

The equation above in  $k$  and  $t$  parametrizes a curve of genus zero with a non-singular rational point. We parametrize this curve and find that

$$k = \frac{u^3(-2u - 3v)}{(u + v)^3(u - 3v)}.$$

for some integers  $u$  and  $v$  with  $\gcd(u, v) = 1$ .

Now, we apply Lemma 3. Let  $\psi(x) = -2x^4 - 3x^3$  and  $\phi(x) = (x + 1)^3(x - 3)$ . We find there exist  $F(x)$  and  $G(x)$  that  $F(x)\phi(x) + G(x)\psi(x) = 1$  and obtain that

$$\gcd(v^4\phi(u/v), v^4\psi(u/v))$$

divides 243.

Since we assume that  $k$  is an integer, it follows that  $(u + v)^3(u - 3v)$  divides the numerator, and hence  $(u + v)^3(u - 3v)$  divides 243. For each divisor  $d$  of 243, we use Magma to solve the Thue equation  $(u + v)^3(u - 3v) = d$ . Using this process, we find that the only integral values of  $k$  of the form

$$\frac{u^3(-2u - 3v)}{(u + v)^3(u - 3v)}$$

are  $k = -72$ ,  $k = -2$ ,  $k = 0$ ,  $k = 2058$ , and  $k = 56000$ . However,  $k$  having this form is a necessary, but not sufficient condition for  $E_{1,k}$  or  $E_{3,k}$  to have a rational point of order 3. For these values of  $k$  we compute the torsion subgroup of  $E_{1,k}$  and  $E_{3,k}$  and obtain the desired result.

**Case 4:**  $E_{1,k}(\mathbb{Q})$  or  $E_{3,k}(\mathbb{Q})$  contains a rational point of order 4.

We will show that this only occurs if  $k = 135$ , and we previously found  $C_{135}(\mathbb{Q})$  in Case 2.

An elliptic curve has a point of order 4 if and only if it can be put in the form  $y^2 + xy + ty = x^3 + ty^2$  for some  $t \in \mathbb{Q}$ . So if  $k$  is an integer and  $E_{1,k}$  or  $E_{3,k}$  has a point of order 4, then

$$j(E_{1,k}) = j(E_{3,k}) = \frac{-3^9}{16k^2 + 27k} = \frac{-256(t^2 - t + 1/16)^3}{t^4(t - 1/16)}$$

for some rational number  $t$ .

This equation in  $k$  and  $t$  defines a genus zero curve which is birational to  $\mathbb{P}^1$  and it follows that  $k$  must have the form

$$k = \frac{-\frac{27}{16}t^6 + \frac{243}{16}t^4}{(t^2 - 3)^3}.$$

Let  $\mathbb{Q}(t)$  be the collection of rational functions. We wish to determine the  $x$ -coordinates of points of  $E_{1,k}$  and  $E_{3,k}$  with order 4 in  $\mathbb{Q}(t)$ . We will use the Magma command `DivisionPolynomial(E,4)` to compute these and find that the roots for  $E_{3,k}$  are  $(18t^2 - 54t)/(t^2 - 3)$  and  $(18t^2 + 54t)/(t^2 - 3)$ . The roots for  $E_{1,k}$  are  $(-3/2t^2 - 9/2t)/(t^2 - 3)$  and  $(-3/2t^2 + 9/2t)/(t^2 - 3)$ .

Write  $t = a/b$  for integers  $a$  and  $b$  with  $\gcd(a, b) = 1$ . We get

$$k = \frac{-\frac{27a^6}{16b^6} + \frac{243a^4}{16b^4}}{(\frac{a^2}{b^2} - 3)^3} = \frac{-\frac{27}{16}a^6 + \frac{243}{16}a^4b^2}{(a^2 - 3b^2)^3} = \frac{-27a^6 + 243a^4b^2}{16(a^2 - 3b^2)^3}.$$

Since  $k$  should be an integer,  $16(a^2 - 3b^2)^3 \mid (-27a^6 + 243a^4b^2)$ . Now we consider two situations.

If  $3 \nmid a$ , then because  $\gcd(a, b) = 1$ , one can show that  $a^2 - 3b^2 \mid a^2 - 9b^2$  implies  $a^2 - 3b^2 \mid 2$ . Considering the congruence situation modulo 3, the only possibilities are  $a^2 - 3b^2 = 1$  and  $a^2 - 3b^2 = -2$ . For  $a^2 - 3b^2 = -2$ , both  $a$  and  $b$  should be odd numbers. Otherwise, either  $a^2 - 3b^2$  is odd or  $\gcd(a, b)$  is greater than 1. If  $a^2 - 3b^2$  is odd, then  $-128 \mid (-27a^4(a^2 - 9b^2))$  and  $-128 \mid (-2 - 6b^2)$ . Then  $64 \mid 1 - 3b^2$ . Because  $b$  is an odd number,  $1 - 3b^2 \equiv 6 \pmod{8}$  and this shows that  $64 \nmid 1 - 3b^2$ . The conclusion is if  $3 \nmid a$ , then  $a^2 - 3b^2 = 1$ .

If  $3 \mid a$ , then let  $a = 3c$ . Because  $\gcd(a, b) = 1$ ,  $3 \nmid b$  and  $\gcd(b, c) = 1$ . In a similar way, one can derive  $3c^2 - b^2 = -1$  or  $-2$  and so  $a^2 - 3b^2 = -3$  or  $a^2 - 3b^2 = -6$ . This shows that if  $3 \mid a$ , then  $a^2 - 3b^2 = -3$  or  $a^2 - 3b^2 = 6$ .

In particular, in any case we have  $a^2 - 3b^2 \in \{1, -3, 6\}$ .

The conditions on  $k$  and the rational number  $t$  ensures that  $E_{1,k}$  and  $E_{3,k}$  have points of order 4 whose  $x$ -coordinate is rational. The condition that the  $y$ -coordinate is rational gives rise to an elliptic curve. In the case of  $E_{3,k}$ , there is a rational number  $u$  so that  $u^2 = 3(t - 1)(t - 3)(t^2 - 3)$  or  $u^2 = 3(t + 1)(t + 3)(t^2 - 3)$ . In the first case

the elliptic curve has rank zero and torsion subgroup  $\mathbb{Z}/2\mathbb{Z}$ . This means that the only solutions are  $t = 1$  and  $t = 3$ . The former gives  $k = -27/16$ , which is not integral, and the latter gives  $k = 0$ , and we assume  $k \neq 0$ . The second case is similar; the elliptic curve has rank zero and torsion subgroup  $\mathbb{Z}/2\mathbb{Z}$  and the two points correspond to  $k = -27/16$  and  $k = 0$  again. So we have shown that there are no integral values of  $k$  for which  $E_{3,k}$  has a point of order 4.

In the case of  $E_{1,k}$ , the  $x$ -coordinate of a point of order 4 is  $\frac{-(3/2)t^2+(9/2)}{t^2-3}$  or  $\frac{-(3/2)t^2-(9/2)}{t^2-3}$ . In order for the  $y$ -coordinate to be rational, we need a rational solution to  $u^2 = -(t+1)(t+3)(t^2-3)$ , or  $u^2 = -(t-1)(t-3)(t^2-3)$  in the other case. Both cases give rise to the elliptic curve  $y^2 = x^3 - 12x$ , which has rank 1. Also, we can change  $u^2 = -(t-1)(t-3)(t^2-3)$  to  $u^2 = -(-t+1)(-t+3)(t^2-3)$ . Negating  $t$  will not affect the value  $k$  because the powers of  $t$  in the polynomial are even, we only focus on  $u^2 = -(t+1)(t+3)(t^2-3)$ .

This can be transformed into  $y^2 = x^3 - 12x$  where  $x = \frac{2a+6b}{a+b}$ . Rather than use the technique of Corollary IX.3.2.2, we will use an elementary argument based on Pythagorean triples to find the integral points on  $y^2 = x^3 - 12x$  for which  $k$  is an integer.

We have that

$$y^2 = x^3 - 12x = \frac{-16(a+b)(a+3b)(a^2-3b^2)}{(a+b)^4} = \frac{(-16 \text{ or } 48 \text{ or } -96)(a+b)(a+3b)}{(a+b)^4}$$

is a perfect square using that  $a^2 - 3b^2 = 1, -3$ , or  $6$ .

Since it is a perfect square, we can use Pythagorean triples to show that there are only finitely many options for the coprime integers  $a$  and  $b$ .

First, if  $a^2 - 3b^2 = 1$ , then  $\frac{-16(a+b)(a+3b)}{(a+b)^4}$  is a perfect square. Write  $-(a+b)(a+3b) = u^2$  for a positive integer  $u$ . So  $-(a^2 + 4ab + 3b^2) = b^2 - (a + 2b)^2 = u^2$ . Then  $a + 2b, u, b$  is a Pythagorean triple. By Theorem 8.7 on page 101 of [8], there exist relatively prime positive integers  $s$  and  $t$ , one even and one odd, such that  $b = s^2 + t^2$ , and  $\{u, a + 2b\} = \{2st, s^2 - t^2\}$ .

If  $a$  is even then  $a + 2b = 2st$ , and so  $a = 2st - 2s^2 - 2t^2$ . So  $a^2 - 3b^2 = s^4 - 8s^3t + 6s^2t^2 - 8st^3 + t^4 = 1$ . By using the Thue function in Magma, we find  $(s, t)$  can be  $(1, 0)$ ,  $(-1, 0)$ ,  $(0, 1)$ , or  $(0, -1)$ . In this situation,  $a = -2$  and  $b = 1$  and so  $t = -2$  and this yields  $k = 135$ .

If  $a$  is odd then  $a + 2b = s^2 - t^2$ ,  $a = s^2 - t^2 - 2s^2 - 2t^2 = -s^2 - 3t^2$ . In this situation,  $1 = a^2 - 3b^2 = -2s^4 + 6t^4$  which is an even number, which is a contradiction.

Second, if  $a^2 - 3b^2 = -3$ , then  $\frac{48(a+b)(a+3b)}{(a+b)^4}$  is a perfect square. Then  $3(a+b)(a+3b) = 3a^2 + 12b^2 + 9b^2 = (2a+3b)^2 - a^2$ . Let  $(2a+3b)^2 - a^2 = u^2$ . Since  $\gcd(a, b) = 1$ , one can also deduce that  $\gcd(2a+3b, a) = 1$  and  $(2a+3b, u, a)$  is a Pythagorean triple. There are relatively prime integers  $s$  and  $t$  such that  $2a+3b = s^2 + t^2$ . In this situation  $b$  is odd.

If  $a$  is odd, then  $a = s^2 - t^2$ . Since  $2a+3b = s^2 + t^2$ ,  $b = t^2 - \frac{s^2}{3}$ . Then  $a^2 - 3b^2 = \frac{2}{3}s^4 - 2t^4$  is equal to  $-3$ . This implies that  $2s^4 - 6t^4 = -9$ , which is a contradiction since  $2s^4 - 6t^4$  is even.

If  $a$  is an even number, then  $a = 2st$ , so  $b = \frac{s^2 + t^2 - 4st}{3}$ . Then  $a^2 - 3b^2 = 4s^2t^2 - \frac{1}{3}(s^4 - 8s^3t + 18s^2t^2 - 8st^3 + t^4) = -\frac{1}{3}s^4 + \frac{8}{3}s^3t - 2s^2t^2 + \frac{8}{3}st^3 - \frac{1}{3}t^4$  is equal to  $-3$ . This shows that  $s^4 - 8s^3t + 6s^2t^2 - 8st^3 + t^4$  is equal to 9. By using the Thue function in Magma, we find that there are no solutions in this case.

Third, if  $a^2 - 3b^2 = 6$ , then  $\frac{-96(a+b)(a+3b)}{(a+b)^4}$  is a perfect square. One can let  $-6(a+b)(a+3b)(a+b)^4 = d^2$ . In the previous proof, we showed that both  $a$  and  $b$  are odd and  $\gcd(a, b) = 1$ . Let  $k = \gcd(a+b, a+3b)$ . Then  $k \mid a+3b - (a+b)$  which means  $k \mid 2b$ . Also,  $k \mid 2(a+b) - 2b$  means  $k \mid 2a$ . This shows that  $k \mid 2b$  and  $k \mid 2a$  and  $\gcd(a, b) = 1$ . We get  $k = 2$  and  $\gcd(\frac{a+b}{2}, \frac{a+3b}{2}) = 1$ . We also get the equation  $-6((a+b)/2)((a+3b)/2) = (d/2)^2$ .

Because  $\gcd(\frac{a+b}{2}, \frac{a+3b}{2}) = 1$  and  $-6((a+b)/2)((a+3b)/2) = (d/2)^2$ ,  $3 \mid \frac{a+3b}{2}$  and there are four possible options: either  $\frac{a+b}{2}$  is a square, the negative of a square, twice a square, or the negative of twice a square.

We can represent the four situations as  $\frac{a+b}{2} = cs^2$  and  $\frac{a+3b}{2} = -(\frac{6}{c})t^2$  with  $c \in \{\pm 1, \pm 2\}$ . Then  $b = \frac{a+3b}{2} - \frac{a+b}{2} = (\frac{6}{c})t^2 - cs^2$  and  $a = 3(\frac{a+b}{2}) - \frac{a+3b}{2} = 3cs^2 + (\frac{6}{c})t^2$ . We also know  $a^2 - 3b^2 = 6$  in this situation. This gives  $a^2 - 3b^2 = 6c^2s^4 - \frac{72}{c^2}t^4 = 6$ . Since  $c \in \{\pm 1, \pm 2\}$  we get  $s^4 - 12t^4 = 1$  or  $4s^4 - 3t^4 = 1$ .

Using the Thue function in Magma, we find that the only solutions are  $(c, a, b) = (1, 3, -1)$ ,  $(-1, -3, 1)$ ,  $(2, 9, -5)$ , and  $(-2, -9, 5)$ . These yield  $t = -3$  or  $t = -9/5$ . These lead to  $k = 0$  and  $k = 59049/8$ .

In conclusion, the only solution we found which yields a value of  $K$  which is a nonzero integer was  $(a, b) = (-2, 1)$ . In this case  $k = 135$  and  $E_{1,k}$  has a rational point of order 4.

**Case 5:**  $E_{1,k}(\mathbb{Q})$  or  $E_{3,k}(\mathbb{Q})$  contains a rational point of order 5.

We will show that this only occurs if  $k = 864$ . We have  $864 = 2 \cdot (-6)^3 + (-6)^4$ . We have that  $E_{3,k}(\mathbb{Q})$  is trivial, and therefore every rational point  $P \in C_{864}(\mathbb{Q})$  maps to

$(0 : 1 : 0)$  on  $E_{3,k}$ . The calculation in Case 1 shows that  $C_{864}(\mathbb{Q}) = \{(1 : 0 : 0), (0 : 1 : 0), (-6 : -6 : 1)\}$ .

If  $E_{1,k}$  or  $E_{3,k}$  has a rational point of order 5, then there is a rational number  $t$  so that

$$j(E_{1,k}) = j(E_{3,k}) = \frac{-3^9}{16k^2 + 27k} = \frac{-(t^4 + 12t^3 + 14t^2 - 12t + 1)^3}{t^5(t^2 + 11t - 1)}$$

The equation above in  $k$  and  $t$  parametrizes a curve of genus zero with a non-singular rational point. Parametrizing this curve shows that

$$16k^2 + 27k = \frac{3^9(u^7v^5 + 11u^6v^6 - u^5v^7)}{(u^4 + 12u^3v + 14u^2v^2 - 12uv^3 + v^4)^3}$$

for some integers  $u$  and  $v$  with  $\gcd(u, v) = 1$ .

Now, we apply Lemma 3. Let  $\psi(x) = 3^9(x^7 + 11x^6 - x^5)$  and  $\phi(x) = (x^4 + 12x^3 + 14x^2 - 12x - 1)^3$ . We find that there exist  $F(x)$  and  $G(x)$  such that  $F(x)\phi(x) + G(x)\psi(x) = 1$  and obtain that

$$\gcd(v^{12}\phi(u/v), v^{12}\psi(u/v))$$

divides 2460375.

Since we assume that  $k$  is an integer, it follows that  $(u^4 + 12u^3v + 14u^2v^2 - 12uv^3 + v^4)^3$  divides the numerator, and hence  $(u^4 + 12u^3v + 14u^2v^2 - 12uv^3 + v^4)^3$  divides 2460375. For each divisor  $d$  of 2460375 which is a cube, we use Magma to solve the Thue equation  $(u^4 + 12u^3v + 14u^2v^2 - 12uv^3 + v^4)^3 = d$ . Using this process, we find that the only integral values of  $k$  of the form

$$\frac{3^9(u^7v^5 + 11u^6v^6 - u^5v^7)}{(u^4 + 12u^3v + 14u^2v^2 - 12uv^3 + v^4)^3}$$

should be the solution of the equation  $16k^2 + 27k = 3^9 \cdot 19 \cdot 2^5 = 11967264$ . By solving the quadratic equation, we find  $k = 13581/16$  or  $k = 864$ . The only integral value is therefore  $k = 864$  and for this value of  $k$  we find that  $E_{1,k}$  has a rational point of order 5 and  $E_{3,k}$  does not.

**Case 6:**  $E_{1,k}(\mathbb{Q})$  or  $E_{3,k}(\mathbb{Q})$  contains a rational point of order 7.

We will show that this does not occur. If  $E_{1,k}$  or  $E_{3,k}$  has a rational point of order 7, then there is a rational number  $t$  so that

$$j(E_{1,k}) = j(E_{3,k}) = \frac{-3^9}{16k^2 + 27k} = \frac{(t^2 + 5t + 1)^3(t^2 + 13t + 49)}{t}$$

The equation above in  $k$  and  $t$  parametrizes a curve of genus zero with a non-singular rational point. Parametrizing this curve shows that

$$16k^2 + 27k = \frac{-3^9(uv^7)}{(u^2 + 5uv + b^2)^3(u^2 + 13uv + 49v^2)}$$

for some integers  $u$  and  $v$  with  $\gcd(u, v) = 1$ .

Now, we apply Lemma 3. Let  $\psi(x) = 3^9x$  and  $\phi(x) = (x^2 + 5x + 1)^3(x^2 + 13x + 49)$ . We find that there exist  $F(x)$  and  $G(x)$  that  $F(x)\phi(x) + G(x)\psi(x) = 1$  and obtain that

$$\gcd(v^8\phi(u/v), v^8\psi(u/v))$$

divides 964467.

Since we assume that  $k$  is an integer, it follows that  $(u^2 + 5uv + b^2)^3(u^2 + 13uv + 49v^2)$  divides the numerator, and hence  $(u^2 + 5uv + b^2)^3(u^2 + 13uv + 49v^2)$  divides 964467. For each divisor  $d$  of 964477, we use Magma to solve the Thue equation  $(u^2 + 5uv + b^2)^3(u^2 + 13uv + 49v^2) = d$ . Using this process, we find that the only integral values of  $k$  of the form

$$\frac{-3^9(uv^7)}{(u^2 + 5uv + b^2)^3(u^2 + 13uv + 49v^2)}$$

should be the solution of the equation  $16k^2 + 27k = 3^7 \cdot 5 = 10935$ . By solving the quadratic equation, we find  $k = 13581/16$  and  $k = -27$ . The only integral  $k$  is  $-27$ . For this value of  $k$ ,  $E_{1,k}$  and  $E_{3,k}$  both have rational 7-isogenies, but no rational point of order 7.

Thus, there are no values of  $k$  for which  $E_{1,k}$  or  $E_{3,k}$  has a rational point of order 7, and this concludes the proof of Theorem 1.

We conclude with a table below of the special cases of  $k$  that arose in the proof above.

$k$	Is $k = 2a^3 + a^4$ ?	$E_{1,k}(\mathbb{Q})$	$E_{2,k}(\mathbb{Q})$	$E_{3,k}(\mathbb{Q})$	Some points $(x : y : 1) \in C_k(\mathbb{Q})$
-1	Yes, $a = -1$	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	$\{(0 : -1 : 1), (-1 : -1 : 1), (-1 : 0 : 1)\}$
-2	No	Trivial	$\mathbb{Z}/6\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z}$	None.
-17	No	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$	$\{(2 : -5 : 1), (1 : -3 : 1), (-5 : 2 : 1), (-3 : 1 : 1)\}$
-72	No	$\mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}$	Trivial	None.
135	Yes, $a = 3$	$\mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z}$	$\{(3 : -6 : 1), (3 : 3 : 1), (-6 : 3 : 1)\}$
368	No	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z} \times \mathbb{Z}$	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z} \times \mathbb{Z}$	$\{(6 : 2 : 1) : (2 : 6 : 1)\}$
864	Yes, $a = -6$	$\mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}$	Trivial	$\{(-6 : -6 : 1)\}$ .

## REFERENCES

- [1] Jennifer Balakrishnan, Netan Dogra, J. Steffen Müller, Jan Tuitman, and Jan Vonk. Explicit Chabauty-Kim for the split Cartan modular curve of level 13. *Ann. of Math. (2)*, 189(3):885–944, 2019.
- [2] Jennifer S. Balakrishnan and Netan Dogra. Quadratic Chabauty and rational points, I:  $p$ -adic heights. *Duke Math. J.*, 167(11):1981–2038, 2018. With an appendix by J. Steffen Müller.
- [3] Jennifer S. Balakrishnan and Netan Dogra. Quadratic Chabauty and rational points II: Generalised height functions on Selmer varieties. *Int. Math. Res. Not. IMRN*, (15):11923–12008, 2021.
- [4] Yuri Bilu and Guillaume Hanrot. Solving Thue equations of high degree. *J. Number Theory*, 60(2):373–392, 1996.

- [5] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [6] Sinnou David. Minorations de formes linéaires de logarithmes elliptiques. *Mém. Soc. Math. France (N.S.)*, (62):iv+143, 1995.
- [7] Daniel Sion Kubert. Universal bounds on the torsion of elliptic curves. *Proc. London Math. Soc.* (3), 33(2):193–237, 1976.
- [8] David C. Marshall, Edward Odell, and Michael Starbird. *Number theory through inquiry*. MAA Textbooks. Mathematical Association of America, Washington, DC, 2007.
- [9] B. Mazur. Modular curves and the Eisenstein ideal. *Inst. Hautes Études Sci. Publ. Math.*, (47):33–186 (1978), 1977. With an appendix by Mazur and M. Rapoport.
- [10] J. S. Milne. Abelian varieties. In *Arithmetic geometry (Storrs, Conn., 1984)*, pages 103–150. Springer, New York, 1986.
- [11] Joseph H. Silverman. *The arithmetic of elliptic curves*, volume 106 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1992. Corrected reprint of the 1986 original.
- [12] Joseph H. Silverman and John Tate. *Rational points on elliptic curves*. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1992.
- [13] R. J. Stroeker and N. Tzanakis. Solving elliptic Diophantine equations by estimating linear forms in elliptic logarithms. *Acta Arith.*, 67(2):177–196, 1994.
- [14] Andrew V. Sutherland and David Zywina. Modular curves of prime-power level with infinitely many rational points. *Algebra Number Theory*, 11(5):1199–1229, 2017.
- [15] Jan Tuitman. Counting points on curves using a map to  $\mathbf{P}^1$ . *Math. Comp.*, 85(298):961–981, 2016.
- [16] Jan Tuitman. Counting points on curves using a map to  $\mathbf{P}^1$ , II. *Finite Fields Appl.*, 45:301–322, 2017.