

# Fourier-Stieltjes algebras, decomposable Fourier multipliers and amenability

Cédric Arhancet - Christoph Kriegler

## Abstract

We prove that the Fourier-Stieltjes algebra  $B(G)$  of a discrete group  $G$  is isometrically isomorphic to the algebra  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers on the group von Neumann algebra  $VN(G)$ . With sharp contrast, we prove that  $\mathfrak{M}^{\infty, \text{dec}}(G) \neq B(G)$  for some class of non-discrete locally compact groups and that  $\mathfrak{M}^{\infty, \text{dec}}(G) = B(G)$  for any (second countable unimodular) inner amenable locally compact group. To prove these results, we connect to groupoid theory and to the problem of the existence of a contractive projection preserving complete positivity from the space of normal completely bounded operators on  $VN(G)$  onto the space  $\mathfrak{M}^{\infty, \text{cb}}(G)$  of completely bounded Fourier multipliers, that we solve affirmatively in the inner amenable case. In addition, we are able to show that such projections does not exist for (second countable) non-amenable connected locally compact groups. Indeed, we examine more generally if the space  $\mathfrak{M}^{p, \text{cb}}(G)$  of completely bounded Fourier multipliers on the noncommutative  $L^p$ -space  $L^p(VN(G))$  is complemented in the space of completely bounded operators, where  $1 \leq p \leq \infty$ . Using doubling metrics on Lie groups and structure theory from the solution to Hilbert's fifth problem, we establish that any (second countable unimodular) finite-dimensional amenable locally compact group admits compatible bounded projections at level  $p = 1$  and  $p = \infty$ . Finally, we also state a related new characterization of amenability for (second countable unimodular) locally compact groups and we also give applications to decomposable (=regular) multipliers.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Fourier-Stieltjes algebras and decomposable multipliers on <math>VN(G)</math></b>	<b>6</b>
2.1	Preliminaries . . . . .	6
2.2	Fourier-Stieltjes algebras and decomposable multipliers on $VN(G)$ . . . . .	11
<b>3</b>	<b>Inner amenability</b>	<b>17</b>
3.1	Preliminaries on inner amenability and amenability . . . . .	17
3.2	Some characterizations of inner amenability . . . . .	19
<b>4</b>	<b>Projections on the space of completely bounded Fourier multipliers</b>	<b>22</b>
4.1	Preliminaries . . . . .	22
4.2	Description of the approach . . . . .	24
4.3	Step 1: the mappings $P_j(T)$ . . . . .	27

2020 *Mathematics subject classification*: 46L51, 43A15, 46L07, 43A07.

*Key words*: noncommutative  $L^p$ -spaces, Fourier-Stieltjes algebras, decomposable operators, Fourier multipliers, Schur multipliers, inner amenability, amenability, complementations, groupoids, operator spaces.

4.4	Step 2: the symbol of $P_j(T)$ is Herz-Schur if $G$ is inner amenable . . . . .	33
4.5	Step 2: the symbol of $P_j(T)$ for a Fourier multiplier $T$ if $p = \infty$ or $p = 1$ . . . . .	35
4.6	Step 2: convergence of the symbols for a multiplier $T$ with arbitrary symbol . . . . .	36
4.7	Step 2: the case of finite-dimensional unimodular amenable locally compact groups . . . . .	37
4.8	Step 3: the projection $Q: \mathfrak{M}_G^{p,cb} \rightarrow \mathfrak{M}_G^{p,cb,HS}$ on the space of Herz-Schur multipliers . . . . .	43
4.9	Steps 1-3 put together: theorems of complementation . . . . .	47
<b>5</b>	<b>Final remarks</b> . . . . .	<b>51</b>
5.1	A characterization of the amenability of unimodular locally compact groups . . . . .	51
5.2	A description of decomposable Fourier multipliers on some locally compact groups . . . . .	52
5.3	$CB_{w^*}$ -strongly non decomposable multipliers . . . . .	52
5.4	Multipliers on reduced $C^*$ -algebras vs multipliers on von Neumann algebras . . . . .	54
	<b>Bibliography</b> . . . . .	<b>55</b>

# 1 Introduction

The Fourier-Stieltjes algebra  $B(G)$  of a locally compact  $G$  is a generalization for non-abelian groups of the algebra of bounded complex measures of an abelian locally compact group. Since its introduction by Eymard in [Eym], this commutative algebra became a central object in noncommutative harmonic analysis, related to the unitary representation theory of  $G$ . More precisely, the elements of  $B(G)$  are exactly the matrix entries of continuous unitary representations of  $G$  (see (2.1)) and the operations of this algebra are pointwise multiplication and addition. Also note that  $B(G)$  is a complete invariant of  $G$ , i.e.  $B(G_1)$  and  $B(G_2)$  are isometrically isomorphic as Banach algebras if and only if  $G_1$  and  $G_2$  are topologically isomorphic as locally compact groups as proved by Walter in [Wal4] (see also [KaL1, Theorem 3.2.5]).

Decomposable maps is a class of operators between  $C^*$ -algebras generalizing completely positive maps. The class of decomposable maps is maybe the most general class of tractable operators. If  $A$  and  $B$  are  $C^*$ -algebras, recall that a linear map  $T: A \rightarrow B$  is decomposable [Haa] if there exist linear maps  $v_1, v_2: A \rightarrow B$  such that the linear map

$$(1.1) \quad \Phi = \begin{bmatrix} v_1 & T \\ T^\circ & v_2 \end{bmatrix} : M_2(A) \rightarrow M_2(B), \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mapsto \begin{bmatrix} v_1(a) & T(b) \\ T^\circ(c) & v_2(d) \end{bmatrix}$$

is completely positive where  $T^\circ(c) \stackrel{\text{def}}{=} T(c^*)^*$ . In this case,  $v_1$  and  $v_2$  are completely positive and the decomposable norm of  $T$  is defined by

$$(1.2) \quad \|T\|_{\text{dec}, A \rightarrow B} \stackrel{\text{def}}{=} \inf \{ \max\{\|v_1\|, \|v_2\|\} \},$$

where the infimum is taken over all maps  $v_1$  and  $v_2$ . See the books [BLM], [ER] and [Pis7] for more information on this classical notion and to [ArK1] and [JuR] for the analogue notion for operators acting on noncommutative  $L^p$ -spaces.

Our first result (see Corollary 2.20) is the following.

**Theorem 1.1** *The Fourier-Stieltjes algebra  $B(G)$  of a discrete group  $G$  is canonically isometrically isomorphic to the algebra  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers on the group von Neumann algebra  $VN(G)$ .*

This identification strengthens the ubiquity of the Fourier-Stieltjes algebra  $B(G)$  of a discrete group  $G$ . Indeed, we will show in Proposition 2.10 that for any locally compact group  $G$  there exists a well-defined injective *contractive* map from the Fourier-Stieltjes algebra  $B(G)$  into the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers and we will also examine the surjectivity of this map. We will show that the following property<sup>1</sup> plays an important role in this problem.

**Definition 1.2** *Let  $G$  be a locally compact group. We say that  $G$  has property  $(\kappa_\infty)$  if there exists a bounded projection  $P_G^\infty: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  preserving the complete positivity onto the space  $\mathfrak{M}^{\infty, \text{cb}}(G)$  of completely bounded Fourier multipliers on  $\text{VN}(G)$ . In this case, the infimum of bounds of such projections will be denoted  $\kappa_\infty(G)$ :*

$$(1.3) \quad \kappa_\infty(G) \stackrel{\text{def}}{=} \inf \|P_G^\infty\|_{\text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))}.$$

It is variant of the relative projection constant of a closed subspace of a Banach space, e.g. [Wot1, page 112]. This property means that the space  $\mathfrak{M}^{\infty, \text{cb}}(G)$ <sup>2</sup> is complemented in the space  $\text{CB}_{w^*}(\text{VN}(G))$ . Indeed, in Proposition 2.14, we will prove that this property suffices to ensure that the previous inclusion  $B(G) \hookrightarrow \mathfrak{M}^{\infty, \text{dec}}(G)$  is a bijection. In order to prove that this map is an isometry, we need a matricial generalization of property  $(\kappa_\infty)$  (satisfied for any discrete group  $G$ ) and surprisingly (at first sight) the use of results on locally compact *groupoids*. Note also that the existence of non-discrete locally compact groups with  $(\kappa_\infty)$  (and even with the stronger property  $(\kappa)$  of Definition 1.7) was a rather surprising result of [ArK1] since the proof of property  $(\kappa_\infty)$  of a discrete group  $G$  is an average argument relying on the compactness of the compact quantum group  $(\text{VN}(G), \Delta)$  defined by the von Neumann algebra  $\text{VN}(G)$  and its canonical coproduct  $\Delta$ . By [ArK1], a second countable pro-discrete locally compact group  $G$  satisfies  $\kappa_\infty(G) = 1$ . With sharp contrast, we will observe in this paper (as announced in [ArK1]) that the unimodular locally compact group  $G = \text{SL}_2(\mathbb{R})$  does not have  $(\kappa_\infty)$ .

If  $G$  is a locally compact group, with our result we can insert the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers on  $\text{VN}(G)$  in the classical contractive inclusion  $B(G) \subset \mathfrak{M}^{\infty, \text{cb}}(G)$ :

$$(1.4) \quad B(G) \subset \mathfrak{M}^{\infty, \text{dec}}(G) \subset \mathfrak{M}^{\infty, \text{cb}}(G).$$

It is known [Pis6, p. 54] that  $B(G) = \mathfrak{M}^{\infty, \text{cb}}(G)$  is a characterization of amenability for locally compact groups. This observation allows us to revisit an another nice characterization of amenability of Lau and Paterson [LaP2, Corollary 3.2] which says that a second countable locally compact group  $G$  is amenable if and only if  $\text{VN}(G)$  is injective and if  $G$  is inner amenable. We state the following conjectures.

**Conjecture 1.3** *Let  $G$  be a locally compact group.*

1.  *$G$  is inner amenable if and only if we have  $B(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$ .*
2. *The von Neumann algebra  $\text{VN}(G)$  is injective if and only if we have  $\mathfrak{M}^{\infty, \text{dec}}(G) = \mathfrak{M}^{\infty, \text{cb}}(G)$ .*

We will prove the “only if” part of the first assertion for unimodular groups by showing that inner amenability implies  $\kappa_\infty(G) = 1$  hence  $B(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$ . We refer to Section 4.2 for a detailed presentation of our approach. This is our first main result.

**Theorem 1.4** *Let  $G$  be a second countable unimodular locally compact group. If  $G$  is inner amenable then we have  $B(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$ .*

1. The subscript  $w^*$  means “weakly\* continuous”.

2. This space is denoted sometimes  $M_0A(G)$  or  $M_{\text{cb}}A(G)$ .

The “only if” part of the second assertion of Conjecture 1.3 is true by a result of Haagerup [Haa]. A consequence of our results is that the second point of Conjecture 1.3 is true for discrete groups, see Theorem 2.22, and also for second countable unimodular inner amenable locally compact groups (see Corollary 5.4).

**Theorem 1.5** *Let  $G$  be a discrete group or a second countable unimodular inner amenable locally compact group. Then the von Neumann algebra  $\text{VN}(G)$  is injective if and only if we have  $\mathfrak{M}^{\infty, \text{dec}}(G) = \mathfrak{M}^{\infty, \text{cb}}(G)$ .*

We also obtain a new characterization of amenability, see Theorem 5.1.

**Theorem 1.6** *Let  $G$  be a second countable unimodular locally compact group. Then the following are equivalent.*

1.  $\text{VN}(G)$  is injective and  $G$  has  $(\kappa_{\infty})$ .
2.  $G$  is amenable.

Finally, we will observe in this paper that the first inclusion of (1.4) can also be strict, e.g. for  $G = \text{SL}_2(\mathbb{R})$ . The converses of these results will need further investigations. We also give other characterizations of inner amenability for unimodular locally compact groups and we will use one of these in the proofs of our results.

On noncommutative  $L^p$ -spaces with  $1 < p < \infty$ , we will prove in Corollary 4.33 that the space  $\mathfrak{M}^{p, \text{cb}}(G)$  of completely bounded Fourier multipliers on the noncommutative  $L^p$ -space  $L^p(\text{VN}(G))$  is contractively complemented by a contractive projection preserving the complete positivity for any (second countable unimodular) amenable locally compact group  $G$ . Note that this property does not characterizes amenability if  $1 < p < \infty$  since a discrete group  $G$  such that the von Neumann algebra  $\text{VN}(G)$  is QWEP<sup>3</sup> also satisfies this property, see [ArK1, Theorem 4.2].

Finally, we describe a new class of locally compact groups with the following property introduced<sup>4</sup> in [ArK1, Definition 1.1] which asks a bounded projection at the level  $p = \infty$  and a compatible<sup>5</sup> bounded projection at the level  $p = 1$ . This compatibility property is crucial in Section 5.2 in order to describe the decomposable norm of Fourier multipliers on noncommutative  $L^p$ -spaces associated with group von Neumann algebras.

**Definition 1.7** *A locally compact group  $G$  has property  $(\kappa)$  if there exist compatible bounded projections  $P_G^{\infty}: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  and  $P_G^1: \text{CB}(L^1(\text{VN}(G))) \rightarrow \text{CB}(L^1(\text{VN}(G)))$  on  $\mathfrak{M}^{\infty, \text{cb}}(G)$  and  $\mathfrak{M}^{1, \text{cb}}(G)$  preserving the complete positivity. In this case, we introduce the constant*

$$(1.5) \quad \kappa(G) \stackrel{\text{def}}{=} \inf \max \left\{ \|P_G^{\infty}\|_{\text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))}, \|P_G^1\|_{\text{CB}(L^1(\text{VN}(G))) \rightarrow \text{CB}(L^1(\text{VN}(G)))} \right\}$$

where the infimum is taken on all admissible couples  $(P_G^{\infty}, P_G^1)$  of projections.

3. It is not clear if this assumption is removable or not. It is required by the use of vector-valued noncommutative  $L^p$ -spaces and the most general known theory needs the QWEP assumption.

4. The notation “CB” means completely bounded. The compatibility is taken in the sense of interpolation theory [BeL] [Tri].

5. If  $P_G^{\infty}: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  is a bounded projection on  $\mathfrak{M}^{\infty, \text{cb}}(G)$  then we can define a map  $P_G^1: \text{CB}(L^1(\text{VN}(G))) \rightarrow \text{CB}(L^1(\text{VN}(G)))$  by

$$P_G^1(T) \stackrel{\text{def}}{=} (P_G^{\infty}(T^*)^*), \quad T \in \text{CB}(L^1(\text{VN}(G))).$$

It is then easy to check that  $P_G^1$  is a bounded projection preserving complete positivity onto the space  $\mathfrak{M}^{1, \text{cb}}(G)$  and its norm is equal to the one of  $P_G^{\infty}$ . It is important to note that there is no evidence that  $P_G^{\infty}$  and  $P_G^1$  are compatible in the sense of interpolation.

The average trick of Haagerup essentially implies that  $\kappa(G) = 1$  for any discrete group  $G$ . In [ArK1], it is proved that an abelian locally compact group satisfies  $\kappa(G) = 1$  and that a second countable pro-discrete locally compact group  $G$  satisfies  $\kappa(G) = 1$ . It is equally proved in [ArK1] that some class of second countable unimodular locally compact groups approximable by lattice subgroups have  $(\kappa)$ .

Our last main result is the following result, see Corollary 4.35.

**Theorem 1.8** *Let  $G$  be a second countable unimodular finite-dimensional amenable locally compact group. Then  $G$  has  $(\kappa)$ .*

In particular, we obtain that a second countable semisimple connected Lie group  $G$  or a second countable connected nilpotent Lie group  $G$  satisfies  $\kappa(G) < \infty$ . An upper estimate of  $\kappa(G)$  is possible for some groups. For example, in the case of a second countable unimodular totally disconnected amenable locally compact group, our method gives  $\kappa(G) = 1$  which is a sharp result. Note that these results complement the result of [ArK1] which does not need any *amenability* assumption. We refer to Section 4.2 for a detailed presentation of our approach.

The proof relies on the structure of finite-dimensional locally compact groups extract from the solution to Hilbert’s fifth problem. More precisely, we use essentially a version of Iwasawa’s local splitting theorem to reduce the problem to connected Lie groups and totally disconnected groups. It allows us to use doubling constants of Carnot-Carathéodory metric of connected Lie groups for small balls for construct special “noncommutative functions”. From this point of view, totally disconnected groups behave better than Lie groups, phenomenon that we already noticed in [ArK1].

In the case where the unimodular locally compact group  $G$  is amenable and with  $1 < p < \infty$  such that  $\frac{p}{p^*}$  is rational, using some other special “noncommutative functions”, we obtain a *contractive* linear map  $P_p^{(1)} : \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$  which is better than the *boundedness* of the previous case, but only for *one* value of  $p$  (see Theorem 4.33). By comparison to Theorem 1.8, this result can not be used to prove the “isometric” result” of Proposition 5.5.

**Structure of the paper** To facilitate access to the individual topics, the chapters are rendered as self-contained as possible. The paper is organized as follows.

The next section 2.1 gives background on Fourier-Stieltjes algebras, groupoids and operator algebras. In Section 2.2, we prove that for any locally compact group  $G$  there exists a well-defined injective contractive map from the Fourier-Stieltjes algebra  $B(G)$  into the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers. We also show that that the Fourier-Stieltjes algebra  $B(G)$  of a discrete group  $G$  is isometrically isomorphic to the algebra  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers on the group von Neumann algebra  $\text{VN}(G)$ . We equally describe some links between the equality  $B(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$  and property  $(\kappa_\infty)$  and a matricial variant  $(\kappa_{\infty, 2})$ . Finally, we show that the space  $\text{Dec}(A, B)$  of decomposable operators admits a canonical structure of operator space.

In Section 3.1, we give background on inner amenability and amenability. In Section 3.2, we prove different characterizations of inner amenability for unimodular locally compact groups. We will use one of these in Section 4.4.

In Section 4.1, we give background on measurable Schur multipliers and Plancherel weights. In the next section 4.2, we describe the approach of this chapter which is somewhat technical. Section 4.3 contains a construction of some Schur multipliers from completely bounded maps. In Section 4.4, we show that the symbol can be chosen to be Herz-Schur if the group is inner amenable. Section 4.5, we investigate the symbol of the Schur multipliers in the case  $p = 1$  or  $p = \infty$ . In Section 4.6, we study the convergence of the symbols of the Schur multipliers.

In Section 4.7, we prove some results in order to prove in Section 4.9 that finite-dimensional unimodular amenable locally compact groups have property  $(\kappa)$ . In Section 4.8, we construct a contractive projection from the space of completely bounded Schur multipliers  $\mathfrak{M}_G^{p,\text{cb}}$  onto the subspace  $\mathfrak{M}_G^{p,\text{cb,HS}}$  of Herz-Schur multipliers. Section 4.9, we state our main results of complementation.

In Section 5.1, we give a new characterization of amenability for unimodular locally compact groups. Section 5.2 contains a description of decomposable Fourier multipliers on noncommutative  $L^p$ -spaces in the spirit of the ones of [ArK1]. In Section 5.3, we introduce the notion of  $\text{CB}_{w^*}$ -strongly non decomposable operator and we show that this notion is connected in the case of Fourier multipliers to a famous work [DCH] of de Cannière and Haagerup. Section 5.4, we show that the notions of decomposable Fourier multiplier on a group von Neumann algebra and on a reduced  $C^*$ -algebras are identical for a discrete group.

## 2 Fourier-Stieltjes algebras and decomposable multipliers on $\text{VN}(G)$

### 2.1 Preliminaries

**Full group  $C^*$ -algebras** Let  $G$  be a locally compact group equipped with a left Haar measure  $\mu_G$ . Let  $P(G)$  (resp.  $P_1(G)$ ) denote the set of continuous, positive definite functions  $\varphi$  (resp. with  $\varphi(e) = 1$ ). Then the Gelfand-Naimark construction provides for each  $\varphi$  in  $P_1(G)$  a continuous unitary representation  $\pi_\varphi$ . The universal representation of  $G$  is  $U = \bigoplus_{\varphi \in P_1(G)} \pi_\varphi$  on the Hilbert space  $H$ . We define  $C^*(G)$  to be the norm closure in  $\mathcal{B}(H)$  of  $U(C_c(G))$  where here  $U: C_c(G) \rightarrow \mathcal{B}(H)$ ,  $f \mapsto \int_G f(s)U_s d\mu_G(s)$  denotes the integrated representation associated to  $U$  where the latter integral is understood in the weak operator sense. We identify  $C_c(G)$  as a subspace of  $C^*(G)$ . Finally, recall that there is a one-to-one correspondence between the continuous unitary representations of  $G$  and the nondegenerate representations of the  $C^*$ -algebra  $C^*(G)$ . We refer to [BeH1], [Ern1], [Mil1], [Wal1], [Wal2] and [Wal3] for more information.

**Fourier-Stieltjes algebras** Recall that the Fourier-Stieltjes algebra  $B(G)$  of a locally compact group  $G$  is the complex linear span of the set of all continuous positive-definite functions on  $G$ , see [KaL1, Def. 2.1.5]. It coincides with the set of all matrix coefficients of continuous unitary representations of  $G$ , i.e.

$$(2.1) \quad B(G) = \{ \langle \pi(\cdot)\xi, \eta \rangle_H : (\pi, H) \text{ is a unitary representation of } G \text{ and } \xi, \eta \in H \}.$$

The norm is defined by

$$(2.2) \quad \|\varphi\|_{B(G)} \stackrel{\text{def}}{=} \inf \|\xi\| \|\eta\|$$

where the infimum is taken on  $\pi, \xi, \eta$  such that  $\varphi = \langle \pi(\cdot)\xi, \eta \rangle_H$ . Equipped with pointwise multiplication  $B(G)$  becomes a commutative unital Banach algebra by [KaL1, Th. 2.1.11].

Let  $\mu_G$  be a left Haar measure of  $G$ . If  $\varphi \in B(G)$  then the linear form  $\omega_\varphi: C_c(G) \rightarrow \mathbb{C}$  defined by

$$(2.3) \quad \omega_\varphi(f) \stackrel{\text{def}}{=} \int_G \varphi(s)f(s) d\mu_G(s)$$

extends to a bounded linear form  $\omega_\varphi: C^*(G) \rightarrow \mathbb{C}$  with  $\|\varphi\|_{B(G)} = \|\omega_\varphi\|$ . It is well-known that each bounded linear form on  $C^*(G)$  satisfies this description, i.e. we have  $B(G) = C^*(G)^*$

isometrically, see [KaL1, page 40]. Moreover, essentially by [Fel1, Lemma 1.4] and [KaL1, Th. 1.6.1] the linear form  $C_c(G) \rightarrow \mathbb{C}$ ,  $f \mapsto \int_G \varphi(s)f(s) d\mu_G(s)$  extends to a positive linear form on  $C^*(G)$  if and only if  $\varphi$  is a continuous positive definite function. Finally, we denote by  $W^*(G)$  the enveloping von Neumann algebra of the  $C^*$ -algebra  $C^*(G)$ .

**Example 2.1** If the locally compact group  $G$  is *abelian*, recall that the Fourier transform  $\hat{\mu}: G \rightarrow \mathbb{C}$  of a bounded complex measure  $\mu \in M(\hat{G})$  on the dual group  $\hat{G}$  is given by  $\hat{\mu}(s) \stackrel{\text{def}}{=} \int_{\hat{G}} \chi(s) d\mu(\chi)$  where  $s \in G$ . In this case,  $B(G)$  is the space of Fourier transforms  $\varphi = \hat{\mu}$  of bounded complex measures  $\mu \in M(\hat{G})$  and  $\|\varphi\|_{B(G)} = \|\mu\|_{M(\hat{G})}$ .

**Fell's absorption principle** Let  $G$  be a locally compact group. Let  $\pi: G \rightarrow \mathcal{B}(H)$  be any continuous unitary representation of  $G$ . Recall Fell's absorption principle, e. g. [KaL1, Lemma 5.5.3]. If  $1_H: G \rightarrow \mathcal{B}(H)$ ,  $s \mapsto \text{Id}_H$  is the identity representation, we have a unitary equivalence

$$(2.4) \quad \lambda \otimes \pi \approx \lambda \otimes 1_H.$$

**Groupoids** A groupoid is a set  $G$  together with a distinguished subset  $G^{(2)} \subset G \times G$ , a multiplication map  $G^{(2)} \rightarrow G$ ,  $(s, t) \mapsto st$  and an inverse map  $G \rightarrow G$ ,  $s \mapsto s^{-1}$  such that

1. for any  $s \in G$  we have  $(s^{-1})^{-1} = s$ ,
2. If  $(s, t), (t, r) \in G^{(2)}$  then  $(st, r)$  and  $(s, tr)$  belong to  $G^{(2)}$  and  $(st)r = s(tr)$ ,
3. for any  $s \in G$  we have  $(s, s^{-1}) \in G^{(2)}$  and if  $s, r \in G$  satisfies  $(s, r) \in G^{(2)}$ , we have  $s^{-1}(sr) = r$  and  $(sr)r^{-1} = s$ .

We say that  $G^{(2)}$  is the set of composable pairs. Axiom (2) shows that for products of three groupoid elements, there is no ambiguity in dropping the parentheses (as we do in groups), and simply writing  $str$  for  $(st)r$ .

Given a groupoid  $G$  we shall write  $G^{(0)} \stackrel{\text{def}}{=} \{s^{-1}s : s \in G\}$  and refer to elements of  $G^{(0)}$  as units and to  $G^{(2)}$  itself as the unit space. Since  $(s^{-1})^{-1} = s$  for any  $s \in G$ , we also have  $G^{(0)} = \{ss^{-1} : s \in G\}$ . We define the range and domain maps  $r, d: G \rightarrow G^{(0)}$  by

$$r(s) \stackrel{\text{def}}{=} ss^{-1} \quad \text{and} \quad d(s) \stackrel{\text{def}}{=} s^{-1}s, \quad s \in G.$$

For any  $s, t \in G$  we have  $(s, t) \in G^{(2)}$  if and only if  $d(s) = r(t)$ . For any unit  $u \in G^{(0)}$ , we let  $G^u \stackrel{\text{def}}{=} r^{-1}(\{u\})$  and  $G_u \stackrel{\text{def}}{=} d^{-1}(\{u\})$ .

A locally compact groupoid is a groupoid  $G$  endowed with a locally compact topology such that the map  $s \mapsto s^{-1}$  are continuous and the map  $(s, t) \rightarrow st$  is continuous with respect to the relative topology on  $G^{(2)}$  as a subset of  $G \times G$ .

Recall, that by [Ren2, Def. 2.2] [Muh1, Def 2.28] a left Haar system for  $G$  is a family  $(\nu^u)_{u \in G^{(0)}}$  of positive Radon measures on  $G$  such that

1. the support  $\text{supp } \nu^u$  of the measure  $\nu^u$  is  $G^u$ ,
2. for any function  $f \in C_c(G)$ , the map  $G^{(0)} \rightarrow \mathbb{C}$ ,  $u \mapsto \int_G f d\nu^u$  is continuous,
3. for any function  $f \in C_c(G)$  and any  $s \in G$  we have  $\int_G f(st) d\nu^{d(s)}(t) = \int_G f(t) d\nu^{r(s)}(t)$ .

Roughly speaking, each unit  $u$  has its own measure  $\nu^u$  is  $G^u$  which lives on the little piece  $G^u$ .

In this case, the space  $C_c(G)$  of continuous functions with compact support is a  $*$ -algebra [Pat3, p. 38] equipped with the operations

$$(f * g)(s) \stackrel{\text{def}}{=} \int_G f(t)g(t^{-1}s) d\nu^{r(s)}(t), \quad f^*(s) \stackrel{\text{def}}{=} \overline{f(s^{-1})}.$$

Let  $G$  be a locally compact groupoid with Haar system  $(\nu^u)_{u \in G^{(0)}}$  and  $\mu$  be a positive Radon measure on  $G^{(0)}$ . The measure  $\nu^\mu$  induced on  $G$  by  $\mu$  [Ren2, Def. 3.1] [Pat3, p. 86] is defined by

$$\int_G f(\xi) d\nu^\mu(\xi) \stackrel{\text{def}}{=} \int_{G^{(0)}} \int_G f(\xi) d\nu^u(\xi) d\mu(u), \quad f \in C_c(G).$$

The measure  $\mu$  is said to be quasi-invariant if  $\nu^\mu$  is equivalent to its image  $\nu_\mu$  by  $s \mapsto s^{-1}$ . A measured groupoid  $(G, \nu, \mu)$  [Ren2, p. 23] is a locally compact groupoid equipped with a left Haar system  $\nu$  and a quasi-invariant measure  $\mu^6$ . The function  $\Delta \stackrel{\text{def}}{=} \frac{d\nu^\mu}{d\nu_\mu} : G \rightarrow \mathbb{R}_+^\times$  is called the modular function of  $\mu$ .

**Example 2.2** Every locally compact group  $G$  can be viewed as a locally compact groupoid, with  $G^{(0)} = \{e\}$ , multiplication given by the group operation, and inversion by the usual group inverse. We obtain a measured groupoid with a left Haar measure and the Dirac measure as a quasi-invariant measure on  $G^{(0)}$ . A groupoid  $G$  is a group if and only if its unit space  $G^{(0)}$  is a singleton.

**Example 2.3** Let  $X$  be a locally compact space. Set  $G \stackrel{\text{def}}{=} X \times X$  and  $G^{(2)} \stackrel{\text{def}}{=} \{(x, y), (y, z) : x, y, z \in X\}$ . Moreover, for any  $x, y, z \in X$  we define  $(x, y)(y, z) \stackrel{\text{def}}{=} (x, z)$  and  $(x, y)^{-1} \stackrel{\text{def}}{=} (y, x)$ . We obtain the pair groupoid (or Brandt groupoid). We have  $G^0 = \{(x, x) : x \in X\}$  which can be identified with  $X$ . Moreover, for any  $x, y \in X$  we have  $r(x, y) = x$  and  $d(x, y) = y$ . For any unit  $u \in X$ , we have  $G_u = X \times \{u\}$  and  $G^u = \{u\} \times X$ .

If we equip  $X$  with a positive Radon measure  $\mu$ , we can define for any unit  $u \in X$  the measure  $\nu^u \stackrel{\text{def}}{=} \delta_u \otimes \mu$  on  $G$  where  $\delta_u$  is the unit measure at  $u$ . In this case, the measure  $\mu$  is quasi-invariant. If we consider the discrete space  $X = \{1, \dots, n\}$  equipped with the counting measure  $\mu_n$ , we denote by  $P_n$  the associated measured groupoid.

**Example 2.4** If  $G_1$  and  $G_2$  are groupoids, it is clear that the product  $G_1 \times G_2$  has a canonical structure of groupoid with  $(G_1 \times G_2)^{(2)} = \{(x_1, y_1), (x_2, y_2) : (x_1, x_2) \in G_1^{(2)}, (y_1, y_2) \in G_2^{(2)}\}$ ,  $(x, y)^{-1} = (x^{-1}, y^{-1})$ ,  $(x_1, y_1)(x_2, y_2) = (x_1 x_2, y_1 y_2)$ . We have  $(G_1 \times G_2)^{(0)} = G_1^{(0)} \times G_2^{(0)}$ ,  $d(x, y) = (d(x), d(y))$  and  $r(x, y) = (r(x), r(y))$ .

If  $G_1 = G$  is a group and  $G_2 = P_2$ , an element of  $P_2 \times G$  can be written  $(i, j, s)$  with  $s \in G$ ,  $i, j \in \{1, 2\}$ . We can see a function  $F : P_2 \times G \rightarrow \mathbb{C}$  as a  $2 \times 2$  matrix valued function  $\begin{bmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{bmatrix}$  on the group  $G$  where  $F_{ij} : G \rightarrow \mathbb{C}$ ,  $s \mapsto F(i, j, s)$ .

We refer to [Bun1], [Hah1], [Muh1], [Pat1], [Pat2], [Pat3], [Ren1] [Ren2], [RaW1], [RaW2] and [Ren3] for more information.

---

6. Strictly speaking, only the class on the measure is important. But we do not need this point in our paper.



**von Neumann algebras and multipliers** Let  $(G, \nu, \mu)$  be a measured groupoid. The von Neumann algebra of  $(G, \lambda, \mu)$  is the von Neumann algebra generated by  $\lambda(C_c(G))$  where  $\lambda$  is the regular representation [Ren2, p. 55] [Pat3, p. 93-94] of the measured groupoid  $(G, \nu, \mu)$ .

Following [Ren1, Definition 3.1], we say that a function  $\varphi \in L^\infty(G)$  induces a bounded multiplier if pointwise multiplication by  $\varphi$  induces a weakly\* continuous<sup>7</sup> operator  $VN(G) \rightarrow VN(G)$ ,  $\lambda(f) \mapsto \lambda(\varphi f)$ .

**Positive definite functions** Let  $(G, \nu, \mu)$  be measured groupoid. By [Ren1, Prop. 1.1 and Def. 1.1], a function  $\varphi \in L^\infty(G)$  is positive definite if and only if for any positive integer  $n$  and any  $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ , the inequality

$$(2.5) \quad \sum_{k,l=1}^n \alpha_k \overline{\alpha_l} \varphi(\gamma_k^{-1} \gamma_l) \geq 0$$

holds for  $\mu$ -almost every  $u \in G^{(0)}$  and  $\nu^u$ -almost every  $\gamma_1, \dots, \gamma_n \in G^u$ . Now, we connect (with no surprise) this condition with [ArK1, Prop. 8.4] (the proof of the last result is true for a locally compact group and a trivial cocycle).

**Lemma 2.5** *Let  $G$  be a locally compact group. The matrix  $F = \begin{bmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{bmatrix} \in L^\infty(\mathbb{P}_2 \times G)$  defines a positive definite function on the groupoid  $\mathbb{P}_2 \times G$  if and only if for any integer  $n \in \mathbb{N}$ , any elements  $i_1, \dots, i_n \in \{1, 2\}$ , any  $s_1, \dots, s_n \in G$  and any complex numbers  $\alpha_1, \dots, \alpha_n$  we have*

$$(2.6) \quad \sum_{k,l=1}^n \alpha_k \overline{\alpha_l} F_{i_k i_l}(s_k^{-1} s_l) \geq 0.$$

*Proof* : If  $\gamma_k = (j', i_k, s_k)$  and  $\gamma_l = (j', j_l, s_l)$  where  $s_k, s_l \in G$ ,  $j', i_k \in \{1, 2\}$  then we have

$$\gamma_k^{-1} \gamma_l = (j', i_k, s_k)^{-1} (j', j_l, s_l) = (i_k, j', s_k^{-1})(j', j_l, s_l) = (i_k, j_l, s_k^{-1} s_l).$$

So the condition (2.5) translates to

$$(2.7) \quad \sum_{k,l=1}^n \alpha_k \overline{\alpha_l} F(i_k, j_l, s_k^{-1} s_l) \geq 0 \quad \text{i.e.} \quad \sum_{k,l=1}^n \alpha_k \overline{\alpha_l} F_{i_k, j_l}(s_k^{-1} s_l) \geq 0$$

■

**Fourier-Stieltjes algebras** In the case of groupoids, we warn the reader that there exists three notions of Fourier-Stieltjes algebra introduced in the papers [Pat2], [Ren1] and [RaW1]. See the beautiful survey [Pat1] for more information. We need a particular case of a result on Fourier-Stieltjes algebras associated with measured groupoids. A careful study of the proof of [Ren1, Prop. 1.3 and Lemma 1.1] and [Pat2, Prop. 5] gives indeed the following result.

**Proposition 2.6** *Suppose that  $G$  is a locally compact group. A function  $\varphi: G \rightarrow \mathbb{C}$  belongs to the Fourier-Stieltjes algebra  $B(G)$  if and only if there exists continuous positive definite*

7. In [Ren1, Prop 3.1], we must replace “bounded” by “weakly\* continuous” in order to obtain a correct statement.

functions  $\psi_1, \psi_2$  such that the matrix  $\begin{bmatrix} \psi_1 & \varphi \\ \overline{\varphi} & \psi_2 \end{bmatrix}$  defines a continuous positive definite function on the groupoid  $P_2 \times G$ . In this case, we have

$$(2.8) \quad \|\varphi\|_{\mathbb{B}(G)} = \inf \|\psi_1\|_{\infty} \|\psi_2\|_{\infty}$$

where the infimum is taken over all  $\psi_1$  and  $\psi_2$  satisfying the previous condition.

**Decomposable maps** The following is folklore. See [Arh9, Lemma 2.2] for a proof.

**Lemma 2.7** *Let  $M$  be a von Neumann algebra acting on a Hilbert space  $H$ . Then  $M$  equipped with the canonical locally convex structure which gives the  $\sigma$ -strong\* topology is quasi-complete.*

We will use the following elementary lemma.

**Lemma 2.8** *Let  $M_1, M_2$  and  $N$  be von Neumann algebras and  $T: M_1 \rightarrow M_2$  be a weakly\* continuous decomposable map. Then we have a well-defined weakly\* continuous decomposable map  $\text{Id}_N \otimes T: N \overline{\otimes} M_1 \rightarrow N \overline{\otimes} M_2$  and*

$$(2.9) \quad \|\text{Id}_N \otimes T\|_{\text{dec}, N \overline{\otimes} M_1 \rightarrow N \overline{\otimes} M_2} \leq \|T\|_{\text{dec}, M_1 \rightarrow M_2}.$$

*Proof* : Note that  $T$  is completely bounded by [Haa, Proposition 1.3 3)]. By [DCH, Lemma 1.5 b)], we infer that the map  $\text{Id}_N \otimes T: N \overline{\otimes} M_1 \rightarrow N \overline{\otimes} M_2$  is a well-defined weakly\* continuous completely bounded map. By [Haa, Remark 1.5], there exist some maps  $v_1, v_2: M_1 \rightarrow M_2$  such that the map  $\begin{bmatrix} v_1 & T \\ T^\circ & v_2 \end{bmatrix}: M_2(M_1) \rightarrow M_2(M_2)$  is completely positive with  $\max\{\|v_1\|, \|v_2\|\} = \|T\|_{\text{dec}, M_1 \rightarrow M_2}$ . It is not difficult to see that we can suppose that  $v_1$  and  $v_2$  are weakly\* continuous by using [ArK1, Proposition 3.1]. Then by [Bin, proposition 4.3.7] the map

$$\begin{bmatrix} \text{Id}_N \otimes v_1 & \text{Id}_N \otimes T \\ \text{Id}_N \otimes T^\circ & \text{Id}_N \otimes v_2 \end{bmatrix} = \text{Id}_{M_2(N)} \otimes \begin{bmatrix} v_1 & T \\ T^\circ & v_2 \end{bmatrix}: M_2(N \overline{\otimes} M_1) \rightarrow M_2(N \overline{\otimes} M_2)$$

is also completely positive. We deduce that the map  $\text{Id}_N \otimes T: N \overline{\otimes} M_1 \rightarrow N \overline{\otimes} M_2$  is decomposable with  $\|\text{Id}_N \otimes T\|_{\text{dec}} \leq \max\{\|\text{Id}_N \otimes v_1\|, \|\text{Id}_N \otimes v_2\|\} \leq \max\{\|v_1\|_{\text{cb}}, \|v_2\|_{\text{cb}}\} = \max\{\|v_1\|, \|v_2\|\} = \|T\|_{\text{dec}, M_1 \rightarrow M_2}$  where we use in the first equality the complete positivity of  $v_1$  and  $v_2$  combined with [Pau, Prop. 3.6].  $\blacksquare$

We will use the following well-known result.

**Proposition 2.9** *Let  $M$  and  $N$  be two von Neumann algebras equipped with normal semifinite faithful traces. Suppose  $1 \leq p \leq \infty$ . Let  $T: L^p(M) \rightarrow L^p(N)$  be a completely positive map. Then  $T$  is decomposable and*

$$(2.10) \quad \|T\|_{\text{dec}, L^p(M) \rightarrow L^p(N)} \leq \|T\|_{L^p(M) \rightarrow L^p(N)}.$$

Let  $M_1, M_2$  and  $M_3$  be von Neumann algebras equipped with normal semifinite faithful traces. Suppose  $1 \leq p \leq \infty$ . Let  $T_1: L^p(M_1) \rightarrow L^p(M_2)$  and  $T_2: L^p(M_2) \rightarrow L^p(M_3)$  be some decomposable maps. It is easy to see that the composition  $T_2 \circ T_1$  is decomposable and that

$$(2.11) \quad \|T_2 \circ T_1\|_{\text{dec}} \leq \|T_2\|_{\text{dec}} \|T_1\|_{\text{dec}}.$$

## 2.2 Fourier-Stieltjes algebras and decomposable multipliers on $\text{VN}(G)$

Let  $G$  be a locally compact group. Recall that by [KaL1, Corollary 5.4.11] we have a contractive inclusion  $\text{B}(G) \subset \mathfrak{M}^{\infty, \text{cb}}(G)$ . In the next result, we improve this result by replacing the space  $\mathfrak{M}^{\infty, \text{cb}}(G)$  of completely bounded Fourier multipliers by the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers. The inclusion  $\text{B}(G) \subset \mathfrak{M}^{\infty, \text{dec}}(G)$  is elementary<sup>8</sup> but the *contractivity* of the inclusion  $\text{B}(G) \subset \mathfrak{M}^{\infty, \text{dec}}(G)$  is new even in the discrete case.

**Proposition 2.10** *Let  $G$  be a locally compact group. The map  $\text{B}(G) \rightarrow \mathfrak{M}^{\infty, \text{dec}}(G)$ ,  $\varphi \mapsto M_\varphi$  is a well-defined injective contraction from the Fourier-Stieltjes algebra  $\text{B}(G)$  in the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable Fourier multipliers.*

*Proof* : We will give two very different proofs. We start with a “purely group” argument. Let  $\varphi \in \text{B}(G)$ . By homogeneity, we can suppose that  $\|\varphi\|_{\text{B}(G)} = 1$ . We will use the associated linear form  $\omega_\varphi: C^*(G) \rightarrow \mathbb{C}$ ,  $\int_G f(s)U_s d\mu_G(s) \mapsto \int_G \varphi(s)f(s) d\mu_G(s)$  of (2.3). By [BLM, Lemma A.2.2], we can consider the unique weakly\* continuous extension  $\tilde{\omega}_\varphi: W^*(G) \rightarrow \mathbb{C}$  on the von Neumann algebra  $W^*(G)$  where we use here (and only in this step) the identification  $W^*(G) = C^*(G)^{**}$  of [Dix1, page 265]. Note that for any  $s \in G$  the element  $U_s$  belong to the von Neumann algebra  $W^*(G)$ . We will prove that

$$(2.12) \quad \tilde{\omega}_\varphi(U(s)) = \varphi(s), \quad s \in G.$$

Let  $s \in G$  and let  $\mathfrak{B}$  be a neighbourhood basis at  $s$  constituted of compact neighbourhoods. For any  $V \in \mathfrak{B}$ , consider a positive continuous function  $f_V: G \rightarrow \mathbb{R}^+$  on  $G$  with  $\int_G f_V d\mu_G = 1$  with support contained in  $V$ . Then by [Bou3, VIII.17 Corollary 3] the net  $(\int_G f_V(t)U_t d\mu_G(t))$  converges to  $U_s$  for the strong operator topology, hence for the weak operator topology. Moreover, by [Bou3, VIII.15] for any  $V$  we have the estimate

$$\left\| \int_G f_V(t)U_t d\mu_G(t) \right\| \leq \int_G f_V d\mu_G = 1.$$

So the net  $(\int_G f_V(t)U_t d\mu_G(t))_V$  is bounded. We deduce that the net  $(\int_G f_V(t)U_t d\mu_G(t))_V$  converges to  $U_s$  for the weak\* topology. On the one hand, we infer that the net  $(\tilde{\omega}_\varphi(\int_G f_V(t)U_t d\mu_G(t)))_V$  converges to  $\tilde{\omega}_\varphi(U_s)$ . On the other hand, using [Bou3, VIII.15] and the continuity of  $\varphi$  we have

$$\tilde{\omega}_\varphi\left(\int_G f_V(t)U_t d\mu_G(t)\right) = \omega_\varphi\left(\int_G f_V(t)U_t d\mu_G(t)\right) = \int_G \varphi(t)f_V(t) d\mu_G(t) \xrightarrow{V} \varphi(s).$$

By uniqueness of the limit, we conclude that (2.12) is true.

By Fell’s absorption principle (2.4) applied to the universal representation  $U: G \rightarrow \mathcal{B}(H)$  instead of  $\pi$ , there exists a unitary  $W: L^2(G, H) \rightarrow L^2(G, H)$  such that for any  $s \in G$

$$W(\lambda_s \otimes \text{Id}_H)W^* = \lambda_s \otimes U_s.$$

By [Dix2, p. 9 and 25], we deduce that there exists a normal \*-homomorphism  $\Delta: \text{VN}(G) \rightarrow \text{VN}(G) \overline{\otimes} W^*(G)$ ,  $\lambda_s \mapsto \lambda_s \otimes U_s$ . By [Bou2, Corollary 2, III page 38] and Lemma 2.7, note that

8. Indeed, if  $\varphi \in \text{B}(G)$  then we can write  $\varphi = \varphi_1 - \varphi_2 + i\varphi_3 - \varphi_4$  where each  $\varphi_i$  is a continuous positive definite function. By [KaL1, Proposition 5.4.9], each Fourier multiplier  $M_{\varphi_i}: \text{VN}(G) \rightarrow \text{VN}(G)$  is completely positive. Then it is immediate that the Fourier multiplier

$$M_\varphi = M_{\varphi_1 - \varphi_2 + i(\varphi_3 - \varphi_4)} = M_{\varphi_1} - M_{\varphi_2} + i(M_{\varphi_3} - M_{\varphi_4})$$

is decomposable.

for any  $f \in C_c(G)$ , the integral  $\int_G f(s)\lambda_s d\mu_G(s)$  is a well-defined weak\* integral. Using the weak\* continuity of  $\Delta$  together with [Bou3, VI.3 Proposition 1] in the first equality, we deduce that

$$(2.13) \quad \Delta\left(\int_G f(s)\lambda_s d\mu_G(s)\right) = \int_G f(s)\Delta(\lambda_s) d\mu_G(s) = \int_G f(s)(\lambda_s \otimes U_s) d\mu_G(s).$$

Now, for any  $s \in G$ , we obtain again with [Bou3, VI.3 Proposition 1] that

$$\begin{aligned} (\text{Id} \otimes \tilde{\omega}_\varphi) \circ \Delta\left(\int_G f(s)\lambda_s d\mu_G(s)\right) &\stackrel{(2.13)}{=} (\text{Id} \otimes \tilde{\omega}_\varphi)\left(\int_G f(s)(\lambda_s \otimes U_s) d\mu_G(s)\right) \\ &= \left(\int_G f(s)(\text{Id} \otimes \tilde{\omega}_\varphi)(\lambda_s \otimes U_s) d\mu_G(s)\right) \stackrel{(2.12)}{=} \int_G \varphi(s)f(s)\lambda_s d\mu_G(s). \end{aligned}$$

We conclude that the weakly\* continuous map  $(\text{Id} \otimes \tilde{\omega}_\varphi) \circ \Delta$  is the Fourier multiplier  $M_\varphi$  of symbol  $\varphi$ .

Note that by Proposition 2.9, the \*-homomorphism  $\Delta$  is decomposable. According to [ER, Lemma 5.4.3], the linear form  $\tilde{\omega}_\varphi$  is equally decomposable with  $\|\tilde{\omega}_\varphi\|_{\text{dec}} = \|\tilde{\omega}_\varphi\| = 1$ . By Lemma 2.8, we deduce that we have a well-defined weakly\* continuous map  $\text{Id} \otimes \tilde{\omega}_\varphi: \text{VN}(G) \overline{\otimes} \mathbb{W}^*(G) \rightarrow \text{VN}(G)$  which is also decomposable. We conclude by composition that the map  $M_\varphi = (\text{Id} \otimes \tilde{\omega}_\varphi) \circ \Delta: \text{VN}(G) \rightarrow \text{VN}(G)$  is decomposable and that

$$\|M_\varphi\|_{\text{dec}} = \|(\text{Id} \otimes \tilde{\omega}_\varphi) \circ \Delta\|_{\text{dec}} \stackrel{(2.11)}{\leq} \|\text{Id} \otimes \tilde{\omega}_\varphi\|_{\text{dec}} \|\Delta\|_{\text{dec}} \stackrel{(2.9)}{\leq} \|\tilde{\omega}_\varphi\|_{\text{dec}} \|\Delta\|_{\text{dec}} = \|\Delta\|_{\text{dec}} \stackrel{(2.10)}{\leq} 1.$$

It is well-known that the map  $\text{B}(G) \rightarrow \mathfrak{M}^\infty(G)$ ,  $\psi \mapsto M_\psi$  is injective. So the map  $\text{B}(G) \rightarrow \mathfrak{M}^{\infty, \text{dec}}(G)$ ,  $\varphi \mapsto M_\varphi$  is also injective.

Now, we give a second proof using groupoids. Let  $\varphi \in \text{B}(G)$  and  $\varepsilon > 0$ . By Proposition 2.6, there exists continuous positive definite functions  $\psi_1, \psi_2$  (hence bounded by [BHV, Proposition C.4.2]) such that the matrix  $\begin{bmatrix} \psi_1 & \varphi \\ \frac{\varphi}{\psi_1} & \psi_2 \end{bmatrix}$  defines a continuous positive definite function  $F$  on the groupoid  $\text{P}_2 \times G$  with

$$(2.14) \quad \|\psi_1\|_\infty \|\psi_2\|_\infty \leq \|\varphi\|_{\text{B}(G)} + \varepsilon.$$

Note that  $\text{VN}(\text{P}_2 \times G) = \text{VN}(\text{P}_2) \overline{\otimes} \text{VN}(G) = \text{M}_2 \overline{\otimes} \text{VN}(G) = \text{M}_2(\text{VN}(G))$ . By generalizing the very transparent argument of [BrO, Prop. 5.6.16] with [Arh10, Prop. 4.12 and Remark 4.12],  $F$  induces a completely positive multiplier on the von Neumann algebra  $\text{VN}(\text{P}_2 \times G)$ . This completely positive multiplier identifies to the map  $\begin{bmatrix} M_{\psi_1} & M_\varphi \\ M_{\frac{\varphi}{\psi_1}} & M_{\psi_2} \end{bmatrix} = \begin{bmatrix} M_{\psi_1} & M_\varphi \\ M_\varphi^\circ & M_{\psi_2} \end{bmatrix}: \text{M}_2(\text{VN}(G)) \rightarrow \text{M}_2(\text{VN}(G))$ . Note that  $M_{\psi_1}$  and  $M_{\psi_2}$  are completely positive. We conclude that the Fourier multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  is decomposable with

$$\begin{aligned} \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)} &\leq \|M_{\psi_1}\|_{\text{VN}(G) \rightarrow \text{VN}(G)} \|M_{\psi_2}\|_{\text{VN}(G) \rightarrow \text{VN}(G)} \\ &\stackrel{(2.14)}{=} \|\psi_1\|_\infty \|\psi_2\|_\infty \leq \|\varphi\|_{\text{B}(G)} + \varepsilon. \end{aligned}$$

■

Now, we study the converse of Proposition 2.10 in Proposition 2.14 and in Theorem 2.19. We need the two following results. The next observation is written without proof in [Wall, page 212] and in [Eym, page 188]. For the sake of completeness, we give a proof.

**Proposition 2.11** *Let  $G$  be a locally compact group. Let  $\varphi: G \rightarrow \mathbb{C}$  be a continuous positive definite function. We have*

$$(2.15) \quad \|\varphi\|_{\mathbb{B}(G)} = \varphi(e).$$

*Proof :* *First proof if  $G$  is discrete.* Let  $G$  be a discrete group. By [Dix1, 13.9.2], the full  $C^*$ -algebra  $C^*(G)$  of  $G$  is unital<sup>9</sup>. By (2.3), we have a positive linear form  $\omega_\varphi: C^*(G) \rightarrow \mathbb{C}$ ,  $U(s) \mapsto \varphi(s)$ . So using [KaR1, Theorem 4.3.2] in the second equality, we conclude that  $\|\varphi\|_{\mathbb{B}(G)} = \|\omega_\varphi\|_{C^*(G)^*} = \omega_\varphi(1) = \omega_\varphi(U(e)) = \varphi(e)$ .

*Second proof if  $G$  is locally compact.* Using [KaL1, Remark 2.1.10 p. 43] we have  $\|\varphi\|_{\mathbb{B}(G)} \geq \|\varphi\|_\infty \geq \varphi(e)$ . Furthermore with [KaL1, Prop. 1.4.17], we can write  $\varphi = \langle \pi(\cdot)\xi, \xi \rangle_H$  where  $\pi$  is a continuous unitary representation on  $H$  and  $\xi \in H$ . We deduce that  $\varphi(e) = \langle \xi, \xi \rangle_H = \|\xi\|^2 \stackrel{(2.2)}{\geq} \|\varphi\|_{\mathbb{B}(G)}$ . ■

Now, we give a description of the norm of the Fourier-Stieltjes algebra  $\mathbb{B}(G)$  for some suitable functions.

**Proposition 2.12** *Let  $G$  be a locally compact group. Let  $\varphi \in \mathbb{B}(G)$  such that  $\check{\varphi} = \overline{\varphi}$ . We have*

$$(2.16) \quad \|\varphi\|_{\mathbb{B}(G)} = \inf \{ \varphi_1(e) + \varphi_2(e) : \varphi = \varphi_1 - \varphi_2, \varphi_1, \varphi_2 \in \mathbb{P}(G) \}.$$

*Proof :* Suppose that  $\varphi = \varphi_1 - \varphi_2$  with  $\varphi_1, \varphi_2 \in \mathbb{P}(G)$ . We have

$$\|\varphi\|_{\mathbb{B}(G)} = \|\varphi_1 - \varphi_2\|_{\mathbb{B}(G)} \leq \|\varphi_1\|_{\mathbb{B}(G)} + \|\varphi_2\|_{\mathbb{B}(G)} \stackrel{(2.15)}{=} \varphi_1(e) + \varphi_2(e).$$

Passing to the infimum, we obtain that  $\|\varphi\|_{\mathbb{B}(G)} \leq \inf \{ \varphi_1(e) + \varphi_2(e) : \varphi = \varphi_1 - \varphi_2, \varphi_1, \varphi_2 \in \mathbb{P}(G) \}$ . Indeed, by [Eym, (2.7)] (or [KaL1, page 41]<sup>10</sup>) we have an equality in this last inequality and the infimum is a minimum. ■

**Remark 2.13** Suppose that the locally compact  $G$  is *abelian*. For any real bounded measure  $\mu$  on the dual group  $\hat{G}$ , the previous result combined with Example 2.1 and (2.15) says that

$$(2.17) \quad \|\mu\|_{\mathbb{M}(\hat{G})} = \inf \{ \|\mu_1\|_{\mathbb{M}(\hat{G})} + \|\mu_2\|_{\mathbb{M}(\hat{G})} : \mu = \mu_1 - \mu_2, \mu_1, \mu_2 \geq 0 \}.$$

We can replace the group  $\hat{G}$  by a locally compact space  $X$ . Indeed, for any *real* bounded measure  $\mu$ , we can decompose  $\mu$  by [Bou2, III §1. 8 Cor. 2] as  $\mu = \mu^+ - \mu^-$  for some bounded positive measures  $\mu_+$  and  $\mu_-$  on  $X$  with  $\|\mu\|_{\mathbb{M}(X)} = \|\mu^+\|_{\mathbb{M}(X)} + \|\mu^-\|_{\mathbb{M}(X)}$  and we can use a similar reasoning.

If we add property  $(\kappa_\infty)$  on the group  $G$ , we obtain a partial converse to Proposition 2.10.

**Proposition 2.14** *Let  $G$  be a locally compact group. If  $G$  has  $(\kappa_\infty)$ , then the map  $\mathbb{B}(G) \rightarrow \mathfrak{M}^{\infty, \text{dec}}(G)$ ,  $\varphi \mapsto M_\varphi$  is a bijection from  $\mathbb{B}(G)$  onto the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable multipliers. Moreover, if  $\kappa_\infty(G) = 1$  and if  $\varphi$  belongs to  $\mathbb{B}(G)$  and satisfies  $\check{\varphi} = \overline{\varphi}$ , we have  $\|\varphi\|_{\mathbb{B}(G)} = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}$ .*

9. Actually, by [Mil1] the full  $C^*$ -algebra  $C^*(G)$  of a locally compact group  $G$  is unital if and only if  $G$  is discrete.  
10. Note that in this reference, the assumption “ $\check{u} = \overline{u}$ ” is missing.

*Proof* : In Proposition 2.10, we have seen that we have a contractive inclusion  $B(G) \subset \mathfrak{M}^{\infty, \text{dec}}(G)$ . First, we show the reverse inclusion. Suppose that  $M_\varphi : \text{VN}(G) \rightarrow \text{VN}(G)$  is a decomposable Fourier multiplier (hence weakly\* continuous) with symbol  $\varphi$ . We can write  $M_\varphi = T_1 + T_2 + i(T_3 - T_4)$  for some completely positive maps  $T_1, T_2, T_3, T_4 : \text{VN}(G) \rightarrow \text{VN}(G)$ . By using the contractive projection  $P_{w^*} : B(\text{VN}(G)) \rightarrow B(\text{VN}(G))$  of [ArK1, Proposition 3.1] which preserves the complete positivity, we can suppose that these maps  $T_1, T_2, T_3, T_4$  are weakly\* continuous since  $P_{w^*}(M_\varphi) = M_\varphi$ . Using the bounded projection  $P_G^\infty : \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  provided by  $\kappa_\infty(G)$ , we obtain

$$M_\varphi = P_G^\infty(M_\varphi) = P_G^\infty(T_1 - T_2 + i(T_3 - T_4)) = P_G^\infty(T_1) - P_G^\infty(T_2) + i(P_G^\infty(T_3) - P_G^\infty(T_4))$$

where each  $P_G^\infty(T_i) : \text{VN}(G) \rightarrow \text{VN}(G)$  is a completely positive Fourier multiplier with symbol  $\varphi_i$ , i.e.  $P_G^\infty(T_i) = M_{\varphi_i}$ . By [DCH, p. 458 and Prop. 4.2], the function  $\varphi_i$  is continuous and positive definite. We deduce that

$$M_\varphi = M_{\varphi_1} - M_{\varphi_2} + i(M_{\varphi_3} - M_{\varphi_4}) = M_{\varphi_1 - \varphi_2 + i(\varphi_3 - \varphi_4)}.$$

We infer that  $\varphi = \varphi_1 - \varphi_2 + i\varphi_3 - \varphi_4$ . So  $\varphi$  belongs to the Fourier-Stieltjes algebra  $B(G)$ .

Suppose that  $\varphi$  belongs to  $B(G)$  and satisfies  $\check{\varphi} = \overline{\varphi}$ . This last condition means that  $M_\varphi : \text{VN}(G) \rightarrow \text{VN}(G)$  is adjoint preserving. Let  $\varepsilon > 0$ . By [Haa, Proposition 1.3 (1)] and [Haa, p. 184], there exists some completely positive operators  $T_1, T_2 : \text{VN}(G) \rightarrow \text{VN}(G)$  such that  $M_\varphi = T_1 - T_2$  with  $\|T_1 + T_2\| = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}$ . By using the contractive projection  $P_{w^*} : B(\text{VN}(G)) \rightarrow B(\text{VN}(G))$  of [ArK1, Proposition 3.1] which preserves the complete positivity, it is easy to check that we can suppose that the maps  $T_1$  and  $T_2$  are weakly\* continuous since  $P_{w^*}(M_\varphi) = M_\varphi$ . Using  $\kappa_\infty(G)$ , we deduce that

$$(2.18) \quad M_\varphi = P_G^\infty(M_\varphi) = P_G^\infty(T_1 - T_2) = P_G^\infty(T_1) - P_G^\infty(T_2).$$

We denote by  $\varphi_1$  and  $\varphi_2$  the symbols of the completely positive Fourier multipliers  $P_G^\infty(T_1)$  and  $P_G^\infty(T_2)$ . These functions are continuous and definite positive again by [DCH, p. 458 and Prop. 4.2]. The equality (2.18) gives  $\varphi = \varphi_1 - \varphi_2$ . By using [DCH, Proposition 4.3] in the second equality, we see that

$$\begin{aligned} \|\varphi\|_{B(G)} &\stackrel{(2.16)}{\leq} \varphi_1(e) + \varphi_2(e) = (\varphi_1 + \varphi_2)(e) = \|M_{\varphi_1 + \varphi_2}\|_{\text{VN}(G) \rightarrow \text{VN}(G)} \\ &= \|M_{\varphi_1} + M_{\varphi_2}\|_{\text{VN}(G) \rightarrow \text{VN}(G)} = \|P_G^\infty(T_1) + P_G^\infty(T_2)\| = \|P_G^\infty(T_1 + T_2)\| \\ &\leq \|T_1 + T_2\| = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}. \end{aligned}$$

Combining with Proposition 2.10, we conclude that  $\|\varphi\|_{B(G)} = \|M_\varphi\|_{\text{dec}}$ . ■

Now, we show that the first inclusion of (1.4) can be strict.

**Theorem 2.15** *Let  $G$  be a non-amenable locally compact group such that the von Neumann algebra  $\text{VN}(G)$  is injective. Then there exists a decomposable Fourier multiplier  $T : \text{VN}(G) \rightarrow \text{VN}(G)$  which is not induced by an element  $\varphi \in B(G)$ .*

*Proof* : Since the von Neumann algebra  $\text{VN}(G)$  is injective, we have by [Haa, Theorem 1.6] the equality  $\mathfrak{M}^{\infty, \text{cb}}(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$  isometrically. Since  $G$  is not amenable, we deduce by an unpublished result of Ruan [Pis6, p. 54] [Spr1, p. 190] that  $B(G) \subsetneq \mathfrak{M}^{\infty, \text{cb}}(G)$ . We conclude that  $B(G) \subsetneq \mathfrak{M}^{\infty, \text{dec}}(G)$ . ■

**Example 2.16** By [Con2, Corollary 7], the von Neumann algebra  $\text{VN}(G)$  of a second countable connected locally compact group  $G$  is injective. Hence this result applies for example to the locally compact group  $\text{SL}_2(\mathbb{R})$  which is non-amenable by [BHV, Example G.2.4 (i)]. We conclude that  $\text{B}(\text{SL}_2(\mathbb{R})) \not\subseteq \mathfrak{M}^{\infty, \text{dec}}(\text{SL}_2(\mathbb{R}))$ .

Suppose  $1 \leq p \leq \infty$ . We denote by  $\mathfrak{M}_2^{p, \text{cb}}(G)$  the space of (normal if  $p = \infty$ ) completely bounded operators  $\Phi: L^p(\text{M}_2(\text{VN}(G))) \rightarrow L^p(\text{M}_2(\text{VN}(G)))$  such that  $\Phi = [M_{a_{ij}}]_{1 \leq i, j \leq 2}$  for some multipliers  $M_{a_{ij}}$  in  $\mathfrak{M}^{p, \text{cb}}(G)$ . See [ArK1, Section 4.1] for a particular case.

**Definition 2.17** Let  $G$  be a locally compact group with  $(\kappa_\infty)$ . We say that  $G$  has property  $(\kappa_{\infty, 2})$  if

$$P_{2,G}^\infty = \text{Id} \otimes P_G^\infty: \text{M}_2(\text{CB}_{w^*}(\text{VN}(G))) \mapsto \text{M}_2(\text{CB}_{w^*}(\text{VN}(G)))$$

onto  $\mathfrak{M}_2^{\infty, \text{cb}}(G)$  preserves the complete positivity. We write

$$\kappa_{\infty, 2}(G) = \begin{cases} \kappa_\infty(G) & : G \text{ has } (\kappa_{\infty, 2}) \\ \infty & : G \text{ does not have } (\kappa_{\infty, 2}). \end{cases}$$

**Remark 2.18** The result [ArK1, Theorem 4.2] says in particular  $\kappa_{\infty, 2}(G) = 1$  for any discrete group  $G$ .

Replacing  $\kappa_\infty(G)$  by  $\kappa_{\infty, 2}(G) = 1$ , we obtain an *isometric* statement for all functions, which is an improvement of Proposition 2.14.

**Theorem 2.19** Let  $G$  be a locally compact group with  $\kappa_{\infty, 2}(G) = 1$ . The map  $\text{B}(G) \rightarrow \mathfrak{M}^{\infty, \text{dec}}(G)$ ,  $\varphi \mapsto M_\varphi$  is an isometric bijection from  $\text{B}(G)$  onto the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable multipliers.

*Proof* : In Proposition 2.10, we have seen that we have a contractive inclusion  $\text{B}(G) \subset \mathfrak{M}^{\infty, \text{dec}}(G)$ . It suffices to show the reverse inclusion. Suppose that the multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  is decomposable. By [Haa, Remark 1.5], there exist some linear maps  $v_1, v_2: \text{VN}(G) \rightarrow \text{VN}(G)$  such that the linear map  $\begin{bmatrix} v_1 & M_\varphi \\ M_{\bar{\varphi}} & v_2 \end{bmatrix}: \text{M}_2(\text{VN}(G)) \rightarrow \text{M}_2(\text{VN}(G))$  is completely positive with  $\max\{\|v_1\|, \|v_2\|\} = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}$ . We can suppose that  $v_1$  and  $v_2$  are in addition weakly\* continuous by using [ArK1, Proposition 3.1]. Now, we can apply the amplification projection  $\text{Id}_{\text{M}_2} \otimes P_G^\infty: \text{M}_2(\text{CB}_{w^*}(\text{VN}(G))) \rightarrow \text{M}_2(\text{CB}_{w^*}(\text{VN}(G)))$  of Definition 2.17. We obtain that

$$(2.19) \quad \begin{bmatrix} P_G^\infty(v_1) & M_\varphi \\ M_{\bar{\varphi}} & P_G^\infty(v_2) \end{bmatrix} = \begin{bmatrix} P_G^\infty(v_1) & P_G^\infty(M_\varphi) \\ P_G^\infty(M_{\bar{\varphi}}) & P_G^\infty(v_2) \end{bmatrix} = (\text{Id}_{\text{M}_2} \otimes P_G^\infty) \left( \begin{bmatrix} v_1 & M_\varphi \\ M_{\bar{\varphi}} & v_2 \end{bmatrix} \right)$$

is completely positive. Moreover, we have

$$(2.20) \quad \max\{\|P_G^\infty(v_1)\|, \|P_G^\infty(v_2)\|\} \leq \|P_G^\infty\| \max\{\|v_1\|, \|v_2\|\} = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}.$$

We can write  $\begin{bmatrix} P_G^\infty(v_1) & M_\varphi \\ M_{\bar{\varphi}} & P_G^\infty(v_2) \end{bmatrix}$  as  $\begin{bmatrix} M_{\psi_1} & M_\varphi \\ M_{\bar{\varphi}} & M_{\psi_2} \end{bmatrix}$  for some continuous positive definite functions  $\psi_1, \psi_2: G \rightarrow \mathbb{C}$ . By [ArK1, Prop. 8.4], the condition (2.6) is satisfied with  $\begin{bmatrix} \psi_1 & \varphi \\ \varphi^\circ & \psi_2 \end{bmatrix}$  instead of  $\begin{bmatrix} F_{11} & F_{12} \\ F_{21} & F_{22} \end{bmatrix}$ . By Lemma 2.5, we conclude that  $F = \begin{bmatrix} \psi_1 & \varphi \\ \varphi^\circ & \psi_2 \end{bmatrix}$  identifies to a definite positive function on the groupoid  $\text{P}_2 \times G$ . By Proposition 2.6, we obtain that  $\varphi$  belongs

to the Fourier-Stieltjes algebra  $B(G)$ . Moreover, using the well-known contractive inclusion  $\mathfrak{M}^\infty(G) \subset L^\infty(G)$  in the second inequality we infer that

$$\|\varphi\|_{B(G)} \stackrel{(2.8)}{\leq} \|\psi_1\|_\infty \|\psi_2\|_\infty \leq \max \{ \|P_G^\infty(v_1)\|, \|P_G^\infty(v_2)\| \} \stackrel{(2.20)}{\leq} \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}.$$

■

In particular, using [ArK1, Th. 4.2] we obtain the following result.

**Corollary 2.20** *Let  $G$  be a discrete group. The map  $B(G) \rightarrow \mathfrak{M}^{\infty, \text{dec}}(G)$ ,  $\varphi \mapsto M_\varphi$  is an isometric isomorphism from the Fourier-Stieltjes  $B(G)$  onto the algebra  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of decomposable multipliers.*

On the one hand, if  $G$  is a locally compact group, we can equip the Fourier-Stieltjes algebra  $B(G)$  with the dual operator space structure induced by the equality  $C^*(G)^* = B(G)$ , which is also by [Spr1, Prop. 4.2] (proved in [Spr2, Prop 2.3.2]) the operator space structure induced on  $B(G)$  when we consider  $B(G)$  as a subspace of  $\text{CB}(C^*(G))$ .

On the other hand, if  $A$  and  $B$  are  $C^*$ -algebras such that  $B$  is unital we will observe that we can equip the space  $\text{Dec}(A, B)$  of decomposable maps with an operator space structure. To see how this is done suppose that  $[T_{ij}]$  belongs to  $M_{m,n}(\text{Dec}(A, B))$ . We identify the matrix  $[T_{ij}]$  with the map  $A \mapsto M_{m,n}(B)$ ,  $x \mapsto [T_{ij}(x)]$ . We define a norm on  $M_{m,n}(\text{Dec}(A, B))$  by setting

$$(2.21) \quad \|[T_{ij}]\|_{M_{m,n}(\text{Dec}(A, B))} \stackrel{\text{def}}{=} \|x \mapsto [T_{ij}(x)]\|_{\text{Dec}(A, M_{m,n}(B))}.$$

In short, we make the identification  $M_{m,n}(\text{Dec}(A, B)) = \text{Dec}(A, M_{m,n}(B))$ .

**Proposition 2.21** *Let  $A$  and  $B$  be  $C^*$ -algebras such that  $B$  is unital. Equipped with the matricial norms (2.21),  $\text{Dec}(A, B)$  is an operator space.*

*Proof* : Let  $X, Y \in M_n$  and  $[T_{ij}] \in M_n(\text{Dec}(A, B))$ . Note that by [Pis7, Exercise 12.1] the map  $u: M_n(B) \rightarrow M_n(B)$ ,  $y \mapsto (X \otimes 1_B)y(Y \otimes 1_B)$  is decomposable with  $\|u\|_{\text{dec}} \leq \|X \otimes 1_B\|_{M_n(B)} \|Y \otimes 1_B\|_{M_n(B)}$ . Using this observation in the first equality, we have

$$\begin{aligned} \|X[T_{ij}]Y\|_{M_n(\text{Dec}(A, B))} &\stackrel{(2.21)}{=} \|x \mapsto X[T_{ij}(x)]Y\|_{\text{Dec}(A, M_n(B))} \\ &= \|x \mapsto (X \otimes 1_B)[T_{ij}(x)](Y \otimes 1_B)\|_{\text{Dec}(A, M_n(B))} \\ &\leq \|X \otimes 1_B\|_{M_n(B)} \|Y \otimes 1_B\|_{M_n(B)} \|x \mapsto [T_{ij}(x)]\|_{\text{Dec}(A, M_n(B))} \\ &\stackrel{(2.21)}{=} \|X\|_{M_n} \|[T_{ij}]\|_{M_n(\text{Dec}(A, B))} \|Y\|_{M_n}. \end{aligned}$$

Let  $[T_{ij}] \in M_n(\text{Dec}(A, B))$  and  $[S_{kl}] \in M_m(\text{Dec}(A, B))$ . Using [Pis17, Lemma 6.8] in the second equality, we have

$$\begin{aligned} \|[T_{ij}] \oplus [S_{kl}]\|_{M_{n+m}(\text{Dec}(A, B))} &\stackrel{(2.21)}{=} \|x \mapsto [T_{ij}(x)] \oplus [S_{kl}(x)]\|_{\text{Dec}(A, M_{n+m}(B))} \\ &= \max \left\{ \|x \mapsto [T_{ij}(x)]\|_{\text{Dec}(A, M_n(B))}, \|x \mapsto [S_{kl}(x)]\|_{\text{Dec}(A, M_m(B))} \right\} \\ &\stackrel{(2.21)}{=} \max \left\{ \|[T_{ij}]\|_{M_n(\text{Dec}(A, B))} + \|[S_{kl}]\|_{M_m(\text{Dec}(A, B))} \right\}. \end{aligned}$$

Now, it suffices to use Ruan's theorem [Pis7, page 35].

■



Then  $\mathfrak{M}^{\infty, \text{dec}}(G)$  can be equipped with the operator space structure induced by the inclusion  $\mathfrak{M}^{\infty, \text{dec}}(G) \subset \text{Dec}(\text{VN}(G))$ . It is likely that we can prove that the isometry of Corollary 2.20 is a complete isometry. Details are left to the interested reader.

Finally, we prove the second part of Conjecture 1.3 in the discrete case. This result improves [ArK1, Proposition 3.32 (1)] which says that the second inclusion of (1.4) is strict for any non-amenable weakly amenable discrete group  $G$ .

The following result can be seen as a new characterization of amenability for discrete groups.

**Theorem 2.22** *Let  $G$  be a discrete group. The von Neumann algebra  $\text{VN}(G)$  is injective if and only if we have  $\mathfrak{M}^{\infty, \text{dec}}(G) = \mathfrak{M}^{\infty, \text{cb}}(G)$ .*

*Proof* : By Corollary 2.20, we have an isometry  $\mathfrak{M}^{\infty, \text{dec}}(G) = \text{B}(G)$ . It suffices to use the result [Pis6, p. 54] which says that  $\text{B}(G) = \mathfrak{M}^{\infty, \text{cb}}(G)$  if and only if  $G$  is amenable. For a discrete group  $G$ , the last point is equivalent to the injectivity of the von Neumann algebra  $\text{VN}(G)$  by [SS, Theorem 3.8.2] (or Theorem 3.2). ■

## 3 Inner amenability

### 3.1 Preliminaries on inner amenability and amenability

We warn the reader that different notions of inner amenability coexist in the literature, see [Pat, page 84] for more information. We say that a locally compact group  $G$  is inner amenable if there exists a state  $m$  on  $L^\infty(G)$  such that

$$(3.1) \quad m(\text{int}_s f) = m(f)$$

for any  $s \in G$  where

$$(3.2) \quad (\text{int}_s f)(t) \stackrel{\text{def}}{=} f(s^{-1}ts), \quad s, t \in G.$$

**Example 3.1** Every amenable group is obviously amenable. Recall that a locally compact group  $G$  is said to have an invariant neighborhood (IN) [Pal2, page 1273] if there exists a compact neighbourhood  $V$  of the identity  $e$  in  $G$  such  $V$  is stable under all inner automorphisms of  $G$ . By [Pal2, Proposition 12.1.9], an IN-group  $G$  is unimodular. Note that if  $\mu_G$  is a Haar measure on an IN-group  $G$  it is clear using [Bou3, (31) and (33) VII.13] that the map  $m: L^\infty(G) \rightarrow \mathbb{C}$ ,  $f \mapsto \frac{1}{\mu_G(V)} \int_V f$  satisfies (3.1). Hence an IN-group is inner amenable<sup>11</sup>. By [ArK1, Proposition 6.36]<sup>12</sup>, pro-discrete locally compact groups are IN. Moreover, by [Pal2, Proposition 12.1.9], compact groups and locally compact abelian groups and discrete groups are IN. These groups are therefore all inner amenable. So in general, inner amenability is much weaker than amenability.

Recall the following characterization of amenability of [LaP2, Corollary 3.2] and [Pat, page 85] which we will use<sup>13</sup>.

**Theorem 3.2** *Let  $G$  be a second countable locally compact group. The following are equivalent:*

- 
- 11. This observation generalizes [FSW, Remark 2.6 (i)].
  - 12. A locally compact group  $G$  is pro-discrete if and only if it admits a basis  $(X_j)$  of neighborhoods of the identity  $e$  consisting of open compact normal subgroups.
  - 13. We added the assumption “second countable” since this equivalence uses the equivalence between “injective” and property  $P$  for von Neumann algebras with separable preduals.

1. the group von Neumann algebra  $\text{VN}(G)$  is injective and  $G$  is inner amenable.
2.  $G$  is amenable.

**Example 3.3** Recall that a topological group  $G$  is of type I [BeH1, Def. 6.D.1 and Prop. 7.C.I] if for any continuous unitary representation  $\pi$  of  $G$ , the von Neumann algebra  $\pi(G)''$  is of type I, hence injective by [Str, Prop. 10.23]. In particular, by Theorem 3.2 a second countable locally compact group  $G$  of type I is inner amenable if and only if it is amenable. We refer to [BeH1, Th 6.E.19] for a list of groups of type I.

In particular, the group  $\text{SL}_2(\mathbb{Q}_p)$  is a non-inner amenable totally disconnected locally compact group since it is non-amenable by [BHV, Example G.3.6 (iii)].

**Example 3.4** If a locally compact group  $G$  is almost connected<sup>14</sup> then its von Neumann algebra  $\text{VN}(G)$  is injective by [Pat4, page 228]. Again, by Theorem 3.2 such a group is inner amenable if and only if it is amenable. This result in the connected case was first proved by Losert and Rindler in [LoR1, Theorem 1] and proven again in [LaP2, Corollary 3.4].

If  $A$  and  $B$  are two subsets of a set  $E$ , the notation  $A\Delta B \stackrel{\text{def}}{=} (A - B) \cup (B - A)$  denotes here the symmetric difference of  $A$  and  $B$ . Recall that

$$(3.3) \quad |1_A - 1_B| = 1_{A\Delta B}.$$

We will use the following reformulation of [CPPR, Lemma 8.6 p. 43] which is actually and essentially a classical trick in amenability theory [EiW1, page 364-365] [BHV, p. 441-442]. We give the two lines of calculus for the sake of completeness.

**Lemma 3.5** *Let  $G$  be a locally compact group equipped with a left Haar measure  $\mu$ . Let  $\varepsilon > 0$  and consider some positive functions  $f, g_1, \dots, g_n$  of  $L^1(G)$  satisfying  $\sum_{k=1}^n \|f - g_k\|_{L^1(G)} < \varepsilon$  and  $\|f\|_{L^1(G)} = 1$ . Then there exists  $t > 0$  such that*

$$(3.4) \quad \sum_{k=1}^n \mu(\{f > t\} \Delta \{g_k > t\}) < \varepsilon \mu(\{f > t\}).$$

*Proof* : For any  $s \in G$  and  $1 \leq k \leq n$ , we have by [BHV, Lemma G.5.2] and (3.3) the equalities

$$(3.5) \quad \|f\|_{L^1(G)} = \int_0^\infty \mu(\{f > t\}) dt \quad \text{and} \quad \|f - g_k\|_{L^1(G)} = \int_0^\infty \mu(\{f > t\} \Delta \{g_k > t\}) dt.$$

We deduce that

$$\begin{aligned} \int_0^\infty \sum_{k=1}^n \mu(\{f > t\} \Delta \{g_k > t\}) dt &= \sum_{k=1}^n \int_0^\infty \mu(\{f > t\} \Delta \{g_k > t\}) dt \\ &\stackrel{(3.5)}{=} \sum_{k=1}^n \|f - g_k\|_{L^1(G)} < \varepsilon \|f\|_{L^1(G)} \stackrel{(3.5)}{=} \varepsilon \int_0^\infty \mu(\{f > t\}) dt. \end{aligned}$$

The conclusion is obvious. ■

---

14. That means that  $G/G_e$  is compact where  $G_e$  is the connected component of the identity  $e$ .

### 3.2 Some characterizations of inner amenability

Now, we introduce the following definition which is an “inner”-variant of a well-known definition in amenability theory.

**Definition 3.6** *Let  $G$  be a locally compact group. Then  $G$  is called inner Følner (in short  $G \in (\text{IF})$ ) if for every finite subset  $F$  of  $G$  there exists a net  $(V_j^F)_j$  of measurable subsets of  $G$  such that  $\mu(V_j^F) \in (0, \infty)$ , with the property that for all  $s \in F$ ,*

$$(3.6) \quad \frac{\mu(V_j^F \Delta (s^{-1}V_j^F s))}{\mu(V_j^F)} \xrightarrow{j \rightarrow \infty} 0.$$

Now, we give different characterizations of inner amenability for unimodular locally compact groups. The equivalence between the first and the second point is sketched in [LoR1, Prop 1]. For the sake of completeness, we give a complete proof.

**Theorem 3.7** *Let  $G$  be a unimodular locally compact group. The following are equivalent.*

1.  $G$  is inner amenable.
2. There exists an asymptotically central net  $(f_j)$  of functions of  $L^1(G)$ , i.e. for any  $s \in G$ , we have

$$(3.7) \quad \frac{\|f_j - \text{int}_s f_j\|_{L^1(G)}}{\|f_j\|_{L^1(G)}} \xrightarrow{j} 0.$$

3.  $G$  is inner Følner.
4. There exists a net  $(f_j)$  of positive functions of  $L^1(G)$  with  $\int_G f_j d\mu = 1$  such that for all  $s \in G$ , we have  $\|f_j - \text{int}_s f_j\|_1 \xrightarrow{j} 0$ .
5. The same property as before, but the  $f_j$ 's belong in addition to  $C_c(G)$  and are positive definite.
6.  $G$  is inner Følner and in addition the sets  $V_j$  can be chosen to be symmetric, open and containing  $e$ .

Moreover, if  $G$  is second countable, then the nets  $(f_j)$  can be chosen to be sequences. Finally, the net  $(V_j^F)_j$  in the previous definition of inner Følner can be chosen to be a sequence.

*Proof* : 1.  $\implies$  4.: Let  $m \in L^\infty(G)^*$  be an inner invariant mean. By [Pie, Proposition 3.3] (see also [EiW1, Lemma 10.16]), we can approximate  $m$  in the weak\* topology by a net  $(f_j)$  in  $L^1(G)$  with  $f_j \geq 0$   $\|f_j\|_{L^1(G)} = 1$ . For any  $s \in G$  and any  $g \in L^\infty(G)$ , we have

$$\langle \text{int}_s(f_j), g \rangle_{L^1(G), L^\infty(G)} = \langle f_j, \text{int}_{s^{-1}}(g) \rangle_{L^1(G), L^\infty(G)} \xrightarrow{j \rightarrow \infty} \langle m, \text{int}_{s^{-1}}(g) \rangle = \langle m, g \rangle.$$

and  $\langle f_j, g \rangle_{L^1(G), L^\infty(G)} \rightarrow \langle m, g \rangle_{L^\infty(G)^*, L^\infty(G)}$ . With a  $\frac{\varepsilon}{2}$ -argument, it follows that for any  $s \in G$  we have  $\lim_j (\text{int}_s f_j - f_j) = 0$ .

Since for convex sets the weak closure coincides with the norm closure [Meg1, Theorem 2.5.16], we can replace  $f_j$  by some convex combinations to get  $\lim_j \|\text{int}_s(f_j) - f_j\|_{L^1(G)} = 0$ .

This replacement can be seen in the following way. Let  $F \stackrel{\text{def}}{=} \{s_1, \dots, s_n\}$  be a finite set

of  $G$ . According to the above,  $(0, \dots, 0)$  belongs to the weak-closure of the convex hull of  $\{(\text{int}_{s_1}(f_j) - f_j, \text{int}_{s_2}(f_j) - f_j, \dots, \text{int}_{s_n}(f_j) - f_j) : j\}$ , hence to the  $L^1(G)^n$  norm closure of this convex hull. Thus there exists a sequence  $(g_k)_k$  in this convex hull converging to  $(0, \dots, 0)$  in norm. For any  $k \in \mathbb{N}$ , we can write

$$\begin{aligned} g_k &= \sum_{\ell=1}^L \lambda_\ell (\text{int}_{s_1}(f_{j_\ell}) - f_{j_\ell}, \dots, \text{int}_{s_n}(f_{j_\ell}) - f_{j_\ell}) \\ &= \left( \text{int}_{s_1} \left( \sum_{\ell=1}^L \lambda_\ell f_{j_\ell} \right) - \sum_{\ell=1}^L \lambda_\ell f_{j_\ell}, \dots, \text{int}_{s_n} \left( \sum_{\ell=1}^L \lambda_\ell f_{j_\ell} \right) - \sum_{\ell=1}^L \lambda_\ell f_{j_\ell} \right) \\ &= (\text{int}_{s_1}(h_k) - h_k, \dots, \text{int}_{s_n}(h_k) - h_k), \end{aligned}$$

where  $\lambda_\ell \geq 0$ ,  $\sum_{\ell=1}^L \lambda_\ell = 1$  and where  $h_k \stackrel{\text{def}}{=} \sum_{\ell=1}^L \lambda_\ell f_{j_\ell}$  still is a positive normalized element of  $L^1(G)$ . We can suppose that  $\|\text{int}_s(h_k) - h_k\|_1 \leq \frac{1}{k}$  for any  $s \in F$ . Now write  $h_k = h_{k,F}$ , let  $F$  vary in the set of finite subsets of  $G$  directed by inclusion, so that  $(h_{k,F})_{k,F}$  becomes a net in  $L^1(G)$  such that  $\|\text{int}_s(h_{k,F}) - h_{k,F}\|_1 \rightarrow 0$  as  $(k, F) \rightarrow \infty$  for any  $s \in G$ .

4.  $\implies$  1.: Note that we have an isometric inclusion  $L^1(G) \subset L^\infty(G)^*$ . Consider a cluster point  $m \in L^\infty(G)^*$  of this net for the weak\* topology which is positive and satisfies clearly  $m(1) = 1$ . For any  $f \in L^\infty(G)$  and any  $s \in G$ , we have

$$\begin{aligned} \left| \langle f_j, f \rangle_{L^1(G), L^\infty(G)} - \langle f_j, \text{int}_s f \rangle_{L^1(G), L^\infty(G)} \right| &= |\langle f_j, f \rangle - \langle \text{int}_{s^{-1}} f_j, f \rangle| = |\langle f_j - \text{int}_{s^{-1}} f_j, f \rangle| \\ &\leq \|f_j - \text{int}_{s^{-1}} f_j\|_{L^1(G)} \|f\|_{L^\infty(G)} \xrightarrow{j} 0. \end{aligned}$$

With a  $\frac{\varepsilon}{3}$ -argument we conclude that (3.1) is satisfied.

3.  $\implies$  2.: Let  $F$  be a finite subset of  $G$ . According to the assumption, there exists a subset  $V_j^F$  such that  $\frac{\mu(V_j^F \Delta (s^{-1} V_j^F s))}{\mu(V_j^F)} \leq \frac{1}{j}$  for any  $s \in F$ . Putting  $f_j^F \stackrel{\text{def}}{=} 1_{V_j^F}$  and using (3.3), we obtain from Definition 3.6 that  $\|f_j^F - \text{int}_s f_j^F\|_1 / \|f_j^F\|_1 \leq \frac{1}{j}$  for any  $s \in F$ . Directing the subsets  $F$  by inclusion and the  $j$  in the usual manner, we obtain a net of positive functions  $f_j^F$  of  $L^1(G)$  as in (3.7).

2.  $\implies$  4.: Using a normalization, it suffices to see that by the elementary inequality  $\int_G |f| - |g| \leq \int_G |f - g|$ , the  $f_j$ 's in (3.7) can be chosen positive.

4.  $\implies$  5.: Let  $(f_j)_j$  be the net as in 4. For  $\varepsilon > 0$ , choose some  $f_{j,\varepsilon} \in C_c(G)$  such that  $f_{j,\varepsilon} \geq 0$ , and  $\|f_{j,\varepsilon} - f_j\|_1 < \varepsilon$ . Since the map  $\text{int}_s : L^1(G) \rightarrow L^1(G)$  is isometric, for any  $s \in G$  we have

$$\|f_{j,\varepsilon} - \text{int}_s f_{j,\varepsilon}\|_1 \leq \|f_{j,\varepsilon} - f_j\|_1 + \|f_j - \text{int}_s f_j\|_1 + \|\text{int}_s f_j - \text{int}_s f_{j,\varepsilon}\|_1 \leq \varepsilon + \|f_j - \text{int}_s f_j\|_1 + \varepsilon.$$

We can suppose  $\int_G f_{j,\varepsilon} d\mu = 1$ . Replacing the index  $j$  by  $(j, \varepsilon)$  and equipping it with the suitable order, we obtain a net  $(f_{j,\varepsilon})_{j,\varepsilon}$  of positive normalized continuous functions with compact support with the convergence property from 3. We may and do thus assume now that the net  $(f_j)_j$  in the third point consists of continuous functions with compact support.

For any  $j$ , put now  $g_j \stackrel{\text{def}}{=} f_j * \check{f}_j$  where  $\check{f}_j(s) \stackrel{\text{def}}{=} f_j(s^{-1})$ . For any  $s \in G$  we have

$$(3.8) \quad g_j(s) = (f_j * \check{f}_j)(s) \stackrel{(4.5)}{=} \int_G f_j(st^{-1}) \check{f}_j(t) d\mu(t) = \int_G f_j(sr) f_j(r) d\mu(r) \geq 0.$$

Then for any  $j$

$$\begin{aligned}\|g_j\|_1 &= \int_G g_j(s) \, d\mu(s) \stackrel{(3.8)}{=} \int_G \int_G f_j(sr) f_j(r) \, d\mu(r) \, d\mu(s) = \int_G \left( \int_G f_j(sr) \, d\mu(s) \right) f_j(r) \, d\mu(r) \\ &= \int_G \left( \int_G f_j(u) \, d\mu(u) \right) f_j(r) \, d\mu(r) = \|f_j\|_1^2 = 1.\end{aligned}$$

Moreover, for any  $t \in G$ , we have

$$(3.9) \quad (\text{int}_t g_j)(s) \stackrel{(3.2)}{=} g_j(t^{-1}st) \stackrel{(3.8)}{=} \int_G f_j(t^{-1}str) f_j(r) \, d\mu(r) = \int_G f_j(t^{-1}srt) f_j(t^{-1}rt) \, d\mu(r).$$

Thus we obtain

$$\begin{aligned}\|g_j - \text{int}_t g_j\|_1 &= \int_G |g_j(s) - (\text{int}_t g_j)(s)| \, d\mu(s) \\ &\stackrel{(3.8)(3.9)}{=} \int_G \left| \int_G f_j(sr) f_j(r) - f_j(t^{-1}srt) f_j(t^{-1}rt) \, d\mu(r) \right| \, d\mu(s) \\ &\leq \int_G \left| \int_G f_j(sr) [f_j(r) - f_j(t^{-1}rt)] \, d\mu(r) \right| + \left| \int_G [f_j(sr) - f_j(t^{-1}srt)] f_j(t^{-1}rt) \, d\mu(r) \right| \, d\mu(s) \\ &\leq \int_{G \times G} f_j(sr) |f_j(r) - f_j(t^{-1}rt)| \, d\mu(r) \, d\mu(s) + \int_{G \times G} |f_j(sr) - f_j(t^{-1}srt)| f_j(t^{-1}rt) \, d\mu(r) \, d\mu(s) \\ &\leq \|f_j\|_1 \|f_j - \text{int}_t f_j\|_1 + \|f_j - \text{int}_t f_j\|_1 \|\text{int}_t f_j\|_1 = 2 \|f_j - \text{int}_t f_j\|_1.\end{aligned}$$

This shows that the  $g_j$ 's have the same normalisation and convergence property than the  $f_j$ 's. Moreover, by [HeR2, page 281] the  $g_j$ 's are continuous positive definite functions with compact support.

5.  $\implies$  6.: Let  $F = \{s_1, \dots, s_n\}$  be a finite subset of  $G$  and  $\varepsilon > 0$ . According to the fifth point and (3.7), choose some positive definite functions  $f_j \in C_c(G)$  such that  $\|f_j - \text{int}_s f_j\|_1 / \|f_j\| < \varepsilon / \text{card } F$  for all  $s \in F$ . Using Lemma 3.5 with  $n = \text{card } F$ ,  $f = f_j$  and  $g_k = \text{int}_{s_k^{-1}} f_j$  and the subset  $V \stackrel{\text{def}}{=} \{f_j > t\}$  of  $G$ , we deduce that for some suitable  $t > 0$  and  $s \in F$ ,

$$\sum_{s \in F} \mu(V \Delta (s^{-1}Vs)) = \sum_{s \in F} \mu(\{f_j > t\} \Delta \{\text{int}_{s^{-1}} f_j > t\}) \stackrel{(3.4)}{<} \varepsilon \mu(\{f_j > t\}) = \varepsilon \mu(V).$$

Therefore,  $G$  is inner Følner. Moreover, since  $f_j$  is continuous,  $V$  is an open subset of  $G$ . Furthermore, since the function  $f_j$  is positive definite, we have  $\|f_j\|_\infty = f_j(e_G)$  by [KaL1, page 23]. We deduce that  $e_G$  belongs to  $V$  since otherwise we would have  $V = \emptyset$  and the previous strict inequality could not hold. Finally, [KaL1, Proposition 1.4.16 (ii)], we have  $f_j = \check{f}_j$  since  $f_j \geq 0$ . We conclude that  $V$  is symmetric.

6.  $\implies$  3.: trivial.

The proofs assertions concerning second countable groups uses standard tricks and are left to the reader.  $\blacksquare$

**Remark 3.8** By [CrT1, Proposition 3.2], a locally compact group  $G$  is inner amenable if and only if there exists a state  $m$  on  $\text{VN}(G)$  such that

$$(3.10) \quad m(\lambda_s^* x \lambda_s) = m(x), \quad s \in G, x \in \text{VN}(G).$$

## 4 Projections on the space of completely bounded Fourier multipliers

### 4.1 Preliminaries

**Hilbert-Schmidt operators** Let  $\Omega$  be a  $\sigma$ -finite measure space. We will use the space  $S_\Omega^\infty \stackrel{\text{def}}{=} S^\infty(L^2(\Omega))$  of compact operators, its dual  $S_\Omega^1$  and the space  $B(L^2(\Omega))$  of bounded operators on the Hilbert space  $L^2(\Omega)$ . If  $f \in L^2(\Omega \times \Omega)$ , we denote the associated Hilbert-Schmidt operator by

$$(4.1) \quad \begin{array}{ccc} K_f: & L^2(\Omega) & \longrightarrow & L^2(\Omega) \\ & \xi & \longmapsto & \int_X f(\cdot, y)\xi(y) \, dy \end{array} .$$

Using the notation  $\check{f}(x, y) \stackrel{\text{def}}{=} f(y, x)$ , we have  $(K_f)^* = K_{\check{f}}$ . Since  $L^2(\Omega \times \Omega) \rightarrow S_\Omega^2$ ,  $f \mapsto K_f$  is an isometry from the Hilbert space  $L^2(\Omega \times \Omega)$  onto the Hilbert space  $S_\Omega^2$  of Hilbert-Schmidt operators on  $L^2(\Omega)$ , that means that

$$(4.2) \quad \text{Tr}(K_f K_g) = \int_{\Omega \times \Omega} f \check{g}, \quad f, g \in L^2(\Omega \times \Omega).$$

**Measurable Schur multipliers** Let  $\Omega$  be a  $\sigma$ -finite measure space. We say that  $\varphi \in L^\infty(\Omega \times \Omega)$  induces a measurable Schur multiplier on  $B(L^2(\Omega))$  if the map  $S_\Omega^2 \mapsto S_\Omega^2$ ,  $K_f \mapsto K_{\varphi f}$  is bounded. In this case, the operator  $S_\Omega^2 \mapsto B(L^2(\Omega))$ ,  $K_f \mapsto K_{\varphi f}$  admits by [BLM, Lemma A.2.2] a unique weakly\* continuous extension  $M_\varphi: B(L^2(\Omega)) \rightarrow B(L^2(\Omega))$  called the measurable Schur multiplier associated with  $\varphi$ . We say that  $\varphi$  is the symbol of the operator  $M_\varphi$ . See [Duq22, Lemma 3.1].

**Example 4.1** If  $X = \{1, \dots, n\}$  is equipped with the counting measure, we can identify  $B(L^2(X))$  with  $M_n$ . Then each  $K_f$  identifies to the matrix  $[f(i, j)]$ , the Schur multiplier is the map  $M_\varphi: M_n \rightarrow M_n$ ,  $[f(i, j)] \mapsto [\varphi(i, j)f(i, j)]$ .

Let  $X$  be a  $\sigma$ -finite measure space. The map  $S_X^2 \rightarrow S_X^2$ ,  $K_f \mapsto K_{\check{f}}$  extends to an involutive normal \*-antiautomorphism  $R: B(L^2(X)) \rightarrow B(L^2(X))$ .

We introduce the following duality bracket

$$(4.3) \quad \langle z, y \rangle_{B(L^2(X)), S_X^1} \stackrel{\text{def}}{=} \text{Tr}(R(z)y), \quad z \in B(L^2(X)), y \in S_X^1$$

which is more suitable than the bracket  $\langle z, y \rangle = \text{Tr}(zy)$ .

$$(4.4) \quad \langle M_\varphi(z), y \rangle_{B(L^2(X)), S_X^1} = \langle z, M_\varphi(y) \rangle_{B(L^2(X)), S_X^1} .$$

**$S_\Omega^p$ -Schur multipliers** Suppose  $1 < p < \infty$ . We say that a measurable function  $\varphi: \Omega \times \Omega \rightarrow \mathbb{C}$  induces a bounded Schur multiplier on  $S_\Omega^p$  if for any  $f \in L^2(\Omega \times \Omega)$  satisfying  $K_f \in S_\Omega^p$  we have  $K_{\varphi f} \in S_\Omega^p$  and if the map  $S_\Omega^2 \cap S_\Omega^p \rightarrow S_\Omega^p$ ,  $K_f \mapsto K_{\varphi f}$  extends to a bounded map  $M_\varphi$  from  $S_\Omega^p$  into  $S_\Omega^p$  called the Schur multiplier associated with  $\varphi$ . We denote by  $\mathfrak{M}_\Omega^{p, \text{cb}}$  the space of completely bounded Schur multipliers on  $S_\Omega^p$ . We denote by  $\mathfrak{M}_G^{p, \text{cb}, \text{HS}}$  the subspace of  $\mathfrak{M}_G^{p, \text{cb}}$  of completely bounded Herz-Schur multipliers.

**Convolution** If  $G$  is a *unimodular* locally compact group equipped with a Haar measure  $\mu_G$ , recall that the convolution product of two functions  $f$  and  $g$  is given, when it exists, by

$$(4.5) \quad (f * g)(s) \stackrel{\text{def}}{=} \int_G f(r)g(r^{-1}s) d\mu_G(r) = \int_G f(sr^{-1})g(r) d\mu_G(r).$$

**Plancherel weights** Let  $G$  be a locally compact group. A function  $g \in L^2(G)$  is called left bounded [Haa2, Definition 2.1] if the convolution operator  $\lambda(g)$  induces a bounded operator on  $L^2(G)$ . The Plancherel weight  $\tau_G: \text{VN}(G)^+ \rightarrow [0, \infty]$  is<sup>15</sup> defined by the formula

$$\tau_G(x) = \begin{cases} \|g\|_{L^2(G)}^2 & \text{if } x^{\frac{1}{2}} = \lambda(g) \text{ for some left bounded function } g \in L^2(G) \\ +\infty & \text{otherwise} \end{cases}.$$

By [Haa2, Proposition 2.9] (see also [Ped, Theorem 7.2.7]), the canonical left ideal  $\mathfrak{n}_{\tau_G} = \{x \in \text{VN}(G) : \tau_G(x^*x) < \infty\}$  is given by

$$\mathfrak{n}_{\tau_G} = \{\lambda(g) : g \in L^2(G) \text{ is left bounded}\}.$$

Recall that  $\mathfrak{m}_{\tau_G}^+$  denotes the set  $\{x \in \text{VN}(G)^+ : \tau_G(x) < \infty\}$  and that  $\mathfrak{m}_{\tau_G}$  is the complex linear span of  $\mathfrak{m}_{\tau_G}^+$  which is a two-sided ideal of  $\text{VN}(G)$ . By [Haa2, Proposition 2.9] and [Str, Proposition page 280], we have

$$\mathfrak{m}_{\tau_G}^+ = \{\lambda(g) : g \in L^2(G) \text{ continuous and left bounded, } \lambda(g) \geq 0\}.$$

By [Haa2, page 125] or [Ped, Proposition 7.2.8], the Plancherel weight  $\tau_G$  on  $\text{VN}(G)$  is tracial if and only if  $G$  is unimodular, which means that the left Haar measure of  $G$  and the right Haar measure of  $G$  coincide. Now, in the sequel, we suppose that the locally compact group  $G$  is unimodular.

We will use the involution  $f^*(t) = \overline{f(t^{-1})}$ . By [Kum, Theorem 4], if  $f, g \in L^2(G)$  are left bounded then  $f * g$  and  $f^*$  are left bounded and we have

$$(4.6) \quad \lambda(f)\lambda(g) = \lambda(f * g) \quad \text{and} \quad \lambda(f)^* = \lambda(f^*).$$

If  $f, g \in L^2(G)$  it is well-known [Bou3, VIII p. 39] that the function  $f * g$  is continuous and that we have  $(f * g)(e_G) = (g * f)(e_G) = \int_G \check{g}f d\mu_G$  where  $e_G$  denotes the identity element of  $G$  and where  $\check{g}(s) \stackrel{\text{def}}{=} g(s^{-1})$ . By [StZ, (4) page 282], if  $f, g \in L^2(G)$  are left bounded, the operator  $\lambda(g)^*\lambda(f)$  belongs to  $\mathfrak{m}_{\tau_G}$  and we have the fundamental “noncommutative Plancherel formula”

$$(4.7) \quad \tau_G(\lambda(g)^*\lambda(f)) = \langle g, f \rangle_{L^2(G)} \quad \text{which gives} \quad \tau_G(\lambda(g)\lambda(f)) = \int_G \check{g}f d\mu_G = (g * f)(e_G).$$

In particular, this formula can be used with any functions of  $L^1(G) \cap L^2(G)$ .

If we consider the subset  $C_e(G) = \text{span}\{g^* * f : g, f \in L^2(G) \text{ left bounded}\}$  of  $C(G)$ , we have

$$(4.8) \quad \mathfrak{m}_{\tau_G} = \lambda(C_e(G))$$

and we can see  $\tau_G$  as the functional that evaluates functions of  $C_e(G)$  at  $e_G \in G$ . Although the formula  $\tau_G(\lambda(h)) = h(e)$  seems to make sense for every function  $h$  in  $C_c(G)$ , we warn the reader that it is not true in general that  $\lambda(C_c(G)) \subset \mathfrak{m}_{\tau_G}$  contrary to what is unfortunately too often written in the literature.

15. This is the natural weight associated with the left Hilbert algebra  $C_c(G)$ .

**Operator theory** Let  $T: L^2(M) \rightarrow L^2(M)$  be any bounded operator. We will denote by  $T^*$  the adjoint of  $T$  defined by

$$\tau(T(x)y) = \tau(xT^*(y)), \quad x, y \in L^2(M).$$

For any  $1 \leq p \leq \infty$  and any  $T: L^p(M) \rightarrow L^p(M)$ , we can consider the map  $T^\circ: L^p(M) \rightarrow L^p(M)$  defined by

$$(4.9) \quad T^\circ(x) \stackrel{\text{def}}{=} T(x^*)^*, \quad x \in L^p(M).$$

If  $p = 2$  and if we denote by  $T^\dagger$  the adjoint of  $T$  in the usual sense of Hilbertian operator theory, that is

$$\tau(T(x)y^*) = \tau(x(T^\dagger(y))^*), \quad x, y \in L^2(M),$$

we see that

$$(4.10) \quad T^\dagger = T^{*\circ}.$$

In particular, the selfadjointness of  $T: L^2(M) \rightarrow L^2(M)$  means that  $T^* = T^\circ$ .

## 4.2 Description of the approach

Suppose  $1 \leq p \leq \infty$  and let  $G$  be a locally compact group. In this section, we present an approach for obtaining some bounded projections  $P_G^p: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(L^p(\text{VN}(G)))$  from the space  $\text{CB}(L^p(\text{VN}(G)))$  of completely bounded operators on  $L^p(\text{VN}(G))$  onto the subspace  $\mathfrak{M}^{p,\text{cb}}(G)$  of completely bounded Fourier multipliers on  $L^p(\text{VN}(G))$ , beyond the case of discrete groups, for a suitable locally compact group. The methods are different from the ones of [ArK1] and complement the results of this paper. If  $G$  is a locally compact group, we will use the fundamental unitary  $W: L^2(G \times G) \rightarrow L^2(G \times G)$  of  $\text{B}(L^2(G) \overline{\otimes} \text{VN}(G))$  and its inverse  $W^{-1}$  defined in [Vae1, Example 2.2.10 p. 26] (see also [Kus05, Remark 5.16 p. 150]) by

$$(4.11) \quad (W\xi)(s, t) \stackrel{\text{def}}{=} \xi(s, s^{-1}t), \quad (W^{-1}\xi)(s, t) = \xi(s, st), \quad s, t \in G, \xi \in L^2(G \times G).$$

Before going into the details, let us shortly present the roadmap of the proof of results of Section 4.9.

Suppose that the group  $G$  is discrete and recall the well-known construction. Let  $\Delta: \text{VN}(G) \rightarrow \text{VN}(G) \overline{\otimes} \text{VN}(G)$ ,  $\lambda_s \mapsto \lambda_s \otimes \lambda_s$  be the coproduct. This *trace preserving* normal unital injective  $*$ -homomorphism extends to a completely positive isometric map  $\Delta_p: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G) \overline{\otimes} \text{VN}(G))$  for any  $1 \leq p \leq \infty$ . With the adjoint  $(\Delta_p^*)^*: L^p(\text{VN}(G) \overline{\otimes} \text{VN}(G)) \rightarrow L^p(\text{VN}(G))$ , the map  $P_G^p: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(\text{VN}(G))$  defined by

$$(4.12) \quad P_G^p(T) = (\Delta_p^*)^*(\text{Id}_{L^p(\text{VN}(G))} \otimes T)\Delta_p, \quad T \in \text{CB}(L^p(\text{VN}(G)))$$

is a contractive projection from the Banach space  $\text{CB}(L^p(\text{VN}(G)))$  onto the subspace  $\mathfrak{M}^{p,\text{cb}}(G)$  of completely bounded Fourier multipliers acting on  $L^p(\text{VN}(G))$  which preserves the complete positivity (in the case  $p = \infty$  replace  $\text{CB}(L^p(\text{VN}(G)))$  by the space  $\text{CB}_{w^*}(\text{VN}(G))$ ).

By [Vae1, page 26] and [Str1, p. 267], we can factorize<sup>16</sup> the coproduct as

$$\Delta(x) = W(x \otimes 1)W^{-1}, \quad x \in \text{VN}(G).$$

---

16. Indeed, this factorization is the definition of the coproduct.



If  $u, v \in \text{VN}(G)$ , we can therefore rewrite the formula (4.12) as

$$\langle P_G(T)u, v \rangle_{L^p(\text{VN}(G), L^{p^*}(\text{VN}(G)))} = \langle (\text{Id} \otimes T)\Delta_p(u), \Delta_{p^*}(v) \rangle_{L^p, L^{p^*}}$$

and finally

$$(4.13) \quad \langle P_G(T)u, v \rangle = \langle (\text{Id} \otimes T)(W(u \otimes 1)W^{-1}), W(v \otimes 1)W^{-1} \rangle_{L^p, L^{p^*}}.$$

Now, if  $G$  is a (second countable unimodular) locally compact group and if  $T: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))$  is again a completely bounded map, we wish replace one or both units 1 of the formula (4.13) by suitable sequences  $(x_j)$  and  $(y_j)$  of elements which approximate 1 in some sense. Actually, we start by replacing in (4.13) the elements 1 by elements  $x, y \in L^1(\text{VN}(G)) \cap \text{VN}(G)$  and  $u, v$  by elements of  $u \in S_G^p$  and  $v \in S_G^{p^*}$ . We will show that there exists a completely bounded Schur multiplier  $M_{x,y,T}: S_G^p \rightarrow S_G^p$  (replace the Schatten class  $S_G^p$  by the von Neumann algebra  $B(L^2(G))$  if  $p = \infty$ ) such that

$$(4.14) \quad \langle M_{x,y,T}(u), v \rangle_{S_G^p, S_G^{p^*}} = \langle (\text{Id} \otimes T)(W(u \otimes x)W^{-1}), W(v \otimes y)W^{-1} \rangle_{S_G^p(L^p(\text{VN}(G))), S_G^{p^*}(L^{p^*}(\text{VN}(G)))}$$

for any suitable elements  $u \in S_G^p$  and  $v \in S_G^{p^*}$ . Note that  $x \in L^p(\text{VN}(G))$ ,  $y \in L^{p^*}(\text{VN}(G))$  and that  $W, W^{-1} \in B(L^2(G)) \overline{\otimes} \text{VN}(G)$ . Moreover, we will compute the symbol  $\varphi_{x,y,T}$ , belonging to  $L^\infty(G \times G)$ , of the Schur multiplier  $M_{x,y,T}$  and we will get

$$(4.15) \quad \varphi_{x,y,T}(s, t) = \tau_G(\lambda_t y \lambda_{s^{-1}} T(\lambda_s x \lambda_{t^{-1}})) \quad s, t \in G.$$

In the particular case of *finite* groups, these assertions are straightforward and we refer to the end of this section for a short proof of (4.14) and (4.15).

For the case of a locally compact group, this step unfortunately uses a painful approximation procedure described in Section 4.3 relying on a sequence  $(M_{\phi_n})$  of completely bounded Fourier multipliers  $M_{\phi_n}: L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  which allows us to consider the completely bounded maps  $M_{\phi_n}T: L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  in order to reduce the problem to the level  $p = 2$ .

We therefore obtain a map  $P_{x,y}: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$ ,  $T \mapsto M_{x,y,T}$  and it is easy to check that this map preserves the complete positivity. Introducing suitable sequences  $(x_j)$  and  $(y_j)$  of elements of  $L^1(\text{VN}(G)) \cap \text{VN}(G)$  which approximate the element 1 we obtain a sequence  $(P_j)$  of linear maps  $P_j \stackrel{\text{def}}{=} P_{x_j, y_j}: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$ . One of the difficulties in this area is to construct suitable sequences with the chosen assumptions on the group  $G$ .

Essentially, in the sequel we capture a cluster point of the bounded family  $(P_j)$  and we obtain a bounded map  $P^{(1)}: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$ . Each  $P^{(1)}(T)$  is a completely bounded Schur multiplier.

**Case where  $G$  is inner amenable and  $p = \infty$**  With a *suitable* choice of the sequences  $(x_j)$  and  $(y_j)$  provided by the inner amenability of  $G$ , the map  $P^{(1)}: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}(B(L^2(G)))$  is *contractive* and the Schur multiplier  $P^{(1)}(T): B(L^2(G)) \rightarrow B(L^2(G))$  is a *Herz-Schur* multiplier for all normal completely bounded map  $T: \text{VN}(G) \rightarrow \text{VN}(G)$ . So, we can see the linear map  $P^{(1)}$  as a map  $P^{(1)}: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \mathfrak{M}_G^{\infty, \text{HS}} = \mathfrak{M}_G^{\infty, \text{cb}, \text{HS}}$ . Now, it suffices to identify (completely) bounded Herz-Schur multipliers acting on the space  $B(L^2(G))$  isometrically with completely bounded Fourier multipliers acting on  $\text{VN}(G)$ , while preserving the complete positivity. This step is well-known [BoF] and true for any locally compact group  $G$  *without* amenability assumption. Denoting  $I: \mathfrak{M}_G^{\infty, \text{HS}} \rightarrow \text{CB}(\text{VN}(G))$  the associated isometry with range  $\mathfrak{M}^{\infty, \text{cb}}(G)$ , the final contractive projection will be  $P_G^\infty \stackrel{\text{def}}{=} I \circ P^{(1)}$ .

Indeed, in the case where  $T = M_\phi: \text{VN}(G) \rightarrow \text{VN}(G)$  is a Fourier multiplier we will prove that the symbol  $\phi_{j,T}$  of the Schur multiplier  $P_j(M_\phi): \text{B}(\text{L}^2(G)) \rightarrow \text{B}(\text{L}^2(G))$  is equal to the symbol  $\phi^{\text{HS}}: (s, t) \mapsto \phi(st^{-1})$  for any  $j$ . By passing the limit,  $P^{(1)}(M_\phi) = M_{\phi^{\text{HS}}}$  and finally

$$P_G^\infty(M_\phi) = I \circ P^{(1)}(M_\phi) = I(M_{\phi^{\text{HS}}}) = M_\phi.$$

So we obtain the property  $(\kappa_\infty)$  of Definition 1.2 for these groups with constant  $\kappa_\infty(G) = 1$ .

**Case where  $G$  is amenable and finite-dimensional and simultaneous cases  $p = 1$  and  $p = \infty$**  In the case where the group  $G$  is in addition finite-dimensional and amenable, replacing the sequences  $(x_j)$  and  $(y_j)$  of the proof of the last case by new ones, we obtain linear map  $P_p^{(1)}: \text{CB}(\text{L}^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$  for  $p = 1$  and  $p = \infty$  (replace  $\text{CB}(\text{L}^p(\text{VN}(G)))$  by  $\text{CB}_{w^*}(\text{VN}(G))$  here an the sequel), that we see as maps  $P_p^{(1)}: \text{CB}(\text{L}^p(\text{VN}(G))) \rightarrow \mathfrak{M}_G^{p,\text{cb}}$ . The cost of this replacement of sequences is the non-contractivity of  $P_p^{(1)}$  but we obtain the compatibility of the maps  $P_\infty^{(1)}(T)$  and  $P_1^{(1)}(T)$ . For the construction of the sequences  $(x_j)$  and  $(y_j)$ , our approach relies on the structure of finite-dimensional locally compact groups from the solution to Hilbert's fifth problem which makes appear connected Lie groups in this context and the use of Carnot-Carathéodory metrics on connected Lie groups.

Now, we construct and use a contractive map  $Q_p: \mathfrak{M}_G^{p,\text{cb}} \rightarrow \mathfrak{M}_G^{p,\text{cb,HS}}$  from the space  $\mathfrak{M}_G^{p,\text{cb}}$  of Schur multipliers onto the subspace of Herz-Schur multipliers which preserves the complete positivity and the Herz-Schur multipliers<sup>17</sup>. In this essentially folklore step, we need the amenability of the group  $G$  in sharp contrast with our previous work [ArK1]. Then put  $P_p^{(2)} \stackrel{\text{def}}{=} Q_p \circ P_p^{(1)}: \text{CB}(\text{L}^p(\text{VN}(G))) \rightarrow \mathfrak{M}_G^{p,\text{cb,HS}}$ .

At present, it suffices with [BoF] to identify completely bounded Herz-Schur multipliers isometrically with completely bounded Fourier multipliers, preserving the complete positivity. Denoting  $I_p: \mathfrak{M}_G^{p,\text{cb,HS}} \rightarrow \text{CB}(\text{L}^p(\text{VN}(G)))$  the associated isometry with range  $\mathfrak{M}^{p,\text{cb}}(G)$ , the final contractive projection will be  $P_G^p \stackrel{\text{def}}{=} I_p \circ P_p^{(2)} = I_p \circ Q_p \circ P_p^{(1)}$ .

In the case where  $T = M_\phi$  is a Fourier multiplier, we will prove that the symbol  $\phi_{j,T}$ , element of  $\text{L}^\infty(G \times G)$ , of the completely bounded Schur multiplier  $P_j(T)$  converges to the symbol  $\phi^{\text{HS}}: (s, t) \mapsto \phi(st^{-1})$  for the weak\* topology of  $\text{L}^\infty(G \times G)$ , using the continuity of the symbol  $\phi$ . From this, we deduce that the limit  $P_p^{(1)}(M_\phi)$  of the sequence  $(P_j(T))$  also admits the symbol  $\phi^{\text{HS}}$ . We conclude that

$$P_G^p(M_\phi) = I_p \circ Q_p \circ P_p^{(1)}(M_\phi) = I_p \circ Q_p(M_{\phi^{\text{HS}}}) = I_p(M_{\phi^{\text{HS}}}) = M_\phi.$$

We conclude that we obtain the property  $(\kappa)$  of Definition 1.7 for these groups. For totally disconnected groups, the method gives the sharp result  $\kappa(G) = 1$ .

**Case where  $G$  is amenable and  $1 < p < \infty$  such that  $\frac{p}{p^*}$  is rational.** In the case where the group  $G$  is amenable, using some sequences  $(x_j)$  and  $(y_j)$ , we obtain a *contractive* linear map  $P_p^{(1)}: \text{CB}(\text{L}^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$  which is better than the *boundedness* of the previous case, but only for *one* value of  $p$ . The method is similar to the previous case but we use [CDS] (see also [NR]) instead of [BoF] to identify completely bounded Herz-Schur multipliers isometrically with completely bounded Fourier multipliers (which require the amenability of  $G$  once again). Consequently, we only obtain a map  $P_G^p: \text{CB}(\text{L}^p(\text{VN}(G))) \rightarrow \text{CB}(\text{L}^p(\text{VN}(G)))$  whose range is included in  $\mathfrak{M}_G^{p,\text{cb}}$  such that  $P_G^p(M_\phi) = M_\phi$  if the symbol  $\phi$  is continuous.

17. We can see  $Q$  as a contractive projection  $Q_p: \mathfrak{M}_G^{p,\text{cb}} \rightarrow \mathfrak{M}_G^{p,\text{cb,HS}}$  onto the subspace  $\mathfrak{M}_G^{p,\text{cb,HS}}$  of completely bounded Herz-Schur multipliers.

We give an example which shows that we cannot hope for a compatibility for two distinct values of  $p$  with this choice of sequences  $(x_j)$  and  $(y_j)$  in a next version of this preprint. We also examine the case where  $\frac{p}{p^*}$  is not rational.

**Particular case of finite groups: proof of (4.14) and (4.15)** If the group  $G$  is finite and if  $(e_i)$  is an orthonormal basis of the Hilbert space  $\ell_G^2$  then (4.11) translates to

$$(4.16) \quad W(e_t \otimes e_r) = e_t \otimes e_{tr}, \quad W^{-1}(e_t \otimes e_r) = e_t \otimes e_{t^{-1}r}, \quad t, r \in G.$$

For any  $i, j, s, t, u \in G$ , we have

$$\begin{aligned} W(e_{st} \otimes \lambda_u)W^{-1}(e_i \otimes e_j) &\stackrel{(4.16)}{=} W(e_{st} \otimes \lambda_u)(e_i \otimes e_{i^{-1}j}) = W(e_{st}e_i \otimes \lambda_u(e_{i^{-1}j})) \\ &= \delta_{t=i}W(e_s \otimes e_{ui^{-1}j}) \stackrel{(4.16)}{=} \delta_{t=i}e_s \otimes e_{sut^{-1}j}. \end{aligned}$$

Hence in  $B(\ell^2(G)) \overline{\otimes} \text{VN}(G)$ , we have

$$(4.17) \quad W(e_{st} \otimes \lambda_u)W^{-1} = e_{st} \otimes \lambda_{sut^{-1}}.$$

We deduce that

$$(4.18) \quad (\text{Id} \otimes T)(W(e_{st} \otimes \lambda_u)W^{-1}) = e_{st} \otimes T(\lambda_{sut^{-1}}).$$

We infer that

$$\begin{aligned} &(\text{Tr} \otimes \tau_G)[(\text{Id} \otimes T)(W(e_{st} \otimes \lambda_u)W^{-1})(W(e_{ij} \otimes \lambda_r)W^{-1})] \\ &\stackrel{(4.18)(4.17)}{=} (\text{Tr} \otimes \tau_G)[(e_{st} \otimes T(\lambda_{sut^{-1}}))(e_{ij} \otimes \lambda_{irj^{-1}})] \\ &= \text{Tr}(e_{st}e_{ij})\tau_G(T(\lambda_{sut^{-1}})\lambda_{irj^{-1}}) = \delta_{t=i}\delta_{s=j}\tau_G(\lambda_i\lambda_r\lambda_{j^{-1}}T(\lambda_{sut^{-1}})). \end{aligned}$$

By linearity, we deduce on the one hand for any  $x \in L^p(\text{VN}(G))$  and any  $y \in L^{p^*}(\text{VN}(G))$

$$(\text{Tr} \otimes \tau_G)[(\text{Id} \otimes T)(W(e_{st} \otimes x)W^{-1})(W(e_{ij} \otimes y)W^{-1})] = \delta_{t=i}\delta_{s=j}\tau_G(\lambda_i y \lambda_{j^{-1}} T(\lambda_s x \lambda_{t^{-1}}))$$

On the other hand, if we consider the Schur multiplier  $M_{x,y,T}: S_G^p \rightarrow S_G^{p^*}$  with symbol (4.15), we have

$$\langle M_{x,y,T}(e_{st}), e_{ij} \rangle_{S_G^p, S_G^{p^*}} \stackrel{(4.15)}{=} \delta_{t=i}\delta_{s=j}\tau_G(\lambda_t y \lambda_{s^{-1}} T(\lambda_s x \lambda_{t^{-1}})).$$

**Remark 4.2** Note that with  $x = y = 1$ ,  $M_{x,y,T}$  is a Herz-Schur multiplier. See Section 4.4 for a generalization of this crucial observation.

### 4.3 Step 1: the mappings $P_j(T)$

In this section we will show (4.14) and (4.15). Unfortunately, we warn the reader that this part is unfortunately technically involved but the idea is quite simple. We reduce the computation to the level  $p = 2$  where we can use Parseval's identity.

Let  $G$  be a unimodular locally compact group. We denote by  $\text{Tr}_G$  and  $\tau_G$  the canonical traces  $\text{Tr}_G$  of the von Neumann algebras  $B(L^2(G))$  and  $\text{VN}(G)$ . Suppose that  $G$  and fix an orthonormal basis  $(e_i)$  of the Hilbert space  $L^2(G)$  such that each function  $e_i$  is continuous with compact

support<sup>18</sup>. Note that by [BLM, page 40] we have a canonical identification  $B(L^2(G)) \overline{\otimes} VN(G) = M_\infty(VN(G))$ . That means that an element  $X$  belonging to the von Neumann tensor product  $B(L^2(G)) \overline{\otimes} VN(G)$  identifies to a matrix  $[x_{ij}]$  with entries in  $VN(G)$ . For any  $h \in L^2(G)$  and any integer  $k$ , note that in  $L^2(G)$

$$(4.19) \quad (x_{kk}(h))(w) = \int_G (X(e_k \otimes h))(s, w) \overline{e_k(s)} d\mu_G(s), \quad h \in L^2(G), \text{ a.e. } w \in G.$$

If  $1 \leq p < \infty$ , we have by [Pis5] a similar isometry  $L^p(B(L^2(G)) \overline{\otimes} VN(G)) = S_G^p(L^p(VN(G)))$ . Moreover, if  $X$  belongs to the intersection  $L^1(B(L^2(G)) \overline{\otimes} VN(G)) \cap [B(L^2(G)) \overline{\otimes} VN(G)]$  we have  $x_{kk} \in L^1(VN(G)) \cap VN(G)$  for any integer  $k$  and

$$(4.20) \quad (\text{Tr}_G \otimes \tau_G)(X) = \sum_{k=1}^{\infty} \tau_G(x_{kk}).$$

In the next result, we use the operator  $W$  of  $B(L^2(G)) \overline{\otimes} VN(G)$  and its inverse from (4.11).

**Lemma 4.3** *Let  $G$  be a second countable unimodular locally compact group.*

1. *Let  $\phi \in L^2(G \times G)$  such that  $K_\phi$  belongs to  $S_G^1$  and  $x \in L^1(VN(G)) \cap VN(G)$ . Then  $W(K_\phi \otimes x)W^{-1}$  belongs to  $L^1(B(L^2(G)) \overline{\otimes} VN(G))$  and to  $B(L^2(G)) \overline{\otimes} VN(G)$ .*
2. *If  $f$  belongs to  $C_c(G)$  and if  $g$  belongs to  $C_c(G) * C_c(G)$ , we have for any integer  $i, j$*

$$(4.21) \quad (\text{Tr}_G \otimes \tau_G)[W(K_\phi \otimes \lambda(g))W^{-1} \cdot (e_{ij}^* \otimes \lambda(f))] \\ = \int_G \int_G \phi(s, t) \tau_G[\lambda_s \lambda(g) \lambda_{t^{-1}} \lambda(f)] \overline{e_i(s)} e_j(t) d\mu_G(s) d\mu_G(t).$$

*Proof* : 1. The element  $K_\phi \otimes x$  belongs to  $S_G^1 \otimes [L^1(VN(G)) \cap VN(G)]$ , hence to the space  $L^1(B(L^2(G)) \overline{\otimes} VN(G)) \cap [B(L^2(G)) \overline{\otimes} VN(G)]$ . Then the claim follows since  $W$  and  $W^{-1}$  belong to the space  $B(L^2(G)) \overline{\otimes} VN(G)$  and since  $L^1(B(L^2(G)) \overline{\otimes} VN(G)) \cap [B(L^2(G)) \overline{\otimes} VN(G)]$  is an ideal of the von Neumann algebra  $B(L^2(G)) \overline{\otimes} VN(G)$ .

2. By the first part, observe that the element  $X \stackrel{\text{def}}{=} W(K_\phi \otimes \lambda(g))W^{-1} \cdot (e_{ij}^* \otimes \lambda(f))$  belongs to the intersection  $L^1(B(L^2(G)) \overline{\otimes} VN(G)) \cap [B(L^2(G)) \overline{\otimes} VN(G)]$ . According to (4.20), we have

$$(\text{Tr}_G \otimes \tau_G)[W(K_\phi \otimes \lambda(g))W^{-1}(e_{ij}^* \otimes \lambda(f))] \stackrel{(4.20)}{=} \sum_{k=1}^{\infty} \tau_G(x_{kk})$$

Now, we want to compute  $\tau_G(x_{kk})$  with (4.19). If  $k \neq i$ , we have

$$X(e_k \otimes h) = W(K_\phi \otimes \lambda(g))W^{-1}(e_{ji} \otimes \lambda(f))(e_k \otimes h) = 0.$$

Hence  $x_{kk} = 0$  in this case and therefore  $\tau_G(x_{kk}) = 0$ . Thus, we only need to consider  $k = i$  in

---

18. To see that such an orthonormal basis exists, consider a sequence of continuous functions with compact support which is dense in  $L^2(G)$  and apply the Gram Schmidt procedure to it.

the sequel. Then replacing  $r$  by  $tv^{-1}s^{-1}r$  in the last equation for  $h \in C_c(G)$

$$\begin{aligned}
(4.22) \quad (X(e_i \otimes h))(s, w) &= (W(K_\phi \otimes \lambda(g))W^{-1}(e_{ji} \otimes \lambda(f))(e_i \otimes h))(s, w) \\
&= (W(K_\phi \otimes \lambda(g))W^{-1}(e_j \otimes \lambda(f)h))(s, w) \\
&\stackrel{(4.11)}{=} ((K_\phi \otimes \lambda(g))W^{-1}(e_j \otimes \lambda(f)h))(s, s^{-1}w) \\
&\stackrel{(4.1)(4.5)}{=} \int_G \int_G \phi(s, t)g(v)W^{-1}(e_j \otimes \lambda(f)h)(t, v^{-1}s^{-1}w) d\mu_G(t) d\mu_G(v) \\
&\stackrel{(4.11)}{=} \int_G \int_G \phi(s, t)g(v)(e_j \otimes \lambda(f)h)(t, tv^{-1}s^{-1}w) d\mu_G(t) d\mu_G(v) \\
&\stackrel{(4.5)}{=} \int_G \int_G \int_G \phi(s, t)e_j(t)g(v)f(r)h(r^{-1}tv^{-1}s^{-1}w) d\mu_G(t) d\mu_G(v) d\mu_G(r) \\
&= \int_G \int_G \int_G \phi(s, t)g(v)f(tv^{-1}s^{-1}r)h(r^{-1}w)e_j(t) d\mu_G(t) d\mu_G(v) d\mu_G(r).
\end{aligned}$$

Hence for almost all  $t \in G$

$$\begin{aligned}
(x_{ii}(h))(w) &\stackrel{(4.19)}{=} \int_G (X(e_i \otimes h))(s, w)\overline{e_i(s)} d\mu_G(s) \\
&\stackrel{(4.22)}{=} \int_G \int_G \int_G \int_G \phi(s, t)g(v)f(tv^{-1}s^{-1}r)h(r^{-1}w)\overline{e_i(s)}e_j(t) d\mu_G(t) d\mu_G(v) d\mu_G(r) d\mu_G(s).
\end{aligned}$$

So we obtain

$$x_{ii} = \int_G \int_G \int_G \int_G \phi(s, t)g(v)f(tv^{-1}s^{-1}r)\overline{e_i(s)}e_j(t)\lambda_{r^{-1}} d\mu_G(t) d\mu_G(v) d\mu_G(r) d\mu_G(s).$$

which identifies to the convolution operator  $\lambda(k)$  where  $k$  is the function defined by

$$(4.23) \quad k(r) \stackrel{\text{def}}{=} \int_G \int_G \int_G \phi(s, t)g(v)f(tv^{-1}s^{-1}r)\overline{e_i(s)}e_j(t) d\mu_G(t) d\mu_G(v) d\mu_G(s).$$

We can easily evaluate the trace of the diagonal entry  $x_{ii}$  which is an element of  $L^1(\text{VN}(G)) \cap \text{VN}(G)$ . Indeed, replacing  $v$  by  $s^{-1}v$  in the second equality, we have

$$\begin{aligned}
\tau_G(x_{ii}) &= k(e) \stackrel{(4.23)}{=} \int_G \int_G \int_G \phi(s, t)g(v)f(tv^{-1}s^{-1})\overline{e_i(s)}e_j(t) d\mu_G(t) d\mu_G(v) d\mu_G(s) \\
&= \int_G \int_G \int_G \phi(s, t)g(s^{-1}v)f(tv^{-1})\overline{e_i(s)}e_j(t) d\mu_G(t) d\mu_G(v) d\mu_G(s) \\
&= \int_G \int_G \phi(s, t) \left( \int_G g(s^{-1}v)f(tv^{-1}) d\mu_G(v) \right) \overline{e_i(s)}e_j(t) d\mu_G(t) d\mu_G(s) \\
&\stackrel{(4.7)}{=} \int_G \int_G \phi(s, t)\tau_G[\lambda_s \lambda(g)\lambda_{t^{-1}} \lambda(f)]\overline{e_i(s)}e_j(t) d\mu_G(t) d\mu_G(s).
\end{aligned}$$

■

In a similar way to Lemma 4.3, we have the following result.

**Lemma 4.4** *Let  $G$  be a second countable unimodular locally compact group. Let  $\phi \in L^2(G \times G)$  such that  $K_\phi \in S_G^1$  and  $x \in L^1(\text{VN}(G)) \cap \text{VN}(G)$ . Suppose  $1 \leq p \leq 2$ . Let  $T: L^p(\text{VN}(G)) \rightarrow$*

$L^2(\text{VN}(G))$  be a completely bounded operator. Then  $(\text{Id} \otimes T)(W(K_\phi \otimes x)W^{-1})$  belongs to  $L^2(\text{B}(L^2(G)) \overline{\otimes} \text{VN}(G))$ . If  $g \in C_c(G) * C_c(G)$  we have for any integer  $i, j$  and any  $f \in C_c(G)$

$$(4.24) \quad (\text{Tr}_G \otimes \tau_G)[(\text{Id} \otimes T)(W(K_\phi \otimes \lambda(g))W^{-1}) \cdot (e_{ij}^* \otimes \lambda(f))] \\ = \int_G \int_G \phi(s, t) \tau_G[\lambda_s \lambda(g) \lambda_{t^{-1}} T^*(\lambda(f))] \overline{e_i(s)} e_j(t) d\mu_G(s) d\mu_G(t).$$

*Proof* : By Lemma 4.3, the element  $W(K_\phi \otimes \lambda(g))W^{-1}$  belongs to  $L^1(\text{B}(L^2(G)) \overline{\otimes} \text{VN}(G)) \cap [\text{B}(L^2(G)) \overline{\otimes} \text{VN}(G)]$ , hence to the space  $L^p(\text{B}(L^2(G)) \overline{\otimes} \text{VN}(G)) = S_G^p(L^p(\text{VN}(G)))$ . By the complete boundedness of  $T: L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  and [Pis5, Lemma 1.7], we infer that the element  $(\text{Id}_{S_G^p} \otimes T)(W(K_\phi \otimes x)W^{-1})$  belongs to the space  $S_G^p(L^2(\text{VN}(G)))$ . Since  $p \leq 2$ , it belongs to the space  $S_G^2(L^2(\text{VN}(G))) = L^2(\text{B}(L^2(G)) \overline{\otimes} \text{VN}(G))$ . We have immediately

$$(\text{Tr}_G \otimes \tau_G)[(\text{Id} \otimes T)(W(K_\phi \otimes \lambda(g))W^{-1}) \cdot (e_{ij}^* \otimes \lambda(f))] \\ = (\text{Tr}_G \otimes \tau_G)[(W(K_\phi \otimes \lambda(g))W^{-1}) \cdot (e_{ij}^* \otimes T^* \lambda(f))]$$

where  $R: \text{VN}(G) \rightarrow \text{VN}(G)$ ,  $\lambda_s \rightarrow \lambda_{s^{-1}}$  is the antipode. Now, it suffices to show that (4.21) holds for generic elements of  $L^{p^*}(\text{VN}(G))$  instead of  $\lambda(f)$ . This is indeed the case by density since both sides of (4.21) are continuous as functions in  $\lambda(f) \in L^{p^*}(\text{VN}(G))$ . ■

**Remark 4.5** The two previous lemma are true without the assumption ‘‘second countable’’.

**Lemma 4.6** *Let  $G$  be a second countable unimodular locally compact group. Let  $1 \leq p \leq 2$  and  $T: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))$  be a completely bounded map. There exists a sequence  $(M_{\phi_n})$  of bounded Fourier multipliers  $M_{\phi_n}: \text{VN}(G) \rightarrow \text{VN}(G)$  such that  $\phi_n \in C_c(G)$ ,  $\|\phi_n\|_\infty \leq 1$ ,  $M_{\phi_n}: L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  is completely bounded, satisfying for any  $g \in C_c(G)$ , any  $\phi \in C_c(G \times G)$  such that  $K_\phi \in S_G^1$  and any sufficiently large  $n$*

$$M_{\phi_n}(\lambda(g)) = \lambda(g) \quad \text{and} \quad (\text{Id} \otimes M_{\phi_n})(W(K_\phi \otimes \lambda(g))W^{-1}) = W(K_\phi \otimes \lambda(g))W^{-1}.$$

*Proof* : Since  $G$  is second countable, we can consider a sequence  $(K_n)$  of symmetric compacts in  $G$  such that for any compact  $K$  of  $G$ , one has  $K \subset K_n$  for sufficiently large enough  $n$ . By [KaL1, Proposition 2.3.2], for any  $n$  there exists a function  $\phi_n: G \rightarrow \mathbb{C}$  which is a finite linear combination of continuous definite positive functions with compact support with  $0 \leq \phi_n \leq 1$  and  $\phi_n(s) = 1$  for any  $s \in K_n$ . By essentially [KaL1, Proposition 5.4.9] each function induces a completely bounded Fourier multiplier  $M_{\phi_n}: \text{VN}(G) \rightarrow \text{VN}(G)$ . Furthermore, since  $\phi_n \in L^2(G)$  each  $M_{\phi_n}: L^1(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  is completely bounded by [GJP17, Remark 2.4] and duality. Now, it suffices to interpolate with  $M_{\phi_n}: L^2(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  to obtain a completely bounded Fourier multiplier  $M_{\phi_n}: L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$ .

Now, let  $g \in C_c(G)$  and  $\phi \in C_c(G \times G)$  such that  $K_\phi \in S_G^1$ . Consider the compact  $K \stackrel{\text{def}}{=} \text{supp } g$  and some compacts  $L_1, L_2$  of  $G$  such that  $\text{supp } \phi \subset L_1 \times L_2$  and let  $L \stackrel{\text{def}}{=} L_1 \cdot K \cdot L_2^{-1}$ , which is also compact. Then for any sufficiently large enough  $n$  such that  $K \subset K_n$ ,

$$M_{\phi_n}(\lambda(g)) = \lambda(\phi_n g) = \lambda(g).$$

Moreover, consider some sufficiently large enough  $n$  such that  $L \subset K_n = \check{K}_n$ . For any  $s \in L_1$  and any  $t \in L_2$ , the element  $\lambda_s \lambda(g) \lambda_{t^{-1}}$  has its Fourier support in  $L_1 \cdot K \cdot L_2^{-1} = L$ . Thus,  $M_{\check{\phi}_n}(\lambda_s \lambda(g) \lambda_{t^{-1}}) = \lambda_s \lambda(g) \lambda_{t^{-1}}$ . Then for any integer  $i, j$  and any  $f \in C_c(G) * C_c(G)$  we have

$M_{\phi_n}^* = M_{\check{\phi}_n}$ . Hence

$$\begin{aligned}
& (\mathrm{Tr}_G \otimes \tau_G) [(\mathrm{Id} \otimes M_{\phi_n})(W(K_\phi \otimes \lambda(g))W^{-1}) \cdot (e_{ij}^* \otimes \lambda(f))] \\
& \stackrel{(4.24)}{=} \int_G \int_G \phi(s, t) \tau_G [\lambda_s \lambda(g) \lambda_{t^{-1}} M_{\phi_n}^*(\lambda(f))] \overline{e_i(s)} e_j(t) \, d\mu_G(s) \, d\mu_G(t). \\
& = \int_{L_1} \int_{L_2} \phi(s, t) \tau_G [M_{\check{\phi}_n}(\lambda_s \lambda(g) \lambda_{t^{-1}}) \lambda(f)] \overline{e_i(s)} e_j(t) \, d\mu_G(s) \, d\mu_G(t) \\
& = \int_{L_1} \int_{L_2} \phi(s, t) \tau_G [\lambda_s \lambda(g) \lambda_{t^{-1}} \lambda(f)] \overline{e_i(s)} e_j(t) \, d\mu_G(s) \, d\mu_G(t) \\
& \stackrel{(4.21)}{=} (\mathrm{Tr}_G \otimes \tau_G) [W(K_\phi \otimes \lambda(g))W^{-1} \cdot (e_{ij}^* \otimes \lambda(f))].
\end{aligned}$$

By density of the  $e_{ij}^* \otimes \lambda(f)$ 's, we infer that  $(\mathrm{Id} \otimes M_{\phi_n})(W(K_\phi \otimes \lambda(g))W^{-1}) = W(K_\phi \otimes \lambda(g))W^{-1}$ .  $\blacksquare$

Recall that  $(e_n)$  is an orthonormal basis of  $L^2(G)$  such that each function  $e_n$  is continuous with compact support. So  $(\lambda(e_k))$  is an orthonormal basis of  $L^2(\mathrm{VN}(G))$  and  $(e_{ij} \otimes \lambda(e_k))_{i,j,k}$  is an orthonormal basis of  $L^2(\mathrm{B}(L^2(G)) \overline{\otimes} \mathrm{VN}(G))$ .

**Proposition 4.7** *Let  $G$  be a second countable unimodular locally compact group. Suppose  $1 \leq p \leq \infty$ . Let  $T: L^p(\mathrm{VN}(G)) \rightarrow L^p(\mathrm{VN}(G))$  be a completely bounded operator (normal if  $p = \infty$ ). Let  $\phi, \psi \in L^2(G \times G)$  such that  $K_\phi, K_\psi \in S_G^1$ , and  $x, y \in L^1(\mathrm{VN}(G)) \cap \mathrm{VN}(G)$ . With the symbol*

$$(4.25) \quad \varphi_{x,y,T}(s, t) \stackrel{\mathrm{def}}{=} \tau_G(\lambda_t y \lambda_{s^{-1}} T(\lambda_s x \lambda_{t^{-1}})).$$

we have

$$(4.26) \quad \langle (\mathrm{Id} \otimes T)(W(u \otimes x)W^{-1}), W(v \otimes y)W^{-1} \rangle_{S_G^p(L^p(\mathrm{VN}(G))), S_G^{p*}(L^{p*}(\mathrm{VN}(G)))} = \langle M_{x,y,T}(u), v \rangle_{S_G^p, S_G^{p*}}.$$

Finally, if  $p = \infty$ , the same holds for  $x \in \mathrm{VN}(G)$  and if  $p = 1$ , the same holds for  $y \in \mathrm{VN}(G)$ .

*Proof* : Note first that by a simple duality argument, we can suppose that  $1 \leq p \leq 2$  and that the functions  $\phi, \psi$  belongs to  $C_c(G \times G)$ .

We start with the case where  $T$  also induces a completely bounded map  $T: L^p(\mathrm{VN}(G)) \rightarrow L^2(\mathrm{VN}(G))$ . Then by Lemma 4.3 and Lemma 4.4, the elements  $W(K_\psi \otimes y)W^{-1}$  and  $(\mathrm{Id} \otimes T)(W(K_\phi \otimes x)W^{-1})$  belong to the Hilbert space  $L^2(\mathrm{B}(L^2(G)) \overline{\otimes} \mathrm{VN}(G))$ . So the left-hand side of (4.27) is well-defined, and can be calculated with Parseval's formula and the orthonormal basis  $(e_{ij} \otimes \lambda(e_k))_{i,j,k}$ . With Lemma 4.4, we get

$$\begin{aligned}
& \langle (\mathrm{Id} \otimes T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle_{S_G^2(L^2(\mathrm{VN}(G)))} \\
& = \sum_{i,j,k} (\mathrm{Tr}_G \otimes \tau_G) [(\mathrm{Id} \otimes T)(W(K_\phi \otimes x)W^{-1})(e_{ij}^* \otimes \lambda(e_k^*))] \\
& \quad \times \overline{(\mathrm{Tr}_G \otimes \tau_G) [W(K_\psi \otimes y)W^{-1}(e_{ij}^* \otimes \lambda(e_k^*))]} \\
& \stackrel{(4.24)(4.21)}{=} \sum_{i,j,k} \int_G \int_G \phi(s, t) \tau_G [\lambda_s x \lambda_{t^{-1}} T^*(\lambda(e_k^*))] \overline{e_i(s)} e_j(t) \, d\mu_G(s) \, d\mu_G(t) \\
& \quad \times \int_G \int_G \psi(s, t) \tau_G [\lambda_s y \lambda_{t^{-1}} \lambda(e_k^*)] \overline{e_i(s)} e_j(t) \, d\mu_G(s) \, d\mu_G(t).
\end{aligned}$$

Since the functions  $(s, t) \mapsto \tau_G[\lambda_s x \lambda_{t-1} T^*(\lambda(e_k^*))] \phi(s, t)$  and  $(s, t) \mapsto \tau_G[\lambda_s y \lambda_{t-1} \lambda(e_k^*)] \psi(s, t)$  belong to the Hilbert space  $L^2(G \times G)$ , we can use the orthonormal basis  $(\bar{e}_i \otimes e_j)_{i,j}$  of  $L^2(G \times G)$  and reduce in the previous expression the sum over  $i, j$  and the integral over  $s, t$ , which then becomes

$$\sum_k \int_G \int_G \phi(s, t) \tau_G[\lambda_s x \lambda_{t-1} T^*(\lambda(e_k^*))] \overline{\tau_G[\lambda_s y \lambda_{t-1} \lambda(e_k^*)]} d\mu_G(s) d\mu_G(t).$$

Recall that  $T^{*\circ} = T^\dagger$  where  $T^\dagger$  is the hilbertian adjoint. We fix  $s, t \in G$  and calculate using Parseval's identity in the fourth equality

$$\begin{aligned} \sum_k \tau_G[\lambda_s x \lambda_{t-1} T^*(\lambda(e_k^*))] \overline{\tau_G[\lambda_s y \lambda_{t-1} \lambda(e_k^*)]} &= \sum_k \langle \lambda_s x \lambda_{t-1}, (T^*(\lambda(e_k^*))^*)^* \rangle_{L^2} \overline{\langle \lambda_s y \lambda_{t-1}, \lambda(e_k) \rangle_{L^2}} \\ &= \sum_k \langle \lambda_s x \lambda_{t-1}, T^\dagger(\lambda(e_k)) \rangle_{L^2} \overline{\langle \lambda_s y \lambda_{t-1}, \lambda(e_k) \rangle_{L^2}} = \sum_k \langle T(\lambda_s x \lambda_{t-1}), \lambda(e_k) \rangle \overline{\langle \lambda_s y \lambda_{t-1}, \lambda(e_k) \rangle} \\ &= \langle T(\lambda_s x \lambda_{t-1}), \lambda_s y \lambda_{t-1} \rangle_{L^2(\text{VN}(G))} = \tau_G(\lambda_t y^* \lambda_{s-1} T(\lambda_s x \lambda_{t-1})). \end{aligned}$$

Note that  $\|T(\lambda_s x \lambda_{t-1})\|_2 \leq C$  and  $\|\lambda_s y \lambda_{t-1}\|_2 \leq C$  justifies that we can integrate over  $s$  and  $t$ . Consequently

$$\begin{aligned} &\langle (\text{Id} \otimes T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle_{S_G^2(L^2(\text{VN}(G)))} \\ &= \int_{G \times G} \tau_G(\lambda_t y^* \lambda_{s-1} T(\lambda_s x \lambda_{t-1})) \phi(s, t) \overline{\psi(s, t)} d\mu_G(s) d\mu_G(t) \\ &\stackrel{(4.2)}{=} \text{Tr}(K_{\varphi_{x,y,T}\phi} K_\psi^*) = \langle M_{\varphi_{x,y,T}}(K_\phi), K_\psi \rangle_{S_G^2}. \end{aligned}$$

Thus we have shown the formula

$$(4.27) \quad \langle (\text{Id} \otimes T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle_{S_G^2(L^2(\text{VN}(G)))} = \langle M_{\varphi_{x,y,T}}(K_\phi), K_\psi \rangle_{S_G^2}.$$

under the additional assumption that the linear map  $T$  defines a completely bounded operator  $L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$ . Replacing the both brackets, this formula translates to (4.26).

For the general case, we use the first part of Lemma 4.6 with the approximation sequence  $(M_{\phi_n})$  of Fourier multipliers  $M_{\phi_n} : \text{VN}(G) \rightarrow \text{VN}(G)$ . By a density argument, we can assume that  $y \in \lambda(C_c(G))$ . According to Lemma 4.6, the composition  $M_{\phi_n} T : L^p(\text{VN}(G)) \rightarrow L^2(\text{VN}(G))$  is completely bounded. So the first part of the proof applies to this operator. We obtain

$$\langle (\text{Id} \otimes M_{\phi_n} T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle = \langle M_{\varphi_n}(K_\phi), K_\psi \rangle_{S_G^2}$$

with  $\varphi_n(s, t) \stackrel{\text{def}}{=} \tau_G(\lambda_t y^* \lambda_{s-1} M_{\phi_n} T(\lambda_s x \lambda_{t-1}))$ . By the approximation from Lemma 4.6, we have

$$\begin{aligned} &\langle (\text{Id} \otimes M_{\phi_n} T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle \\ &= \langle (\text{Id} \otimes T)(W(K_\phi \otimes x)W^{-1}), (\text{Id} \otimes M_{\phi_n}^*)(W(K_\psi \otimes y)W^{-1}) \rangle \\ &\xrightarrow{n \rightarrow \infty} \langle (\text{Id} \otimes T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle. \end{aligned}$$

Again with Lemma 4.6, we have

$$\begin{aligned} \varphi_n(s, t) &= \langle M_{\phi_n} T(\lambda_s x \lambda_{t-1}), \lambda_s y \lambda_{t-1} \rangle \\ &= \langle T(\lambda_s x \lambda_{t-1}), M_{\phi_n}^*(\lambda_s y \lambda_{t-1}) \rangle \xrightarrow{n \rightarrow \infty} \langle T(\lambda_s x \lambda_{t-1}), \lambda_s y \lambda_{t-1} \rangle \stackrel{(4.25)}{=} \varphi_{x,y,T}(s, t) \end{aligned}$$



pointwise in  $s, t \in G$ . Even stronger, if  $s, t$  vary in given compacts,  $\varphi_n(s, t) = \varphi(s, t)$  for  $n$  sufficiently large. Since  $\phi, \psi$  are supposed to live in  $C_c(G \times G)$ , we obtain then  $\langle M_{\varphi_n}(K_\phi), K_\psi \rangle = \langle M_\varphi(K_\phi), K_\psi \rangle$  if  $n$  sufficiently large. In all, we have shown

$$\langle (\text{Id} \otimes T)(W(K_\phi \otimes x)W^{-1}), W(K_\psi \otimes y)W^{-1} \rangle = \langle M_\varphi(K_\phi), K_\psi \rangle.$$

■

**Lemma 4.8** *Suppose  $1 \leq p \leq \infty$ . We have*

$$(4.28) \quad \|M_{\varphi_{x,y,T}}\|_{\text{cb}, S_G^p \rightarrow S_G^p} \leq \|T\|_{\text{cb}, L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))} \|x\|_p \|y\|_{p^*}$$

with the usual convention if  $p = 1$  or  $p = \infty$ . Moreover, if  $T$  is completely positive then the Schur multiplier  $M_{\varphi_{x,y,T}}$  is also completely positive.

*Proof* : By [Pis4, Definition 2.1], the duality [Pis5, Theorem 4.7] and Plancherel formula (4.7), we have

$$\begin{aligned} & \|M_{\varphi_{x,y,T}}\|_{\text{cb}, S_G^p \rightarrow S_G^p} \\ & \leq \sup \left\{ \left| \sum_{ij} \langle (\text{Id} \otimes T)(W(K_{\phi_{ij}} \otimes x)W^{-1}), W(K_{\psi_{ij}} \otimes y)W^{-1} \rangle \right| : \| [K_{\phi_{ij}} ] \|_p, \| [K_{\psi_{ij}} ] \|_{p^*} \leq 1 \right\} \\ & \leq \| \text{Id} \otimes \text{Id} \otimes T \|_{\text{B}(S^p(S_G^p(L^p(\text{VN}(G)))))} \|W\|_\infty \|x\|_p \|W^{-1}\|_\infty \|W\|_\infty \|y\|_{p^*} \|W^{-1}\|_\infty \\ & \leq \|T\|_{\text{cb}, L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))} \|x\|_p \|y\|_{p^*}. \end{aligned}$$

If  $T: \text{VN}(G) \rightarrow \text{VN}(G)$  is normal and completely positive, then for positive  $[K_{\phi_{ij}}]$  and  $[K_{\psi_{ij}}]$  we have

$$\sum_{ij} \langle M_{x,y,T} K_{\phi_{ij}}, K_{\psi_{ij}} \rangle = \sum_{ij} \langle (\text{Id} \otimes T)(W(K_{\phi_{ij}} \otimes x)W^{-1}), W(K_{\psi_{ij}} \otimes y)W^{-1} \rangle \geq 0.$$

Indeed,  $\text{Id} \otimes \text{Id} \otimes T$  preserves the positivity and  $[W(K_{\phi_{ij}} \otimes x)W^{-1}]$  and  $[W(K_{\psi_{ij}} \otimes y)W^{-1}]$  are positive. We infer by [ArK1, Lemma 2.6] that  $[M_{\varphi_{x,y,T}} K_{\phi_{ij}}]$  is positive, hence  $M_{\varphi_{x,y,T}}$  is completely positive. ■

#### 4.4 Step 2: the symbol of $P_j(T)$ is Herz-Schur if $G$ is inner amenable

**Lemma 4.9** *Let  $G$  be a second countable unimodular locally compact group. Suppose that  $G$  is inner amenable. Let  $F$  be finite subset of  $G$  and  $(V_j^F)_j$  be a sequence of subsets  $G$  satisfying the last point of Theorem 3.7. Consider a weakly\* continuous completely bounded map  $T: \text{VN}(G) \rightarrow \text{VN}(G)$ . With the notation (4.25), we let*

$$(4.29) \quad y_j^F \stackrel{\text{def}}{=} c_j^F |\lambda(1_{V_j^F})|^2 \quad \text{and} \quad \phi_{j,T}^F \stackrel{\text{def}}{=} \phi_{1,y_j^F,T}$$

where  $c_j^F > 0$  is the normalisation to have  $\|y_j^F\|_{L^1(\text{VN}(G))} = 1$ . Then any weak\* cluster point  $\phi_T^F$  of the net  $(\phi_{j,T}^F)_j$  satisfies

$$\phi_T^F(sr, tr) = \phi_T^F(s, t), \quad s, t \in G, r \in F.$$

Moreover, any weak\* cluster point of such  $(\phi_T^F)_F$ , where the finite subsets  $F \subset G$  are directed by inclusion, is also a Herz-Schur symbol.

*Proof* : For any  $s, t \in G$  and  $r \in F$ , we have

$$\begin{aligned}
\phi_{j,T}^F(sr, tr) - \phi_{j,T}^F(s, t) &\stackrel{(4.29)}{=} \phi_{1, y_j^F, T}(sr, tr) - \phi_{1, y_j^F, T}(s, t) \\
&\stackrel{(4.25)}{=} \tau_G(\lambda_{tr} y_j^F \lambda_{(sr)^{-1}} T(\lambda_{sr} \lambda_{(tr)^{-1}})) - \tau_G(\lambda_t y_j^F \lambda_{s^{-1}} T(\lambda_s \lambda_{t^{-1}})) \\
&= \tau_G(\lambda_t \lambda_r y_j^F \lambda_{r^{-1}} \lambda_{s^{-1}} T(\lambda_{st^{-1}})) - \tau_G(\lambda_t y_j^F \lambda_{s^{-1}} T(\lambda_{st^{-1}})) \\
&= \tau_G(\lambda_t (\lambda_r y_j^F \lambda_{r^{-1}} - y_j^F) \lambda_{s^{-1}} T(\lambda_{st^{-1}})).
\end{aligned}$$

If we can show that

$$(4.30) \quad \|\lambda_r y_j^F \lambda_{r^{-1}} - y_j^F\|_1 \xrightarrow{j \rightarrow \infty} 0,$$

then we will obtain the pointwise convergence  $\phi_{j,T}^F(sr, tr) - \phi_{j,T}^F(s, t) \rightarrow 0$  as  $j \rightarrow \infty$ , for fixed  $s, t \in G$  and  $r \in F$ . Since  $\phi_{j,T}^F(sr, tr) - \phi_{j,T}^F(s, t)$  is uniformly bounded in  $L^\infty(G \times G)$ , by dominated convergence, it follows that this sequence converges for the weak\* topology to 0 in  $L^\infty(G \times G)$ . Thus, if  $\phi_T^F$  is a cluster point of  $\phi_{j,T}^F$ , it is easy to check by a  $\frac{\varepsilon}{3}$ -argument using the weak\* continuity of translations on  $L^\infty$  that  $\phi_T^F(sr, tr) - \phi_T^F(s, t)$  for any  $s, t \in G$  and  $r \in F$ . It remains to show (4.30).

First, for any  $j$  we have

$$(4.31) \quad (c_j^F)^{-1} = \|\lambda(1_{V_j^F})\|_1^2 = \|\lambda(1_{V_j^F})\|_2^2 = \|1_{V_j^F}\|_{L^2(G)}^2 = \mu(V_j^F).$$

Now, observe by unimodularity in the second equality

$$\begin{aligned}
&\|\lambda_r |\lambda(1_{V_j^F})|^2 \lambda_{r^{-1}} - |\lambda(1_{V_j^F})|^2\|_1 \stackrel{(4.6)}{=} \|\lambda_r \lambda(1_{V_j^F} * 1_{V_j^F}) \lambda_{r^{-1}} - \lambda(1_{V_j^F} * 1_{V_j^F})\|_1 \\
&= \|\lambda(1_{V_j^F} * 1_{V_j^F}(r^{-1}(\cdot)r) - 1_{V_j^F} * 1_{V_j^F})\|_1 \\
&\leq \|\lambda(1_{V_j^F} * 1_{V_j^F}(r^{-1}(\cdot)r) - 1_{V_j^F} * 1_{rV_j^F r^{-1}} + 1_{V_j^F} * 1_{rV_j^F r^{-1}} - 1_{V_j^F} * 1_{V_j^F})\|_1 \\
&\leq \|\lambda(1_{V_j^F} * 1_{rV_j^F r^{-1}} - 1_{V_j^F} * 1_{V_j^F}(r^{-1}(\cdot)r))\|_1 + \|\lambda(1_{V_j^F} * (1_{rV_j^F r^{-1}} - 1_{V_j^F}))\|_1.
\end{aligned}$$

We estimate the second summand with unimodularity by

$$\|1_{V_j^F}\|_{L^2(G)} \|1_{rV_j^F r^{-1}} - 1_{V_j^F}\|_{L^2(G)} \stackrel{(3.3)}{=} \mu(V_j^F)^{\frac{1}{2}} \mu(rV_j^F r^{-1} \Delta V_j^F)^{\frac{1}{2}}.$$

Now, we manipulate the first summand. By replacing  $t$  by  $tr^{-1}$  and  $t$  by  $r^{-1}t$  in the fourth equality, we obtain

$$\begin{aligned}
(4.32) \quad &1_{V_j^F} * 1_{rV_j^F r^{-1}}(s) - 1_{V_j^F} * 1_{V_j^F}(r^{-1}sr) \\
&\stackrel{(4.5)}{=} \int_G 1_{V_j^F}(t) 1_{rV_j^F r^{-1}}(t^{-1}s) - 1_{V_j^F}(t) 1_{V_j^F}(t^{-1}r^{-1}sr) \, d\mu_G(t) \\
&= \int_G 1_{V_j^F}(t) 1_{V_j^F r^{-1}}(r^{-1}t^{-1}s) \, d\mu_G(t) - \int_G 1_{V_j^F}(t) 1_{V_j^F r^{-1}}(t^{-1}r^{-1}s) \, d\mu_G(t) \\
&= \int_G 1_{V_j^F}(tr^{-1}) 1_{V_j^F r^{-1}}(t^{-1}s) \, d\mu_G(t) - \int_G 1_{V_j^F}(r^{-1}t) 1_{V_j^F r^{-1}}(t^{-1}s) \, d\mu_G(t) \\
&= \int_G (1_{V_j^F r} - 1_{rV_j^F})(t) 1_{V_j^F r^{-1}}(t^{-1}s) \, d\mu_G(t) \stackrel{(4.5)}{=} (1_{V_j^F r} - 1_{rV_j^F}) * 1_{V_j^F r^{-1}}(s).
\end{aligned}$$

Using the invariance of  $\mu_G$ , we therefore obtain the following estimate for the first summand:

$$\begin{aligned} \|\lambda(1_{V_j^F} * 1_{rV_j^F r^{-1}} - 1_{V_j^F} * 1_{V_j^F}(r^{-1}(\cdot)r))\|_1 &\stackrel{(4.32)}{\leq} \|1_{V_j^F r} - 1_{rV_j^F}\|_{L^2(G)} \|1_{V_j^F r^{-1}}\|_{L^2(G)} \\ &\stackrel{(3.3)}{=} \mu(V_j^F r \Delta r V_j^F)^{\frac{1}{2}} \mu(V_j^F r^{-1})^{\frac{1}{2}} = \mu(V_j^F \Delta r V_j^F r^{-1})^{\frac{1}{2}} \mu(V_j^F)^{\frac{1}{2}}. \end{aligned}$$

Combining the two estimates, we obtain

$$\|\lambda_r y_j \lambda_r^{-1} - y_j\|_1 \leq 2c_j^F \mu(V_j^F \Delta r V_j^F r^{-1})^{\frac{1}{2}} \mu(V_j^F)^{\frac{1}{2}} \stackrel{(4.31)}{=} 2 \left[ \frac{\mu(V_j^F \Delta r V_j^F r^{-1})}{\mu(V_j^F)} \right]^{\frac{1}{2}} \stackrel{(3.6)}{=} \frac{0}{j} \rightarrow 0$$

thanks to the inner amenability assumption.

Now fix for any finite subset  $F$  of  $G$  a weak\* cluster point  $\phi_T^F$  of  $(\phi_{j,T}^F)_j$ . Let  $\phi_T$  be a weak\* cluster point of  $(\phi_T^F)_F$ . Then for any  $f \in L^1(G \times G)$  and  $r \in G$ , the function  $f(\cdot r^{-1}, \cdot r^{-1})$  belongs to  $L^1(G)$ . Moreover, using unimodularity in the first and third steps and the first part of the proof valid for  $F$  containing  $\{r\}$  in the fourth step, we obtain

$$\begin{aligned} \langle \phi_T(\cdot r, \cdot r), f \rangle_{L^\infty(G \times G), L^1(G \times G)} &= \langle \phi_T, f(\cdot r^{-1}, \cdot r^{-1}) \rangle_{L^\infty, L^1} = \lim_{F \rightarrow \infty} \langle \phi_T^F, f(\cdot r^{-1}, \cdot r^{-1}) \rangle_{L^\infty, L^1} \\ &= \lim_{F \rightarrow \infty} \langle \phi_T^F(\cdot r, \cdot r), f \rangle_{L^\infty, L^1} = \lim_{F \rightarrow \infty} \langle \phi_T^F, f \rangle_{L^\infty, L^1} = \langle \phi_T, f \rangle_{L^\infty(G \times G), L^1(G \times G)}, \end{aligned}$$

We deduce that  $\phi_T$  is a Herz-Schur symbol. ■

#### 4.5 Step 2: the symbol of $P_j(T)$ for a Fourier multiplier $T$ if $p = \infty$ or $p = 1$

Let  $G$  be a second countable unimodular locally compact group. We start with the case  $p = \infty$ . Let  $T = M_\phi: \text{VN}(G) \rightarrow \text{VN}(G)$  be a completely bounded Fourier multiplier. If  $x \in \text{VN}(G)$  and  $y \in L^1(\text{VN}(G)) \cap \text{VN}(G)$ , recall that the symbol  $\varphi_{x,y,T}$  is defined in (4.25).

**Lemma 4.10** *Let  $y$  be a positive element of  $L^1(\text{VN}(G)) \cap \text{VN}(G)$  such that  $\tau_G(y) = 1$ . We have*

$$(4.33) \quad \varphi_{1,y,T}(s,t) = \phi(st^{-1}), \quad s, t \in G.$$

*Proof* : For any  $s, t \in G$ , we have

$$\begin{aligned} \varphi_{1,y,T}(s,t) &\stackrel{(4.25)}{=} \tau_G(y \lambda_{s^{-1}} M_\phi(\lambda_s \lambda_{t^{-1}}) \lambda_t) = \phi(st^{-1}) \tau_G(\lambda_t y \lambda_{s^{-1}} \lambda_s \lambda_{t^{-1}}) \\ &= \phi(st^{-1}) \tau_G(y) = \phi(st^{-1}). \end{aligned}$$

■

**Example 4.11** Let  $g$  be a continuous functions with compact support on  $G$  with  $\|g\|_{L^2(G)} = 1$ . With  $y \stackrel{\text{def}}{=} \lambda(g^* * g)$ , the assumptions of Lemma 4.10 are satisfied by (4.6) and since  $\tau_G(y) \stackrel{(4.6)}{=} \tau_G(\lambda(g)^* \lambda(g)) \stackrel{(4.7)}{=} \|g\|_{L^2(G)}^2 = 1$ .

We continue with the case  $p = 1$ . Let  $T = M_\phi: L^1(\text{VN}(G)) \rightarrow L^1(\text{VN}(G))$  be a completely bounded Fourier multiplier. If  $x \in L^1(\text{VN}(G)) \cap \text{VN}(G)$  and  $y \in \text{VN}(G)$ , recall again that the symbol  $\varphi_{x,y,T}$  is defined in (4.25). We can prove the following similar result.

**Lemma 4.12** *Let  $x$  be a positive element of  $L^1(\text{VN}(G)) \cap \text{VN}(G)$  such that  $\tau_G(x) = 1$ . We have*

$$(4.34) \quad \varphi_{x,1,T}(s, t) = \phi(st^{-1}), \quad s, t \in G.$$

*Proof* : For any  $s, t \in G$ , we have

$$\begin{aligned} \varphi_{x,1,T}(s, t) &\stackrel{(4.25)}{=} \tau_G(\lambda_t \lambda_{s^{-1}} M_\phi(\lambda_s x \lambda_{t^{-1}})) = \tau_G(\lambda_t \lambda_{s^{-1}} M_\phi(\lambda_s x \lambda_{t^{-1}})) \\ &= \tau_G(M_\phi(\lambda_{ts^{-1}} \lambda_s x \lambda_{t^{-1}})) = \phi(st^{-1}) \tau_G(\lambda_{ts^{-1}} \lambda_s x \lambda_{t^{-1}}) = \phi(st^{-1}) \tau_G(x) = \phi(st^{-1}). \end{aligned}$$

■

## 4.6 Step 2: convergence of the symbols for a multiplier $T$ with arbitrary symbol

**Proposition 4.13** *Let  $G$  be a second countable unimodular locally compact group. Suppose  $1 \leq p \leq \infty$ . Consider some completely bounded Fourier multiplier  $T = M_\phi: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))$ . Let  $(f_j)$  and  $(g_j)$  be sequences of positive functions with compact support belonging to  $C_c(G)$  such that if  $x_j \stackrel{\text{def}}{=} \lambda(f_j)$ ,  $y_j \stackrel{\text{def}}{=} \lambda(g_j)$  we have*

- $\|x_j\|_p \|y_j\|_{p^*} \leq C$  for all  $j$  for some positive constant  $C$ ,
- $\tau_G(x_j y_j) = 1$  for all  $j$ ,
- $\text{supp } f_j \rightarrow \{e\}$  or  $\text{supp } g_j \rightarrow \{e\}$ .

Moreover, let

$$(4.35) \quad \phi_{j,T}(s, t) \stackrel{\text{def}}{=} \tau_G(\lambda_t y_j \lambda_{s^{-1}} T(\lambda_s x_j \lambda_{t^{-1}})), \quad s, t \in G.$$

Then the sequence  $(\phi_{j,T})_j$  of  $L^\infty(G \times G)$  converges for the weak\* topology to the function  $\phi^{\text{HS}}: (s, t) \mapsto \phi(st^{-1})$ .

*Proof* : For any  $j$  and almost all  $s, t \in G$ , we have using a change of variables in the last equality

$$\begin{aligned} (4.36) \quad \phi_{j,T}(s, t) &\stackrel{(4.35)}{=} \tau_G(y_j \lambda_{s^{-1}} T(\lambda_s x_j \lambda_{t^{-1}}) \lambda_t) = \tau_G(\lambda_t \lambda(g_j) \lambda_{s^{-1}} T(\lambda_s \lambda(f_j) \lambda_{t^{-1}})) \\ &= \tau_G(\lambda(g_j(t^{-1} \cdot s)) M_\phi(\lambda(f_j(s^{-1} \cdot t))) = \tau_G(\lambda(g_j(t^{-1} \cdot s)) \lambda(\phi f_j(s^{-1} \cdot t))) \\ &\stackrel{(4.7)}{=} \int_G g_j(t^{-1} u^{-1} s) \phi(u) f_j(s^{-1} u t) d\mu_G(u) = \int_G \phi(s u t^{-1}) g_j(u^{-1}) f_j(u) d\mu_G(u). \end{aligned}$$

By Lemma 4.8 and (4.15), we deduce that  $\|M_{\phi_{j,T}}\|_{\text{cb}, S_G^p \rightarrow S_G^p} \leq C$  for any  $j$ . By [LaS, Remark 1.4], we infer that the sequence  $(\phi_{j,T})_j$  of functions is uniformly bounded in the Banach space  $L^\infty(G \times G)$ . Thus to check the claimed weak\* convergence, it suffices to test against a function  $h \in C_c(G \times G)$ . We suppose that  $\text{supp } f_j \rightarrow \{e\}$ . Since  $\int_G \check{g}_j f_j d\mu_G \stackrel{(4.7)}{=} \tau(\lambda(g_j) \lambda(f_j)) = \tau(x_j y_j) = 1$ , we have, using unimodularity in a change of variables, and with the notation

$$K_j = \text{supp } h \cup \{(s, t) : \exists u \in \text{supp } f_j : (su^{-1}, t) \in \text{supp } h\}$$

$$\begin{aligned} & \left| \int_{G \times G} (\phi_{j,T}(s, t) - \phi^{\text{HS}}(s, t)) h(s, t) \, d\mu_G(s) \, d\mu_G(t) \right| \\ & \stackrel{(4.36)}{=} \left| \int_G \int_G \int_G (\phi(st^{-1}) - \phi(st^{-1})) \check{g}_j(u) f_j(u) h(s, t) \, d\mu_G(u) \, d\mu_G(s) \, d\mu_G(t) \right| \\ & = \left| \int_G \int_G \int_G \phi(st^{-1}) \check{g}_j(u) f_j(u) (h(su^{-1}, t) - h(s, t)) \, d\mu_G(u) \, d\mu_G(s) \, d\mu_G(t) \right| \\ & \leq \sup \{ |h(su^{-1}, t) - h(s, t)| : u \in \text{supp } f_j, (s, t) \in \text{supp } h, (su^{-1}, t) \in \text{supp } h \} \cdot \\ & \quad \cdot \int_{K_j} \int_G |\phi(st^{-1})| f_j(u) \check{g}_j(u) \, d\mu_G(u) \, d\mu_G(s) \, d\mu_G(t) \\ & = \sup \{ |h(su^{-1}, t) - h(s, t)| : u \in \text{supp } f_j, (s, t) \in \text{supp } h, (su^{-1}, t) \in \text{supp } h \} \cdot \\ & \quad \cdot \int_{K_j} |\phi(st^{-1})| \, d\mu_G(s) \, d\mu_G(t) \xrightarrow{j} 0 \end{aligned}$$

since  $K_j$  is contained in a fixed compact, so that the last integral is uniformly bounded in  $j$ , and  $h$  was supposed to be continuous. We can use a similar reasoning if  $\text{supp } g_j \rightarrow \{e\}$ . ■

**Example 4.14** Let  $G$  be a second countable unimodular locally compact group. Let  $1 < p < \infty$ . Suppose that  $\frac{p}{p^*}$  is rational. That is,  $p = \frac{p}{p^*} + 1$  is rational, and thus also  $\frac{1}{p}$  and  $\frac{1}{p^*}$  are. Therefore there exist integers  $r, k, l \in \mathbb{N}$  such that  $\frac{1}{p} = \frac{k}{r}$  and  $\frac{1}{p^*} = \frac{l}{r}$ . Note that  $\frac{r}{p} = k$  and  $\frac{r}{p^*} = l$  are then integers. For any  $j$ , we use the function  $h_j \stackrel{\text{def}}{=} k_j^* * k_j$  for some positive function  $k_j \in C_c(G)$  with  $\text{supp } k_j \rightarrow \{e\}$ . We can suppose that  $\|\lambda(h_j)\|_{2r} = 1$ . Define  $x_j \stackrel{\text{def}}{=} \lambda((h_j^* * h_j))^k \stackrel{(4.6)}{=} |\lambda(h_j)|^{\frac{2r}{p}}$  and  $y_j \stackrel{\text{def}}{=} \lambda((h_j^* * h_j))^l$ . Note that these functions belong to the space  $C_c(G)$ . Then the sequences  $(x_j)$  and  $(y_j)$  satisfy the assumptions of Proposition 4.13. Note that these nets depend on  $p$ .

*Proof* : We have

$$\|x_j\|_p = \|\lambda((h_j^* * h_j))^k\|_p = \||\lambda(h_j)|^{\frac{2r}{p}}\|_p = \|\lambda(h_j)\|_{2r}^{\frac{2r}{p}} = 1^{\frac{2r}{p}} = 1$$

and similarly

$$\|y_j\|_{p^*} = \|\lambda((h_j^* * h_j))^l\|_{p^*} = \||\lambda(h_j)|^{\frac{2r}{p^*}}\|_{p^*} = \|\lambda(h_j)\|_{2r}^{\frac{2r}{p^*}} = 1.$$

Finally, we have

$$\tau_G(x_j y_j) = \tau_G(|\lambda(h_j)|^{\frac{2r}{p}} |\lambda(h_j)|^{\frac{2r}{p^*}}) = \tau_G(|\lambda(h_j)|^{2r}) = \|\lambda(h_j)\|_{2r}^{2r} = 1. \quad \blacksquare$$

## 4.7 Step 2: the case of finite-dimensional unimodular amenable locally compact groups

In the case where one strives for a complementation that satisfies a compatibility of the resulting Fourier multiplier  $P_G^p$  for different values of  $p$ , one needs in view of the last sentence of Example 4.14 to find a different choice of the sequences  $(x_j)$  and  $(y_j)$  defined in Remark 4.14.

This will be achieved in Corollary 4.23 in the case where the locally compact group  $G$  is finite-dimensional. Recall that three notions of dimension of a *suitable* topological space  $X$  exist (and defined e.g. in [Eng, Chap. 7]): the small inductive dimension, the large inductive dimension and the covering dimension. By [Eng, Theorem 7.3.3 p. 404], these notions coincide if  $X$  is metrizable and separable. Note that a second countable locally compact group  $G$  satisfies this property<sup>19</sup>. Indeed, Arhangel'skii and Pasynkov showed in [Ar1] and [Pas1] that these notions coincide for an *arbitrary* locally compact group  $G$ . We refer to the survey [ArM1] and to the books [Eng] and [Pea1] for more information.

Recall the definition of small inductive dimension. Let  $X$  be a regular topological space. We say that  $\text{ind } X = -1$  if  $X$  is empty. If  $n$  is a natural number, then we say that  $\text{ind } X \leq n$  if for every point  $x \in X$  and every neighborhood  $V$  of  $x$  in  $X$  there exists a open set  $U$  included in  $V$  such that  $x \in U$  and such that the boundary  $\partial U$  satisfies  $\text{ind } \partial U \leq n - 1$ . We say that  $\text{ind } X = n$  if  $\text{ind } X \leq n$  and  $\text{ind } X \leq n - 1$  does not hold. Finally, we say that  $\text{ind } X = \infty$  if the inequality  $\text{ind } X \leq n$  does not hold for any integer  $n$ .

If two regular topological spaces  $X$  and  $Y$  are homeomorphic then  $\text{ind } X = \text{ind } Y$ .

**Example 4.15** By [Eng, pages 363-380], a zero-dimensional space topological space  $X$  is a non-empty  $T_1$ -space which has a basis of open-and-closed subsets. By [Eng, Theorem 6.2.1], such a space is totally disconnected<sup>20</sup>, i.e. it does not contain any connected subspace of cardinality larger than 1. By [Eng, Theorem 6.2.10], a non-empty locally compact space  $X$  is zero-dimensional if and only if it is totally disconnected.

**Example 4.16** By [Stro1, Remark 39.5 (d)], a finite-dimensional locally compact group  $G$  is a Lie group if and only if it is locally connected<sup>21</sup>. See also [Pon1, Th. 70, p. 337] for the compact case.

We also need background on local isomorphisms.

**Local isomorphisms** Recall that two topological groups  $G$  and  $H$  are said to be locally isomorphic [Bou4, p. 224] if there exists open neighborhoods  $V, W$  of  $e_G$  and  $e_H$  and a homeomorphism  $f: V \rightarrow W$  satisfying  $f(xy) = f(x)f(y)$  for all  $x, y \in V$  such that  $xy \in V$  and if  $g$  is the mapping inverse to  $f$ , then for each pair of points  $x', y'$  of  $W$  such that  $x'y' \in W$ , we have  $g(x'y') = g(x')g(y')$ . We say that  $f$  is a local isomorphism of  $G$  with  $H$ . As observed in [Bou4, p. 224] if  $f$  is a local isomorphism of  $G$  with  $H$ , then every restriction of  $f$  to a neighbourhood  $V_0$  of the identity element of  $G$  such that  $V_0^2 \subset V$  is again a local isomorphism of  $G$  with  $H$ .

The following result is [Bou3, page 18-19] which describes the link between Haar measures and local isomorphisms.

**Lemma 4.17** *Let  $G$  and  $G'$  be locally isomorphic locally compact groups via a local homeomorphism  $f: V \rightarrow W$ . Consider a left Haar measures  $\mu_G$  of  $G$  and its restriction  $\mu_G^V$  on  $V$ . Then  $f(\mu_G^V)$  is the restriction of a unique left Haar measure on  $G'$ .*

We warn the reader that the property “unimodular” is not stable under local isomorphisms. See [Bou3, Exercise 5 page 78].

19. Such a group is metrizable by [CoH1, p. 20] and second-countable topological spaces are separable by [Eng, 1.3.8 Corollary]. See also [CoH1, Theorem 2.A.10].

20. In [Eng, page 360], this notion is called hereditarily disconnected.

21. This result is stronger than [Tao1, Ex. 1.6.9] which says without proof that a locally compact group  $G$  is a Lie group if and only if it is first-countable, locally connected and finite-dimensional. Moreover, the notion of dimension of [Tao1, Ex. 1.6.9] is different.

**Carnot-Carathéodory metrics on connected Lie groups** Let  $G$  be a connected Lie group with neutral element  $e$ . We consider a finite sequence  $X \stackrel{\text{def}}{=} (X_1, \dots, X_m)$  of left invariant vector fields whose the generated Lie algebra is the Lie algebra  $\mathfrak{g}$  of the group  $G$  such that the vectors  $X_1(e), \dots, X_m(e)$  are linearly independent. We say that it is a family of left-invariant Hörmander vector fields.

Let  $\gamma: [0, 1] \rightarrow G$  be an absolutely continuous path such that  $\dot{\gamma}(t)$  belongs to the subspace  $\text{span}\{X_1|_{\gamma(t)}, \dots, X_m|_{\gamma(t)}\}$  for almost every  $t \in [0, 1]$ . If  $\dot{\gamma}(t) = \sum_{k=1}^m \gamma_k(t) X_k|_{\gamma(t)}$  for almost every  $t \in [0, 1]$ , where  $\dot{\gamma}_k(t) \in \mathbb{R}$  and where each  $\dot{\gamma}_k$  is measurable, we can define the length of  $\gamma$  by

$$\ell(\gamma) \stackrel{\text{def}}{=} \int_0^1 \left( \sum_{k=1}^m |\dot{\gamma}_k(t)|^2 \right)^{1/2} dt$$

which belongs to  $[0, \infty]$ . For any  $s, s' \in G$  there exists such a path  $\gamma: [0, 1] \rightarrow G$  with finite length with  $\gamma(0) = s$  and  $\gamma(1) = s'$ . If  $s, s' \in G$  then we define the Carnot-Carathéodory distance  $\text{dist}_{\text{CC}}(s, s')$  between  $s$  and  $s'$  to be the infimum of the length of all such paths with  $\gamma(0) = s$  and  $\gamma(1) = s'$ :

$$(4.37) \quad \text{dist}_{\text{CC}}(s, s') \stackrel{\text{def}}{=} \inf_{\gamma(0)=s, \gamma(1)=s'} \ell(\gamma).$$

See [VSCC92, p. 39] and [DtER03, p. 22]. Then it is known that  $\text{dist}_{\text{CC}}$  is a left invariant metric on  $G$  inducing the same topology that the one of  $G$ , see [VSCC92, Proposition III.4.1] and [DtER03, p. 22-23].

For any  $s \in G$  and any  $r > 0$ , we denote by  $B(s, r)$  the open ball with respect to the Carnot-Carathéodory metric centered at  $s$  and of radius  $r$ . By [VSCC92, p. 124] there exist  $c_1, c_2 > 0$  and  $d \in \mathbb{N}$  such that for all  $r \in (0, 1]$  we have

$$(4.38) \quad c_1 r^d \leq \mu_G(B(e, r)) \leq c_2 r^d$$

where  $\mu_G$  is a left Haar measure of  $G$ . The integer  $d$  is sometimes called the local dimension of  $G$  with respect to  $(X_1, \dots, X_m)$ . According to (4.38), we have  $\mu_G(B(e, r)) \cong r^d$  for any  $r \in (0, 1]$ . We infer that there exists  $c > 0$

$$(4.39) \quad \mu_G(B(e, 2r)) \leq c \mu_G(B(e, r)), \quad r \in (0, 1/2].$$

The smallest constant which satisfies this inequality is the “local” doubling constant of  $G$  with respect to  $\mu_G$  and  $\text{dist}_{\text{CC}}$ .

**Example 4.18** It is known that the local doubling constant of the Heisenberg group is equal to 16.

A Borel regular measure  $\mu$  on a metric space  $(X, d)$  is called a doubling measure [HKST15, p. 76] if every ball in  $X$  has positive and finite measure and there exists a constant  $c \geq 1$  such that

$$(4.40) \quad \mu(B(x, 2r)) \leq c \mu(B(x, r)), \quad x \in X, r > 0.$$

We call the triple  $(X, d, \mu)$  a doubling metric measure space if  $\mu$  is a doubling measure on  $X$ .

Here, we list several typical examples satisfying the doubling condition. [BEM13, Proposition 2.4 p. 199] a Lie group  $G$  equipped with a left Haar measure  $\mu_G$  has polynomial growth if and only if the metric measure space  $(G, \text{dist}_{\text{CC}}, \mu_G)$  has the doubling property.

Let  $G$  be a compactly generated group of polynomial growth. Consider a symmetric compact generating subset  $V$ . The word metric defined by

$$d(s, t) \stackrel{\text{def}}{=} \inf\{n \in \mathbb{N} : s^{-1}t \in V^n\}, \quad s, t \in G$$

satisfies (4.40).

Let  $G$  be a compactly generated group of polynomial growth and let  $d$  be a metric on  $G$ . If  $d$  is invariant under a cocompact subgroup of  $G$  and if  $d$  satisfies a weak kind of the existence of geodesics axiom (see [Bre14, Definition 4.1]), then  $(G, d)$  satisfies (4.40). We refer to [Nev06, Sections 4 and 5] for more examples.

We will use the weaker notion of doubling measure for small balls replacing (4.40) by

$$(4.41) \quad \mu(B(x, 2r)) \leq c\mu(B(x, r)), \quad x \in X, r \in (0, 1/2].$$

Note we can use the following lemma with a Lie group by considering the connected component of  $G$  equipped with the Carnot-Carathéodory distance (4.37) with respect to some fixed sequence of left invariant vector fields. Next, we consider the left invariant metric on the Lie group  $G$ . Recall that the connected component of a Lie group is second countable.

**Lemma 4.19** *Let  $G$  be a second countable locally compact group equipped with a left invariant distance and a doubling left Haar measure  $\mu_G$  for small balls. There exists a sequence  $(B_j)$  of open ball  $B_j \stackrel{\text{def}}{=} B(e, r_j)$  with decreasing radius  $r_j \rightarrow 0$  satisfying*

$$(4.42) \quad \mu_G(B_j)^3 \leq c^3 \int_{B_j} \mu_G(B_j \cap sB_j)^2 d\mu_G(s), \quad j \in \mathbb{N}$$

where  $c$  is a constant satisfying (4.41).

*Proof* : Let  $1 > \varepsilon > 0$ . For any integer  $j$ , we put  $B_j \stackrel{\text{def}}{=} B(e, \frac{\varepsilon}{j})$ . We have

$$(4.43) \quad \mu_G(B_j) = \mu_G\left(B\left(e, \frac{\varepsilon}{j}\right)\right) \stackrel{(4.41)}{\leq} c \mu_G\left(B\left(e, \frac{\varepsilon}{2j}\right)\right) = c \mu_G(B_{2j}).$$

For any element  $s$  of  $B_{2j} = B(e, \frac{\varepsilon}{2j})$ , we will show that

$$(4.44) \quad B\left(e, \frac{\varepsilon}{2j}\right) \subset B\left(e, \frac{\varepsilon}{j}\right) \cap sB\left(e, \frac{\varepsilon}{j}\right), \quad \text{i.e.} \quad B_{2j} \subset B_j \cap sB_j.$$

Indeed, if  $r \in G$  satisfies  $\text{dist}_{\text{CC}}(e, r) < \frac{\varepsilon}{2j}$  we have obviously  $r \in B(e, \frac{\varepsilon}{j})$  and by left invariance of the distance we obtain

$$\text{dist}_{\text{CC}}(e, s^{-1}r) = \text{dist}_{\text{CC}}(s, r) \leq \text{dist}_{\text{CC}}(s, e) + \text{dist}_{\text{CC}}(e, r) < \frac{\varepsilon}{2j} + \frac{\varepsilon}{2j} = \frac{\varepsilon}{j}.$$

So  $s^{-1}r \in B(e, \frac{\varepsilon}{j})$  and consequently  $r \in sB(e, \frac{\varepsilon}{j})$ . For any  $j$ , we deduce that

$$\begin{aligned} c^3 \int_{B_j} \mu_G^2(B_j \cap sB_j) d\mu_G(s) &\geq c^3 \int_{B_{2j}} \mu_G^2(B_j \cap sB_j) d\mu_G(s) \\ &\stackrel{(4.44)}{\geq} c^3 \int_{B_{2j}} \mu_G^2(B_{2j}) d\mu_G(s) = c^3 \mu_G^3(B_{2j}) \stackrel{(4.43)}{\geq} \mu_G^3(B_j). \end{aligned}$$

■



**Remark 4.20** Recall that by [CoH1, Theorem 2.A.10 p. 15], a locally compact space  $X$  is second-countable if and only if  $X$  is metrizable and separable. Since a doubling metric measure space  $X$  is separable as a metric space by [HKST15, Lemma 3.3.30], so the assumption "G is second-countable" is useless in Lemma 4.19 if the measure is doubling.

We continue with the totally disconnected case.

**Lemma 4.21** *Let  $G$  be a second countable totally disconnected locally compact group equipped with a left Haar measure  $\mu_G$ . Then there exists a basis  $(K_j)$  of symmetric open compact neighborhoods  $K_j$  of  $e$  such that*

$$(4.45) \quad \mu_G(K_j)^3 = \int_{K_j} \mu_G(K_j \cap sK_j)^2 d\mu_G(s), \quad j \in \mathbb{N}.$$

*Proof* : According to Van Dantzig's theorem [HR, (7.7) Theorem p. 62],  $G$  admits a basis  $(K_j)$  of open compact subgroups. Clearly, the  $K_j$  are symmetric being groups, neighborhoods being open, and of finite measure being compact. Since we have assumed  $G$  to be second countable, one can easily see from the proof of [HR, (7.7) Theorem p. 62] that  $(K_j)$  can be chosen to be a sequence. Since  $K_j$  is a subgroup, we have  $K_j \cap sK_j = \emptyset$  for  $s \notin K_j$  and  $K_j \cap sK_j = K_j$  for  $s \in K_j$ . Thus,

$$\int_{K_j} \mu_G(K_j \cap sK_j)^2 d\mu_G(s) = \int_{K_j} \mu_G(K_j)^2 d\mu_G(s) = \mu_G(K_j)^3.$$

■

Using a version of Iwasawa's local splitting theorem, we are now able to obtain a result for *finite-dimensional* locally compact groups.

**Lemma 4.22** *Let  $G$  be a second countable locally compact group equipped with a left Haar measure  $\mu_G$ . Suppose that  $G$  is finite-dimensional. Then there exists a basis  $(V_j)$  of symmetric open neighborhoods  $V_j$  of  $e$  and a constant  $c > 0$  such that*

$$(4.46) \quad \mu_G(V_j)^3 \leq c^3 \int_{V_j} \mu_G(V_j \cap sV_j)^2 d\mu_G(s), \quad j \in \mathbb{N}.$$

*Proof* : We denote by  $n$  the dimension of  $G$ . According to [Glu1, Theorem B p. 92], [Tao1, Ex 1.6.8 p.122], [Stro1, Remark 39.5 (a) p. 283] and [MZ, p. 237]<sup>22</sup> a  $n$ -dimensional first countable locally compact group is locally isomorphic to the product of a compact totally disconnected group  $K$  and a Lie group  $L$  of dimension  $n$ . So there exists a neighborhood  $V$  of  $e_G$  which is homeomorphic the direct product  $W \times U$  of a neighborhood  $W$  of the neutral element  $e_K$  and an open neighborhood  $U$  of  $e_L$ . By restriction, we can suppose that  $U$  is connected and, by considering the connected component of  $L$ , that  $L$  is connected. Finally, we identify  $V$  with  $W \times U$ .

Next, we consider the left invariant metric on the Lie group  $L$  given by the Carnot-Caratheodory distance (4.37) with respect to some fixed sequence of left vector fields.

By Lemma 4.17, we can choose left Haar measures  $\mu_K, \mu_L$  on the groups  $K$  and  $L$  such that

$$(4.47) \quad \mu_G(A \times B) = \mu_K(A)\mu_L(B), \quad A \subset W, B \subset U.$$

<sup>22</sup>. See also [Pon1, Th. 70, p. 337] for the case of compact groups.

Let  $(K_j)$  a basis of neighbourhoods consisting as in Lemma 4.21 and  $(B_j)$  as in Lemma 4.42. For any  $j$ , we put

$$(4.48) \quad V_j \stackrel{\text{def}}{=} K_j \times B_j$$

and we can suppose that  $K_j \subset W$  for any  $j$  and that  $B_j \subset U$ . Being a subgroup,  $K_j$  is symmetric and  $B_j$  is also symmetric, being a ball with respect to a left invariant metric<sup>23</sup>. We conclude that  $V_j$  is symmetric as well.

In view of the previous local product structure of the Haar measure from (4.47), we deduce that for any  $j$

$$\begin{aligned} c^3 \int_{V_j} \mu_G^2(V_j \cap sV_j) d\mu_G(s) &= c^3 \int_{K_j \times B_j} \mu_G^2(V_j \cap sV_j) d\mu_{K \times L}(s) \\ &\stackrel{(4.48)}{=} c^3 \int_{K_j \times B_j} \mu_G^2((K_j \times B_j) \cap (r, t)(K_j \times B_j)) d\mu_K(r) d\mu_L(t) \\ &\stackrel{(4.47)}{=} c^3 \int_{K_j \times B_j} \mu_K^2(K_j \cap rK_j) \mu_L^2(B_j \cap tB_j) d\mu_K(r) d\mu_L(t) \\ &= c^3 \int_{K_j} \mu_K^2(K_j \cap rK_j) d\mu_K(r) \int_{B_j} \mu_L^2(B_j \cap tB_j) d\mu_L(t) \\ &\stackrel{(4.42)(4.45)}{\geq} \mu_K^3(K_j) \mu_G^3(B_j) \stackrel{(4.47)}{=} \mu_G^3(K_j \times B_j) \stackrel{(4.48)}{=} \mu_G^3(V_j). \end{aligned}$$

■

**Corollary 4.23** *Let  $G$  be an second countable unimodular amenable locally compact group. Suppose that  $G$  is finite-dimensional. Let  $1 \leq p \leq \infty$  and let  $T = M_\phi: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))$  be a completely bounded Fourier multiplier. Let  $(V_j)$  be the sequence of nested neighborhoods as in Lemma 4.22. Moreover, put*

$$(4.49) \quad f_j \stackrel{\text{def}}{=} 1_{V_j} * 1_{V_j}, \quad x_j \stackrel{\text{def}}{=} a_j \lambda(f_j) \quad \text{and} \quad y_j \stackrel{\text{def}}{=} b_j \lambda(f_j)$$

with

$$(4.50) \quad a_j \stackrel{\text{def}}{=} \|\lambda(f_j)\|_{p^*} \|\lambda(f_j)\|_2^{-2} \quad \text{and} \quad b_j \stackrel{\text{def}}{=} \|\lambda(f_j)\|_{p^*}^{-1}.$$

Then the sequences  $(x_j)$  and  $(y_j)$  satisfy the assumptions from Proposition 4.13. More precisely, we have  $\|x_j\|_p \leq c^3$ ,  $\|y_j\|_{p^*} = 1$ .

*Proof* : Consider any measurable subset  $V$  of  $G$  with  $\mu_G(V) \in (0, \infty)$ . Using Young's inequality in the second inequality, we have<sup>24</sup>

$$\begin{aligned} \|\lambda(1_{V^{-1}} * 1_V)\|_p &\stackrel{(4.6)}{=} \|\lambda(1_V) * \lambda(1_V)\|_p = \|\lambda(1_V)\|_p^2 = \|\lambda(1_V)\|_{2p}^2 \leq \|\lambda(1_V)\|_{p^*}^{\frac{2}{p}} \|\lambda(1_V)\|_2^{\frac{2}{p}} \\ &= \|\lambda(1_V)\|_{L^2(G) \rightarrow L^2(G)}^{\frac{2}{p^*}} \|\lambda(1_V)\|_2^{\frac{2}{p}} \leq \mu_G(V)^{\frac{2}{p^*}} \cdot \mu_G(V)^{\frac{1}{p}} = \mu_G(V)^{1 + \frac{1}{p^*}}. \end{aligned}$$

23. Note that  $\text{dist}_{\text{CC}}(e, s) = \text{dist}_{\text{CC}}(s^{-1}, e)$ .

24. In the first inequality, we use  $\|\cdot\|_{(X, Y)_\theta} \leq \|\cdot\|_X^{1-\theta} \|\cdot\|_Y^\theta$  and  $L^{2p} = (L^\infty, L^2)_{\frac{2}{p}}$ .

Taking  $V = V_j$  and using the previous calculation also for  $p^*$  in place of  $p$ , we deduce with Lemma 4.22 that

$$(4.51) \quad \begin{aligned} \|\lambda(1_{V_j} * 1_{V_j})\|_p \cdot \|\lambda(1_{V_j} * 1_{V_j})\|_{p^*} &\leq \mu_G(V_j)^{1+\frac{1}{p^*}+1+\frac{1}{p}} \\ &= \mu_G(V_j)^3 \stackrel{(4.46)}{\leq} c^3 \int_{V_j} \mu_G(V_j \cap sV_j)^2 d\mu_G(s). \end{aligned}$$

On the other hand, using  $V_j = V_j^{-1}$ , we obtain

$$(4.52) \quad \begin{aligned} \|\lambda(1_{V_j} * 1_{V_j})\|_2^2 &\stackrel{(4.5)}{=} \int_G \left| \int_G 1_{V_j}(t) 1_{V_j}(t^{-1}s) d\mu_G(t) \right|^2 d\mu_G(s) \\ &= \int_G \left| \int_G 1_{V_j}(t) 1_{V_j s^{-1}}(t^{-1}) d\mu_G(t) \right|^2 d\mu_G(s) = \int_G \left| \int_G 1_{V_j}(t) 1_{sV_j}(t) d\mu_G(t) \right|^2 d\mu_G(s) \\ &= \int_G \mu_G^2(V_j \cap sV_j) d\mu_G(s). \end{aligned}$$

Combining (4.51) and (4.52), we see that

$$(4.53) \quad \|\lambda(1_{V_j} * 1_{V_j})\|_p \cdot \|\lambda(1_{V_j} * 1_{V_j})\|_{p^*} \leq c^3 \|\lambda(1_{V_j} * 1_{V_j})\|_2^2.$$

Note that with the choice of  $a_j$  and  $b_j$ , we finally obtain

$$\|x_j\|_p \stackrel{(4.49)}{=} \|a_j \lambda(f_j)\|_p \stackrel{(4.50)}{=} \|\lambda(f_j)\|_{p^*} \|\lambda(f_j)\|_2^{-2} \|\lambda(f_j)\|_p \stackrel{(4.49)(4.53)}{\leq} c^3$$

and  $\|y_j\|_{p^*} \stackrel{(4.49)}{=} \|b_j \lambda(f_j)\|_{p^*} = 1$ , as well as  $\tau_G(x_j y_j) \stackrel{(4.50)}{=} \|\lambda(f_j)\|_2^{-2} \tau_G(\lambda(f_j)^2) = 1$ .  $\blacksquare$

**Remark 4.24** Suppose  $1 \leq p \leq \infty$  and consider a second countable totally disconnected locally compact group  $G$ . Then the sequences  $(x_j)$  and  $(y_j)$  from Corollary 4.23, with respect to the choice of  $V_j = K_j$ , where  $K_j$  is defined in Lemma 4.21, satisfy  $\|x_j\|_p \leq 1$  and  $\|y_j\|_{p^*} = 1$ .

### 4.8 Step 3: the projection $Q: \mathfrak{M}_G^{p,\text{cb}} \rightarrow \mathfrak{M}_G^{p,\text{cb,HS}}$ on the space of Herz-Schur multipliers

The first part of the following result says that the unit ball of the space  $\mathfrak{M}_\Omega^p$  of measurable Schur multipliers is closed for the weak\* topology of  $L^\infty(\Omega \times \Omega)$ .

**Lemma 4.25** *Let  $\Omega$  be a  $\sigma$ -finite measure space. Suppose  $1 \leq p \leq \infty$ . Let  $(M_{\phi_j})$  be a bounded net of measurable Schur multipliers on  $S_\Omega^p$  and suppose that  $\phi$  is an element of  $L^\infty(\Omega \times \Omega)$  such that  $(\phi_j)$  converges to  $\phi$  for the weak\* topology of  $L^\infty(\Omega \times \Omega)$ . Then  $\phi$  induces a measurable Schur multiplier on  $S_\Omega^p$ . Moreover, the net  $(M_{\phi_j})$  converges to  $M_\phi$  in  $\mathfrak{M}_\Omega^p$  for the weak operator topology of  $B(S_\Omega^p)$  and*

$$\|M_\phi\|_{S_\Omega^p \rightarrow S_\Omega^p} \leq \liminf_{j \rightarrow \infty} \|M_{\phi_j}\|_{S_\Omega^p \rightarrow S_\Omega^p}.$$

*A similar statement is true by replacing ‘‘bounded’’ by ‘‘completely bounded’’ and the norms by  $\|\cdot\|_{\text{cb}, S_\Omega^p \rightarrow S_\Omega^p}$ .*

*Proof* : For any  $f, g \in L^2(\Omega \times \Omega)$ , note that we have  $f\check{g} \in L^1(\Omega \times \Omega)$ . Consider some  $f, g \in L^2(\Omega \times \Omega)$  such that  $K_f \in S_\Omega^p, K_g \in S_\Omega^{p*}$ . For any  $j$ , we have

$$\begin{aligned} \left| \int_{\Omega \times \Omega} \phi_j f \check{g} \right| &\stackrel{(4.2)}{=} \left| \langle M_{\phi_j}(K_f), K_g \rangle_{S_\Omega^p, S_\Omega^{p*}} \right| \leq \|M_{\phi_j}(K_f)\|_{S_\Omega^p} \|K_g\|_{S_\Omega^{p*}} \\ &\leq \|M_{\phi_j}\|_{S_\Omega^p \rightarrow S_\Omega^p} \|K_f\|_{S_\Omega^p} \|K_g\|_{S_\Omega^{p*}}. \end{aligned}$$

Passing to the limit, we obtain

$$\left| \langle K_{\phi_f}, K_g \rangle_{S_\Omega^p, S_\Omega^{p*}} \right| \stackrel{(4.2)}{=} \left| \int_{\Omega \times \Omega} \phi f \check{g} \right| \leq \liminf_{j \rightarrow \infty} \|M_{\phi_j}\|_{S_\Omega^p \rightarrow S_\Omega^p} \|K_f\|_{S_\Omega^p} \|K_g\|_{S_\Omega^{p*}}.$$

By density, we conclude that  $\phi$  induces a bounded measurable Schur multiplier on  $S_\Omega^p$  with the estimate on the norm. Using again the weak\* convergence of the net  $(\phi_j)$ , we see that for any  $f, g \in L^2(\Omega \times \Omega)$  such that  $K_f \in S_\Omega^p$  and  $K_g \in S_\Omega^{p*}$

$$\begin{aligned} \text{Tr}((M_\phi - M_{\phi_j})(K_f)K_g) &= \text{Tr}(K_{(\phi - \phi_j)f}K_g) \stackrel{(4.2)}{=} \iint_{\Omega \times \Omega} (\phi - \phi_j) f \check{g} \\ &= \langle \phi - \phi_j, f \check{g} \rangle_{L^\infty(\Omega \times \Omega), L^1(\Omega \times \Omega)} \xrightarrow{j} 0. \end{aligned}$$

By density, using a  $\frac{\varepsilon}{4}$ -argument and the boundedness of the net, we conclude<sup>25</sup> the proof.

Now, we prove the last sentence, it suffices to show that  $\phi$  induces a completely bounded Schur multiplier. For any  $f_{kl}, g_{kl} \in L^2(\Omega \times \Omega)$  where  $1 \leq k, l \leq N$ , we have for  $f_{kl}\check{g}_{kl} \in L^1(\Omega \times \Omega)$ . For any  $j$  by [Pis5, Theorem 4.7], we infer that

$$\left| \langle [M_{\phi_j}(K_{f_{kl}})], [K_{g_{kl}}] \rangle_{M_N(S_\Omega^p), S_N^1(S_\Omega^{p*})} \right| \leq \|M_{\phi_j}\|_{\text{cb}, S_\Omega^p \rightarrow S_\Omega^p} \|[K_{f_{kl}}]\|_{M_N(S_\Omega^p)} \|[K_{g_{kl}}]\|_{S_N^1(S_\Omega^{p*})},$$

that is,

$$\left| \sum_{k,l=1}^N \int_{\Omega \times \Omega} \phi_j f_{kl} \check{g}_{kl} \right| \leq \|M_{\phi_j}\|_{\text{cb}, S_\Omega^p \rightarrow S_\Omega^p} \|[K_{f_{kl}}]\|_{M_N(S_\Omega^p)} \|[K_{g_{kl}}]\|_{S_N^1(S_\Omega^{p*})}.$$

Passing to the limit, we obtain

$$\left| \sum_{k,l=1}^N \int_{\Omega \times \Omega} \phi f_{kl} \check{g}_{kl} \right| \leq \liminf_{j \rightarrow \infty} \|M_{\phi_j}\|_{\text{cb}, S_\Omega^p \rightarrow S_\Omega^p} \|[K_{f_{kl}}]\|_{M_N(S_\Omega^p)} \|[K_{g_{kl}}]\|_{S_N^1(S_\Omega^{p*})}.$$

We deduce that  $\phi$  induces a completely bounded Schur multiplier on  $S_\Omega^p$  with the suitable estimate on the completely bounded norm.  $\blacksquare$

If  $1 \leq p < \infty$ , note that  $S_\Omega^p$  is a dual space. So the Banach space  $\text{CB}(S_\Omega^p)$  is also a dual space with predual  $S_\Omega^p \widehat{\otimes} S_\Omega^{p*}$ , where  $\widehat{\otimes}$  denotes the operator space projective tensor product and the duality bracket is given by

$$(4.54) \quad \langle T, x \otimes y \rangle_{\text{CB}(S_\Omega^p), S_\Omega^p \widehat{\otimes} S_\Omega^{p*}} = \langle T(x), y \rangle_{S_\Omega^p, S_\Omega^{p*}}.$$

25. More precisely, if  $X$  is a Banach space, if  $E_1$  is dense subset of  $X$ , if  $E_2$  is a dense subset of  $X^*$  and if  $(T_j)$  is a bounded net of  $\text{B}(X)$  with an element  $T$  of  $\text{B}(X)$  such that  $\langle T_j(x), x^* \rangle \xrightarrow{i \rightarrow +\infty} \langle T(x), x^* \rangle$  for any  $x \in E_1$  and any  $x^* \in E_2$ , then the net  $(T_j)$  converges to  $T$  for the weak operator topology of  $\text{B}(X)$ .

**Lemma 4.26** *Let  $\Omega$  be a  $\sigma$ -finite measure space.*

1. *Let  $1 \leq p < \infty$ . Then  $\mathfrak{M}_\Omega^{p,\text{cb}}$  is weak\* closed in  $\text{CB}(S_\Omega^p)$  and  $\mathfrak{M}_\Omega^p$  is weak\* closed in  $\text{B}(S_\Omega^p)$ .*
2. *The space  $\mathfrak{M}_\Omega^{\infty,\text{cb}}$  is weak\* closed in  $\text{CB}(S_\Omega^\infty, \text{B}(L^2(\Omega)))$ .*

*Proof* : By the Banach-Dieudonné theorem [Hol1, page 154], it suffices to show that the closed unit ball of the space  $\mathfrak{M}_\Omega^{p,\text{cb}}$  is weak\* closed in  $\text{CB}(S_\Omega^p)$ . Let  $(M_{\phi_j})$  be a net in that unit ball converging for the weak\* topology to some completely bounded map  $T: S_\Omega^p \rightarrow S_\Omega^p$ . By [LaS, Remark 1.4], we have for any  $j$  the inequality

$$\|\phi_j\|_{L^\infty(\Omega \times \Omega)} \leq \|M_{\phi_j}\|_{\text{cb}, S_\Omega^p \rightarrow S_\Omega^p} \leq 1.$$

By Banach-Alaoglu theorem there exists a subnet of  $(\phi_j)$  converging for the weak\* topology to some  $\phi \in L^\infty(\Omega \times \Omega)$ . It remains to show that  $T = M_\phi$ . By (4.54), we have  $\langle M_{\phi_j}(x), y \rangle \xrightarrow{j} \langle T(x), y \rangle$  for any  $x \in S_\Omega^p$  and any  $y \in S_\Omega^{p*}$ . That means that the net  $(M_{\phi_j})$  converges to  $T$  for the weak operator topology. By Lemma 4.25, the net  $(M_{\phi_j})$  converges to  $M_\phi$ . We conclude by uniqueness of the limit that  $T = M_\phi$ .

The statement on the space  $\mathfrak{M}_\Omega^p$  can be proved in a similar manner, using the predual  $S_\Omega^p \hat{\otimes} S_\Omega^{p*}$  of  $\text{B}(S_\Omega^p)$  where  $\hat{\otimes}$  denotes the Banach space projective tensor product. The second point is also similar.  $\blacksquare$

The following is essentially folklore. We sketch a proof since it is very important for us.

**Proposition 4.27** *Let  $G$  be an amenable locally compact group. Suppose  $1 \leq p \leq \infty$ . Then there exists a contractive projection  $Q: \mathfrak{M}_G^{p,\text{cb}} \rightarrow \mathfrak{M}_G^{p,\text{cb}}$  onto the space of completely bounded Herz-Schur multipliers  $\mathfrak{M}_G^{p,\text{cb,HS}}$  over  $S_G^p$  ( $\text{B}(L^2(G))$  if  $p = \infty$ ) which preserves the complete positivity. Moreover, the obtained projections are compatible for all different values of  $1 \leq p \leq \infty$ .*

*Proof* : Let  $(F_j)$  be a Følner net in  $G$  provided by the amenability of  $G$ . For any  $K_f \in S_G^2 \cap S_G^p$ , the map  $G \rightarrow S_G^p$ ,  $R \mapsto \text{Ad}(\rho_r)(K_f)$  is continuous since the composition of operators is strongly continuous on bounded sets by [EFHN, Proposition C.19]. Similarly for any  $M_\phi \in \mathfrak{M}_G^{p,\text{cb}}$ , the map  $G \rightarrow S_G^p$ ,  $r \mapsto [\text{Ad}(\rho_r^*)M_\phi \text{Ad}(\rho_r)](K_f)$  is also continuous, hence Bochner integrable on the compact  $F_j$ . Now, for any  $K_f \in S_G^2 \cap S_G^p$  and any  $M_\phi \in \mathfrak{M}_G^{p,\text{cb}}$ , put

$$(4.55) \quad Q_j(M_\phi)(K_f) = \frac{1}{\mu_G(F_j)} \int_{F_j} [\text{Ad}(\rho_r^*)M_\phi \text{Ad}(\rho_r)](K_f) d\mu_G(r).$$

For any  $K_f \in S_G^2 \cap S_G^p$  and any Schur multiplier  $M_\phi \in \mathfrak{M}_G^{p,\text{cb}}$ , we have

$$\begin{aligned} \|Q_j(M_\phi)(K_f)\|_{S_G^p} &= \frac{1}{\mu_G(F_j)} \left\| \int_{F_j} [\text{Ad}(\rho_r^*)M_\phi \text{Ad}(\rho_r)](K_f) d\mu_G(r) \right\|_{S_G^p} \\ &\leq \frac{1}{\mu_G(F_j)} \int_{F_j} \|[\text{Ad}(\rho_r^*)M_\phi \text{Ad}(\rho_r)](K_f)\|_{S_G^p} d\mu_G(r) \leq \|M_\phi\|_{S_G^p \rightarrow S_G^p} \|K_f\|_{S_G^p}. \end{aligned}$$

A similar argument shows that  $\|Q_j(M_\phi)\|_{\text{cb}, S_G^p \rightarrow S_G^p} \leq \|M_\phi\|_{\text{cb}, S_G^p \rightarrow S_G^p}$ . Consequently, we have a well-defined contractive map  $Q_j: \mathfrak{M}_G^{p,\text{cb}} \rightarrow \text{CB}(S_G^p)$ ,  $M_\phi \mapsto \frac{1}{\mu_G(F_j)} \int_{F_j} \text{Ad}(\rho_r^*)M_\phi \text{Ad}(\rho_r) d\mu_G(r)$ . If  $M_\phi$  is completely positive then  $[\text{Ad}(\rho_r^*)M_\phi \text{Ad}(\rho_r)]$  is also completely positive. Thus the map

$Q_j$  preserves complete positivity. It is easy to check that  $Q_j(M_\phi)$  is a Schur multiplier with symbol

$$(4.56) \quad \phi_j(s, t) \stackrel{\text{def}}{=} \frac{1}{\mu_G(F_j)} \int_{F_j} \phi(sr, tr) d\mu_G(r)$$

(Gelfand integral in  $L^\infty(G \times G)$ ).

We continue with the case  $1 \leq p < \infty$ . Since  $\mathfrak{M}_G^{p, \text{cb}}$  is weak\* closed in  $\text{CB}(S_\Omega^p)$ , the space  $\mathfrak{M}_G^{p, \text{cb}}$  is a dual space. Hence  $\text{B}(\mathfrak{M}_G^{p, \text{cb}}, \mathfrak{M}_G^{p, \text{cb}})$  is a dual space. By Banach-Alaoglu's theorem, the uniformly bounded net  $(Q_j)$  admits a weak\* accumulation point that we denote by  $Q$  which obviously a contraction. So we can suppose that  $Q_j \rightarrow Q$  for the weak\* topology. So, for each completely bounded Schur multiplier  $M_\phi: S_G^p \rightarrow S_G^p$  this implies that  $Q(M_\phi) = \lim_j Q_j(M_\phi)$  in the weak operator topology. Recall that the weak\* topology on  $\text{CB}(S_G^p)$  coincides on bounded subsets with the weak operator topology. Since  $Q_j(M_\phi)$  belongs to  $\mathfrak{M}_G^{p, \text{cb}}$  and since the latter space is weak\* closed in  $\text{CB}(S_G^p)$  according to Lemma 4.26, we obtain that  $Q(M_\phi)$  also belongs to  $\mathfrak{M}_G^{p, \text{cb}}$ . Since each map  $Q_j$  preserves complete positivity, by [ArK1, Lemma 2.10 2.], the map  $Q$  also preserves complete positivity.

For any completely bounded Schur multiplier  $M_\phi: S_G^p \rightarrow S_G^p$ , it remains to show that  $Q(M_\phi)$  is in addition a Herz-Schur multiplier. That is, for any  $r_0 \in G$  we have to show that  $\lim_j M_{\phi_j(sr_0, tr_0)} = \lim_j M_{\phi_j(s, t)}$ . Fix some  $r_0 \in G$  and  $j$ . Using the Følner condition in the last line, we have

$$\begin{aligned} & \left\| M_{\phi_j(s, t)} - M_{\phi_j(sr_0, tr_0)} \right\|_{\text{cb}, S_G^p \rightarrow S_G^p} \\ & \stackrel{(4.56)}{=} \left\| \frac{1}{\mu_G(F_j)} \int_{F_j} M_{\phi(sr, tr)} d\mu_G(r) - \frac{1}{\mu(F_j)} \int_{F_j} M_{\phi(sr_0u, tr_0u)} d\mu_G(u) \right\|_{\text{cb}, S_G^p \rightarrow S_G^p} \\ & = \frac{1}{\mu_G(F_j)} \left\| \int_{F_j} M_{\phi(sr, tr)} d\mu_G(r) - \int_{r_0^{-1}F_j} M_{\phi(sr, tr)} d\mu_G(r) \right\|_{\text{cb}, S_G^p \rightarrow S_G^p} \\ & \leq \frac{1}{\mu_G(F_j)} \int_{F_j \Delta r_0 F_j} \|M_{\phi(sr, tr)}\|_{\text{cb}, S_G^p \rightarrow S_G^p} d\mu_G(r) \\ & = \frac{\mu_G(F_j \Delta r_0 F_j)}{\mu_G(F_j)} \|M_\phi\|_{\text{cb}, S_G^p \rightarrow S_G^p} \xrightarrow{j \rightarrow \infty} 0. \end{aligned}$$

Using the weak\* lower semicontinuity of the norm [Meg1, Th. 2.6.14], we infer that

$$\left\| Q(M_\phi) - Q(M_{\phi(\cdot r_0, \cdot r_0)}) \right\|_{\text{cb}, S_G^p \rightarrow S_G^p} \leq \liminf_j \left\| M_{\phi_j(s, t)} - M_{\phi_j(sr_0, tr_0)} \right\|_{\text{cb}, S_G^p \rightarrow S_G^p} = 0.$$

Finally, it is easy to see that  $Q(M_{\phi(\cdot r_0, \cdot r_0)}) = Q(M_\phi)_{(\cdot r_0, \cdot r_0)}$ . The case  $p = \infty$  is similar.

In order to obtain the mappings  $Q^{(p)}: \mathfrak{M}_G^{p, \text{cb}} \rightarrow \mathfrak{M}_G^{p, \text{cb}}$  compatible for different values of  $1 \leq p \leq \infty$ , it suffices to observe that we can choose the indices  $j'$  in the converging subnet  $Q_{j'}^{(p)}$  independent of  $p$ , in the same manner as done in the proof of Corollary 4.35 below by means of a Tychonoff argument. The proof is complete.  $\blacksquare$

**Remark 4.28** We have a similar result for spaces of bounded Schur multipliers.

**Remark 4.29** If  $G$  is compact, the proof is more simple. We do not need to use an approximation procedure.

**Remark 4.30** Consider a locally compact group  $G$  and  $1 \leq p \leq \infty$ . To know if the amenability of  $G$  is characterized by the property of Proposition 4.27 is a well-known open intriguing question.

## 4.9 Steps 1-3 put together: theorems of complementation

Let  $G$  be a locally compact group. Note that  $B(\text{CB}(\text{VN}(G)), \text{CB}(S_G^\infty, B(L^2(G))))$  is a dual Banach space and admits the predual

$$(4.57) \quad \text{CB}(\text{VN}(G)) \hat{\otimes} (S_G^\infty \hat{\otimes} S_G^1),$$

where  $\hat{\otimes}$  denotes the Banach space projective tensor product and where  $\widehat{\otimes}$  denotes the operator space projective tensor product. The duality bracket is given by

$$(4.58) \quad \langle P, T \otimes (x \otimes y) \rangle = \langle P(T)x, y \rangle_{B(L^2(G)), S_G^1}.$$

**Theorem 4.31** *Let  $G$  be a second countable unimodular inner amenable locally compact group. Then  $G$  has  $(\kappa_\infty)$  with  $\kappa_\infty(G) = 1$ , that means that there exists a contractive projection  $P_G^\infty: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  preserving the complete positivity onto the space  $\mathfrak{M}^{\infty, \text{cb}}(G)$  of completely bounded Fourier multipliers on the von Neumann algebra  $\text{VN}(G)$ .*

*Proof* : Fix some finite subset  $F$  of  $G$ . We can consider a net  $(V_j^F)_j$  provided by Definition 3.6. As in Lemma 4.9, we consider the elements  $y_j^F \stackrel{\text{def}}{=} c_j^F |\lambda(1_{V_j^F})|^2$  of  $L^1(\text{VN}(G)) \cap \text{VN}(G)$  and the symbol  $\phi_{j,T}^F$  if  $T: \text{VN}(G) \rightarrow \text{VN}(G)$  is a normal completely bounded map.

**Step 1** Consider the mapping  $P_j^F: \text{CB}(\text{VN}(G)) \rightarrow \text{CB}(S_G^\infty, B(L^2(G)))$ ,  $T \mapsto M_{\phi_{j,T}^F}$  where  $P_{w^*}$  is defined in [ArK1, proposition 3.1]. By Lemma 4.8, we have

$$\|M_{\phi_{j,T}^F}\|_{\text{cb}, S_G^\infty \rightarrow B(L^2(G))} \leq \|T\|_{\text{cb}, \text{VN}(G) \rightarrow \text{VN}(G)}.$$

Hence the linear maps  $P_j^F$  are in the unit ball of the space  $B(\text{CB}(\text{VN}(G)), \text{CB}(S_G^\infty, B(L^2(G))))$ . By Banach-Alaoglu theorem, we capture a weak\* accumulation point  $P^F: \text{CB}(\text{VN}(G)) \rightarrow \text{CB}(S_G^\infty, B(L^2(G)))$ . So, we have a net  $(P_{j(k)}^F)$  which converges to  $P^F$  for the weak\* topology. By (4.58), this implies that the net  $(P_{j(k)}^F(T))$ , that is  $(M_{\phi_{j(k), P_{w^*}^*(T)}^F})$ , converges for the point weak\* topology of  $\text{CB}(S_G^\infty, B(L^2(G)))$  to  $P^F(T)$ . Since the weak\* topology on  $\text{CB}(S_G^\infty, B(L^2(G)))$  coincides, essentially by the same argument than the one of [Pau, Lemma 7.2], on bounded subsets with the point weak\* topology, we conclude by the second part of Lemma 4.26 that  $P^F(T): S_G^\infty \rightarrow B(L^2(G))$  is itself a Schur multiplier. Note that by [Meg1, Theorem 2.6.14], we have

$$\|P^F\|_{\text{CB}(\text{VN}(G)) \rightarrow \text{CB}(S_G^\infty, B(L^2(G)))} \leq \liminf_{k \rightarrow \infty} \|P_{j(k)}^F\|_{\text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^\infty, B(L^2(G)))} \leq 1.$$

We next show that  $P^F$  preserves the complete positivity. Suppose that  $T$  is completely positive. Using Lemma 4.8, we see that  $M_{\phi_{j, P_{w^*}^*(T)}^F}$  is completely positive. Since  $P^F(T)$  is the limit in the point weak\* topology of the  $M_{\phi_{j(k), P_{w^*}^*(T)}^F}$ 's, the complete positivity of  $M_{\phi_{j, P_{w^*}^*(T)}^F}$  carries over to that of  $P^F(T)$  by [ArK1, Lemma 2.10].

We consider a weak\* accumulation point  $P^{(1)}: \text{CB}(\text{VN}(G)) \rightarrow \text{CB}(S_G^\infty, B(L^2(G)))$  of the net  $(P^F)_F$  and by the same reasoning as before, the map  $P^{(1)}(T): S_G^\infty \rightarrow B(L^2(G))$  is again a completely bounded Schur multiplier and preserves the complete positivity.

**Step 2** For any normal completely bounded map  $T: \text{VN}(G) \rightarrow \text{VN}(G)$ , we claim that the map  $P^{(1)}(T): S_G^\infty \rightarrow \text{B}(\text{L}^2(G))$  is in fact a Herz-Schur multiplier. The weak\* convergence of a subnet of  $(M_{\phi_{j, P_{w^*}^*(T)}}^F)_j$  to  $M_{\varphi^F} \stackrel{\text{def}}{=} P^F(T)$  implies that  $\varphi^F$  is a cluster point of  $(\phi_{j, P_{w^*}^*(T)}^F)_j$  for the weak\* topology of  $\text{L}^\infty(G \times G)$ . In the same manner, the symbol  $\varphi$  of  $P^{(1)}(T)$  is a cluster point of  $(\varphi^F)_F$  for the weak\* topology of  $\text{L}^\infty(G \times G)$ . Thus according to Lemma 4.9,  $\varphi$  is a Herz-Schur symbol. This is the step where we use that  $G$  is inner amenable.

**Step 3** By [BoF] [Spr1], we have an isometric map  $I: \mathfrak{M}_G^{\infty, \text{cb}, \text{HS}} \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  with range  $\mathfrak{M}^{\infty, \text{cb}}(G)$  preserving the complete positivity. We introduce the map  $P \stackrel{\text{def}}{=} I \circ P^{(1)}: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  with values in  $\mathfrak{M}^{\infty, \text{cb}}(G)$ . By composition, this map is contractive and preserves the complete positivity.

Finally, if  $T = M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  is a completely bounded Fourier multiplier, then for any  $j$  the symbol  $\phi_{j, T}^F$  of the Schur multiplier  $P_j^F(T): S_G^\infty \rightarrow \text{B}(\text{L}^2(G))$  is given by

$$\begin{aligned} \phi_{j, T}^F(s, t) &\stackrel{(4.29)}{=} \varphi_{1, y_j^F}(s, t) \stackrel{(4.25)}{=} \tau_G(y_j^F \lambda_{s^{-1}} T(\lambda_{st^{-1}}) \lambda_t) = \varphi(st^{-1}) \tau_G(y_j^F \lambda_{s^{-1}} \lambda_{st^{-1}} \lambda_t) \\ &= \varphi(st^{-1}) \tau_G(y_j^F) = \varphi(st^{-1}). \end{aligned}$$

Thus, also  $P^{(1)}(T) = P^F(T) = M_\varphi^{\text{HS}}$  and  $P(T) = I(M_\varphi^{\text{HS}}) = M_\varphi = T$ . ■

Now, we state the following general theorem of complementation onto the space of completely bounded Fourier multipliers. The proof uses in a crucial way that completely bounded Herz-Schur on  $S_G^p$  multipliers are in one-to-one (linear, norm and order) correspondence with completely bounded Fourier  $\text{L}^p$ -Fourier multipliers thanks to the amenability of  $G$ . Note that the construction of the projection in the case  $p = \infty$  of the following result is different of the one of Theorem 4.31.

**Theorem 4.32** *Let  $G$  be a second countable unimodular amenable locally compact group. Suppose  $1 < p < \infty$ . Assume that there are nets  $(x_j)$  and  $(y_j)$  of elements of  $\text{L}^1(\text{VN}(G)) \cap \text{VN}(G)$  with  $x_j, y_j \geq 0$  such that*

- $\|x_j\|_p, \|y_j\|_{p^*} \leq C$  for all  $j$ ,
- $\tau_G(x_j y_j) = 1$  for all  $j$ ,

*Then there exists a bounded projection*

$$P: \text{CB}(\text{L}^p(\text{VN}(G))) \rightarrow \text{CB}(\text{L}^p(\text{VN}(G)))$$

*onto the space  $\mathfrak{M}^{p, \text{cb}}(G)$  of completely bounded Fourier multipliers with the properties*

1.  $\|P\| \leq C^2$ ,
2.  $P(T)$  is completely positive whenever  $T$  is completely positive.

*Finally, let  $p = \infty$  or  $p = 1$ . Let  $(x_j)$  be a net of positive elements of  $\text{L}^1(\text{VN}(G)) \cap \text{VN}(G)$  such that  $\tau_G(x_j) = 1$  for all  $j$ . Then there exists contractive projections*

$$P: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G)) \quad \text{and} \quad P: \text{CB}(\text{L}^1(\text{VN}(G))) \rightarrow \text{CB}(\text{L}^1(\text{VN}(G)))$$

*on  $\mathfrak{M}^{\infty, \text{cb}}(G)$  and  $\mathfrak{M}^{1, \text{cb}}(G)$  satisfying the previous second property.*

*Proof :*



**Step 1** Consider the mapping  $P_j: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(S_G^p)$ ,  $T \mapsto M_{\phi_j, T}$  (restrict to normal operators in case  $p = \infty$ ). Using Lemma 4.8, we obtain the estimate

$$\|P_j(T)\|_{\text{cb}, S_G^p \rightarrow S_G^p} \leq \|T\|_{\text{cb}, p \rightarrow p} \|x_j\|_p \|y_j\|_{p^*} \leq C^2 \|T\|_{\text{cb}, L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))},$$

according to the assumptions of the theorem. In the case  $p = \infty$  or  $p = 1$ , we put  $P^{(1)} = P_{j_0}$  for some fixed index  $j_0$ . In the case  $1 < p < \infty$ , we capture  $P^{(1)}$  as a limit. Namely, then  $(P_j)$  is a bounded net in the space  $\text{B}(\text{CB}(L^p(\text{VN}(G))), \text{CB}(S_G^p))$ , which is a dual space so admits by Banach-Alaoglu's theorem a weak\* accumulation point, which we denote  $P^{(1)}$  and fix for the rest of the proof. Since  $\mathfrak{M}_G^{p, \text{cb}}$  is weak\* closed in  $\text{CB}(S_G^p)$  according to Lemma 4.26, it is easy to check that  $P^{(1)}$  takes its image in  $\mathfrak{M}_G^{p, \text{cb}}$ . This is clear by construction in the case  $p = \infty$  or  $p = 1$ .

Lemma 4.8 says that the Schur multiplier  $P_j(T): S_G^p \rightarrow S_G^p$  is completely positive. By [ArK1, Lemma 2.10], the point weak\* limit  $P^{(1)}(T)$  is again completely positive (clear in case  $p = \infty$  or  $p = 1$ ). If  $T = M_\varphi$  is itself a Fourier multiplier, the weak\* limit  $P^{(1)}(T)$  of the net  $(P_j(T))$  has symbol  $(s, t) \mapsto \varphi(st^{-1})$  (clear in the case  $p = \infty$  or  $p = 1$ ).

**Step 2** We will use the contractive projection onto the space  $\mathfrak{M}_G^{p, \text{cb}, \text{HS}}$  of completely bounded Herz-Schur multipliers of Proposition 4.27. Seeing this projection as a map  $Q: \mathfrak{M}_G^{p, \text{cb}} \rightarrow \mathfrak{M}_G^{p, \text{cb}, \text{HS}}$ , we let  $P^{(2)} \stackrel{\text{def}}{=} Q \circ P^{(1)}: \text{CB}(L^p(\text{VN}(G))) \rightarrow \mathfrak{M}_G^{p, \text{cb}, \text{HS}}$ . Then by this proposition together with Step 1,  $P^{(2)}$  satisfies  $\|P^{(2)}\| \leq C^2$  and preserves the complete positivity and  $P^{(2)}(M_\varphi)$  has again symbol  $(s, t) \mapsto \varphi(st^{-1})$ .

**Step 3** By [CDS] and [NR], we can consider the isometric map  $I: \mathfrak{M}_G^{p, \text{cb}, \text{HS}} \rightarrow \text{CB}(L^p(\text{VN}(G)))$  onto the space  $\mathfrak{M}^{p, \text{cb}}(G)$  of completely bounded Fourier multipliers. Note that if the Herz-Schur multiplier  $M_\varphi^{\text{HS}}: S_G^p \rightarrow S_G^p$  is completely positive, it is easy to see with [ArK1, Lemma 2.10] that the Fourier multiplier  $I(M_\varphi^{\text{HS}})$  is completely positive as a limit for the weak operator topology of completely positive maps.

We finally put  $P \stackrel{\text{def}}{=} I \circ P^{(2)}: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(L^p(\text{VN}(G)))$  and need to check the claimed properties. First,  $\|P\| \leq \|I\| \|P^{(2)}\| \leq C^2$ . Second, if  $T$  is completely positive, then so is  $P^{(2)}(T)$ , so also  $P(T) = I \circ P^{(2)}(T)$ . Third, if  $T = M_\varphi$  is itself a multiplier, then  $P^{(2)}(T)$  is the Herz-Schur multiplier with symbol  $(s, t) \mapsto \varphi(st^{-1})$ . So  $P(T)$  is the Fourier multiplier with symbol  $\varphi$  again.  $\blacksquare$

Using the Theorem 4.32, we obtain the following result.

**Corollary 4.33** *Let  $G$  be a second countable unimodular amenable locally compact group. Let  $1 \leq p \leq \infty$  such that  $\frac{p}{p^*}$  is rational. Then there exists a contractive projection*

$$P^p: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(L^p(\text{VN}(G)))$$

*on the space  $\mathfrak{M}^{p, \text{cb}}(G)$  such that  $P^p(T)$  is completely positive whenever  $T$  is completely positive (with the usual convention if  $p = \infty$ ).*

**Remark 4.34** Going through the different steps of the proof, one sees that the projection in the Corollary 4.33 has the property that  $P^p(T)^* = P^{p^*}(T^*)$  for any completely bounded map  $T: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))$  and any  $1 \leq p < \infty$ . Here we emphasize that  $P = P^p$  depends on  $1 \leq p \leq \infty$ . However, it is not clear that  $P^p(T)$  and  $P^q(T)$  coincide for  $1 \leq p, q \leq \infty$  and a  $T$  acting on both  $L^p(\text{VN}(G))$  and  $L^q(\text{VN}(G))$ .

Using Theorem 4.32, we obtain the following important property of compatibility of projections which is achieved through a different method.

**Corollary 4.35** *Let  $G$  be a second countable unimodular amenable locally compact group. Suppose that  $G$  is finite-dimensional. Then  $G$  has  $(\kappa)$ . More precisely, for all  $1 \leq p \leq \infty$  there exists a bounded projection*

$$P^p: \text{CB}(L^p(\text{VN}(G))) \rightarrow \text{CB}(L^p(\text{VN}(G)))$$

on  $\mathfrak{M}^{p,\text{cb}}(G)$  (resp  $P^\infty: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  on  $\mathfrak{M}^{\infty,\text{cb}}(G)$ ) with the properties

1.  $\|P^p\| \leq C$ , where the constant  $C$  depends on  $G$  but not on  $p$ .
2.  $P^p(T)$  is completely positive whenever  $T$  is completely positive.
3. If  $T$  belongs to  $\text{CB}(L^p(\text{VN}(G)))$  and to  $\text{CB}(L^q(\text{VN}(G)))$  for two values  $1 \leq p, q \leq \infty$ , then the Fourier multipliers  $P^p(T)$  and  $P^q(T)$  are compatible mappings coinciding on  $L^p(\text{VN}(G)) \cap L^q(\text{VN}(G))$ , i.e. have the same symbol.

*Proof* : Consider first the case  $1 < p < \infty$ . It suffices to pick the sequence  $(f_j)$  in  $C_c(G)$  and the associated  $x_j = a_j \lambda(f_j)$ ,  $y_j = b_j \lambda(f_j)$  from Corollary 4.23. Then apply this corollary together with Theorem 4.32 to deduce the two first points. Let us show the last point 3. By the particular choice of  $x_j = a_j \lambda(f_j)$  and  $y_j = b_j \lambda(f_j)$  with  $a_j \cdot b_j$  independent of  $p$  (see Corollary 4.23), we deduce that  $P_j^p(T)K_\phi = P_j^q(T)K_\phi$  for  $K_\phi \in S^p \cap S^q$ , where  $P_j^p, P_j^q$  denote the mappings from Step 1 of the proof of Theorem 4.32.

Consider  $X = \prod_{1 < p < \infty} \text{B}(\text{CB}(L^p(\text{VN}(G))), \text{CB}(S_G^p))$  which is a topological space when equipped with the product topology of the weak\* topology. Then  $P_j = (P_j^p)_{1 < p < \infty}$  lies in a compact subspace of  $X$  for all  $j$ , since  $\|P_j^p\| \leq C$  with a constant independent of  $p$  and Tychonoff's theorem for the product of compact spaces applies here. Thus,  $(P_j)_j$  admits an accumulation point in  $X$ , that is to say that for the same subnet  $j'$  for all  $p$ , we have  $P^{(1),p} = \lim_{j'} P_{j'}^p$ . We infer that  $P^{(1),p}(T)K_\phi = P^{(1),q}(T)K_\phi$  for any  $K_\phi \in S^p \cap S^q$ , where  $P^{(1),p}, P^{(1),q}$  denote the mappings from Step 1 of the proof of Theorem 4.32. Since the mapping  $Q$  from Proposition 4.27 is compatible for different values  $p$  and  $q$ , also  $P^{(2),p}$  and  $P^{(2),q}$  from Step 2 of the proof of Theorem 4.32 are compatible. Finally, since the mapping  $I$  from Step 3 of the proof of Theorem 4.32 is compatible for different values  $p$  and  $q$  (being the identity), also  $P^p$  and  $P^q$  are compatible in the sense of point 4 of the corollary.

Now turn to the case  $p \in \{1, \infty\}$ . We let again  $x_j = a_j \lambda(f_j)$  and  $y_j = b_j \lambda(f_j)$  as in Corollary 4.23. Let us go through steps 1,2,3 of the proof of Theorem 4.32, but with this new choice of  $x_j$  and  $y_j$  which is due to the compatibility of point 4 of the statement, and where these steps have to be adapted here. We let  $\langle P_j^p(T)K_\phi, K_\psi \rangle = \langle M_\varphi(K_\phi), K_\psi \rangle$  as in Theorem 4.32. As in step 1 in its proof, we obtain that  $P_j^p(T)$  belongs to  $\mathfrak{M}_G^{p,\text{cb}}$  with norm control  $\|x_j\|_p \|y_j\|_{p^*} \|T\|_{\text{cb}, p \rightarrow p}$ . In case  $p = 1$ , by the same compactness argument as above, we find that for some subnet  $j'$ , there is a weak\* limit  $P^p(T) = \lim_{j'} P_{j'}^p(T)$  belonging to  $\mathfrak{M}_G^{p,\text{cb}}$  according to Lemma 4.26. In case  $p = \infty$ , view  $P_j^p$  as elements of  $\text{B}(\text{CB}(\text{VN}(G)), \text{CB}(S^\infty, \text{B}(L^2(G))))$  and argue in the same way. Since all  $P_j^p(T)$  are compatible also for  $1 \leq p \leq \infty$ , also the  $P^p(T)$  are compatible.

The only difference to the case  $1 < p < \infty$  is that if  $T = M_\varphi$  belongs to  $\mathfrak{M}^{p,\text{cb}}(G)$ , we cannot rely on Theorem 4.32 any more to ensure that  $P^p(M_\varphi) = M_{\varphi(st^{-1})}$ . Instead, we observe that  $M_\varphi \in \mathfrak{M}^{p,\text{cb}}(G) \subset \mathfrak{M}^{2,\text{cb}}(G)$  and that  $P^p(M_\varphi) = P^2(M_\varphi) = M_{\varphi(st^{-1})}$ , the first equality being a consequence of the compatibility.

Now steps 2 and 3 go through as in the first part of the proof to show statements 1-4 for general  $1 \leq p \leq \infty$ . ■

**Example 4.36** Let  $G$  be a second countable unimodular amenable locally compact group. Suppose that  $G$  is in addition totally disconnected, hence zero-dimensional by Remark 4.15. Consequently, Corollary 4.35 applies. Then the projections  $P^p$  in that result are in fact contractions.

Indeed, an inspection of the proof of Corollary 4.35 (see also Step 1 of the proof of Theorem 4.32) shows that  $\|P^p\| \leq \sup_j \|x_j\|_p \|y_j\|_{p^*}$ . According to Remark 4.24, the right-hand side is less than 1.

Therefore, Corollary 4.35 gives a variant of [ArK1, Theorem 6.38] stated for (second countable) pro-discrete groups for (second countable unimodular) amenable totally disconnected groups. Note that Corollary 4.35 needs that the group  $G$  is amenable in the cases  $p = \infty$  and  $p = 1$  in contrast to [ArK1, Theorem 6.38]. We conjecture that inner amenability is equivalent to  $(\kappa_\infty)$ . Hence the amenability seems necessary by the last observation of Example 3.3.

**Example 4.37** Recall that a semisimple connected Lie group is unimodular by [Fol, page 47] and amenable if and only if it is compact by [Pie, Proposition 14.10]. A connected nilpotent Lie group is unimodular and amenable by [Pal2, page 1487]. So Corollary 4.35 applies to second countable groups belonging to these classes.

**Remark 4.38** We have resisted the temptation to write a matricial version of Theorem 4.31 in the spirit of [ArK1, Theorem 4.2] in order to obtain the property  $(\kappa_{\infty,2})$  of Definition 2.17. It is very likely that the same method works.

**Remark 4.39** We make no attempt with non-unimodular locally compact groups. It is likely that the same strategy works in the case  $p = \infty$ . We leave this case as an exercise for the reader. At the level of quantum groups, the first author has an unpublished which is a generalization of a result of [ArK1].

## 5 Final remarks

### 5.1 A characterization of the amenability of unimodular locally compact groups

We give a new characterization of amenability which is similar to the one of Theorem 3.2. Recall that  $(\kappa_\infty)$  is defined in Definition 1.2.

**Theorem 5.1** *Let  $G$  be a second countable unimodular locally compact group. Then the following are equivalent.*

1.  $\text{VN}(G)$  is injective and  $G$  has  $(\kappa_\infty)$ .
2.  $G$  is amenable.

Moreover, the implication 1.  $\Rightarrow$  2. is true without the assumption “unimodular”.

*Proof* : 1.  $\Rightarrow$  2. Since the group  $G$  has  $(\kappa_\infty)$ , Proposition 2.14 gives

$$\mathfrak{M}^{\infty, \text{dec}}(G) = \text{B}(G).$$

Since  $\text{VN}(G)$  is injective, by [Haa, Theorem 1.6] each completely bounded operator  $T: \text{VN}(G) \rightarrow \text{VN}(G)$  is decomposable with  $\|T\|_{\text{cb}, \text{VN}(G) \rightarrow \text{VN}(G)} = \|T\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}$ . In particular, we have  $\mathfrak{M}^{\infty, \text{cb}}(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$  isometrically. We deduce that

$$\mathfrak{M}^{\infty, \text{cb}}(G) = \text{B}(G).$$

Using [Pis6, page 54] (see also [Spr1, page 190] and [KaL1]), we conclude that  $G$  is amenable.

2.  $\Rightarrow$  1. If  $G$  is locally compact and amenable, then by Theorem 3.2 the von Neumann algebra  $\text{VN}(G)$  is injective. For the second property, it suffices to use Theorem 4.31. ■

Now, we prove that there exists locally compact groups  $G$  such the space  $\mathfrak{M}^{\infty, \text{dec}}(G)$  of completely bounded Fourier multipliers on the von Neumann algebra  $\text{VN}(G)$  is not complemented in the space  $\text{CB}_{w^*}(\text{VN}(G))$ .

**Corollary 5.2** *Any non-amenable second countable connected locally compact group  $G$  has not property  $(\kappa_\infty)$ .*

*Proof* : By [Con2, Corollary 7], the von Neumann algebra  $\text{VN}(G)$  of such a group is injective. Suppose that  $G$  has  $(\kappa_\infty)$ . By the implication 1.  $\Rightarrow$  2. of Theorem 5.1, we obtain that  $G$  is amenable, i.e. a contradiction. ■

**Example 5.3** This result applies for example to the locally compact group  $\text{SL}_2(\mathbb{R})$  which is non-amenable by [BHV, Example G.2.4 (i)] contradicting<sup>26</sup>. [DFSW, Remark 7.6] stated for unimodular locally compact quantum groups since  $\text{SL}_2(\mathbb{R})$  is unimodular by [Lan1, page 4].

The following result is a generalization of Theorem 2.22 if we do not take account the second countability.

**Corollary 5.4** *Let  $G$  be a second countable unimodular inner amenable locally compact group. Then the von Neumann algebra  $\text{VN}(G)$  is injective if and only if we have  $\mathfrak{M}^{\infty, \text{dec}}(G) = \mathfrak{M}^{\infty, \text{cb}}(G)$ .*

*Proof* : The “only if” part is true by a result of Haagerup [Haa]. By Proposition 2.14 and Theorem 4.31, we have a bijection from  $\mathfrak{M}^{\infty, \text{cb}}(G)$  onto  $\text{B}(G)$ . So  $G$  is amenable by [Pis6, p. 54]. We conclude that  $\text{VN}(G)$  is injective by Theorem 3.2. ■

## 5.2 A description of decomposable Fourier multipliers on some locally compact groups

The following is a variant of [ArK1, Theorem 6.45] which can be proved with similar arguments.

**Theorem 5.5** *Let  $G$  be a second countable amenable unimodular totally disconnected locally compact group. Suppose  $1 < p < \infty$ . Then a measurable function  $\phi: G \rightarrow \mathbb{C}$  induces a decomposable Fourier multiplier  $M_\phi: L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))$  if and only if it induces a (completely) bounded Fourier multiplier on  $\text{VN}(G)$ . In this case, we have*

$$(5.1) \quad \|M_\phi\|_{\text{VN}(G) \rightarrow \text{VN}(G)} = \|M_\phi\|_{\text{dec}, L^p(\text{VN}(G)) \rightarrow L^p(\text{VN}(G))}.$$

## 5.3 $\text{CB}_{w^*}$ -strongly non decomposable multipliers

The following is a variant of [ArK1, Definition 7.2].

**Definition 5.6** *We say that a normal completely bounded operator  $T: M \rightarrow M$  is  $\text{CB}_{w^*}$ -strongly non decomposable if  $T$  does not belong to the closure  $\overline{\text{Dec}_{w^*}(M)}^{\text{CB}}$  of the space  $\text{Dec}_{w^*}(M)$  of weak\* continuous decomposable operators with respect to the completely bounded norm  $\|\cdot\|_{\text{cb}, M \rightarrow M}$ .*

<sup>26</sup> We would like to thank Adam Skalski for his confirmation of this problem in this small remark by email on his own initiative. The results of the nice paper [DFSW] remains correct.

It means that  $T$  cannot be approximated by normal decomposable operators. Of course, a  $\text{CB}_{w^*}$ -strongly non decomposable operator is not decomposable. Note [Haa, Corollary 2.8] which says that if  $M$  is injective, it is impossible to find such operator.

Let  $G$  be a discrete group. By the discussion of the introduction, we know that  $G$  is amenable if and only if  $\mathfrak{M}^{\infty, \text{cb}}(G) = \mathfrak{M}^{\infty, \text{dec}}(G)$ . So the following question is natural.

**Question 5.7** *Let  $G$  be a non-amenable discrete group. Does there exists a  $\text{CB}_{w^*}$ -strongly non decomposable completely bounded Fourier multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  ?*

The result [ArK1, Proposition 7.25] solves an  $L^p$ -version of this problem with  $1 < p < \infty$  and relies on the use of *infinite abelian* groups. So we cannot hope to solve Question 5.7 with the methods of [ArK1].

**Proposition 5.8** *Let  $G$  be a locally compact group with  $\kappa_\infty(G) = 1$ . Let  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  be a completely bounded Fourier multiplier. We have*

$$(5.2) \quad \text{dist}_{\text{CB}(\text{VN}(G))}(M_\varphi, \text{Dec}_{w^*}(\text{VN}(G))) \geq \text{dist}_{\text{CB}(\text{VN}(G))}(M_\varphi, \mathfrak{M}^{\infty, \text{dec}}(G)).$$

*Proof* : Let  $R: \text{VN}(G) \rightarrow \text{VN}(G)$  be a normal decomposable operator. By [Haa, page 175], we can write

$$R = R_1 - R_2 + i(R_3 - R_4)$$

where each  $R_j$  is a completely positive map on  $\text{VN}(G)$ . We can suppose each  $R_j$  normal by the classical argument using [ArK1, Proposition 3.1]. Using the contractive projection  $P_G^\infty: \text{CB}_{w^*}(\text{VN}(G)) \rightarrow \text{CB}_{w^*}(\text{VN}(G))$  of Definition 1.2, we obtain

$$P_G^\infty(R) = P_G^\infty(R_1 - R_2 + i(R_3 - R_4)) = P_G^\infty(R_1) - P_G^\infty(R_2) + i(P_G^\infty(R_3) - P_G^\infty(R_4)).$$

By [Haa, page 175], we infer that the Fourier multiplier  $P_G^\infty(R)$  is decomposable on  $\text{VN}(G)$  and in particular completely bounded by [Haa, Proposition 1.3 (3)]. Now, using the contractivity of the projection  $P_G^\infty$ , we obtain that

$$\begin{aligned} \|M_\varphi - R\|_{\text{cb}, \text{VN}(G) \rightarrow \text{VN}(G)} &\geq \|P_G^\infty(M_\varphi - R)\|_{\text{cb}, \text{VN}(G) \rightarrow \text{VN}(G)} \\ &= \|P_G^\infty(M_\varphi) - P_G^\infty(R)\|_{\text{cb}, \text{VN}(G) \rightarrow \text{VN}(G)} = \|M_\varphi - P_G^\infty(R)\|_{\text{cb}, \text{VN}(G) \rightarrow \text{VN}(G)} \\ &\geq \text{dist}_{\text{CB}(\text{VN}(G))}(M_\varphi, \mathfrak{M}^{\infty, \text{dec}}(G)). \end{aligned}$$

Hence, passing to the infimum we deduce (5.2). ■

We denote by  $\overline{\text{B}(G)}^{\text{cb}}$  the closure of the Fourier-Stieltjes algebra  $\text{B}(G)$  in the space  $\mathfrak{M}^{\infty, \text{cb}}(G)$  of completely bounded Fourier multipliers on  $\text{VN}(G)$ . See [BrF1] for related things.

**Corollary 5.9** *Let  $G$  be a discrete group. Any completely bounded Fourier multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  such that the function  $\varphi$  does not belong to the closure  $\overline{\text{B}(G)}^{\text{cb}}$  is a  $\text{CB}_{w^*}$ -strongly non decomposable operator.*

*Proof* : By Corollary 2.20, we have an isometry  $\mathfrak{M}^{\infty, \text{dec}}(G) = \text{B}(G)$ . Now, it suffices to use Proposition 5.8. ■

So, it is natural to state the following conjecture.

**Conjecture 5.10** *Let  $G = \mathbb{F}_n$  be the free group with  $n \geq 2$ . There exists a completely bounded Fourier multiplier  $M_\varphi: \text{VN}(\mathbb{F}_n) \rightarrow \text{VN}(\mathbb{F}_n)$  which does not belong to the closure  $\overline{\text{B}(G)}^{\text{cb}}$ .*

Note that [DCH, Corollary 3.10] says that the contractive inclusion  $\text{B}(G) \subset \mathfrak{M}^{\infty, \text{cb}}(G)$  is *strict* if  $G = \mathbb{F}_n$  is the free group with  $2 \leq n \leq \infty$  ( $n$  countable). Maybe, it is possible to improve the method of [DCH, Corollary 3.10]. See also [DCH, Corollary 2.5] for the case of other groups.

## 5.4 Multipliers on reduced $C^*$ -algebras vs multipliers on von Neumann algebras

Let  $G$  be a locally compact group and  $\varphi: G \rightarrow \mathbb{C}$  be a function. It is well known that by [DCH, Proposition 1.2 and Definition 1.3] the multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  is completely bounded if and only if  $M_\varphi: C_\lambda^*(G) \rightarrow C_\lambda^*(G)$  is completely bounded. So the following is a natural question.

**Proposition 5.11** *Let  $G$  be a discrete group and  $\varphi: G \rightarrow \mathbb{C}$  be a function. Then  $\varphi$  induces a decomposable Fourier multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  if and only if  $\varphi$  induces a decomposable Fourier multiplier  $M_\varphi: C_\lambda^*(G) \rightarrow C_\lambda^*(G)$ . In this case, we have*

$$\|M_\varphi\|_{\text{dec}, C_\lambda^*(G) \rightarrow C_\lambda^*(G)} = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}.$$

*Proof* :  $\Rightarrow$ : Suppose that the multiplier  $M_\varphi: \text{VN}(G) \rightarrow \text{VN}(G)$  is decomposable. By [Haa, Remark 1.5], there exist some linear maps  $v_1, v_2: \text{VN}(G) \rightarrow \text{VN}(G)$  such that the linear map  $\begin{bmatrix} v_1 & M_\varphi \\ M_{\check{\varphi}} & v_2 \end{bmatrix}: \text{M}_2(\text{VN}(G)) \rightarrow \text{M}_2(\text{VN}(G))$  is completely positive with  $\max\{\|v_1\|, \|v_2\|\} = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}$ . We can suppose that  $v_1$  and  $v_2$  are in addition weak\* continuous by using [ArK1, Proposition 3.1]. Now, we can apply the amplification projection  $\text{Id}_{\text{M}_2} \otimes P_G^\infty: \text{M}_2(\text{CB}_{w^*}(\text{VN}(G))) \rightarrow \text{M}_2(\text{CB}_{w^*}(\text{VN}(G)))$  of [ArK1, Theorem 4.2]. We obtain that

$$(5.3) \quad \begin{bmatrix} P_G^\infty(v_1) & M_\varphi \\ M_{\check{\varphi}} & P_G^\infty(v_2) \end{bmatrix} = \begin{bmatrix} P_G^\infty(v_1) & P_G^\infty(M_\varphi) \\ P_G^\infty(M_{\check{\varphi}}) & P_G^\infty(v_2) \end{bmatrix} = (\text{Id}_{\text{M}_2} \otimes P_G^\infty) \left( \begin{bmatrix} v_1 & M_\varphi \\ M_{\check{\varphi}} & v_2 \end{bmatrix} \right)$$

is completely positive. Moreover, we have

$$\max\{\|P_G^\infty(v_1)\|, \|P_G^\infty(v_2)\|\} \leq \|P_G^\infty\| \max\{\|v_1\|, \|v_2\|\} = \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}.$$

The restriction to  $\text{M}_2(C_\lambda^*(G))$  of the map (5.3) shows that the function  $\varphi$  induces a decomposable multiplier  $M_\varphi: C_\lambda^*(G) \rightarrow C_\lambda^*(G)$  with

$$\|M_\varphi\|_{\text{dec}, C_\lambda^*(G) \rightarrow C_\lambda^*(G)} \leq \|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)}.$$

$\Leftarrow$  : Suppose that the multiplier  $M_\varphi: C_\lambda^*(G) \rightarrow C_\lambda^*(G)$  is decomposable. Let  $\varepsilon > 0$ . There exist some maps  $v_1, v_2: C_\lambda^*(G) \rightarrow C_\lambda^*(G)$  such that the map  $\begin{bmatrix} v_1 & M_\varphi \\ M_{\check{\varphi}} & v_2 \end{bmatrix}: \text{M}_2(C_\lambda^*(G)) \rightarrow \text{M}_2(C_\lambda^*(G))$  is completely positive with  $\max\{\|v_1\|, \|v_2\|\} \leq \|M_\varphi\|_{\text{dec}, C_\lambda^*(G) \rightarrow C_\lambda^*(G)} + \varepsilon$ . By an easy adaptation of [ArK1, Theorem 4.2] and [Haa3, Lemma 2.5] to the  $C^*$ -setting, we have a contractive projection  $\text{Id}_{\text{M}_2} \otimes P_G^\infty: \text{M}_2(\text{CB}(C_\lambda^*(G))) \rightarrow \text{M}_2(\text{CB}(C_\lambda^*(G)))$  which preserves the complete positivity. We obtain that

$$(5.4) \quad \begin{bmatrix} P_G^\infty(v_1) & M_\varphi \\ M_{\check{\varphi}} & P_G^\infty(v_2) \end{bmatrix} = \begin{bmatrix} P_G^\infty(v_1) & P_G^\infty(M_\varphi) \\ P_G^\infty(M_{\check{\varphi}}) & P_G^\infty(v_2) \end{bmatrix} = (\text{Id}_{\text{M}_2} \otimes P_G^\infty) \left( \begin{bmatrix} v_1 & M_\varphi \\ M_{\check{\varphi}} & v_2 \end{bmatrix} \right)$$

is completely positive. Moreover, we have

$$\begin{aligned} \max\{\|P_G^\infty(v_1)\|_{C_\lambda^*(G) \rightarrow C_\lambda^*(G)}, \|P_G^\infty(v_2)\|_{C_\lambda^*(G) \rightarrow C_\lambda^*(G)}\} &\leq \|P_G^\infty\| \max\{\|v_1\|, \|v_2\|\} \\ &\leq \|M_\varphi\|_{\text{dec}, C_\lambda^*(G) \rightarrow C_\lambda^*(G)} + \varepsilon. \end{aligned}$$

By extending each block as a Fourier multiplier on  $\text{VN}(G)$ , we have a weak\* continuous extension  $M_2(\text{VN}(G)) \rightarrow M_2(\text{VN}(G))$  of the map (5.4). Using Kaplansky's theorem, it is easy to see that this map is completely positive. This shows that the function  $\varphi$  induces a decomposable multiplier  $M_\varphi : \text{VN}(G) \rightarrow \text{VN}(G)$ .

By [DCH, page 461], we have the equalities  $\|P_G^\infty(v_1)\|_{\text{VN}(G) \rightarrow \text{VN}(G)} = \|P_G^\infty(v_1)\|_{C_\lambda^*(G) \rightarrow C_\lambda^*(G)}$  and  $\|P_G^\infty(v_2)\|_{\text{VN}(G) \rightarrow \text{VN}(G)} = \|P_G^\infty(v_2)\|_{C_\lambda^*(G) \rightarrow C_\lambda^*(G)}$ . We conclude that

$$\|M_\varphi\|_{\text{dec}, \text{VN}(G) \rightarrow \text{VN}(G)} \leq \|M_\varphi\|_{\text{dec}, C_\lambda^*(G) \rightarrow C_\lambda^*(G)} .$$

■

**Remark 5.12** The same argument seems correct for locally compact groups with a  $C^*$ -variation of the property  $(\kappa_{\infty,2})$  of Definition 2.17.

**Acknowledgment.** The authors are greatly indebted to the anonymous referee of [ArK1]. Indeed, his report contained an interesting research line which combined with unpublished results, news ideas and some work has given this paper. We thank Claire Anantharaman-Delaroche and Martin Walter for very short discussions, Matthew Daws, Adam Skalski and Ami Viselter for old discussions and encouragements and finally Marco Peloso for a discussion on (4.38) in the non-unimodular case.

Both authors were supported by the grant ANR-18-CE40-0021 (project HASCON). The second author was partly supported by the grant ANR-17-CE40-0021 of the French National Research Agency ANR (project Front).

## Bibliography

- [Ar1] A. Arhangel'skii. On the identity of the dimension  $\text{ind } G$  and  $\text{dim } G$  for locally bicomact groups. *Dokl. Akad. Nauk SSSR* 132 980–981 (Russian); translated as *Soviet Math. Dokl.* 1 1960 670–671. [38](#)
- [ArM1] A. V. Arhangel'skii and J. van Mill. Some aspects of dimension theory for topological groups. *Indag. Math. (N.S.)* 29 (2018), no. 1, 202–225. [38](#)
- [Arh9] C. Arhancet. Dilations of markovian semigroups of Fourier multipliers on locally compact groups. *Proceedings of the American Mathematical Society* (2020), 2551–2563. [10](#)
- [Arh10] C. Arhancet. Dilations of markovian semigroups of measurable Schur multipliers. Preprint. [12](#)
- [ArK1] C. Arhancet and C. Kriegler. Projections, multipliers and decomposable maps on noncommutative  $L^p$ -spaces. To appear in *Mémoires de la Société Mathématique de France*. Preprint on <https://arxiv.org/abs/1707.05591> [2](#), [3](#), [4](#), [5](#), [6](#), [9](#), [10](#), [14](#), [15](#), [16](#), [17](#), [24](#), [26](#), [33](#), [46](#), [47](#), [49](#), [51](#), [52](#), [53](#), [54](#), [55](#)
- [BEM13] L. Bandara, A. F. M. ter Elst and A. McIntosh. Square roots of perturbed subelliptic operators on Lie groups. *Studia Math.* 216 (2013), no. 3, 193–217. [39](#)
- [BHV] B. Bekka, P. de la Harpe, Pierre and A. Valette. *Kazhdan's property (T)*. New Mathematical Monographs, 11. Cambridge University Press, Cambridge, 2008. [12](#), [15](#), [18](#), [52](#)
- [BeH1] B. Bekka and P. de la Harpe. Unitary representations of groups, duals, and characters. Preprint, [arXiv:1912.07262](https://arxiv.org/abs/1912.07262). [6](#), [18](#)
- [BeL] J. Bergh, J. Löfström. *Interpolation spaces. An Introduction*. Springer-Verlag, Berlin, Heidelberg, New York, 1976. [4](#)
- [Bin] L. Bing-Ren. *Introduction to operator algebras*. World Scientific Publishing, 1992. [10](#)
- [BLM] D. Blecher and C. Le Merdy. *Operator algebras and their modules—an operator space approach*. London Mathematical Society Monographs. New Series, 30. Oxford Science Publications. The Clarendon Press, Oxford University Press, Oxford, 2004. [2](#), [11](#), [22](#), [28](#)

- [Bou2] N. Bourbaki. Integration. I. Chapters 1–6. Translated from the 1959, 1965 and 1967 French originals by Sterling K. Berberian. Elements of Mathematics. Springer-Verlag, Berlin, 2004. [11](#), [13](#)
- [Bou3] N. Bourbaki. Integration. II. Chapters 7–9. Translated from the 1963 and 1969 French originals by Sterling K. Berberian. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004. [11](#), [12](#), [17](#), [23](#), [38](#)
- [Bou4] N. Bourbaki. General topology. Chapters 1–4. Translated from the French. Reprint of the 1989 English translation. Elements of Mathematics. (Berlin). Springer-Verlag, Berlin, 1998. [38](#)
- [BoF] M. Bożejko and G. Fendler. Herz-Schur multipliers and completely bounded multipliers of the Fourier algebra of a locally compact group. Boll. Un. Mat. Ital. A (6) 3 (1984), no. 2, 297–302. [25](#), [26](#), [48](#)
- [BrF1] M. Brannan and B. Forrest. Extending multipliers of the Fourier algebra from a subgroup. Proc. Amer. Math. Soc. 142 (2014), no. 4, 1181–1191. [53](#)
- [Bre14] E. Breuillard. Geometry of locally compact groups of polynomial growth and shape of large balls. Groups Geom. Dyn. 8 (2014), no. 3, 669–732. [40](#)
- [BrO] N. P. Brown and N. Ozawa.  $C^*$ -algebras and finite-dimensional approximations. Graduate Studies in Mathematics, 88. American Mathematical Society, Providence, RI, 2008. [12](#)
- [Bun1] M. R. Buneci. Groupoid  $C^*$ -algebras. Surv. Math. Appl. 1 (2006), 71–98. [8](#)
- [CDS] M. Caspers and M. de la Salle. Schur and Fourier multipliers of an amenable group acting on non-commutative  $L^p$ -spaces. Trans. Amer. Math. Soc. 367 (2015), no. 10, 6997–7013. [26](#), [49](#)
- [CPPR] M. Caspers, J. Parcet, M. Perrin and É. Ricard. Noncommutative de Leeuw theorems. Forum Math. Sigma 3 (2015), e21. [18](#)
- [Con2] A. Connes. Classification of injective factors. Cases  $II_1$ ,  $II_\infty$ ,  $III_\lambda$ ,  $\lambda \neq 1$ . Ann. of Math. (2) 104 (1976), no. 1, 73–115. [15](#), [52](#)
- [CrT1] J. Crann and Z. Tanko. On the operator homology of the Fourier algebra and its  $cb$ -multiplier completion. J. Funct. Anal. 273 (2017), no. 7, 2521–2545. [21](#)
- [CoH1] Y. Cornulier and P. de la Harpe. Metric Geometry of Locally Compact Groups. European Mathematical Society, 2016. [38](#), [41](#)
- [DFSW] M. Daws, P. Fima, A. Skalski and S. White. The Haagerup property for locally compact quantum groups. J. Reine Angew. Math. 711 (2016), 189–229. [52](#)
- [DCH] J. de Cannière and U. Haagerup. Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. Amer. J. Math. 107 (1985), no. 2, 455–500. [6](#), [10](#), [14](#), [53](#), [54](#), [55](#)
- [Dix1] J. Dixmier.  $C^*$ -algebras. Translated from the French by Francis Jellet. North-Holland Mathematical Library, Vol. 15. North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977. [11](#), [13](#)
- [Dix2] J. Dixmier. Von Neumann algebras. With a preface by E. C. Lance. Translated from the second French edition by F. Jellet. North-Holland Mathematical Library, 27. North-Holland Publishing Co., Amsterdam-New York, 1981. [11](#)
- [Duq22] C. Duquet. Dilation properties of measurable Schur multipliers and Fourier multipliers. Preprint, arXiv:2202.06575. [22](#)
- [DtER03] N. Dungey, A.F.M. ter Elst and D. Robinson. Analysis on Lie groups with polynomial growth. Progress in Mathematics, 214. Birkhäuser Boston, Inc., Boston, MA, 2003. viii+312 pp. [39](#)
- [ER] E. Effros and Z.-J. Ruan. Operator spaces. Oxford University Press, 2000. [2](#), [12](#)
- [EFHN] T. Eisner, B. Farkas, M. Haase and R. Nagel. Operator theoretic aspects of ergodic theory. Graduate Texts in Mathematics, 272. Springer, Cham, 2015. [45](#)
- [EiW1] M. Eidsiedler and T. Ward. Functional analysis, spectral theory, and applications. Graduate Texts in Mathematics, 276. Springer, Cham, 2017. [18](#), [19](#)
- [Eng] R. Engelking. General topology. Translated from the Polish by the author. Second edition. Sigma Series in Pure Mathematics, 6. Heldermann Verlag, Berlin, 1989. [38](#)
- [Ern1] J. Ernest. A new group algebra for locally compact groups. Amer. J. Math. 86 (1964), 467–492. [6](#)



- [Eym] P. Eymard. L'algèbre de Fourier d'un groupe localement compact (French). Bull. Soc. Math. France 92 (1964), 181–236. [2](#), [12](#), [13](#)
- [Fel1] J. M. G. Fell. The dual spaces of  $C^*$ -algebras. Trans. Amer. Math. Soc. 94 (1960), 365–403. [7](#)
- [Fol] G. Folland. A course in abstract harmonic analysis. Second edition. Textbooks in Mathematics. CRC Press, Boca Raton, FL, 2016. xiii+305 pp.+loose errata. [51](#)
- [FSW] Forrest, Spronk and Wiersma. Existence of tracial states on reduced group  $C^*$ -algebras. Preprint, arXiv:1706.05354. [17](#)
- [GJP17] A. González-Pérez, M. Junge and J. Parcet. Smooth Fourier multipliers in group algebras via Sobolev dimension. Ann. Sci. Éc. Norm. Supér. (4) 50 (2017), no. 4, 879–925. [30](#)
- [Glu1] V. M. Gluskov. The structure of locally compact groups and Hilbert's fifth problem. Amer. Math. Soc. Transl. (2) 15 1960 55–93. [41](#)
- [Haa2] U. Haagerup. On the dual weights for crossed products of von Neumann algebras. II. Application of operator-valued weights. Math. Scand. 43 (1978/79), no. 1, 119–140. [23](#)
- [Haa] U. Haagerup. Injectivity and decomposition of completely bounded maps. Operator algebras and their connections with topology and ergodic theory (Buşteni, 1983), 170–222, Lecture Notes in Math., 1132, Springer, Berlin, 1985. [2](#), [4](#), [10](#), [14](#), [15](#), [51](#), [52](#), [53](#), [54](#)
- [Haa3] U. Haagerup. Group  $C^*$ -algebras without the completely bounded approximation property. J. Lie Theory 26 (2016), no. 3, 861–887. [54](#)
- [Hah1] P. Hahn. The regular representations of measure groupoids. Trans. Amer. Math. Soc. 242 (1978), 35–72. [8](#)
- [HKST15] J. Heinonen, P. Koskela, N. Shanmugalingam and J. T. Tyson. Sobolev spaces on metric measure spaces. An approach based on upper gradients. New Mathematical Monographs, 27. Cambridge University Press, Cambridge, 2015. [39](#), [41](#)
- [HR] E. Hewitt and K. A. Ross. Abstract harmonic analysis. Vol. I. Structure of topological groups, integration theory, group representations. Second edition. Grundlehren der Mathematischen Wissenschaften, 115. Springer-Verlag, Berlin-New York, 1979. [41](#)
- [HeR2] E. Hewitt and K. A. Ross. Abstract harmonic analysis. Vol. II: Structure and analysis for compact groups. Analysis on locally compact Abelian groups. Die Grundlehren der mathematischen Wissenschaften, Band 152, Springer-Verlag, New York-Berlin, 1970. [21](#)
- [Hol1] R. B. Holmes. Geometric functional analysis and its applications. Graduate Texts in Mathematics, No. 24. Springer-Verlag, New York-Heidelberg, 1975. [45](#)
- [JuR] M. Junge and Z.-J. Ruan. Decomposable maps on non-commutative  $L_p$ -spaces. Operator algebras, quantization, and noncommutative geometry, 355–381, Contemp. Math., 365, Amer. Math. Soc., Providence, RI, 2004. [2](#)
- [KaR1] R. V. Kadison and J. R. Ringrose. Fundamentals of the theory of operator algebras. Vol. I. Elementary theory. Reprint of the 1983 original. Graduate Studies in Mathematics, 15. American Mathematical Society, Providence, RI, 1997. [13](#)
- [KaL1] E. Kaniuth and A. T.-M. Lau. Fourier and Fourier-Stieltjes algebras on locally compact groups. Mathematical Surveys and Monographs, 231. American Mathematical Society, Providence, RI, 2018. [2](#), [6](#), [7](#), [11](#), [13](#), [21](#), [30](#), [52](#)
- [Kus05] J. Kustermans. Locally compact quantum groups. Quantum independent increment processes. I, 99–180, Lecture Notes in Math., 1865, Springer, Berlin, 2005. [24](#)
- [Kun] R. A. Kunze.  $L_p$  Fourier transforms on locally compact unimodular groups. Trans. Amer. Math. Soc. 89 (1958), 519–540. [23](#)
- [LaS] V. Lafforgue and M. de la Salle. Noncommutative  $L^p$ -spaces without the completely bounded approximation property. Duke Math. J. 160 (2011), no. 1, 71–116. [36](#), [45](#)
- [Lan1] S. Lang.  $SL_2(\mathbb{R})$ . Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1975. [52](#)
- [LaP2] A. T.-M. Lau and A. L. T. Paterson. Inner amenable locally compact groups. Trans. Amer. Math. Soc. 325 (1991), no. 1, 155–169. [3](#), [17](#), [18](#)
- [LoR1] V. Losert and H. Rindler. Conjugation-invariant means. Colloq. Math. 51 (1987), 221–225. [18](#), [19](#)

- [Meg1] R. E. Megginson. An introduction to Banach space theory. Graduate Texts in Mathematics, 183. Springer-Verlag, New York, 1998. [19](#), [46](#), [47](#)
- [Mil1] P. Milnes. Identities of group algebras. Proc. Amer. Math. Soc. 29 (1971), 421–422. [6](#), [13](#)
- [MZ] D. Montgomery and L. Zippin. Topological transformation groups. Interscience Publishers, New York-London, 1955. xi+282 pp. [41](#)
- [Mos1] R. D. Mosak. Central functions in group algebras. Proc. Amer. Math. Soc. 29 (1971), 613–616.
- [Muh1] P. S. Muhly. Coordinates in operator algebras. Preprint. [7](#), [8](#)
- [NR] S. Neuwirth and É. Ricard. Transfer of Fourier multipliers into Schur multipliers and sumsets in a discrete group. Canad. J. Math. 63 (2011), no. 5, 1161–1187. [26](#), [49](#)
- [Nev06] A. Nevo. Pointwise ergodic theorems for actions of groups. Handbook of dynamical systems. Vol. 1B, 871–982, Elsevier B. V., Amsterdam, 2006. [40](#)
- [Pal2] T. W. Palmer. Banach algebras and the general theory of  $*$ -algebras. Vol. 2. Encyclopedia of Mathematics and its Applications, 79. Cambridge University Press, Cambridge, 2001. [17](#), [51](#)
- [Pau] V. Paulsen. Completely bounded maps and operator algebras. Cambridge Univ. Press, 2002. [10](#), [47](#)
- [Pas1] B. Pasynkov. The coincidence of various definitions of dimensionality for locally bicomact groups. Dokl. Akad. Nauk SSSR 132 1035–1037 (Russian); translated as Soviet Math. Dokl. 1 1960 720–722. [38](#)
- [Pat] A. L. T. Paterson. Amenability. Mathematical Surveys and Monographs, 29. American Mathematical Society, Providence, RI, 1988. [17](#)
- [Pat1] A. L. T. Paterson. The Fourier-Stieltjes and Fourier algebras for locally compact groupoids. Trends in Banach spaces and operator theory (Memphis, TN, 2001), 223–237, Contemp. Math., 321, Amer. Math. Soc., Providence, RI, 2003. [8](#), [9](#)
- [Pat2] A. L. T. Paterson. The Fourier algebra for locally compact groupoids. Canad. J. Math. 56 (2004), no. 6, 1259–1289. [8](#), [9](#)
- [Pat3] A. L. T. Paterson. Groupoids, inverse semigroups, and their operator algebras. Progress in Mathematics, 170. Birkhäuser Boston, Inc., Boston, MA, 1999. [8](#), [9](#)
- [Pat4] A. L. T. Paterson. The class of locally compact groups  $G$  for which  $C^*(G)$  is amenable. Harmonic analysis (Luxembourg, 1987), 226–237, Lecture Notes in Math., 1359, Springer, Berlin, 1988. [18](#)
- [Pea1] A. R. Pears. Dimension theory of general spaces. Cambridge University Press, Cambridge, England-New York-Melbourne, 1975. [38](#)
- [Ped] G. K. Pedersen.  $C^*$ -algebras and their automorphism groups. London Mathematical Society Monographs, 14. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1979. [23](#)
- [Pie] J.-P. Pier. Amenable locally compact groups. Pure and Applied Mathematics (New York). A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1984. [19](#), [51](#)
- [Pis4] G. Pisier. Regular operators between non-commutative  $L_p$ -spaces. Bull. Sci. Math. 119 (1995), no. 2, 95–118. [33](#)
- [Pis5] G. Pisier. Non-commutative vector valued  $L_p$ -spaces and completely  $p$ -summing maps. Astérisque, 247, 1998. [28](#), [30](#), [33](#), [44](#)
- [Pis6] G. Pisier. Similarity problems and completely bounded maps, volume 1618 of Lecture Notes in Mathematics. Springer-Verlag, expanded edition, 2001. [3](#), [14](#), [17](#), [52](#)
- [Pis7] G. Pisier. Introduction to operator space theory. Cambridge University Press, Cambridge, 2003. [2](#), [16](#)
- [Pis17] G. Pisier. Tensor Products of  $C^*$ -Algebras and Operator Spaces. Book. [16](#)
- [Pon1] L. S. Pontryagin. Topological groups. Second Edition. Gordon and Breach, 1966. [38](#), [41](#)
- [RaW1] A. Ramsay and M. E. Walter. Fourier-Stieltjes algebras of locally compact groupoids. J. Funct. Anal. 148 (1997), no. 2, 314–367. [8](#), [9](#)
- [RaW2] A. Ramsay and M. E. Walter. Selfadjoint and nonselfadjoint operator algebras and operator theory (Fort Worth, TX, 1990). Contemp. Math., 120, Amer. Math. Soc., Providence, RI, 1991, 143–156. [8](#)
- [Ren1] J. Renault. The Fourier algebra of a measured groupoid and its multipliers. J. Funct. Anal. 145 (1997), no. 2, 455–490. [8](#), [9](#)

- [Ren2] J. Renault. A groupoid approach to  $C^*$ -algebras. Lecture Notes in Mathematics, 793. Springer, Berlin, 1980. [7](#), [8](#), [9](#)
- [Ren3] J. Renault. Groupoid cocycles and derivations. *Ann. Funct. Anal.* 3 (2012), no. 2, 1–20. [8](#)
- [Spr1] N. Spronk. Measurable Schur multipliers and completely bounded multipliers of the Fourier algebras. *Proc. London Math. Soc.* (3) 89 (2004), no. 1, 161–192. [14](#), [16](#), [48](#), [52](#)
- [Spr2] N. Spronk. On Multipliers of the Fourier Algebra of a Locally Compact Group. PhD Thesis. Waterloo, 2002. [16](#)
- [SS] A. Sinclair and R. Smith. Finite von Neumann algebras and masas. London Mathematical Society Lecture Note Series, 351. Cambridge University Press, Cambridge, 2008. [17](#)
- [Str] S. Stratila, Modular theory in operator algebras. Translated from the Romanian by the author. Editura Academiei Republicii Socialiste România, Bucharest, Abacus Press, Tunbridge Wells, 1981. [18](#), [23](#)
- [StZ] S. Stratila and L. Zsido. Lectures on von Neumann algebras. Translated from the Romanian by Silviu Teleman. Editura Academiei, Bucharest; Abacus Press, Tunbridge Wells, 1975. [23](#)
- [Stro1] M. Stroppel. Locally compact groups. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2006. [38](#), [41](#)
- [Str1] R. A. Struble. Metrics in locally compact groups. *Compositio Math.* 28 (1974), 217–222. [24](#)
- [Tao1] T. Tao. Hilbert’s fifth problem and related topics. Graduate Studies in Mathematics, 153. American Mathematical Society, Providence, RI, 2014. [38](#), [41](#)
- [Tri] H. Triebel. Interpolation theory, function spaces, differential operators. Second edition. Johann Ambrosius Barth, Heidelberg, 1995. [4](#)
- [Vae1] S. Vaes. Locally compact quantum groups. With Johan Kustermans, Leonid Vainerman, Alfons Van Daele and Stanislaw Woronowicz. Lecture Notes School/Conference on Noncommutative Geometry and Quantum groups, Warsaw, 2001, Banach Centre Publications, to appear. Preliminary version. Internet. [24](#)
- [VSCC92] N. Varopoulos, L. Saloff-Coste and T. Coulhon. Analysis and geometry on groups. Cambridge Tracts in Mathematics, 100. Cambridge University Press, Cambridge, 1992. [39](#)
- [Wal1] M. E. Walter. On a new method for defining the norm of Fourier-Stieltjes algebras. *Pacific J. Math.* 137 (1989), no. 1, 209–223. [6](#), [12](#)
- [Wal2] M. E. Walter. Dual algebras. *Math. Scand.* 58 (1986), no. 1, 77–104. [6](#)
- [Wal3] M. E. Walter. Group duality and isomorphisms of Fourier and Fourier-Stieltjes algebras from a  $W^*$ -algebra point of view. *Bull. Amer. Math. Soc.* 76 (1970), 1321–1325. [6](#)
- [Wal4] M. E. Walter. A duality between locally compact groups and certain Banach algebras. *J. Funct. Anal.* 17 (1974), 131–160. [2](#)
- [Wot1] P. Wojtaszczyk. Banach spaces for analysts. Cambridge Studies in Advanced Mathematics, 25. Cambridge University Press, Cambridge, 1991. [3](#)

Cédric Arhancet  
 13 rue Didier Daurat, 81000 Albi, France  
 URL: <https://sites.google.com/site/cedricarhancet>  
 cedric.arhancet@protonmail.com

Christoph Kriegler  
 Université Clermont Auvergne  
 CNRS  
 LMBP  
 F-63000 CLERMONT-FERRAND  
 FRANCE  
 URL: <https://lmbp.uca.fr/~kriegler/indexenglish.html>  
 christoph.kriegler@uca.fr