

Points of Bounded Height on Weighted Projective Spaces Over Global Function Fields

Tristan Phillips

Abstract

In this note we give an asymptotic formula for the number of rational points of bounded height on weighted projective spaces over global function fields.

1 Introduction

In 1979 Schanuel proved a beautiful asymptotic formula for the number of rational points of bounded height on projective space \mathbb{P}^n over a number field K [Sch79]. In 1998 Schanuel's result was generalized to the case of weighted projective spaces by Deng [Den98] (see also [BN20, Theorem 3.7]). Recently Darda has given a new proof of Deng's result using height zeta functions [Dar21].

Theorem 1.1 (Schanuel-Deng-Darda). *Let K be a number field of degree d over \mathbb{Q} , with class number h , regulator R , discriminant $\Delta(K)$, r_1 real places, r_2 complex places, Dedekind zeta function $\zeta_K(s)$, and which contains ω roots of unity. Let $\mathbf{w} = (w_0, \dots, w_n)$ be an $(n+1)$ -tuple of positive integers and set $|\mathbf{w}| := w_0 + w_1 + \dots + w_n$ and*

$$\gcd(\mathbf{w}, \omega) := \gcd\{w_0, w_1, \dots, w_n, \omega\}.$$

Then the number of K -rational points on the weighted projective space $\mathbb{P}(\mathbf{w})$ of height at most B is

$$\frac{hR(2^{r_1}(2\pi)^{r_2})^{n+1}|w|^{r_1+r_2-1} \gcd(\mathbf{w}, \omega)}{\zeta_K(|\mathbf{w}|)|\Delta(K)|^{(n+1)/2}\omega} B^{|\mathbf{w}|} + \begin{cases} O(B \log(B)) & \text{if } \mathbb{P}(\mathbf{w})(K) = \mathbb{P}^1(\mathbb{Q}) \\ O(B^{|\mathbf{w}| - \min_i\{w_i\}/d}) & \text{otherwise.} \end{cases}$$

Schanuel's Theorem, which is the case of Theorem 1.1 with $\mathbf{w} = (1, \dots, 1)$, was independently extended to global function fields by DiPippo [Dip90] and Wan [Wan92]. In order to state their result, we set some notation.

Let X be an absolutely irreducible smooth projective curve of genus g over \mathbb{F}_q . Let $\mathbb{F}_q(X)$ denote the function field of X and h the class number of $\mathbb{F}_q(X)$.

Let $\text{Div}(X)$ denote the set of divisors on X and let

$$\text{Div}^+(X) := \left\{ \sum_P n_P P \in \text{Div}(X) : n_P \geq 0 \text{ for all } P \right\}$$

denote the set of effective divisors on X . Define by

$$Z(X, t) := \exp \left(\sum_{d=1}^{\infty} \#X(\mathbb{F}_{q^d}) \frac{t^d}{d} \right) = \sum_{D \in \text{Div}^+(X)} t^{\deg(D)}$$

the classical zeta function of X . Let $\zeta_X(s) = Z(X, q^{-s})$ denote the Dedekind zeta function of X .

We define the *height* of a $\mathbb{F}_q(X)$ -rational point of projective space $x = [x_0 : x_1 : \cdots : x_n] \in \mathbb{P}^n(\mathbb{F}_q(X))$ by

$$\text{ht}_X(x) := -\deg(\inf_i(x_i))$$

where $\inf_i(x_i)$ denotes the greatest divisor D of X such that $D \leq (x_i)$ for all i . Let

$$A_d(\mathbb{P}^n) := \{x \in \mathbb{P}^n(\mathbb{F}_q(X)) : \text{ht}_X(x) = d\}$$

denote the number of $\mathbb{F}_q(X)$ -rational points on \mathbb{P}^n of height d .

Theorem 1.2 (DiPippo-Wan). *With notation as above, for any $\varepsilon > 0$ there exist polynomials $b_i(d)$ in d , whose coefficients are algebraic numbers, and algebraic integers α_i , with $|\alpha_i| = \sqrt{q}$, such that*

$$\begin{aligned} A_d(\mathbb{P}^n) &= \frac{hq^{(n+1)(1-g)}}{\zeta_X(n+1)(q-1)} q^{(n+1)d} + \sum_{i=1}^{2g} b_i(d) \alpha_i^d \\ &= \frac{hq^{(n+1)(1-g)}}{\zeta_X(n+1)(q-1)} q^{(n+1)d} + O(q^{d/2+\varepsilon}). \end{aligned}$$

In this article we extend the Theorem of DiPippo and Wan to the case of weighted projective spaces, giving an analog of Theorem 1.1 for global function fields. Let $A_d(\mathbf{w})$ denote then number of $\mathbb{F}_q(X)$ -rational points of bounded height on the weighted projective space $\mathbb{P}(\mathbf{w})$.

Theorem 1.3. *Let notation be as above and set $w_{\min} := \min_i\{w_i\}$. Then for any $\varepsilon > 0$ there exist polynomials $b_i(d)$ in d , whose coefficients are algebraic numbers, and algebraic integers α_i , with $|\alpha_i| = \sqrt{q}$, such that the number of K -rational points on the weighted projective space $\mathbb{P}(\mathbf{w})$ of height d is*

$$\begin{aligned} A_d(\mathbf{w}) &= \sum_{\substack{\mathbf{u} \in 2^{\mathbf{w}} \\ |\mathbf{u}| \geq 2}} \sum_{\substack{\mathbf{v} \in 2^{\mathbf{w}} \\ \mathbf{u} \subseteq \mathbf{v}}} \frac{\gcd(\mathbf{v}, q-1) (-1)^{\#\mathbf{v} - \#\mathbf{u}} hq^{\#\mathbf{u}(1-g)}}{(q-1)\zeta_X(|\mathbf{u}|)} q^{d\#\mathbf{u}} + \sum_{i=1}^{2g} b_i(d) \alpha_i^d \\ &= \frac{hq^{\#\mathbf{w}(1-g)} \gcd(\mathbf{w}, q-1)}{\zeta_X(|\mathbf{w}|)(q-1)} q^{|\mathbf{w}|d} + O(q^{d(|\mathbf{w}|-w_{\min})}). \end{aligned}$$

One motivation for counting points on weighted projective spaces is that they are an example of a stack. In particular, this result fits into the more general framework of counting points of bounded height on stacks, as outlined in [ESZB21]. Just as Schanuel's theorem was a catalyst for the Batyrev-Manin Conjecture (for counting points on projective varieties) we hope that this note will motivate further work on counting points in weighted projective varieties. A weak version of the Batyrev-Manin conjecture for stacks has been proposed in [ESZB21, Conjecture 4.14], and it seems likely that a stronger version should hold if one is willing to restrict attention to stacks that arise as weighted projective varieties.

2 Weighted projective spaces

Definition 2.1 (weighted projective space). Given an $(n+1)$ -tuple of positive integers $\mathbf{w} = (w_0, w_1, \dots, w_n)$, the *weighted projective space* $\mathbb{P}(\mathbf{w})$ is defined to be the quotient stack

$$\mathbb{P}(\mathbf{w}) = [(\mathbb{A}^{n+1} - \{0\})/\mathbb{G}_m],$$

with respect to the action

$$\begin{aligned} *_{\mathbf{w}} : \mathbb{G}_m \times (\mathbb{A}^{n+1} - \{0\}) &\rightarrow (\mathbb{A}^{n+1} - \{0\}) \\ (\lambda, (x_0, \dots, x_n)) &\mapsto \lambda *_{\mathbf{w}}(x_0, \dots, x_n) := (\lambda^{w_0}x_0, \dots, \lambda^{w_n}x_n). \end{aligned}$$

Note that the weighted projective space with weights $\mathbf{w} = (1, 1, \dots, 1)$ is the same as projective n -space, \mathbb{P}^n . Over fields of characteristic p the weighted projective space $\mathbb{P}(w_0, \dots, w_n)$ is a Deligne-Mumford stack if and only if $p \nmid w_i$ for all i .

Definition 2.2 (weighted divisor). Let $y \in \mathbb{F}_q(X)$ and $w \in \mathbb{Z}_{>0}$. The w -weighted divisor of y is defined to be

$$(y)_w := \sum_{P \in X} \left\lfloor \frac{\text{ord}_P(y)}{w} \right\rfloor P$$

where P ranges over the points of X .

Definition 2.3 ((logarithmic) height). The (logarithmic) height of a $\mathbb{F}_q(X)$ -rational point, $y = [y_0 : y_1 : \dots : y_n] \in \mathbb{P}(\mathbf{w})(\mathbb{F}_q(X))$, of weighted projective space is defined as

$$\text{ht}_{\mathbf{w}, X}(y) := -\deg \left(\inf_i ((y_i)_{w_i}) \right).$$

Remark 2.4. This height can also be realized as the *stacky height* associated to the tautological bundle $\mathcal{O}_{\mathbb{P}(\mathbf{w})}(1)$ of $\mathbb{P}(\mathbf{w})$ [ESZB21].

We shall often abbreviate $\text{ht}_{\mathbf{w}, X}$ to $\text{ht}_{\mathbf{w}}$ when the function field is clear from context.

Define a counting function

$$A_d(\mathbf{w}) := \#\{y \in \mathbb{P}(\mathbf{w})(\mathbb{F}_q(X)) : \text{ht}_{\mathbf{w}, X}(y) = d\}$$

which counts the number of $\mathbb{F}_q(X)$ -rational points of height d on $\mathbb{P}(\mathbf{w})$. Note that

$$A_d(\underbrace{1, 1, \dots, 1}_{n+1}) = A_d(\mathbb{P}^n).$$

We now define weighted versions of linear equivalence, Picard groups, and Riemann-Roch spaces.

Definition 2.5 (w -linearly equivalent divisors). Let $w \in \mathbb{Z}_{>0}$. Two divisors $D, D' \in \text{Div}(X)$ are said to be w -linearly equivalent if $D - D' = (f)_w$ for some $f \in \mathbb{F}_q(X)$.

Definition 2.6 (w -Picard group). The w -Picard group is defined to be the group of w -equivalence classes of divisors on X , and is denoted $\text{Pic}_w(X)$.

Definition 2.7 (w -Riemann-Roch space). For $D \in \text{Div}(X)$ we define the w -Riemann-Roch space attached to D as

$$\mathcal{L}_w(D) := \{f \in \mathbb{F}_q(X) : (f)_w + D \geq 0\} \cup \{0\}.$$

Set $\ell_w(D) := \dim_{\mathbb{F}_q}(\mathcal{L}_w(D))$.

When $w = 1$ we recover the usual definitions of Picard group and Riemann-Roch space. In these cases we will write $\text{Pic}(X)$ for $\text{Pic}_1(X)$, $\mathcal{L}(D)$ for $\mathcal{L}_1(D)$, and $\ell(D)$ for $\ell_1(D)$.

3 Weighted divisor zeta functions

Let $\mathbf{w} = (w_0, \dots, w_n)$ be an $(n+1)$ -tuple of positive integers. Let $\#\mathbf{w}$ denote the length of \mathbf{w} , let $|\mathbf{w}|$ denote the sum of components $w_0 + w_1 + \dots + w_n$, and let $2^{\mathbf{w}}$ denote the set of sub-tuples of \mathbf{w} . Denote by $\gcd(\mathbf{w}, q-1)$ the greatest common divisor of the set $\{w_0, w_1, \dots, w_n, q-1\}$.

Consider the \mathbf{w} -weighted action of \mathbb{F}_q^\times on $\mathbb{F}_q(X)^{\#\mathbf{w}} - \{0\}$ induced by $*_{\mathbf{w}}$. With respect to this action, define the following set of \mathbb{F}_q^\times -orbits of points in the affine cone of $\mathbb{P}(\mathbf{w})$:

$$B_{\mathbf{w}}(D) := \{x = (x_0, \dots, x_n) \in \frac{\mathbb{F}_q(X)^{\#\mathbf{w}} - \{0\}}{\mathbb{F}_q^\times} : -\inf_i (x_i)_{w_i} \leq D\}.$$

Choose a degree one divisor D^* on X once and for all.

Definition 3.1 (weighted divisor zeta function). Let D_1, \dots, D_h generate the torsion part of the divisor class group. For $\mathbf{w} = (w_0, \dots, w_n)$ define the \mathbf{w} -weighted divisor zeta function of X to be

$$Z_{\mathbf{w}}(X, t) := \sum_{j=1}^h \sum_{d=0}^{\infty} \#B_{\mathbf{w}}(D_j + dD^*) t^d.$$

Theorem 3.2 (Rationality of weighted divisor zeta functions). *Let X be a non-singular genus g curve and let $\mathbf{w} = (w_0, \dots, w_n)$ be an $(n+1)$ -tuple of positive integers. Then, for each $\mathbf{u} \in 2^{\mathbf{w}}$, there exists an integral polynomial $P_{\mathbf{u}}(t)$ of degree at most $1 + (2g-2)/\min_i(u_i)$ such that*

$$Z_{\mathbf{w}}(X, t) = \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \sum_{\mathbf{u} \in 2^{\mathbf{v}}} \frac{\gcd(\mathbf{v}, q-1) (-1)^{\#\mathbf{v} - \#\mathbf{u}} P_{\mathbf{u}}(t)}{(q-1)(1 - q^{|\mathbf{u}|}t)}.$$

Proof. For $\mathbf{u} = (u_0, \dots, u_r) \in \mathbb{Z}_{>0}^{r+1}$, set

$$Z(\mathbf{u}, t) := \sum_{j=1}^h \sum_{d=0}^{\infty} \left(\prod_{u_i \in \mathbf{u}} q^{\ell(u_i(D_j + dD^*))} \right) t^d.$$

Riemann-Roch implies that for any divisor D of degree one, if $u_i d > 2g-2$ then $\ell(u_i d D) = u_i d - g + 1$. Therefore there are integral polynomials $Q_{\mathbf{u}}(t)$ and $P_{\mathbf{u}}(t)$, of degrees at most $(2g-2)/\min_i(u_i)$ and $1 + (2g-2)/\min_i(u_i)$ respectively, such that

$$\begin{aligned} Z(\mathbf{u}, t) &= Q_{\mathbf{u}}(t) + \sum_{d=0}^{\infty} h \prod_{j=0}^{\#\mathbf{u}-1} q^{u_j d - g + 1} t^d \\ &= Q_{\mathbf{u}}(t) + q^{\#\mathbf{u}(1-g)} h \sum_{d=0}^{\infty} q^{|\mathbf{u}|d} t^d \\ &= Q_{\mathbf{u}}(t) + \frac{h q^{\#\mathbf{u}(1-g)}}{1 - q^{|\mathbf{u}|}t} \\ &= \frac{P_{\mathbf{u}}(t)}{1 - q^{|\mathbf{u}|}t}. \end{aligned}$$

Let $x = (x_0, \dots, x_n) \in B_{\mathbf{w}}(D)$ and let $\mathbf{v} \subset 2^{\mathbf{w}}$ be the subtuple of \mathbf{w} corresponding to the non-zero coordinates of x (e.g., if $\mathbf{w} = (1, 2, 3)$ and $x = (2, 0, 7)$, then $\mathbf{v} = (1, 3)$). Then the number of

elements of $\mathbb{F}_q(X)^{\#\mathbf{w}} - \{0\}$ contained in x is $\gcd(\mathbf{v}, q-1)/(q-1)$. It follows that

$$\begin{aligned} \#B_{\mathbf{w}}(D) &= \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \frac{\gcd(\mathbf{v}, q-1)}{q-1} \prod_{v_i \in \mathbf{v}} (\#\mathcal{L}_{v_i}(D) - 1) \\ &= \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \frac{\gcd(\mathbf{v}, q-1)}{q-1} \sum_{\mathbf{u} \in 2^{\mathbf{v}}} (-1)^{\#\mathbf{v} - \#\mathbf{u}} \prod_{u_i \in \mathbf{u}} q^{\ell_{u_i}(D)}. \end{aligned}$$

It follows that

$$\begin{aligned} Z_{\mathbf{w}}(X, t) &= \sum_{j=1}^h \sum_{d=0}^{\infty} \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \frac{\gcd(\mathbf{v}, q-1)}{q-1} \sum_{\mathbf{u} \in 2^{\mathbf{v}}} (-1)^{\#\mathbf{v} - \#\mathbf{u}} \left(\prod_{u_i \in \mathbf{u}} q^{\ell_{u_i}(D_j + dD^*)} \right) t^d \\ &= \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \frac{\gcd(\mathbf{v}, q-1)}{q-1} \sum_{\mathbf{u} \in 2^{\mathbf{v}}} (-1)^{\#\mathbf{v} - \#\mathbf{u}} Z(\mathbf{u}, t) \\ &= \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \sum_{\mathbf{u} \in 2^{\mathbf{v}}} \frac{\gcd(\mathbf{v}, q-1) (-1)^{\#\mathbf{v} - \#\mathbf{u}} P_{\mathbf{u}}(t)}{(q-1)(1 - q^{|\mathbf{u}|}t)}. \end{aligned}$$

□

4 Height zeta functions of weighted projective spaces

Definition 4.1 (height zeta function). Define the *height zeta function* of the weighted projective space $\mathbb{P}(\mathbf{w})$ defined over the function field $\mathbb{F}_q(X)$ of X to be

$$Z(\mathbb{P}(\mathbf{w})(\mathbb{F}_q(X)), t) := \sum_{d=0}^{\infty} \#A_d(\mathbf{w}) t^d.$$

Theorem 4.2. *The height zeta function is related to the divisor zeta function as follows,*

$$Z(\mathbb{P}(\mathbf{w})(\mathbb{F}_q(X)), t) = \frac{Z_{\mathbf{w}}(X, t)}{Z(X, t)}.$$

Proof. For a divisor $D \in \text{Div}(X)$ define the set

$$N(D) := \{D' \in \text{Div}^+(X) : [D] = [D'] \text{ in } \text{Pic}(X)\}$$

of effective divisors linearly equivalent to D .

We now show that there is a bijection

$$\bigsqcup_{\substack{[D'] \in \text{Div}^+(X) \\ \deg(D') = d' \leq d}} N(D') \times A_{d-d'}(W) \leftrightarrow \bigsqcup_{\substack{[D] \in \text{Pic}(X) \\ \deg(D) = d}} B_W(D).$$

Elements on the left can be specified by pairs (D'', z) , where $D'' \in N(D')$ and $z \in A_{d-d'}(W)$. Elements on the right can be specified by pairs (D, y) , where D is a degree d divisor and $y \in B_W(D)$. The bijection is then given by

$$\begin{aligned} (D'', z) &\mapsto (D'' - \inf_i (z_i)_{w_i}, z) \\ (D + \inf_i (y_i)_{w_i}, y) &\leftarrow (D, y). \end{aligned}$$

From this it follows that

$$Z(X, t) \cdot Z(\mathbb{P}(W)(\mathbb{F}_q(X)), t) = Z_{\mathbf{w}}(X, t).$$

□

Theorem 4.2 together with Theorem 3.2 shows that $Z(\mathbb{P}(W)(\mathbb{F}_q(X)), t)$ is a rational function. From this we can derive an exact formula for $A_d(\mathbf{w})$:

Theorem 1.3. *Let X be a non-singular projective curve of genus g whose global function, $\mathbb{F}_q(X)$, has class number h . Let $\zeta_X(s) = Z(X, q^{-s})$ be the Dedekind zeta function of X . Then there exist algebraic integers α_i with $|\alpha_i| = \sqrt{q}$ and polynomials $b_i(d)$ in d whose coefficients are algebraic numbers such that*

$$\#A_d(\mathbf{w}) = \sum_{\substack{\mathbf{u} \in 2^{\mathbf{w}} \\ |\mathbf{u}| \geq 2}} \sum_{\substack{\mathbf{v} \in 2^{\mathbf{w}} \\ \mathbf{u} \subseteq \mathbf{v}}} \frac{\gcd(\mathbf{v}, q-1)(-1)^{\#\mathbf{v}-\#\mathbf{u}} h q^{\#\mathbf{u}(1-g)}}{(q-1)\zeta_X(|\mathbf{u}|)} q^{d\#\mathbf{u}} + \sum_{i=1}^{2g} b_i(d) \alpha_i^d.$$

Proof. It is well known that the classical zeta function $Z(X, t)$ is a rational function and can be written as

$$Z(X, t) = \frac{P(t)}{(1-t)(1-q^t)},$$

where $P(t) = \prod_{i=1}^{2g} (1 - \alpha_i t)$ with α_i algebraic integers satisfying $|\alpha_i| = \sqrt{q}$. Combining this fact with Theorem 3.2 we have that

$$\frac{Z_{\mathbf{w}}(X, t)}{Z(X, t)} = \sum_{\mathbf{v} \in 2^{\mathbf{w}}} \sum_{\mathbf{u} \in 2^{\mathbf{v}}} \frac{\gcd(\mathbf{v}, q-1)(-1)^{\#\mathbf{v}-\#\mathbf{u}} (1-t)(1-qt) P_{\mathbf{u}}(t)}{(q-1)(1-q^{|\mathbf{u}|}t) P(t)}. \quad (4.1)$$

Note that this is a rational function which has only simple poles. These poles can only possibly occur at values of t in the following set,

$$\{q^{-|\mathbf{u}|} : \mathbf{u} \in 2^{\mathbf{w}} \text{ with } |\mathbf{u}| \geq 2\}.$$

Note also that the degree of the numerator in (4.1) is at most $2g$ since $\deg(P_{\mathbf{u}}(t)) \leq 2g - 2$. By Theorem 4.2 and a partial fractions decomposition we have that there are rational numbers $a_{\mathbf{u}}$ for $\mathbf{u} \in 2^{\mathbf{w}}$, and polynomials $b_i(d)$ for $1 \leq i \leq 2g$, such that for all $d \geq 1$,

$$\#A_d(\mathbf{w}) = \sum_{\substack{\mathbf{u} \in 2^{\mathbf{w}} \\ |\mathbf{u}| \geq 2}} a_{\mathbf{u}} q^{d\#\mathbf{u}} + \sum_{i=1}^{2g} b_i(d) \alpha_i^d.$$

Noting that

$$a_{\mathbf{u}} = \sum_{\substack{\mathbf{v} \in 2^{\mathbf{w}} \\ \mathbf{u} \subseteq \mathbf{v}}} \frac{\gcd(\mathbf{v}, q-1)(-1)^{\#\mathbf{v}-\#\mathbf{u}} h q^{\#\mathbf{u}(1-g)}}{(q-1)\zeta_X(|\mathbf{u}|)},$$

we obtain the desired formula. □

Corollary 4.4. *For $\mathbf{w} = (w_0, \dots, w_n)$ set $w_{\min} := \min_i \{w_i\}$. Then we have the following asymptotic in d :*

$$A_d(\mathbf{w}) = \frac{h q^{\#\mathbf{w}(1-g)} \gcd(\mathbf{w}, q-1)}{\zeta_X(|\mathbf{w}|)(q-1)} q^{d|\mathbf{w}|} + O(q^{d(|\mathbf{w}|-w_{\min})}).$$

5 Acknowledgements

The author would like to thank Bryden Cais and C. Douglas Haessig for their encouragement during this project.

References

- [BN20] Peter Bruin and Filip Najman. Counting elliptic curves with prescribed level structures over number fields, 2020. Preprint, [arXiv:2008.05280](#).
- [Dar21] Ratko Darda. Rational points of bounded height on weighted projective stacks, 2021. Preprint, [arXiv:2106.10120](#).
- [Den98] An-Wen Deng. Rational points on weighted projective spaces, 1998. Preprint, [arXiv:9812082](#).
- [Dip90] Stephen Ascanio Dipippo. *Spaces of rational functions on curves over finite fields*. ProQuest LLC, Ann Arbor, MI, 1990. Thesis (Ph.D.)—Harvard University.
- [ESZB21] Jordan S. Ellenberg, Matthew Satriano, and David Zureick-Brown. Heights on stacks and a generalized batyrev-manin-malle conjecture, 2021. Preprint, [arXiv:92106.11340](#).
- [Sch79] Stephen Hoel Schanuel. Heights in number fields. *Bull. Soc. Math. France*, 107(4):433–449, 1979.
- [Wan92] Da Qing Wan. Heights and zeta functions in function fields. In *The arithmetic of function fields (Columbus, OH, 1991)*, volume 2 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 455–463. de Gruyter, Berlin, 1992.