

**A note about extension of functions
belonging to Sobolev - Grand Lebesgue Spaces.**

M.R.Formica, E.Ostrovsky, L.Sirota.

Università degli Studi di Napoli Parthenope, via Generale Parisi 13, Palazzo Pacanowsky, 80132, Napoli, Italy.
e-mail: mara.formica@uniparthenope.it

Department of Mathematics and Statistics, Bar-Ilan University, 59200, Ramat Gan, Israel.
e-mail: eugostrovsky@list.ru

Department of Mathematics and Statistics, Bar-Ilan University, 59200, Ramat Gan, Israel.
e-mail: sirota3@bezeqint.net

Abstract

We deduce an extension theorem for the so - called Sobolev - Grand Lebesgue Spaces defined on the suitable subsets of the whole finite - dimensional Euclidean space, and estimate the norms of correspondent extension operator, which may be choosed as linear.

Key words and phrases. Lebesgue - Riesz, Yudovich, ordinary Sobolev and Sobolev - Grand Lebesgue norm and spaces, Euclidean space, Lipschitz domain, ordinary and linear extension, linear bounded operator, norm, measurable functions, semi - space, generating function, estimation.

1 Introduction. Previous results.

Let $B(G)$, where G is non - trivial subset of an Euclidean space $R^d : G \subset R^d$ be a family of Banach spaces defined on the class of functions defined in turn on the support G ; the norm in this space will be denoted $\|f\|_{B(G)}$.

It will be presumed henceforth that all the considered domains G are closures of the non - empty open sets and are Lipschitzian. The case when $G = R^d$ is trivial for us and may be excluded.

Put also for definiteness $B = B(R^d)$.

For instance, the classical Lebesgue - Riesz $L_p(G)$ spaces equipped with the ordinary norm

$$\|f\|_p(G) = \|f\|_{L_p(G)} := \left[\int_G |f(x)|^p dx \right]^{1/p}, \quad p \in [1, \infty), \quad f : G \rightarrow R; \quad x \in G,$$

as well as the famous Sobolev spaces $W_p^m(G)$, $m = 1, 2, \dots$:

$$\|f\|_{W_p^m(G)} \stackrel{def}{=} \max_{\alpha, |\alpha| \leq m} \|D^\alpha f\|_{L_p(G)}, \quad (1)$$

$\|f\|_{W_p^m} := \|f\|_{W_p^m(R^d)}$; see e.g. [1], [24], [25], [26], [30], [31].
Here as ordinary

$$\alpha = \vec{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_d); \quad \alpha_j = 0, 1, 2, \dots;$$

$$|\alpha| := \sum_{j=1}^d \alpha_j; \quad D^\alpha f := \frac{D^{|\alpha|} f}{\partial x_1^{\alpha_1} \partial x_2^{\alpha_2} \dots \partial x_d^{\alpha_d}}$$

and all the derivatives are understood in the weak (Sobolev) sense. Of course, $D^0 f = f$.

Let the function $f : G \rightarrow R$ belongs to some space $B(G)$. By definition, the function $\tilde{f} : R^d \rightarrow R$ is named as an *extension* of the function f (from the set G), iff

$$\forall x \in G \Rightarrow \tilde{f}(x) = f(x)$$

and wherein $\|\tilde{f}\|_B < \infty$.

If there exists a *linear bounded* operator $L : \tilde{f} = Lf = L_G f$, $f \in B(G)$, where again \tilde{f} is an extension for f , such that

$$K = K(G) = K(B, G) \stackrel{def}{=} \sup_{0 \neq f \in B(G)} \left\{ \frac{\|Lf\|_B}{\|f\|_{B(G)}} \right\} < \infty,$$

then the extension $\tilde{f} = Lf = L_G f$ is named linear.

It is known, see e.g. [3], [4], [11], [31] that under imposed conditions on the Lipschitz domain G and for the Sobolev's spaces $B(G) = W_p^m(G)$, $m \geq 1$, $p \geq 1$ (the case $m = 0$ is trivial) there exist a linear extension operator. See also the works [18], [19], [25], [26], [29] etc.

We will ground in offered report that this linear operator there exists also for the so - called Sobolev - Grand Lebesgue Spaces.

2 Main result.

SOBOLEV - GRAND LEBESGUE SPACES.

Let $(a, b) = \text{const}, 1 \leq a < b \leq \infty$. Let also $\psi = \psi(p), p \in (a, b)$ be certain numerical valued measurable strictly positive: $\inf_{p \in (a, b)} \psi(p) > 0$ function, not necessary to be bounded. Denotation:

$$(a, b) := \text{supp}(\psi); \Psi[a, b] := \{ \psi : \text{supp}(\psi) = (a, b) \},$$

$$\Psi \stackrel{\text{def}}{=} \cup_{(a, b): 1 \leq a < b \leq \infty} \Psi[a, b].$$

Definition 2.1., see e.g. [28]. The Sobolev - Grand Lebesgue Space $S[m, G, \psi]$ based on the set $G, G \subset R^d$ is defined as a set of all (measurable) functions having a finite norm

$$\|f\|S[m, G, \psi] \stackrel{\text{def}}{=} \sup_{p \in (a, b)} \left\{ \frac{\|f\|W_p^m(G)}{\psi(p)} \right\}. \quad (2)$$

The particular case $m = 0$, i.e. when

$$\|f\|G\psi = \|f\|G\psi(a, b) \stackrel{\text{def}}{=} \|f\|S[0, G, \psi] = \sup_{p \in (a, b)} \left\{ \frac{\|f\|L_p(G)}{\psi(p)} \right\} \quad (3)$$

and under some additional restrictions on the *generating function* $\psi = \psi(p)$ correspondent to the so - called *Yudovich spaces*, see [32], [33]. These spaces was applied at first in the theory of Partial Differential Equations (PDE), see [7], [8].

A general case of these spaces when $m = 0$ are named as the classical Grand Lebesgue Spaces (GLS) $G\psi, \psi \in \Psi$. These spaces are investigated in many works, see e.g. [9], [10], [12], [13], [14], [15], [16], [17], [20], [21], [22], [23], [27]. The general case of Sobolev - Grand Lebesgue Spaces appears at first perhaps in the article [28], where was investigated the modulus of continuity of the functions belonging to these spaces.

Theorem 2.1. Assume that all the formulated before restrictions are satisfied, indeed: that the Lipschitz domain G is closure of the non - empty open subset of whole Euclidean space R^d . We propose that for arbitrary Sobolev - Grand Lebesgue Space $S[m, G, \psi]$ there exists a *linear bounded* extension operator $L = L_G$.

Proof. One can suppose $d \geq 2$ and that $G = \vec{x} = \{x_j\} = \{x_1, x_2, \dots, x_{d-1}, x_d\}$, where $\vec{x} = (\tilde{x}, x_d)$; $\tilde{x} = \{x_1, x_2, \dots, x_{d-1}\}$; so that $\vec{x} \in G \Leftrightarrow x_d \geq 0$; on the other words, upper semi - space; see e.g. [3], [4], [11]. Let us define the following extension operator $Lf(x) := f(x), x \in G, f(\cdot) \in S[m, G, \psi]$;

$$Lf(x) := \sum_{k=1}^{d+1} c_k f(\tilde{x}, -kx_d), \quad x_d < 0.$$

The coefficients $\{c_k\}$ may be uniquely determined from the following system of linear equations

$$\sum_{k=1}^{m+1} (-k)^l c_k = 1; \quad l = 0, 1, \dots, d.$$

Suppose that $f \in S[m, G, \psi]$; one can assume without loss of generality $\|f\|_{S[m, G, \psi]} = 1$. Then for all the values $p \in (a, b)$

$$\|f\|_{W_p^m} \leq \psi(p), \quad p \in (a, b),$$

therefore

$$\forall p \in (a, b), \quad \forall \alpha : |\alpha| \leq m \Rightarrow \|D^\alpha f\|_p(G) \leq \psi(p).$$

Introduce the functions

$$g_k(\tilde{x}, y) := f(\tilde{x}, -ky), \quad y \leq 0;$$

then

$$D^\alpha g_k = (-k)^{\alpha_d} f^{(\alpha)}(\tilde{x}, -ky),$$

$$\|D^\alpha g_k\|_p(G) = k^{\alpha_d - 1/p} \|D^\alpha f\|_p(G) \leq k^{\alpha_d} \|D^\alpha f\|_p(G),$$

therefore

$$\begin{aligned} \forall \alpha : |\alpha| \leq m \Rightarrow \sum_{k=1}^{m+1} |c_k| k^m \cdot \|D^\alpha f\|_p(G) \leq \\ \sum_{k=1}^{m+1} |c_k| k^m \cdot \psi(p), \quad p \in (a, b). \end{aligned}$$

Following, by virtue of triangle inequality for Lebesgue - Riesz spaces

$$\|L[f]\|_{W_p^m(R^d \setminus G)} \leq C(d, m) \psi(p), \quad C(d, m) < \infty,$$

$$\|Lf\|_{S[m, R^d, \psi]} \leq 1 + C(d, m) < \infty,$$

Q.E.D.

3 Concluding remarks.

It is interest in our opinion to compute the exact value of extension constant for Sobolev - Grand Lebesgue Spaces, as well as to generalize the extension theorem on the anisotropic spaces.

Acknowledgement. The first author has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and by Università degli Studi di Napoli Parthenope through the project “sostegno alla Ricerca individuale”.

References

- [1] **Adams, R.A.** (1975), *Sobolev spaces*. Academic Press, New York.
- [2] **V.V. Buldygin V.V., D.I.Mushtary, E.I.Ostrovsky, M.I.Pushalsky.** *New Trends in Probability Theory and Statistics*. Mokslas, (1992), V.1, p. 78 - 92; Amsterdam, Utrecht, New York, Tokyo.
- [3] **Burenkov V. I.** *On a certain method for extending differentiable functions.* (Russian). Trudy Mat. Inst. Steklov 140 (1976), 27 - 67; English translation.: Proc. of Steklov Inst. Math.; **140**, (1976).
- [4] **Burenkov V. I.** *On the extension of functions with preservation of semi-norm.* (Russian). Dokl. Akad. Nauk SSSR, **228**, (1976), 779 - 782; English translation.: Soviet Math. Dokl. **17**, (1976). MR 0412796
- [5] **Capone C, Formica M.R, Giova R.** *Grand Lebesgue spaces with respect to measurable functions.* Nonlinear Analysis 2013; 85: 125 - 131.
- [6] **Capone C, and Fiorenza A.** *On small Lebesgue spaces. Journal of function spaces and applications.* 2005; 3; 73 - 89.
- [7] **Qionglei Chen, Changxing Miao and Xiaoxin Zheng.** *The two - dimensional Euler equation in Yudovich and bmo - type spaces.* arXiv:1311.0934v4 [math.AP] 10 Jan 2019
- [8] **Gianluca Crippa and Giorgio Stefani.** *An elementary proof of existence and uniqueness for the Euler flow in localized Yudovich spaces.* arXiv:2110.15648v1 [math.AP] 29 Oct 2021
- [9] **S. V. Ermakov, E. I. Ostrovsky.** *Central limit theorem for weakly dependent Banach - space valued random variables.* Theory Probab. Appl., **30**, 2, (1986), 391 - 394.
- [10] **S.V.Ermakov, and E. I. Ostrovsky.** *Continuity Conditions, Exponential Estimates, and the Central Limit Theorem for Random Fields.* Moscow, VINITI, (1986), (in Russian).
- [11] **Charles L.Fefferman, Arie Israel, and Garving K. Luli.** *Sobolev extension by linear operators.* Dordrecht, (2012).
- [12] **Fiorenza A., and Karadzhov G.E.** *Grand and small Lebesgue spaces and their analogs.* Consiglio Nazionale Delle Ricerche, Istituto per le Applicazioni del Calcolo Mauro Picone, Sezione di Napoli, Rapporto tecnicon. 272/03, (2005).

- [13] **A. Fiorenza, M. R. Formica and A. Gogatishvili.** *On grand and small Lebesgue and Sobolev spaces and some applications to PDE's.* *Differ. Equ. Appl.* **10** (2018), no. 1, 21–46.
- [14] **A. Fiorenza, M. R. Formica, A. Gogatishvili, T. Kopaliani and J. M. Rakotoson.** *Characterization of interpolation between grand, small or classical Lebesgue spaces.* Preprint arXiv:1709.05892, *Nonlinear Anal.*, to appear.
- [15] **A. Fiorenza, M. R. Formica and J. M. Rakotoson.** *Pointwise estimates for GT-functions and applications.* *Differential Integral Equations* **30** (2017), no. 11-12, 809–824.
- [16] **M. R. Formica and R. Giova.** *Boyd indices in generalized grand Lebesgue spaces and applications.* *Mediterr. J. Math.* **12** (2015), no. 3, 987–995.
- [17] **M.R.Formica, E.Ostrovsky, L.Sirota** *Central Limit Theorem in Lebesgue-Riesz spaces for weakly dependent random sequences.*
arXiv:1912.00338v2 [math.PR] 3 Dec 2019
- [18] **Miguel Garcia, Ia - Bravo, Tapio Rajala, and Jurki Takanen.** *Two - sides boundary points of Sobolev - extension domains on Euclidean spaces.*
arXiv:2111.01079v1 [math.CA] 1 Nov 2021
- [19] **P. Koskela, D.Yang, and Y. Zhou.** *A Jordan Sobolev extension domain.* *Ann. Acad. Sci. Fenn. Math.* **35**, (2010), no. 1, 309 - 320.
- [20] **Yu.V. Kozachenko and E.I. Ostrovsky.** *Banach spaces of random variables of subgaussian type.* *Theory Probab. Math. Stat.*, Kiev, (1985), v. 43, 42 - 56, (in Russian).
- [21] **Kozachenko Yu.V., Ostrovsky E., Sirota L.** *Relations between exponential tails, moments and moment generating functions for random variables and vectors.*
arXiv:1701.01901v1 [math.FA] 8 Jan 2017
- [22] **Kozachenko Yu.V., Ostrovsky E., Sirota L.** *Equivalence between tails, Grand Lebesgue Spaces and Orlicz norms for random variables without Kramer's condition.* *Bulletin of KSU*, Kiev, 2018, **4**, pp. 20 - 29.
- [23] **E.Liflyand, E.Ostrovsky, L.Sirota.** *Structural Properties of Bilateral Grand Lebesgue Spaces.* *Turk. J. Math.*; **34**, (2010), 207 - 219.
- [24] **Leoni, Giovanni.** (2009). *A First Course in Sobolev Spaces.* Graduate Studies in Mathematics, **105**. American Mathematical Society. pp. xvi + 607.
ISBN 978-0-8218-4768-8. MR 2527916. Zbl 1180.46001.
- [25] **V. Mazya.** *Extension of functions from Sobolev spaces.* *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, **113**, (1981), 231 - 236.
- [26] **Maz'ja, Vladimir G.** (1985). *Sobolev Spaces.* Springer Series in Soviet Mathematics, Berlin-HeidelbergNew York: Springer-Verlag, pp. xix+486,
doi:10.1007/978-3-662-09922-3, ISBN 0-387-13589-8, MR 0817985, Zbl 0692.46023
- [27] **E.I. Ostrovsky.** *Exponential Estimations for Random Fields.* Moscow - Obninsk, OINPE, (1999), in Russian.
- [28] **Ostrovsky E., Sirota L.** *Module of continuity for the functions belonging to the Sobolev-Grand Lebesgue Spaces.*
arXiv:1006.4177v1 [math.FA] 21 Jun 2010
- [29] **A.S.Romanov.** *On the extension of functions that belong to Sobolev spaces.* *Sibirsk. Mat. Zh.*, **34**, (1993), 149 - 152; English transl. in *Siberian Math. J.* **34**, (1993), 723 - 726.
- [30] **Sobolev, S.L.** (1963). *Some applications of functional analysis in mathematical physics.* Amer. Math. Soc.
- [31] **Steve Taylor.** *An introduction to Sobolev spaces.* (2015), American Mathematical Society. pp. xvi + 769.

- [32] **V. I. Yudovich.** *Nonstationary flow of an ideal incompressible liquid.* Zh. Vych. Mat., **3**, (1963), 1032 - 1066.
- [33] **V. I. Yudovich.** *Uniqueness theorem for the basic nonstationary problem in the dynamics of an ideal incompressible fluid.* Mathematical Research Letters, **2**. (1995), 27 - 38.