

Jacobi's Inversion Problem for Genus Two Hyperelliptic Integral II

Kazuyasu Shigemoto¹

Tezukayama University, Tezukayama 7, Nara 631, Japan

Abstract

In the previous paper, we reviewed the Rosenhain's paper to the Jacobi's inversion problem for the genus two hyperelliptic integral. In this paper, we review the Göpel's paper to the Jacobi's inversion problem for the genus two hyperelliptic integral.

1 Introduction

The doubly periodic function, the elliptic function, is obtained as the ratio of the one variable theta function. While, according to Abel, the elliptic function is obtained as the solution of the inversion problem of the elliptic integral. Similarly, the multiply periodic function such as the genus two hyperelliptic function, is obtained as the ratio of the two variables theta function. If we obtain such multiply periodic function as the inversion problem of the genus two hyperelliptic integral, Jacobi noticed that the naive solution of the inversion problem provides the non-single valued multiply periodic function [1]. Jacobi found that the symmetric combination of the hyperelliptic integral becomes the single valued function. Thus, Jacobi proposed the following Jacobi's inversion problem

$$u = \int^x \frac{dx}{\sqrt{f_5(x)}} + \int^{x'} \frac{dx'}{\sqrt{f_5(x')}}, \quad v = \int^x \frac{x dx}{\sqrt{f_5(x)}} + \int^{x'} \frac{x' dx'}{\sqrt{f_5(x')}},$$

for the fifth-degree polynomial function $f_5(x)$. For this inversion problem, Jacobi conjectured that the multiply periodic function, which is the solution of the Jacobi's inversion problem, is given by the symmetric combination such as $x + x'$, xx' , which become the single valued functions of u and v ; furthermore, $x + x'$, xx' are expressed as the ratio of the two variable theta functions [2].

The Jacobi's inversion problem for the genus two case is solved by Göpel [3, 4] and independently by Rosenhain [5, 6].

In Rosenhain's approach, the Riemann type addition formula of the hyperelliptic theta function is used. The key identities are the three quadratic theta identities, which parametrize five ratios of the square of hyperelliptic functions, which naturally gives the fifth-degree polynomial function $f_5(x)$. While, in Göpel's approach, the theta formula by the duplication method of the hyperelliptic theta function is used. The key identities are three quartic theta identities. One of these identities is the Kummer's quartic identity, i.e., Kummer surface relation.

¹E-mail address: shigemot@tezukayama-u.ac.jp

In the previous paper, we reviewed the Rosenhain's paper [7]. In this paper, we review Göpel's paper.

2 The addition formulae of the genus two hyperelliptic theta functions -Duplication method-

The quadratic addition formula, which is called the duplication method, was first introduced by Jacobi [8] in 1828 for the the genus one elliptic theta functions. Göpel [3, 4] used this addition formula for the genus two hyperelliptic theta functions in 1847. Using this addition formula, we provided the full addition formula for the genus two hyperelliptic functions instead of the hyperelliptic theta functions [9].

The quartic addition formula, which is called the Riemann's theta formula, was first introduced by Jacobi [10] in 1838 for the genus one elliptic theta functions. Rosenhain [5, 6] used this addition formula for the genus two hyperelliptic theta functions in 1850. Prym [11] named such addition formula as *the Riemann's theta formula* because he writes down the proof according to the direct suggestion by Riemann [12] at Pisa in 1865. Using this addition formula, the full addition formula for the genus two hyperelliptic functions instead of the hyperelliptic theta functions are given by Kossak [13, 14].

The theta functions with two variables are defined by

$$\begin{aligned} \vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y; \tau_{11}, \tau_{22}, \tau_{12}) \\ = \sum_{m, n \in \mathbb{Z}} \exp \left\{ \pi i \left(\tau_{11} \left(m + \frac{a}{2} \right)^2 + \tau_{22} \left(n + \frac{c}{2} \right)^2 + 2\tau_{12} \left(m + \frac{a}{2} \right) \left(n + \frac{c}{2} \right) \right) \right. \\ \left. + 2\pi i \left(\left(m + \frac{a}{2} \right) \left(x + \frac{b}{2} \right) + \left(n + \frac{c}{2} \right) \left(y + \frac{d}{2} \right) \right) \right\}, \end{aligned} \quad (2.1)$$

where we assume that $\text{Im}\tau_{11} > 0$, $\text{Im}\tau_{22} > 0$, $(\text{Im}\tau_{11})(\text{Im}\tau_{22}) - (\text{Im}\tau_{12})^2 > 0$ in order that the summation of $m, n \in \mathbb{Z}$ becomes convergent. Renaming $m \rightarrow m$, $n \rightarrow -n$, we can always choose $\text{Im}\tau_{12} > 0$, thus we assume $\text{Im}\tau_{12} > 0$.

We simply denote $\vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y) = \vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y; \tau_{11}, \tau_{22}, \tau_{12})$.

We also use the notation $\Theta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y) = \vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y; 2\tau_{11}, 2\tau_{22}, 2\tau_{12})$. These have the following properties

$$\begin{aligned} \vartheta \left[\begin{array}{c} a+2 \ c \\ b \ d \end{array} \right] (x, y) &= \vartheta \left[\begin{array}{c} a \ c+2 \\ b \ d \end{array} \right] (x, y) = \vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y), \\ \vartheta \left[\begin{array}{c} a \ c \\ b+2 \ d \end{array} \right] (x, y) &= (\sqrt{-1})^{2a} \vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y), \quad \vartheta \left[\begin{array}{c} a \ c \\ b \ d+2 \end{array} \right] (x, y) = (\sqrt{-1})^{2c} \vartheta \left[\begin{array}{c} a \ c \\ b \ d \end{array} \right] (x, y). \end{aligned}$$

In order to obtain the addition formula according to the duplication method, we consider

the product of the ϑ function in the form

$$\begin{aligned}
& \vartheta \begin{bmatrix} a & c \\ b & d \end{bmatrix} (x_1, y_1) \vartheta \begin{bmatrix} e & g \\ f & h \end{bmatrix} (x_2, y_2) \\
&= \sum_{m,n,p,q \in \mathbb{Z}} \exp \left\{ \pi i \left(\tau_{11} \left(m + \frac{a}{2} \right)^2 + \tau_{22} \left(n + \frac{c}{2} \right)^2 + 2\tau_{12} \left(m + \frac{a}{2} \right) \left(n + \frac{c}{2} \right) \right. \right. \\
&\quad \left. \left. + \tau_{11} \left(p + \frac{e}{2} \right)^2 + \tau_{22} \left(q + \frac{g}{2} \right)^2 + 2\tau_{12} \left(p + \frac{e}{2} \right) \left(q + \frac{g}{2} \right) \right) \right. \\
&\quad \left. + 2\pi i \left(\left(m + \frac{a}{2} \right) \left(x_1 + \frac{b}{2} \right) + \left(n + \frac{c}{2} \right) \left(y_1 + \frac{d}{2} \right) + \left(p + \frac{e}{2} \right) \left(x_2 + \frac{f}{2} \right) + \left(q + \frac{g}{2} \right) \left(y_2 + \frac{h}{2} \right) \right) \right\}, \\
&= \sum_{m_1, n_1, m_2, n_2 \in \mathbb{Z}} \exp \left\{ \frac{\pi i}{2} \left(\tau_{11} \left(m_1 + \frac{a+e}{2} \right)^2 + \tau_{22} \left(n_1 + \frac{c+g}{2} \right)^2 + 2\tau_{12} \left(m_1 + \frac{a+e}{2} \right) \left(n_1 + \frac{c+g}{2} \right) \right. \right. \\
&\quad \left. \left. + \tau_{11} \left(m_2 + \frac{a-e}{2} \right)^2 + \tau_{22} \left(n_2 + \frac{c-g}{2} \right)^2 + 2\tau_{12} \left(m_2 + \frac{a-e}{2} \right) \left(n_2 + \frac{c-g}{2} \right) \right) \right. \\
&\quad \left. + \pi i \left(\left(m_1 + \frac{a+e}{2} \right) \left(x_1 + x_2 + \frac{b+f}{2} \right) + \left(n_1 + \frac{c+g}{2} \right) \left(y_1 + y_2 + \frac{d+h}{2} \right) \right. \right. \\
&\quad \left. \left. + \left(m_2 + \frac{a-e}{2} \right) \left(x_1 - x_2 + \frac{b-f}{2} \right) + \left(n_2 + \frac{c-g}{2} \right) \left(y_1 - y_2 + \frac{d-h}{2} \right) \right) \right\},
\end{aligned}$$

where we use $m_1 = m + p, m_2 = m - p, n_1 = n + q, n_2 = n - q$. With $m_1 - m_2 = 2p =$ (even number), $\{m_1, m_2\}$ are both even number or odd number. The pair $\{n_1, n_2\}$ are also both even number or odd number. Hence, we obtain 4 cases i) m_1, m_2 : even number; n_1, n_2 : even number, ii) m_1, m_2 : even number; n_1, n_2 : odd number, iii) m_1, m_2 : odd number; n_1, n_2 : even number, iv) m_1, m_2 : odd number; n_1, n_2 : odd number. Therefore, we obtain

$$\begin{aligned}
& \vartheta \begin{bmatrix} a & c \\ b & d \end{bmatrix} (x_1, y_1) \vartheta \begin{bmatrix} e & g \\ f & h \end{bmatrix} (x_2, y_2) \\
&= \Theta \begin{bmatrix} \frac{a+e}{2} & \frac{c+g}{2} \\ b+f & d+h \end{bmatrix} (x_1 + x_2, y_1 + y_2) \Theta \begin{bmatrix} \frac{a-e}{2} & \frac{c-g}{2} \\ b-f & d-h \end{bmatrix} (x_1 - x_2, y_1 - y_2) \\
&+ \Theta \begin{bmatrix} \frac{a+e}{2} & \frac{c+g+2}{2} \\ b+f & d+h \end{bmatrix} (x_1 + x_2, y_1 + y_2) \Theta \begin{bmatrix} \frac{a-e}{2} & \frac{c-g+2}{2} \\ b-f & d-h \end{bmatrix} (x_1 - x_2, y_1 - y_2) \\
&+ \Theta \begin{bmatrix} \frac{a+e+2}{2} & \frac{c+g}{2} \\ b+f & d+h \end{bmatrix} (x_1 + x_2, y_1 + y_2) \Theta \begin{bmatrix} \frac{a-e+2}{2} & \frac{c-g}{2} \\ b-f & d-h \end{bmatrix} (x_1 - x_2, y_1 - y_2) \\
&+ \Theta \begin{bmatrix} \frac{a+e+2}{2} & \frac{c+g+2}{2} \\ b+f & d+h \end{bmatrix} (x_1 + x_2, y_1 + y_2) \Theta \begin{bmatrix} \frac{a-e+2}{2} & \frac{c-g+2}{2} \\ b-f & d-h \end{bmatrix} (x_1 - x_2, y_1 - y_2).
\end{aligned}$$

Putting $x_1 = u_1 + u_2, x_2 = u_1 - u_2, y_1 = v_1 + v_2, y_2 = v_1 - v_2$, we express the above formula

in the form

$$\begin{aligned}
& \vartheta \begin{bmatrix} a & c \\ b & d \end{bmatrix} (u_1 + u_2, v_1 + v_2) \vartheta \begin{bmatrix} e & g \\ f & h \end{bmatrix} (u_1 - u_2, v_1 - v_2) \\
&= \Theta \begin{bmatrix} \frac{a+e}{2} & \frac{c+g}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e}{2} & \frac{c-g}{2} \\ b-f & d-h \end{bmatrix} (2u_2, 2v_2) \\
&+ \Theta \begin{bmatrix} \frac{a+e}{2} & \frac{c+g+2}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e}{2} & \frac{c-g+2}{2} \\ b-f & d-h \end{bmatrix} (2u_2, 2v_2) \\
&+ \Theta \begin{bmatrix} \frac{a+e+2}{2} & \frac{c+g}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e+2}{2} & \frac{c-g}{2} \\ b-f & d-h \end{bmatrix} (2u_2, 2v_2) \\
&+ \Theta \begin{bmatrix} \frac{a+e+2}{2} & \frac{c+g+2}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e+2}{2} & \frac{c-g+2}{2} \\ b-f & d-h \end{bmatrix} (2u_2, 2v_2). \tag{2.2}
\end{aligned}$$

In the special case of $u_2 = 0, v_2 = 0$, we obtain

$$\begin{aligned}
& \vartheta \begin{bmatrix} a & c \\ b & d \end{bmatrix} (u_1, v_1) \vartheta \begin{bmatrix} e & g \\ f & h \end{bmatrix} (u_1, v_1) \\
&= \Theta \begin{bmatrix} \frac{a+e}{2} & \frac{c+g}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e}{2} & \frac{c-g}{2} \\ b-f & d-h \end{bmatrix} (0, 0) \\
&+ \Theta \begin{bmatrix} \frac{a+e}{2} & \frac{c+g+2}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e}{2} & \frac{c-g+2}{2} \\ b-f & d-h \end{bmatrix} (0, 0) \\
&+ \Theta \begin{bmatrix} \frac{a+e+2}{2} & \frac{c+g}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e+2}{2} & \frac{c-g}{2} \\ b-f & d-h \end{bmatrix} (0, 0) \\
&+ \Theta \begin{bmatrix} \frac{a+e+2}{2} & \frac{c+g+2}{2} \\ b+f & d+h \end{bmatrix} (2u_1, 2v_1) \Theta \begin{bmatrix} \frac{a-e+2}{2} & \frac{c-g+2}{2} \\ b-f & d-h \end{bmatrix} (0, 0). \tag{2.3}
\end{aligned}$$

We frequently use in the following form

$$\begin{pmatrix} \vartheta \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix}^2(u, v) \\ \vartheta \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}^2(u, v) \\ \vartheta \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix}^2(u, v) \\ \vartheta \begin{bmatrix} a & b \\ 1 & 1 \end{bmatrix}^2(u, v) \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} \Theta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} (0, 0) \\ \Theta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a & b-1 \\ 0 & 0 \end{bmatrix} (0, 0) \\ \Theta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a-1 & b \\ 0 & 0 \end{bmatrix} (0, 0) \\ \Theta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a-1 & b-1 \\ 0 & 0 \end{bmatrix} (0, 0) \end{pmatrix}.$$

Here the Riemann matrix, which is the self-adjoint orthogonal matrix, emerges. We inversely solve the above in the following matrix form

$$\begin{pmatrix} \Theta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} (0, 0) \\ \Theta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a & b-1 \\ 0 & 0 \end{bmatrix} (0, 0) \\ \Theta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a-1 & b \\ 0 & 0 \end{bmatrix} (0, 0) \\ \Theta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} a-1 & b-1 \\ 0 & 0 \end{bmatrix} (0, 0) \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} \vartheta \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix}^2(u, v) \\ \vartheta \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}^2(u, v) \\ \vartheta \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix}^2(u, v) \\ \vartheta \begin{bmatrix} a & b \\ 1 & 1 \end{bmatrix}^2(u, v) \end{pmatrix}.$$

We use the Göpel's simplified notation of the form

$$\begin{aligned}
P_0 &= \vartheta \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}(u, v), & P_1 &= \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v), & P_2 &= \vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}(u, v), & P_3 &= \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}(u, v), \\
Q_0 &= \vartheta \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}(u, v), & Q_1 &= \vartheta \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}(u, v), & Q_2 &= \vartheta \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}(u, v), & Q_3 &= \vartheta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(u, v), \\
R_0 &= \vartheta \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}(u, v), & R_1 &= \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}(u, v), & R_2 &= \vartheta \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}(u, v), & R_3 &= \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(u, v), \\
S_0 &= \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}(u, v), & S_1 &= \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u, v), & S_2 &= \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}(u, v), & S_3 &= \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}(u, v), \\
T &= \Theta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}(2u, 2v), & U &= \Theta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(2u, 2v), & V &= \Theta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(2u, 2v), & W &= \Theta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}(2u, 2v), \\
t &= \Theta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}(0, 0), & u &= \Theta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(0, 0), & v &= \Theta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(0, 0), & w &= \Theta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}(0, 0).
\end{aligned} \tag{2.4}$$

Using the Göpel's notation, we obtain

$$\begin{pmatrix} P_3^2 \\ P_1^2 \\ P_2^2 \\ P_0^2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} tT \\ vV \\ uU \\ wW \end{pmatrix}, \quad \begin{pmatrix} tT \\ vV \\ uU \\ wW \\ P_0^2 \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} P_3^2 \\ P_1^2 \\ P_2^2 \\ P_0^2 \end{pmatrix}, \tag{2.5}$$

$$\begin{pmatrix} Q_3^2 \\ Q_1^2 \\ Q_2^2 \\ Q_0^2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} uT \\ wV \\ tU \\ vW \end{pmatrix}, \quad \begin{pmatrix} uT \\ wV \\ tU \\ vW \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} Q_3^2 \\ Q_1^2 \\ Q_2^2 \\ Q_0^2 \end{pmatrix}, \tag{2.6}$$

$$\begin{pmatrix} R_3^2 \\ R_1^2 \\ R_2^2 \\ R_0^2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} vT \\ tV \\ wU \\ uW \end{pmatrix}, \quad \begin{pmatrix} vT \\ tV \\ wU \\ uW \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} R_3^2 \\ R_1^2 \\ R_2^2 \\ R_0^2 \end{pmatrix}, \tag{2.7}$$

$$\begin{pmatrix} S_3^2 \\ S_1^2 \\ S_2^2 \\ S_0^2 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} wT \\ uV \\ vU \\ tW \end{pmatrix}, \quad \begin{pmatrix} wT \\ uV \\ vU \\ tW \end{pmatrix} = \frac{1}{4} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} S_3^2 \\ S_1^2 \\ S_2^2 \\ S_0^2 \end{pmatrix}. \tag{2.8}$$

3 The Kummer surface relation

We use the Kummer's quartic relation, i.e., Kummer surface relation, to solve the Jacobi's inversion relation. Thus, we derive the Kummer's quartic relation. By using the addition formula of P_1 , S_1 , P_2 and S_2 , we obtain

$$P_1^2 = tT + uU - vV - wW, \quad P_2^2 = tT - uU + vV - wW, \tag{3.1}$$

$$S_1^2 = wT + vU - uV - tW, \quad S_2^2 = wT - vU + uV - tW. \tag{3.2}$$

We inversely solve in the form

$$T = \frac{t(P_1^2 + P_2^2) - w(S_1^2 + S_2^2)}{2(t^2 - w^2)}, \quad U = \frac{u(P_1^2 - P_2^2) - v(S_1^2 - S_2^2)}{2(u^2 - v^2)}, \quad (3.3)$$

$$V = \frac{v(P_1^2 - P_2^2) - u(S_1^2 - S_2^2)}{2(u^2 - v^2)}, \quad W = \frac{w(P_1^2 + P_2^2) - t(S_1^2 + S_2^2)}{2(t^2 - w^2)}. \quad (3.4)$$

Later, we use the following

$$Q_1^2 = uT + tU - wV - vW, \quad R_1^2 = vT + wU - tV - uW. \quad (3.5)$$

3.1 Proof of $Q_1R_1 = aP_1S_1 + bP_2S_2$

We will demonstrate that $Q_1R_1 = aP_1S_1 + bP_2S_2$, ($a, b = \text{const.}$). Considering the square of this relation, we obtain $Q_1^2R_1^2 = (aP_1S_1 + bP_2S_2)^2$. Using Eqs.(3.3)-(3.5), the left-hand side term $Q_1^2R_1^2$ is expressed as the polynomial of P_1, S_1, P_2, S_2 . While the right-hand side term $(aP_1S_1 + bP_2S_2)^2$ is of course the polynomial of P_1, S_1, P_2, S_2 . Hence, $Q_1^2R_1^2 = (aP_1S_1 + bP_2S_2)^2$ gives the Kummer's quartic relations of P_1, S_1, P_2, S_2 . If we can find one of the expressions, which is expressed as the linear combination of P_1S_1 and P_2S_2 , it suffices for our purpose. For example, instead of using Q_1R_1 , we can use Q_2R_2 because this also becomes the linear combination of P_1S_1 and P_2S_2 . By using the addition formula, we obtain

$$\begin{aligned} Q_1R_1 &= \vartheta \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} (u, v) \\ &= \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 2 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) + \Theta \begin{bmatrix} \frac{1}{2} & \frac{3}{2} \\ 0 & 2 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &\quad + \Theta \begin{bmatrix} \frac{3}{2} & \frac{1}{2} \\ 0 & 2 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) + \Theta \begin{bmatrix} \frac{3}{2} & \frac{3}{2} \\ 0 & 2 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} \frac{3}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0), \end{aligned}$$

which provides

$$\begin{aligned} Q_1R_1 &= i\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) - i\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &\quad + i\Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) - i\Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &= i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &\quad - i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0), \end{aligned} \quad (3.6)$$

where we used

$$\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) = \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0), \quad \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) = \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0). \quad (3.7)$$

Similarly, we obtain

$$\begin{aligned}
P_1 S_1 &= \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} (u, v) \\
&= i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\
&\quad - i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0), \tag{3.8}
\end{aligned}$$

$$\begin{aligned}
P_2 S_2 &= \vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} (u, v) \\
&= i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\
&\quad + i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0). \tag{3.9}
\end{aligned}$$

Combining these relations, we obtain three coupled equations

$$Q_1 R_1 = iBX - iAY, \tag{3.10}$$

$$P_1 S_1 = iAX - iBY, \tag{3.11}$$

$$P_2 S_2 = iAX + iBY, \tag{3.12}$$

$$X = \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right), \tag{3.13}$$

$$Y = \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right), \tag{3.14}$$

$$A = \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0), \quad B = \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0). \tag{3.15}$$

Expressing X and Y by $P_1 S_1$ and $P_2 S_2$, and substituting into the right-hand side of $Q_1 R_1$, we complete the proof of the form

$$Q_1 R_1 = aP_1 S_1 + bP_2 S_2, \quad a = \frac{A^2 + B^2}{2AB}, \quad b = \frac{A^2 - B^2}{2AB}, \quad a^2 - b^2 = 1. \tag{3.16}$$

3.2 The expression of a and b with t, u, v, w

In order to express A and B as the function of t, u, v, w , we consider the quantity

$$\begin{aligned}
P_3 S_3 &= \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} (u, v) \\
&= \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\
&\quad + \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0). \tag{3.17}
\end{aligned}$$

By putting $u = 0, v = 0$, we obtain

$$\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} (0, 0) \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} (0, 0) = 2\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)^2 + 2\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)^2 = 2(A^2 + B^2). \tag{3.18}$$

Similarly, we consider the quantity

$$\begin{aligned}
Q_3 R_3 &= \vartheta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(u, v) \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(u, v) \\
&= \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(0, 0) \\
&\quad + \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(0, 0).
\end{aligned} \tag{3.19}$$

By putting $u = 0, v = 0$, we obtain

$$\vartheta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(0, 0) \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(0, 0) = 4\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(0, 0) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(0, 0) = 4AB. \tag{3.20}$$

Therefore, the constant a is solved by the function of $\vartheta \begin{bmatrix} a & c \\ b & d \end{bmatrix}(0, 0)$ in the form

$$a = \frac{A^2 + B^2}{2AB} = \frac{\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}(0, 0) \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}(0, 0)}{\vartheta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(0, 0) \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(0, 0)}. \tag{3.21}$$

Taking the square and use the addition formula, $S_3(0, 0)^2 = \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}(0, 0)^2 = 2(tw + uv)$,
 $P_3(0, 0)^2 = \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}(0, 0)^2 = t^2 + u^2 + v^2 + w^2$, $Q_3(0, 0)^2 = \vartheta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(0, 0)^2 = 2(tu + vw)$,
 $R_3(0, 0)^3 = \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(0, 0)^2 = 2(tv + uw)$, we obtain the expression of a^2 with t, u, v, w

$$a^2 = \frac{\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}(0, 0)^2 \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}(0, 0)^2}{\vartheta \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}(0, 0)^2 \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}(0, 0)^2} = \frac{(tw + uv)(t^2 + u^2 + v^2 + w^2)}{2(tu + vw)(tv + uw)}. \tag{3.22}$$

By using $a^2 - b^2 = 1$, we obtain b^2 in the form

$$b^2 = a^2 - 1 = \frac{(t^2 - u^2 - v^2 + w^2)(tw - uv)}{2(tu + vw)(tv + uw)}, \tag{3.23}$$

which provides

$$a = \sqrt{\frac{(t^2 + u^2 + v^2 + w^2)(tw + uv)}{2(tu + vw)(tv + uw)}}, \quad b = \sqrt{\frac{(t^2 - u^2 - v^2 + w^2)(tw - uv)}{2(tu + vw)(tv + uw)}}. \tag{3.24}$$

3.3 The Kummer surface relation of P_1, S_1, P_2, S_2

The Kummer's quartic relation for P_1, S_1, P_2, S_2 is given by considering the square of $Q_1 R_1 = aP_1 S_1 + bP_2 S_2$ in the form $Q_1^2 R_1^2 = a^2 P_1^2 S_1^2 + b^2 P_2^2 S_2^2 + 2abP_1 S_1 P_2 S_2$ via Eqs.(3.3)-(3.5). After

the straightforward calculation, we obtain the following Kummer's quartic relation

$$P_1^4 + S_1^4 + P_2^4 + S_2^4 - 2F(P_1^2P_2^2 + S_1^2S_2^2) + 2C(P_1^2S_2^2 + P_2^2S_1^2) - 2E(P_1^2S_1^2 + P_2^2S_2^2) - 4DP_1P_2S_1S_2 = 0, \quad (3.25)$$

$$E = \frac{t^2u^2 + t^2v^2 + u^2w^2 + v^2w^2 + 4tuvw}{2(tu + vw)(tv + uw)}, \quad (3.26)$$

$$F = \frac{t^4 + u^4 + v^4 + w^4 - 2t^2w^2 - 2u^2v^2}{(t^2 + u^2 - v^2 - w^2)(t^2 - u^2 + v^2 - w^2)}, \quad (3.27)$$

$$C = \frac{t^2u^2 + t^2v^2 + u^2w^2 + v^2w^2 - 4tuvw}{2(tu - vw)(tv - uw)}, \quad (3.28)$$

$$D = \frac{(t^2 - w^2)^2(u^2 - v^2)^2 \sqrt{(t^2 + u^2 + v^2 + w^2)(t^2 - u^2 - v^2 + w^2)(t^2w^2 - u^2v^2)}}{(t^2 + u^2 - v^2 - w^2)(t^2 - u^2 + v^2 - w^2)(t^2u^2 - v^2w^2)(t^2v^2 - u^2w^2)}, \quad (3.29)$$

where $C^2 - D^2 + E^2 + F^2 - 2CEF = 1$ is satisfied.

If we use $P_3(0, 0) = a$, $S_3(0, 0) = b$, $P_0(0, 0) = c$, $S_0(0, 0) = d$, we have the relations

$$a^2 = t^2 + u^2 + v^2 + w^2, \quad b^2 = 2(tw + uv), \quad c^2 = t^2 - u^2 - v^2 + w^2, \\ d^2 = 2(tw - uv),$$

by using the addition formula. Thus, we obtain the Kummer surface relation in the following Hudson's standard form [15]

$$P_1^4 + S_1^4 + P_2^4 + S_2^4 + A_1(P_1^2S_2^2 + P_2^2S_1^2) + B_1(P_1^2P_2^2 + S_1^2S_2^2) + C_1(P_1^2S_1^2 + P_2^2S_2^2) + 2D_1P_1P_2S_1S_2 = 0, \quad (3.30)$$

$$A_1 = \frac{b^4 + c^4 - a^4 - d^4}{a^2d^2 - b^2c^2}, \quad B_1 = \frac{a^4 + c^4 - b^4 - d^4}{b^2d^2 - a^2c^2}, \quad C_1 = \frac{a^4 + b^4 - c^4 - d^4}{c^2d^2 - a^2b^2}, \\ D_1 = \frac{abcd(a^2 + d^2 - b^2 - c^2)(b^2 + d^2 - a^2 - c^2)(c^2 + d^2 - a^2 - b^2)(a^2 + b^2 + c^2 + d^2)}{(a^2d^2 - b^2c^2)(b^2d^2 - a^2c^2)(c^2d^2 - a^2b^2)}, \quad (3.31)$$

where $A_1^2 + B_1^2 + C_1^2 - D_1^2 - A_1B_1C_1 = 4$ is satisfied.

4 The differential equation (Step I)

4.1 The derivative formula

In this section, by using $p = S_1/P_1$, $q = S_2/P_2$ and $s = P_1/P_2$ of the form

$$p = \frac{S_1}{P_1} = \frac{\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u, v)}{\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v)}, \quad q = \frac{S_2}{P_2} = \frac{\vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}(u, v)}{\vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}(u, v)}, \quad s = \frac{P_1}{P_2} = \frac{\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v)}{\vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}(u, v)}, \quad (4.1)$$

we first derive the differential equation of $p = S_1/P_1$ and $q = S_2/P_2$. By noticing the relation

$$dp = \frac{1}{P_1^2}(dS_1P_1 - dP_1S_1), \quad dq = \frac{1}{P_2^2}(dS_2P_2 - dP_2S_2), \quad (4.2)$$

we consider the combination

$$I = (dS_1P_1 - dP_1S_1) = d\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u, v)\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v) - d\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v)\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u, v), \quad (4.3)$$

$$J = (dS_2P_2 - dP_2S_2) = d\vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}(u, v)\vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}(u, v) - d\vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}(u, v)\vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}(u, v). \quad (4.4)$$

We derive the derivative formula of the theta function by using the addition formula of the theta function. Thus, we consider

$$\begin{aligned} I_1 &= \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u_1 + u_2, v_1 + v_2)\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u_1 - u_2, v_1 - v_2) \\ &\quad - \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u_1 + u_2, v_1 + v_2)\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u_1 - u_2, v_1 - v_2) \\ &= i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_1, 2v_1) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_1, 2v_1) \right) \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) \right) \\ &\quad - i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_1, 2v_1) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_1, 2v_1) \right) \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) \right). \end{aligned}$$

If we notice that

$$\begin{aligned} &\left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) \right), \\ &\left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u_2, 2v_2) \right), \end{aligned}$$

are odd functions, these odd functions become zero at $u_2 = 0, v_2 = 0$. Hence, by putting $u_2 = du_1, v_2 = 0$, and replacing $u_1 \rightarrow u, v_1 \rightarrow v$, we obtain $\partial\vartheta/\partial u$ in the form

$$\begin{aligned} &\left(\frac{\partial}{\partial u} \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u, v)\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v) - \frac{\partial}{\partial u} \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}(u, v)\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}(u, v) \right) \\ &= i\alpha \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) \right) \\ &\quad - i\beta \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) \right) \\ &= i\alpha f(2u, 2v) - i\beta g(2u, 2v), \end{aligned} \quad (4.5)$$

where

$$f(2u, 2v) = \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) \right), \quad (4.6)$$

$$g(2u, 2v) = \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(2u, 2v) \right), \quad (4.7)$$

$$\alpha = \frac{d}{du} \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(u, 0) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(u, 0) \right) \Big|_{u=0}, \quad (4.8)$$

$$\beta = \frac{d}{du} \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix}(u, 0) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix}(u, 0) \right) \Big|_{u=0}. \quad (4.9)$$

Similarly, putting $u_2 = 0$, $v_2 = dv_1$, and replacing $u_1 \rightarrow u$, $v_1 \rightarrow v$, we obtain $\partial\vartheta/\partial v$ in the form

$$\begin{aligned} & \left(\frac{\partial}{\partial v} \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} (u, v) - \frac{\partial}{\partial v} \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} (u, v) \right) \\ &= i\gamma \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \\ &- i\delta \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \\ &= i\gamma f(2u, 2v) - i\delta g(2u, 2v), \end{aligned} \quad (4.10)$$

$$\gamma = \frac{d}{dv} \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, v) - \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, v) \right) \Big|_{v=0}, \quad (4.11)$$

$$\delta = \frac{d}{dv} \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, v) - \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, v) \right) \Big|_{v=0}. \quad (4.12)$$

Thus, we obtain the derivative formula from the addition formula of the theta function in the form

$$\begin{aligned} I &= (dS_1P_1 - dP_1S_1) = du(\partial_u S_1P_1 - \partial_u P_1S_1) + dv(\partial_v S_1P_1 - \partial_v P_1S_1) \\ &= if(2u, 2v)(\alpha du + \gamma dv) - ig(2u, 2v)(\beta du + \delta dv). \end{aligned} \quad (4.13)$$

We can obtain J by the replacement of $u \rightarrow u + 1$, $v \rightarrow v + 1$ in Eq.(4.13). In the right-hand side of Eq.(4.13), we obtain

$$\begin{aligned} I &= (dS_1P_1 - dP_1S_1) = d\vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} (u, v) - d\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} (u, v) \\ &\rightarrow d\vartheta \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 1 & 2 \end{bmatrix} (u, v) - d\vartheta \begin{bmatrix} 0 & 0 \\ 1 & 2 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} (u, v) \\ &= (-1) \left(d\vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} (u, v) - d\vartheta \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} (u, v) \right) = -J. \end{aligned} \quad (4.14)$$

While, in the right-hand side of Eq.(4.13), we obtain

$$\begin{aligned} & if(2u, 2v)(\alpha du + \gamma dv) - ig(2u, 2v)(\beta du + \delta dv) \\ &= i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) (\alpha du + \gamma dv) \\ &- i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) (\beta du + \delta dv) \\ &\rightarrow i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) \right) (\alpha du + \gamma dv) \\ &- i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) \right) (\beta du + \delta dv) \\ &= -i \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) (\alpha du + \gamma dv) \\ &- i \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) (\beta du + \delta dv) \\ &= -if(2u, 2v)(\alpha du + \gamma dv) - ig(2u, 2v)(\beta du + \delta dv). \end{aligned} \quad (4.15)$$

Thus, we obtain the derivative formula of J in the form²

$$\begin{aligned} J &= (dS_2P_2 - dP_2S_2) = du(\partial_u S_2P_2 - \partial_u P_2S_2) + dv(\partial_v S_2P_2 - \partial_v P_2S_2), \\ &= if(2u, 2v)(\alpha du + \gamma dv) + ig(2u, 2v)(\beta du + \delta dv). \end{aligned} \quad (4.16)$$

Therefore, the necessary derivative formulae provide

$$(dS_1P_1 - dP_1S_1) = if(2u, 2v)(\alpha du + \gamma dv) - ig(2u, 2v)(\beta du + \delta dv), \quad (4.17)$$

$$(dS_2P_2 - dP_2S_2) = if(2u, 2v)(\alpha du + \gamma dv) + ig(2u, 2v)(\beta du + \delta dv). \quad (4.18)$$

Next, we express $f(2u, 2v)$, $g(2u, 2v)$ with the product of the original theta function $P_0, P_1, \dots, S_2, S_3$. Using Eq.(3.17), we obtain

$$\begin{aligned} P_3S_3 &= \vartheta \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} (u, v) \\ &= \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &+ \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &= \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) f(2u, 2v) + \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) g(2u, 2v). \end{aligned} \quad (4.19)$$

By replacing $u \rightarrow u + 1, v \rightarrow v + 1$ in Eq.(4.19), we obtain

$$\begin{aligned} P_0S_0 &= \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} (u, v) \vartheta \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} (u, v) \\ &= \left(\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &+ \left(\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) + \Theta \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ 2 & 2 \end{bmatrix} (2u, 2v) \right) \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) \\ &= -\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) f(2u, 2v) + \Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0) g(2u, 2v). \end{aligned} \quad (4.20)$$

Thus, we can express $f(2u, 2v)$ and $g(2u, 2v)$ with the product of the original theta functions in the form

$$f(2u, 2v) = \frac{P_3S_3 - P_0S_0}{2\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)}, \quad g(2u, 2v) = \frac{P_3S_3 + P_0S_0}{2\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)}. \quad (4.21)$$

4.2 The differential equation (Step I)

By using Eqs.(4.17), (4.18), and (4.21), we obtain the following differential equation

$$(dS_1P_1 - dP_1S_1) = i \frac{P_3S_3 - P_0S_0}{2\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)} (\alpha du + \gamma dv) - i \frac{P_3S_3 + P_0S_0}{2\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)} (\beta du + \delta dv), \quad (4.22)$$

$$(dS_2P_2 - dP_2S_2) = i \frac{P_3S_3 - P_0S_0}{2\Theta \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)} (\alpha du + \gamma dv) + i \frac{P_3S_3 + P_0S_0}{2\Theta \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{bmatrix} (0, 0)} (\beta du + \delta dv). \quad (4.23)$$

²In the Göpel's expression, there is the overall minus sign.

We can verify that Eq.(4.23) is obtained from Eq.(4.22) by replacing $u \rightarrow u + 1, v \rightarrow v + 1$.³ Under such replacement, $(dS_1P_1 - dP_1S_1) \rightarrow -(dS_2P_2 - dP_2S_2)$, $(P_3S_3 + P_0S_0) \rightarrow (P_3S_3 + P_0S_0)$, $(P_3S_3 - P_0S_0) \rightarrow -(P_3S_3 - P_0S_0)$, which provides the proof that Eq.(4.23) is obtained from Eq.(4.22). Thus, we obtain the following differential equation

$$\frac{(dS_1P_1 - dP_1S_1) - (dS_2P_2 - dP_2S_2)}{P_3S_3 + P_0S_0} = \frac{-i(\beta du + \delta dv)}{\Theta\left[\begin{smallmatrix} \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 \end{smallmatrix}\right](0,0)} = d\mu, \quad (4.24)$$

$$\frac{(dS_1P_1 - dP_1S_1) + (dS_2P_2 - dP_2S_2)}{P_3S_3 - P_0S_0} = \frac{i(\alpha du + \gamma dv)}{\Theta\left[\begin{smallmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{smallmatrix}\right](0,0)} = d\nu. \quad (4.25)$$

Which can be expressed in the form

$$\frac{(P_1^2 dp - P_2^2 dq)/P_1P_2}{(P_3S_3 + P_0S_0)/P_1P_2} = \frac{sdp - dq/s}{\varphi} = d\mu, \quad (4.26)$$

$$\frac{(P_1^2 dp + P_2^2 dq)/P_1P_2}{(P_3S_3 - P_0S_0)/P_1P_2} = \frac{sdp + dq/s}{\psi} = d\nu, \quad (4.27)$$

$$s = \frac{P_1}{P_2}, \quad \psi = \frac{P_3S_3 - P_0S_0}{P_1P_2}, \quad \varphi = \frac{P_3S_3 + P_0S_0}{P_1P_2}. \quad (4.28)$$

Therefore, we obtain the starting differential equation⁴

$$\frac{sdp - dq/s}{\varphi} = d\mu, \quad \frac{sdp + dq/s}{\psi} = d\nu, \quad (4.29)$$

$$p = \frac{S_1}{P_1}, \quad q = \frac{S_2}{P_2}, \quad s = \frac{P_1}{P_2} \\ \varphi = \frac{P_3S_3 + P_0S_0}{P_1P_2}, \quad \psi = \frac{P_3S_3 - P_0S_0}{P_1P_2}. \quad (4.30)$$

5 The differential equation with p and q (Step II)

5.1 The elimination of s -dependence in $sdp \mp \frac{dq}{s}$

By dividing $P_1^2P_2^2$ in the Kummer surface relation Eq.(3.25), we obtain

$$(1 - 2Ep^2 + p^4)s^4 + (1 - 2Eq^2 + q^4) - 2\left(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq\right)s^2 = 0, \quad (5.1)$$

which provides

$$(1 - 2Ep^2 + p^4)s^2 + \frac{(1 - 2Eq^2 + q^4)}{s^2} = 2\left(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq\right). \quad (5.2)$$

This is given in the form

$$\left(\sqrt{1 - 2Ep^2 + p^4} s \pm \frac{\sqrt{1 - 2Eq^2 + q^4}}{s}\right)^2 \\ = 2\left(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq\right) \pm 2\sqrt{1 - 2Ep^2 + p^4}\sqrt{1 - 2Eq^2 + q^4}. \quad (5.3)$$

³In the Göpel's expression, in the right-hand side of Eq.(4.23), there is the overall minus sign.

⁴In Göpel's expression, (± 1) sign of $sdp \mp dq/s$ is in the opposite

Introducing the function Δ of the form $\Delta(x) = \sqrt{1 - 2Ex^2 + x^4}$, we can express the above in the form

$$\left(\Delta(p) s \pm \frac{\Delta(q)}{s} \right)^2 = 2G(p, q) \pm 2\Delta(p)\Delta(q), \quad (5.4)$$

$$G(p, q) = \left(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq \right). \quad (5.5)$$

Thus, we obtain the relation

$$\Delta(p) s \pm \frac{\Delta(q)}{s} = \sqrt{2G(p, q) \pm 2\Delta(p)\Delta(q)}. \quad (5.6)$$

Therefore, in the combination of $(\Delta(p)s \pm \Delta(q)/s)$, s -dependence is eliminated. In order to eliminate s -dependence of the differential, we rearrange $(\Delta(p)s \pm \Delta(q)/s)$ in the form

$$\begin{aligned} & s \, dp - \frac{dq}{s} \\ &= \left(s\Delta(p) - \frac{\Delta(q)}{s} \right) \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) + \left(s\Delta(p) + \frac{\Delta(q)}{s} \right) \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) \\ &= \sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) \\ &+ \sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right), \end{aligned} \quad (5.7)$$

$$\begin{aligned} & s \, dp + \frac{dq}{s} \\ &= \left(s\Delta(p) + \frac{\Delta(q)}{s} \right) \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) + \left(s\Delta(p) - \frac{\Delta(q)}{s} \right) \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) \\ &= \sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) \\ &+ \sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right). \end{aligned} \quad (5.8)$$

Thus, we can express $sdp \mp \frac{dq}{s}$ only with p and q .

5.2 The expression of ψ and φ with p and q

a) The expression of $(P_3^2 S_3^2 + P_0^2 S_0^2)/P_1^2 P_2^2$ with p and q

Next, we express $(P_3^2 S_3^2 + P_0^2 S_0^2)/P_1^2 P_2^2$ as the function of p and q . By using the addition formula, we first express $P_3^2, S_3^2, P_0^2, S_0^2, P_1^2, P_2^2$, as the function of T, U, V, W via Eqs.(2.5)-(2.8); in addition, by using Eq.(3.3) and Eq.(3.4), $P_3^3 S_3^2, P_0^3 S_0^2$ and $P_1^3 S_1^2$ are expressed as the function of $P_1^2, S_1^2, P_2^2, S_2^2$. Thus, $(P_3^2 S_3^2 + P_0^2 S_0^2)/P_1^2 P_2^2$ is expressed as the function of p^2, q^2, s^2 in the form

$$\begin{aligned} \frac{1}{2}(\varphi^2 + \psi^2) &= \frac{P_3^2 S_3^2 + P_0^2 S_0^2}{P_1^2 P_2^2} = \frac{(t^2 u^2 - v^2 w^2)(t^2 v^2 - u^2 w^2)}{(t^2 - w^2)^2 (u^2 - v^2)^2} \\ &\times \left(2C \left((1 - 2Ep^2 + p^4)s^2 + \frac{(1 - 2Eq^2 + q^4)}{s^2} \right) - 4E(1 + p^2q^2) + 4(p^2 + q^2) \right). \end{aligned} \quad (5.9)$$

The s -dependence is eliminated by using the Kummer's quartic relation Eq.(5.2) of the form

$$(1 - 2Ep^2 + p^4)s^2 + \frac{(1 - 2Eq^2 + q^4)}{s^2} = 2\left(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq\right) = 2G(p, q).$$

Thus, we obtain the s -independent expression of $(P_3^2S_3^2 + P_0^2S_0^2)/P_1^2P_2^2$ in the form

$$\begin{aligned} \frac{1}{2}(\varphi^2 + \psi^2) &= \frac{P_3^2S_3^2 + P_0^2S_0^2}{P_1^2P_2^2} = \frac{(t^2u^2 - v^2w^2)(t^2v^2 - u^2w^2)}{(t^2 - w^2)^2(u^2 - v^2)^2} \\ &\times \left(4C(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq) - 4E(1 + p^2q^2) + 4(p^2 + q^2)\right) \\ &= b(1 + p^2q^2) - a(p^2 + q^2) + 2cpq, \end{aligned} \quad (5.10)$$

where

$$a = \frac{C^2 - 1}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \quad b = \frac{CF - E}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \quad c = \frac{CD}{\sqrt{(C^2 - 1)(E^2 - 1)}},$$

and we used $\sqrt{(C^2 - 1)(E^2 - 1)} = \frac{(t^2 - w^2)^2(u^2 - v^2)^2}{4(t^2u^2 - v^2w^2)(t^2v^2 - u^2w^2)}$. Thus, we obtain

$$\frac{1}{2}(\varphi^2 + \psi^2) = \frac{P_3^2S_3^2 + P_0^2S_0^2}{P_1^2P_2^2} = b(1 + p^2q^2) - a(p^2 + q^2) + 2cpq. \quad (5.11)$$

b) The expression of $P_3^2S_3^2P_0^2S_0^2/P_1^4P_2^4$ with p and q

Similarly, we calculate

$$\begin{aligned} \varphi\psi &= \frac{P_3^2S_3^2 - P_0^2S_0^2}{P_1^2P_2^2} = \frac{(tu + vw)(tv + uw)}{(t^2 - w^2)(u^2 - v^2)} \\ &\times \left((1 - 2Ep^2 + p^4)s^2 - \frac{(1 - 2Eq^2 + q^4)}{s^2} \right) \\ &= \frac{1}{2\sqrt{E^2 - 1}} \left((1 - 2Ep^2 + p^4)s^2 - \frac{(1 - 2Eq^2 + q^4)}{s^2} \right), \end{aligned} \quad (5.12)$$

where $2\sqrt{E^2 - 1} = \frac{(t^2 - w^2)(u^2 - v^2)}{(tu + vw)(tv + uw)}$ is used. Using Eq.(5.2) and Eq.(5.12), we obtain

$$\left((1 - 2Ep^2 + p^4)s^2 + \frac{(1 - 2Eq^2 + q^4)}{s^2} \right) = 2G(p, q), \quad (5.13)$$

$$\left((1 - 2Ep^2 + p^4)s^2 - \frac{(1 - 2Eq^2 + q^4)}{s^2} \right) = 2\sqrt{E^2 - 1}\varphi\psi, \quad (5.14)$$

which provides

$$s^2 = \frac{G(p, q) + \sqrt{E^2 - 1}\varphi\psi}{1 - 2Ep^2 + p^4}, \quad \frac{1}{s^2} = \frac{G(p, q) - \sqrt{E^2 - 1}\varphi\psi}{1 - 2Eq^2 + q^4}, \quad (5.15)$$

Multiplying the first and the second term, we obtain

$$1 = \frac{G(p, q)^2 - (E^2 - 1)\varphi^2\psi^2}{(1 - 2Ep^2 + p^4)(1 - 2Eq^2 + q^4)}.$$

Thus, we obtain $\varphi^2\psi^2$ expressed with p and q

$$(E^2 - 1)\varphi^2\psi^2 = G(p, q)^2 - (1 - 2Ep^2 + p^4)(1 - 2Eq^2 + q^4) = G(p, q)^2 - \Delta(p)^2\Delta(q)^2.$$

Hence, we obtain $\varphi\psi$ as the function of p and q

$$\varphi\psi = \frac{P_3^2 S_3^2 - P_0^2 S_0^2}{P_1^2 P_2^2} = \sqrt{\frac{G(p, q)^2 - \Delta(p)^2\Delta(q)^2}{E^2 - 1}}. \quad (5.16)$$

Using Eq.(5.11) and Eq.(5.16), we obtain

$$K_1 = \frac{P_3^2 S_3^2 + P_0^2 S_0^2}{P_1^2 P_2^2} = \frac{1}{2}(\varphi^2 + \psi^2) = b(1 + p^2 q^2) - a(p^2 + q^2) + 2cpq, \quad (5.17)$$

$$K_2 = \frac{P_3^2 S_3^2 - P_0^2 S_0^2}{P_1^2 P_2^2} = \varphi\psi = \sqrt{\frac{G(p, q)^2 - \Delta(p)^2\Delta(q)^2}{E^2 - 1}}. \quad (5.18)$$

Thus, we obtain $P_3^2 S_3^2 P_0^2 S_0^2 / P_1^4 P_2^4$ in the form

$$\begin{aligned} \frac{P_3^2 S_3^2 P_0^2 S_0^2}{P_1^4 P_2^4} &= \frac{K_1^2 - K_2^2}{4} \\ &= \frac{1}{4} \left((b(1 + p^2 q^2) - a(p^2 + q^2) + 2cpq)^2 - \frac{G(p, q)^2 - \Delta(p)^2\Delta(q)^2}{E^2 - 1} \right) \\ &= \left(\frac{1}{2}b_1(1 + p^2 q^2) - c_1 pq \right)^2, \\ b_1 &= \frac{D}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \quad c_1 = \frac{CE - F}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \end{aligned} \quad (5.19)$$

where we used $C^2 - D^2 + E^2 + F^2 - 2CEF = 1$. Thus, we obtain

$$\frac{1}{4}(\varphi^2 - \psi^2) = \frac{P_3 S_3 P_0 S_0}{P_1^2 P_2^2} = \pm \left(\frac{1}{2}b_1(1 + p^2 q^2) - c_1 pq \right). \quad (5.20)$$

For the sign ambiguity, we take (+1) sign. Therefore, we obtain φ^2 and ψ^2 as the function of p and q

$$\varphi^2 = \frac{P_3^2 S_3^2 + P_0^2 S_0^2 + 2P_3 S_3 P_0 S_0}{P_1^2 P_2^2} = (b + b_1)(1 + p^2 q^2) - a(p^2 + q^2) + 2(c - c_1)pq, \quad (5.21)$$

$$\psi^2 = \frac{P_3^2 S_3^2 + P_0^2 S_0^2 - 2P_3 S_3 P_0 S_0}{P_1^2 P_2^2} = (b - b_1)(1 + p^2 q^2) - a(p^2 + q^2) + 2(c + c_1)pq. \quad (5.22)$$

Thus, φ and ψ are expressed by the function of p and q in the form

$$\varphi = \sqrt{(b + b_1)(1 + p^2 q^2) - a(p^2 + q^2) + 2(c - c_1)pq}, \quad (5.23)$$

$$\psi = \sqrt{(b - b_1)(1 + p^2 q^2) - a(p^2 + q^2) + 2(c + c_1)pq}. \quad (5.24)$$

5.3 The differential equation with p and q (Step II)

Combining the previous results, we obtain the following differential equation, expressed with only p and q

$$\frac{1}{\varphi} \left\{ \sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) + \sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) \right\} = d\mu, \quad (5.25)$$

$$\frac{1}{\psi} \left\{ \sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) + \sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) \right\} = d\nu, \quad (5.26)$$

$$\varphi = \sqrt{(b + b_1)(1 + p^2q^2) - a(p^2 + q^2) + 2(c - c_1)pq}, \quad (5.27)$$

$$\psi = \sqrt{(b - b_1)(1 + p^2q^2) - a(p^2 + q^2) + 2(c + c_1)pq}, \quad (5.28)$$

$$G(p, q) = F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq, \quad \Delta(x) = \sqrt{1 - 2Ex^2 + x^4}, \quad (5.29)$$

$$b = \frac{CF - E}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \quad a = \frac{C^2 - 1}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \quad c = \frac{CD}{\sqrt{(C^2 - 1)(E^2 - 1)}},$$

$$b_1 = \frac{D}{\sqrt{(C^2 - 1)(E^2 - 1)}}, \quad c_1 = \frac{CE - F}{\sqrt{(C^2 - 1)(E^2 - 1)}}.$$

6 The differential equation with y and z (Step III)

6.1 The change of the functions from p, q to y, z

Next, we change the function from p, q into y, z in such a way as the differential equation becomes separable. Because the combination of $\frac{dp}{\Delta(p)} \pm \frac{dq}{\Delta(q)}$ emerges, we change from p, q to y, z in such a way as y, z satisfy the following differential equation

$$\frac{dp}{\Delta(p)} = \frac{dy}{\Delta(y)} + \frac{dz}{\Delta(z)}, \quad \frac{dq}{\Delta(q)} = \frac{dy}{\Delta(y)} - \frac{dz}{\Delta(z)}. \quad (6.1)$$

This is the differential equation, which provides the addition formula of the elliptic function for the Jacobi type the elliptic curve $y^2 = 1 - 2Ex^2 + x^4 = \Delta(x)^2$. Thus, the addition formula gives

$$p = \frac{y\Delta(z) + z\Delta(y)}{1 - y^2z^2}, \quad q = \frac{y\Delta(z) - z\Delta(y)}{1 - y^2z^2}. \quad (6.2)$$

For more general Jacobi type elliptic curve $y^2 = 1 + \lambda_2x^2 + \lambda_4x^4$, we put $\Delta(x) = \sqrt{1 + \lambda_2x^2 + \lambda_4x^4}$ and the differential equation $\frac{dp}{\Delta(p)} = \frac{dy}{\Delta(y)} + \frac{dz}{\Delta(z)}$ provides the addition formula $p = \frac{y\Delta(z) + y\Delta(z)}{1 - \lambda_4y^2z^2}$.

Using these functions, we obtain

$$\frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) = \frac{dy}{\Delta(y)}, \quad \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) = \frac{dz}{\Delta(z)}. \quad (6.3)$$

6.2 $2G(p, q) \pm \Delta(p)\Delta(q)$ as the function of y and z

Next, we calculate the necessary symmetric function of p and q

$$\begin{aligned} pq &= \frac{y^2\Delta(z)^2 - z^2\Delta(y)^2}{(1 - y^2z^2)^2} = \frac{y^2(1 - 2Ez^2 + z^4) - z^2(1 - 2Ey^2 + y^4)}{(1 - y^2z^2)^2} \\ &= \frac{y^2 - z^2}{1 - y^2z^2}, \end{aligned} \quad (6.4)$$

$$\begin{aligned} p^2 + q^2 &= \frac{2\left(y^2\Delta(z)^2 + z^2\Delta(y)^2\right)}{(1 - y^2z^2)^2} = \frac{2\left(y^2(1 - 2Ez^2 + z^4) + z^2(1 - 2Ey^2 + y^4)\right)}{(1 - y^2z^2)^2} \\ &= \frac{2\left((1 + y^2z^2)(y^2 + z^2) - 4Ey^2z^2\right)}{1 - y^2z^2}, \end{aligned} \quad (6.5)$$

$$\Delta(p)\Delta(q) = \mp \frac{(1 + y^2z^2)^2 - 2E(1 + y^2z^2)(y^2 + z^2) + (y^2 + z^2)^2}{1 - y^2z^2}. \quad (6.6)$$

For the ambiguity of sign in Eq.(6.6), we take (-1) sign.⁵ Substituting the above expressions into

$$2G(p, q) \pm 2\Delta(p)\Delta(q) = 2\left(F(1 + p^2q^2) - C(p^2 + q^2) + 2Dpq\right) \pm 2\Delta(p)\Delta(q),$$

we obtain $2G(p, q) \pm 2\Delta(p)\Delta(q)$ as the function of y and z

$$\begin{aligned} &2G(p, q) \pm 2\Delta(p)\Delta(q) \\ &= \frac{2(F \mp 1)}{(1 - y^2z^2)^2} \left(1 - 2\frac{(C \mp E - D)y^2}{(F \mp 1)} + y^4\right) \left(1 - 2\frac{(C \mp E + D)z^2}{(F \mp 1)} + z^4\right), \end{aligned} \quad (6.7)$$

where we use the identity $C^2 - D^2 + E^2 + F^2 - 2CEF = 1$. It is quite surprising that the y dependence and the z dependence becomes separable in the second and the third term of Eq.(6.7). Thus, we obtain $2G(p, q) \pm 2\Delta(p)\Delta(q)$ as the function of y and z

$$\begin{aligned} &\sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \\ &= \frac{\sqrt{2(F+1)}}{(1 - y^2z^2)} \sqrt{\left(1 - 2\frac{(C+E-D)y^2}{(F+1)} + y^4\right)\left(1 - 2\frac{(C+E+D)z^2}{(F+1)} + z^4\right)} \\ &= \frac{\sqrt{2(F+1)}}{(1 - y^2z^2)} \sqrt{(1 - 2E_2y^2 + y^4)(1 - 2E_3z^2 + z^4)}, \end{aligned} \quad (6.8)$$

$$\begin{aligned} &\sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \\ &= \frac{\sqrt{2(F-1)}}{(1 - y^2z^2)} \sqrt{\left(1 - 2\frac{(C-E-D)y^2}{(F-1)} + y^4\right)\left(1 - 2\frac{(C-E+D)z^2}{(F-1)} + z^4\right)} \\ &= \frac{\sqrt{2(F-1)}}{(1 - y^2z^2)} \sqrt{(1 - 2E_1y^2 + y^4)(1 - 2E_4z^2 + z^4)}, \end{aligned} \quad (6.9)$$

$$E_1 = \frac{(C - E - D)}{(F - 1)}, E_2 = \frac{(C + E - D)}{(F + 1)}, E_3 = \frac{(C + E + D)}{(F + 1)}, E_4 = \frac{(C - E + D)}{(F - 1)},$$

where we used the identity $C^2 - D^2 + E^2 + F^2 - 2CEF = 1$.

⁵Göpel takes $(+1)$ sign. In such case, in connection with the sign of Eq.(4.16), we will see that the differential equation does not become of separable type.

6.3 The expression of φ and ψ with y and z

Next, we calculate φ^2 and ψ^2 as the function of y and z . First, we obtain

$$\begin{aligned}
\varphi^2 &= (b + b_1)(1 + p^2q^2) - a(p^2 + q^2) + 2(c - c_1)pq \\
&= (b + b_1) \left(1 + p^2q^2 - \frac{(C^2 - 1)}{(CF - E + D)}(p^2 + q^2) + \frac{2(CD - CE + F)}{(CF - E + D)}pq \right) \\
&= \frac{b + b_1}{(1 - y^2z^2)^2} \left(1 - \frac{2(C - E - D)}{(F - 1)}y^2 + y^4 \right) \left(1 - \frac{2(C + E + D)}{(F + 1)}z^2 + z^4 \right) \\
&= \frac{b + b_1}{(1 - y^2z^2)^2} (1 - 2E_1y^2 + y^4) (1 - 2E_3z^2 + z^4). \tag{6.10}
\end{aligned}$$

Similarly, we obtain

$$\begin{aligned}
\psi^2 &= (b - b_1)(1 + p^2q^2) - a(p^2 + q^2) + 2(c + c_1)pq \\
&= (b - b_1) \left(1 + p^2q^2 - \frac{(C^2 - 1)}{(CF - E - D)}(p^2 + q^2) + \frac{2(CD + CE - F)}{(CF - E - D)}pq \right) \\
&= \frac{b + b_1}{(1 - y^2z^2)^2} \left(1 - \frac{2(C + E - D)}{(F + 1)}y^2 + y^4 \right) \left(1 - \frac{2(C - E + D)}{(F - 1)}z^2 + z^4 \right) \\
&= \frac{b + b_1}{(1 - y^2z^2)^2} (1 - 2E_2y^2 + y^4) (1 - 2E_4z^2 + z^4), \tag{6.11}
\end{aligned}$$

where we used the identity $C^2 - D^2 + E^2 + F^2 - 2CEF = 1$. Thus, we obtain φ and ψ as the function of y and z

$$\varphi = \frac{\sqrt{b + b_1}}{(1 - y^2z^2)} \sqrt{(1 - 2E_1y^2 + y^4)(1 - 2E_3z^2 + z^4)}, \tag{6.12}$$

$$\psi = \frac{\sqrt{b - b_1}}{(1 - y^2z^2)} \sqrt{(1 - 2E_2y^2 + y^4)(1 - 2E_4z^2 + z^4)}. \tag{6.13}$$

6.4 The differential equation with y and z (Step III)

Using Eqs.(6.3), (6.8), (6.9), (6.12) and (6.13), we can express the necessary quantities with y and z

$$\frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) = \frac{dy}{\Delta(y)}, \quad \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) = \frac{dz}{\Delta(z)}, \tag{6.14}$$

$$\begin{aligned}
&\frac{\sqrt{2G(p, q) - 2\Delta(p)\Delta(q)}}{(1 - y^2z^2)} \\
&= \frac{\sqrt{2(F + 1)}}{(1 - y^2z^2)} \sqrt{(1 - 2E_2y^2 + y^4)(1 - 2E_3z^2 + z^4)}, \tag{6.15}
\end{aligned}$$

$$\begin{aligned}
&\frac{\sqrt{2G(p, q) + 2\Delta(p)\Delta(q)}}{(1 - y^2z^2)} \\
&= \frac{\sqrt{2(F - 1)}}{(1 - y^2z^2)} \sqrt{(1 - 2E_1y^2 + y^4)(1 - 2E_4z^2 + z^4)}, \tag{6.16}
\end{aligned}$$

$$\varphi = \frac{\sqrt{b + b_1}}{(1 - y^2z^2)} \sqrt{(1 - 2E_1y^2 + y^4)(1 - 2E_3z^2 + z^4)}, \tag{6.17}$$

$$\psi = \frac{\sqrt{b - b_1}}{(1 - y^2z^2)} \sqrt{(1 - 2E_2y^2 + y^4)(1 - 2E_4z^2 + z^4)}. \tag{6.18}$$

Substituting these expressions into the following differential equations,

$$\frac{1}{\varphi} \left\{ \sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) + \sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) \right\} = d\mu, \quad (6.19)$$

$$\frac{1}{\psi} \left\{ \sqrt{2G(p, q) + 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} + \frac{dq}{\Delta(q)} \right) + \sqrt{2G(p, q) - 2\Delta(p)\Delta(q)} \frac{1}{2} \left(\frac{dp}{\Delta(p)} - \frac{dq}{\Delta(q)} \right) \right\} = d\nu, \quad (6.20)$$

we obtain the separable differential equation of y and z in the form

$$\begin{aligned} d\mu &= \frac{1 - y^2 z^2}{\sqrt{b + b_1}} \frac{1}{\sqrt{(1 - 2E_1 y^2 + y^4)(1 - 2E_3 z^2 + z^4)}} \\ &\times \left(\frac{\sqrt{2(F + 1)}}{1 - y^2 z^2} \sqrt{(1 - 2E_2 y^2 + y^4)(1 - 2E_3 z^2 + z^4)} \frac{dy}{\Delta(y)} \right. \\ &\left. + \frac{\sqrt{2(F - 1)}}{1 - y^2 z^2} \sqrt{(1 - 2E_1 y^2 + y^4)(1 - 2E_4 z^2 + z^4)} \frac{dz}{\Delta(z)} \right) \\ &= \frac{1}{\sqrt{b + b_1}} \left(\sqrt{\frac{2(F + 1)(1 - 2E_2 y^2 + y^4)}{(1 - 2E y^2 + y^4)(1 - 2E_1 y^2 + y^4)}} dy \right) \\ &\left. + \sqrt{\frac{2(F - 1)(1 - 2E_4 z^2 + z^4)}{(1 - 2E z^2 + z^4)(1 - 2E_3 z^2 + z^4)}} dz \right). \end{aligned} \quad (6.21)$$

Similarly, we obtain

$$\begin{aligned} d\nu &= \frac{(1 - y^2 z^2)}{\sqrt{b - b_1}} \frac{1}{\sqrt{(1 - 2E_2 y^2 + y^4)(1 - 2E_4 z^2 + z^4)}} \\ &\times \left(\frac{\sqrt{2(F - 1)}}{(1 - y^2 z^2)} \sqrt{(1 - 2E_1 y^2 + y^4)(1 - 2E_4 z^2 + z^4)} \frac{dy}{\Delta(y)} \right. \\ &\left. + \frac{\sqrt{2(F + 1)}}{(1 - y^2 z^2)} \sqrt{(1 - 2E_2 y^2 + y^4)(1 - 2E_3 z^2 + z^4)} \frac{dz}{\Delta(z)} \right) \\ &= \frac{1}{\sqrt{b - b_1}} \left(\sqrt{\frac{2(F - 1)(1 - 2E_1 y^2 + y^4)}{(1 - 2E y^2 + y^4)(1 - 2E_2 y^2 + y^4)}} dy \right) \\ &\left. + \sqrt{\frac{2(F + 1)(1 - 2E_3 z^2 + z^4)}{(1 - 2E z^2 + z^4)(1 - 2E_4 z^2 + z^4)}} dz \right). \end{aligned} \quad (6.22)$$

It is quite surprising that, if we use the functions y and z , the differential equations become of the separable type.

7 The differential equation with y and y' (Step IV)

Next, we change the function in order that it provide the same type of Abelian differential.

7.1 The change of the function from z to y'

The differential equation in the previous section provides

$$d\mu = \frac{1}{\sqrt{b+b_1}} \left(\frac{\sqrt{2(F+1)}(1-2E_2y^2+y^4)}{\sqrt{(1-2Ey^2+y^4)(1-2E_1y^2+y^4)(1-2E_2y^2+y^4)}} dy \right. \\ \left. + \frac{\sqrt{2(F-1)}(1-2E_4z^2+z^4)}{\sqrt{(1-2Ez^2+z^4)(1-2E_3z^2+z^4)(1-2E_4z^2+z^4)}} dz \right), \quad (7.1)$$

$$d\nu = \frac{1}{\sqrt{b-b_1}} \left(\frac{\sqrt{2(F-1)}(1-2E_1y^2+y^4)}{\sqrt{(1-2Ey^2+y^4)(1-2E_1y^2+y^4)(1-2E_2y^2+y^4)}} dy \right. \\ \left. + \frac{\sqrt{2(F+1)}(1-2E_3z^2+z^4)}{\sqrt{(1-2Ez^2+z^4)(1-2E_3z^2+z^4)(1-2E_4z^2+z^4)}} dz \right). \quad (7.2)$$

For the function y , we obtain the Abelian differential of the type

$$\frac{dy}{\sqrt{(1-2Ey^2+y^4)(1-2E_1y^2+y^4)(1-2E_2y^2+y^4)}}.$$

While, for the function z , we obtain the Abelian differential of the type

$$\frac{dz}{\sqrt{(1-2Ez^2+z^4)(1-2E_3z^2+z^4)(1-2E_4z^2+z^4)}}.$$

However, the type of the Abelian differential is different.

We keep y to be the same function, yet we change the function z into y' , which cause the Möbius transformation of z^2 , and make the Abel function of the same type. We parametrize $2E$ with α in the form $2E = \alpha^2 + 1/\alpha^2$. For the constant e , we obtain $\Delta(e) = 1 - 2Ee^2 + e^4 = (1 - e^2/\alpha^2)(1 - e^2/\beta^2)$, $\alpha\beta = 1$. We change the function from y, z to y, y' (y is not changed) by the following differential equation

$$\frac{dz}{\Delta(z)} = \frac{dy'}{\Delta(y')} + \frac{de}{\Delta(e)} = \frac{dy'}{\Delta(y')}, \quad (7.3)$$

where we use $de = 0$ because e is the constant. This provides

$$z = \frac{y'\Delta(e) + e\Delta(y')}{1 - e^2y'^2}. \quad (7.4)$$

Next, we choose $e = \alpha$, thus we obtain $\Delta(e) = \Delta(\alpha) = 0$, hence z is given as the function of y'

$$z = \frac{\alpha\Delta(y')}{1 - \alpha^2y'^2}. \quad (7.5)$$

Considering the square of z , we obtain

$$z^2 = \frac{\alpha^2 - y'^2}{1 - \alpha^2y'^2}. \quad (7.6)$$

Thus, y'^2 is the Möbius transformation of z^2 . Hence, we obtain

$$\Delta(z)^2 = \left(1 - \frac{z^2}{\alpha^2}\right)\left(1 - \frac{z^2}{\alpha^2}\right) = \left(\frac{(1 - \alpha^2)y'}{\alpha(1 - \alpha^2 y'^2)}\right)^2, \quad (7.7)$$

which provides $\Delta(z)$ as the function of y in the following form

$$\Delta(z) = \frac{(1 - \alpha^2)y'}{\alpha(1 - \alpha^2 y'^2)}. \quad (7.8)$$

Using Eq.(7.5) and Eq.(7.8) , we obtain the necessary quantities as the function of y and y'

$$p = \frac{y\Delta(z) + z\Delta(y)}{1 - y^2 z^2} = \frac{\beta(\alpha^4 - 1)yy' + \alpha\Delta(y)\Delta(y')}{1 - \alpha^2(y^2 + y'^2) + y^2 y'^2}, \quad (7.9)$$

$$q = \frac{y\Delta(z) - z\Delta(y)}{1 - y^2 z^2} = \frac{\beta(\alpha^4 - 1)yy' - \alpha\Delta(y)\Delta(y')}{1 - \alpha^2(y^2 + y'^2) + y^2 y'^2}, \quad (7.10)$$

$$pq = \frac{y^2 - z^2}{1 - y^2 z^2} = \frac{y^2 + y'^2 - \alpha^2(1 + y^2 y'^2)}{1 - \alpha^2(y^2 + y'^2) + y^2 y'^2}. \quad (7.11)$$

We will use these relations in the next section.

7.2 The differential equation with y and y' (Step IV)

We calculate the following quantity as the function of y'

$$\frac{1}{2}\left(z^2 + \frac{1}{z^2}\right) = \frac{1}{2}\left(\frac{\alpha^2 - y'^2}{1 - \alpha^2 y'^2} + \frac{1 - \alpha^2 y'^2}{\alpha^2 - y'^2}\right) = \frac{E - 2y'^2 + Ey'^4}{1 - 2Ey'^2 + y'^4}.$$

Thus, we obtain

$$\frac{1 - 2E_3 z^2 + z^4}{2z^2} = \frac{1}{2}\left(z^2 + \frac{1}{z^2}\right) - E_3 = \frac{(E - E_3) - 2(1 - EE_3)y'^2 + (E - E_3)y'^4}{1 - 2Ey'^2 + y'^4}. \quad (7.12)$$

Similarly, we obtain

$$\frac{1 - 2E_4 z^2 + z^4}{2z^2} = \frac{1}{2}\left(z^2 + \frac{1}{z^2}\right) - E_4 = \frac{(E - E_4) - 2(1 - EE_4)y'^2 + (E - E_4)y'^4}{1 - 2Ey'^2 + y'^4}. \quad (7.13)$$

By using the identity

$$1 - E(E_1 + E_3) + E_1 E_3 = 0, \quad 1 - E(E_2 + E_4) + E_2 E_4 = 0, \quad \frac{E - E_4}{E - E_3} = \frac{F + 1}{F - 1},$$

we obtain in the form

$$\begin{aligned} \frac{1 - 2E_3 z^2 + z^4}{2z^2} &= \frac{(E - E_3) - 2(1 - EE_3)y'^2 + (E - E_3)y'^4}{1 - 2Ey'^2 + y'^4} \\ &= \frac{(E - E_3)(1 - 2E_1 y'^2 + y'^4)}{1 - 2Ey'^2 + y'^4}, \end{aligned} \quad (7.14)$$

$$\begin{aligned} \frac{1 - 2E_4 z^2 + z^4}{2z^2} &= \frac{(E - E_4) - 2(1 - EE_4)y'^2 + (E - E_4)y'^4}{1 - 2Ey'^2 + y'^4} \\ &= \frac{(E - E_4)(1 - 2E_2 y'^2 + y'^4)}{1 - 2Ey'^2 + y'^4}. \end{aligned} \quad (7.15)$$

Taking the ratio of Eq.(7.14) and Eq.(7.15), we obtain the expression

$$\frac{1 - 2E_3z^2 + z^4}{1 - 2E_4z^2 + z^4} = \left(\frac{F-1}{F+1}\right) \left(\frac{1 - 2E_1y'^2 + y'^4}{1 - 2E_2y'^2 + y'^4}\right), \quad (7.16)$$

$$\frac{1 - 2E_4z^2 + z^4}{1 - 2E_3z^2 + z^4} = \left(\frac{F+1}{F-1}\right) \left(\frac{1 - 2E_2y'^2 + y'^4}{1 - 2E_1y'^2 + y'^4}\right). \quad (7.17)$$

Therefore, we obtain the differential equation of the same type Abelian differential

$$\begin{aligned} d\mu &= \frac{1}{\sqrt{b+b_1}} \left(\sqrt{\frac{2(F+1)(1-2E_2y^2+y^4)}{(1-2Ey^2+y^4)(1-2E_1y^2+y^4)}} dy + \sqrt{\frac{2(F-1)(1-2E_4z^2+z^4)}{(1-2Ez^2+z^4)(1-2E_3z^2+z^4)}} dz \right) \\ &= \sqrt{\frac{2(F+1)}{b+b_1}} \left(\sqrt{\frac{(1-2E_2y^2+y^4)}{(1-2Ey^2+y^4)(1-2E_1y^2+y^4)}} dy \right. \\ &\quad \left. + \sqrt{\frac{(1-2E_2y'^2+y'^4)}{(1-2Ey'^2+y'^4)(1-2E_1y'^2+y'^4)}} dy' \right). \end{aligned} \quad (7.18)$$

Similarly, we obtain

$$\begin{aligned} d\nu &= \frac{1}{\sqrt{b-b_1}} \left(\sqrt{\frac{2(F-1)(1-2E_1y^2+y^4)}{(1-2Ey^2+y^4)(1-2E_2y^2+y^4)}} dy + \sqrt{\frac{2(F+1)(1-2E_3z^2+z^4)}{(1-2Ez^2+z^4)(1-2E_4z^2+z^4)}} dz \right) \\ &= \sqrt{\frac{2(F-1)}{b-b_1}} \left(\sqrt{\frac{(1-2E_1y^2+y^4)}{(1-2Ey^2+y^4)(1-2E_2y^2+y^4)}} dy \right. \\ &\quad \left. + \sqrt{\frac{(1-2E_1y'^2+y'^4)}{(1-2Ey'^2+y'^4)(1-2E_2y'^2+y'^4)}} dy' \right). \end{aligned} \quad (7.19)$$

Thus, for dy and dy' , we have the same type Abelian differential of the form

$$\frac{dx}{\sqrt{(1-2Ex^2+x^4)(1-2E_1x^2+x^4)(1-2E_2x^2+x^4)}}.$$

8 The differential equation with x and x' (Step V)

8.1 The differential equation of the genus two Jacobi's inversion problem

By changing the functions y, y' into x, x' in such a way as dx and dx' becomes the same Abelian differential of the fifth-degree hyperelliptic curve. For that purpose, we change the function in the form

$$x = \left(\frac{1-y^2}{1+y^2}\right)^2, \quad x' = \left(\frac{1-y'^2}{1+y'^2}\right)^2. \quad (8.1)$$

We denote in the following

$$m = \frac{E+1}{E-1}, \quad m_1 = \frac{E_1+1}{E_1-1}, \quad m_2 = \frac{E_2+1}{E_2-1}. \quad (8.2)$$

Thus, we obtain the simple expression

$$\frac{1 - m_2x}{1 - m_1x} = \left(\frac{E_1 - 1}{E_2 - 1} \right) \left(\frac{1 - 2E_2y^2 + y^4}{1 - 2E_1y^2 + y^4} \right), \quad \frac{1 - m_1x}{1 - m_2x} = \left(\frac{E_2 - 1}{E_1 - 1} \right) \left(\frac{1 - 2E_1y^2 + y^4}{1 - 2E_2y^2 + y^4} \right). \quad (8.3)$$

Therefore, we obtain the simple expression

$$\sqrt{\frac{1 - 2E_1y^2 + y^4}{1 - 2E_2y^2 + y^4}} = \sqrt{\frac{E_1 - 1}{E_2 - 1}} \sqrt{\frac{1 - m_1x}{1 - m_2x}}, \quad \sqrt{\frac{1 - 2E_1y'^2 + y'^4}{1 - 2E_2y'^2 + y'^4}} = \sqrt{\frac{E_1 - 1}{E_2 - 1}} \sqrt{\frac{1 - m_1x'}{1 - m_2x'}}, \quad (8.4)$$

and

$$\sqrt{\frac{1 - 2E_2y^2 + y^4}{1 - 2E_1y^2 + y^4}} = \sqrt{\frac{E_2 - 1}{E_1 - 1}} \sqrt{\frac{1 - m_2x}{1 - m_1x}}, \quad \sqrt{\frac{1 - 2E_2y'^2 + y'^4}{1 - 2E_1y'^2 + y'^4}} = \sqrt{\frac{E_2 - 1}{E_1 - 1}} \sqrt{\frac{1 - m_2x'}{1 - m_1x'}}. \quad (8.5)$$

By differentiating y , we obtain

$$\begin{aligned} dx &= -\frac{8((1 - y^2)ydy)}{(1 + y^2)^3}, \quad \sqrt{x} = \frac{1 - y^2}{1 + y^2}, \quad \sqrt{1 - x} = \frac{2y}{1 + y^2}, \\ \sqrt{1 - mx} &= \sqrt{\frac{-2}{E - 1}} \frac{\sqrt{1 - 2Ey^2 + y^4}}{1 + y^2}. \end{aligned} \quad (8.6)$$

Combining these relations, we obtain the connection of the differential of dx and dy

$$\frac{dx}{\sqrt{x(1 - x)(1 - mx)}} = \frac{2\sqrt{2}\sqrt{-1}\sqrt{E - 1} dy}{\sqrt{1 - 2Ey^2 + y^4}}. \quad (8.7)$$

Similarly, we obtain

$$\frac{dx'}{\sqrt{x'(1 - x')(1 - mx')}} = \frac{2\sqrt{2}\sqrt{-1}\sqrt{E - 1} dy'}{\sqrt{1 - 2Ey'^2 + y'^4}}. \quad (8.8)$$

Using Eqs.(7.18), (7.19), (8.7) and (8.8), we finally obtain the simplified differential equation

$$\begin{aligned} &\sqrt{\frac{2(F + 1)}{b + b_1}} \left(\sqrt{\frac{(1 - 2E_2y^2 + y^4)}{(1 - 2Ey^2 + y^4)(1 - 2E_1y^2 + y^4)}} dy + \sqrt{\frac{(1 - 2E_2y'^2 + y'^4)}{(1 - 2Ey'^2 + y'^4)(1 - 2E_1y'^2 + y'^4)}} dy' \right) \\ &= \frac{1}{2\sqrt{b + b_1}} \sqrt{\frac{(F + 1)(1 - E_2)}{(1 - E)(1 - E_1)}} \left(\sqrt{\frac{(1 - m_2x)}{x(1 - x)(1 - mx)}} dx + \sqrt{\frac{(1 - m_2x')}{x'(1 - x')(1 - mx')}} dx' \right) = d\mu, \end{aligned} \quad (8.9)$$

$$\begin{aligned} &\sqrt{\frac{2(F - 1)}{b - b_1}} \left(\sqrt{\frac{(1 - 2E_1y^2 + y^4)}{(1 - 2Ey^2 + y^4)(1 - 2E_2y^2 + y^4)}} dy + \sqrt{\frac{(1 - 2E_1y'^2 + y'^4)}{(1 - 2Ey'^2 + y'^4)(1 - 2E_2y'^2 + y'^4)}} dy' \right) \\ &= \frac{1}{2\sqrt{b + b_1}} \sqrt{\frac{(F + 1)(1 - E_2)}{(1 - E)(1 - E_1)}} \left(\sqrt{\frac{(1 - m_2x)}{x(1 - x)(1 - mx)}} dx + \sqrt{\frac{(1 - m_2x')}{x'(1 - x')(1 - mx')}} dx' \right) = d\nu. \end{aligned} \quad (8.10)$$

Therefore, we obtain the differential equation of the genus two Jacobi's inversion problem

$$\frac{(1 - m_2x)dx}{\sqrt{f_5(x)}} + \frac{(1 - m_2x')dx'}{\sqrt{f_5(x')}} = 2\sqrt{b + b_1} \sqrt{\frac{(1 - E)(1 - E_1)}{(F + 1)(1 - E_2)}} d\mu = d\hat{\mu}, \quad (8.11)$$

$$\frac{(1 - m_1x)dx}{\sqrt{f_5(x)}} + \frac{(1 - m_1x')dx'}{\sqrt{f_5(x')}} = 2\sqrt{b - b_1} \sqrt{\frac{(1 - E)(1 - E_2)}{(F - 1)(1 - E_1)}} d\nu = d\hat{\nu}, \quad (8.12)$$

$$f(x) = x(1 - x)(1 - mx)(1 - m_1x)(1 - m_2x). \quad (8.13)$$

8.2 The expression of x and x' as the function of p and q

Next, we express x, x' with p, q . From Eqs.(7.9)-(7.11), we obtain the necessary quantities as the function of y and y'

$$p + q = \frac{2\beta(\alpha^4 - 1)yy'}{1 - \alpha^2(y^2 + y'^2) + y^2y'^2}, \quad pq = \frac{y^2 + y'^2 - \alpha^2(1 + y^2y'^2)}{1 - \alpha^2(y^2 + y'^2) + y^2y'^2},$$

which provides

$$1 + \alpha^2pq = \frac{(1 - \alpha^4)(1 + y^2y'^2)}{1 + y^2y'^2 - \alpha^2(y^2 + y'^2)}, \quad 1 + \beta^2pq = \frac{(\beta^2 - \alpha^2)(y^2 + y'^2)}{1 + y^2y'^2 - \alpha^2(y^2 + y'^2)}. \quad (8.14)$$

Thus, we obtain

$$\frac{p + q}{1 + \alpha^2pq} = -\frac{2\beta yy'}{(1 + y^2y'^2)}, \quad \frac{p + q}{1 + \beta^2pq} = -\frac{2\alpha yy'}{(y^2 + y'^2)}. \quad (8.15)$$

Hence, we can, in principle, express $y^2 + y'^2$ and yy' as the function of p and q , which implies that $x = (1 - y^2)^2/(1 + y^2)^2$ and $x' = (1 - y'^2)^2/(1 + y'^2)^2$ can be expressed as the function of p and q . For our purpose, we make the following combination

$$f = \left(\frac{1 - yy'}{1 + yy'}\right)^2, \quad g = \left(\frac{y - y'}{y + y'}\right)^2, \quad (8.16)$$

which provides the following nice factorization property

$$f = \frac{(1 + \alpha p)(1 + \alpha q)}{(1 - \alpha p)(1 - \alpha q)}, \quad g = \frac{(1 + \beta p)(1 + \beta q)}{(1 - \beta p)(1 - \beta q)}. \quad (8.17)$$

Later, we use $f + g, 1 + fg, \sqrt{fg}$. Thus, we calculate these quantities

$$\begin{aligned} f + g &= \frac{2(p^2q^2 + (\alpha - \beta)^2pq - (p^2 + q^2) + 1)}{(1 - \alpha p)(1 - \alpha q)(1 - \beta p)(1 - \beta q)} \\ &= \frac{2(p^2q^2 + (2E - 2)pq - (p^2 + q^2) + 1)}{(1 - \alpha p)(1 - \alpha q)(1 - \beta p)(1 - \beta q)}, \end{aligned} \quad (8.18)$$

$$\begin{aligned} 1 + fg &= \frac{2(p^2q^2 + (\alpha + \beta)^2pq + (p^2 + q^2) + 1)}{(1 - \alpha p)(1 - \alpha q)(1 - \beta p)(1 - \beta q)} \\ &= \frac{2(p^2q^2 + (2E + 2)pq + (p^2 + q^2) + 1)}{(1 - \alpha p)(1 - \alpha q)(1 - \beta p)(1 - \beta q)}, \end{aligned} \quad (8.19)$$

$$\sqrt{fg} = \pm \frac{\Delta(p)\Delta(q)}{(1 - \alpha p)(1 - \alpha q)(1 - \beta p)(1 - \beta q)}, \quad (8.20)$$

where we used $\alpha^2 + \beta^2 = 2E, \alpha\beta = 1$. We take (+1) sign in Eq.(8.20).⁶ Using Eq.(8.16), we obtain

$$\left(\frac{1 - yy'}{1 + yy'}\right) = \sqrt{f}, \quad \left(\frac{y - y'}{y + y'}\right) = \sqrt{g}, \quad (8.21)$$

⁶Göpel take (-1) sign in Eq.(8.20).

which provides yy' , y/y' as the function of \sqrt{f} , \sqrt{g}

$$yy' = \frac{1 - \sqrt{f}}{1 + \sqrt{f}}, \quad \frac{y}{y'} = \frac{1 + \sqrt{g}}{1 - \sqrt{g}}. \quad (8.22)$$

Multiplying and dividing the first and the second term of Eq.(8.22), we obtain

$$y^2 = \frac{(1 - \sqrt{f})(1 + \sqrt{g})}{(1 + \sqrt{f})(1 - \sqrt{g})}, \quad y'^2 = \frac{(1 - \sqrt{f})(1 - \sqrt{g})}{(1 + \sqrt{f})(1 + \sqrt{g})}. \quad (8.23)$$

Thus, we obtain the desired form

$$\frac{1 - y^2}{1 + y^2} = \frac{\sqrt{f} - \sqrt{g}}{1 - \sqrt{fg}}, \quad \frac{1 - y'^2}{1 + y'^2} = \frac{\sqrt{f} + \sqrt{g}}{1 + \sqrt{fg}}. \quad (8.24)$$

Therefore, x and x' are expressed as the function of p and q

$$x = \left(\frac{1 - y^2}{1 + y^2} \right)^2 = \frac{f + g - 2\sqrt{fg}}{1 + fg - 2\sqrt{fg}} = \frac{E(1 + p^2q^2) - (p^2 + q^2) - \Delta(p)\Delta(q)}{(E + 1)(pq + 1)^2}, \quad (8.25)$$

$$x' = \left(\frac{1 - y'^2}{1 + y'^2} \right)^2 = \frac{f + g + 2\sqrt{fg}}{1 + fg + 2\sqrt{fg}} = \frac{E(1 + p^2q^2) - (p^2 + q^2) + \Delta(p)\Delta(q)}{(E + 1)(pq + 1)^2}. \quad (8.26)$$

8.3 The solution of the genus two Jacobi's inversion problem

From the differential equation

$$\frac{(1 - m_2x)dx}{\sqrt{f_5(x)}} + \frac{(1 - m_2x')dx'}{\sqrt{f_5(x')}} = d\hat{\mu}, \quad \frac{(1 - m_1x)dx}{\sqrt{f_5(x)}} + \frac{(1 - m_1x')dx'}{\sqrt{f_5(x')}} = d\hat{\nu},$$

we can rearrange in the standard Jacobi's inversion problem of the form

$$\frac{dx}{\sqrt{f_5(x)}} + \frac{dx'}{\sqrt{f_5(x')}} = dU_1, \quad \frac{xdx}{\sqrt{f_5(x)}} + \frac{x'dx'}{\sqrt{f_5(x')}} = dU_2, \quad (8.27)$$

where we obtain the expression $U_1 = k_1u_1 + k_2u_2$, $U_2 = k_3u_1 + k_4u_2$. We inversely express $u_1 = \ell_1U_1 + \ell_2U_2$, $u_2 = \ell_3U_1 + \ell_4U_2$. Thus, the solution of the Jacobi's inversion problem is given by

$$\begin{aligned} \wp_{22}(U_1, U_2) &= x + x' = \frac{2(E(1 + p^2q^2) - (p^2 + q^2))}{(E + 1)(pq + 1)^2} \\ &= \frac{2E}{(E + 1)} - \frac{2(p^2 + q^2 - 2Epq)}{(E + 1)(1 + pq)^2}, \end{aligned} \quad (8.28)$$

$$\wp_{12}(U_1, U_2) = -xx' = -\frac{(E - 1)}{(E + 1)} + \frac{4(E - 1)pq}{(E + 1)(1 + pq)^2}, \quad (8.29)$$

where

$$p = \frac{S_1(u_1, u_2)}{P_1(u_1, u_2)} = \frac{\vartheta \left[\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix} \right] (\ell_1U_1 + \ell_2U_2, \ell_3U_1 + \ell_4U_2)}{\vartheta \left[\begin{smallmatrix} 0 & 0 \\ 0 & 1 \end{smallmatrix} \right] (\ell_1U_1 + \ell_2U_2, \ell_3U_1 + \ell_4U_2)}, \quad (8.30)$$

$$q = \frac{S_2(u_1, u_2)}{P_2(u_1, u_2)} = \frac{\vartheta \left[\begin{smallmatrix} 1 & 1 \\ 1 & 0 \end{smallmatrix} \right] (\ell_1U_1 + \ell_2U_2, \ell_3U_1 + \ell_4U_2)}{\vartheta \left[\begin{smallmatrix} 0 & 0 \\ 1 & 0 \end{smallmatrix} \right] (\ell_1U_1 + \ell_2U_2, \ell_3U_1 + \ell_4U_2)}. \quad (8.31)$$

For the explicit functional form for $\wp_{22}(U_1, U_2), \wp_{12}(U_1, U_2)$, we must carefully determine $\ell_i, (i = 1, 2, 3, 4)$ by the $u_1 = 0, u_2 = 0$ value of various theta functions and derivative of the various theta functions.

9 Summary and Discussions

In the previous paper, we reviewed the Rosenhain's paper to the Jacobi's inversion problem for the genus two hyperelliptic integral. In this paper, we have reviewed the Göpel's paper to the Jacobi's inversion problem for the genus two hyperelliptic integral.

In the Rosenhain's approach, the Riemann's addition formula of the hyperelliptic theta function is used. The key identity of the Rosenhain's approach is the three quadratic theta identities. In the Göpel's approach, the addition formula by the duplication method is used. The key identity of the Göpel's approach is the three quartic theta identities, Eq.(3.25), Eq.(5.11), Eq.(5.20). One of these identities, Eq.(3.25), is the quartic Kummer surface relation. Three quartic identities are given in the form

$$\begin{aligned} P_1^4 + S_1^4 + P_2^4 + S_2^4 - 2F(P_1^2 P_2^2 + S_1^2 S_2^2) + 2C(P_1^2 S_2^2 + P_2^2 S_1^2) \\ - 2E(P_1^2 S_1^2 + P_2^2 S_2^2) - 4DP_1 S_1 P_2 S_2 = 0, \\ P_3^2 S_3^2 + P_0^2 S_0^2 - b(P_1^2 P_2^2 + S_1^2 S_2^2) + a(P_1^2 S_2^2 + P_2^2 S_1^2) - 2cP_1 S_1 P_2 S_2 = 0, \\ 2P_3 S_3 P_0 S_0 - b_1(P_1^2 P_2^2 + S_1^2 S_2^2) + 2c_1 P_1 S_1 P_2 S_2 = 0. \end{aligned}$$

Starting from the genus two hyperelliptic theta functions, Göpel takes several steps to obtain the differential equation of the genus two Jacobi's inversion equation. In step I, by using the addition formula of the genus two hyperelliptic theta functions, the derivative formula is obtained; thus the starting differential equation is obtained. In step II, by using three quartic theta identities, the differential equation of only p and q is obtained. In step III, by using the addition formula of the elliptic function of the form $\frac{dp}{\Delta(p)} = \frac{dy}{\Delta(y)} + \frac{dz}{\Delta(z)}$ and $\frac{dq}{\Delta(q)} = \frac{dy}{\Delta(y)} - \frac{dz}{\Delta(z)}$, the separable type differential equation of y and z is obtained. In step IV, by using the addition formula of the elliptic function of the form $\frac{dz}{\Delta(z)} = \frac{dy'}{\Delta(y')} + \frac{de}{\Delta(e)} = \frac{dy'}{\Delta(y')}$ with some special constant e , the differential equation of y and y' with the same type Abelian differential is obtained. In step V, via $x = \left(\frac{1-y^2}{1+y^2}\right)^2, x' = \left(\frac{1-y'^2}{1+y'^2}\right)^2$, the differential equation of x and x' with the Abelian differential of the fifth-degree hyperelliptic curve is obtained, i.e., the differential equation of the genus two Jacobi's inversion problem is obtained.

In the paper on the comments of the Göpel's paper, Jacobi pointed out the issue of the number of modules [16]. For the genus g hyperelliptic curve $y^2 = \lambda_{2g+1}x^{2g+1} + \lambda_{2g}x^{2g} + \dots + \lambda_1x + \lambda_0$, we have the $2g + 2$ constant coefficients. However, we have the freedom to change, i) constant scale of x , ii) constant shift of x , iii) take the ratio of the coefficients, in order to provide the standard form. For example, in the $g = 1$ case, i) provides $y^2 = 4x^3 + \lambda_2x^2 + \lambda_1x + \lambda_0$. While, ii) provide $y^2 = 4x^3 - g_2x - g_3$. In the step iii), $g_2^3/(g_2^3 - 27g_3^2)$ provides $\tau = \omega_3/\omega_1$ from ω_3 and ω_1 by taking the ratio of the constant coefficients. Thus, the number of the modules from the hyperelliptic curve is $2g + 2 - 3 = 2g - 1$. While, in the g -variable hyperelliptic theta function, in general, there are modules of the form $\tau_{ij}(= \tau_{ji}), (i, j = 1, 2, \dots, g)$; thus, the number of the modules is $g(g + 1)/2$. Hence, for

$g \geq 3$, the number of the module for the hyperelliptic function $g(g+1)/2$ becomes greater than that of the hyperelliptic curve $2g-1$.

References

- [1] C.G.J. Jacobi, “ De Functionibus Duarum Variabilium Quadrupliciter Periodicis, Quibus Theoria Transcendentium Abelianarum Innititur”, J. reine angew. Math., **13** (1835), 55-78.
- [2] C.G.J. Jacobi, “ Note sur les Fonctions Abéliennes”, J. reine angew. Math., **30** (1846), 183-184.
- [3] A. Göpel, “Theoriae transcendentium Abelianarum primi ordinis adumbratio levis”, J. reine angew. Math., **35** (1847), 277-312.
- [4] A. Göpel edited by H. Weber and A. Witting, “Entwurf einer Theorie der Abel’schen Transcendenten earster Ordnung”, (Leipzig, W. Engelmann,1895).
- [5] G. Rosenhain edited by H. Weber and A. Witting “Abhandlung über die Functionen zweiter Variabler mit vier Perioden, welche die Inversen sind der ultra-elliptischen Integrale erster klasse”, (Leipzig, W. Engelmann, 1895).
- [6] G. Rosenhain, “ Auszug mehrerer Schreiben des Dr. Rosenhain an Herrn Professor Jacobi über die hyperelliptischen Transcendenten”, J. reine angew. Math., **40** (1850), 319-360.
- [7] K. Shigemoto, ”Jacobi’s Inversion Problem for Genus Two Hyperelliptic Integral”, Tezukayama Academic Review, **20** (2014), 1-45, [[arXiv:1603.02508v2\[math-ph\]](#)].
- [8] C.G.J. Jacobi, “ Suite des notices sur les fonctions elliptiques”, J. reine angew. Math., **3** (1828), 303-310.
- [9] K. Shigemoto, “Full Addition Formulae of Genus Two Hyperelliptic Functions by the Duplication Method”, [[arXiv:2010.15505v1\[math.CA\]](#)] (2020).
- [10] C.G.J. Jacobi, “Theorie der elliptischen Functionen, aus den Eigenschaften der Thetareihen abgeleitet”(1838), hrsg. von Adolf Kneser, Ostwald’s Klassiker der exakten Wissenschaften Nr. 224, (Leipzig, Akademische Verlagsgesellschaft, 1927).
- [11] F. Prym, ”Untersuchungen über die Riemann’sche Thetaformel und Riemann’sche Charakteristikentheorie”, (Drug und Verlag von B.O. Teubner, Leipzig, 1882).
- [12] B. Riemann, “Ueber das Verschwinden der ϑ -Functionen”, J. reine angew. Math., **65** (1866), 161-172.
- [13] E. Kossak, ”Das Additionstheorem Der Ultra-Elliptischen Functionen Erster Ordnung”, (Berlin, Nicolai’sche Verlagsbuchhandlung, 1871).

- [14] K. Shigemoto, “Integrability Condition in the Statistical Model and the Addition Formula of $g = 2$ Hyperelliptic Function”, Tezukayama Academic Review **21**, 1-25 (2015), [arXiv:1603.02131v2[math-ph]].
- [15] R. W. H. T. Hudson, *Kummer’s Quartic Surface*, (Cambridge University Press, Cambridge,1990), p.81.
- [16] C.G.J. Jacobi, “Notiz über A. Göpel”, J. reine angew. Math., **35** (1847), 313-317.