

**PROOF OF VOGAN'S CONJECTURE ON ARTHUR PACKETS: IRREDUCIBLE
PARAMETERS OF p -ADIC GENERAL LINEAR GROUPS**

CLIFTON CUNNINGHAM AND MISHTY RAY

ABSTRACT. In this paper we prove Vogan's conjecture on local Arthur packets, as recalled in [CFM⁺22, Section 8.3, Conjecture 1(a)], for irreducible Arthur parameters of p -adic general linear groups. This result shows that these Arthur packets may be characterized by properties of simple perverse sheaves on a moduli space of Langlands parameters.

1. INTRODUCTION

The local Langlands correspondence for a connected reductive algebraic group G over a p -adic field F partitions the set of equivalence classes of smooth irreducible representations of $G(F)$ into L -packets using equivalence classes of Langlands parameters. Following [Vog93, ABV92], the correspondence may be viewed as a bijection between smooth irreducible representations that share a common infinitesimal parameter of a p -adic group $G(F)$, along with its pure inner forms, and simple objects in the category of equivariant perverse sheaves on a moduli space of Langlands parameters. This perspective leads to the notion of an ABV-packet, as articulated in [CFM⁺22, Section 8], which conjecturally generalizes the local Arthur packet - we call this as Vogan's conjecture on local Arthur packets. In this paper, we prove this conjecture for irreducible Arthur parameters of the p -adic group $GL_n(F)$.

1.1. Simple and irreducible parameters. Let F be a non-archimedean local field with residue characteristic p and residue field \mathbb{F}_q , so $q = p^f$. Let W_F be the Weil group of F and W'_F denote the group $W_F \times \mathrm{SL}_2(\mathbb{C})$. Recall that a Langlands parameter is a homomorphism

$$\phi : W'_F \rightarrow {}^L G = \widehat{G} \rtimes W_F$$

commuting with the natural projections to W_F , satisfying conditions adapted from [Bor79]. In particular, this means that all the information of a Langlands parameter ϕ is captured by a 1-cocycle

$$\phi^0 : W'_F \rightarrow \widehat{G}$$

such that $\phi(w, g) = \phi^0(g, w) \rtimes w$. In the case $G = \mathrm{GL}_n$, the cocycle ϕ^0 is a representation of the group W'_F .

Now set $W''_F := W_F \times \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C})$ and recall that an Arthur parameter [Art13, Chapter 1] is a homomorphism

$$\psi : W''_F \rightarrow {}^L G = \widehat{G} \rtimes W_F,$$

likewise defined by a 1-cocycle

$$\psi^0 : W''_F \rightarrow \widehat{G}$$

satisfying certain conditions, notably that the restriction of ψ^0 to W_F is bounded. Again, when $G = \mathrm{GL}_n$, ψ^0 is a representation. We say that ψ is a *simple* Arthur parameter of $\mathrm{GL}_n(F)$ if ψ^0 is

an irreducible representation and the restriction of ψ^0 to W_F is trivial. We say that ψ is *irreducible* if ψ^0 is an irreducible representation of W_F'' .

Every Arthur parameter determines a Langlands parameter by the rule

$$\phi_\psi(w, x) = \psi \left(w, x, \begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{-\frac{1}{2}} \end{pmatrix} \right).$$

The infinitesimal parameter λ_ϕ of a Langlands parameter ϕ is a homomorphism

$$\lambda_\phi : W_F \rightarrow {}^L G$$

defined by the rule

$$\lambda_\phi(w) = \phi \left(w, \begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{-\frac{1}{2}} \end{pmatrix} \right).$$

See [CFM⁺22, Section 4.1] for definition. As before, it is determined by

$$\lambda_\phi^0 : W_F \rightarrow \widehat{G}$$

which is a representation when $G = \mathrm{GL}_n$. The function $\phi \mapsto \lambda_\phi$ defined on \widehat{G} -equivalence classes of these parameters is not injective in general. The infinitesimal parameters λ_{ϕ_ψ} for a simple Arthur parameter ψ is unramified as a representation of W_F , *i.e.*, it are trivial on the inertia group I_F .

1.2. Local Langlands correspondence. The local Langlands conjecture for a connected reductive algebraic group G over F gives a map between the set $\Pi(G(F))$ of equivalence classes of smooth irreducible representations of $G(F)$ and the set $\Phi({}^L G)$ of equivalence classes of Langlands parameters. This map can be refined [Vog93, ABV92] as a bijection between $\Pi(G(F))$ and the set $\Xi({}^L G)$ of equivalence classes of pairs $\xi = (\phi, \epsilon)$, called enhanced Langlands parameters, where ϵ is an irreducible representation of the finite group

$$S_\phi := Z_{\widehat{G}}(\phi) / Z_{\widehat{G}}(\phi)^0 Z(\widehat{G})^{\Gamma_F},$$

where Γ_F is the absolute Galois group of F . We emphasize the distinction between Langlands parameters and L -parameters - Langlands parameters are maps and L -parameters are \widehat{G} -equivalence classes of Langlands parameters. These L -parameters partition $\Pi(G(F))$ into disjoint finite sets $\Pi_\phi(G(F))$ called L -packets. This correspondence must satisfy a list of functoriality properties and should be compatible with L -functions and ϵ -factors on both sides (See [Bor79] or [Kal16], for example). When $G = \mathrm{GL}_n$, the local Langlands correspondence is a theorem [Hen00, HT01]. In this case, $S_\phi = 1$ (see [Art13], for example) and the correspondence can be viewed as a bijection

$$\Pi(G(F)) \rightarrow \Phi(G),$$

where $\Phi(G)$ is the set of L -parameters for G . Thus, L -packets are singletons for GL_n .

1.3. A-packets. In 1989, Arthur introduced [Art89] what are now known as Arthur packets, or A -packets, which enjoy some nice properties that non-tempered L -packets fail to satisfy. This was a global theory and the local analogue was clarified by Arthur's subsequent work [Art13] in 2013. Local A -packets $\Pi_\psi(G(F))$ are finite sets of smooth irreducible representations of $G(F)$ which are not necessarily disjoint and consist of unitary admissible representations. The representations that appear in A -packets are said to be of Arthur type. Every A -packet $\Pi_\psi(G(F))$ contains a distinguished L -packet $\Pi_{\phi_\psi}(G(F))$ through the injection $\psi \mapsto \phi_\psi$. In the 1990s, Adams, Barbasch, and Vogan suggested [ABV92] a geometric characterization of A -packets by using microlocal analysis on

certain stratified complex varieties built out of Langlands parameters. When $G = \mathrm{GL}_n$, A -packets are singletons [Art13]; in particular, the A -packet for ψ coincides with the L -packet for ϕ_ψ .

1.4. Vogan-Langlands correspondence. Vogan's perspective [Vog93] on the local Langlands correspondence suggests a beautiful geometric approach to A -packets. This perspective concerns a moduli space X_λ of Langlands parameters that have the same infinitesimal parameter λ . Vogan observed that simple objects in the category $\mathrm{Per}_{\widehat{G}}(X_\lambda)$ of equivariant perverse sheaves on X_λ are naturally identified with the pairs (ϕ, ϵ) , when we properly interpret the finite group attached to ϕ and consider representations of $G(F)$ together with its pure inner forms. Viewed in this way, the local Langlands correspondence takes the form of a bijection

$$\Pi_\lambda(G/F) \rightarrow \mathrm{Per}_{H_\lambda}(V_\lambda)_{/iso}^{\mathrm{simple}}$$

between the set $\Pi_\lambda(G/F)$ of equivalence classes of irreducible objects in the category $\mathrm{Rep}_\lambda(G/F)$ of smooth representations of $G(F)$ and its pure rational forms which have a matching infinitesimal parameter λ and the set $\mathrm{Per}_{H_\lambda}(V_\lambda)_{/iso}^{\mathrm{simple}}$ of isomorphism classes of simple objects in the category $\mathrm{Per}_{\widehat{G}}(X_\lambda)$.

Instead of working with $\mathrm{Per}_{\widehat{G}}(X_\lambda)$, we work with an equivalent category $\mathrm{Per}_{H_\lambda}(V_\lambda)$ (see [CFM⁺22, Section 4.5]). The variety V_λ of Langlands parameters and the group H_λ are described in 2. In the case $G = \mathrm{GL}_n$, the correspondence can be viewed as a bijection

$$\Pi_\lambda(G(F)) \rightarrow \mathrm{Per}_{H_\lambda}(V_\lambda)_{/iso}^{\mathrm{simple}},$$

where $\Pi_\lambda(G(F))$ is the set of equivalence classes of smooth irreducible representations of $G(F)$ with a matching infinitesimal parameter λ and

$$\mathrm{Per}_{H_\lambda}(V_\lambda)_{/iso}^{\mathrm{simple}} = \{\mathcal{IC}(\mathbb{1}_C) : C \subseteq V_\lambda \text{ is an } H_\lambda\text{-orbit}\},$$

where $\mathbb{1}_C$ denotes the constant local system on C . In this way, every $\pi \in \Pi_\lambda(G(F))$ corresponds to $\mathcal{P}(\pi) = \mathcal{IC}(\mathbb{1}_C) \in \mathrm{Per}_{H_\lambda}(V_\lambda)_{/iso}^{\mathrm{simple}}$, for a unique H_λ -orbit $C \subseteq V_\lambda$. In other words, in this case there is a bijection between elements of $\Pi_\lambda(G(F))$ and H_λ -orbits C in V_λ .

1.5. Vogan's conjecture on A-packets. Following an approach similar to [ABV92], Vogan attached to any Langlands parameter ϕ the set of irreducible representations of $G(F)$ and its pure inner forms for which the characteristic cycles of the corresponding simple perverse sheaf contains the conormal bundle of the \widehat{G} -orbit of ϕ in the moduli space mentioned above. This notion is revised in [CFM⁺22] in terms of vanishing cycles. This set is referred to as the ABV-packet $\Pi_\phi^{\mathrm{ABV}}(G/F)$.

Vogan's perspective on the local Langlands correspondence introduces pure A-packets $\Pi_\psi(G/F)$ which are unions of A -packets for G and its pure inner forms. This perspective also identifies $\Pi_\psi(G/F)$ with a finite set of simple perverse sheaves on V_{λ_ψ} , where λ_ψ is the infinitesimal parameter of ϕ_ψ . What property do these perverse sheaves share, and how are they determined by the Arthur parameter ψ ? This question is answered by the functor

$$\mathrm{Evs}_\psi : \mathrm{Per}_{H_{\lambda_\psi}}(V_{\lambda_\psi}) \rightarrow \mathrm{Rep}(A_\psi)$$

where $A_\psi = Z_{\widehat{G}}(\psi)/Z_{\widehat{G}}(\psi)^0$, constructed in [CFM⁺22, Section 7.10]. In fact, this functor is defined in terms of a more general functor $\mathrm{Ev}_{C_\psi} : \mathrm{D}_{H_{\lambda_\psi}}(V_{\lambda_\psi}) \rightarrow \mathrm{D}_{H_{\lambda_\psi}}(\Lambda_{C_\psi}^{\mathrm{reg}})$ where $\Lambda_{C_\psi} \subset \Lambda_{\lambda_\psi}$ is the conormal bundle above C_ψ in the conormal variety for V_λ [CFM⁺22, Section 7.3]. This Ev_{C_ψ} functor is defined in such a way that

$$(\mathrm{Ev}_{C_\psi} \mathcal{P})_{(x,\xi)} = (\mathrm{R}\Phi_\xi \mathcal{P})_x,$$

for all $(x, \xi) \in \Lambda_{C_\psi}^{\text{reg}}$, where $R\Phi_\xi$ is Deligne's vanishing cycles functor for $\xi : V_{\lambda_\psi} \rightarrow \mathbb{C}$ [CFM⁺22, Equation 7.6]. When restricted from $D_{H_{\lambda_\psi}}(V_{\lambda_\psi})$ to $\text{Per}_{H_{\lambda_\psi}}(V_{\lambda_\psi})$, the functor Ev_{C_ψ} lands in $\text{Per}_{H_{\lambda_\psi}}(\Lambda_{C_\psi}^{\text{reg}})$, which is equivalent to the semisimple category $\text{Loc}_{H_{\lambda_\psi}}(\Lambda_{C_\psi}^{\text{reg}})$ of equivariant local systems on $\Lambda_{C_\psi}^{\text{reg}}$, by [CFM⁺22, Proposition 6.9]. This proposition also gives an equivalence

$$\text{Loc}_{H_{\lambda_\psi}}(\Lambda_{C_\psi}^{\text{reg}}) \cong \text{Rep}(A_\psi);$$

see [CFM⁺22, Equation 7.30]. As explained in [CFM⁺22, Equations 7.22, 7.24], the precise relation between Evs_ψ and Ev_{C_ψ} is given by

$$(1) \quad \text{Evs}_\psi \mathcal{P} = \mathcal{H}_{(x_\psi, \xi_\psi)}^{1+\text{codim } C_\psi} (\text{Ev}_{C_\psi} \mathcal{P}),$$

as a vector space with A_ψ -action, where $(x_\psi, \xi_\psi) \in \Lambda_{\lambda_\psi}$ is defined in [CFM⁺22, Equation 6.12, 6.13]. For $G = \text{GL}_n$, $A_\psi = 1$ by [Art13, Chapter 1], for example; so in this case, $\text{Rep}(A_\psi)$ is simply the category of vector spaces. Following [CFM⁺22], the ABV-packet for ψ is defined by

$$(2) \quad \Pi_{\phi_\psi}^{\text{ABV}}(G(F)) := \{\pi \in \Pi_{\lambda_\psi}(G(F)) : \text{Evs}_\psi(\mathcal{P}(\pi)) \neq 0\},$$

where $\mathcal{P}(\pi)$ is the simple perverse sheaf on V_{λ_ψ} corresponding to $\pi \in \Pi_{\lambda_\psi}(G(F))$.

The main result of this paper, below, gives a proof of [CFM⁺22, Section 8.3, Conjecture 1(a)] adapted to general linear groups, for irreducible Arthur parameters:

Theorem 1 (Theorem 6.4). *Let ψ be an irreducible Arthur parameter of $G = \text{GL}_n(F)$, let ϕ_ψ be its corresponding Langlands parameter, let $\Pi_\psi(G(F))$ be the Arthur packet attached to ψ . Then*

$$\Pi_\psi(G(F)) = \Pi_{\phi_\psi}^{\text{ABV}}(G(F)).$$

We briefly overview the structure of the paper. Henceforth, $G = \text{GL}_n$. In Section 2, we write down the *Vogan variety* V_λ , the group H_λ , and the explicit action of H_λ on V_λ . Our perspective on the local Langlands correspondence is that of representations corresponding to orbits in the Vogan variety. Both sides of this correspondence are encoded in combinatorial objects - multisegments for representations and rank triangles for orbits. We reiterate these notions and write down an algorithm to go from multisegments to rank triangles and vice-versa in Section 3. In Section 4, we discuss the Zelevinsky involution on multisegments and explain how it corresponds to an involution on orbits. We write down the Mœglin-Waldspurger algorithm to compute the involution on a multisegment, and use this notion to prove Lemma 4.8. In section 5, we adapt Lemma 4.8 to a statement about orbits in Lemma 5.1. Using this lemma and the properties of the functor Evs_ψ , we prove the conjecture for simple A -parameters in Theorem 5.3. In section 6, we use the process of hyper-unramification to generalize the result to an irreducible A -parameter.

In upcoming work, we expect to generalize Theorem 1 to arbitrary A -parameters for $\text{GL}_n(F)$ for a non-archimedean field F .

Since ABV-packets are expected to generalize Arthur packets, it is natural to ask - are all ABV-packets are singletons for GL_n ? The answer is no; one can see an ABV-packet of size 2 for a non Arthur-type Langlands parameter of GL_{16} in [CFK22].

Acknowledgements. The authors would like to acknowledge Andrew Fiori for sharing his insights on the combinatorics of multisegments, which was crucial for Lemma 4.8. The second named author would like to extend her gratitude to Kristaps Balodis and James Steele for helpful discussions.

2. A MODULI SPACE OF LANGLANDS PARAMETERS

Recall that F is a non-archimedean local field of residue characteristic $q = p^f$ with p -adic absolute value denoted by $|\cdot|$ and the uniformizer denoted by ϖ . The Weil group is $W_F = I_F \rtimes \mathbb{Z}$, where I_F is the inertia subgroup and \mathbb{Z} is identified with the cyclic subgroup generated by the Frobenius. From local class field theory, we have the local Artin map $\alpha : W_F \rightarrow F^\times$ that maps I_F to the group of units \mathfrak{o}^\times in F^\times and the geometric Frobenius \mathfrak{f} to $1/\varpi$. We abuse notation and use $|\cdot|$ to denote the absolute value on W_F , where $|w| := |\alpha(w)|$. Thus, $|\mathfrak{f}| = |1/\varpi| = q$. We have that $G = \mathrm{GL}_n(F)$, and we can work with $\widehat{G} = \mathrm{GL}_n(\mathbb{C})$ instead of ${}^L G$.

We consider *simple* Arthur parameters, *i.e.*, of the form

$$(3) \quad \begin{aligned} \psi : W_F'' &\longrightarrow \mathrm{GL}_n(\mathbb{C}) \\ (w, x, y) &\mapsto \mathrm{Sym}^d(x) \otimes \mathrm{Sym}^a(y), \end{aligned}$$

where a and d are positive integers¹. For $w \in W_F$, let d_w denote the matrix $\begin{pmatrix} |w|^{\frac{1}{2}} & 0 \\ 0 & |w|^{-\frac{1}{2}} \end{pmatrix}$. The corresponding Langlands parameter is given by

$$(4) \quad \begin{aligned} \phi_\psi : W_F' &\longrightarrow \mathrm{GL}_n(\mathbb{C}) \\ (w, x) &\mapsto \psi(w, x, d_w). \end{aligned}$$

One can simplify $\psi(w, x, d_w)$ to get

$$\phi_\psi(w, x) = |w|^{\frac{a}{2}} \mathrm{Sym}^d(x) \oplus |w|^{\frac{a-2}{2}} \mathrm{Sym}^d(x) \cdots \oplus |w|^{\frac{-a}{2}} \mathrm{Sym}^d(x).$$

The infinitesimal parameter associated to this Langlands parameter is given by

$$(5) \quad \begin{aligned} \lambda_{\phi_\psi} : W_F &\longrightarrow \mathrm{GL}_n(\mathbb{C}) \\ (w, x) &\mapsto \phi_\psi(w, d_w). \end{aligned}$$

One can simplify $\phi_\psi(w, d_w)$ to get

$$(6) \quad \lambda_{\phi_\psi}(w) = \bigoplus_{i=0}^a \bigoplus_{j=0}^d |w|^{\frac{a+d}{2} - (i+j)}$$

We fix this infinitesimal parameter for the rest of this work and call it λ for simplicity. The element $\lambda(\mathfrak{f}) \in \mathrm{GL}_n(\mathbb{C})$ is semisimple with eigenvalues $\lambda_i := q^{e_i}$ occurring with multiplicities m_i , where $0 \leq i \leq k$, with $e_i = e_{i-1} - 1$. Let us rewrite the matrix $\lambda(\mathfrak{f})$ in a more instructive way:

$$(7) \quad \lambda(\mathfrak{f}) = \begin{bmatrix} q^{e_0} I_{m_0} & 0 & \cdots \\ 0 & q^{e_1} I_{m_1} & \cdots \\ \vdots & \ddots & \\ 0 & & q^{e_k} I_{m_k} \end{bmatrix}.$$

We denote the corresponding eigenspaces by E_i . If we consider the n -dimensional vector space which is the standard representation of $\mathrm{Lie}(\widehat{G})$ spanned by the usual basis $\{f_1, \dots, f_n\}$, then E_0 is spanned by $\{f_1, \dots, f_{m_0}\}$, E_1 by $\{f_{m_0+1}, \dots, f_{m_1}\}$, and in general E_i by $\{f_{m_{i-1}+1}, \dots, f_{m_i}\}$.

¹Here we use d and a in our notation to refer to the fact that the first SL_2 in $W_F'' = W_F \times \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C})$ is the ‘‘Deligne- SL_2 ’’, *i.e.*, it comes from Weil-Deligne representations, and the second SL_2 is the ‘‘Arthur- SL_2 ’’, *i.e.*, it comes from the definition of Arthur parameters.

Following [Vog93] and [CFM⁺22], we describe the variety V_λ and group H_λ below:

$$(8) \quad V_\lambda := \{X \in \mathfrak{gl}_n(\mathbb{C}) : \text{Ad}(\lambda(w))X = |w|X, \forall w \in W_F\}.$$

As the absolute value on I_F is trivial, any $X \in V_\lambda$ is determined by the equation

$$(9) \quad \text{Ad}(\lambda(\mathfrak{f}))X = qX.$$

V_λ is referred to as the *Vogan variety*. In this case, it is easy to see the structure of this variety. Any matrix in V_λ is strictly upper triangular or trace zero so it can be identified as a vector subspace of the \mathfrak{sl}_n . $\lambda(\mathfrak{f})$ is an element inside the maximal torus in SL_n . Using the Cartan decomposition of \mathfrak{sl}_n , V_λ can be identified as a sum of root spaces of \mathfrak{sl}_n such that the roots evaluate to q at $\lambda(\mathfrak{f})$. This shows that V_λ is simply an affine space. This fact is true more generally, as seen in [CFM⁺22, Lemma 5.5].

Lemma 2.1. The Vogan variety V_λ can be decomposed as

$$(10) \quad V_\lambda = \text{Hom}(E_k, E_{k-1}) \times \text{Hom}(E_{k-1}, E_{k-2}) \times \cdots \times \text{Hom}(E_1, E_0).$$

Proof. Let $X \in V_\lambda$ and $v \in E_i$. By definition, $\lambda(\mathfrak{f})v = q^{e_i}v$ and $\lambda(\mathfrak{f})X\lambda(\mathfrak{f})^{-1} = qX$. We show that $Xv \in E_{i-1}$. Indeed

$$\begin{aligned} \lambda(\mathfrak{f})Xv &= \lambda(\mathfrak{f})X\lambda(\mathfrak{f})^{-1} \cdot \lambda(\mathfrak{f})v, \\ &= qX \cdot q^{e_i}v, \\ &= q^{e_i+1}Xv. \end{aligned}$$

As E_{i-1} is the eigenspace for q^{e_i+1} ,

$$V_\lambda \subseteq \text{Hom}(E_k, E_{k-1}) \times \text{Hom}(E_{k-1}, E_{k-2}) \times \cdots \times \text{Hom}(E_1, E_0).$$

Now we show the reverse containment. Let $T \in \text{Hom}(E_i, E_{i-1})$. If we view this as an endomorphism of \mathbb{C}^n , then the $m_{i-1} \times m_i$ matrix associated with T can be thought of as an $n \times n$ matrix (a_{ij}) such that $a_{ij} \neq 0$ if $m_{i-2} + 1 \leq i \leq m_{i-1}$ and $m_{i-1} + 1 \leq j \leq m_i$, and $a_{ij} = 0$ otherwise. However, the adjoint (conjugation) action of $\lambda(\mathfrak{f})$ simply multiplies all the nonzero entries a_{ij} by q . Thus, $\text{Ad}(\lambda(\mathfrak{f})) \cdot (a_{ij}) = q \cdot (a_{ij})$ and the matrix (a_{ij}) can be identified with an element of V_λ . This shows the reverse containment. \square

The group H_λ is $Z_{\widehat{G}}(\lambda)$, the centralizer of the image of λ in \widehat{G} . Again, any $g \in H_\lambda$ is determined by the equation

$$(11) \quad \text{Ad}(\lambda(\mathfrak{f})) \cdot g = g.$$

Using the description of $\lambda(\mathfrak{f})$ in (7), we can directly see that

$$Z_{\widehat{G}}(\lambda(\mathfrak{f})) = \begin{bmatrix} \text{GL}_{m_0}(\mathbb{C}) & 0 & \cdots \\ 0 & \text{GL}_{m_1}(\mathbb{C}) & \cdots \\ \vdots & \ddots & \\ 0 & & \text{GL}_{m_k}(\mathbb{C}) \end{bmatrix}$$

We may identify H_λ as follows.

$$(12) \quad H_\lambda = \text{Aut}(E_k) \times \text{Aut}(E_{k-1}) \times \cdots \times \text{Aut}(E_0)$$

H_λ acts on V_λ via conjugation. Using the interpretation from (10) and (12), we describe the action as follows. For elements in V_λ let us use the notation

$$x = (x_{kk-1}, \cdots, x_{21}, x_{10}), \text{ where } x_{ii-1} \in \text{Mat}_{m_{i-1}, m_i}(\mathbb{C}).$$

For elements of H_λ , we write

$$h = (h_k, \dots, h_1, h_0), \text{ where } h_i \in \text{GL}_{m_i}(\mathbb{C}).$$

In this notation, the action H_λ on V_λ is given by

$$(13) \quad (h_k, \dots, h_1, h_0) \cdot (x_{kk-1}, \dots, x_{21}, x_{10}) = (h_{k-1}x_{kk-1}h_k^{-1}, \dots, h_1x_{21}h_2^{-1}, h_0x_{10}h_1^{-1}).$$

Observe that $\text{rank}(h_{i-1}x_{ii-1}h_i^{-1}) = \text{rank}(x_{ii-1})$, thus any orbit preserves the ranks of x_{ii-1} s. More generally every orbit is determined by the ranks of x_{ii-1} s and all their permitted products. Realizing elements of V_λ as a matrix as in (10), we have

$$x = \begin{bmatrix} 0 & x_{10} & 0 & \dots \\ 0 & 0 & x_{21} & \dots \\ \vdots & & \ddots & \\ 0 & 0 & \dots & x_{kk-1} \\ & & & 0 \end{bmatrix}.$$

The orbit of any x is determined by its Jordan canonical form. Notice that 0 is the only eigenvalue, so calculating dimensions of the generalized eigenspaces will involve computing ranks of powers of this matrix x , which in turn involves computing ranks of all permitted products

$$(14) \quad x_{ij} := x_{ii-1}x_{i-1i-2} \cdots x_{j+1j}$$

where where $0 \leq j < i - 1 \leq k$. Writing down these rank equations exactly realizes an orbit as an algebraic variety.

With emphasis on the fact that λ determines H_λ and V_λ , we drop the subscript and use H and V instead.

Example 2.2. For GL_4 , set $\psi(w, x, y) = \text{Sym}^1(x) \otimes \text{Sym}^1(y)$. Then $\phi_\psi = \psi(w, x, d_w) = |w|^{1/2}x \oplus |w|^{-1/2}x$ and $\lambda(w) = \psi(w, d_w, d_w)$, so

$$\lambda(\mathbf{f}) = \begin{bmatrix} q & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & q^{-1} \end{bmatrix}.$$

By direct calculation using (9) and (11)

$$V = \left\{ \begin{bmatrix} 0 & u_1 & u_2 & 0 \\ 0 & 0 & 0 & v_1 \\ 0 & 0 & 0 & v_2 \\ 0 & 0 & 0 & 0 \end{bmatrix} : u_1, u_2, v_1, v_2 \in \mathbb{C} \right\} \simeq \mathbb{A}_{\mathbb{C}}^4,$$

$$H = \left\{ \begin{bmatrix} t_1 & 0 & 0 & 0 \\ 0 & t_2 & r & 0 \\ 0 & s & t_3 & 0 \\ 0 & 0 & 0 & t_4 \end{bmatrix} : t_1, t_4 \in \mathbb{C}^\times, \begin{bmatrix} t_2 & r \\ s & t_3 \end{bmatrix} \in \text{GL}_2(\mathbb{C}) \right\} \simeq \text{GL}_1(\mathbb{C}) \times \text{GL}_2(\mathbb{C}) \times \text{GL}_1(\mathbb{C}).$$

The eigenvalues of $\lambda(\mathbf{f})$ are $q, 1, q^{-1}$ with multiplicities 1, 2, and 1 and eigenspaces $E_0 = \text{span}\{(1, 0, 0, 0)\}$, $E_1 = \text{span}\{(0, 1, 0, 0), (0, 0, 1, 0)\}$ and $E_2 = \text{span}\{(0, 0, 0, 1)\}$, respectively.

In the formulation of (10) and (12), we may write

$$\begin{aligned} V &= \mathrm{Hom}(E_2, E_1) \times \mathrm{Hom}(E_1, E_0), \\ H &= \mathrm{Aut}(E_2) \times \mathrm{Aut}(E_1) \times \mathrm{Aut}(E_0). \end{aligned}$$

Following (13), the action of H on V is given by

$$\left(t_4, \begin{bmatrix} t_2 & r \\ s & t_3 \end{bmatrix}, t_1 \right) \cdot \left(\begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, [u_1, u_2] \right) = \left(\begin{bmatrix} t_2 & r \\ s & t_3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} t_4^{-1}, t_1 [u_1, u_2] \begin{bmatrix} t_2 & r \\ s & t_3 \end{bmatrix}^{-1} \right).$$

3. FROM MULTISEGEMENTS TO RANK TRIANGLES AND VICE-VERSA

As outlined in Section 1.4, there is a bijection between equivalence classes of smooth irreducible representations that share the infinitesimal parameter λ and H -orbits in V . We know that smooth irreducible representations of G can be parameterized using multisegments [Zel80]. The relationship between multisegments and orbits appears in [Zel81]. In this section, we introduce the combinatorial gadget of rank triangles to parametrize the orbits and write down an algorithm compute the multisegment corresponding to a rank triangle and vice-versa. This has been worked out for some specific cases in GL_{16} in [CFK22]; here we present it with updated notation for GL_n .

3.1. Multisegments. The Langlands classification using multisegments is summarized beautifully in [Kud94, Section 1]. We recall this classification in more detail following the notation in [Kud94]. For any representation π of $\mathrm{GL}_m(F)$, let $\pi(i) := |\det(\cdot)|^i \pi$. For a partition $n = \underbrace{m + m + \dots + m}_{r\text{-times}}$

and a supercuspidal representation σ of $\mathrm{GL}_m(F)$, we call

$$(15) \quad (\sigma, \sigma(1), \dots, \sigma(r-1)) = [\sigma, \sigma(r-1)] = \Delta$$

a segment. This segment determines a representation on a parabolic subgroup of G whose levi is $\underbrace{\mathrm{GL}_m(F) \times \mathrm{GL}_m(F) \times \dots \times \mathrm{GL}_m(F)}_{r\text{-times}}$. More precisely, we have the representation $\sigma \otimes \sigma(1) \otimes \dots \otimes$

$\sigma(r-1)$ on the levi, inflated to a representation of the parabolic, which is trivial on the unipotent. We can then carry out parabolic induction to obtain the induced representation $I_P^G(\Delta)$ of G which has a unique irreducible quotient denoted by $Q(\Delta)$.

A multisegment is a collection of segments with repetitions allowed. The Langlands classification theorem tells us that any smooth irreducible representation of G occurs as a unique irreducible quotient inside a parabolically induced representation determined by a multisegment $\{\Delta_1, \Delta_2, \dots, \Delta_r\}$ subject to certain conditions - we denote that quotient by $Q(\Delta_1, \Delta_2, \dots, \Delta_r)$. Next, for integers $i < j$ we introduce the notation

$$(16) \quad [i, j] := (|\cdot|^i, |\cdot|^{i+1}, \dots, |\cdot|^j)$$

for a segment which is the special case of (15) when we consider the partition $1 + 1 + \dots + 1$ and σ to be the character $|\cdot|^i$ of F^\times . This notation may be extended to half integers $i < j$ as long as $j - i + 1$ is a positive integer (this is the length of the segment). A segment of length 1 of the form $\{|\cdot|^i\}$ is just denoted $[i]$. There is a partial order on multisegments which we recall below.

Definition 3.1. Let α and β be any two multisegments. We say that $\alpha \leq \beta$ if we can form β by performing elementary operations on segments in α . More precisely, $\alpha \leq \beta$ if we can form β by

replacing any two segments Δ_1 and Δ_2 in α with

$$\begin{cases} \Delta_1 \cup \Delta_2 \text{ and } \Delta_1 \cap \Delta_2 & \text{if } \Delta_1 \cap \Delta_2 \neq \emptyset, \\ \Delta_1 \cup \Delta_2 & \text{if } \Delta_1 \cup \Delta_2 = \emptyset \text{ and } \Delta_1 \cup \Delta_2 \text{ is a segment,} \\ \Delta_1 \text{ and } \Delta_2 & \text{otherwise.} \end{cases}$$

Example 3.2. In the case of GL_4 , the Steinberg representation can be written as $Q([-\frac{3}{2}, \frac{3}{2}])$ and the trivial representation is given by $Q([\frac{3}{2}], [\frac{1}{2}], [-\frac{1}{2}], [-\frac{3}{2}])$. The multisegment for ϕ_ψ from Example 2.2 is $\{[0,1], [-1,0]\}$.

3.2. Rank triangles. We know from Section 2 that any orbit of $(x_{kk-1}, \dots, x_{21}, x_{10}) \in V$ is determined by the ranks of x_{ij} s as defined in (14). We arrange these ranks in a rank triangle. This arrangement helps us write down a visual algorithm to compute the correspondence between multisegments and rank triangles in Section 3.3. To that end we set

$$r_{ij} := \text{rank}(x_{ij}),$$

where $0 \leq j < i \leq k$. Recall that E_i is an eigenspace for $\lambda(\mathfrak{f})$ with eigenvalue q^{e_i} with multiplicity m_i . We arrange these exponents, multiplicities, and ranks into a triangle to reflect the corresponding combinations of the x_{ij} and refer to this as a *rank triangle*:

$$\begin{array}{cccccc} e_k & e_{k-1} & \cdots & e_1 & e_0 & \\ \hline m_k & m_{k-1} & \cdots & m_1 & m_0 & \\ \hline & r_{kk-1} & r_{(k-1)(k-2)} \cdots & r_{21} & r_{10} & \\ & & r_{k(k-2)} & \cdots & r_{20} & \\ & & & \cdots & \cdots & \\ & & & & r_{k0} & \end{array}$$

From left to right, the values in the top row of the rank triangle correspond, respectively, to the exponents e_i of the eigenvalues q^{e_i} . The values in the second row correspond respectively to the multiplicities m_i of the eigenvalues q^{e_i} . The row below these multiplicities shows ranks r_{ij} of x_{ij} , which are subject to the condition that every rank is less than or equal to the two eigenvalue multiplicities above it. For the other rows, the ranks are subject to exactly two conditions:

- Every rank is less than or equal to the two ranks above it. This is a consequence of a basic fact from linear algebra: if A and B are matrices such that AB and BC are defined then $\text{rank}(AB) \leq \text{rank}(A)$ and $\text{rank}(AB) \leq \text{rank}(B)$.
- The ranks satisfy

$$r_{is} + r_{lj} \leq r_{ij} + r_{ls}, \quad j \leq s < l \leq i.$$

This condition is a consequence of the Frobenius inequality from linear algebra: if A, B and C are matrices such that AB, BC and ABC are defined then $\text{rank}(AB) + \text{rank}(BC) \leq \text{rank}(ABC) + \text{rank}(B)$.

The set of H -orbits in V is naturally in bijection with rank triangles subject to these conditions. The set of H -orbits in V carries a partial order defined by the Zariski topology:

Definition 3.3. There is a partial order on the set of orbits in V_λ : For any two orbits C and C' , $C \leq C'$ if and only if $C \subseteq \overline{C'}$.

This partial order can be read easily from the corresponding rank triangles as follows: If the ranks for C are r_{ij} and C' are r'_{ij} then $C \leq C'$ if and only if $r_{ij} \leq r'_{ij}$ for all i, j .

Example 3.4. For λ in Example 2.2, the template for the rank triangles will look as follows.

$$\begin{array}{ccc} e_2 & e_1 & e_0 \\ \hline m_2 & m_1 & m_0 \\ \hline r_{21} & r_{10} & \\ & r_{20} & \end{array}$$

The orbit corresponding to the case $r_{21} = r_{10} = 1$ and $r_{20} = 0$ will have the following rank triangle.

$$\begin{array}{ccc} -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 1 & 1 & \\ & 0 & \end{array}$$

3.3. Correspondence between rank triangles and multisegments. We write down an algorithm to compute a rank triangle from a multisegment and vice-versa.

To each segment $\Delta = [e_j, e_i]$ we associate the rank triangle T_Δ with

$$r_{ls} = \begin{cases} 1, & j \leq s < l \leq i; \\ 0, & \text{otherwise.} \end{cases} \quad m_s = \begin{cases} 1, & j \leq s \leq i; \\ 0, & \text{otherwise.} \end{cases}$$

A multisegment \underline{m} then determines the triangle

$$T_{\underline{m}} = \sum_{\Delta \in \underline{m}} T_\Delta.$$

We now explain how to pass from a rank triangle with eigenvalues q^{e_i} and multiplicities (top row) m_i to a multisegment with support \underline{m}_λ by the following inductive rule:

- (1) Set $\underline{m} := \emptyset$.
- (2) Let r_{ij} be the lowest and most to the left non-zero entry in the rank triangle. Note that $j < i$.
- (3) Add the segment $\Delta = [e_i, e_j]$ to \underline{m} .
- (4) Replace r_{kl} by $r_{kl} - 1$ for each value of k, l with $j \leq l < k \leq i$. Replace m_k by $m_k - 1$ for each $j \leq k \leq i$. That is, subtract T_Δ from the rank triangle.
- (5) Repeat steps (ii) through (iv) until all r_{ij} are zero.
- (6) If any $m_k \neq 0$, add the singleton $\{e_k\}$ with multiplicity m_k to \underline{m} .

Now \underline{m} is the multisegment determined by the rank triangle. These procedures, $\underline{m} \leftrightarrow T_{\underline{m}}$ establish a bijection between rank triangles and multisegments.

In Definition 3.1 we define a partial order on multisegments and in Definition 3.3 we have a partial order on orbits (and therefore rank triangles). From [Zel81] (Theorem 2.2, Contiguity theorem) these partial orders are compatible under the correspondence. More precisely, let $\underline{m}, T_{\underline{m}}$, and $C_{\underline{m}}$ be a multisegment, its corresponding rank triangle and orbit respectively. Then the above statement can be rephrased as: If α and β are multisegments, then $\alpha \leq \beta$ if and only if $C_\alpha \leq C_\beta$ if and only if $T_\alpha \leq T_\beta$.

Example 3.5. Let us compute the multisegment for the rank triangle from Example 3.4.

$$\begin{array}{ccc}
 \begin{array}{c} \begin{array}{|c|c|c|} \hline -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 1 & & 1 \\ \hline & 0 & \\ \hline \end{array} \\ \Downarrow \\ [-1, 0] \end{array} & \longrightarrow & \begin{array}{c} \begin{array}{|c|c|c|} \hline -1 & 0 & 1 \\ \hline 0 & 1 & 1 \\ \hline 0 & & 1 \\ \hline & 0 & \\ \hline \end{array} \\ \Downarrow \\ [0, 1] \end{array} & \longrightarrow & \begin{array}{c} \begin{array}{|c|c|c|} \hline -1 & 0 & 1 \\ \hline 0 & 0 & 0 \\ \hline 0 & & 0 \\ \hline & 0 & \\ \hline \end{array} \\ \Downarrow \\ \text{Terminate} \end{array}
 \end{array}$$

The resulting multisegment is $\{[0, 1], [-1, 0]\}$. This orbit corresponds to the representation $Q([0, 1], [-1, 0])$. In this way, we can compute the multisegments for all orbits in the Vogan variety V from in Example 2.2.

$\Pi_\lambda(\mathrm{GL}_4(F))$	Rank triangle
$Q([-1, 1], [0])$	$ \begin{array}{ c c c } \hline -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 1 & & 1 \\ \hline & 1 & \\ \hline & & 1 \\ \hline \end{array} $
$Q([0, 1], [-1, 0])$	$ \begin{array}{ c c c } \hline -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 1 & & 1 \\ \hline & 0 & \\ \hline \end{array} $
$Q([1], [0], [-1, 0])$	$ \begin{array}{ c c c } \hline -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 1 & & 0 \\ \hline & 0 & \\ \hline \end{array} $
$Q([0, 1], [0], [-1])$	$ \begin{array}{ c c c } \hline -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 0 & 1 & \\ \hline & 0 & \\ \hline \end{array} $
$Q([1], [0], [0], [-1])$	$ \begin{array}{ c c c } \hline -1 & 0 & 1 \\ \hline 1 & 2 & 1 \\ \hline 0 & 0 & \\ \hline & 0 & \\ \hline \end{array} $

The open orbit in V consists of elements of full rank. From the table above, we see that the open orbit corresponds to the representation $Q([-1, 1], [0])$. On the other hand, having directly computed the multisegment $\{[-1, 0], [0, 1]\}$ for ϕ_ψ from Example 2.2, we now have its rank triangle.

4. ZELEVINSKY INVOLUTION ON MULTISEGMENTS AND ORBITS

We now work with an involution on orbits, often referred to as the Zelevinsky involution, which interchanges the Steinberg and trivial representations. This appears in [Zel80], [Zel81], and is expanded upon in [MW86]. We first write down an algorithm to compute the involution $\alpha \mapsto \tilde{\alpha}$ following [MW86] and then prove an important combinatorial lemma (Lemma 4.8). Finally, we see the analogous notions on orbits.

Let α denote a multisegment. This means that α is a collection of segments of the form $[b, e]$, where b and e are integers or half integers and are referred to as the base value and end values of the segment respectively.

4.1. Moeglin-Waldspurger algorithm. In this section, write down the Moeglin-Waldspurger algorithm [MW86] to compute the Zelevinsky involution on multisegments for GL_n . For this we recall what it means for one segment to precede the other. Let $\Delta_1 = [b_1, e_1]$ and $\Delta_2 = [b_2, e_2]$ be two segments.

Definition 4.1. Δ_1 is said to *precede* Δ_2 if and only if

$$b_1 < b_2, e_1 < e_2, \text{ and } b_2 \leq e_1 + 1.$$

We list the steps to compute $\tilde{\alpha}$ below.

First, we compute the segment $M(\alpha)$ associated to α :

- (1) Set e to be the largest end value that appears in any segment. Set $m := e$.
- (2) Consider all segments in α with end value m . Among these, choose a segment with the largest base value and call it Δ_m .
- (3) Consider the set of all segments in α that precede Δ_m with end value $m - 1$. If this is empty, go to step (5). Otherwise, choose a segment from this set with the largest base value and call it Δ_{m-1} .
- (4) Set $m := m - 1$ and go to step (3).
- (5) Return $M(\alpha) := [m, e]$.

Next, we inherit the following notation from the above procedure: for any $i \in \{m, m + 1, \dots, e\}$, set Δ_i as the segment whose base value is the highest among segments with end value i . Define $\alpha \setminus M(\alpha)$ to be the multisegment obtained by removing i from Δ_i for all $m \leq i \leq e$. Then, the dual $\tilde{\alpha}$ is recursively defined via

$$\tilde{\alpha} := (M(\alpha), \widetilde{\alpha \setminus M(\alpha)}).$$

The fact that this procedure gives you an involution comes from [MW86]. In particular, we have the useful property that $\tilde{\tilde{\alpha}} = \alpha$.

Example 4.2. Let us continue working in the setting of Example 2.2 and compute the involution on multisegments that appear:

α	$\tilde{\alpha}$
$\{-1, 1, [0]\}$	$\{-1, [0], [0], [1]\}$
$\{-1, 0, [0, 1]\}$	$\{-1, 0, [0, 1]\}$
$\{[1], [0], [-1, 0]\}$	$\{[0, 1], [0], [-1]\}$

4.2. Lemma about simple multisegments. The aim of this section is to prove Lemma 4.8. We establish some terminology and notation associated to a multisegment α . The length of a segment $[b, e]$ is simply the value $e - b + 1$.

L_α := Length of the longest segment in α .

n_α := number of segments in α

c_α := minimum number of segments that $\bigcup_{\Delta \in \alpha} \Delta$ can be broken into.

Lemma 4.3. Let α and β be any multisegments. If $\alpha \leq \beta$, then

- (1) $L_\alpha \leq L_\beta$.
- (2) $n_\alpha \geq n_\beta$.
- (3) $n_{\tilde{\alpha}} \geq L_\alpha$ and $n_\alpha \geq L_{\tilde{\alpha}}$.

Proof. (1) Since β is obtained by performing elementary operations on segments in α , the longest segment in β is either a segment from α or a union of two segments in α . Therefore, $L_\beta \geq L_\alpha$.

(2) Every pair of segments in α is replaced by at most two segments to form β . Therefore $n_\beta \leq n_\alpha$.

(3) Fix a segment in α . From the Moeclin-Waldspurger algorithm, one can observe that each element of that segment appears in a distinct segment in $\tilde{\alpha}$. In particular, this is true for the longest segment of α . Thus, we get $n_{\tilde{\alpha}} \geq L_\alpha$. Since $\tilde{\tilde{\alpha}} = \alpha$, we get the other inequality. \square

Definition 4.4. α is a simple multisegment if it is of the form

$$\{[b, e], [b + 1, e + 1], \dots, [b + n - 1, e + n - 1]\}.$$

Observe that n in the definition above is equal to n_α .

Lemma 4.5. If α is a simple multisegment, then

- (1) $\tilde{\alpha}$ is simple.
- (2) $n_{\tilde{\alpha}} = L_\alpha$.
- (3) $c_\alpha = 1$.

Proof. (1) Applying the Moeclin-Waldspurger algorithm, one can explicitly compute

$$\tilde{\alpha} = \{[e - k, e + n_\alpha - k - 1] : 0 \leq k \leq e - b\}.$$

Clearly, it is simple.

- (2) Observe that the length of every segment in α is the same, and $L_\alpha = e - b + 1$. Observing how the index k varies in part (1) tells us that $n_{\tilde{\alpha}} = e - b + 1$.
- (3) $c_\alpha = 1$ because

$$\bigcup_{\Delta \in \alpha} \Delta = [b, e + n - 1].$$

\square

Lemma 4.6. If $L_{\tilde{\alpha}} = n_\alpha$, $\alpha \leq \beta$, and $\tilde{\alpha} \leq \tilde{\beta}$, then

$$n_\alpha = n_\beta = L_{\tilde{\alpha}} = L_{\tilde{\beta}}.$$

The above statement also holds true if α is interchanged with $\tilde{\alpha}$ and β with $\tilde{\beta}$.

Proof. By hypothesis, $\tilde{\alpha} \leq \tilde{\beta}$. Therefore,

$$\begin{aligned} L_{\tilde{\alpha}} &\leq L_{\tilde{\beta}}, \text{ (from Lemma 4.3)} \\ &\leq n_{\tilde{\beta}}, \text{ (from Lemma 4.3)} \\ &= n_\beta, \text{ (since } \tilde{\tilde{\beta}} = \beta) \\ &\leq n_\alpha, \text{ (hypothesis and Lemma 4.3)} \\ &= L_{\tilde{\alpha}}. \text{ (hypothesis)} \end{aligned}$$

\square

Lemma 4.7. Let α be a simple multisegment and β any multisegment such that $\alpha \leq \beta$ and $L_\alpha \leq L_\beta$. Then, $\alpha = \beta$.

Proof. Suppose $\alpha < \beta$. Then, there is a segment in β of the form $\Delta_1 \cup \Delta_2$ strictly containing Δ_1 and Δ_2 , where $\Delta_1, \Delta_2 \in \alpha$. Now α is simple, so every segment has the same length and that length is equal to L_α . Thus, the strict containment above implies

$$L_\beta \geq \text{length of } \Delta_1 \cup \Delta_2 > \text{length of } \Delta_1 = L_\alpha,$$

which contradicts the hypothesis. \square

Now we are ready to state the primary result of this section.

Lemma 4.8. Suppose α is a simple multisegment and β is any multisegment satisfying $\alpha \leq \beta$ and $\tilde{\alpha} \leq \tilde{\beta}$, then $\alpha = \beta$.

Proof. α is simple, so from Lemma 4.5 $n_{\tilde{\alpha}} = L_\alpha$. Lemma 4.6 implies $L_\alpha = L_\beta$. The hypothesis of Lemma 4.7 is satisfied and thus $\alpha = \beta$. \square

4.3. Involution on orbits. Following [CFM⁺22, Section 6], we explain the notion of a dual orbit which is used to define an analogous involution on orbits $C \mapsto \hat{C}$ of V_λ .

From (8), we may write

$$V = \{x \in \mathfrak{gl}_n(\mathbb{C}) : \text{Ad}(\lambda(f))x = qx\}.$$

Using the killing form on $\mathfrak{gl}_n(\mathbb{C})$, the dual variety V^* may be realized as

$$V^* := \{x \in \mathfrak{gl}_n(\mathbb{C}) : \text{Ad}(\lambda(f))x = q^{-1}x\}.$$

We have a presentation of V^* analogous to (10):

$$V^* = \text{Hom}(E_1, E_0) \times \text{Hom}(E_{k-1}, E_{k-2}) \times \cdots \times \text{Hom}(E_k, E_{k-1}),$$

with an H -action

$$(h_k, \dots, h_1, h_0) \cdots (y_{10}, y_{21}, \dots, y_{kk-1}) = (h_0 y_{10} h_1^{-1}, h_1 y_{21} h_2^{-1}, \dots, h_{k-1} y_{kk-1} h_k^{-1}),$$

where $(h_k, \dots, h_1, h_0) \in H$ and $(y_{10}, y_{21}, \dots, y_{kk-1}) \in V^*$.

The cotangent variety $T^*(V)$ for V is simply $V \times V^*$. For $x \in V$ and $y \in V^*$, $[x, y] = xy - yx$ is the usual Lie bracket from $\mathfrak{gl}_n(\mathbb{C})$. The conormal variety, $\Lambda \subseteq T^*(V)$, is defined as the kernel of the map $[\cdot, \cdot] : T^*(V) \rightarrow \text{Lie}(H)$.

By [CFM⁺22, Proposition 6.3.1], for each H -orbit C in V ,

$$(17) \quad \Lambda_C := \{(x, y) \in \Lambda : x \in C\}$$

is the conormal bundle to C . By identifying $V^{**} \simeq V$, for an H -orbit C^* in V^* we may write

$$\Lambda_{C^*} = \{(x, y) \in \Lambda : x \in C^*\}$$

where Λ_{C^*} is the conormal bundle to C^* .

For any C , there exists a unique orbit C^* in V^* with $\bar{\Lambda}_C \simeq \bar{\Lambda}_{C^*}$; see [Pja75, Corollary 2]. The map $C \mapsto C^*$ defines a bijection the set of H -orbits in V and the set of H -orbits in V^* . Using this, we may define

$$\hat{C} := {}^t C^*,$$

where $y \mapsto {}^t y$ is matrix transposition. Then $C \mapsto \hat{C}$ defines an involution on the orbits.

In section 4.1, we wrote down the Zelevinsky involution on multisegments and in this section we see an involution on orbits. Since multisegments and orbits correspond to each other via rank triangles as seen in Section 3.3, it is natural to ask whether this involution is compatible under this correspondence. Crucially, it is. More precisely, if the multisegment α corresponds to the orbit C , then $\tilde{\alpha}$ corresponds to \hat{C} . See [KZ96b] for more details.

Example 4.9. From Example 4.2, we see that the involution applied to $\{-1, 1, [0]\}$ gives $\{[1], [0], [0], [-1]\}$. From Example 3.5, we see that the involution interchanges the open orbit (with elements of full rank) with the closed point (origin).

5. VOGAN'S CONJECTURE FOR SIMPLE ARTHUR PARAMETERS

Recall the simple Arthur parameter ψ defined in (3), the Langlands parameter ϕ_ψ coming from it defined in (4), and its infinitesimal parameter λ defined in (5). We remind the reader that $G = \mathrm{GL}_n$. Now, $\pi \in \Pi(G(F))$ is said to have the infinitesimal parameter λ if its corresponding Langlands parameter ϕ has the infinitesimal parameter λ . Let $\Pi_\lambda(G(F))$ denote all such π and $\Phi_\lambda(G(F))$ denote equivalence classes of all such ϕ . As outlined in sections 1.4 and 2, equivalence classes of smooth irreducible representations of $G(F)$ with infinitesimal parameter λ correspond to equivalence classes of simple objects in $\mathrm{Per}_H(V)$. This bijection is summarized below:

$$\begin{array}{ccccccc} \Pi_\lambda(G(F)) & \longrightarrow & \Phi_\lambda(G(F)) & \longrightarrow & H\text{-orbits in } V & \longrightarrow & \mathrm{Perv}_H(V)_{\mathrm{iso}}^{\mathrm{simple}}, \\ \pi & \longmapsto & \phi & \longmapsto & C_\phi & \longmapsto & \mathcal{P}(\pi) = \mathcal{IC}(\mathbb{1}_{C_\phi}). \end{array}$$

Thus, there is a unique orbit C_{ϕ_ψ} in V attached to ϕ_ψ . We shorten this notation to C_ψ . This orbit in turn corresponds to a simple multisegment - one can see this directly by writing down the multisegment for the Langlands parameter ϕ_ψ when $\psi(w, x, y) = \mathrm{Sym}^d(x) \otimes \mathrm{Sym}^a(y)$:

$$(18) \quad \left\{ \left[\frac{-(d+a)}{2}, \frac{-(d+a)}{2} + d \right], \left[\frac{-(d+a)}{2} + 1, \frac{-(d+a)}{2} + d + 1 \right], \dots, \left[\frac{-(d+a)}{2} + a, \frac{-(d+a)}{2} + d + a \right] \right\}.$$

The dictionary between Langlands parameters and multisegments is written down in many places, see [Kud94, Section 4.2], for example. In Section 3.3 we saw that rank triangles and multisegments correspond to each other and this correspondence respects the partial order on both sides. In section 4.3, we see that this correspondence respects the involution on both sides. Thus, we may adapt Lemma 4.8 as follows:

Lemma 5.1. Let C_ψ be the orbit corresponding to a simple Arthur parameter as explained above. Let C be any H -orbit in V . If $C_\psi \leq C$ and $\widehat{C}_\psi \leq \widehat{C}$, then $C_\psi = C$.

Lemma 5.2. Let C_ψ be the orbit corresponding to a simple Arthur parameter. Then

$$\widehat{C}_\psi = C_{\widehat{\psi}},$$

where $\widehat{\psi}$ is the simple Arthur parameter obtained by interchanging the role of the two $\mathrm{SL}_2(\mathbb{C})$'s in W_F'' :

$$\widehat{\psi}(w, x, y) := \psi(w, y, x).$$

Proof. Since ψ is simple, C_ψ corresponds to the multisegment written in (18). The dual of this multisegment, in the sense of 4.1, is

$$\left\{ \left[\frac{-(d+a)}{2}, \frac{-(d+a)}{2} + a \right], \left[\frac{-(d+a)}{2} + 1, \frac{-(d+a)}{2} + a + 1 \right], \dots, \left[\frac{-(d+a)}{2} + d, \frac{-(d+a)}{2} + d + a \right] \right\}.$$

By inspection, this multisegment comes from the simple Arthur parameter $(w, x, y) \mapsto \mathrm{Sym}^a(x) \otimes \mathrm{Sym}^d(y)$. From the discussion in 4.3, this multisegment also corresponds to \widehat{C}_ψ , as required. \square

We recall the functor defined in (1.5):

$$\text{Evs}_\psi : \text{Per}_{H_{\lambda_\psi}}(V_{\lambda_\psi}) \rightarrow \text{Rep}(A_\psi), .$$

For the Langlands parameter ϕ_ψ , we recall the ABV-packet from (2):

$$\Pi_{\phi_\psi}^{\text{ABV}}(G(F)) := \{\pi \in \Pi_\lambda(G(F)) : \text{Evs}_\psi(\mathcal{P}(\pi)) \neq 0\}.$$

We now state and prove the main theorem of this paper.

Theorem 5.3. *Let ψ be a simple Arthur parameter of $G = \text{GL}_n$, ϕ_ψ its corresponding Langlands parameter, $\Pi_\psi(G(F))$ the Arthur packet attached to ψ , and $\Pi_{\phi_\psi}^{\text{ABV}}(G(F))$ the ABV-packet attached to ϕ_ψ . Then,*

$$\Pi_{\phi_\psi}^{\text{ABV}}(G(F)) = \Pi_\psi(G(F)).$$

Proof. The local Langlands correspondence tells us that L -packets are singletons. As discussed in 1.3, the Arthur packet $\Pi_\psi(G(F))$ is a singleton and it coincides with the L -packet for ϕ_ψ . We may write

$$\Pi_\psi(G(F)) = \Pi_{\phi_\psi}(G(F)) = \{\pi_\psi\}.$$

Let C_ψ be the unique orbit associated to the Langlands parameter ϕ_ψ . By [CFM⁺22, Theorem 7.19], $\text{Evs}_\psi \mathcal{IC}(\mathbb{1}_{C_\psi}) \neq 0$. By definition, this implies

$$\Pi_{\phi_\psi}(G(F)) \subseteq \Pi_{\phi_\psi}^{\text{ABV}}(G(F)),$$

and therefore

$$\Pi_\psi(G(F)) \subseteq \Pi_{\phi_\psi}^{\text{ABV}}(G(F)).$$

To prove the theorem, it is sufficient to prove that $\Pi_{\phi_\psi}^{\text{ABV}}(G(F))$ is a singleton, *i.e.*, it contains no representation other than $\{\pi_\psi\}$.

Suppose we have a representation $\pi \in \Pi_{\phi_\psi}^{\text{ABV}}(G(F))$. By definition, this means

$$(19) \quad \text{Evs}_\psi(\mathcal{P}(\pi)) \neq 0.$$

From the discussion in sections 1.4 and 2, we know that π corresponds to an H -orbit C in V and $\mathcal{P}(\pi) = \mathcal{IC}(\mathbb{1}_C)$. We may rephrase (19) as

$$(20) \quad \text{Evs}_\psi \mathcal{IC}(\mathbb{1}_C) \neq 0.$$

Using [CFK22, Proposition 3.2.1] and Lemma 5.2, (19) also implies that

$$(21) \quad \text{Evs}_{\widehat{\psi}} \mathcal{IC}(\mathbb{1}_{\widehat{C}}) \neq 0.$$

Again from [CFK22, Proposition 3.2.1], (20) implies that $C_\psi \leq C$ and (21) implies $\widehat{C}_\psi \leq \widehat{C}$. The hypothesis of Lemma 5.1 is satisfied, so we may conclude that $C_\psi = C$. The corresponding representations are the same, as required. □

6. VOGAN'S CONJECTURE FOR IRREDUCIBLE ARTHUR PARAMETERS

We now extend the result in Theorem 5.3 to an *irreducible* Arthur parameter of GL_n , *i.e.*, of the form

$$(22) \quad \psi(w, x, y) = \rho(w) \otimes \mathrm{Sym}^d(x) \mathrm{Sym}^a(y),$$

where ρ is an irreducible admissible representation of W_F of dimension m , and a and d are non-negative integers. First note that $\rho = \rho_0 \otimes |\cdot|^s$, where ρ_0 is of Galois type and $|\cdot|$ is the absolute value on the Weil group as explained in section 2.

We apply the procedure of hyper-unramification as explained in [CFM⁺22, Section 5] on the infinitesimal parameter λ . Note that $\lambda(w) = \rho_0(w) \otimes |w|^s \otimes \mathrm{Sym}^d(d_w) \otimes \mathrm{Sym}^a(d_w)$ where d_w is defined in Section 2. For any $X \in \mathrm{SL}_2(\mathbb{C})$, set $\eta(X) := \mathrm{Sym}^d(X) \otimes \mathrm{Sym}^a(X)$.

Lemma 6.1. The semisimple element $\lambda(\mathfrak{f})$ has the decomposition $\lambda(\mathfrak{f}) = s_\lambda t_\lambda$ where

$$s_\lambda = \rho_0 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes |\mathfrak{f}|^{\mathrm{Re}(s)} \otimes \eta(d_{\mathfrak{f}})$$

is the hyperbolic part and

$$t_\lambda = \rho_0(\mathfrak{f}) \otimes |\mathfrak{f}|^{i \mathrm{Im}(s)} \otimes \eta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

is the elliptic part.

Proof. Clearly, $s_\lambda t_\lambda = t_\lambda s_\lambda = \lambda(\mathfrak{f})$. To show that s_λ is hyperbolic, it suffices to show that its eigenvalues are positive real numbers. Indeed, after s_λ is identified with a matrix, its eigenvalues are of the form $q^{\mathrm{Re} s + \frac{a+d}{2} + j}$ where j is an integer. This can be seen from direct calculation aided by Equation 6. To show that t_λ is elliptic, it suffices to show that its eigenvalues have complex norm 1. Indeed, t_λ can be identified with the (block) matrix

$$\begin{bmatrix} \rho_0(\mathfrak{f})q^{i \mathrm{Im} s} & 0 & \cdots & 0 \\ 0 & \rho_0(\mathfrak{f})q^{i \mathrm{Im} s} & \cdots & 0 \\ \vdots & & \ddots & \\ 0 & \cdots & 0 & \rho_0(\mathfrak{f})q^{i \mathrm{Im} s} \end{bmatrix},$$

so complex norm of its eigenvalues are determined by those of $\rho_0(\mathfrak{f})$. However, ρ_0 is of Galois-type which means it has finite image. Thus $\rho_0(\mathfrak{f})$ is a matrix of finite order, whose eigenvalues must be roots of unity. □

The process of hyper-unramification described in [CFM⁺22, Section 5.3] adapted to our case gives us an unramified parameter $\lambda_{nr} : W_F \rightarrow J_\lambda$, where J_λ is $Z_{\hat{G}}(\lambda|_{I_F}, t_\lambda)$ whose image of Frobenius is the hyperbolic element s_λ . Using [CFM⁺22, Lemma 5.3] we have that

$$V_{\lambda_{nr}} = V_\lambda, \text{ and } H_{\lambda_{nr}} = H_\lambda^0.$$

Lemma 6.2. Suppose ψ is an irreducible Arthur parameter as defined in Equation 22. Then,

- (1) $J_\lambda = \mathrm{GL}_{n/m}(\mathbb{C})$.
- (2) $H_{\lambda_{nr}} = H_\lambda$

Proof. (1) By definition, $J_\lambda = Z_{\hat{G}}(\lambda|_{I_F}, t_\lambda)$. Now $\lambda(w)|_{I_F} = \rho(w)|_{I_F} \otimes \eta(d_w)|_{I_F}$. Since I_F consists of elements of absolute value 1, $\lambda(w)|_{I_F} = \rho_0(w)|_{I_F} \otimes \eta \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. As ρ_0 is of

Galois-type, it is determined by a finite list of elements. In particular, there are elements $\sigma_1, \sigma_2, \dots, \sigma_k$ in I_F so that their images (block matrices) of the form

$$\begin{bmatrix} \rho_0(\sigma_i) & 0 & \cdots & 0 \\ 0 & \rho_0(\sigma_i) & \cdots & 0 \\ \vdots & & \ddots & \\ 0 & 0 & \cdots & \rho_0(\sigma_i) \end{bmatrix}$$

determine $\rho_0|_{I_F}$.

$$\begin{aligned} J_\lambda &= Z_{\mathrm{GL}_n(\mathbb{C})}(\lambda(w)|_{I_F}, t_\lambda) \\ &= \{X \in \mathrm{GL}_n(\mathbb{C}) : t_\lambda X t_\lambda^{-1} = X \text{ and } \lambda(\sigma_i) X \lambda(\sigma_i)^{-1} = X \text{ for } 1 \leq i \leq k\} \\ &= \{X = (X_{rs})_{r,s} \in \mathrm{GL}_n(\mathbb{C}) \text{ where } X_{rs} \in \mathrm{Mat}_m(\mathbb{C}) : \rho_0(\mathfrak{f}) X_{rs} \rho_0(\mathfrak{f})^{-1} = X_{rs} \text{ and } \rho_0(\sigma_i) X_{rs} \rho_0(\sigma_i)^{-1} = X_{rs}\} \\ &= \{X = (X_{rs})_{r,s} \in \mathrm{GL}_n(\mathbb{C}) \text{ where } X_{rs} \in \mathrm{Mat}_m(\mathbb{C}) : X_{rs} \in Z_{\mathrm{Mat}_m(\mathbb{C})}(\rho_0)\} \\ &= \{X = (c_{rs} \mathrm{Id}_m)_{r,s} \in \mathrm{GL}_n(\mathbb{C}) \text{ where } c_{rs} \in \mathbb{C}\} \\ &\simeq \mathrm{GL}_{n/m}(\mathbb{C}). \end{aligned}$$

The first equality is by definition, the second by the fact that $\lambda|_{I_F}$ is determined by $\sigma_i s$ as explained above, the third follows by writing down X, t_λ , and $\lambda(\sigma_i)s$ as block matrices, the fourth follows from the fact that ρ_0 is completely determined by \mathfrak{f} and $\sigma_i s$, and the fifth follows from Schur's lemma applied to the irreducible representation ρ_0 for every r, s . The isomorphism in the sixth step comes from realizing $(c_{rs} \mathrm{Id}_m)_{r,s}$ as $(c_{rs})_{r,s} \otimes \mathrm{Id}_m$ whose determinant is $\det((c_{rs})_{r,s})^m$ using the determinant of the Kronecker product of two matrices.

(2) As centralizers in $\mathrm{GL}_n(\mathbb{C})$ are connected, H_λ is connected so $H_{\lambda_{nr}} = H_\lambda^0 = H_\lambda$. \square

The above lemma shows that if we replace J_λ with the isomorphic general linear group, we may interpret the unramified infinitesimal parameter as a homomorphism $W_F \rightarrow \mathrm{GL}_{n/m}(\mathbb{C})$ whose image at the Frobenius can now be thought of as $|\mathfrak{f}|^{\mathrm{Res}} \otimes \eta(d_{\mathfrak{f}})$, and whose Vogan variety and group acting on it remain the same as that of λ . Also observe that removing the factor $|\mathfrak{f}|^{\mathrm{Res}}$ from this infinitesimal parameter does not change the variety and the group acting on it for trivial reasons. We collect these observations on hyper-unramification of ψ as defined in 22 in the following lemma.

Lemma 6.3. Suppose ψ is an *irreducible* Arthur parameter of GL_n , *i.e.*,

$$\psi(w, x, y) = \rho(w) \otimes \mathrm{Sym}^d(x) \otimes \mathrm{Sym}^a(y),$$

where ρ is an irreducible representation of W_F in $\mathrm{GL}_m(\mathbb{C})$. We obtain the unramified parameter

$$\lambda_{nr} : W_F \rightarrow \mathrm{GL}_{n/m}(\mathbb{C})$$

defined via $w \mapsto \mathrm{Sym}^d(d_w) \otimes \mathrm{Sym}^a(d_w)$, with

$$V_{\lambda_{nr}} \simeq V_\lambda, \text{ and } H_{\lambda_{nr}} \simeq H_\lambda,$$

where these isomorphisms are obtained by an application of Schur's lemma, as in the proof of Lemma 6.2 (1).

The upshot is that λ_{nr} can be thought of as the infinitesimal parameter of $\psi_{nr}(w, x, y) := \mathrm{Sym}^d(x) \otimes \mathrm{Sym}^a(y)$, after an application of the isomorphisms $V_{\lambda_{nr}} \simeq V_\lambda$ and $H_{\lambda_{nr}} \simeq H_\lambda$ that are induced by taking scalar matrices to their respective scalars in a smaller matrix group. Since the

variety and group action are unchanged, the category $\text{Per}_{H_{\lambda_{nr}}}(V_{\lambda_{nr}}) = \text{Per}_{H_{\lambda}}(V_{\lambda})$. As a result, we directly see that the isomorphisms give us a bijection $C \mapsto C_{nr}$ of orbits which preserves the ordering and the Zelevinsky involution. The geometric side of the Vogan-Langlands correspondence can thus be reduced to the study of the unramified infinitesimal parameter. This paves the way for generalizing Theorem 5.3 from simple Arthur parameters to irreducible Arthur parameters.

Theorem 6.4. *Suppose ψ is an irreducible Arthur parameter of $G = \text{GL}_n$, i.e.,*

$$\psi(w, x, y) = \rho(w) \otimes \text{Sym}^d(x) \otimes \text{Sym}^a(y),$$

where ρ is an irreducible representation of W_F on $\text{GL}_m(\mathbb{C})$. Let $\Pi_{\psi}(G(F))$ denote the local Arthur packet of ψ and $\Pi_{\phi_{\psi}}^{\text{ABV}}(G(F))$ denote the ABV-packet attached to ϕ_{ψ} . Then,

$$\Pi_{\psi}(G(F)) = \Pi_{\phi_{\psi}}^{\text{ABV}}(G(F)).$$

Proof. From Lemma 6.3, $V_{\lambda} \simeq V_{\lambda_{nr}}$ and $H_{\lambda} \simeq H_{\lambda_{nr}}$. We have a bijection $C \mapsto C_{nr}$ between orbits in V_{λ} and $V_{\lambda_{nr}}$ which preserves partial order and the Zelevinsky involution on orbits. This gives a bijection $\pi \mapsto \pi_{nr}$ between $\Pi_{\lambda}(G(F))$ and $\Pi_{\lambda_{nr}}(G(F))$ preserving the partial order and involution on multisegments such that $(C_{\pi})_{nr} = C_{\pi_{nr}}$. In particular, $(C_{\psi})_{nr} = C_{\psi_{nr}}$. Since Evs_{ψ} is purely dependent on the geometry, we have that $\text{Evs}_{\psi} \mathcal{IC}(\mathbb{1}_{C_{\pi}}) \neq 0$ for $\pi \in \Pi_{\lambda}(G(F))$ if and only if $\text{Evs}_{\psi_{nr}} \mathcal{IC}(\mathbb{1}_{C_{\pi_{nr}}}) \neq 0$ for the corresponding $\pi_{nr} \in \Pi_{\lambda_{nr}}(G(F))$. Since the ABV-packet for ψ_{nr} is a singleton from Theorem 5.3, same is true for the ABV packet for ψ . \square

Remark 6.5. The process of hyper-unramification taking $\psi \mapsto \psi_{nr}$ gives us a bijection between $\Pi_{\lambda}(G(F))$ and $\Pi_{\lambda_{nr}}(G(F))$ which can be written down explicitly in terms of multisegments. Let π_{ρ} be the supercuspidal representation of $\text{GL}_m(F)$ corresponding to the irreducible representation ρ . The correspondence at the level of multisegments is induced by taking any segment of the form $[\pi_{\rho}(b), \pi_{\rho}(e)]$ to the segment $[b, e]$.

REFERENCES

- [ABV92] Jeffrey Adams, Dan Barbasch, and David A. Vogan Jr., *The Langlands classification and irreducible characters for real reductive groups*, Progress in Mathematics, vol. 104, Birkhäuser Boston, Inc., Boston, MA, 1992.
- [Art13] James Arthur, *The endoscopic classification of representations*, American Mathematical Society Colloquium Publications, vol. 61, American Mathematical Society, Providence, RI, 2013. Orthogonal and symplectic groups.
- [Art90] ———, *Unipotent automorphic representations: global motivation*, Automorphic forms, Shimura varieties, and L -functions, Vol. I (Ann Arbor, MI, 1988), Perspect. Math., vol. 10, Academic Press, Boston, MA, 1990, pp. 1–75. MR1044818
- [Art89] ———, *Unipotent automorphic representations: conjectures*, Astérisque, 1989.
- [Bor79] A Borel, *Automorphic L -functions*, Automorphic forms, representations, and L -functions Part 2 (Proc. Sympos. Pure Math, Corvallis XXXIII), American Mathematical Society, Providence, RI, 1979. p. 27-61.
- [CFK22] Clifton Cunningham, Andrew Fiori, and Nicole Kitt, *Appearance of the Kashiwara-Saito singularity in the representation theory of p -adic GL_{16}* , Pacific Journal of Mathematics (2022). To appear. Available at <https://arxiv.org/abs/2103.04538>.
- [CFM⁺22] Clifton Cunningham, Andrew Fiori, Ahmed Moussaoui, James Mracek, and Bin Xu, *Arthur packets for p -adic groups by way of microlocal vanishing cycles of perverse sheaves, with examples*, Memoirs of the American Mathematical Society, vol. 276, AMS, 2022.
- [HT01] Michael Harris and Richard Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001. With an appendix by Vladimir G. Berkovich.
- [Hen00] Guy Henniart, *Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique*, Inventiones mathematicae **139** (2000), no. 2, 439–455.

- [Kal16] Tasho Kaletha, *The Local Langlands Conjectures for Non-quasi-split Groups* (Werner Müller, Sug Woo Shin, and Nicolas Templier, eds.), Springer International Publishing, 2016.
- [KZ96a] Harold Knight and Andrei Zelevinsky, *Representations of quivers of type A and the multisegment duality*, Adv. Math. **117** (1996), no. 2, 273–293.
- [KZ96b] ———, *Representations of quivers of type A and the multisegment duality*, Advances in mathematics **117** (1996), no. 2, 273–293.
- [Kud94] Stephen S Kudla, *The local Langlands correspondence: the non-archimedean case*, Motives (Seattle, WA, 1991) **55** (1994), no. Part 2, 365–391.
- [Lus91] George Lusztig, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), no. 2, 365–421.
- [Lus95] ———, *Classification of unipotent representations of simple p -adic groups*, Internat. Math. Res. Notices **11** (1995), 517–589.
- [MW86] Colette Mœglin and Jean-Loup Waldspurger, *Sur l’involution de Zelevinski*, J. Reine Angew. Math. **372** (1986), 136–177.
- [Pja75] V. S. Pjaseckii, *Linear Lie groups that act with a finite number of orbits*, Akademiya Nauk SSSR. Funkcional’nyi Analiz i ego Priloženija **9** (1975), no. 4, 85–86.
- [Vog93] David A. Vogan Jr., *The local Langlands conjecture*, Representation theory of groups and algebras, Contemp. Math., vol. 145, Amer. Math. Soc., Providence, RI, 1993, pp. 305–379.
- [Zel80] Andrei V. Zelevinsky, *Induced representations of reductive p -adic groups. II. On irreducible representations of $GL(n)$* , Ann. Sci. École Norm. Sup. (4) **13** (1980), no. 2, 165–210.
- [Zel81] Andrei Vladlenovich Zelevinskii, *p -adic analog of the kazhdan-Lusztig hypothesis*, Functional Analysis and Its Applications **15** (1981), no. 2, 83–92.