

# Non-virtually abelian discontinuous group actions vs. proper $SL(2, \mathbb{R})$ -actions on homogeneous spaces

Maciej Bocheński, Willem A. de Graaf, Piotr Jastrzębski,

Aleksy Tralle

October 25, 2022

## Abstract

We develop algorithms and computer programs which verify criteria of properness of discrete group actions on semisimple homogeneous spaces. We apply these algorithms to find new examples of non-virtually abelian discontinuous group actions on homogeneous spaces which do not admit proper  $SL(2, \mathbb{R})$ -actions.

*Key words and phrases:* proper action, homogeneous space, semisimple Lie algebra, GAP4, computing with simple Lie algebras.

*2020 Mathematics Subject Classification:* 17B05, 17B10, 53C30, 57S25, 57S30.

## 1 Proper actions of non-virtually abelian discrete subgroups

Assume that  $G$  is a linear connected semisimple real Lie group and  $H \subset G$  is a closed reductive subgroup of  $G$  with finitely many connected components such that  $G/H$  is non-compact. Denote by  $\mathfrak{g}, \mathfrak{h}$  the Lie algebras of  $G, H$  respectively. Recall that a discrete group is non-virtually abelian if it does not contain an abelian subgroup of finite index. Consider the following three conditions:

- C1 := the space  $G/H$  admits a properly discontinuous action of an infinite discrete subgroup of  $G$ ,

- C2 := the space  $G/H$  admits a properly discontinuous action of a non-virtually abelian infinite discrete subgroup of  $G$ ,
- C3 := the space  $G/H$  admits a proper action of a subgroup  $L \subset G$  locally isomorphic to  $SL(2, \mathbb{R})$ .

We will say that  $G/H$  is a  $C_i$  space,  $i = 1, 2, 3$  if  $G/H$  fulfills the condition  $C_i$ . If  $H$  is compact then every closed subgroup of  $G$  acts discontinuously on  $G/H$  and all the three conditions above hold. However this is not the case for non-compact  $H$ . For instance we have the Calabi-Markus phenomenon (see, for example [22], Corollary 4.4):

$$G/H \text{ is } C1 \text{ if and only if } \text{rank}_{\mathbb{R}} \mathfrak{g} > \text{rank}_{\mathbb{R}} \mathfrak{h}.$$

One of the reasons for considering these properties comes from the idea of the "continuous analogue" due to T. Kobayashi: under the above mentioned assumptions a homogeneous space  $G/H$  admits a proper action of an infinite discontinuous group if it admits a proper action of a one-dimensional non-compact closed subgroup in  $G$ . Going along this line of thinking, it was shown in [26] that a semisimple symmetric space admits a non-virtually abelian properly discontinuous action of a subgroup  $\Gamma \subset G$  if and only if it admits a proper action of a three-dimensional simple Lie subgroup of  $G$ . Thus, in our terminology, for the class of symmetric spaces, C2 and C3 are equivalent ([26], Theorem 2.2). Some other classes of homogeneous spaces of class C3 were described in [10] and of class C2 in [7]. In [1] a necessary and sufficient condition for  $G/H$  to satisfy C2 was found. Let us explain it in some detail. Let  $K \subset G$  be a maximal compact subgroup of  $G$  and denote by  $\mathfrak{k}$  the Lie algebra of  $K$ . Let  $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$  be a Cartan decomposition of  $\mathfrak{g}$  induced by a Cartan involution fixing  $\mathfrak{k}$  and choose a maximal abelian subspace  $\mathfrak{a}$  of  $\mathfrak{p}$ . The little Weyl group is defined as  $W := N_K(\mathfrak{a})/Z_K(\mathfrak{a})$  where  $N_K(\mathfrak{a})$  and  $Z_K(\mathfrak{a})$  denote respectively the normalizer and the centralizer of  $\mathfrak{a}$  in  $K$ . We fix a positive system of the restricted root system of  $\mathfrak{g}$  determined by  $\mathfrak{a}$ . The finite group  $W$  acts on  $\mathfrak{a}$  by orthogonal transformations as a finite group generated by reflections in the hyperplanes determined by simple roots of the restricted root system of  $\mathfrak{g}$ . Let  $w_0 \in W$  be the longest element and put

$$\mathfrak{b} := \{X \in \mathfrak{a} \mid -w_0(X) = X\}.$$

We may assume that  $\theta|_{\mathfrak{h}}$  is a Cartan involution of  $\mathfrak{h}$  so that  $\mathfrak{h} = \mathfrak{k}_h + \mathfrak{p}_h$  is a Cartan decomposition of  $\mathfrak{h}$  such that  $\mathfrak{k}_h \subset \mathfrak{k}$ ,  $\mathfrak{p}_h \subset \mathfrak{p}$ . We also may assume that a maximal abelian subspace  $\mathfrak{a}_h$  of  $\mathfrak{p}_h$  belongs to  $\mathfrak{a}$  (see [22]). The following theorem yields the required criterion:

**Theorem 1** ([1], Theorem 1).  $G/H$  is C2 if and only if for every  $w$  in  $W$ ,  $w \cdot \mathfrak{a}_h$  does not contain  $\mathfrak{b}$ . In this case, one can choose a discrete subgroup of  $G$  acting properly discontinuously on  $G/H$  to be free and Zariski dense in  $G$ .

For example the space  $SL(3, \mathbb{R})/SL(2, \mathbb{R})$  fulfills the condition C1 but does not fulfill the condition C2 and therefore the condition C3 (see [1], Example 1). For many important classes of homogeneous spaces C2 and C3 are equivalent. For example, apart from irreducible symmetric spaces, this holds for some strongly regular homogeneous spaces ([2], Theorem 2 and Corollary 1). On the other hand, there are known exactly two examples of spaces which fulfill the condition C2 but not C3 [2, 27]. Thus for a space of reductive type we only have the following:

$$C1 \Leftrightarrow C2 \Leftrightarrow C3.$$

The criterion for  $G/H$  to be C3 is given in [22] in a general setting of the properness of the action of a Lie subgroup  $L \subset G$  on  $G/H$ . We describe it in Subsection 2.2. Studying conditions C2, C3 and relations between them is of great importance. We refer to a recent paper [23] for a more detailed account. However, let us mention two challenging problems from [23].

**Problem 1.** *Determine all pairs  $(G, H)$  such that  $G/H$  admits co-compact discontinuous actions of discrete subgroups  $\Gamma \subset G$ .*

This is a long standing open problem [20, 21, 22]. There are many partial results, for example, [4, 5, 6, 9, 25], however, the general case is not settled.

**Problem 2.** *Find a necessary and sufficient condition for  $G/H$  to admit a discontinuous action of a group  $\Gamma \subset G$  isomorphic to a surface group  $\pi_1(\Sigma_g)$  with  $g \geq 2$ .*

Note that this problem is the same as C3 for the class of symmetric spaces [26]. The general case is still open. Some partial results on homogeneous spaces which are C3 are obtained in [10]. Also, in this context an important question arises.

**Question 1.** *What is the relation between classes C2 and C3?*

The known two examples of homogeneous spaces  $G/H$  which are C2 but not C3 are obtained in [2] and [27] by applying the criterion of the properness of the Lie subgroup action on  $G/H$  (see Subsection 2.2) and Theorem 1. One shows that there is an appropriate subspace  $\mathfrak{a}_m \subset \mathfrak{a}$  such that  $W\mathfrak{a}_m$  does not contain  $\mathfrak{b}$  and that (by Kobayashi's criterion of proper actions and the classification of nilpotent orbits) for any  $\mathfrak{sl}(2, \mathbb{R}) \hookrightarrow \mathfrak{g}$  there exists  $g \in G$  such that

$$\text{Ad}_g(\mathfrak{sl}(2, \mathbb{R})) \cap \mathfrak{a}_m \neq \{0\}.$$

In this case  $G/A_m$ , where the subgroup  $A_m \subset G$  corresponds to  $\mathfrak{a}_m$ , is a C2 space but not a C3 space. Notice that these examples are quotients obtained by explicit calculations. The calculations are fairly easy because in these cases the description of the embedding  $\mathfrak{h} \hookrightarrow \mathfrak{g}$  in terms of the root systems is simple. This description enables one to write down the action of the little Weyl group on  $\mathfrak{a}_m$  and use the known theoretical results. Thus, these examples are not sufficient for a qualitative understanding of the relation between C2 and C3. On the other hand, one can see that important Problems 1 and 2 theoretically are settled in terms of root systems and the action of the little Weyl group, by Theorems 1 and 3. The known examples of  $G/H$  which are C2 but not C3 are obtained in the same way. Therefore, from the computational point of view these become problems which could and should be attacked in an algorithmic fashion. We pose the following.

**Problem 3.** *Create computer algorithms and programs which verify C2 and C3 for general semisimple homogeneous spaces.*

Our aim is to solve this problem. We create algorithms verifying C2 versus C3 and implement them in the computational algebra system `GAP4` [15]. One can see that Theorem 1 yields a sufficient condition for the *non-existence* of compact quotients  $\Gamma \backslash G/H$ . Our approach yields methods of computer checking this as well. Note that a substantial amount of work related to the algorithms and computer programs for calculations in Lie algebras and a classification of semisimple subalgebras in semisimple Lie algebras relevant to this article was done in [8], [14], [17], [18].

The main application of our solution to Problem 3 is finding new examples of semisimple homogeneous spaces which are C2 but not C3. Consider the class of homogeneous spaces  $G/H$ , where  $G$  is linear and simple, and  $H$  is a maximal proper semisimple subgroup of  $G$  and assume that  $\mathfrak{h}^c$  is maximal in  $\mathfrak{g}^c$  and  $\text{rank } G \leq 8$  (here  $\mathfrak{h}^c, \mathfrak{g}^c$  denote the complexifications of  $\mathfrak{h}, \mathfrak{g}$  respectively). The corresponding pairs  $(\mathfrak{g}, \mathfrak{h})$  are classified in [18]. This database will be denoted by  $\mathcal{GM}$ .

**Theorem 2.** *For any  $(\mathfrak{g}, \mathfrak{h}) \in \mathcal{GM}$  such that  $\text{rank } \mathfrak{g} \leq 6$  conditions C2 and C3 are equivalent. There are two new examples of  $(\mathfrak{g}, \mathfrak{h}) \in \mathcal{GM}$  with split  $\mathfrak{g}$  and  $\mathfrak{h}$  such that  $\text{rank } \mathfrak{g} = 7, 8$  which belong to C2 but not C3:*

$$(\mathfrak{e}_{7(7)}, \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{f}_{4(4)}), (\mathfrak{e}_{8(8)}, \mathfrak{f}_{4(4)} \oplus \mathfrak{g}_{2(2)}).$$

This theorem is proved by computation, using the implementation of the algorithms given in this paper. The arxiv version of this paper (available on

<https://arxiv.org/abs/2206.01069>) has an ancillary file which is a short GAP program that checks that the two homogeneous spaces listed in the theorem indeed do not satisfy C3.

Notice that the assumption of  $H$  being maximal in  $G$  is very interesting in the context of searching for homogeneous spaces which are C2 but not C3. On one hand, one should expect that homogeneous spaces which are not C3 are given by “large” semisimple subgroups of  $G$  (to be more specific, by subgroups which have as large real rank as possible). On the other hand we are restricted by the condition C2 (so we cannot take  $\text{rank}_{\mathbb{R}} \mathfrak{g} = \text{rank}_{\mathbb{R}} \mathfrak{h}$ ).

Finally, let us stress that the main results of this article are algorithms and their implementation which check C2 and C3, as well as Theorem 2. These use theoretical results contained in Theorem 1 (Benoist), Theorem 3 (Kobayashi), Theorem 4 and Proposition 1 (Okuda).

## 2 Lie theory approach

Let us note that our notation and terminology is close to the sources [28, 29]. Throughout this section  $G$  is a real reductive linear connected Lie group with the Lie algebra  $\mathfrak{g}$ .

### 2.1 $\mathfrak{sl}(2, \mathbb{R})$ -triples and antipodal hyperbolic orbits

We say that an element  $X \in \mathfrak{g}$  is *hyperbolic*, if  $X$  is semisimple (that is,  $ad_X$  is diagonalizable) and all eigenvalues of  $ad_X$  are real.

**Definition 1.** An adjoint orbit  $O_X := Ad_G X$  is said to be hyperbolic if  $X$  (and therefore every element of  $O_X$ ) is hyperbolic. An orbit  $O_Y$  is antipodal if  $-Y \in O_Y$  (and therefore for every  $Z \in O_Y$ ,  $-Z \in O_Y$ ).

**Proposition 1** ([26], Proposition 4.5). *Let  $X \in \mathfrak{g}$  be hyperbolic. The intersection of the complex hyperbolic orbit of  $X$  in  $\mathfrak{g}^c$  with  $\mathfrak{g}$  is a single adjoint orbit  $O_X$  in  $\mathfrak{g}$ .*

A triple  $(h, e, f)$  of vectors in  $\mathfrak{g}$  is called an  $\mathfrak{sl}(2, \mathbb{R})$ -triple if

$$[h, e] = 2e, \quad [h, f] = -2f, \quad \text{and} \quad [e, f] = h.$$

In what follows we will write  $\mathfrak{sl}_2$ -triple instead of  $\mathfrak{sl}(2, \mathbb{R})$ -triple, and call  $h$  a neutral element. One can show that  $e$  is nilpotent and the adjoint orbit  $Ad_G h$  is antipodal and hyperbolic (see [11]). Furthermore there is the homomorphism of Lie groups

$$\Phi : SL(2, \mathbb{R}) \rightarrow G$$

defined by  $d\Phi \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = h$ ,  $d\Phi \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = e$ ,  $d\Phi \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = f$ .

## 2.2 Proper $SL(2, \mathbb{R})$ -actions and a properness criterion

Let  $H$  and  $L$  be reductive subgroups in  $G$ . We may assume that there exists a Cartan involution  $\theta$  of  $\mathfrak{g}$  such that  $\theta|_{\mathfrak{h}}$ ,  $\theta|_{\mathfrak{l}}$  are Cartan involutions of  $\mathfrak{h}$ ,  $\mathfrak{l}$ , respectively. We may also assume that there are maximal abelian subalgebras  $\mathfrak{a}$ ,  $\mathfrak{a}_h$ ,  $\mathfrak{a}_l$  in  $\mathfrak{p}$ ,  $\mathfrak{p}_h$  and  $\mathfrak{p}_l$  satisfying the inclusions

$$\mathfrak{a}_h \subset \mathfrak{a}, \mathfrak{a}_l \subset \mathfrak{a}.$$

**Theorem 3** ([22], Theorem 4.1). *In the above settings, the following conditions are equivalent:*

- (i)  $H$  acts on  $G/L$  properly.
- (ii)  $L$  acts on  $G/H$  properly.
- (iii)  $\mathfrak{a}_h \cap W\mathfrak{a}_l = \{0\}$ .

We can restate this conditions in the language of hyperbolic orbits for the action of  $L = SL(2, \mathbb{R})$ .

**Theorem 4** ([26], Corollary 5.5). *Let  $H$  be a reductive subgroup of  $G$  and denote by  $\mathfrak{h}$  the Lie algebra of  $H$ . Let  $\Phi : SL(2, \mathbb{R}) \rightarrow G$  be a Lie group homomorphism, and denote its differential by  $\phi = d\Phi : \mathfrak{sl}(2, \mathbb{R}) \rightarrow \mathfrak{g}$ . We put*

$$h_\phi := \phi \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \in \mathfrak{g}.$$

*Then  $SL(2, \mathbb{R})$  acts on  $G/H$  properly via  $\Phi$  if and only if the real antipodal adjoint orbit through  $h_\phi$  in  $\mathfrak{g}$  does not meet  $\mathfrak{a}_h$ .*

Notice that, by the classification of nilpotent orbits, the number of subspaces  $\mathbb{R}h_\phi \subset \mathfrak{a}$  is finite. Also, all  $h_\phi \in \mathfrak{a}$  are known (see [11] for details).

## 2.3 a-hyperbolic rank

Define the *a-hyperbolic rank* of a reductive Lie algebra  $\mathfrak{g}$ , denoted by  $\text{rank}_{\text{a-hyp}} \mathfrak{g}$ , as the dimension of  $\mathfrak{b}$ . The a-hyperbolic rank of a reductive Lie algebra equals the sum of a-hyperbolic ranks of all its simple ideals. The a-hyperbolic rank of an absolutely simple Lie algebras can be calculated using Table 1 (see [3] for more details about the a-hyperbolic rank).

This invariant will be used in the proof of Theorem 2.

$\mathfrak{g}$	$\text{rank}_{\mathfrak{a}\text{-hyp}}$
$\mathfrak{sl}(2k, \mathbb{R})$ $k \geq 1$	$k$
$\mathfrak{sl}(2k+1, \mathbb{R})$ $k \geq 1$	$k$
$\mathfrak{su}^*(4k)$ $k \geq 1$	$k$
$\mathfrak{su}^*(4k+2)$ $k \geq 1$	$k$
$\mathfrak{so}(2k+1, 2k+1)$ $k \geq 2$	$2k$
$\mathfrak{e}_{6(6)}$	$4$
$\mathfrak{e}_{6(-26)}$	$1$

Table 1: Real forms of simple Lie algebras  $\mathfrak{g}^c$ , with  $\text{rank}_{\mathbb{R}}(\mathfrak{g}) \neq \text{rank}_{\mathfrak{a}\text{-hyp}}(\mathfrak{g})$ .

### 3 Checking C3 and C2

In this section we describe computational methods for checking whether a given homogeneous space  $G/H$  is C3 and C2. We assume that  $G$  and  $H$  are given by their Lie algebras  $\mathfrak{g}$  and  $\mathfrak{h}$ . For the use of weighted Dynkin diagrams one can consult [3], [6].

#### 3.1 Checking C3

The following theorem follows directly from Theorem 4. It underpins our method for checking whether  $G/H$  is C3.

**Theorem 5.** *Let  $G \supset H$  be semisimple Lie groups with Lie algebras  $\mathfrak{g}$ ,  $\mathfrak{h}$ . Let  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$  be a Cartan decomposition of  $\mathfrak{g}$  and assume (as we may) that  $\mathfrak{h} = (\mathfrak{h} \cap \mathfrak{k}) \oplus (\mathfrak{h} \cap \mathfrak{p})$  is a Cartan decomposition of  $\mathfrak{h}$ . Let  $\mathfrak{a} \subset \mathfrak{p}$  be a maximal abelian subspace. Let  $\mathfrak{a}_h \subset \mathfrak{h} \cap \mathfrak{p}$  be a maximal abelian subspace with  $\mathfrak{a}_h \subset \mathfrak{a}$ . Let  $L \subset G$  be a subgroup isomorphic to  $\text{SL}(2, \mathbb{R})$ , with Lie algebra  $\mathfrak{l}$  spanned by  $h, e, f$ , with the usual commutation relations. Let  $W$  be the little Weyl group of  $\mathfrak{g}$  with respect to  $\mathfrak{a}$ . Then  $G/H$  admits a proper action of  $L$  if and only if the orbit  $W \cdot h$  has no point in  $\mathfrak{a}_h$ .*

We want to check whether there is a subgroup  $L$  as in the theorem acting properly on  $G/H$ . We assume that we are given the Lie algebras  $\mathfrak{g}$ ,  $\mathfrak{h}$ , along with a Cartan decomposition of  $\mathfrak{g}$  as in the theorem such that  $\mathfrak{h} = \mathfrak{k}_h \oplus \mathfrak{p}_h$ , where  $\mathfrak{k}_h = \mathfrak{k} \cap \mathfrak{h}$ ,  $\mathfrak{p}_h = \mathfrak{p} \cap \mathfrak{h}$ . We also fix a maximal abelian subspace  $\mathfrak{a} \subset \mathfrak{p}$  as in the theorem.

The first problem is to find a conjugate  $\mathfrak{a}'$  of  $\mathfrak{a}$  such that  $\mathfrak{a}' \cap \mathfrak{h}$  is a maximal abelian subspace of  $\mathfrak{p}_h$ . We do this in the following way.

1. Let  $\mathfrak{a}_h \subset \mathfrak{p}_h$  be a maximal abelian subspace. Set  $\hat{\mathfrak{a}} = \mathfrak{a}_h$  and while  $\dim \hat{\mathfrak{a}} < \dim \mathfrak{a}$  do the following
  - Compute the centralizer  $\mathfrak{z} = \mathfrak{c}_{\mathfrak{g}}(\hat{\mathfrak{a}})$ .
  - Find  $x \in \mathfrak{z} \cap \mathfrak{p}$  with  $x \notin \hat{\mathfrak{a}}$  and replace  $\hat{\mathfrak{a}}$  by the subalgebra spanned by  $\hat{\mathfrak{a}}$  and  $x$ .
2. Set  $\mathfrak{a}' = \hat{\mathfrak{a}}$ .

We note that this works because any abelian subspace of  $\mathfrak{p}$  lies in a maximally abelian subspace of  $\mathfrak{p}$ . So the spaces  $\mathfrak{z}$  always contain a maximally abelian subspace.

Now we write  $\mathfrak{a}$  instead of  $\mathfrak{a}'$ , and assume that  $\mathfrak{a}_h = \mathfrak{a} \cap \mathfrak{h}$  is a maximally abelian subspace of  $\mathfrak{p}_h$ .

We note that the subalgebras of  $\mathfrak{g}$  that are isomorphic to  $\mathfrak{sl}(2, \mathbb{R})$  are given by the nilpotent orbits of  $\mathfrak{g}$ . More precisely, we have the following. The nilpotent orbits in  $\mathfrak{g}^c$  are parametrized by weighted Dynkin diagrams. Each weighted Dynkin diagram uniquely determines an element  $h$  in a given Cartan subalgebra of  $\mathfrak{g}^c$ . This  $h$  lies in a non-uniquely determined  $\mathfrak{sl}_2$ -triple  $(h, e, f)$ . Let  $G_{\text{ad}}^c$  denote the adjoint group of  $\mathfrak{g}^c$ . Then the above construction gives a bijective correspondence between the list of weighted Dynkin diagrams and the  $G_{\text{ad}}^c$ -conjugacy classes of subalgebras of  $\mathfrak{g}^c$  that are isomorphic to  $\mathfrak{sl}_2$ .

By [26, Proposition 7.8] we have that a nilpotent orbit in  $\mathfrak{g}^c$  has points in  $\mathfrak{g}$  if and only if its weighted Dynkin diagram matches the Satake diagram of  $\mathfrak{g}$ . Moreover, these are precisely the nilpotent orbits having  $\mathfrak{sl}_2$ -triples lying in  $\mathfrak{g}$ . So in the next part of our procedure we perform the following steps:

1. Obtain the list of weighted Dynkin diagrams of  $\mathfrak{g}^c$ .
2. For the weighted Dynkin diagrams that match the Satake diagram of  $\mathfrak{g}$  compute an  $\mathfrak{sl}_2$ -triple  $(h, e, f)$  with  $h \in \mathfrak{a}$ . Let  $\mathcal{H}$  denote the set of all elements  $h$  that are obtained in this way.

By [26, Proposition 4.5] we have that  $\mathcal{H}$  is a set of representatives of the  $G$ -orbits of elements  $h' \in \mathfrak{g}$  that lie in an  $\mathfrak{sl}_2$ -triple  $(h', e', f')$ .

In order to apply Theorem 5 we need to construct the action of the little Weyl group  $W$  on  $\mathfrak{a}$ . Let  $\Sigma \subset \mathfrak{a}^*$  denote the set of roots of  $\mathfrak{g}$  with respect to  $\mathfrak{a}$  (see [29, Chapter 4, §4]). For each root  $\alpha \in \Sigma$  let  $P_\alpha = \{a \in \mathfrak{a} \mid \alpha(a) = 0\}$ . We define  $s_\alpha : \mathfrak{a} \rightarrow \mathfrak{a}$  to be the reflection in the hyperplane  $P_\alpha$ , where the inner product on  $\mathfrak{a}$  is given by the Killing form. Then the group generated by the  $s_\alpha$  is isomorphic to  $W$  ([29, Chapter 4, Proposition 4.2]).

Now for each  $h \in \mathcal{H}$  we check whether  $W \cdot h \cap \mathfrak{a}_h = \emptyset$ . If this holds then the subgroup of  $G$  corresponding to the subalgebra spanned by an  $\mathfrak{sl}_2$ -triple  $(h, e, f)$  acts properly on  $G/H$ . If the intersection is not empty for all  $h \in \mathcal{H}$  then  $G/H$  does not admit a proper action of a subgroup isomorphic to  $\text{SL}(2, \mathbb{R})$ .

## 3.2 Checking C2

Our procedure for checking whether  $G/H$  is a C2 space follows directly from Theorem 1. The main ingredients have already been described above. Indeed, we can compute the little Weyl group  $W$  and hence its longest element. It is then a straightforward task of linear algebra to compute a basis of the space  $\mathfrak{b}$ . By running over  $W$  we can check whether there is a  $w \in W$  such that  $w \cdot \mathfrak{a}_h$  contains  $\mathfrak{b}$ . The space  $G/H$  is C2 if and only if there is no such  $w$ .

## 3.3 Implementation

We have implemented the two procedures of this section in the computational algebra system GAP4 [15], using the package CoReLG [12]. This package has a database  $\mathcal{GM}$  of maximal reductive subalgebras of simple real Lie algebras, which was computed in [18]. Also we use the SLA package which has lists of weighted Dynkin diagrams for the nilpotent orbits of the semisimple complex Lie algebras.

Here we comment on two main computational problems that occur. Firstly, in order to compute the elements of  $W$  we need to compute the root system of  $\mathfrak{g}$  with respect to the space  $\mathfrak{a}$ . However, in some (but not many) cases this required eigenvalues that did not lie in the ground field that CoReLG uses. (This field is denoted `SqrtField`; here we do not go into the details.) So for a few cases our procedures could not be used.

Secondly, when  $\mathfrak{g}$  is a split form of type  $E_7$  or  $E_8$ , and the little Weyl group is equal to the usual Weyl group, the orbits  $W \cdot h$  can be very large. In this case we can handle large orbits by using an algorithm due to Snow ([30], see also [16, §8.6]) for running over an orbit of the Weyl group without computing all of its elements. This algorithm has been implemented in the core system of GAP4. Also we note two things:

- If we find  $h$  such that  $W \cdot h \cap \mathfrak{a}_h = \emptyset$  then we can stop.
- If  $W \cdot h \cap \mathfrak{a}_h \neq \emptyset$  then when enumerating the orbit  $W \cdot h$  we can reasonably hope to find an element in the intersection rather quickly.

So in practice it usually suffices to enumerate only part of the orbits.

## 3.4 Computational results

The package CoReLG contains a database  $\mathcal{GM}$  of maximal reductive subalgebras of the simple real Lie algebras up to rank 8. We have used our implementation to check whether  $G/H$  is C2 and C3 where the Lie algebras  $\mathfrak{g}$ ,  $\mathfrak{h}$  are taken from this database. (If the subalgebra  $\mathfrak{h}$  is reductive rather than semisimple we have taken its derived algebra instead.) We have done this for

non-compact  $\mathfrak{g}$  of rank up to 6, and for  $\mathfrak{g}$  equal to the split forms of type  $E_7$  and  $E_8$ . We have obtained the following results

- For the simple  $\mathfrak{g}$  up to rank 6 there are 842 maximal subalgebras in total. We found no examples of  $G/H$  that were C2 but not C3. Of these subalgebras 13 could not be dealt with by our programs for the reasons in Section 3.3. We list them in Table 2. Analyzing this table, we see that 9 of the spaces from this list do not yield  $G/H$  which is C2 but not C3 by theoretical considerations. This is because for such spaces either  $\text{rank}_{\mathbb{R}} \mathfrak{g} = \text{rank}_{\mathbb{R}} \mathfrak{h}$ , or  $\text{rank}_{\mathbb{R}} \mathfrak{h} = 1$ , and the result follows from the classification of nilpotent orbits, according to Section 4.2. The following homogeneous spaces from Table 2 are not covered by these arguments:

$$(\mathfrak{sp}(3, 3), \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{su}(1, 3)), (\mathfrak{so}(4, 6), \mathfrak{so}(2, 3)), (\mathfrak{e}_{6(2)}, \mathfrak{sl}(3, \mathbb{R})), (\mathfrak{e}_{6(2)}, \mathfrak{g}_{2(2)}).$$

Here we apply one more procedure to verify that these spaces are C3. We will show the following.

**Proposition 2.** *Consider a simple real Lie algebra  $\mathfrak{g}$  satisfying the assumption that  $\text{rank}_{a\text{-hyp}} \mathfrak{g} > 1$ . Assume that  $\mathfrak{h}$  is a reductive subalgebra such that  $\text{rank}_{\mathbb{R}} \mathfrak{h} = 2$ . Let  $\mathcal{P}$  denote the set of 2-planes in  $\mathfrak{a}$  with the following property:  $P \in \mathcal{P}$  if and only if it is generated by a pair of vectors  $w_1 h_1, w_2 h_2, w_i \in W$  and  $h_i$  are neutral elements of some  $\mathfrak{sl}_2$ -triple,  $i = 1, 2$ . Then*

- either  $G/H$  is C3, or
- $\mathfrak{a}_h \in \mathcal{P}$ .

*Proof.* If there is no  $\mathfrak{sl}_2$  orbit meeting  $\mathfrak{a}_h$  then  $G/H$  is C3. So assume that there is one  $w_1 h_1 \in \mathfrak{a}_h$ . Also assume that  $\mathfrak{a}_h \notin \mathcal{P}$  and  $G/H$  is not C3. Then each  $W$ -orbit of any neutral element must intersect  $\mathfrak{a}_h$ . It follows that this orbit must intersect the line  $\langle w_1 h_1 \rangle$ , because otherwise one would get  $\mathfrak{a}_h = \langle w_1 h_1, w_2 h_2 \rangle$  which is not the case. Hence, each  $W$ -orbit belongs to  $\langle w_1 h_1 \rangle$ . Since we can assume that  $\mathbb{R}^+(w_1 h_1) \subset C^+$ , where  $C^+$  is the positive Weyl chamber, we get  $\text{rank}_{a\text{-hyp}} \mathfrak{g} = 1$ , which contradicts the assumption. Thus, either  $G/H$  is C3, or  $\mathfrak{a}_h \in \mathcal{P}$ .  $\square$

This result implies the following.

**Proposition 3.** *Assume that  $\mathfrak{g} = \mathfrak{sp}(3, 3)$  or  $\mathfrak{g} = \mathfrak{e}_{6(2)}$  and let  $\mathfrak{h} \subset \mathfrak{g}$  be an arbitrary reductive subalgebra with  $\text{rank}_{\mathbb{R}} \mathfrak{h} = 2$ . Then  $(\mathfrak{g}, \mathfrak{h})$  corresponds to a homogeneous space  $G/H$  which is C3.*

*Proof.* Note that for both Lie algebras,  $\mathfrak{g} = \mathfrak{sp}(3, 3)$  and  $\mathfrak{g} = \mathfrak{e}_{6(2)}$ , the  $a$ -hyperbolic rank is greater than 1. By Proposition 2, it is sufficient to show that for any  $P \in \mathcal{P}$ , C3 condition holds. This is verified by the

(optimized) procedure described below. It is implemented and executed with the required result that all  $P \in \mathcal{P}$  are C3.  $\square$

**Remark 1.** Here we do not assume that  $(\mathfrak{g}, \mathfrak{h}) \in \mathcal{GM}$ , since the procedure which checks C3 is general and can be applied to all  $P \in \mathcal{P}$ .

### Simplified procedure

1. Write down all weighted Dynkin diagrams for  $\mathfrak{g}^c$  which match the Satake diagram for  $\mathfrak{g}$ .
2. Write down vectors  $A_1, \dots, A_t \in \mathfrak{a}$  whose coordinates are read off the weights of the weighted Dynkin diagram (here  $t$  is the number of the non-trivial weighted Dynkin diagrams of  $\mathfrak{g}^c$  which match the Satake diagram).
3. If some of  $A_1, \dots, A_t$  are on the same line, then choose only one of these, getting  $A_1, \dots, A_k$ .
4. Check, if there exist  $w_1, \dots, w_k \in W$  such that  $w_1 A_1, \dots, w_k A_k$  lie in a 2-plane.
5. If  $w_1, \dots, w_k \in W$  do not exist, then  $G/H$  cannot be "not C3".

The above procedure does not require the knowledge of the embedding  $\mathfrak{a}_h \subset \mathfrak{a}$ . By Theorem 4  $G/H$  is of class C3, if and only if at least one  $W$ -orbit of the element  $h_\phi$  does not meet  $\mathfrak{a}_h$ . In our case  $\dim \mathfrak{a}_h = 2$ , therefore, Step 4 of the algorithm is a necessary condition to get  $G/H$  not belonging to C3. This procedure cannot be implemented directly, because the number of operations is too big (it exceeds  $10^{60}$ ). However, we can optimize it to settle the cases of our 4 spaces.

### Optimized procedure

1. Compute the following sets of vectors in  $\mathfrak{a}$ :

$$WA_j = \{w_s A_j \mid w_s \in W\}, \quad X = \cup_{j=1}^k WA_j$$

2. Create the set  $\mathcal{P}$  of 2-planes in  $\mathfrak{a}$  spanned by  $A_1$  and a vector from  $X$ , so

$$P \in \mathcal{P} \Leftrightarrow P = \langle A_1, x \rangle, x \in X.$$

3. For each  $P \in \mathcal{P}$  verify C3 (according to Section 3.1).
4. If a given  $P \in \mathcal{P}$  is not C3, then stop. In such case there exists  $\mathfrak{h}$  reductive in  $\mathfrak{g}$  such that  $\text{rank}_{\mathbb{R}} \mathfrak{h} = 2$  and  $G/H$  is not C3 (one can take  $\mathfrak{h} := P$ ).
5. If every  $P \in \mathcal{P}$  is C3, then  $G/H$  is C3.

One can roughly estimate the number of operations considering the case of the biggest little Weyl group of  $E_{6(2)}$ . The number of  $A_j$  is 14 (see [11], Section 8.4), the number of elements in the little Weyl group is 1152. Take into account, that  $A_1$  is fixed, so  $|\mathcal{P}| \leq |X| - 1$  (in our case,  $\leq 15 \cdot 1151$ ).

- Let  $\mathfrak{g}$  be split of type  $E_7$  or  $E_8$ . We also checked the maximal semisimple subalgebras of these Lie algebras. Again a few subalgebras could not be dealt with by our programs. However, we did find two examples of homogeneous spaces that are C2 but not C3:

$$(\mathfrak{e}_{7(7)}, \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{f}_{4(4)}), (\mathfrak{e}_{8(8)}, \mathfrak{f}_{4(4)} \oplus \mathfrak{g}_{2(2)}).$$

$(\mathfrak{g}, \mathfrak{h})$	$(\mathfrak{g}, \mathfrak{h})$
$(\mathfrak{su}(3, 3), \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{su}(3))$	$(\mathfrak{su}(2, 5), \mathfrak{so}(2, 5))$
$(\mathfrak{so}(6, 5), \mathfrak{sl}(2, \mathbb{R}))$	$(\mathfrak{so}(6, 7), \mathfrak{sl}(2, \mathbb{R}))$
$(\mathfrak{sp}(4, \mathbb{R}), \mathfrak{sl}(2, \mathbb{R}))$	$(\mathfrak{sp}(5, \mathbb{R}), \mathfrak{sl}(2, \mathbb{R}))$
$(\mathfrak{sp}(3, 3), \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{su}(1, 3))$	$(\mathfrak{so}(6, \mathbb{R}), \mathfrak{sl}(2, \mathbb{R}))$
$(\mathfrak{so}(4, 6), \mathfrak{so}(2, 3))$	$(\mathfrak{e}_{6(2)}, \mathfrak{su}(1, 2))$
$(\mathfrak{e}_{6(2)}, \mathfrak{sl}(3, \mathbb{R}))$	$(\mathfrak{e}_{6(2)}, \mathfrak{g}_{2(2)})$
$(\mathfrak{f}_{4(4)}, \mathfrak{sl}(2, \mathbb{R}))$	

Table 2: Spaces with rank  $\mathfrak{g} \leq 6$  not covered by Sect. 3.

## 4 Proof of Theorem 2

The algorithms checking C2 and C3 for the spaces in the database are applied after filtering it with respect to certain conditions (see Subsection 4.2). Also, the database [18] which we use yields  $\mathfrak{h} \subset \mathfrak{g}$  up to an abstract isomorphism. Therefore, our first aim is to show that this classification is sufficient.

### 4.1 The sufficiency of the database

In what follows we need the notion of a linear equivalence of subalgebras. The definition can be found in [17].

**Proposition 4.** *Let  $G/H_i \in \mathcal{GM}$  ( $i = 1, 2$ ). Assume that  $\text{rank } G \leq 6$ . If  $\mathfrak{h}_1 \cong \mathfrak{h}_2$  then  $G/H_1$  belongs to C2 if and only if  $G/H_2$  belongs to C2. Analogously,  $G/H_1$  is of class C3 if and only if  $G/H_2$  is of class C3.*

*Proof.* By [24], Theorem 3, Theorem 7 and Example 1, for the spaces in Table 2 the linear equivalence is the same as  $\text{Ad}(g^c)$ -equivalence, with the following two exceptions:

$$(\mathfrak{e}_{6(2)}, \mathfrak{sl}(3, \mathbb{R})), \text{ and } (\mathfrak{e}_{6(2)}, \mathfrak{g}_{2(2)})$$

which correspond to complex embeddings

$$A_2 \subset E_6, G_2 \subset E_6.$$

Therefore, we consider the cases of  $\text{Ad}(g^c)$ -conjugate and non- $\text{Ad}(g^c)$ -conjugate subalgebras separately.

**Case 1.** There exists  $g^c \in G^c$  such that  $\text{Ad } g^c(\mathfrak{h}_1^c) = \mathfrak{h}_2^c$ . After conjugation we may assume that  $\mathfrak{h}_1^c = \mathfrak{h}_2^c$  in  $\mathfrak{g}^c$ . Now we want to show that this implies that there exists  $g \in G$  such that  $\text{Ad } g(\mathfrak{a}_h^1) = \mathfrak{a}_h^2$ . Assume that  $\mathfrak{a}_h^1$  is not conjugate to  $\mathfrak{a}_h^2$  in  $G$ . It follows that

$$\mathfrak{a}_h^1 \not\subset \text{Ad}_G(\mathfrak{a}_h^2).$$

Therefore, there exists  $0 \neq h \in \mathfrak{a}_h^1$  such that  $h \notin \text{Ad}_G(\mathfrak{a}_h^2)$ . Therefore, the hyperbolic orbit  $\text{Ad}_G(h)$  meets  $\mathfrak{h}_1$  but does not meet  $\mathfrak{h}_2$ . By [26], Theorem 7.4 (as  $\mathfrak{h}_1$  and  $\mathfrak{h}_2$  are isomorphic)  $\text{Ad}_{H^c}(h)$  meets  $\mathfrak{h}_2$ . Thus, there exists  $Z \in \mathfrak{h}_2 \cap \text{Ad}_{G^c}(h)$  such that

$$\text{Ad}_G(h) \neq \text{Ad}_G(Z), \text{ but } \text{Ad}_{G^c}(h) = \text{Ad}_{G^c}(Z).$$

This contradicts Proposition 1. We complete the proof by noticing that the condition  $\text{Ad } g(\mathfrak{a}_h^1) = \mathfrak{a}_h^2$  implies that  $G/H_1$  and  $G/H_2$  simultaneously satisfy or do not satisfy C2 and C3, which follows from Theorem 1 and Theorem 3, since these criteria are invariant with respect to the adjoint action of  $G$ .


**Case 2.** By [13], (Theorem 11.1) in each of these cases there are two embeddings

$$j_1 : A_2 \rightarrow E_6, j_2 : A_2 \rightarrow E_6, i_1 : G_2 \rightarrow E_6, i_2 : G_2 \rightarrow E_6$$

such that

$$j_2(A_2) = \varphi(j_1(A_2)), i_2 = \varphi(i_1(G_2))$$

where  $\varphi$  is an outer automorphism of  $\mathfrak{e}_6$ . Consider the case of embeddings  $j_1$  and  $j_2$  (the case of  $i_1$  and  $i_2$  can be treated in the same manner). Looking at the Satake diagram of  $\mathfrak{e}_{6(2)}$  one can easily see that  $\varphi$  fixes every point of  $\mathfrak{a}$ .



Satake diagram of  $\mathfrak{e}_{6(2)}$ .

□

## 4.2 Filtering of the database

We exclude some cases when considering the homogeneous spaces from the database  $\mathcal{GM}$ . Our filtering is based on the following theoretical results.

**Theorem 6** ([3], Theorem 8). *Under the assumptions of Theorem 2 the following holds:*

1. *if  $\text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{h} = \text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g}$ , then  $G/H$  does not admit discontinuous actions of non-virtually abelian discrete subgroups,*
2. *if  $\text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g} > \text{rank}_{\mathbb{R}} \mathfrak{h}$  than  $G/H$  admits a discontinuous action of non-virtually abelian discrete subgroup.*

**Proposition 5.** *Assume that  $\text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g} > 1$ . Then there exist two copies of  $\mathfrak{sl}(2, \mathbb{R})$  in  $\mathfrak{g}$ , with neutral elements  $h_1, h_2 \in \mathfrak{a}$  such that  $\mathbb{R}h_1$  does not meet the orbit  $W(h_2)$ .*

*Proof.* The proof follows from the analysis of the tables of nilpotent orbits [11] in exceptional real Lie algebras. It is clear that the Weyl group cannot send a vector lying on the wall of a Weyl chamber to the interior of a Weyl chamber. Checking the table for  $\text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g} > 1$  we see that in each case there are two neutral vectors, one in the interior.  $\square$

**Corollary 1.** *If  $\text{rank}_{\mathbb{R}} \mathfrak{h} = 1 < \text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g}$  then  $(\mathfrak{g}, \mathfrak{h})$  is C3.*

*Proof.* Assume that  $(\mathfrak{g}, \mathfrak{h})$  is not C3. By Theorem 4 there exists  $w_1 \in W$  such that  $\mathbb{R}w_1h_1 = \mathfrak{a}_h$  (as  $\mathfrak{a}_h$  is 1-dimensional) and  $w_2 \in W$  such that  $w_2h_2 \in \mathfrak{a}_h$ . But this contradicts Proposition 5.  $\square$

Theorem 6 and the Calabi-Markus phenomenon eliminate the necessity to consider the cases

1.  $\text{rank}_{\mathbb{R}} \mathfrak{g} = \text{rank}_{\mathbb{R}} \mathfrak{h}$ ,
2.  $\text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g} = \text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{h}$

since they are not C2. Finally, we eliminate the case  $\text{rank}_{\mathbb{R}} \mathfrak{h} = 1 < \text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g}$ . This follows from Corollary 1.

**Remark 2.** Looking for  $G/H$  which are C2 but not C3, we consider the cases

$$\text{rank}_{\mathfrak{a}\text{-hyp}} \mathfrak{g} > \text{rank}_{\mathbb{R}} \mathfrak{h}$$

since C2 is satisfied by Theorem 6, and C3 is checked by the general procedure.

### 4.3 Completion of the proof of Theorem 2

We complete the proof of Theorem 2 running the computer program which checks the remaining cases in the database.

**Acknowledgment.** The authors are grateful to the anonymous referees for valuable comments which improved the presentation of the results. The first named author was supported by the National Science Center, Poland, grant NCN 2018/31/D/ST1/00083. The fourth named author was supported by the National Science Center, Poland, grant 2018/31/B/ST1/00053.

## References

- [1] Y. Benoist, *Actions propres sur les espaces homogènes réductifs*, Ann. Math. 144 (1996), 315-347.
- [2] M. Bocheński, *Proper actions on strongly regular homogeneous spaces*, Asian J. Math. 21 (2017), 1121-1134.
- [3] M. Bocheński, A. Tralle, *Clifford-Klein forms and  $a$ -hyperbolic rank*, Int. Math. Res. Notices IMRN 5 (2015), 6267-6285.
- [4] M. Bocheński, A. Tralle, *On solvable compact Clifford-Klein forms*, Proc. Amer. Math. Soc. 145 (2017), 1819-1832
- [5] M. Bocheński, A. Tralle, *On the Hirzebruch-Kobayashi-Ono proportionality principle and the non-existence of compact solvable Clifford-Klein forms of certain homogeneous spaces*, Forum Math., 32 (2020), 471-477.
- [6] M. Bocheński, P. Jastrzębski, A. Tralle, *Non-existence of standard compact Clifford-Klein forms of homogeneous spaces of exceptional Lie groups*, Math. Comp. 89 (2020), 1487-1499.
- [7] M. Bocheński, P. Jastrzębski, A. Tralle, *Homogeneous spaces of real simple Lie groups with proper actions of non virtually abelian discrete subgroups: A computational approach*, J. Symb. Comp. 113 (2022), 171-180.
- [8] M. Bocheński, P. Jastrzębski, A. Tralle, *CKForms*. A GAP Package. Available online (<https://pjastr.github.io/CKForms/>).
- [9] M. Bocheński, P. Jastrzębski, A. Szczepkowska, A. Tralle, A. Woike, *Semisimple subalgebras in simple Lie algebras and a computational approach to the compact Clifford-Klein forms problem*, Exp. Math. 30 (2021), 86-94.
- [10] M. Bocheński, P. Jastrzębski, T. Okuda, A. Tralle, *Proper  $SL(2, \mathbb{R})$ -actions on homogeneous spaces*, Int. J. Math. 27 (2016), No. 13, 1-10.
- [11] D. H. Collingwood, W. M. McGovern, *Nilpotent Orbits in Semisimple Lie Algebras*, Van Nostrand Reinhold, New York (1993).

- [12] H. Dietrich, P. Faccin, and W. de Graaf, *CoReLG, computing with real Lie algebras, Version 1.54*, <https://gap-packages.github.io/corelg/> <https://gap-packages.github.io/corelg/>, Jan 2020, Refereed GAP package.
- [13] E. B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras* Mat. Sbornik N.S. 30 (72) (1952), 349-462.
- [14] P. Faccin, W. de Graaf, *Constructing semisimple subalgebras of real semisimple Lie algebras*, in: Lie Algebras and Related Topics, 75-89, Contemp. Math. 652 (2015), AMS, RI.
- [15] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.11.1*, 2021.
- [16] W. A. de Graaf *Lie algebras: theory and algorithms*. North-Holland Mathematical Library, 56. North-Holland Publishing Co., Amsterdam, 2000.
- [17] W. de Graaf, *Constructing semisimple subalgebras of semisimple Lie algebras*, J. Algebra 325 (2011), 416-430.
- [18] W. de Graaf, A. Marrani, *Real forms of embeddings of maximal reductive subalgebras of the complex simple Lie algebras of rank up to 8*, J. Phys. A 53 (2020), no. 15, 155203.
- [19] W. de Graaf, *SLA, computing with simple Lie algebras, Version 1.5.3*, <https://gap-packages.github.io/sla/>, Nov 2019, Refereed GAP package.
- [20] T. Kobayashi, T. Yoshino *Compact Clifford-Klein forms of symmetric spaces revisited*, Pure Appl. Math. Quart 1 (2005), 603-684.
- [21] T. Kobayashi *Discontinuous Groups for Non-Riemannian Homogeneous Spaces*, In: Engquist B., Schmid W. (eds) Mathematics Unlimited - 2001 and Beyond. Springer, Berlin 2001.
- [22] T. Kobayashi, *Proper actions on a homogeneous space of reductive type*, Math. Ann. 285 (1989), 249-263
- [23] T. Kobayashi, *Conjectures on reductive homogeneous spaces*, arxiv: 2204.0885 [math.RT].
- [24] A. N. Minchenko, *Semisimple subalgebras of exceptional Lie algebras*, Trans. Moscow Math. 2006, 225-259.
- [25] Y. Morita, *A topological necessary condition for the existence of compact Clifford-Klein forms*, J. Diff. Geom. 100 (2015), 533-545.
- [26] T. Okuda, *Classification of Semisimple Symmetric Spaces with Proper  $SL(2, \mathbb{R})$ -Actions*, J. Differential Geometry 2 (2013), 301-342.
- [27] T. Okuda, *Homogeneous space with non-virtually abelian abelian discontinuous groups but without any proper  $SL(2, \mathbb{R})$ -action*, Internat. J. Math. 27 (2016), art. no. 1650018.
- [28] A. L. Onishchik, E. B. Vinberg, *Lie groups and algebraic groups*, Springer (1990).

- [29] A. L. Onishchik, E. B. Vinberg *Lie Groups and Lie algebras III*, Springer (1994).
- [30] D. M. Snow, *Weyl group orbits*. ACM Trans. Math. Software 16 (1990), no. 1, 94-108.

Department of Mathematics,  
University of Trento,  
Via Sommarive 14,  
I-38123 Povo (Trento), Italy  
e-mail address: [degraaf@science.unitn.it](mailto:degraaf@science.unitn.it)

Faculty of Mathematics and Computer Science  
University of Warmia and Mazury  
Słoneczna 54, 10-710 Olsztyn, Poland  
e-mail addresses:  
[mabo@matman.uwm.edu.pl](mailto:mabo@matman.uwm.edu.pl) (MB)  
[piojas@matman.uwm.edu.pl](mailto:piojas@matman.uwm.edu.pl) (PJ)  
[tralle@matman.uwm.edu.pl](mailto:tralle@matman.uwm.edu.pl) (AT)