
Optimal Activation Functions for the Random Features Regression Model

Jianxin Wang
Department of Computer Science
Boston College
wangccy@bc.edu

José Bento
Department of Computer Science
Boston College
bentoayr@bc.edu

Abstract

The asymptotic mean squared test error and sensitivity of the Random Features Regression model (RFR) have been recently studied. We build on this work and identify in closed-form the family of Activation Functions (AFs) that minimize a combination of the test error and sensitivity of the RFR under different notions of functional parsimony. We find scenarios under which the optimal AFs are linear, saturated linear functions, or expressible in terms of Hermite polynomials. Finally, we show how using optimal AFs impacts well established properties of the RFR model, such as its double descent curve, and the dependency of its optimal regularization parameter on the observation noise level.

1 Introduction

For many neural network architectures, the test error does not monotonically increase as a model's complexity increases but can go down with the training error both at low and high complexity levels. This phenomenon, the *double descent curve*, defies intuition and has motivated new frameworks to explain it. Explanations have been advanced involving linear regression with random covariates [1, 2], kernel regression [3, 4], the *neural tangent kernel* model [5], and the *Random Features Regression* (RFR) model [6]. These frameworks allow queries beyond the generalization power of neural networks. For example, they have been used to study networks' robustness properties [7, 8].

One aspect within reach and unstudied to this day is finding optimal Activation Functions (AFs) for these models. It is known that AFs affect a network's approximation accuracy and efforts to optimize AFs have been undertaken. Previous work has justified the choice of AFs empirically, e.g., [9], or provided numerical procedures to learn AF parameters, sometimes jointly with models' parameters, e.g. [10]. See [11] for commonly used AFs and Section 2 for how AFs have been previously derived.

We derive for the first time closed-form optimal AFs such that an explicit objective function involving the asymptotic test error and sensitivity of a model is minimized. Setting aside empirical and principled but numerical methods, all past principled and analytical approaches to design AFs focus on non accuracy related considerations, e.g. [12]. We focus on AFs for the RFR model and expand its theoretical understanding. We preview a few surprising conclusions extracted from our main results:

1. The optimal AF can be linear, in which case the RFR model is a linear model. For example, if no regularization is used for training, and for low complexity models, a linear AF is often preferred if we want to minimize test error. For high complexity models a non-linear AF is often better;
2. A linear optimal AF can destroy the double descent curve behaviour and achieve small test error with much fewer samples than e.g. a ReLU;
3. When, apart from the test error, the sensitivity of a model becomes important, optimal AFs that without sensitivity considerations were linear can become non-linear, and vice-versa;
4. Using an optimal AF with an arbitrary regularization during training can lead to the same, or better, test error as using a non-optimal AF, e.g. ReLU, and optimal regularization.

1.1 Problem set up

We consider the effect of AFs on finding an approximation f to a square-integrable function f_d on the d -dimensional sphere $\mathbb{S}^{d-1}(\sqrt{d})$, the function f_d having been randomly generated. The approximation f is to be learnt from training data $\mathcal{D} = \{\mathbf{x}_i, y_i\}_{i=1}^n$ where $\mathbf{x}_i \in \mathbb{S}^{d-1}(\sqrt{d})$, the variables $\{\mathbf{x}_i\}_{i=1}^n$ are i.i.d. uniformly sampled from $\mathbb{S}^{d-1}(\sqrt{d})$, and $y_i = f_d(\mathbf{x}_i) + \epsilon_i$, where the noise variables $\{\epsilon_i\}_{i=1}^n$ are i.i.d. with $\mathbb{E}(\epsilon_i) = 0$, $\mathbb{E}(\epsilon_i^2) = \tau^2$, and $\mathbb{E}(\epsilon_i^4) < \infty$.

The approximation f is defined according to the RFR model. The RFR model can be viewed as a two-layer neural network with random first-layer weights encoded by a matrix $\Theta \in \mathbb{R}^{N \times d}$ with i th row $\theta_i \in \mathbb{R}^d$ satisfying $\|\theta_i\| = \sqrt{d}$, with $\{\theta_i\}$ i.i.d. uniform on $\mathbb{S}^{d-1}(\sqrt{d})$, and with to-be-learned second-layer weights encoded by a vector $\mathbf{a} = [a_i]_{i=1}^N \in \mathbb{R}^N$. Unless specified otherwise, the norm $\|\cdot\|$ denotes the Euclidean norm. The RFR model defines $f_{\mathbf{a}, \Theta} : \mathbb{S}^{d-1}(\sqrt{d}) \mapsto \mathbb{R}$ such that

$$f_{\mathbf{a}, \Theta}(\mathbf{x}) = \sum_{i=1}^N a_i \sigma(\langle \theta_i, \mathbf{x} \rangle / \sqrt{d}). \quad (1)$$

where $\sigma(\cdot)$ is the AF that is the target of our study and $\langle x, y \rangle$ denotes the inner product between vectors x and y . When clear from the context, we write $f_{\mathbf{a}, \Theta}$ as f , omitting the model's parameters. The optimal weights \mathbf{a}^* are learnt using ridge regression with regularization parameter $\lambda \geq 0$, namely,

$$\mathbf{a}^* = \mathbf{a}^*(\lambda, \mathcal{D}) = \arg \min_{\mathbf{a} \in \mathbb{R}^N} \left\{ \frac{1}{n} \sum_{j=1}^n \left(y_j - \sum_{i=1}^N a_i \sigma(\langle \theta_i, \mathbf{x}_j \rangle / \sqrt{d}) \right)^2 + \frac{N\lambda}{d} \|\mathbf{a}\|^2 \right\}. \quad (2)$$

We will tackle this question: *What is the simplest σ that leads to the best approximation of f_d ?*

We quantify the simplicity of an AF σ with its norm in different functional spaces. Namely, either

$$\|\sigma\|_1 \triangleq \mathbb{E}(|\sigma'(Z)|), \quad \text{or} \quad (3) \quad \|\sigma\|_2 \triangleq \sqrt{\mathbb{E}((\sigma'(Z))^2)}, \quad (4)$$

where σ' is the derivative of σ and the expectations are with respect to a normal random variable Z with zero mean and unit variance, i.e. $Z \sim \mathcal{N}(0, 1)$. We quantify the quality with which $f = f_{\mathbf{a}^*, \Theta}$ approximates f_d via O , a linear combination of the mean squared error and the sensitivity of f to perturbations in its input (cf. [13]). For $\alpha \in [0, 1]$ and \mathbf{x} uniform on $\mathbb{S}^{d-1}(\sqrt{d})$, we define

$$O \triangleq (1-\alpha)\mathcal{E} + \alpha\mathcal{S}, \quad (5) \quad \text{where } \mathcal{E} \triangleq \mathbb{E}((f(\mathbf{x}) - f_d(\mathbf{x}))^2), \quad (6) \quad \text{and } \mathcal{S} \triangleq \mathbb{E}(\|\nabla_{\mathbf{x}} f(\mathbf{x})\|^2). \quad (7)$$

Like in [6, 14], we operate in an asymptotic setting where $n, d, N \rightarrow \infty$, and have constant ratios between them, namely, $N/d \rightarrow \psi_1$ and $n/d \rightarrow \psi_2$. In this asymptotic setting, it does not matter if in defining (6) and (7), in addition to taking the expectation with respect to the test data \mathbf{x} , independently of \mathcal{D} , we also take expectations over \mathcal{D} and the random features in RFR. This is because when $n, d, N \rightarrow \infty$ with the ratios defined above, \mathcal{E} and \mathcal{S} will concentrate around their means [6, 14].

Mathematically, denoting by $\|\sigma\|$ either (4) or (3), our goal is to study the solutions of the problem

$$\min_{\sigma^*} \|\sigma^*\| \text{ subject to } \sigma^* \in \arg \min_{\sigma} O(\sigma). \quad (8)$$

Notice that the outer optimization only affects the selection of optimal AF in so far as the inner optimization does not uniquely define σ^* , which, as we will later see, it does not.

2 Related work

First attempts to optimize AFs include [15], [16], and [17], where genetic and evolutionary algorithms were used to learn how to numerically combine different AFs from a library into the same network. More recently, [9] used reinforcement learning to empirically discover AFs that minimize test accuracy. Their search was done over AFs that were a combination of basic units. This work produced the *Swish* AF. Similarly, [18] defined AFs as the weighted sum of a pre-defined basis and searched for optimal weights via training. The authors in [10] provided a theoretical foundation to simultaneously learn a neural network's weights and continuous piecewise-linear AFs. They showed that learning in their framework is compatible with learning in current existing deep-ReLU, parametric ReLU, APL (adaptive piecewise-linear) and MaxOut architectures. The work [19] parameterized continuous piece-wise linear AFs and numerically learnt their parameters to improve both accuracy

and robustness to adversarial perturbations. They numerically compared the performance of their SPLASH framework with that of using ReLUs, leaky-ReLUs [20], PReLU [21], tanh units, sigmoid units, ELUs [22], maxout units [23], Swish units, and APL units [24]. Similarly, [25] parametrized AFs as piece-wise linear units and learnt the AFs parameters to optimize solving different tasks. [26] proposed an empirical method to learn variations of ReLUs. [27] studied two-layer neural networks and gave a condition on the Lipschitz constant of a polynomial AF for the network to perfectly fit data. They related this condition to the model’s parameter-size and robustness. They numerically investigated the relationship between the number of ReLUs in the model and its robustness.

Several papers proposed new AFs and empirically studied their performance without systematically tuning them. In [12] the authors identified ReLU and Swish as naturally arising components of a statistical mechanics inspired model for a neural network. The authors in [28] introduced a “tent”-shaped AF that improves robustness without adversarial training, while not hurting the accuracy on non-adversarial examples. [29] proposed a new AF called SRS that can overcome the non-zero mean, negative missing, and unbounded output in ReLUs. Their work was purely empirical. [30] developed the Squared Natural Law AF. [31] proposed the Piece-wise Linear Unit AF.

The RFR model was introduced by [32] as a way to project input data into a low dimensional random features space and it has since then been studied considerably. A great part of the literature has drawn connections between the expressive power of neural networks and that of the RFR model, often via the study of Gaussian processes. For example, the work [33] did this in the context of shallow but infinitely wide neural network and the works [34, 35, 36, 37] did this for deep networks. Other works [38, 39] connected the RFR model to training a neural network with gradient descent.

In addition to [6, 14], already discussed, other papers studied the approximation properties of the RFR model. The authors in [40] studied both the RFR model and the neural tangent kernel model and provided conditions under which these models can fit polynomials in the raw features up to a maximum degree. These conditions were provided under two regimes, when $n \rightarrow \infty$ and N, d large but finite, or when $N \rightarrow \infty$ and n, d large but finite. Their results hold under weak assumptions on the AFs. The work [8] used the RFR model to compute how robust the test error is to distribution shifts between training and test data. This was done in a high-dimensional asymptotic limit when random features and training data are normal distributed. The derivations hold for a generic AF that satisfies some mild assumptions similar to the assumptions in this paper. The recent work of [7] characterized the role of overparametrization on the adversarial robustness for the RFR under an asymptotic regime when learning a linear function using a RFR model with normal-distributed random weights and normal samples. In their work, the AF was fixed to a shifted ReLU.

3 Background on the asymptotic properties of the RFR model

We review recently derived closed-form expressions for the asymptotic (model’s dimensions to infinity) mean squared error and sensitivity of the RFR model, which are the starting point of our work. We first need the following assumptions, which we assume hold in the theorems in this section.

Assumption 1. *We assume that the AF σ is weakly differentiable with weak derivative σ' , it satisfies $|\sigma(u)|, |\sigma'(u)| \leq c_0 e^{c_1|u|} \forall u \in \mathbb{R}$ for some constants $0 < c_0, c_1 < \infty$, and that it also satisfies*

$$\mu_0 = \mathbb{E}\{\sigma(Z)\}, \quad \mu_1 = \mathbb{E}\{Z\sigma(Z)\}, \quad \mu_2 = \mathbb{E}\{\sigma(Z)^2\}, \quad \mu_\star^2 = \mu_2 - \mu_0^2 - \mu_1^2, \quad \zeta = \mu_1/\mu_\star, \quad (9)$$

for some $\mu_0, \mu_1, \mu_2 \in \mathbb{R}$, where the expectations are with respect to $Z \sim \mathcal{N}(0, 1)$.

Assumption 2. *We assume that $N = N(d)$ and $n = n(d)$ such that the following limits exist in $(0, \infty)$: $\lim_{d \rightarrow \infty} N(d)/d = \psi_1$ and $\lim_{d \rightarrow \infty} n(d)/d = \psi_2$.*

Assumption 3. *We assume that $y_i = f_d(\mathbf{x}_i) + \epsilon_i$, where $\{\epsilon_i\}_{i \leq n} \sim i.i.d. \mathbb{P}_\epsilon$ are independent of $\{\mathbf{x}_i\}_{i \leq n}$, with $\mathbb{E}(\epsilon_1) = 0$, $\mathbb{E}(\epsilon_1^2) = \tau^2$, $\mathbb{E}(\epsilon_1^4) < \infty$, expectations with respect to $\{\epsilon_i\}$. Furthermore,*

$$f_d(\mathbf{x}) = \beta_{d,0} + \langle \beta_{d,1}, \mathbf{x} \rangle + f_d^{\text{NL}}(\mathbf{x}), \quad (10)$$

where $\beta_{d,0} \in \mathbb{R}$, $\beta_{d,1} \in \mathbb{R}^d$ are deterministic with $\lim_{d \rightarrow \infty} \beta_{d,0}^2 = F_0^2$, $\lim_{d \rightarrow \infty} \|\beta_{d,1}\|_2^2 = F_1^2 > 0$. The non-linear f_d^{NL} is a centered Gaussian process indexed by $\mathbf{x} \in \mathbb{S}^{d-1}(\sqrt{d})$, with covariance

$$\mathbb{E}_{f_d^{\text{NL}}}\{f_d^{\text{NL}}(\mathbf{x}_1)f_d^{\text{NL}}(\mathbf{x}_2)\} = \Sigma_d(\langle \mathbf{x}_1, \mathbf{x}_2 \rangle / d), \quad (11)$$

where $\Sigma_d(\cdot)$ satisfies $\mathbb{E}_{\mathbf{x} \sim \text{Unif}(\mathbb{S}^{d-1}(\sqrt{d}))}\{\Sigma_d(x_1/\sqrt{d})\} = 0$, $\mathbb{E}_{\mathbf{x} \sim \text{Unif}(\mathbb{S}^{d-1}(\sqrt{d}))}\{\Sigma_d(x_1/\sqrt{d})x_1\} = 0$, where x_1 is the 1st component of \mathbf{x} . We define the Signal to Noise Ratio (SNR) ρ by

$$\rho = F_1^2 / (F_\star^2 + \tau^2), \quad \text{where } F_\star^2 \triangleq \lim_{d \rightarrow \infty} \Sigma_d(1). \quad (12)$$

Informally, μ_\star quantifies how non-linear the AF is (cf. Lemma 4.1), ψ_1 quantifies the complexity of the RFR model relative of the dimension d , ψ_2 quantifies the amount of data used for training relative to d , τ^2 is the variance of the observation noise, F_1 is the magnitude of the linear component of our target function f_d , which is controlled by $\beta_{d,1}$, F_\star is the magnitude of the non-linear component f_d^{NL} in the target function, and ρ is the ratio between the magnitude of the linear component and the magnitude of all of the sources of randomness in the noisy function $f_d + \epsilon$. Recall that all of our results will be derived in the asymptotic regime when $d \rightarrow \infty$.

To obtain closed-form expressions in our main results, we will focus our analysis on the following three different important regimes: R_1 : *Ridgeless limit* regime, when $\lambda \rightarrow 0^+$; R_2 : *Highly overparameterized limit*, when $\psi_1 \rightarrow \infty$; R_3 : *Large sample limit*, when $\psi_2 \rightarrow \infty$. In the context of the RFR model, these regimes were introduced and discussed in [6]. For what follows we define $\bar{\lambda} \triangleq \lambda/\mu_\star^2$. Any “ $\lim_{d \rightarrow \infty} X = Y$ ” should be interpreted as X converging to Y in probability with respect to the training data \mathcal{D} , the random features Θ , and the random target function f_d as $d \rightarrow \infty$.

3.1 Asymptotic mean squared test error of the RFR model

The following theorems are a specialization of a more general theorem, Theorem 12 [6], which we include in the Appendix A for completeness.

Theorem 1 (Theorem 3 [6]). *The asymptotic test error (6) for regime R_1 equals*

$$\mathcal{E}_{R_1}^\infty \equiv \lim_{\lambda \rightarrow 0^+} \lim_{d \rightarrow \infty} \mathcal{E} = F_1^2 \mathcal{B}_{\text{rless}}(\zeta, \psi_1, \psi_2) + (\tau^2 + F_\star^2) \mathcal{V}_{\text{rless}}(\zeta, \psi_1, \psi_2) + F_\star^2, \quad (13)$$

where $\mathcal{B}_{\text{rless}}(\zeta, \psi_1, \psi_2) \equiv \mathcal{E}_{1,\text{rless}}/\mathcal{E}_{0,\text{rless}}$, $\mathcal{V}_{\text{rless}}(\zeta, \psi_1, \psi_2) \equiv \mathcal{E}_{2,\text{rless}}/\mathcal{E}_{0,\text{rless}}$, and the functions $\mathcal{E}_{0,\text{rless}}$, $\mathcal{E}_{1,\text{rless}}$ and $\mathcal{E}_{2,\text{rless}}$ are defined using $\psi \equiv \min\{\psi_1, \psi_2\}$ as follows,

$$\begin{aligned} \mathcal{E}_{0,\text{rless}}(\zeta, \psi_1, \psi_2) &\equiv -\chi^5 \zeta^6 + 3\chi^4 \zeta^4 + (\psi_1 \psi_2 - \psi_2 - \psi_1 + 1)\chi^3 \zeta^6 - 2\chi^3 \zeta^4 - 3\chi^3 \zeta^2 \\ &+ (\psi_1 + \psi_2 - 3\psi_1 \psi_2 + 1)\chi^2 \zeta^4 + 2\chi^2 \zeta^2 + \chi^2 + 3\psi_1 \psi_2 \chi \zeta^2 - \psi_1 \psi_2, \\ \mathcal{E}_{1,\text{rless}}(\zeta, \psi_1, \psi_2) &\equiv \psi_2 \chi^3 \zeta^4 - \psi_2 \chi^2 \zeta^2 + \psi_1 \psi_2 \chi \zeta^2 - \psi_1 \psi_2, \text{ and} \\ \mathcal{E}_{2,\text{rless}}(\zeta, \psi_1, \psi_2) &\equiv \chi^5 \zeta^6 - 3\chi^4 \zeta^4 + (\psi_1 - 1)\chi^3 \zeta^6 + 2\chi^3 \zeta^4 + 3\chi^3 \zeta^2 + (-\psi_1 - 1)\chi^2 \zeta^4 - 2\chi^2 \zeta^2 - \chi^2, \end{aligned}$$

$$\text{where } \chi \equiv -\left(\sqrt{(\psi \zeta^2 - \zeta^2 - 1)^2 + 4\zeta^2 \psi} + \psi \zeta^2 - \zeta^2 - 1 \right) / (2\zeta^2). \quad (14)$$

Remark 1. As a function of ψ_1 , $\mathcal{E}_{R_1}^\infty$ has a discontinuity at $\psi_1 = \psi_2$ called the *interpolation threshold*. For ψ_2 high enough, and for $\psi_1 < \psi_2$, $\mathcal{E}_{R_1}^\infty$ decreases, reaches a minimum and then explodes approaching ψ_2 . However, past ψ_2 , $\mathcal{E}_{R_1}^\infty$ decreases again with ψ_1 . This *double descent* behavior has been observed in many settings and is the target of many studies, including [6] and references therein.

Theorem 2 (Theorem 4 [6]). *The asymptotic test error (6) for regime R_2 equals*

$$\mathcal{E}_{R_2}^\infty \equiv \lim_{\psi_1 \rightarrow \infty} \lim_{d \rightarrow \infty} \mathcal{E} = F_1^2 \mathcal{B}_{\text{wide}}(\zeta, \psi_2, \bar{\lambda}) + (\tau^2 + F_\star^2) \mathcal{V}_{\text{wide}}(\zeta, \psi_2, \bar{\lambda}) + F_\star^2, \quad (15)$$

where $\mathcal{B}_{\text{wide}}(\zeta, \psi_2, \bar{\lambda}) \equiv (\psi_2 \omega_2 - \psi_2) / ((\psi_2 - 1)\omega_2^3 + (1 - 3\psi_2)\omega_2^2 + 3\psi_2 \omega_2 - \psi_2)$, $\mathcal{V}_{\text{wide}}(\zeta, \psi_2, \bar{\lambda}) \equiv (\omega_2^3 - \omega_2^2) / ((\psi_2 - 1)\omega_2^3 + (1 - 3\psi_2)\omega_2^2 + 3\psi_2 \omega_2 - \psi_2)$ and

$$\omega_2 \equiv -\left(\sqrt{(\psi_2 \zeta^2 - \zeta^2 - \bar{\lambda} \psi_2 - 1)^2 + 4\psi_2 \zeta^2 (\bar{\lambda} \psi_2 + 1)} + \psi_2 \zeta^2 - \zeta^2 - \bar{\lambda} \psi_2 - 1 \right) / (2(\bar{\lambda} \psi_2 + 1)). \quad (16)$$

Theorem 3 (Theorem 5 [6]). *The asymptotic test error (6) for regime R_3 equals*

$$\mathcal{E}_{R_3}^\infty \equiv \lim_{\psi_2 \rightarrow \infty} \lim_{d \rightarrow \infty} \mathcal{E} = F_1^2 \mathcal{B}_{\text{isamp}}(\zeta, \psi_1, \lambda/\mu_\star^2) + F_\star^2, \text{ where} \quad (17)$$

$\mathcal{B}_{\text{isamp}}(\zeta, \psi_1, \bar{\lambda}) \equiv (((\omega_1^3 - \omega_1^2)/\zeta^2) + \psi_1 \omega_1 - \psi_1) / ((\psi_1 - 1)\omega_1^3 + (1 - 3\psi_1)\omega_1^2 + 3\psi_1 \omega_1 - \psi_1)$, and

$$\omega_1 \equiv -\left(\sqrt{(\psi_1 \zeta^2 - \zeta^2 - \bar{\lambda} \psi_1 - 1)^2 + 4\psi_1 \zeta^2 (\bar{\lambda} \psi_1 + 1)} + \psi_1 \zeta^2 - \zeta^2 - \bar{\lambda} \psi_1 - 1 \right) / (2(\bar{\lambda} \psi_1 + 1)). \quad (18)$$

3.2 Asymptotic sensitivity of the RFR model

We derive a sensitivity formula for regimes R_1, R_2, R_3 . Our theorems are a specialization (proofs in Appendix D) of the more general Theorem 13 that we include in the Appendix A for completeness.

Theorem 4. *The sensitivity (7) for regime R_1 equals*

$$\mathcal{S}_{R_1}^\infty \equiv \lim_{\lambda \rightarrow 0^+} \lim_{d \rightarrow \infty} \mathcal{S} = \frac{\mu_1^2}{\mu_\star^2} \left(\frac{F_1^2 \mathcal{D}_{1,\text{rless}}(\zeta, \psi_1, \psi_2)}{(\chi \zeta^2 - 1) \mathcal{D}_{0,\text{rless}}(\zeta, \psi_1, \psi_2)} + \frac{(F_\star^2 + \tau^2) \mathcal{D}_{2,\text{rless}}(\zeta, \psi_1, \psi_2)}{\mathcal{D}_{0,\text{rless}}(\zeta, \psi_1, \psi_2)} \right), \quad (19)$$

$$\text{where } \mathcal{D}_{0,\text{rless}}(\zeta, \psi_1, \psi_2) = \chi^5 \zeta^6 - 3\chi^4 \zeta^4 + (\psi_1 + \psi_2 - \psi_1 \psi_2 - 1) \chi^3 \zeta^6 + 2\chi^3 \zeta^4 + 3\chi^3 \zeta^2 \quad (20)$$

$$+ (3\psi_1 \psi_2 - \psi_2 - \psi_1 - 1) \chi^2 \zeta^4 - 2\chi^2 \zeta^2 - \chi^2 - 3\psi_1 \psi_2 \chi \zeta^2 + \psi_1 \psi_2,$$

$$\mathcal{D}_{1,\text{rless}}(\zeta, \psi_1, \psi_2) = \chi^6 \zeta^6 - 2\chi^5 \zeta^4 - (\psi_1 \psi_2 - \psi_1 - \psi_2 + 1) \chi^4 \zeta^6 + \chi^4 \zeta^4 \quad (21)$$

$$+ \chi^4 \zeta^2 - 2(1 - \psi_1 \psi_2) \chi^3 \zeta^4 - (\psi_1 + \psi_2 + \psi_1 \psi_2 + 1) \chi^2 \zeta^2 - \chi^2,$$

$$\mathcal{D}_{2,\text{rless}}(\zeta, \psi_1, \psi_2) = -(\psi_1 - 1) \chi^3 \zeta^4 - \chi^3 \zeta^2 + (\psi_1 + 1) \chi^2 \zeta^2 + \chi^2, \quad (22)$$

χ is defined as in (14), and $\psi \equiv \min\{\psi_1, \psi_2\}$.

Theorem 5. *Let ω_2 equal (16). The sensitivity (7) for regime R_2 equals*

$$\mathcal{S}_{R_2}^\infty \equiv \lim_{\psi_1 \rightarrow \infty} \lim_{d \rightarrow \infty} \mathcal{S} = \frac{\omega_2^2 ((F_\star^2 + \tau^2)(-1 + \omega_2) + F_1^2(-1 - \psi_2 + \omega_2(-1 + \psi_2)))}{(-1 + \omega_2)(\psi_2 - 2\omega_2 \psi_2 + \omega_2^2(-1 + \psi_2))}. \quad (23)$$

Theorem 6. *Let ω_1 equal (18). The sensitivity (7) for regime R_3 equals*

$$\mathcal{S}_{R_3}^\infty \equiv \lim_{\psi_2 \rightarrow \infty} \lim_{d \rightarrow \infty} \mathcal{S} = F_1^2 (1 + (2/(-1 + \omega_1)) + (\psi_1 / (\psi_1 - 2\omega_1 \psi_1 + \omega_1^2(-1 + \psi_1)))). \quad (24)$$

4 Main results

We will find the simplest AFs that lead to the best trade-off between approximation accuracy and sensitivity for the RFR model. Mathematically, we will solve (8). From the theorems in Section 3 we know that \mathcal{E} and \mathcal{S} , and hence $O = (1 - \alpha)\mathcal{E} + \alpha\mathcal{S}$, only depend on the AF via μ_0, μ_1, μ_2 . Therefore, we will proceed in two steps. In Section 4.1, we will fix μ_0, μ_1, μ_2 , and find σ with associated values μ_0, μ_1, μ_2 that has minimal norm, either (4) or (3). In Section 4.2, we will find values of μ_0, μ_1, μ_2 that minimize $O = (1 - \alpha)\mathcal{E} + \alpha\mathcal{S}$. Together, these specify optimal AFs for the RFR model.

It is the case that properties of the RFR model other than the test error and sensitivity also only depend on the AF via μ_0, μ_1, μ_2 . One example is the robustness of the RFR model to disparities between the training and test data distribution [8]. Although we do not focus on these other properties, the results in Section (4.1) can be used to generate optimal AFs for them as well, as long as, similar to in Section 4.2, we can obtain μ_0, μ_1, μ_2 that optimize these other properties.

We made the decision to, as often as possible, simplify expressions by manipulating them to expose the signal to noise ratio $\rho = F_1^2 / (\tau^2 + F_\star^2)$, $F_1 > 0$, rather than using the variables F_1, τ , and F_\star . The only downside is that conclusions in the regime $\tau = F_\star = 0$ require a bit more of effort to be extracted, often been readable in the limit $\rho \rightarrow \infty$.

The complete proofs of our main results can be found in Appendix D and their main ideas below.

4.1 Optimal activation functions given fixed μ_0, μ_1 , and μ_2

Since one of our goals is knowing when an optimal AF is linear we start with the following lemma.

Lemma 4.1. *The AF σ is linear (almost surely) if and only if $\mu_\star^2 \triangleq \mu_2 - \mu_1^2 - \mu_0^2 = 0$.*

We now state results for the norms (4) and (3). The problem we will solve under both norms is similar. Let $Z \sim \mathcal{N}(0, 1)$. We consider solving the following functional problem, where $i = 1$ or 2 ,

$$\min_{\sigma} \|\sigma\|_i \quad \text{subject to } \mathbb{E}(\sigma(Z)) = \mu_0, \mathbb{E}(Z\sigma(Z)) = \mu_1, \mathbb{E}(\sigma(Z)^2) = \mu_2, \text{ with } Z \sim \mathcal{N}(0, 1). \quad (25)$$

If $i = 2$, we seek solutions over the Gaussian-weighted Lebesgue space of twice weak-differentiable functions that have $\mathbb{E}((\sigma(Z))^2)$ and $\mathbb{E}((\sigma'(Z))^2)$ defined and finite. If $i = 1$, we seek solutions over

the Gaussian-weighted Lebesgue space of weak-differentiable functions that have $\mathbb{E}((\sigma(Z))^2)$ and $\mathbb{E}(|\sigma'(Z)|)$ defined and finite. The derivative σ' is to be understood in a weak sense.

Since σ is a one-dimensional function, the requirement of existence of weak derivative implies that there exists a function v that is absolute continuous and that agrees with σ almost everywhere [41]. Therefore, any specific solution we propose should be understood as an equivalent class of functions that agree with v up to a set of measure zero with respect to the Gaussian measure.

Theorem 7. *The minimizers of (25) for $i = 2$, i.e. $\|\sigma\|^2 = \mathbb{E}((\sigma'(Z))^2)$, are*

$$\sigma(x) = ax^2 + bx + c, \text{ where } a = \pm\mu_\star/\sqrt{2}, b = \mu_1, \text{ and } c = \mu_0 - a. \quad (26)$$

In Theorem 7, if $\mu_\star = 0$ there is only one minimizer, a linear function. If $\mu_\star > 0$, there are exactly two minimizers, both quadratic functions. Note that both minimizers satisfy the growth constraints of Assumption 1, and hence can be used within the analysis of the RFR model. We note that quadratic AFs have been empirically studied in the past, e.g. [30].

Theorem 8. *One minimizer of (25) for $i = 1$, i.e. $\|\sigma\| = \mathbb{E}(|\sigma'(Z)|)$, is*

$$\sigma(x) = \mu_0 + b \max\{\min\{x, -s\}, s\}, \quad (27)$$

where $b = \frac{\mu_1}{\text{erf}(s/\sqrt{2})}$, erf is the Gauss error function, and $s \in \mathbb{R}$ is the unique solution to the equation

$$\zeta^2 \triangleq \mu_1^2/\mu_\star^2 = \left(e^{s^2/2} \text{erf}^2\left(s/\sqrt{2}\right) \right) / \left(e^{s^2/2} \left(1 - \text{erf}\left(s/\sqrt{2}\right) \right) \left(s^2 + \text{erf}\left(s/\sqrt{2}\right) \right) - \sqrt{2/\pi} s \right). \quad (28)$$

if $\mu_\star \neq 0$, and $s = +\infty$ if $\mu_\star = 0$.

When $\|\sigma\| = \mathbb{E}(|\sigma'(Z)|)$, we can characterize the complete solution family to (25). These are AFs of the form $\sigma(x) = a + b \max\{s, \min\{t, x\}\}$, where a, b, s , and t are chosen such that the constraints in (25) hold. It is possible to explicitly write a and b as a function of μ_0, μ_1, s, t , and express s, t as the solution of $E(s, t) = \mu_1^2/\mu_\star^2$, where $E(\cdot, \cdot)$ has explicit form. In this case, for each μ_0, μ_1, μ_2 there are an infinite number of optimal AFs since $E(s, t) = \mu_1^2/\mu_\star^2$ has an infinite number of solutions. ReLU's are included in this family as $t \rightarrow \infty$. The involved lengthy expressions do not bring any new insights, so we state and prove only Theorem 8, which is a specialization of the general theorem to $s = -t$.

Proofs' main ideas: We give the main ideas behind the proof of Theorem 7. The proof of Theorem 8 follows similar techniques. The first-order optimality conditions imply that $-2x\sigma'(x) + 2\sigma''(x) + \lambda_1 + \lambda_2x + \lambda_3\sigma(x) = 0$, where the Lagrange multipliers λ_1, λ_2 , and λ_3 must be later chosen such that $\mathbb{E}\{\sigma(Z)\} = \mu_0, \mathbb{E}\{Z\sigma(Z)\} = \mu_1$, and $\mathbb{E}\{\sigma^2(Z)\} = \mu_2$. Using the change of variable $\sigma(x) = \tilde{\sigma}(x/\sqrt{2}) - \lambda_1/\lambda_3 - x\lambda_2/(\lambda_3 - 1)$ we obtain $-2x\tilde{\sigma}'(x) + \tilde{\sigma}''(x) + \lambda_3\tilde{\sigma}(x) = 0$ which is the *Hermite ODE*, which is well studied in physics, e.g. it appears in the study of the quantum harmonic oscillator. The cases $\lambda_3 \in \{0, 3\}$ require special treatment. Using a finite energy/norm condition we can prove that λ_3 is quantized. In particular $\lambda_3 = 4k, k = 1, 2, \dots$, which implies that $\sigma(x) = -\lambda_1/\lambda_3 - \lambda_2x/(\lambda_3 - 2) + cH_{2k}(x/\sqrt{2})$, where H_i is the i th *Hermite polynomial* and c a constant. The energy/norm is minimal when $k = 1$, which implies a quadratic AF. \square

4.2 Activation function parameters for the optimal trade-off between sensitivity and MSE

We will find AF parameters that minimize a linear combination of sensitivity and test error. We are interested in an asymptotic analytical treatment in the three regimes mentioned in Section 3. To be specific, we will compute

$$\mathcal{U}_{R_i}(\psi_1, \psi_2, \tau, \alpha, F_1, F_\star, \lambda) \equiv \arg \min_{\mu_0, \mu_1, \mu_2} (1 - \alpha)\mathcal{E}_{R_i}^\infty + \alpha\mathcal{S}_{R_i}^\infty, \text{ where } i = 1, 2, \text{ or } 3. \quad (29)$$

We are not aware of previous work explicitly studying the trade-off between \mathcal{E} and \mathcal{S} for the RFR model. For the RFR model, the work of [6] studies only the test error and [14] studies a definition of sensitivity related but different from ours. Other papers have studied trade-offs between robustness and error measures related but different than ours and for other models, e.g. [42, 43].

When $\alpha = 1$, problem (29) reduces to minimizing the sensitivity. This has a trivial solution: the AF, and hence f , must be a constant, and $\mathcal{S} = 0$. Therefore, below we focus on the case when $\alpha \in [0, 1)$.

Special notation: In Theorem 9 we use the following special notation. Given two potential choices for AF parameters, say x and y , we define $x \sqcup y$ to mean that x exists and that y might exist or not, and that $x \sqcup y = y$ if y exists and it leads to a smaller value of $(1 - \alpha)\mathcal{E} + \alpha\mathcal{S}$ than using x , and otherwise $x \sqcup y = x$. Note that $x \sqcup y$ and $y \sqcup x$ make different statements about the existence of x and y . This notation is important to interpret the results of Table 1 in Theorem 9.

Theorem 9. *Let $\alpha \in [0, 1)$, $\psi \equiv \min\{\psi_1, \psi_2\}$ and $\bar{\psi} \equiv \max\{\psi_1, \psi_2\}$. We have that*

$$\mathcal{U}_{R_1} = \{(\mu_0, \mu_1, \mu_2) : x\mu_1^2(-1 + x + \psi) = \mu_2^2(\psi + x)\}, \text{ where } x \text{ is as in Table 1.} \quad (30)$$

	$\beta_1 \leq \bar{\psi}$	$\beta_2 < \bar{\psi} < \beta_1$	$\beta_3 < \bar{\psi} \leq \beta_2$	$\bar{\psi} \leq \beta_3$
$(\alpha < \alpha_L) \wedge E_1$	x_R	$x_R \sqcup x_1$	x_R	x_R
$(\alpha < \alpha_L) \wedge E_2 \wedge (\alpha > \alpha_C)$	x_1	$x_1 \sqcup x_3$	x_1	--
$(\alpha < \alpha_L) \wedge E_2 \wedge (\alpha < \alpha_C)$	x_1	$x_1 \sqcup x_3$	x_1	--
$(\alpha > \alpha_L) \wedge E_1 \wedge (\alpha > \alpha_C)$	--	x_L	x_L	x_L
$(\alpha > \alpha_L) \wedge E_1 \wedge (\alpha < \alpha_C)$	--	x_R	x_R	x_R
$(\alpha > \alpha_L) \wedge E_2$	x_L	$x_L \sqcup x_2$	$x_L \sqcup x_2$	x_L

Table 1: The optimal AFs (30) depends on x according to this table. Cells with "--" never happen. The values of $x_1, x_2, x_3, \beta_1, \beta_2, \beta_3, \alpha_L, \alpha_C, \alpha_R, x_L, x_R$, and the events E_1 and E_2 are specified below.

In Table 1, x_1, x_2 , and x_3 are the smallest, second smallest and third smallest roots of a 4th degree polynomial $p(x)$, specified in Appendix B, in the range $(x_L, x_R) \triangleq (-\psi, \min\{0, 1 - \psi\})$, if these exists. The variables $\beta_1, \beta_2, \beta_3, \alpha_L, \alpha_C, \alpha_R$, and the conditions E_1 and E_2 are as follows.

If $\psi_1 < \psi_2$ then $E_1 = (\alpha < \alpha_R)$, $E_2 = (\alpha > \alpha_R)$,

$$\beta_1 = \min\{\psi_1 - 4, -3\psi_1\} - 8\sqrt{|1 - \psi_1|} \max\{1/\psi_1, \psi_1^{3/2}\} + 8 \max\{1/\psi_1, \psi_1^2\}, \quad (31)$$

$$\beta_2 = \psi_1(\psi_1 + 2)/(\psi_1 + 1), \quad (32) \quad \beta_3 = \psi_1 + |1 - \psi_1| \min\{\psi_1, 1/\psi_1\}, \quad (33)$$

$$\alpha_L = \psi_2/(\psi_2 + 1 + \rho^{-1}), \quad (34) \quad \alpha_C = \psi_2/(2\psi_2 + 1 - \psi_1 + \max\{0, 1 - \psi_1\} + \rho^{-1}), \quad (35)$$

$$\alpha_R = \psi_2/(3\psi_2 + 1 - 2 \min\{1 + \psi_1, 2\psi_1\} + \rho^{-1}). \quad (36)$$

If $\psi_1 > \psi_2$ and $\psi_2 \neq 1$ then $E_1 = (\psi_1 < B) \vee ((\alpha < \alpha_R) \wedge (B < \psi_1 < A))$, $E_2 = (\psi_1 > A) \vee ((\alpha > \alpha_R) \wedge (B < \psi_1 < A))$,

$$\beta_1 : r(\beta_1) = 0, \text{ where } r \text{ is a 4th degree polynomial described in Appendix B,} \quad (37)$$

$$\beta_2 = \psi_2 + (\alpha\psi_2/(\psi_2 + 1 + \rho^{-1})), \quad (38)$$

$$\beta_3 = \psi_2 + (\min\{\psi_2, 1/\psi_2\} \alpha |\psi_2 - 1|^3 / ((\psi_2 - 1)^2 + \rho^{-1}(\psi_2 + 3))), \quad (39)$$

$$\alpha_L = (2\psi_1 - \psi_2)/(2\psi_1 - \psi_2 + 1 + \rho^{-1}), \quad (40)$$

$$\alpha_C = (\rho\psi_1 - (\psi_1 - \psi_2)/|1 - \psi_2|)/(\rho(\max\{0, 1 - \psi_2\} + 2\psi_1 - \psi_2) + 1), \quad (41)$$

$$\alpha_R = \frac{2(\psi_1 - \psi_2) \max\{1, \psi_2\} - \rho(\psi_2 - 1)^2 \psi_2}{(\psi_2 - 1)^2 (\rho(2 \min\{1, \psi_2\} - 2\psi_1 + \psi_2) - \rho - 1)}, \quad (42)$$

$$A = \psi_2 + (\rho \min\{1, \psi_2\}(\psi_2 - 1)^2/2), \text{ and} \quad (43)$$

$$B = \frac{\rho(\psi_2 - 1)^2(2\psi_2 + 1 + 2 \min\{\psi_2 - 1, 0\}) + 2\psi_2 - 1 + 2\psi_2 \max\{\psi_2, 1\} - \psi_2^2}{2\rho(\psi_2 - 1)^2 + 2 \max\{1, \psi_2\}}. \quad (44)$$

If $\psi_1 > \psi_2$ and $\psi_2 = 1$ then $E_1 = \text{False}$, $E_2 = \text{True}$ and eqs. (37)-(44) hold when reading Table 1, except (42), which is not defined, and (41) which becomes $\alpha_C = -\infty$ implying $(\alpha > \alpha_C)$ is True.

Remark 2. Excluding trivial scenarios where the optimal AF is a constant, and hence f is also constant, it follows directly from (30) that the optimal AF is linear if and only if $x = x_R$. With this information and Table 1, we have all the information needed to find exactly when the optimal AF is, or is not, linear. For regime R_1 , changing α alone can change the optimal AF from linear to non-linear and vice-versa (see e.g. 3rd column of Table 1), which justifies the observation 3 in Section 1.

Remark 3. For the cases considered in Table 1, x is unique. When $\alpha \in \{\alpha_R, \alpha_L, \alpha_C\}$, or when $(\psi_1 > \psi_2) \wedge (\psi_1 \in \{A, B\})$, we can lose the uniqueness of x . In these circumstances, we can still explicitly characterize the set of optimal x and the associated set of optimal AFs parameters, but for simplicity we omit these cases from Theorem 9.

Remark 4. Theorem 9's proof gives relationships among Table 1's constants that imply that (1) no two rows/columns simultaneously hold and (2) in some cases some cells might not hold. See App. B.

We do not consider $\psi_1 = \psi_2$ in Theorem 9 because it implies $(1-\alpha)\mathcal{E}_{R_1}^\infty + \alpha\mathcal{S}_{R_1}^\infty$ is not defined. Note that $\psi_1 = \psi_2$ has been called the *interpolation threshold* in the context of studying the generalization properties of the RFR model under vanishing regularization [6]. See Remark 1.

The set (30) of optimal AFs parameters is invariant under scaling and shifting of σ . In particular, let $\mu(\sigma)$ be μ_0, μ_1, μ_2 associated with σ . If $\mu(\sigma) \in \mathcal{U}_{R_1}$ then $\mu(a\sigma + b) \in \mathcal{U}_{R_1}$.

Also, when $x \in \{x_L, x_R\}$ we can compute the optimal value of the objective explicitly. For example, if $\psi_1 < \psi_2$ and $x = x_L$ the optimal value of the objective is $\frac{(1-\alpha)(\psi_2(F_1^2 + F_\star^2) + \psi_1\tau^2)}{\psi_2 - \psi_1}$. If $\psi_1 > \psi_2$ and $x = x_R$ the optimal value of the objective is $\frac{\alpha F_1^2(\psi_1 - \psi_2) + F_\star^2((\alpha-1)\psi_2 - \alpha) + \alpha(\psi_1 - 1)\tau^2 - \psi_1\tau^2}{\psi_1 - \psi_2}$ if $\psi_1 < 1$ and $\frac{F_1^2(\alpha(2\psi_1 - 1)(\psi_1 - \psi_2) + (\psi_1 - 1)\psi_2) + F_\star^2((\alpha-1)\psi_2 - \alpha\psi_1) - \psi_1\tau^2}{\psi_1 - \psi_2}$ if $\psi_1 \geq 1$. This follows from direct substitution.

Theorem 10. Let $\alpha \in [0, 1)$. We have that,

$$\mathcal{U}_{R_2} = \{(\mu_0, \mu_1, \mu_2) : \mu_1^2(-1 + 2\psi_2 - x)(-1 + x) = -2(\mu_\star^2 + \lambda\psi_2)(1 + x)\}, \quad (45)$$

where x is the unique solution to $p(x) = 0$ in the range $x \in (-1, \min\{1, -1 + 2\psi_2\})$, where $p(x) \triangleq p_0 + p_1x + p_2x^2 + p_3x^3 + p_4x^4$ with coefficients,

$$p_0 = 8\psi_2\rho^{-1} + (\alpha + 4\psi_2(-1 + 2\psi_2)(-1 + 2\alpha)), \quad (46) \quad p_1 = 8\psi_2\rho^{-1} + 4((1 - 4\psi_2)\alpha + 2\psi_2^2), \quad (47)$$

$$p_2 = -2(-3\alpha + \psi_2(2 + 4\alpha)), \quad (48) \quad p_3 = 4\alpha, \quad (49) \quad \text{and} \quad p_4 = \alpha. \quad (50)$$

Remark 5. The only way to get $\mu_\star = 0$, and hence a linear optimal AF is if x simultaneously satisfies $\mu_1^2(-1 + 2\psi_2 - x)(-1 + x) = -2\lambda\psi_2(1 + x)$ and $p(x) = 0$. Since the first equation does not depend on α , but the zeros of $p(x) = 0$ change continuously with α , only very special choices of parameters lead to linear AFs, and in general the overparameterized regime R_2 does not have optimal linear AFs.

Theorem 11. Let $\alpha \in [0, 1)$. We have that

$$\mathcal{U}_{R_3} = \begin{cases} \{(\mu_0, \mu_1, \mu_2) : \mu_\star = 0 \wedge \mu_1 = \infty\}, & \text{if } \alpha = 0 \vee (\psi_1 = 1 \wedge 0 < \alpha \leq \frac{1}{4}) \\ \{(\mu_0, \mu_1, \mu_2) : \mu_\star = 0 \wedge \mu_1^2 = \frac{-4\alpha^2\lambda + 3\alpha\lambda + \sqrt{\alpha}\lambda}{16\alpha^2 - 8\alpha + 1}\}, & \text{if } \psi_1 = 1 \wedge \alpha > \frac{1}{4} \\ \{(\mu_0, \mu_1, \mu_2) : \mu_\star = 0 \wedge \mu_1^2(-1 + 2\psi_1 - x)(-1 + x) + 2\lambda\psi_1(1 + x) = 0\}, & \text{if } \psi_1 \neq 1 \end{cases} \quad (51)$$

where x is the unique solution to $p(x) = 0$ in the range $x \in (-1, \min\{1, -1 + 2\psi_2\})$, where $p(x)$ is define like in Theorem 10 but with $\rho \rightarrow \infty$ and with ψ_2 replaced by ψ_1 .

Remark 6. The optimal AF is always linear and independent of the noise variables F_\star and τ .

Remark 7. When $\psi_1 = 1, \alpha \leq \frac{1}{4}$ there is no optimal AF inside our AF search space since no AF can satisfy $\mu_1 = \infty$. Rather, there exists a sequence of valid AFs with decreasing O whose $\mu_1 \rightarrow \infty$.

Remark 8. We can compute the optimal objective in closed-form in some scenarios. When $\alpha = 0$ the optimal objective is F_\star^2 . When $\psi_1 = 1 \wedge 0 < \alpha \leq \frac{1}{4}$, the optimal objective approaches $\alpha F_1^2 + (1-\alpha)F_\star^2$ as $\mu_1 \rightarrow \infty$. When $\psi_1 = 1 \wedge \alpha > \frac{1}{4}$, the optimal objective is $F_1^2(4\sqrt{\alpha} - 1 - 3\alpha) + F_\star^2(1 - \alpha)$.

Proofs' main ideas: We give the main ideas behind the proof of Theorem 10. The proof of Theorems 9 and 11 follows similar techniques but require more care. The objective O only depends on AF parameters via $\omega_2 = \omega_2(\psi_2, \mu_1^2, \mu_\star^2)$. We use the Möbius transformation $x = (1 + \omega_2)/(\omega_2 - 1)$ such that the infinite range $\omega_2 \in [-\infty, 0]$ gets mapped to the finite range $x \in [-1, 1]$. We then focus the rest of the proof on minimizing $O = O(x)$ over the range of x . First we show that given that $\mu_1^2, \mu_\star^2 \geq 0, \psi_1 > 0$, the range of x can be reduced to $x \in [x_L, x_R] \triangleq [-1, \min\{1, -1 + 2\psi_2\}]$. Then we compute dO/dx and d^2O/dx^2 , which turns out to be rational functions of x . We then show that if $x \in [-1, \min\{1, -1 + 2\psi_2\}]$ then $d^2O/dx^2 > 0$, so O is strictly convex. We also show that $dO/dx < 0$ at x_L and $dO/dx > 0$ at x_R , thus x_R and x_L cannot be minimizers. Finally, we show that the zeros of the numerator $p(x)$ of the rational function dO/dx differ from the denominator's zeros. So the optimal x is the unique solution to $p(x)$ in $[x_L, x_R]$. \square

4.3 Important observations

Together, Sections 4.1 and 4.2 explicitly and fully characterize the solutions of (8) in the ridgeless, overparametrized, and large sample regimes. A few important observations follow from our theory. In Appendix C we discuss more on this topic and include details on the observations below.

Observation 1: In regime R_1 , and if $\alpha = 0, \psi_1 < \psi_2$, the optimal AF is linear. This follows from Theorem 9 and Remark 2. Indeed, expressions simplify and we get that $p(x) = \rho\psi_2(\psi_1 - (x + \psi_1)^2)^2$ if $\psi_1 \neq 1$ or $p(x) = -(2 + x)^2\rho\psi_2$ if $\psi_1 = 1$, which implies that x_1 does not exist (since it would be outside of $(-x_L, x_R)$). Hence, the first row of Table 1 always gives $x = x_R$ and the optimal AF is linear. Also, when $\alpha = 0, \psi_1 < \psi_2$, we can explicitly compute the optimal objective (see paragraph before Theorem 10). If furthermore $\tau = F_\star = 0$, we can show that also when $\psi_1 > \psi_2$, x_1 does not exist and $x = x_R$, therefore the optimal objective and optimal AF when $\psi_1 > \psi_2$ have the same formula as when $\psi_1 < \psi_2$. Hence, if $\alpha = \tau = F_\star = 0, \psi_1 < \psi_2$, from the formula one can conclude that choosing an optimal linear AF destroys the double descent curve if $\psi_2 > 1^1$, the test error becoming exactly zero for $\psi_1 \geq 1$. This contrasts with choosing a non-linear, sub-optimal, AF which will exhibit a double descent curve. This justifies observation 1 (low complexity $\psi_1 < \psi_2$) and observation 2 in Section 1. Figure 1-(A,B) illustrates this and details the high-complexity ($\psi_1 > \psi_2$) observation.




Figure 1: (A) Consider the regime R_1 . In a noiseless setting, if $\psi_2 > 1$, the evolution of O versus ψ_1 , when an optimal linear AF σ^\star is used, can achieve 0 test error for $\psi_1 \geq 1$. However, if a non-linear σ_{ReLU} is used, we observe the typical double descent curve. (B) Consider the regime R_1 . If there is observation noise $\tau > 0$, the evolution of O versus ψ_1 with a linear AF σ_{linear} is only optimal for $\psi_1 < \psi_2$. For $\psi_1 > \psi_2$, O is optimal for a linear AF until $\psi_1 < C$ ($C = 5$ for the parameters here). For $\psi_1 > C$ a non-linear AF σ_\star , here close to but different from a ReLU, achieves minimal O . (C) Consider the regime R_2 . When a ReLU is used (green curves), the evolution of O versus λ for both low and high Signal to Noise Ratio (SNR) ρ is only optimal for a special choice of λ , achieving the minimum $O_{\sigma_{\text{ReLU}}}(\lambda^\star)$. However, also for the same low and high SNR settings, when an optimal (non-linear) AF is used (orange curves), we obtain the same, or slightly better, O_{σ^\star} regardless of any careful choice for λ . For low SNR ($\tau^2 = 10$) we have $O_{\sigma^\star} = O_{\sigma_{\text{ReLU}}}(\lambda^\star) = 0.512$ and for high SNR ($\tau^2 = 5$) we get $O_{\sigma_{\text{ReLU}}}(\lambda^\star) = 0.0220 > O_{\sigma^\star} = 0.0217$. (D) In a situation just like in (C) but with even higher SNR, the difference between the minimum O that can be achieved with a particular choice of λ (blue line ordinate value $O_{\sigma_{\text{ReLU}}}(\lambda^\star)$) and the value of O with any choice of λ but with an optimal (non-linear) AF (orange line ordinate value O_{σ^\star}) becomes clearly visible. We include inside of each plot the parameters used.

Observation 2: In regime R_2 , looking at Theorem 2 and Theorem 5, one sees that both \mathcal{E} and \mathcal{S} , and hence the objective O (cf. (5)), only depend on the optimal AF parameters via ω_2 . In particular, we can solve (29) by searching for the ω_2 that achieves the smallest objective. Given the definition of $\omega_2 = \omega_2(\psi_2, \zeta^2, \mu_\star, \lambda)$ in (16), fixing λ and changing ζ or μ_\star always allows one to span a larger range of values for ω_2 than fixing the AF's parameters ζ, μ_\star and changing λ . In particular, a tedious calculation shows that in the first case the achievable range for ω_2 is $[\frac{\psi_2}{\min\{0, -1+\psi_2\}}, 0]$ which contains the range in the second case which is $[\frac{1}{2}(\zeta^2(-\psi_2) - \sqrt{\zeta^2(\psi_2(\zeta^2(\psi_2-2)+2) + \zeta^2+2) + 1 + \zeta^2+1}), 0]$. This implies that while for a fixed AF one needs to tune λ during learning for best performance, if an optimal AF is used, regardless of λ , we always achieve either equal or better performance. This justifies the observation 4 made in Section 1. This is illustrated in Figure 1-(C,D).

5 Conclusion and future work

We solved our task of finding optimal Activation Functions (AFs) for the Random Features Regression (RFR) model and characterized when these are linear, or non-linear. Our theory yielded interesting connections between the best AF to use and the regime in which the RFR model operates: e.g. in

¹If $\psi_2 < 1$ the optimal AF is still linear but the explosion at the interpolation threshold $\psi_1 = \psi_2$ remains.

some regimes using a linear AF is optimal; in other regimes we can avoid hyperparameter tuning by choosing an optimal non-linear AF.

We reduced the gap between the practice and theory of AFs’ design, but parts remain to be closed. A limitation of our work is that to obtain explicit formulas we restricted ourselves to three important regimes only. Although a general and complete explicit treatment is unlikely, we have a plan to extract in future work a few properties of the optimal AF in a general setting, such as determining when the optimal AF will, or will not, be linear. Another limitation is that we could only obtain explicit equations for optimal AFs under two functional norms in the optimization problem from which we extract them. We will explore other norms in the future. One research direction for which we have concrete plans is focusing on the two norms used and adding higher order moment restrictions to the AF. Some of these higher order constraints appear in the theoretical analysis of neural models [40].

In future work we plan to extend our theory to extract AFs for the RFR model that optimize a combination of test error and robustness to test/train distribution shifts and adversarial attacks. The starting point will be [8] and [7] discussed in Section 2, the results of [7] needing to be generalized from a ReLU to general AFs before we can optimize the AFs’ parameters.

References

- [1] Mikhail Belkin, Daniel Hsu, and Ji Xu. Two models of double descent for weak features. *SIAM Journal on Mathematics of Data Science*, 2(4):1167–1180, 2020.
- [2] Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *The Annals of Statistics*, 50(2):949–986, 2022.
- [3] Mikhail Belkin, Alexander Rakhlin, and Alexandre B Tsybakov. Does data interpolation contradict statistical optimality? In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1611–1619. PMLR, 2019.
- [4] Tengyuan Liang and Alexander Rakhlin. Just interpolate: Kernel “ridgeless” regression can generalize. *The Annals of Statistics*, 48(3):1329–1347, 2020.
- [5] Arthur Jacot, Franck Gabriel, and Clément Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.
- [6] Song Mei and Andrea Montanari. The generalization error of random features regression: Precise asymptotics and the double descent curve. *Communications on Pure and Applied Mathematics*, 75(4):667–766, 2022.
- [7] Hamed Hassani and Adel Javanmard. The curse of overparametrization in adversarial training: Precise analysis of robust generalization for random features regression. *arXiv preprint arXiv:2201.05149*, 2022.
- [8] Nilesh Tripathi, Ben Adlam, and Jeffrey Pennington. Overparameterization improves robustness to covariate shift in high dimensions. In A. Beygelzimer, Y. Dauphin, P. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, 2021.
- [9] Prajit Ramachandran, Barret Zoph, and Quoc V. Le. Searching for activation functions, 2017.
- [10] Michael Unser. A representer theorem for deep neural networks, 2019.
- [11] Andriandrasana David Rasamoelina, Fouzia Adjailia, and Peter Sinčák. A review of activation function for artificial neural network. In *2020 IEEE 18th World Symposium on Applied Machine Intelligence and Informatics (SAMII)*, pages 281–286. IEEE, 2020.
- [12] Mirco Milletari, Thiparat Chotibut, and Paolo E. Trevisanutto. Mean field theory of activation functions in deep neural networks, 2019.
- [13] Roman Novak, Yasaman Bahri, Daniel A Abolafia, Jeffrey Pennington, and Jascha Sohl-Dickstein. Sensitivity and generalization in neural networks: an empirical study. *arXiv preprint arXiv:1802.08760*, 2018.

- [14] Alexander D’Amour, Katherine Heller, Dan Moldovan, Ben Adlam, Babak Alipanahi, Alex Beutel, Christina Chen, Jonathan Deaton, Jacob Eisenstein, Matthew D Hoffman, et al. Under-specification presents challenges for credibility in modern machine learning. *arXiv preprint arXiv:2011.03395*, 2020.
- [15] Riccardo Poli. *Parallel distributed genetic programming*. University of Birmingham, Cognitive Science Research Centre, 1996.
- [16] Daniel Weingaertner, Victor K Tatai, Ricardo R Gudwin, and Fernando J Von Zuben. Hierarchical evolution of heterogeneous neural networks. In *Proceedings of the 2002 Congress on Evolutionary Computation. CEC’02 (Cat. No. 02TH8600)*, volume 2, pages 1775–1780. IEEE, 2002.
- [17] Maryam Mahsal Khan, Arbab Masood Ahmad, Gul Muhammad Khan, and Julian F Miller. Fast learning neural networks using cartesian genetic programming. *Neurocomputing*, 121:274–289, 2013.
- [18] Mohit Goyal, Rajan Goyal, and Brejesh Lall. Improved polynomial neural networks with normalised activations. In *2020 International Joint Conference on Neural Networks (IJCNN)*, pages 1–8, 2020.
- [19] Mohammadamin Tavakoli, Forest Agostinelli, and Pierre Baldi. Splash: Learnable activation functions for improving accuracy and adversarial robustness. *Neural Networks*, 140:1–12, 2021.
- [20] Andrew L Maas, Awni Y Hannun, and Andrew Y Ng. Rectifier nonlinearities improve neural network acoustic models. In *Proc. icml*, volume 30, page 3, 2013.
- [21] Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Delving deep into rectifiers: Surpassing human-level performance on imagenet classification. *2015 IEEE International Conference on Computer Vision (ICCV)*, Dec 2015.
- [22] Djork-Arné Clevert, Thomas Unterthiner, and Sepp Hochreiter. Fast and accurate deep network learning by exponential linear units (elus), 2015.
- [23] Ian J Goodfellow, David Warde-Farley, Mehdi Mirza, Aaron Courville, and Yoshua Bengio. Maxout networks. *arXiv preprint arXiv:1302.4389*, 2013.
- [24] Forest Agostinelli, Matthew Hoffman, Peter Sadowski, and Pierre Baldi. Learning activation functions to improve deep neural networks, 2014.
- [25] Yucong Zhou, Zezhou Zhu, and Zhao Zhong. Learning specialized activation functions with the piecewise linear unit, 2021.
- [26] Chaity Banerjee, Tathagata Mukherjee, and Eduardo Pasiliao Jr. An empirical study on generalizations of the relu activation function. In *Proceedings of the 2019 ACM Southeast Conference*, pages 164–167, 2019.
- [27] Sébastien Bubeck, Yuanzhi Li, and Dheeraj Nagaraj. A law of robustness for two-layers neural networks, 2020.
- [28] Andras Rozsa and Terrance E. Boult. Improved adversarial robustness by reducing open space risk via tent activations. 2019.
- [29] Yuan Zhou, Dandan Li, Shuwei Huo, and Sun-Yuan Kung. Soft-root-sign activation function. *arXiv preprint arXiv:2003.00547*, 2020.
- [30] Adedamola Wuraola and Nitish Patel. Ssql: A new computationally efficient activation function. In *2018 International Joint Conference on Neural Networks (IJCNN)*, pages 1–7. IEEE, 2018.
- [31] Andrei Nicolae. Plu: The piecewise linear unit activation function. *arXiv preprint arXiv:1809.09534*, 2018.
- [32] Ali Rahimi and Benjamin Recht. Random features for large-scale kernel machines. In J. Platt, D. Koller, Y. Singer, and S. Roweis, editors, *Advances in Neural Information Processing Systems*, volume 20. Curran Associates, Inc., 2007.

- [33] Christopher Williams. Computing with infinite networks. In M. C. Mozer, M. Jordan, and T. Petsche, editors, *Advances in Neural Information Processing Systems*, volume 9. MIT Press, 1996.
- [34] Adrià Garriga-Alonso, Carl Edward Rasmussen, and Laurence Aitchison. Deep convolutional networks as shallow gaussian processes. In *International Conference on Learning Representations*, 2019.
- [35] Roman Novak, Lechao Xiao, Yasaman Bahri, Jaehoon Lee, Greg Yang, Daniel A. Abolafia, Jeffrey Pennington, and Jascha Sohl-dickstein. Bayesian deep convolutional networks with many channels are gaussian processes. In *International Conference on Learning Representations*, 2019.
- [36] Alexander G. de G. Matthews, Jiri Hron, Mark Rowland, Richard E. Turner, and Zoubin Ghahramani. Gaussian process behaviour in wide deep neural networks. In *International Conference on Learning Representations*, 2018.
- [37] Tamir Hazan and T. Jaakkola. Steps toward deep kernel methods from infinite neural networks. *ArXiv*, abs/1508.05133, 2015.
- [38] Amit Daniely, Roy Frostig, and Yoram Singer. Toward deeper understanding of neural networks: The power of initialization and a dual view on expressivity. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.
- [39] Amit Daniely. Sgd learns the conjugate kernel class of the network. In I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017.
- [40] Behrooz Ghorbani, Song Mei, Theodor Misiakiewicz, and Andrea Montanari. Linearized two-layers neural networks in high dimension. *The Annals of Statistics*, 49(2):1029–1054, 2021.
- [41] Walter Rudin et al. *Principles of mathematical analysis*, volume 3. McGraw-hill New York, 1976.
- [42] Dimitris Tsipras, Shibani Santurkar, Logan Engstrom, Alexander Turner, and Aleksander Madry. Robustness may be at odds with accuracy. *arXiv preprint arXiv:1805.12152*, 2018.
- [43] Hongyang Zhang, Yaodong Yu, Jiantao Jiao, Eric Xing, Laurent El Ghaoui, and Michael Jordan. Theoretically principled trade-off between robustness and accuracy. In *International conference on machine learning*, pages 7472–7482. PMLR, 2019.
- [44] Izrail Moiseevitch Gelfand, Richard A Silverman, et al. *Calculus of variations*. Courier Corporation, 2000.

A General theorems for the asymptotic mean squared test error and sensitivity of the RFR model

This appendix is referenced at the start of Section 3.1 and at the start of Section 3.2.

Theorem 12 (Theorem 2 [6]). *If assumptions 1, 2, 3 hold, and for any value of the regularization parameter $\lambda > 0$, the asymptotic test error (6) for the RFR satisfies*

$$\mathcal{E} \xrightarrow{P} \mathcal{E}^\infty \equiv F_1^2 \mathcal{B}(\zeta, \psi_1, \psi_2, \lambda/\mu_\star^2) + (\tau^2 + F_\star^2) \mathcal{V}(\zeta, \psi_1, \psi_2, \lambda/\mu_\star^2) + F_\star^2, \quad (52)$$

where \xrightarrow{P} denotes convergence in probability when $d \rightarrow \infty$ with respect to the training data \mathcal{D} , the random features Θ , and the random target function f_d , and where $\mathcal{B}(\zeta, \psi_1, \psi_2, \bar{\lambda}) \equiv \mathcal{E}_1(\zeta, \psi_1, \psi_2, \bar{\lambda})/\mathcal{E}_0(\zeta, \psi_1, \psi_2, \bar{\lambda})$, $\mathcal{V}(\zeta, \psi_1, \psi_2, \bar{\lambda}) \equiv \mathcal{E}_2(\zeta, \psi_1, \psi_2, \bar{\lambda})/\mathcal{E}_0(\zeta, \psi_1, \psi_2, \bar{\lambda})$, and the functions \mathcal{E}_0 , \mathcal{E}_1 , \mathcal{E}_2 are defined as follows,

$$\begin{aligned} \mathcal{E}_0(\zeta, \psi_1, \psi_2, \bar{\lambda}) &\equiv -\chi^5 \zeta^6 + 3\chi^4 \zeta^4 + (\psi_1 \psi_2 - \psi_2 - \psi_1 + 1) \chi^3 \zeta^6 - 2\chi^3 \zeta^4 - 3\chi^3 \zeta^2 \\ &\quad + (\psi_1 + \psi_2 - 3\psi_1 \psi_2 + 1) \chi^2 \zeta^4 + 2\chi^2 \zeta^2 + \chi^2 + 3\psi_1 \psi_2 \chi \zeta^2 - \psi_1 \psi_2, \\ \mathcal{E}_1(\zeta, \psi_1, \psi_2, \bar{\lambda}) &\equiv \psi_2 \chi^3 \zeta^4 - \psi_2 \chi^2 \zeta^2 + \psi_1 \psi_2 \chi \zeta^2 - \psi_1 \psi_2, \\ \mathcal{E}_2(\zeta, \psi_1, \psi_2, \bar{\lambda}) &\equiv \chi^5 \zeta^6 - 3\chi^4 \zeta^4 + (\psi_1 - 1) \chi^3 \zeta^6 + 2\chi^3 \zeta^4 + 3\chi^3 \zeta^2 + (-\psi_1 - 1) \chi^2 \zeta^4 - 2\chi^2 \zeta^2 - \chi^2, \end{aligned}$$

where $\chi(\psi_1, \psi_2, \bar{\lambda}) \equiv v_1(i(\psi_1 \psi_2 \bar{\lambda})^{1/2}) \cdot v_2(i(\psi_1 \psi_2 \bar{\lambda})^{1/2})$, and v_1 and v_2 are two functions specified in Def. 1 in [6].

Theorem 13 (Eq. (19) [14]). *If assumptions 1, 2 and 3 hold, and for any value of the regularization parameter $\bar{\lambda} \equiv \lambda/\mu_\star^2 > 0$, the asymptotic sensitivity for the RFR, namely equation (7) with $f = f_{\mathbf{a}, \Theta}$ where $f_{\mathbf{a}, \Theta}$ is defined as in equations (1) and (2), satisfies*

$$\mathcal{S} \xrightarrow{P} \mathcal{S}^\infty \equiv \mu_\star^2 \left(\frac{F_1^2}{\mu_\star^2} \cdot \frac{\mathcal{D}_1(\zeta, \psi_1, \psi_2, \bar{\lambda})}{(\chi \zeta^2 - 1) \mathcal{D}_0(\zeta, \psi_1, \psi_2, \bar{\lambda})} + \frac{\tau^2 + F_\star^2}{\mu_\star^2} \cdot \frac{\mathcal{D}_2(\zeta, \psi_1, \psi_2, \bar{\lambda})}{\mathcal{D}_0(\zeta, \psi_1, \psi_2, \bar{\lambda})} \right), \quad (53)$$

where \xrightarrow{P} denotes convergence in probability when $d \rightarrow \infty$ with respect to the training data \mathbf{X} , ε , the random features Θ , and the random target function f_d , and where

$$\begin{aligned} \mathcal{D}_0(\zeta, \psi_1, \psi_2, \bar{\lambda}) &= \chi^5 \zeta^6 - 3\chi^4 \zeta^4 + (\psi_1 + \psi_2 - \psi_1 \psi_2 - 1) \chi^3 \zeta^6 + 2\chi^3 \zeta^4 + 3\chi^3 \zeta^2 \\ &\quad + (3\psi_1 \psi_2 - \psi_2 - \psi_1 - 1) \chi^2 \zeta^4 - 2\chi^2 \zeta^2 - \chi^2 - 3\psi_1 \psi_2 \chi \zeta^2 + \psi_1 \psi_2, \end{aligned} \quad (54)$$

$$\begin{aligned} \mathcal{D}_1(\zeta, \psi_1, \psi_2, \bar{\lambda}) &= \chi^6 \zeta^6 - 2\chi^5 \zeta^4 - (\psi_1 \psi_2 - \psi_1 - \psi_2 + 1) \chi^4 \zeta^6 + \chi^4 \zeta^4 \\ &\quad + \chi^4 \zeta^2 - 2(1 - \psi_1 \psi_2) \chi^3 \zeta^4 - (\psi_1 + \psi_2 + \psi_1 \psi_2 + 1) \chi^2 \zeta^2 - \chi^2, \end{aligned} \quad (55)$$

$$\mathcal{D}_2(\zeta, \psi_1, \psi_2, \bar{\lambda}) = -(\psi_1 - 1) \chi^3 \zeta^4 - \chi^3 \zeta^2 + (\psi_1 + 1) \chi^2 \zeta^2 + \chi^2, \quad (56)$$

and where χ is defined in (14).

B Details regarding Theorem 9

B.1 Relationship among the constants in Table 1

This appendix is referenced in Remark 4.

The following relationships can be derived either from the proof of Theorem 9, or from a direct computation based on the formulas given in Theorem 9. They do not add critical information to what is proved in Theorem 9 but they give general rules to exclude some cells in Table 1.

Remark 9. Let $\psi_1 < \psi_2$. If $\psi_2 < \min\{2\psi_1, \psi_1 + 1\}$, then $\alpha_L < \alpha_C < \alpha_R$. In this case, the 4th and 5th rows of Table 1 never happen. If $\psi_2 > \min\{2\psi_1, \psi_1 + 1\}$, then $\alpha_R < \alpha_C < \alpha_L$. If $\psi_2 = \min\{2\psi_1, \psi_1 + 1\}$, then $\alpha_L = \alpha_C = \alpha_R$. We always have that $\alpha_L, \alpha_C, \alpha_R \in [0, 1]$, and $0 \leq \beta_3 < \beta_2 < \beta_1$. Since E_1 and E_2 cannot simultaneously hold, no two rows can simultaneously hold. Since $\beta_3 < \beta_2 < \beta_1$, no two columns can simultaneously hold. Remark 9 follows from Lemma D.13 used in the proof of Theorem 9.

Remark 10. Let $\psi_1 > \psi_2$. We always have that $\alpha_L \in [0, 1]$. It also holds that $\alpha_C \in [0, 1]$ if and only if $\psi_1 \leq \psi_2 / (1 - \rho|1 - \psi_2|)$. We have that $\alpha_R \in [0, 1]$ if and only if $\psi_1 \geq \psi_2 + \frac{1}{2}\rho(\psi_2 - 1)^2 \min\{\psi_2, 1\}$ and

$$\psi_1 \leq \frac{-1 + \rho(\psi_2 - 1)^2(2\psi_2 + \min\{2\psi_2 - 1, 1\}) + \psi_2(2 + \max\{\psi_2, 2 - \psi_2\})}{2(\max\{\psi_2, 1\} + \rho(\psi_2 - 1)^2)}.$$

Note that if $\psi_1 > \psi_2 = 1$ then α_R is not defined, see statement of Theorem 9. If we had attempted to use the formulas above we would have gotten that $\alpha \in [0, 1]$ if and only if $1 \leq \psi_1 \leq 1$, this last condition never being met when $\psi_1 > \psi_2 = 1$. Also, $A \leq B$ always, where A and B are given in (43) and (44). Furthermore, $B < \psi_2$ if and only if $1/\rho > \gamma \triangleq \min\{1, \max\{0, 2\psi_2 - 1\}\}$. Since $A \leq B$, it follows that E_1 and E_2 cannot simultaneously hold, and hence no two rows can simultaneously hold. We always have that $\beta_3 < \beta_2 < \beta_1$, therefore no two columns can simultaneously hold. Remark 10 follows from Lemma D.14 used in the proof of Theorem 9.

B.2 Definition of the polynomial $p(x)$ when $\psi_1 < \psi_2$

This appendix is referenced in the statement of Theorem 9 right after Table 1 is introduced.

If $\psi_1 \neq 1$ then $p(x) \triangleq p_0 + p_1x + p_2x^2 + p_3x^3 + p_4x^4 + p_5x^5$ where

$$p_0 = -(\psi_1 - 1)^2\psi_1^2(\rho(\alpha - 4\alpha\psi_1 + (3\alpha - 1)\psi_2) + \alpha), \quad (57)$$

$$p_1 = 2(\psi_1 - 1)\psi_1(\rho(\alpha(\psi_1(9\psi_1 - 6\psi_2 - 4) + \psi_2) + 2\psi_1\psi_2) - 2\alpha\psi_1), \quad (58)$$

$$p_2 = 2\psi_1(\rho(\alpha + 4\alpha\psi_1(4\psi_1 - 3) + \psi_2(4\alpha - 9\alpha\psi_1 + 3\psi_1 - 1)) - \alpha(3\psi_1 - 1)), \quad (59)$$

$$p_3 = 4\psi_1(\rho(\alpha(7\psi_1 - 2) - 3\alpha\psi_2 + \psi_2) - \alpha), \quad (60)$$

$$p_4 = \rho(\alpha(12\psi_1 - 1) - 3\alpha\psi_2 + \psi_2) - \alpha, \quad (61)$$

$$p_5 = 2\alpha\rho, \quad (62)$$

If $\psi_1 = 1$ then $p(x) \triangleq q_0 + q_1x + q_2x^2 + q_3x^3$ where

$$q_0 = \rho(-10\alpha + 10\alpha\psi_2 - 4\psi_2) + 4\alpha, \quad (63) \quad q_1 = \rho(-20\alpha + 12\alpha\psi_2 - 4\psi_2) + 4\alpha, \quad (65)$$

$$q_2 = \rho(-11\alpha + 3\alpha\psi_2 - \psi_2) + \alpha, \quad (64) \quad \text{and } q_3 = -2\alpha\rho. \quad (66)$$

B.3 Definition of the polynomial $p(x)$ when $\psi_1 > \psi_2$

This appendix is referenced in the statement of Theorem 9 right after Table 1 is introduced.

$p(x) \triangleq p_0 + p_1x + p_2x^2 + p_3x^3 + p_4x^4 + p_5x^5$, where (67)

$$p_0 = \psi_2^2(\rho(\psi_2 - 1)^2(2\alpha\psi_1 - (3\alpha + 1)\psi_2 + \alpha) + (\alpha\psi_2^2 - 2(\alpha + 1)\psi_2 + \alpha + 2\psi_1)), \quad (68)$$

$$p_1 = 2\psi_2((\psi_2(2\alpha(\psi_2 - 1) - 1) + \psi_1) - \rho(\psi_2 - 1)((7\alpha + 2)\psi_2^2 - \psi_2(4\alpha\psi_1 + 3\alpha + 1) + \psi_1)), \quad (69)$$

$$p_2 = 2\psi_2(\alpha(3\psi_2 - 1) - \rho((13\alpha + 3)\psi_2^2 - 2\psi_2(3\alpha\psi_1 + 5\alpha + 1) + 2\alpha\psi_1 + \alpha + \psi_1)), \quad (70)$$

$$p_3 = 4\psi_2(\rho(2\alpha(\psi_1 + 1) - (6\alpha + 1)\psi_2) + \alpha), \quad (71)$$

$$p_4 = \rho(2\alpha\psi_1 - (11\alpha + 1)\psi_2 + \alpha) + \alpha, \quad (72)$$

$$p_5 = -2\alpha\rho. \quad (73)$$

B.4 Definition of the polynomial $r(x)$ when $\psi_1 > \psi_2$

This appendix is referenced in the statement of Theorem 9 in equation (37).

The coefficient β_1 is such that $r(\beta_1) = 0$, where $r(x) = r_0 + r_1x + r_2x^2 + r_3x^3 + r_4x^4$ with

$$r_0 = \rho\psi_2^3(\rho\psi_2(\rho\psi_2(16\alpha\rho\psi_2 - 8\alpha(14\alpha\rho + \rho - 6) + \rho) + 8\alpha(\rho((8\alpha(2\alpha + 5) - 1)\rho - 28\alpha + 5) + 6)) + 16\alpha(\alpha\rho + \rho + 1)(-4\alpha\rho + \rho + 1)^2),$$

$$r_1 = 4\rho\psi_2^2(\rho\psi_2(2\alpha(8\alpha\rho(7 - 2(\alpha + 5)\rho) + 3(\rho - 5)\rho - 18) - \rho\psi_2(-56\alpha^2\rho + 12\alpha\rho\psi_2 - 6\alpha(\rho - 6) + \rho)) - 4\alpha(\rho + 1)((2\alpha - 3)\rho - 3)((4\alpha - 1)\rho - 1)),$$

$$r_2 = 2\rho\psi_2(-8\alpha^2\rho(\rho\psi_2(7\rho\psi_2 - 20\rho + 14) + 7(\rho + 1)^2) + 12\alpha(\rho\psi_2 + \rho + 1)(\rho\psi_2) + 2(\rho + 1)^2) + 3\rho^3\psi_2^2),$$

$$r_3 = -8\alpha\rho(\rho\psi_2 + \rho + 1)(\rho\psi_2(2\rho\psi_2 - 3\rho + 4) + 2(\rho + 1)^2) - 4\rho^4\psi_2^2, \text{ and}$$

$$r_4 = \rho^4\psi_2.$$

C Important observations

This appendix is referenced in Section 4.3.

C.1 Reproducing Figure 1

The plots in Figure 1 come directly from our theory. They involve no experiments and can be obtained using many standard mathematical computing software tools. Nonetheless, since it does require some effort to code our equations, we include code to generate Figure 1 in the following anonymous Google Drive folder: https://drive.google.com/drive/folders/1JtV6vo_N2jHg9IF1ITzqqP81Y7PKkcvy?usp=sharing. To generate the plots in Figure 1, run the file named `RunMeToGenerateFigure_1.nb`. It runs using Wolfram Mathematica V12. We ran it in abou using a MacBook Pro with 2.6 GHz 6-Core Intel Core i7 and 32 GB 2667 MHz DDR4. In this machine it takes about 1 second to run.

C.2 Some details about observation 1 in Section 4.3

When $\alpha = 0$ and $\psi_1 < \psi_2$, and using Theorem 9, we can explicitly compute the objective $O = \mathcal{E}_{R_1} = (F_\star^2\psi_2 + F_1^2 \max\{1 - \psi_1, 0\}\psi_2 + \psi_1\tau^2)/(\psi_2 - \psi_1)$ if $\psi_1 < \psi_2$. This follows from making $\alpha = 0$ in the expressions in the paragraph before Theorem 10. Observe that, as function of ψ_1 , the test error \mathcal{E}_{R_1} achieves a minimum of $(F_\star^2\psi_2 + \tau^2)/(-1 + \psi_2)$ at $\psi_1 = 1$ if $\psi_2 > 1 + 1/\rho$ and a minimum of $(F_1^2\psi_2 + F_\star^2\psi_2)/\psi_2$ at $\psi_1 = 0$ otherwise. In particular, if $\tau = F_\star = 0^+$, then the condition $\psi_2 > 1 + 1/\rho$ becomes $\psi_2 > 1$.

When $\alpha = 0$ and $\psi_1 > \psi_2$ the polynomial that determines x_1 becomes

$$p(x) = \psi_2(-4\psi_2x^3 - 2(\psi_1 + \psi_2)(3\psi_2 - 2))x^2 - x^4 - 2(\psi_2 - 1)(\psi_1 + \psi_2)(2\psi_2 - 1)x - (\psi_2 - 1)^2\psi_2^2. \quad (74)$$

One can show that in the range $\psi_1 > \psi_2$ and $x \in (x_L, x_R) = (-\psi_2, \min\{0, 1 - \psi_2\})$, we have $p(x) < 0$ and hence it has no roots. Therefore, x_1 does not exist. Therefore, if furthermore we have $\tau = F_\star = 0$, then condition E_2 is never true, so x must be read from the first row in Table 12, which implies that $x = x_R$, and the optimal AF is linear also when $\psi_1 > \psi_2$.

D Proofs

This appendix is referenced in Section 4.

D.1 Proof of sensitivity properties of Section 3.2

Proof of Theorem 4. The proof follows directly from Theorem 13 by taking the limit when $\lambda \rightarrow +0$. \square

Proof of Theorem 5. The proof follows directly from Theorem 13 by taking the limit when $\psi_1 \rightarrow +\infty$. \square

Proof of Theorem 6. The proof follows directly from Theorem 13 by taking the limit when $\psi_2 \rightarrow +\infty$. \square

D.2 Proof of necessary and sufficient condition for linearity of Section 4.1

Proof of Lemma 4.1. If $\sigma(x)$ is linear function, a direct calculation shows that $\mu_\star = 0$. On the other hand, since $0 \leq \mathbb{E}(\sigma(Z) - \mu_0 - \mu_1 Z)^2 = \mu_\star^2$, we have that $\mu_\star = 0$ implies that $\sigma(Z) = \mu_0 + \mu_1 Z$ almost surely. Since the support of the probability density function of Z is \mathbb{R} , it follows that $\sigma(x) = \mu_0 + \mu_1 x$ except on a set of measure zero in \mathbb{R} . \square

D.3 Proof of Theorem 7

To prove Theorem 7, we first need to state and prove a series of intermediary results.

Lemma D.1. *A necessary condition for optimality of $\sigma(x)$ is that*

$$-2x\sigma'(x) + 2\sigma''(x) + \lambda_1 + \lambda_2 x + \lambda_3 \sigma(x) = 0, \quad (75)$$

where λ_1, λ_2 , and λ_3 satisfy the following constraints,

$$\lambda_1 + \lambda_3 \mu_0 = 0, \quad (76)$$

$$-2\mu_1 + \lambda_2 + \lambda_3 \mu_1 = 0, \quad (77)$$

$$-2\mathbb{E}((\sigma'(Z))^2) + \lambda_1 \mu_0 + \lambda_2 \mu_1 + \lambda_3 \mu_2 = 0, \quad (78)$$

and that

$$\lim_{x \rightarrow +\infty} (\sigma'(x))^2 e^{-\frac{x^2}{2}} = \lim_{x \rightarrow -\infty} (\sigma'(x))^2 e^{-\frac{x^2}{2}} = 0. \quad (79)$$

Remark 11. Note that since (75) is a second order ODE, its solutions are parametrized by two constants in addition to being parametrized by $\lambda_1, \lambda_2, \lambda_3$. These five constants are set by our three constraints (25) together with two boundary conditions from our variational problem, which are $\lim_{x \rightarrow \pm\infty} \sigma'(x) e^{-\frac{x^2}{2}} = 0$. These last two we replace (see proof) by (79).

Remark 12. The lemma implies that knowing μ_0, μ_1, μ_2 and the objective value $\mathbb{E}((\sigma'(Z))^2)$ is enough to determine $\lambda_1, \lambda_2, \lambda_3$, even without solving the ODE.

Proof of Lemma D.1.

Derivation of equation (75): A Lagrangian for (25)-(25) is

$$\mathbb{E}((\sigma'(Z))^2 + \lambda_1(\mu_0 - \sigma(Z)) + \lambda_2(\mu_1 - Z\sigma(Z)) + \lambda_3((1/2)(\mu_2 - \sigma(Z))^2)), \quad (80)$$

which can be written as $\int_{-\infty}^{\infty} L(z, \sigma, \sigma') dz$ where, if we define $p(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}$, the Lagrangian density L is

$$L(z, \sigma, \sigma') = p(z)(\sigma'^2 + \lambda_1(\mu_0 - \sigma) + \lambda_2(\mu_1 - z\sigma) + \lambda_3(1/2)(\mu_2 - \sigma^2)). \quad (81)$$

We use the Euler-Lagrange equation with free boundary conditions [44] to get the necessary condition (75). To do so, we compute in sequence

$$\frac{\partial L}{\partial \sigma} = -p(z)(\lambda_1 + \lambda_2 z + \lambda_3 \sigma), \quad (82)$$

$$\frac{\partial L}{\partial \sigma'} = p(z)(2\sigma'), \quad (83)$$

$$\frac{d}{dz} \frac{\partial L}{\partial \sigma'} = p'(z)(2\sigma') + p(z)(2\sigma'') = p(z)(-2z\sigma' + 2\sigma''). \quad (84)$$

These lead to

$$0 = \frac{d}{dz} \frac{\partial L}{\partial \sigma'} - \frac{\partial L}{\partial \sigma} = p(z)(-2z\sigma' + 2\sigma'' + \lambda_1 + \lambda_2 z + \lambda_3 \sigma) \text{ and} \quad (85)$$

$$0 = \lim_{z \rightarrow \pm\infty} \frac{\partial L}{\partial \sigma'} = \lim_{z \rightarrow \pm\infty} p(z)\sigma'(z), \quad (86)$$

where the last condition follows from the fact that there is no boundary condition on σ . Since $p(z) > 0$, equation (85) implies (75).

Derivation of equations (79): Since we are working with necessary conditions, we can choose not list (86) in our lemma. Rather, we include condition (79), which a consequence of the fact that $\mathbb{E}((\sigma'(Z))^2)$ must be finite. Indeed, $\mathbb{E}((\sigma'(Z))^2) = \int_{-\infty}^{\infty} p(z)(\sigma'(z))^2 < \infty$ implies that $p(z)(\sigma'(z))^2$ must vanish at $\pm\infty$. Although not necessary for this proof, note that, since $p(z)$ goes to zero as $z \rightarrow \pm\infty$, equations (79) imply equations (86).

Derivation of equations (76)-(78):

Equation (75) must hold for all x . Hence, we can replace in it x by a standard normal random variable Z and compute the expected value of both of its sides. This leads to

$$-2\mathbb{E}(Z\sigma'(Z)) + 2\mathbb{E}(\sigma''(Z)) + \lambda_1 + \lambda_3\mu_0 = 0. \quad (87)$$

We can also multiply (75) by x , replace x by a standard normal random variable Z and compute the expected value of both of its sides. This leads to,

$$-2\mathbb{E}(Z^2\sigma'(Z)) + 2\mathbb{E}(Z\sigma''(Z)) + \lambda_2 + \lambda_3\mu_1 = 0. \quad (88)$$

Finally, we multiply (75) by $\sigma(x)$, replace x by a standard normal random variable Z and compute the expected value of both of its sides. This leads to,

$$-2\mathbb{E}(Z\sigma(Z)\sigma'(Z)) + 2\mathbb{E}(\sigma(Z)\sigma''(Z)) + \lambda_1\mu_0 + \lambda_2\mu_1 + \lambda_3\mu_2 = 0. \quad (89)$$

Using integration by parts, and the fact that for $p(z) = \frac{1}{\sqrt{2\pi}}e^{-\frac{z^2}{2}}$ we have that $p'(z) = -zp(z)$ and $p''(z) = (1 - z^2)p(z)$, we derive the following useful relationships

$$\mathbb{E}(\sigma''(Z)) = \mathbb{E}(Z\sigma'(Z)), \quad (90)$$

$$\mathbb{E}(Z\sigma''(Z)) = \mathbb{E}(Z^2\sigma'(Z)) - \mathbb{E}(\sigma'(Z)), \quad (91)$$

$$\mathbb{E}(\sigma'(Z)) = \mu_1, \quad (92)$$

$$\mathbb{E}(\sigma(Z)\sigma''(Z)) = \mathbb{E}(Z\sigma(Z)\sigma'(Z)) - \mathbb{E}((\sigma'(Z))^2). \quad (93)$$

To derive these relationships via integration by parts we made use of the following relationships

$$\lim_{z \rightarrow \pm\infty} \sigma'(z)\sigma(z)p(z) = \lim_{z \rightarrow \pm\infty} \sigma'(z)zp(z) = \lim_{z \rightarrow \pm\infty} \sigma'(z)p(z) = \lim_{z \rightarrow \pm\infty} \sigma(z)p(z) = 0, \quad (94)$$

which can be proved from $\mathbb{E}((\sigma(Z))^2), \mathbb{E}((\sigma'(Z))^2) < \infty$. Also, from $\mathbb{E}((\sigma(Z))^2), \mathbb{E}((\sigma'(Z))^2) < \infty$ and $\mu_1 \in \mathbb{R}$, we can show that each of the expected values in the right-hand-side of (90)-(93) is well-defined. Hence, the left-hand-side of (90)-(93) is well defined.

To finish the proof we replace (90) in (87) to get $\lambda_1 + \lambda_3\mu_0 = 0$, which is equation (76). Then we replace (91) and (92) in (88) to get $-2\mu_1 + \lambda_2 + \lambda_3\mu_1 = 0$, which is equation (77). Lastly, we replace (93) in (89) to get $-2\mathbb{E}((\sigma'(Z))^2) + \lambda_1\mu_0 + \lambda_2\mu_1 + \lambda_3\mu_2 = 0$, which is equation (78). \square

Lemma D.2. *The solutions of (75) are of the form*

$$\sigma(x) = \bar{\sigma}(x) + c_1 H_{\frac{\lambda_3}{2}}\left(\frac{x}{\sqrt{2}}\right) + c_2 F_{\{-\frac{\lambda_3}{4}\}, \{\frac{1}{2}\}}\left(\frac{x^2}{2}\right), \quad (95)$$

where

$$\bar{\sigma}(x) = \frac{\lambda_2 x}{2} - \frac{\lambda_1 x^2}{4} F_{\{1,1\}, \{\frac{3}{2}, 2\}}\left(\frac{x^2}{2}\right), \text{ if } \lambda_3 = 0 \quad (96)$$

$$\bar{\sigma}(x) = -\frac{\lambda_1}{2} - \frac{\lambda_2 x}{2} + \frac{1}{2} \sqrt{\frac{\pi}{2}} \lambda_2 e^{\frac{x^2}{2}} \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right) - \frac{1}{4} \lambda_2 x^3 F_{\{1,1\}, \{\frac{3}{2}, 2\}}\left(\frac{x^2}{2}\right), \text{ if } \lambda_3 = 2 \quad (97)$$

$$\bar{\sigma}(x) = -\frac{\lambda_1}{\lambda_3} - \frac{\lambda_2 x}{\lambda_3 - 2}, \text{ if } \lambda_3 \notin \{0, 2\} \quad (98)$$

where erf is the Gauss error function, $F_{\{a_k\}, \{b_k\}}$ is the generalized hypergeometric function with parameters $\{a_k\}, \{b_k\}$, and H_n is the Hermite polynomial, extended to a possibly non-integer n .

Remark 13. By an extension of $H_n(x)$ to non-integer we mean that $H_n(x) = F_{\{-\frac{1}{2}(n-1), \{\frac{3}{2}\}\}}(x^2)$, which is defined for non-integer n .

Proof of Lemma D.2. Since (75) is a second order ODE, its solutions are spanned by any particular solution $\bar{\sigma}(x)$ plus a linear combination of two solutions to its associated homogeneous ODE. The homogenous ODE associate with (75) is $-2x\sigma'(x) + 2\sigma''(x) + \lambda_3\sigma(x) = 0$. The change of variable $\sigma(x) = \bar{\sigma}(x/\sqrt{2})$ allows us to get $-2x\bar{\sigma}'(x) + \bar{\sigma}''(x) + \lambda_3\bar{\sigma}(x) = 0$, which is a well-known ODE called the *Hermite differential equation*. This implies that the homogeneous solutions of our ODE are spanned by $H_{\frac{\lambda_3}{2}}\left(\frac{x}{\sqrt{2}}\right)$ and $F_{\{-\frac{\lambda_3}{4}, \{\frac{1}{2}\}\}}\left(\frac{x^2}{2}\right)$. The particular solutions $\bar{\sigma}(x)$ can be confirmed by direct substitution. \square

Lemma D.3. *Let σ be of the form (95). If $\lambda_3 = 0$, then $\mathbb{E}((\sigma'(Z))^2) < \infty$ implies that $\sigma(x)$ is of the form*

$$\sigma(x) = c + \frac{x\lambda_2}{2}, \quad (99)$$

for some c , and that $\mathbb{E}((\sigma'(Z))^2) = \frac{\lambda_2^2}{4}$.

Proof of Lemma D.3. If $\lambda_3 = 0$ then based on Lemma D.2 we can re-write that

$$\sigma(x) = \sqrt{\frac{\pi}{2}}c_1 \operatorname{erfi}\left(\frac{x}{\sqrt{2}}\right) + c_2 - \frac{1}{4}x\left(\lambda_1 x F\left(\frac{x^2}{2}\right) - 2\lambda_2\right), \quad (100)$$

for some c_1, c_2 , where erfi is the inverse of erf . From this it follows that,

$$\sigma'(x) = \frac{1}{4}\left(e^{\frac{x^2}{2}}\left(4c_1 - \sqrt{2\pi}\lambda_1 \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right)\right) + 2\lambda_2\right). \quad (101)$$

The objective $\mathbb{E}((\sigma'(Z))^2) < \infty$ implies that $\lim_{x \rightarrow \pm\infty} (\sigma'(x))^2 e^{-\frac{x^2}{2}} = 0$. Since $\lim_{x \rightarrow \pm\infty} \operatorname{erf}(x) = \pm 1$, we have that $\lim_{x \rightarrow \pm\infty} (\sigma'(x))^2 e^{-\frac{x^2}{2}} = 0$ implies that $4c_1 = \pm\sqrt{2\pi}\lambda_1$ for both \pm simultaneously. This implies that $c_1 = \lambda_1 = 0$.

Substituting $c_1 = \lambda_1 = 0$ in (100) and simplifying we get $\sigma(x) = c_2 + \frac{\lambda_2 x}{2}$. From this expression one can then directly compute $\mathbb{E}((\sigma'(Z))^2)$. \square

Lemma D.4. *Let σ be of the form (95). If $\lambda_3 = 2$, then $\mathbb{E}((\sigma'(Z))^2) < \infty$ implies that $\sigma(x)$ is of the form*

$$\sigma(x) = cx - \frac{\lambda_1}{2}, \quad (102)$$

for some c , and that $\mathbb{E}((\sigma'(Z))^2) = c^2$.

Proof of Lemma D.4. If $\lambda_3 = 2$ then based on Lemma D.2 we can re-write that

$$\begin{aligned} \sigma(x) &= \frac{1}{4}\left(-x\left(\lambda_2\left(x^2 F_{\{1,1\}, \{\frac{3}{2}, 2\}}\left(\frac{x^2}{2}\right) + 2\right)\right.\right. \\ &\quad \left.\left.+ 2\sqrt{2}\left(\sqrt{\pi}c_2 \operatorname{erfi}\left(\frac{x}{\sqrt{2}}\right) - 2c_1\right) + e^{\frac{x^2}{2}}\left(\sqrt{2\pi}\lambda_2 \operatorname{erf}\left(\frac{x}{\sqrt{2}}\right) + 4c_2\right) - 2\lambda_1\right)\right) \end{aligned} \quad (103)$$

for some c_1 and c_2 , where erfi is the inverse of erf . From this it follows that

$$\sigma'(x) = \frac{2c_1 - \sqrt{\pi}c_2 \operatorname{erfi}\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{4}\lambda_2 x^2 F_{\{1,1\}, \{\frac{3}{2}, 2\}}\left(\frac{x^2}{2}\right). \quad (104)$$

The objective $\mathbb{E}((\sigma'(Z))^2) < \infty$ implies that $\lim_{x \rightarrow \pm\infty} (\sigma'(x))^2 e^{-\frac{x^2}{2}} = 0$. Since $\lim_{x \rightarrow \pm\infty} \left(\operatorname{erfi}\left(\frac{x}{\sqrt{2}}\right)\right)^2 = \lim_{x \rightarrow \pm\infty} \left(x^2 F_{\{1,1\}, \{\frac{3}{2}, 2\}}\left(\frac{x^2}{2}\right)\right)^2 = \infty$, and since

$$\lim_{x \rightarrow \pm\infty} \frac{x^2 F_{\{1,1\}, \{\frac{3}{2}, 2\}}\left(\frac{x^2}{2}\right)}{\operatorname{erfi}\left(\frac{x}{\sqrt{2}}\right)} = \pm\pi, \quad (105)$$

it follows we only have a finite objective if $\frac{-\sqrt{\pi}c_2}{\sqrt{2}} = \pm \frac{\lambda_2\pi}{4}$ for both \pm simultaneously. But this implies that $c_2 = \lambda_2 = 0$.

Substituting $c_2 = \lambda_2 = 0$ in (103), and simplifying, leads to $\sigma(x) = c_1\sqrt{2}x - \frac{\lambda_1}{2}$. The $\sqrt{2}$ factor can be absorbed by c_1 . From this expression one can compute $\mathbb{E}((\sigma'(Z))^2)$. \square

Lemma D.5. *Let σ be of the form (95). If $\lambda_3 \notin \{0, 2\}$, $c_1 = c_2 = 0$ then*

$$\sigma(x) = -\frac{\lambda_1}{\lambda_3} - \frac{\lambda_2 x}{\lambda_3 - 2}, \quad (106)$$

and

$$\mathbb{E}((\sigma'(Z))^2) = \left(\frac{\lambda_2}{(\lambda_3 - 2)} \right)^2. \quad (107)$$

Proof of Lemma D.5. This follows directly from Lemma D.2. \square

Lemma D.6. *Let σ be of the form (95). If $\lambda_3 \notin \{0, 2\}$, and either c_1 or c_2 are non-zero, then $\mathbb{E}((\sigma'(Z))^2) < \infty$ implies that $\lambda_3 = 4k$ for $k \in \mathbb{Z}^+$.*

Proof of Lemma D.6. If $\lambda_3 \notin \{0, 2\}$, then by Lemma D.2 we have a formula for $\sigma(x)$ from which we get that

$$\sigma'(x) = \frac{c_1 \lambda_3 H_{\frac{\lambda_3}{2}-1} \left(\frac{x}{\sqrt{2}} \right)}{\sqrt{2}} - \frac{1}{2} c_2 \lambda_3 x F_{\{1-\frac{\lambda_3}{4}\}, \{\frac{3}{2}\}} \left(\frac{x^2}{2} \right) - \frac{\lambda_2}{\lambda_3 - 2}, \quad (108)$$

where F and H are as defined in Lemma D.2.

The boundedness of $\mathbb{E}((\sigma'(Z))^2)$ is dependent on how fast $H_{\frac{\lambda_3}{2}-1} \left(\frac{x}{\sqrt{2}} \right)$ and $x F_{\{1-\frac{\lambda_3}{4}\}, \{\frac{3}{2}\}} \left(\frac{x^2}{2} \right)$ grow as $x \rightarrow \pm\infty$.

If $c_1 = 0$ and $c_2 \neq 0$, $\mathbb{E}((\sigma'(Z))^2) < \infty$ only if $\mathbb{E} \left(\left(x F_{\{1-\frac{\lambda_3}{4}\}, \{\frac{3}{2}\}} \left(\frac{x^2}{2} \right) \right)^2 \right) < \infty$, which in turn is true only if $\lim_{x \rightarrow \infty} \left(x F_{\{1-\frac{\lambda_3}{4}\}, \{\frac{3}{2}\}} \left(\frac{x^2}{2} \right) \right)^2 e^{-\frac{x^2}{2}} = 0$. This implies that λ_3 is even.

If $c_1 \neq 0$ and $c_2 = 0$, $\mathbb{E}((\sigma'(Z))^2) < \infty$ only if $\mathbb{E} \left(\left(H_{\frac{\lambda_3}{2}-1} \left(\frac{x}{\sqrt{2}} \right) \right)^2 \right) < \infty$, which in turn is true only if $\lim_{x \rightarrow \infty} \left(H_{\frac{\lambda_3}{2}-1} \left(\frac{x}{\sqrt{2}} \right) \right)^2 e^{-\frac{x^2}{2}} = 0$. This implies that λ_3 is even.

If $c_1, c_2 \neq 0$, $\mathbb{E}((\sigma'(Z))^2) < \infty$ only if $\lim_{x \rightarrow \infty} (\sigma'(x))^2 e^{-\frac{x^2}{2}} = 0$. But $x F_{\{1-\frac{\lambda_3}{4}\}, \{\frac{3}{2}\}} \left(\frac{x^2}{2} \right)$ grows much faster than $H_{\frac{\lambda_3}{2}-1} \left(\frac{x}{\sqrt{2}} \right)$ as $x \rightarrow \infty$, hence it must be that $\lim_{x \rightarrow \infty} \left(x F_{\{1-\frac{\lambda_3}{4}\}, \{\frac{3}{2}\}} \left(\frac{x^2}{2} \right) \right)^2 e^{-\frac{x^2}{2}} = 0$. This implies that λ_3 is even. \square

Lemma D.7. *Let $\lambda_3 = 4k$ for $k \in \mathbb{Z}^+$, and let $\sigma(x)$ be a solution of (75), then*

$$\sigma(x) = -\frac{\lambda_1}{\lambda_3} - \frac{\lambda_2 x}{\lambda_3 - 2} + c H_{2k} \left(\frac{x}{\sqrt{2}} \right) \quad (109)$$

for some c and

$$\mathbb{E}((\sigma'(Z))^2) = \frac{4^k ((2k)!)^2}{(2k-1)!} c^2 + \left(\frac{\lambda_2}{\lambda_3 - 2} \right)^2. \quad (110)$$

Proof of Lemma D.7. If $\lambda_3 = 4k$ then

$$F_{\{\frac{\lambda_3}{4}\}, \{\frac{1}{2}\}}(x^2/2) = \frac{(-1)^k k!}{(2k)!} H_{\lambda_3/2}(x/\sqrt{2}), \quad (111)$$

and hence the homogenous part of $\sigma(x)$ can be written as $cH_{\lambda_3/2}(x)$, from which (109) follows. From this expression it follows that

$$\sigma'(x) = \frac{c\lambda_3 H_{\frac{\lambda_3}{2}-1}\left(\frac{x}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\lambda_2}{\lambda_3 - 2}, \quad (112)$$

and from this expression the value for $\mathbb{E}((\sigma'(Z))^2)$ follows. \square

Lemma D.8. *Let σ be of the form (95). If $\lambda_3 \notin \{0, 2\}$, then $\mathbb{E}((\sigma'(Z))^2) < \infty$ implies that $\sigma(x)$ is of the form*

$$\sigma(x) = -\frac{\lambda_1}{\lambda_3} - \frac{\lambda_2 x}{\lambda_3 - 2} + cH_{\lambda_3/2}\left(\frac{x}{\sqrt{2}}\right) \quad (113)$$

for some c that must be zero if $\lambda_3 \neq 4k$, $k \in \mathbb{Z}^+$, and

$$\mathbb{E}((\sigma'(Z))^2) = \frac{4^k((2k)!)^2}{(2k-1)!} c^2 + \left(\frac{\lambda_2}{\lambda_3 - 2}\right)^2. \quad (114)$$

Proof of Lemma D.8. This follows directly from Lemmas D.5-D.7. \square

Lemma D.9. *If $\lambda_1, \lambda_2, \lambda_3$ satisfy (76)-(78), then*

$$\left(\frac{\lambda_2}{\lambda_3 - 2}\right)^2 = \mu_1^2. \quad (115)$$

Proof of Lemma D.9. Defining $R = \mathbb{E}((\sigma'(Z))^2)$ and solving the linear system (76)-(78) leads to

$$\lambda_1 = \frac{2\mu_0(R - \mu_1^2)}{\mu_0^2 + \mu_1^2 - \mu_2}, \lambda_2 = \frac{2\mu_1(\mu_0^2 - \mu_2 + R)}{\mu_0^2 + \mu_1^2 - \mu_2}, \lambda_3 = -\frac{2(R - \mu_1^2)}{\mu_0^2 + \mu_1^2 - \mu_2}. \quad (116)$$

Computing $\lambda_2/(\lambda_3 - 2)$ we get $-\mu_1$, from which the first relation follows.

If we solve $\lambda_3 = -\frac{2(R - \mu_1^2)}{\mu_0^2 + \mu_1^2 - \mu_2}$ for R and recall that $\mu_\star^2 = \mu_2 - \mu_0^2 - \mu_1^2$, the second relation follows. \square

We are now ready to prove Theorem 7, which we restate here for convenience.

Theorem (7). *The minimizers of (25) are*

$$\sigma(x) = ax^2 + bx + c, \quad (117)$$

where

$$a = \pm \frac{\mu_\star}{\sqrt{2}}, b = \mu_1, \text{ and } c = \mu_0 - a. \quad (118)$$

Proof of Theorem 7. We will show that any minimizer of (25) must be a function $\sigma(x) = ax^2 + bx + c$ for some a, b, c . From this fact, the variational problem can be reduced to a simple quadratic programming problem over a, b, c , from which it is straightforward to derive (117) and (118).

If $\mu_\star = 0$ then by Lemma 4.1, we know that the solution must be linear, and hence we are done.

If $\mu_\star > 0$ then by Lemma 4.1 we know that $\sigma(x)$ cannot be a linear function. Hence, from Lemma D.3 and Lemma D.4 we know that λ_3 cannot be 0 or 2. Therefore, its solution must be of the form specified by Lemma D.8 with $c \neq 0$ and $\lambda_3 = 4k$ for some $k \in \mathbb{Z}^+$.

Define $R = \mathbb{E}((\sigma'(Z))^2)$. From the constraints (25) we know that λ_1, λ_2 and R can be written as a linear function of $\lambda_3 = 4k$. In particular, $R = \mu_1^2 + \frac{\lambda_3}{2}\mu_\star^2$. From (114) and Lemma D.9 we can write

that $R = \frac{4^k((2k)!)^2}{(2k-1)!} c^2 + \mu_1^2$, which implies that $c = \pm \left(\frac{R - \mu_1^2}{\frac{4^k((2k)!)^2}{(2k-1)!}}\right)$ is also a function of λ_3 . Therefore,

the solution to is parametrized by λ_3 alone which must be chosen to minimize $R = \mu_1^2 + \frac{\lambda_3}{2}\mu_\star^2$. Therefore, we must choose $\lambda_3 = 4$, the smallest possible multiple of 4. This implies that $H_{\lambda_3/2}(x/\sqrt{2})$ is a quadratic function, from which it follows that $\sigma(x)$ is also a quadratic function, and hence we are done. \square

D.4 Proof of Theorem 8

Before we prove Theorem 8, we will state and prove a series of intermediary results.

Lemma D.10. *A necessary condition for optimality of $\sigma(x)$ is that*

$$x + \lambda_1 + \lambda_2 x + \lambda_3 \sigma(x) = 0, \text{ for all } x : \sigma'(x) \neq 0, \quad (119)$$

where $\lambda_1, \lambda_2,$ and λ_3 must be such that,

$$\mathbb{E}(\sigma(Z)) = \mu_0, \quad (120)$$

$$\mathbb{E}(\sigma(Z)Z) = \mu_1, \quad (121)$$

$$\mathbb{E}((\sigma(Z))^2) = \mu_2. \quad (122)$$

Proof of Lemma D.10. A Lagrangian for (119) is

$$\mathbb{E}(|\sigma'(Z)| + \lambda_1(\mu_0 - \sigma(Z)) + \lambda_2(\mu_1 - Z\sigma(Z)) + \lambda_3((1/2)(\mu_2 - \sigma(Z))^2)), \quad (123)$$

which can be written as $\int_{-\infty}^{\infty} L(z, \sigma, \sigma') dz$ where, if we define $p(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}$, the Lagrangian density L is

$$L(z, \sigma, \sigma') = p(z)(|\sigma'| + \lambda_1(\mu_0 - \sigma) + \lambda_2(\mu_1 - z\sigma) + \lambda_3(1/2)(\mu_2 - \sigma^2)). \quad (124)$$

Any variation $\sigma(z) + \epsilon\eta(z)$ of an optimal $\sigma(z)$ must yield $\mathbb{E}(|\sigma'(z) + \epsilon\eta'(z)|) \geq \mathbb{E}(|\sigma'(z)|)$. In particular, this must be the case for any variation such that $\eta(z) = 0$ whenever $\sigma'(z) = 0$. If we focus on these variations, from $\mathbb{E}(|\sigma'(z) + \epsilon\eta'(z)|) \geq \mathbb{E}(|\sigma'(z)|)$ the Euler-Lagrange equation can be derived despite $|\cdot|$ not being differentiable at 0. To be specific, it must hold that

$$\frac{d}{dz} \frac{\partial L}{\partial \sigma'} - \frac{\partial L}{\partial \sigma} = 0 \quad \forall z : \sigma'(z) \neq 0. \quad (125)$$

Since there are no fixed boundary conditions on our integration domain $(-\infty, \infty)$, it also needs to hold that $\lim_{z \rightarrow \pm\infty} \frac{\partial L}{\partial \sigma'} = \lim_{z \rightarrow \pm\infty} p(z)\sigma'(z) = 0$, which we choose not to list in our necessary conditions.

Since $\frac{d}{dz} \frac{\partial L}{\partial \sigma'} = 0$ if $\sigma'(z) \neq 0$, equation (125) implies (119). First order optimality conditions imply that the Lagrange multipliers must be choose such that $\mathbb{E}(\sigma(Z)) = \mu_0, \mathbb{E}(\sigma(Z)Z) = \mu_1, \mathbb{E}((\sigma(Z))^2) = \mu_2$. \square

Lemma D.11. *If $\lambda_3 \neq 0$, the solutions of (119) are of the form*

$$\sigma(x) = -\frac{\lambda_1}{\lambda_3} - \frac{1 + \lambda_2}{\lambda_3} \min\{\max\{x, b\}, c\}, \quad (126)$$

for some constants b, c where $b < c$ if $\frac{1+\lambda_2}{\lambda_3} \leq 0$ and $b > c$ otherwise.

Proof of Lemma D.11. Since σ is a one-dimensional function, the requirement of existence of weak derivative implies that there exists v absolute continuous that agrees with σ almost everywhere. We will work with these absolute continuous representations of σ . Other solutions can differ from the absolute continuous solutions only up to a set of measure zero with respect to the Gaussian measure.

From (119) we know that wherever $\sigma'(x) \neq 0$, we have that $\sigma(x) = ax + b$ for the same fixed a and b . Hence, since σ is continuous, σ must be an alternation of flat portions and portions with the same slope a . Because of continuity, we cannot have a flat portion interrupt portion of slope a (unless $a = 0$), as illustrated in Figure 2-right, and σ must be as in the other two cases in Figure 2. These have a form as in (126). \square

Lemma D.12. *Let σ be of the form (126). Then,*

$$\mathbb{E}(|\sigma'(Z)|) = \frac{1}{2} \left| \frac{1 + \lambda_2}{\lambda_3} \right| \mathbb{P}(Z \in [b, c]), \quad (127)$$

$$\mu_1 \triangleq \mathbb{E}(Z\sigma(Z)) = -\frac{1}{2} \frac{1 + \lambda_2}{\lambda_3} \mathbb{P}(Z \in [b, c]), \quad (128)$$

where $[b, c]$ should be interpreted as $[c, b]$ if $c < b$. In particular, $\mathbb{E}(|\sigma'(Z)|) = |\mu_1|$.




Figure 2: The first two functions are the only two possible types of continuous functions that satisfy (119). The right-most function also satisfies (119) but is not continuous.

Proof of Lemma D.12. From Lemma D.11, we have explicit formulas for σ and σ' . The proof boils down to a direct calculation of the expected values, which themselves boil down to computing a few Gaussian integrals. \square

We are now ready to prove Theorem 8, which we restate below for convenience.

Theorem (8). *One minimizer of (3) is*

$$\sigma(x) = \mu_0 + a \min\{\max\{x, -s\}, s\} \quad (129)$$

where $a = \frac{\mu_1}{\text{erf}(s/\sqrt{2})}$, erf is the Gauss error function, and $s \in \mathbb{R}$ is the unique solution to the equation

$$\zeta^2 \triangleq \frac{\mu_1^2}{\mu_\star^2} = \frac{e^{\frac{s^2}{2}} \text{erf}\left(\frac{s}{\sqrt{2}}\right)^2}{e^{\frac{s^2}{2}} \left(1 - \text{erf}\left(\frac{s}{\sqrt{2}}\right)\right) \left(s^2 + \text{erf}\left(\frac{s}{\sqrt{2}}\right)\right) - \sqrt{\frac{2}{\pi}} s}, \quad (130)$$

if $\mu_\star \neq 0$, and $s = \infty$ if $\mu_\star = 0$.

Proof of Theorem 8. From Lemma D.11, we know that if $\lambda_3 \neq 0$, then any minimizer must have the form (126). From Lemma D.12 we know that all of the functions of this form have the same objective. Hence, if $\lambda_3 \neq 0$, all of the functions of the form (126) that satisfy constraints (25) are a global minimizer.

We set $\lambda_3 = 1 \neq 0$. To satisfy the three constraints (25) we have 4 remaining values to play with, namely $\lambda_1, \lambda_2, b, c$. Hence, we set $b = -c$. With a reparameterization, this leads to σ having the form $\sigma(x) = b + a \min\{\max\{x, -s\}, s\}$, where b has a new meaning. That is, σ is flat outside of the interval $[-s/|a|, s/|a|]$, $s \geq 0$, and inside of this interval it has slope a . The goal is to find a, b, s from the constraints (25).

From $\sigma(x) = b + a \min\{\max\{x, -s\}, s\}$ a direct computations leads to

$$\mu_0 = b, \quad (131)$$

$$\mu_1 = a \text{erf}\left(\frac{s}{\sqrt{2}}\right), \quad (132)$$

$$\mu_2 = a^2 (s^2 - 1) \text{erfc}\left(\frac{s}{\sqrt{2}}\right) + a^2 \left(1 - \sqrt{\frac{2}{\pi}} s e^{-\frac{s^2}{2}}\right) + b^2, \quad (133)$$

where $\text{erfc} = 1 - \text{erf}$. We can use the first two equations to write b and a as a function of μ_0, μ_1, s . Substituting a and b with these functions in the third equation, and simplifying, leads to

$$\mu_2 = \mu_0^2 - \frac{\mu_1^2 \left(- (s^2 - 1) \text{erfc}\left(\frac{s}{\sqrt{2}}\right) + \sqrt{\frac{2}{\pi}} e^{-\frac{s^2}{2}} s - 1 \right)}{\text{erf}\left(\frac{s}{\sqrt{2}}\right)^2}. \quad (134)$$

Recalling that $\mu_2 = \mu_\star^2 + \mu_1^2 + \mu_0^2$, replacing this definition into the above equation, and simplifying leads to

$$\frac{\mu_1^2}{\mu_\star^2} \triangleq \zeta^2 = - \frac{e^{\frac{s^2}{2}} \text{erf}\left(\frac{s}{\sqrt{2}}\right)^2}{e^{\frac{s^2}{2}} \left(\text{erf}\left(\frac{s}{\sqrt{2}}\right) - 1\right) \left(s^2 + \text{erf}\left(\frac{s}{\sqrt{2}}\right)\right) + \sqrt{\frac{2}{\pi}} s}. \quad (135)$$

One can show that the function on the right hand side (135) is monotonic increasing in $s \in [-\infty, \infty]$ with range $[0, \infty]$, which implies that there is only one s that solves (135). \square

D.5 Proof of Theorem 9

Proof of Theorem 9. Theorem 9 amounts a statement about the solutions of the optimization problem (29) for regime R_1 .

Its proof amounts to studying the local minima of the objective via its first and second derivatives.

We will prove the theorem for $\psi_1 < \psi_2$ and $\psi_1 > \psi_2$ separately. For $\psi_1 = \psi_2$ the objective is not defined.

In what follows, we let $O = (1 - \alpha)\mathcal{E}_{R_1}^\infty + \alpha\mathcal{S}_{R_1}^\infty$. We will use the fact that O is a one-dimensional function of $\zeta^2 \in [0, +\infty]$. We will use this and the fact that (14) defines a monotonic function between χ and ζ^2 , to express O as a function of χ and study the solutions of the optimization problem (29)

in the variable χ . Notice that (14) can be solved for ζ^2 as $\zeta^2 = \frac{\mu_1^2}{\mu_*^2} = \frac{\chi + \min\{\psi_1, \psi_2\}}{\chi(-1 + \chi + \min\{\psi_1, \psi_2\})}$ and, calling χ by x , this can be arranged to get (30), which connects optimal values of χ with optimal values of μ_0, μ_1, μ_2 . Henceforth, we denote χ by x . We note that since $\zeta^2 \in [0, +\infty]$, from (14) it follows that $x \in [x_L, x_R] \triangleq [-\min\{\psi_1, \psi_2\}, \min\{0, 1 - \min\{\psi_1, \psi_2\}\}]$. Hence, we only need to study the function $O(x)$ in this interval.

Case 1, $\psi_1 < \psi_2$:

For the sake of simplicity, and for the most part, we will assume that $\psi_1 \neq 1$ and omit the argument for $\psi_1 = 1$. The argument for $\psi_1 = 1$ is almost identical to the argument for $\psi_1 \neq 1$ if we work with the extended reals $[-\infty, \infty]$. Furthermore, most conclusions for $\psi_1 = 1$ can be obtained by taking the limit of $\psi_1 \rightarrow 1$ with $\psi_1 \neq 1$. The only situation where this is not the case is that the polynomial $p(x)$, referenced right after Table 1, has an expression when $\psi_1 = 1$ that cannot be obtained as the limit of its expression for $1 \neq \psi_1$.

We start by studying the second derivative of O . A direct computation yields

$$\frac{d^2 O}{dx^2} = \tilde{O}(x) + C, \text{ where } C = \frac{2F_1^2 \alpha}{\psi_1 - \psi_2} \quad (136)$$

$$\text{and } \tilde{O}(x) = -\frac{2\alpha F_1^2 \psi_1 (-3x^2(\psi_1 - 1) - 2x^3 + (\psi_1 - 1)^2 \psi_1)}{(x + \psi_1)^2 - \psi_1^3}.$$

A tedious calculation (omitted) shows that $\frac{d^2 \tilde{O}}{dx^2} \geq 0$ for $x \in [x_L, x_R]$ (i.e. it is convex), that $\tilde{O}(x_L) < \tilde{O}(x_R)$, and that $\frac{d\tilde{O}}{dx}(x = x_L) < 0$. From this it follows that, depending on the value of ψ_2 , the concavity $\frac{d^2 O}{dx^2}$ is positive or negative, as illustrated in the following figure:




Figure 3: Depending on the value of ψ_2 , the function O is convex, convex-concave-convex, concave-convex, or concave respectively. The points A, B, and C will be referenced later in the proof. It is possible to compute closed-form expressions for the y-coordinate of these points.

To be specific, starting from large ψ_2 , i.e. small $-C$, and decreasing its value, i.e. increasing $-C$, we obtain the following four scenarios. While ψ_2 is large and $-C$ is below A_2 , the function is

convex. After ψ_2 reaches a value β_1 at which $-C$ touches A_2 , the function is convex for small x , then concave, and then convex for large x . As ψ_2 keeps decreasing, and after it reaches a value β_2 at which $-C$ touches A_1 , the function is concave for small x , and then convex for large x . Finally, after ψ_2 reaches a value β_3 at which $-C$ touches A_3 , the function is concave. Note that by definition $0 \leq \beta_3 < \beta_2 < \beta_1$.

It is possible to compute closed-form expressions for the y -coordinate of the points A_1, A_2, A_3 , which we denote by A_1, A_2 , and A_3 . Namely, $A_1 = F_1^2 \alpha (2 + \frac{2}{\psi_1})$, $A_2 = F_1^2 \alpha \left(\frac{\sqrt{1-\psi_1}+1}{\psi_1} - \frac{1}{2} \right)$, if $\psi_1 \leq 1$, and $A_2 = F_1^2 \alpha \left(\sqrt{\frac{\psi_1-1}{\psi_1}} - \frac{1}{2\psi_1} + 1 \right)$, if $\psi_1 > 1$, and $A_3 = F_1^2 \alpha \max \left\{ \frac{2}{\psi_1 - \psi_1^2}, \frac{2\psi_1}{\psi_1 - 1} \right\}$. Using the closed-form expression for $-C$, see (136), we find closed-form expressions for β_1, β_2 , and β_3 as $\beta_1 = \psi_2 : A_2(\psi_2) + C(\psi_2) = 0$, $\beta_2 = \psi_2 : A_1(\psi_2) + C(\psi_2) = 0$, and $\beta_3 = \psi_2 : A_3(\psi_2) + C(\psi_2) = 0$. By definition of β_1, β_2 , and β_3 , these equations have a unique solution when $\psi_2 > \psi_1 > 0$ and their explicit expressions are given in (31), (32) and (33) respectively.

Now that we have characterized the curvature of $O(x)$, we are ready to locate its global minimum. To do so, we will use the curvature of $O(x)$ together with the first-order optimality condition $\frac{dO}{dx} = 0$, and the following three extra pieces of information: the sign of the derivative of $O(x)$ at $x = x_L$; the sign of the derivative of $O(x)$ at $x = x_R$; and the sign of $O(x_L) - O(x_R)$.

A direct computation yields

$$\frac{dO}{dx} = \frac{p_0 + p_1x + p_2x^2 + p_3x^3 + p_4x^4 + p_5x^5}{(\psi_1 - (x + \psi_1)^2)^2(\psi_1 - \psi_2)}, \text{ if } \psi_1 \neq 1, \text{ and} \quad (137)$$

$$\frac{dO}{dx} = \frac{q_0 + q_1x + q_2x^2 + q_3x^3}{(2+x)^2(-1+\psi_2)}, \text{ if } \psi_1 = 1. \quad (138)$$

where the coefficients p_0, \dots, p_5 and q_0, \dots, q_3 are, apart from a multiplying constant, given in (57)-(66).

The roots of the first denominator, i.e. $-\sqrt{\psi_1} - \psi_1$ and $\sqrt{\psi_1} - \psi_1$, are not a root of the first numerator when $\psi_1 \neq 1$, and the roots of the second denominator, i.e. -2 , are not a root of the second numerator when $\psi_2 > \psi_1 = 1$. Therefore, the first-order optimality conditions are $p(x) = 0$.

Not all solutions of $p(x) = 0$ minimize $O(x)$. To locate the minimizer, we use the sign of the derivative of $O(x)$ at $x = x_L$; the sign of the derivative of $O(x)$ at $x = x_R$; and the sign of $O(x_L) - O(x_R)$. These signs can be determined using Lemma D.13. Lemma D.13 also proves Remark 9.

Lemma D.13. *If $\psi_1 < \psi_2$, the following relationships hold,*

$$\left. \frac{dO}{dx} \right|_{x=x_L} = -\frac{F_1^2(\alpha + (\alpha - 1)\psi_2) + \alpha(F_\star^2 + \tau^2)}{\psi_1 - \psi_2}, \quad (139)$$

$$\left. \frac{dO}{dx} \right|_{x=x_R} = \frac{F_1^2\psi_2 - \alpha(F_1^2(-4\psi_1 + 3\psi_2 + 1) + F_\star^2 + \tau^2)}{\psi_1 - \psi_2} \text{ if } \psi_1 \leq 1, \quad (140)$$

$$\left. \frac{dO}{dx} \right|_{x=x_R} = \frac{F_1^2(\alpha + 2\alpha\psi_1 - 3\alpha\psi_2 + \psi_2) - \alpha(F_\star^2 + \tau^2)}{\psi_1 - \psi_2} \text{ if } \psi_1 > 1, \quad (141)$$

$$O(x_L) - O(x_R) = \frac{\psi_1(F_1^2(\alpha - 2\alpha\psi_1 + (2\alpha - 1)\psi_2) + \alpha(F_\star^2 + \tau^2))}{\psi_1 - \psi_2} \text{ if } \psi_1 \leq 1, \text{ and} \quad (142)$$

$$O(x_L) - O(x_R) = \frac{\alpha(F_\star^2 + \tau^2) - F_1^2(\alpha\psi_1 - 2\alpha\psi_2 + \psi_2)}{\psi_1 - \psi_2} \text{ if } \psi_1 > 1. \quad (143)$$

Also,

$$\left. \frac{dO}{dx} \right|_{x=x_L} < 0 \text{ if } \alpha < \alpha_L \text{ and } \left. \frac{dO}{dx} \right|_{x=x_L} \geq 0 \text{ if } \alpha \geq \alpha_L, \quad (144)$$

$$\left. \frac{dO}{dx} \right|_{x=x_R} < 0 \text{ if } \alpha < \alpha_R \text{ and } \left. \frac{dO}{dx} \right|_{x=x_R} \geq 0 \text{ if } \alpha \geq \alpha_R, \text{ and} \quad (145)$$

$$O(x_L) - O(x_R) > 0 \text{ if } \alpha < \alpha_C \text{ and } O(x_L) - O(x_R) \leq 0 \text{ if } \alpha \geq \alpha_C, \quad (146)$$

where α_L, α_R and α_C are given in (34), (36), and (35) respectively.

Furthermore, the following are true and help determine from which row in Table 1 to read x . If $\psi_2 < \min\{2\psi_1, \psi_1 + 1\}$, then $\alpha_L < \alpha_C < \alpha_R$. If $\psi_2 > \min\{2\psi_1, \psi_1 + 1\}$, then $\alpha_R < \alpha_C < \alpha_L$. If $\psi_2 = \min\{2\psi_1, \psi_1 + 1\}$, then $\alpha_L = \alpha_C = \alpha_R$. In particular, it follows from this that if $\psi_2 > \beta_1$, then $\alpha_R \leq \alpha_C \leq \alpha_L$ and if $\psi_2 \leq \beta_3$, then $\alpha_L \leq \alpha_C \leq \alpha_R$.

Finally, $\alpha_L, \alpha_R, \alpha_C \in [0, 1]$.

Proof. The derivation of (139)-(143) follows from direct substitution of $x = x_L$ or $x = x_R$ into O (for which we have expressions from Theorems 1 and 4) or its derivative (in eq. (137)-(138)). The derivation of (144)-(146) follows from the observation that the equations (139)-(143) are linear in α , and hence we can easily compute the values of α at which these expressions change from a negative value to a positive value. Once an explicit formula for α_L, α_C , and α_R is obtained, it is easy to find the criteria to decide their relative magnitude by comparing the term under parenthesis in the denominators of (34), (36), and (35). It is also easy to see from their formulas that their value is always in the range $[0, 1]$. \square

To finish the proof of Theorem 9 we consider the different scenarios in Table 1. In what follows, statements about the concavity of O are justified via the explanation accompanying Figure 3, and statements about the slope of O are based on Lemma D.13.

Case 1.1, $\psi_1 < \psi_2 \wedge \psi_2 \geq \beta_1 \wedge \alpha < \alpha_L \wedge \alpha < \alpha_R$: The function O is convex and is decreasing at $x = x_L$ and decreasing at $x = x_R$. Hence its minimum is at $x = x_R$.

Case 1.2, $\psi_1 < \psi_2 \wedge \psi_2 \geq \beta_1 \wedge \alpha_R < \alpha < \alpha_L$: The function O is convex and it is decreasing at $x = x_L$ and increasing at $x = x_R$. Hence it has a unique minimizer at $x = x_1$.

When $\psi_1 < \psi_2 \wedge \psi_2 \geq \beta_1$, we can use Lemma D.13 and a tedious calculation (omitted) to show that $\alpha_R < \alpha_L$, which is why the 4th and 5th rows of the first column of Table 1 are empty.

Case 1.3, $\psi_1 < \psi_2 \wedge \psi_2 \geq \beta_1 \wedge \alpha > \alpha_L \wedge \alpha > \alpha_R$: The function O is convex and it is increasing at both $x = x_L$ and $x = x_R$. Hence its minimum is at $x = x_L$.

Case 1.4, $\psi_1 < \psi_2 \wedge \beta_2 < \psi_2 < \beta_1 \wedge \alpha < \alpha_L \wedge \alpha < \alpha_R$: The function O is first convex, then concave and then convex. It is decreasing at both $x = x_L$ and at $x = x_R$. Hence O has at most one local minimum in the interior of the domain, at $x = x_1$ if it exists, and a local minimum at x_R , the domain being $[x_L, x_R]$. Therefore, the minimum can be expressed as $x_R \sqcup x_1$.

Case 1.5, $\psi_1 < \psi_2 \wedge \beta_2 < \psi_2 < \beta_1 \wedge \alpha_R < \alpha < \alpha_L$: The function O is first convex, then concave and then convex. It is decreasing at $x = x_L$ and increasing at $x = x_R$. Hence O has no local minimum at the end points of the domain $[x_L, x_R]$. Also, O either has exactly one local minimum in the interior of the domain, at $x = x_1$, or, O has exactly two local minimum and one local maximum (resp.) in the interior of the domain, at $x = x_1, x = x_3$ and $x = x_2$ respectively. Therefore, the minimum can be expressed as $x_1 \sqcup x_3$. Note that x_1 always exists but x_3 might not.

Case 1.6, $\psi_1 < \psi_2 \wedge \beta_2 < \psi_2 < \beta_1 \wedge \alpha_L < \alpha < \alpha_R$: The function O is first convex, then concave and then convex. It is increasing at $x = x_L$ and decreasing at $x = x_R$. Hence, the minimum is either at the x_L or x_R , depending whether $\alpha > \alpha_C$ or $\alpha < \alpha_C$. This case is an example where it is clear that allowing $\alpha = \alpha_C$ leads to non-uniqueness in the optimal x . See Remark 9.

Case 1.7, $\psi_1 < \psi_2 \wedge \beta_2 < \psi_2 < \beta_1 \wedge \alpha > \alpha_L \wedge \alpha > \alpha_R$: The function O is first convex, then concave, and then convex. It is increasing at both $x = x_R$ and $x = x_L$. Hence O either has no critical point in the interior of the domain, or it has two critical point in the interior of the domain, at $x = x_1$ (local maximum) and $x = x_2$ (local minimum) if they exists, and always has a local minimum at x_L . Therefore, the minimum can be expressed as $x_L \sqcup x_2$. Note that x_L always exists, but x_2 not always exists.

Case 1.8, $\psi_1 < \psi_2 \wedge \beta_3 < \psi_2 \leq \beta_2 \wedge \alpha < \alpha_L \wedge \alpha < \alpha_R$: The function O is first concave and convex. and is decreasing at both $x = x_L$ and $x = x_R$. Hence its minimum is at $x = x_R$.

Case 1.9, $\psi_1 < \psi_2 \wedge \beta_3 < \psi_2 \leq \beta_2 \wedge \alpha_R < \alpha < \alpha_L$: The function O is first concave and convex. It is decreasing at $x = x_L$ and increasing at $x = x_R$. Hence, O has exactly one local minimum in the interior of the domain, which is at $x = x_1$ and always exists, and which is also a global minimum.

Case 1.10, $\psi_1 < \psi_2 \wedge \beta_3 < \psi_2 \leq \beta_2 \wedge \alpha_L < \alpha < \alpha_R$: The function O is first concave and convex. It is increasing at $x = x_L$ and decreasing at $x = x_R$. Hence, the minimum is either at the x_L or x_R , depending whether $\alpha > \alpha_C$ or $\alpha < \alpha_C$.

Case 1.11, $\psi_1 < \psi_2 \wedge \beta_3 < \psi_2 \leq \beta_2 \wedge \alpha > \alpha_L \wedge \alpha > \alpha_R$: The function O is first concave and convex. It is increasing at both $x = x_L$ and $x = x_R$. Hence O either has no critical point in the interior of the domain or it has two critical point in the interior of the domain, at $x = x_1$ (local maximum) and $x = x_2$ (local minimum) if they exists, and always has a local minimum at x_L . Therefore, the minimum can be expressed as $x_L \sqcup x_2$. Note that x_L always exists, but x_2 not always exists.

Case 1.12, $\psi_1 < \psi_2 \wedge \psi_2 \leq \beta_3 \wedge \alpha < \alpha_L \wedge \alpha < \alpha_R$: The function O is concave. It is decreasing at both $x = x_L$ and $x = x_R$. Hence, its minimum is at $x = x_R$

When $\psi_1 < \psi_2 \wedge \psi_2 \geq \beta_1$, Lemma D.13 and a tedious calculation (omitted) shows that that $\alpha_L < \alpha_R$, which is why the second and third rows of the last column of Table 1 are empty.

Case 1.13, $\psi_1 < \psi_2 \wedge \psi_2 \leq \beta_3 \wedge \alpha_L < \alpha < \alpha_R$: The function O is concave. It is increasing at $x = x_L$ and decreasing at $x = x_R$. Hence, the minimum is either at the x_L or x_R , depending whether $\alpha > \alpha_C$ or $\alpha < \alpha_C$.

Case 1.14, $\psi_1 < \psi_2 \wedge \psi_2 \leq \beta_3 \wedge \alpha > \alpha_L \wedge \alpha > \alpha_R$: The function O is concave. It is increasing at $x = x_L$ and at $x = x_R$. Hence, the minimum is at x_L .

Case 2, $\psi_1 > \psi_2$:

We will deal with the case $\psi_2 = 1$ at the end. For now, we assume that $\psi_2 \neq 1$.

We first prove that the second derivative of O is convex, just like in the case when $\psi_1 < \psi_2$. A direct computation yields

$$\frac{d^2O}{dx^2} = \tilde{O}(x) + C, \text{ where } C = -\frac{2F_1^2\alpha}{\psi_1 - \psi_2} \quad (147)$$

$$\text{and } \tilde{O}(x) = -\frac{2\psi_2(3x^2((F_\star^2 + \tau^2) - F_1^2(\psi_2 - 1)) - 2x^3F_1^2 + 6x(F_\star^2 + \tau^2)\psi_2 + \psi_2(F_1^2(\psi_2 - 1)^2 + (F_\star^2 + \tau^2)(3\psi_2 + 1)))}{((x + \psi_2)^2 - \psi_2)^3}.$$

A tedious calculation (omitted) shows that $\frac{d^2\tilde{O}}{dx^2} \geq 0$ for $x \in [x_L, x_R]$ (i.e. it is convex), that $\tilde{O}(x_L) < \tilde{O}(x_R)$, and that $\frac{d\tilde{O}}{dx}(x = x_L) < 0$. To do this calculation, we recommend the following. First break \tilde{O} into two components. One component proportional to F_1^2 , called \tilde{O}_1 , and one component proportional to $(F_\star^2 + \tau^2)$, called \tilde{O}_\star . Then, show that both components are convex, that $\tilde{O}_\star(x_L) \leq \tilde{O}_\star(x_R)$, that $\tilde{O}_1(x_L) < \tilde{O}_1(x_R)$, that $\frac{d\tilde{O}_\star}{dx}(x = x_L) = 0$, and that $\frac{d\tilde{O}_1}{dx}(x = x_L) < 0$.

From this it follows that, depending on the value of ψ_1 , the concavity $\frac{d^2O}{dx^2}$ is positive or negative. The situation is exactly the same as in the Figure 3 but now the x axis is ψ_1 , the points A_1, A_2 and A_3 are different, and so are the definitions of β_1, β_2 and β_3 . We do however have that, by definition, $0 \leq \beta_3 < \beta_2 < \beta_1$.

With a slight abuse of notation we refer to the y -coordinate value of this points by A_1, A_2 , and A_3 . We now have $A_1 = \frac{2(F_1^2(\psi_2 + 1) + (F_\star^2 + \tau^2))}{\psi_2}$ and $A_3 = -\frac{2\max\{1, \psi_2\}(F_1^2(\psi_2 - 1)^2 + (F_\star^2 + \tau^2)(3\min\{1, \psi_2\} + \max\{1, \psi_2\}))}{(\psi_2 - 1)^3 \min\{1, \psi_2\}}$. Although we are not considering yet the case when $\psi_2 = 1$, notice that when $\psi_2 = 1$ we have that $A_3 = +\infty$, and in fact we also have $O(x_R) = +\infty$. Getting A_2 is a bit more complicated. From the first order condition $\frac{d\tilde{O}}{dx} = 0$ and the convexity of \tilde{O} – recall that \tilde{O} a rational function – we can extract that the x -coordinate of A_2 is the unique root of a 4th degree polynomial $h(x) = h_0 + h_1x + h_2x^2 + h_3x^3 + h_4x^4$ in the range $x \in [x_L, x_R]$, where $h_0 = \psi_2^2(\rho(\psi_2 - 1)^2 + 2(\psi_2 + 1))$, $h_1 = 2\psi_2(\rho(\psi_2 - 1)^2 + (3\psi_2 + 1))$, $h_2 = 6\psi_2$, $h_3 = 2(1 - \rho(\psi_2 - 1))$, $h_4 = -\rho$. Call this root x_{A_2} . We then have $A_2 = \tilde{O}(x_{A_2})$. It turns out that we can write A_2 directly as the solution of $g(A_2)/((\tau^2 + F_\star^2)\psi_2) = 0$, where $g(x) = g_0 + g_1x + g_2x^2 + g_3x^3 + g_4x^4$ with $g_0 = \rho^4$, $g_1 = -4(\rho\psi_2 + \rho + 1)(\rho\psi_2(2\rho\psi_2 - 3\rho + 4) + 2(\rho + 1)^2)$, $g_2 = -4\psi_2^2(\rho\psi_2(7\rho\psi_2 - 20\rho + 14) + 7(\rho + 1)^2)$, $g_3 = -16\psi_2^4(\rho\psi_2 + \rho + 1)$, and $g_4 = 16\psi_2^6$.

Using the expressions for A_1 and A_3 , and the expression for C , see (147), β_2 and β_3 are defined as $\beta_2 = \psi_1 : A_1(\psi_1) + C(\psi_1) = 0$, and $\beta_3 = \psi_1 : A_3(\psi_1) + C(\psi_1) = 0$. By definition of β_2 and β_3 ,

these equations have a unique solution when $\psi_1 > \psi_2 > 0$ and their expressions are given in (38) and (39) respectively. We can also define β_1 as the solution of $\beta_1 = \psi_1 : A_2(\psi_1) + C(\psi_1) = 0$. By definition of β_1 , the solution is unique in the range $\psi_1 > \psi_2 > 0$. We can use the fact that $A_2 = \tilde{O}(x_{A_2})$ to write that $\beta_1 = \psi_2 + \frac{2F_1^2 \alpha}{\tilde{O}(x_{A_2})}$. We can also use the fact that $g(A_2 / ((\tau^2 + F_\star^2)\psi_2)) = 0$, which implies that $g(-C(\psi_1) / ((\tau^2 + F_\star^2)\psi_2)) = 0$ when $\psi_1 = \beta_1$, to write β_1 as the root of a 4th degree polynomial $r(x)$ that is specified in Appendix B.4, which is the way in which we decide to state Theorem 9.

Now that we have characterized the curvature of $O(x)$, we are ready to locate its global minimum. To do so, we will use the curvature of $O(x)$ together with the first-order optimality condition $\frac{dO}{dx} = 0$, and the following three extra pieces of information: the sign of the derivative of $O(x)$ at $x = x_L$; the sign of the derivative of $O(x)$ at $x = x_R$; and the sign of $O(x_L) - O(x_R)$.

A direct computation yields

$$\frac{dO}{dx} = \frac{p_0 + p_1x + p_2x^2 + p_3x^3 + p_4x^4 + p_5x^5}{(\psi_2 - (x + \psi_2)^2)(\psi_1 - \psi_2)}, \quad (148)$$

where the coefficients p_0, \dots, p_5 are, apart from a multiplying constant, given in (68)-(73). The roots of the denominator, i.e. $-\sqrt{\psi_2 - \psi_2}$ and $\sqrt{\psi_2 - \psi_2}$, are not a root of the numerator, therefore, the first order optimality conditions are $p(x) = 0$.

Not all solutions of $p(x) = 0$ minimize $O(x)$. To locate the minimizer, we use the sign of the derivative of $O(x)$ at $x = x_L$; the sign of the derivative of $O(x)$ at $x = x_R$; and the sign of $O(x_L) - O(x_R)$. These signs can be determined using Lemma D.14. Lemma D.14 also proves Remark 10.

Lemma D.14. *If $\psi_1 > \psi_2$ then the following relationships hold.*

$$\left. \frac{dO}{dx} \right|_{x=x_L} = \frac{\alpha (F_\star^2 + \tau^2) + F_1^2 (\alpha (-\psi_2) + 2(\alpha - 1)\psi_1 + \alpha + \psi_2)}{\psi_1 - \psi_2}. \quad (149)$$

$$\text{If } \psi_2 = 1 \text{ then } \left. \frac{dO}{dx} \right|_{x=x_R} = +\infty. \quad (150)$$

$$\text{If } \psi_2 < 1 \text{ then} \quad (151)$$

$$\left. \frac{dO}{dx} \right|_{x=x_R} = \frac{(\alpha\psi_2^2 - 2(\alpha+1)\psi_2 + \alpha + 2\psi_1) (F_\star^2 + \tau^2) + F_1^2 (\psi_2 - 1)^2 (2\alpha\psi_1 - (3\alpha+1)\psi_2 + \alpha)}{(\psi_1 - \psi_2) (\psi_2 - 1)^2}. \quad (152)$$

$$\text{If } \psi_2 > 1 \text{ then} \quad (153)$$

$$\left. \frac{dO}{dx} \right|_{x=x_R} = \frac{F_1^2 (\psi_2 - 1)^2 (-2\alpha\psi_1 + (\alpha+1)\psi_2 + \alpha) - (\psi_2 ((\alpha-2)\psi_2 - 2\alpha + 2\psi_1) + \alpha) (F_\star^2 + \tau^2)}{(\psi_2 - 1)^2 (\psi_2 - \psi_1)}. \quad (154)$$

$$\text{If } \psi_2 = 1 \text{ then } O(x_L) - O(x_R) = -\infty. \quad (155)$$

$$\text{If } \psi_2 < 1 \text{ then} \quad (156)$$

$$O(x_L) - O(x_R) = \frac{\psi_2 (F_1^2 (\psi_2 - 1) ((1-2\alpha)\psi_1 + \alpha (2\psi_2 - 1)) - ((\alpha+1)\psi_2 - \alpha - \psi_1) (F_\star^2 + \tau^2))}{(\psi_1 - \psi_2) (\psi_2 - 1)}. \quad (157)$$

$$\text{If } \psi_2 > 1 \text{ then} \quad (158)$$

$$O(x_L) - O(x_R) = \quad (159)$$

$$\frac{\psi_2 ((\alpha-1) (F_\star^2 + \tau^2) + \alpha F_1^2) + \psi_1 ((2\alpha-1) F_1^2 (\psi_2 - 1) + F_\star^2 + \tau^2) - \alpha (F_\star^2 + \tau^2) - \alpha F_1^2 \psi_2^2}{(\psi_2 - 1) (\psi_2 - \psi_1)}. \quad (160)$$

Also, the following is true.

$$\text{If } \alpha < \alpha_L \text{ then } \left. \frac{dO}{dx} \right|_{x=x_L} < 0; \text{ and if } \alpha \geq \alpha_L \text{ then } \left. \frac{dO}{dx} \right|_{x=x_L} \geq 0. \quad (161)$$

$$\text{Let } \psi_2 \neq 1. \text{ If } E_1 \text{ is true then } \left. \frac{dO}{dx} \right|_{x=x_R} < 0; \text{ and if } E_2 \text{ is true then } \left. \frac{dO}{dx} \right|_{x=x_R} > 0. \quad (162)$$

Conditions E_1 and E_2 are defined in Theorem 9.

Let $\psi_2 \neq 1$. If $\alpha < \alpha_C$ then $O(x_L) - O(x_R) > 0$; and if $\alpha \geq \alpha_C$ then $O(x_L) - O(x_R) \leq 0$ (163)

Constant α_C is defined in equation (41) in Theorem 9.

Furthermore, we always have that $\alpha_L \in [0, 1]$. It also holds that $\alpha_C \in [0, 1]$ if and only if $\psi_1 \leq \psi_2/(1 - \rho|1 - \psi_2|)$. We have that $\alpha_R \in [0, 1]$ if and only if

$$\psi_1 \geq \psi_2 + \frac{1}{2}\rho(\psi_2 - 1)^2 \min\{\psi_2, 1\} \text{ and} \quad (164)$$

$$\psi_1 \leq \frac{-1 + \rho(\psi_2 - 1)^2(2\psi_2 + \min\{2\psi_2 - 1, 1\}) + \psi_2(2 + \max\{\psi_2, 2 - \psi_2\})}{2(\max\{\psi_2, 1\} + \rho(\psi_2 - 1)^2)}. \quad (165)$$

Finally, $A \geq B$ always, where A and B are given in (43) and (44). Furthermore, $B < \psi_2$ if and only if $1/\rho > \gamma \triangleq \min\{1, \max\{0, 2\psi_2 - 1\}\}$.

Proof. Except (155) and (150), the derivation of (149)-(160) follows from direct substitution of $x = x_L$ or $x = x_R$ into O – for which we have expressions from Theorems 1 and 4 – or its derivative in equation 148.

Equation (155), for when $\psi_2 = 1$, is obtained by taking the limit of (157) and (160) as $\psi_2 \uparrow 1$ and $\psi_2 \downarrow 1$ respectively. Equation (150) for when $\psi_2 = 1$ is obtained by taking the limit of (152) and (154) as $\psi_2 \uparrow 1$ and $\psi_2 \downarrow 1$ respectively.

The derivation of condition (161) follows from the observation that the equation (149) is linear in α and is always negative for $\alpha = 0$ and positive for $\alpha = 1$. Hence, we can easily compute a value $\alpha_L \in [0, 1]$ at which the expression changes from a negative value to a positive value. The expression for α_L we obtain is (40).

The derivation of condition (162) can be obtained through following the observations. First notice that $\frac{dO}{dx}|_{x=x_R}$ is a linear increasing function of α . Now focus on the following three implications.

1. If at $\alpha = 1$ we have $\frac{dO}{dx}|_{x=x_R} < 0$, then $\frac{dO}{dx}|_{x=x_R} < 0$ for all $\alpha \in [0, 1]$;
2. If at $\alpha = 0$ we have $\frac{dO}{dx}|_{x=x_R} > 0$, then $\frac{dO}{dx}|_{x=x_R} > 0$ for all $\alpha \in [0, 1]$;
3. If at $\alpha = 0$ we have $\frac{dO}{dx}|_{x=x_R} < 0$ and at $\alpha = 1$ we have $\frac{dO}{dx}|_{x=x_R} > 0$, then there exists $\alpha_R \in [0, 1]$ such that $(\alpha < \alpha_R \Rightarrow \frac{dO}{dx}|_{x=x_R} < 0)$ and $(\alpha > \alpha_R \Rightarrow \frac{dO}{dx}|_{x=x_R} > 0)$;

A direct computation shows that the sufficient condition in the first implication holds if and only if $\psi_1 < B$, where B is given in (44); the sufficient condition in the second implication holds if and only if $\psi_1 > A$, where A is given in (43); the sufficient condition in the the third implication holds if and only if $B < \psi_1 < A$. Therefore, if $E_1 = (\psi_1 < B) \vee ((\alpha < \alpha_R) \wedge (B < \psi_1 < A))$ is true, we can use the first or third implication to conclude that $\frac{dO}{dx}|_{x=x_R} < 0$. Also, if $E_2 = (\psi_1 > A) \vee ((\alpha > \alpha_R) \wedge (B < \psi_1 < A))$ is true, we can use the second or third implication to conclude that $\frac{dO}{dx}|_{x=x_R} > 0$.

A direct calculation shows that $A \geq B$ and that $B < \psi_2$ if and only if $1/\rho > \gamma \triangleq \min\{1, \max\{0, 2\psi_2 - 1\}\}$. A direct calculation also shows that $\alpha_R \in [0, 1]$ if and only if the conditions in (164) and (165) hold.

Condition (163) can be obtained through following the observations. First, notice that both (157) and (160) are linear decreasing functions of α . Second, when $\alpha = 1$ we always have $O(x_L) - O(x_R) < 0$. Therefore, there exists $\alpha_C \leq 1$ such that $(\alpha < \alpha_C \Rightarrow O(x_L) - O(x_R) > 0)$ and $(\alpha > \alpha_C \Rightarrow O(x_L) - O(x_R) < 0)$. The expression for α_C is given by (41). From this expression, a direct calculation shows that $\alpha_C \geq 0$ if and only if $\psi_1 \leq \psi_2/(1 - \rho|1 - \psi_2|)$. \square

To finish the proof of Theorem 9 for $\psi_1 > \psi_2$ we consider the different scenarios in Table 1. These are studied via cases that are exactly the same to the Case 1.1 to Case 1.14 for $\psi_1 < \psi_2$ but with $\alpha < \alpha_R$ replaced by E_1 and $\alpha > \alpha_R$ replaced by E_2 because when $\psi_1 > \psi_2$ it is E_1 and E_2 that determine the sign of $\frac{dO}{dx}|_{x=x_R}$. For example, for the top left-most cell in Table 1, both when $\psi_1 < \psi_2$

or $\psi_1 > \psi_2$, we have that O is convex and its decreasing at x_L and x_R , so its minimum is at $x = x_R$. We omit repeating the arguments for Case 1.2 to Case 1.14. \square

D.6 Proof of Theorem 10

Proof of Theorem 10. The proof amounts to a long calculus exercise, which we shorten by some careful observations.

We first notice that $\mathcal{E}_{R_2}^\infty$ and $\mathcal{S}_{R_2}^\infty$ can both be written as function of ω_2 , and we can use this to reduce the optimization problem (29) to an optimization problem over just one variable.

The variable ω_2 is a function of $\mu_0, \mu_1, \mu_2 \geq 0$, and has range $[-\infty, 0]$, the value $-\infty$ being achieved when $\mu_\star^2 = \mu_2 - \mu_1^2 - \mu_2^2 = 0$.

To avoid having to deal with infinities, we make use of the Mobius transformation $x = \frac{1+\omega_2}{\omega_2-1}$, and instead work with x .

If we substitute $x = \frac{1+\omega_2}{-1+\omega_2}$ into the left hand side (45), and substitute (16) in the resulting expression we confirm that (45) is satisfied.

Furthermore, if we use $x = \frac{1+\omega_2}{-1+\omega_2}$ and (16) to write x as a function of μ_0, μ_1, μ_2 , we can use the fact that $\mu_0, \mu_1, \mu_2 \geq 0$, to conclude that $x \in [-1, \min\{1, -1 + 2\psi_2\}]$.

Our problem is thus equivalent to solving $\min_x (1 - \alpha)\mathcal{E}_{R_2}^\infty + \alpha\mathcal{S}_{R_2}^\infty$ subject to $x \in [-1, \min\{1, -1 + 2\psi_2\}]$. Once we know x , any μ_0, μ_1, μ_2 that satisfies (45) will be a minimizer.

At this point we compute $\frac{d}{dx} [(1 - \alpha)\mathcal{E}_{R_2}^\infty + \alpha\mathcal{S}_{R_2}^\infty]$ and observe that this is a rational function of x . The numerator is, apart from a multiplying constant, $p(x)$, and the denominator is zero if and only if $x = -1 - 2\sqrt{\psi_2}$ or $x = -1 + 2\sqrt{\psi_2}$. Both are outside of the range of x unless $\psi_2 = x = 1$. We treat this case separately soon. Since the zeros of the numerator and of the denominator never cancel out, all of the critical points are given by $p(x) = 0$.

Now we compute the value of the derivative at the extremes of the range of x , namely, $x = -1, 1$, or $-1 + 2\psi_2$. The value of the derivative at $x = -1$ is $F_1^2(-1 + \alpha) < 0$, which implies that $x = -1$ is not a minimizer. The value of the derivative at $x = 1$ is $\alpha F_1^2 + \frac{\psi_2(F_\star^2 + \tau^2)}{(\psi_2 - 1)^2} > 0$, possibly $+\infty$, which implies that $x = 1$ is not a minimizer. The value of the derivative at $x = -1 + 2\psi_2$ is $\alpha F_1^2 + \frac{F_\star^2 + \tau^2}{(\psi_2 - 1)^2} > 0$, possibly $+\infty$, which implies that $x = -1 + 2\psi_2$ is not a minimizer. This calculation implies that we can assume that $x \in (-1, \min\{1, -1 + 2\psi_2\})$, which we will assume from now on.

Finally, we show that the objective is convex in the domain of x , which implies that there is only one critical point – that is $p(x) = 0$ has only one solution in the domain $(-1, \min\{1, -1 + 2\psi_2\})$ – and that this critical point is a global minimum. To show that the objective is convex, we compute its second derivative, which is

$$\frac{8\psi_2 (F_1^2 (4\psi_2^2 + \psi_2(3(x-2)x-5) - x^3 + 3x + 2) + (F_\star^2 + \tau^2) (4\psi_2 + 3(x+1)^2))}{(4\psi_2 - (x+1)^2)^3}. \quad (166)$$

The minimum of denominator for $x \in [-1, \min\{1, -1 + 2\psi_2\}]$ is $\begin{cases} 4(\psi_2 - 1) & , \psi_2 > 1 \\ -4(\psi_2 - 1)\psi_2 & , 0 \leq \psi_2 \leq 1 \end{cases}$, which is always non-negative, and is zero only if $x = 1$, or $x = -1 + 2\psi_2$, which have already been excluded because they are not minimizers. Hence, for $x \in (-1, \min\{1, -1 + 2\psi_2\})$, the denominator is strictly positive.

To show that the numerator is non-negative, we only need to show that $4\psi_2^2 + \psi_2(3(x-2)x-5) - x^3 + 3x + 2 \geq 0$ in the range of x . The minimum of $4\psi_2^2 + \psi_2(3(x-2)x-5) - x^3 + 3x + 2$ for $x \in [-1, \min\{1, -1 + 2\psi_2\}]$ is $\begin{cases} 4(\psi_2 - 1)^2 & , \psi_2 > 1 \\ 4(\psi_2 - 1)^2\psi_2 & , 0 \leq \psi_2 \leq 1, \end{cases}$ which is always non-negative. \square

D.7 Proof of Theorem 11

Proof of Theorem 11. Similar to proof of Theorem 10, the majority of the proof is a long calculus exercise.

We first notice that $\mathcal{E}_{R_3}^\infty$ and $\mathcal{S}_{R_3}^\infty$ can both be written as function of ω_1 and ζ^2 .

The variable ω_1 is a function of $\mu_0, \mu_1, \mu_2 \geq 0$, and has range $[-\infty, 0]$, the value $-\infty$ being achieved when $\mu_\star^2 = \mu_2 - \mu_1^2 - \mu_2^2 = 0$.

To avoid having to deal with infinities, we make use of the Mobius transformation $x = \frac{1+\omega_1}{\omega_1-1}$, and instead work with x . If we use $x = \frac{1+\omega_1}{\omega_1-1}$ and (18) to write x as a function of μ_0, μ_1, μ_2 , we can use the fact that $\mu_0, \mu_1, \mu_2 \geq 0$, to conclude that $x \in [-1, \min\{1, -1 + 2\psi_1\}]$.

Our problem is thus equivalent to solving $\min_x (1 - \alpha)\mathcal{E}_{R_3}^\infty + \alpha\mathcal{S}_{R_3}^\infty$ subject to $x \in [-1, \min\{1, -1 + 2\psi_1\}]$ and $\zeta^2 \geq 0$.

Let $O = (1 - \alpha)\mathcal{E}_{R_3}^\infty + \alpha\mathcal{S}_{R_3}^\infty$. With a direct calculation we can check the following. The derivative $\frac{dO}{dx}|_{x=-1}$ is always negative which implies that there is no local minimum at $x = -1$. The derivatives $\frac{dO}{dx}|_{x=1}$ and $\frac{dO}{dx}|_{x=-1+2\psi_1}$ are always positive, which implies that there is no local minimum at either $x = 1$ or $x = -1 + 2\psi_1$. Therefore, we know that the minimizer has $x \in (-1, \min\{1, -1 + 2\psi_1\})$. For $x \in (-1, \min\{1, -1 + 2\psi_1\})$, the derivative $\frac{dO}{d(\zeta^2)}|_{\zeta^2=0}$ is always positive, which implies that there is no local minimum at $\zeta^2 = 0$. We thus know that the minimizer of O is in the interior of the domain and hence it can be found via $\nabla O = 0$.

The remainder of the proof considers a few different cases depending on the value of α and ψ_1 .

Case when $\alpha = 0$:

In this case $O = \mathcal{E}_{R_3}^\infty$ and $\frac{dO}{d(\zeta^2)} = \frac{F_1^2(x+1)^2}{\zeta^4(x+1)^2-4\psi_1}$. The only way that $\frac{dO}{d(\zeta^2)} = 0$ is if $\zeta^2 = \infty$, which corresponds to $\mu_\star = 0$.

As we noted in the beginning, O is a function of ω_1 and ζ^2 , and since ω_1 is a function ζ^2 and μ_1^2 and ζ^2 is a function of μ_1^2 and μ_\star^2 , we know that O is a function of μ_1^2 and μ_\star^2 . We express O in these variables and compute $\frac{dO}{d(\mu_1^2)}$ when $\mu_\star \rightarrow 0$. We get that

$$\frac{dO}{d(\mu_1^2)} = -\frac{2F_1^2\lambda^2\psi_1^3}{(2\mu_1^2\psi_1(\lambda - \mu_1^2) + \psi_1^2(\lambda + \mu_1^2)^2 + \mu_1^4)^{3/2}}. \quad (167)$$

By minimizing the denominator with respect to $\mu_1 \geq 0$, we conclude that its minimum is $\lambda^2\psi_1^2 > 0$, which implies that $\frac{dO}{d(\mu_1^2)} < 0$, which implies that $\mu_1 \rightarrow \infty$.

In this case, if we express O as a function of μ_1 and μ_\star and take $\mu_1 \rightarrow \infty$ and $\mu_\star \rightarrow 0$, we get $O \rightarrow F_s^2$.

Case when $\psi_1 = 1 \wedge 0 < \alpha \leq \frac{1}{4}$:

In this case $\frac{dO}{d(\zeta^2)} = \frac{(\alpha-1)F_1^2(x+1)^2}{(x-1)(x+3)\zeta^4}$. The only way that $\frac{dO}{d(\zeta^2)} = 0$ is if $\zeta^2 = \infty$, which corresponds to $\mu_\star = 0$.

As we noted in the beginning, O is a function of ω_1 and ζ^2 , and since ω_1 is a function ζ^2 and μ_1^2 and ζ^2 is a function of μ_1^2 and μ_\star^2 , we know that O is a function of μ_1^2 and μ_\star^2 . We express O in these variables and compute $\frac{dO}{d(\mu_1^2)}$ when $\mu_\star \rightarrow 0$. We get that

$$\frac{dO}{d(\mu_1^2)} = \frac{2F_1^2\lambda^2 \left(\frac{4\alpha\lambda(\lambda+4\mu_1^2)}{(\sqrt{\lambda(\lambda+4\mu_1^2)+\lambda})^2} - 1 \right)}{(\lambda(\lambda+4\mu_1^2))^{3/2}}. \quad (168)$$

By maximizing the numerator with respect to $\mu_1^2 \geq 0$, we conclude that its value is always strictly smaller than $2F_1^2\lambda^2(-1+4\alpha) \leq 0$. Hence, $\frac{dO}{d(\mu_1^2)} < 0$, which implies that the optimum is achieved for $\mu_1 \rightarrow \infty$.

In this case, if we express O as a function of μ_1 and μ_\star and take $\mu_1 \rightarrow \infty$ and $\mu_\star \rightarrow 0$, we get $O \rightarrow \alpha F_1^2 + (1 - \alpha) F_s^2$.

Case when $\psi = 1 \wedge \frac{1}{4} < \alpha < 1$:

This case is very similar to the previous case. The only difference being that, because $\frac{1}{4} < \alpha < 1$, when we solve

$$\frac{dO}{d(\mu_1^2)} = \frac{2F_1^2 \lambda^2 \left(\frac{4\alpha\lambda(\lambda+4\mu_1^2)}{(\sqrt{\lambda(\lambda+4\mu_1^2)}+\lambda)^2} - 1 \right)}{(\lambda(\lambda+4\mu_1^2))^{3/2}} = 0, \quad (169)$$

we now get two possible solutions, namely,

$$\mu_1^2 \in \left\{ \frac{-4\alpha^2\lambda + 3\alpha\lambda + (-\sqrt{\alpha})\lambda}{16\alpha^2 - 8\alpha + 1}, \frac{-4\alpha^2\lambda + 3\alpha\lambda + \sqrt{\alpha}\lambda}{16\alpha^2 - 8\alpha + 1} \right\}. \quad (170)$$

First notice that if $\lambda = 0$, both expressions give $\mu_1 = 0$. Let us assume now that $\lambda > 0$. If we maximize the first expression with respect to $\frac{1}{4} < \alpha < 1$, we conclude that its value is always smaller than $-((3\lambda)/16) < 0$, which implies that it is not a valid solution in the range $\mu_1^2 \geq 0$. If we minimize the second expression with respect to $\frac{1}{4} < \alpha < 1$, we conclude that its value is always non-negative, which implies it is a valid solution in the range $\mu_1^2 \geq 0$. Given that this is the only stationary point of O and given that O is bounded from below it must be a global minimum.

If we substitute the optimal values for μ_1^2 and $\mu_\star^2 = 0$ in O , we get $O = (4\sqrt{\alpha} - 1 - 3\alpha) F_1^2 + (1 - \alpha) F_s^2$.

Case when $\psi_1 \neq 1$:

In this case $\frac{dO}{d(\zeta^2)} = \frac{(\alpha-1)F_1^2(x+1)^2}{\zeta^4(4\psi_1-(x+1)^2)}$. The only way that $\frac{dO}{d(\zeta^2)} = 0$ is if $\zeta^2 = \infty$, which corresponds to $\mu_\star = 0$.

If we substitute $x = \frac{1+\omega_1}{\omega_1-1}$ into the left hand side of the last equality in (51), namely, $\mu_1^2(-1 + 2\psi_1 - x)(-1 + x) + 2\lambda\psi_1(1 + x)$, and let $\mu_\star \rightarrow 0$, we get 0, which confirms the condition on x when $\psi_1 \neq 1$.

We take $\zeta \rightarrow \infty$ in O to obtain,

$$O = F_1^2 \left(\frac{\psi_1(x-1)^2}{4\psi_1 - (x+1)^2} + \alpha x \right) - (\alpha-1)F_\star^2, \quad (171)$$

$$\frac{dO}{dx} = F_1^2 \left(\alpha + \frac{4\psi_1(x-1)(2\psi_1-x-1)}{((x+1)^2 - 4\psi_1)^2} \right), \quad (172)$$

$$\frac{d^2O}{dx^2} = \frac{8F_1^2\psi_1(4\psi_1^2 + \psi_1(3(x-2)x-5) - x^3 + 3x + 2)}{(4\psi_1 - (x+1)^2)^3}. \quad (173)$$

If we minimize $\frac{d^2O}{dx^2}$ over $x \in [-1, \min(1, -1 + 2\psi_1)]$ and $\psi_1 \geq 0$, we obtain $F_1^2/8 > 0$. This shows that our objective O is strictly convex in the range $x \in [-1, \min(1, -1 + 2\psi_1)]$ and hence there is only one solution to $\frac{dO}{dx} = 0$.

The rational function $\frac{dO}{dx}$ has a denominator which is zero only if $x = -1 - 2\sqrt{\psi_1}$ or $x = -1 + 2\sqrt{\psi_1}$, both of which are outside the valid range for x . Hence the unique solution to $\frac{dO}{dx} = 0$ comes from the zeros of the numerator of $\frac{dO}{dx}$. This numerator is (a constant times) a polynomial in x whose coefficients are described in Theorem 11.

□