

SUPERCYCLIC VECTORS OF OPERATORS ON NORMED LINEAR SPACES

MOHAMMAD ANSARI

ABSTRACT. We give an affirmative answer to a question asked by Faghih-Ahmadi and Hedayatian regarding supercyclic vectors. We show that if \mathcal{X} is an infinite-dimensional normed linear space and T is a supercyclic operator on \mathcal{X} , then for any supercyclic vector x for T , there exists a strictly increasing sequence $(n_k)_k$ of positive integers such that the closed linear span of the set $\{T^{n_k}x : k \geq 1\}$ is not the whole \mathcal{X} .

1. Introduction

Let \mathcal{X} be an infinite-dimensional normed linear space and $B(\mathcal{X})$ be the space of all bounded linear operators on \mathcal{X} . An operator $T \in B(\mathcal{X})$ is said to be *supercyclic* if there is some $x \in \mathcal{X}$ such that the set $\{cT^n x : c \in \mathbb{C}, n \geq 0\}$ is dense in \mathcal{X} . In this case, x is called a *supercyclic vector* for T . To see detailed information about supercyclic operators we refer the readers to [2] and [4].

For a subset M of \mathcal{X} , by $\bigvee M$ we mean the closed linear span of the set M . It is clear that if x is a supercyclic vector for an operator $T \in B(\mathcal{X})$, then $\bigvee\{T^n x : n \geq 0\} = \mathcal{X}$. A natural question which may be asked here is that whether there is a strictly increasing sequence $(n_k)_k$ of positive integers such that $\bigvee\{T^{n_k} x : k \geq 1\} \neq \mathcal{X}$. In their recently published paper, Faghih-Ahmadi and Hedayatian have proved the following interesting result.

Theorem 1.1 (Theorem 1 of [3]). *Let \mathcal{H} be an infinite-dimensional Hilbert space. If x is a supercyclic vector for $T \in B(\mathcal{H})$, then there is a (strictly increasing) sequence $(n_k)_k$ of positive integers such that $\bigvee\{T^{n_k} x : k \geq 1\} \neq \mathcal{H}$.*

Then they have asked whether the assertion is true for locally convex spaces or at least for Banach spaces [3, Question 1]. In this note, we answer

2020 *Mathematics Subject Classification.* Primary 47A16; Secondary 46B20.

Key words and phrases. supercyclic operator; supercyclic vector; normed linear space.

their question affirmatively for normed linear spaces. To prove our result, we use Lemma 1.3 which is an analogue of the following lemma.

Lemma 1.2 (Lemma 2.3 of [1]). *Let \mathcal{A} be a dense subset of a Banach space \mathcal{X} and e be a fixed element with $\|e\| > 1$. Then, for every finite-dimensional subspace $Y \subset \mathcal{X}$ with $\text{dist}(e, Y) > 1$, for every $\epsilon > 0$ and $y \in Y$, there is an $a \in \mathcal{A}$ such that $\|y - a\| < \epsilon$ and $\text{dist}(e, \bigvee\{Y, a\}) > 1$.*

The proof of the above lemma shows that it can also be stated for infinite-dimensional normed linear spaces. Thus, we can give the following modified version.

Lemma 1.3. *Let \mathcal{A} be a dense subset of an infinite-dimensional normed linear space \mathcal{X} and $e \in \mathcal{X}$ be a fixed element with $\|e\| > 1$. Then, for every finite-dimensional subspace Y of \mathcal{X} with $\text{dist}(e, Y) > 1$, there is an $a \in \mathcal{A}$ such that $\text{dist}(e, \bigvee\{Y, a\}) > 1$.*

2. Main result

Now, we are ready to answer Question 1 of [3] for normed linear spaces.

Theorem 2.1. *Let \mathcal{X} be an infinite-dimensional normed linear space. If x is a supercyclic vector for $T \in B(\mathcal{X})$, then there is a strictly increasing sequence $(n_k)_k$ of positive integers such that $\bigvee\{T^{n_k}x : k \geq 1\} \neq \mathcal{X}$.*

Proof. It is clear that every nonzero scalar multiple of x is also a supercyclic vector for T . On the other hand, since x and Tx are linearly independent vectors, we have $\text{dist}(x, \bigvee\{Tx\}) > 0$. Thus, without loss of generality, we can assume that $\|x\| > 1$ and $\text{dist}(x, \bigvee\{Tx\}) > 1$. Therefore, in view of Lemma 1.3, if we put

$$\mathcal{A} = \{cT^n x : c \in \mathbb{C}, n > 1\}, Y = \bigvee\{Tx\},$$

and $e = x$, then there is some $a = cT^{n_2}x \in \mathcal{A}$ such that

$$\text{dist}(x, \bigvee\{Tx, T^{n_2}x\}) = \text{dist}(x, \bigvee\{Y, a\}) > 1.$$

Now, let

$$\mathcal{A}_2 = \{cT^n x : c \in \mathbb{C}, n > n_2\}, Y_2 = \bigvee\{Tx, T^{n_2}x\}.$$

Then there is some $a_2 = c_2T^{n_3}x \in \mathcal{A}_2$ such that

$$\text{dist}(x, \bigvee\{Tx, T^{n_2}x, T^{n_3}x\}) = \text{dist}(x, \bigvee\{Y_2, a_2\}) > 1.$$

By continuing this construction, assume that for some $k \geq 2$, the dense set \mathcal{A}_k and the finite-dimensional subspace Y_k have been presented and (by using Lemma 1.3) we have found an element $a_k = c_k T^{n_{k+1}} x \in \mathcal{A}_k$ such that

$$\text{dist}(x, \bigvee \{Tx, T^{n_2}x, \dots, T^{n_{k+1}}x\}) = \text{dist}(x, \bigvee \{Y_k, a_k\}) > 1.$$

Then we put

$$\mathcal{A}_{k+1} = \{cT^n x : c \in \mathbb{C}, n > n_{k+1}\}, Y_{k+1} = \bigvee \{Tx, T^{n_2}x, \dots, T^{n_{k+1}}x\}.$$

Again, by Lemma 1.3, there is some $a_{k+1} = c_{k+1} T^{n_{k+2}} x \in \mathcal{A}_{k+1}$ such that

$$\text{dist}(x, \bigvee \{Tx, T^{n_2}x, \dots, T^{n_{k+2}}x\}) = \text{dist}(x, \bigvee \{Y_{k+1}, a_{k+1}\}) > 1.$$

This inductive procedure gives a strictly increasing sequence $(n_k)_k$ (with $n_1 = 1$) and it is easily seen that

$$\text{dist}(x, \bigvee \{T^{n_k}x : k \geq 1\}) \geq 1$$

(see the last paragraph of the proof of Theorem 1 of [3]). This shows that $x \notin \bigvee \{T^{n_k}x : k \geq 1\}$ and we are done. \square

The interested readers are invited to investigate Question 1 of [3] for operators on locally convex spaces.

REFERENCES

- [1] N. Bamerni, V. Kadets, and A. Kilicman, *Hypercyclic operators are subspace hypercyclic*, J. Math. Anal. Appl., **435** (2016), 1812-1815.
- [2] F. Bayart and E. Matheron, *Dynamics of linear operators*, Cambridge University Press, **179**, 2009.
- [3] M. Faghih-Ahmadi and K. Hedayatian, *A note on supercyclic vectors of Hilbert space operators*, J. Math. Anal. Appl., **505** (2022).
- [4] K.-G. Grosse-Erdmann and A. Peris Manguillot, *Linear chaos*, Universitext, Springer-Verlag London Limited, 2011.

Department of Basic Sciences

Azad University of Gachsaran

Gachsaran, Iran

Email Address: mansari@iaug.ac.ir, ansari.moh@gmail.com