

# FIBRATIONS PROPERTY OF EMBEDDING MAPS OF ORBIFOLD CHARTS

Hakeem A. Othman<sup>1</sup> and Santanu Acharjee<sup>2</sup>

<sup>1</sup>Department of Mathematics, AL-Qunfudhah University college, Umm Al-Qura University, KSA.

<sup>1</sup>Department of Mathematics, Rada'a College of Education and Science, Albaydha University, Albaydha, Yemen.

<sup>2</sup>Department of Mathematics, Gauhati University, Guwahati-781014, Assam, India.

e-mails: <sup>1</sup>haoali@uqu.edu.sa, hakim\_albdoie@yahoo.com

<sup>2</sup>sacharjee326@gmail.com

## Abstract

In this paper we introduce the notion of Hurewicz fibrations in the class of embedding maps of orbifold charts by giving the concept of E-fibration embedding. We study the fundamental properties of this concept such as the restriction, product and its relationship with Hurewicz fibration, etc. Furthermore, we introduce the notion of lifting functions of E-fibration embedding and study preserving projection property of these lifting functions.

*Keywords:* Orbifold; embedding; fibration; homotopy.

*2020 AMS classifications:* 55P05, 57R18, 57N35, 14E25, 32C05.

## 1 Introduction

The concept of orbifold has been described using various mathematical constructions and contexts. It was first introduced by Satake [8, 9]. He called them V-manifolds. He defined orbifolds as topological spaces with an atlas of charts. He viewed orbifolds as a generalization of manifolds. Orbifolds and manifolds are described by charts. An orbifold chart under any space  $X$  is defined as a triple  $(X_U, G_U, \Gamma_U)$ , where  $X_U$  is open set in  $R^n$ ,  $G_U$  is a finite group of homeomorphisms of  $X_U$  and  $\Gamma_U$  is a map. One of the issues with the atlas definition is that there is no canonical notion of map between orbifolds. Satake introduced maps of orbifolds [8, 9] and called them embedding maps which are considered as generalizations of smooth maps of manifolds. One may refer to [2, 3, 5, 7] for more about orbifold.

In this paper, Section 2 introduces the concept of E-fibration embedding and studies the fundamental properties of E-fibration embedding such as the restriction property, product property, relationship between E-fibration embedding maps and Hurewicz fibrations. In Section 3, we introduce the notion of lifting functions of E-fibration embedding by giving the concept of E-lifting function and regular E-lifting function. In Section 4, we show preserving projection property for E-lifting functions. Throughout this paper all spaces will be assumed to be Hausdorff spaces. For any space  $X$ ,  $X^I$  denotes the set of all continuous functions since space (paths) from  $I = [0, 1]$  into  $X$ . We take  $X^I$  with the compact-open topology. For all  $x \in X$ , by  $\tilde{x}$  we mean the constant path at a point  $x$ . For two paths  $\alpha, \beta \in X^I$  with  $\alpha(1) = \beta(0)$ , by  $\alpha \star \beta$  we mean the path in  $X$  defined by

$$(\alpha \star \beta)(t) = \begin{cases} \alpha(2t), & 0 \leq t \leq \frac{1}{2}; \\ \beta(2t - 1), & \frac{1}{2} \leq t \leq 1. \end{cases}$$

**Definition 1.1.** [6, 10] For two continuous functions  $f, g : X \rightarrow Y$ , the function  $f$  is called a *homotopic* to  $g$  and write  $f \simeq g$  if there exists a continuous function  $H : X \times I \rightarrow Y$  satisfies  $H(x, 0) = f(x)$  and  $H(x, 1) = g(x)$ .

**Definition 1.2.** [4] A map  $f : X \rightarrow Y$  is called a *Hurewicz fibrations* if for every space  $Z$  and two maps  $h : Z \rightarrow X$  and  $H : Z \times I \rightarrow Y$  with  $H_0 = f \circ h$ , there exists a map  $F : Z \times I \rightarrow X$  such that  $F_0 = h$  and  $f \circ F = H$ .

By a *topological group*  $G$  we mean a group  $G$  together with a topology on  $G$  such that the functions  $(g, g') \rightarrow gg'$  and  $g \rightarrow g^{-1}$  are continuous of a product space  $G \times G$  into a space  $G$  and of a space  $G$  into  $G$ , respectively. The *action* of  $G$  on any space  $X$  is defined as a map  $G \times X \rightarrow X$  denoted by  $(g, x) \rightarrow gx$  such that  $g(g'x) = (gg')x$  and  $1x = x$  for all  $g, g' \in G$  and  $x \in X$ .

For action  $G \times X \rightarrow X$  of a topological group  $G$  on a space  $X$  and for  $x \in X$ , we mean by the *orbit set* of  $x$  is the set  $G(x) = \{gx \in X : g \in G\}$  and the *orbit space*  $X/G$  is the set of all orbits  $G(x)$  in  $X$  endowed with the quotient topology with respect to the *natural orbit map*  $X \rightarrow X/G$ .

**Definition 1.3.** [1] An *orbifold chart* on topological space  $X$  is a triple  $(X_U, G_U, \Gamma_U)$ , where  $X_U$  is an open set in a space  $R^n$ ,  $G_U$  is a finite group of homeomorphisms of  $X_U$  and  $\Gamma_U : X_U \rightarrow X$  is a map defined by  $\Gamma_U = \overline{\Gamma_U} \circ p$ , where  $p : X_U \rightarrow X_U/G_U$  is the orbit map and  $\overline{\Gamma_U} : X_U/G_U \rightarrow X$  is a map that induces a homeomorphism of  $X_U$  onto an open subset  $U$  of  $X$ .

**Definition 1.4.** [1] For a topological space  $X$ , an *embedding*  $f : X_U \rightarrow X_{U'}$  is a smooth injective function from orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$  and it yields a homeomorphism between  $X_U$  and  $f(X_U)$  such that  $\Gamma_{U'} \circ f = \Gamma_U$ .

## 2 E-fibration embedding maps

In this section we introduce *E-fibration* embedding map and study related properties of it.

**Definition 2.1.** An embedding  $f : X_U \rightarrow X_{U'}$  of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$  is called an *E-fibration* if for every space  $Z$  and map  $h : Z \rightarrow X_U$  and  $H : Z \rightarrow X^I$  with  $H_0 = \Gamma_U \circ h$ , there exists a map  $F : Z \rightarrow (X_{U'})^I$  such that  $F_0 = f \circ h$  and  $\Gamma_{U'} \circ F_t = H_t$  for all  $t \in I$ .

By an *E-triple*  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  we mean three spaces  $X_1, X_2$  and  $X$  with three maps  $\Theta_1 : X_1 \rightarrow X$ ,  $\Theta_2 : X_2 \rightarrow X$  and  $\Theta : X_2 \rightarrow X_1$  such that  $\Theta_2 \circ \Theta = \Theta_1$ . We say that an E-triple  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has an *E-fibration property* if for every space  $Z$  and maps  $h : Z \rightarrow X_2$  and  $H : Z \rightarrow X^I$  with  $H_0 = \Theta_2 \circ h$ , there exists a map  $F : Z \rightarrow (X_1)^I$  such that  $F_0 = \Theta \circ h$  and  $\Theta_1 \circ F_t = H_t$  for all  $t \in I$ .

**Theorem 2.2.** An embedding  $f : X_U \rightarrow X_{U'}$  of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$  is an E-fibration embedding if and only if an E-triple  $(\Gamma_U|X_U, \Gamma_{U'}|X_{U'}, X_f)$  has an E-fibration property.

**Example 2.3.** For an embedding  $f : X_U \rightarrow X_{U'}$  of orbifold chart  $(X_U, G_U, \Gamma_U)$  into  $(X_{U'}, G_{U'}, \Gamma_{U'})$ , the E-triple  $(\pi_2|X_U \times X, \pi_2|X_{U'} \times X, X_f \times id_X)$  has an E-fibration property, where  $\pi_2$  is the usual second projection and  $id_X$  is the identity map on  $X$ . Note that If  $Z$  is any space,  $h : Z \rightarrow X_U \times X$  is any map, and  $H : Z \rightarrow X^I$  is a map with  $H_0 = \pi_2 \circ h$ , define the desired a map  $F : Z \rightarrow (X_{U'} \times X)^I$  by

$$F_t = (f \circ \pi_1 \circ h) \times H_t$$

for all  $t \in I$ , where  $\pi_1$  is the usual first projection.

In the following theorem, we show the composition property of E-fibrations and Hurewicz fibrations.

**Theorem 2.4.** Let  $f : X_U \rightarrow X_{U'}$  be an E-fibration embedding and  $f' : X \rightarrow X'$  be a Hurewicz fibration of  $X$  into a space  $X'$ . Then, the E-triple  $(f' \circ \Gamma_U|X_U, f' \circ \Gamma_{U'}|X_{U'}, X'_f)$  has an E-fibration property.

*Proof.* Let  $Z$  be any space and let  $h : Z \rightarrow X_U$  and  $H : Z \rightarrow X'^I$  be any two maps with  $H_0 = (f' \circ \Gamma_U) \circ h = f' \circ (\Gamma_U \circ h)$ . Since  $f'$  is a Hurewicz fibration, then there exists a map  $F' : Z \rightarrow X'^I$  such that  $F'_0 = \Gamma_U \circ h$  and  $f' \circ F'_t = H_t$  for all  $t \in I$ . Since  $f$  is an E-fibration, then there exists a map  $F : Z \rightarrow (X_{U'})^I$  such that  $F_0 = f \circ h$  and  $\Gamma_{U'} \circ F_t = F'_t$  for all  $t \in I$ . Then,

$$(f' \circ \Gamma_{U'}) \circ F_t = f' \circ (\Gamma_{U'} \circ F_t) = f' \circ F'_t = H_t$$

for all  $t \in I$ . Hence the E-triple  $(f' \circ \Gamma_U|X_U, f' \circ \Gamma_{U'}|X_{U'}, X'_f)$  has an E-fibration property.  $\square$

Theorems 2.5, 2.6 and 2.7 give the relations between E-fibrations and Hurewicz fibrations.

**Theorem 2.5.** Let  $f : X_U \rightarrow X_{U'}$  be an E-embedding from orbifold chart  $(X_U, G_U, \Gamma_U)$  into  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . If  $\Gamma_U$  or  $\Gamma_{U'}$  is a Hurewicz fibration then  $f$  is an E-fibration.

*Proof.* Let  $Z$  be any space and let  $h : Z \rightarrow X_U$  and  $H : Z \rightarrow X^I$  be any two maps with  $H_0 = \Gamma_U \circ h$ . If  $\Gamma_U$  is a Hurewicz fibration, then there exists a map  $F' : Z \rightarrow (X_U)^I$  such that  $F'_0 = h$  and  $\Gamma_U \circ F'_t = H_t$  for all  $t \in I$ . Define a map  $F : Z \rightarrow (X_{U'})^I$  by  $F_t = f \circ F'_t$  for all  $t \in I$ . Note that  $F_0 = f \circ F'_0 = f \circ h$  and

$$\Gamma_{U'} \circ F_t = \Gamma_{U'} \circ (f \circ F'_t) = (\Gamma_{U'} \circ f) \circ F'_t = \Gamma_U \circ F'_t = H_t$$

for all  $t \in I$ . Hence,  $f$  is an E-fibration. If  $\Gamma_{U'}$  is a Hurewicz fibration and since

$$H_0 = \Gamma_U \circ h = (\Gamma_{U'} \circ f) \circ h = \Gamma_{U'} \circ (f \circ h),$$

then there exists a map  $F : Z \rightarrow (X_{U'})^I$  such that  $F_0 = f \circ h$  and  $\Gamma_{U'} \circ F_t = H_t$  for all  $t \in I$ . Hence,  $f$  is a E-fibration.  $\square$

**Theorem 2.6.** Let  $f : X_U \rightarrow X_{U'}$  be an E-fibration embedding from orbifold chart  $(X_U, G_U, \Gamma_U)$  into  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . If  $f$  is a surjective, then  $\Gamma_{U'}$  is a Hurewicz fibration.

*Proof.* Let  $Z$  be any space and let  $h : Z \rightarrow X_{U'}$  and  $H : Z \rightarrow X^I$  be any two maps with  $H_0 = \Gamma_{U'} \circ h$ . Since  $f$  is a surjective, then there exists a map  $g : X_{U'} \rightarrow X_U$  such that  $f \circ g = id_{X_{U'}}$ . Hence,

$$H_0 = \Gamma_{U'} \circ h = (\Gamma_U \circ g) \circ h = \Gamma_U \circ (g \circ h).$$

Then, there exists a map  $F : Z \rightarrow (X_{U'})^I$  such that

$$F_0 = f \circ (g \circ h) = (f \circ g) \circ h = id_{X_{U'}} \circ h = h$$

and  $\Gamma_{U'} \circ F_t = H_t$  for all  $t \in I$ . Thus,  $\Gamma_{U'}$  is a Hurewicz fibration.  $\square$

**Theorem 2.7.** Let  $f : X_U \rightarrow X_{U'}$  be an E-fibration embedding from orbifold chart  $(X_U, G_U, \Gamma_U)$  into  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . If  $f$  is a homeomorphism, then  $\Gamma_U$  and  $\Gamma_{U'}$  are Hurewicz fibrations.

*Proof.* Due to theorem 2.6,  $\Gamma_{U'}$  is a Hurewicz fibration. Let  $Z$  be any space and let  $h : Z \rightarrow X_U$  and  $H : Z \rightarrow X^I$  be any two maps with  $H_0 = \Gamma_U \circ h$ . Then, there exists a map  $F' : Z \rightarrow (X_{U'})^I$  such that  $F'_0 = f \circ h$  and  $\Gamma_{U'} \circ F'_t = H_t$  for all  $t \in I$ . Define a map  $F : Z \rightarrow (X_U)^I$  by  $F_t = f^{-1} \circ F'_t$  for all  $t \in I$ . It is easy to note that

$$F_0 = f^{-1} \circ F'_0 = f^{-1} \circ (f \circ h) = h$$

and

$$\Gamma_U \circ F_t = \Gamma_U \circ (f^{-1} \circ F'_t) = (\Gamma_U \circ f^{-1}) \circ F'_t = \Gamma_{U'} \circ F'_t = H_t$$

for all  $t \in I$ . Thus,  $\Gamma_U$  is a Hurewicz fibration.  $\square$

The following theorem shows that the product of two E-fibrations has an E-fibration property.

**Theorem 2.8.** Let  $f : X_U \rightarrow X_{U'}$  and  $g : Y_V \rightarrow Y_{V'}$  be two E-fibration embeddings over two spaces  $X$  and  $Y$ , respectively. Then, the E-triple

$$(\Gamma_U \times \Gamma_V | X_U \times Y_V, \Gamma_{U'} \times \Gamma_{V'} | X_{U'} \times Y_{V'}, (X \times Y)_{f \times g})$$

has an E-fibration property.

*Proof.* Let  $Z$  be any space and let  $h : Z \rightarrow X_U \times Y_V$  and  $H : Z \rightarrow (X \times Y)^I$  be any two maps with  $H_0 = (\Gamma_U \times \Gamma_V) \circ h$ . Define two maps

$$H^X : Z \rightarrow X^I \text{ and } H^Y : Z \rightarrow Y^I$$

by

$$H_t^X = \pi_1 \circ H_t \text{ and } H_t^Y = \pi_2 \circ H_t$$

for all  $t \in I$ , where  $\pi_1$  and  $\pi_2$  are the usual first and second projections. It is easy to note that

$$H_0^X = \pi_1 \circ H_0 = \pi_1 \circ [(\Gamma_U \times \Gamma_V) \circ h] = \Gamma_U \circ (\pi_1 \circ h).$$

Since  $f$  is an E-fibration, then there exists a map  $F' : Z \rightarrow (X_{U'})^I$  such that  $F'_0 = f \circ (\pi_1 \circ h)$  and  $\Gamma_{U'} \circ F'_t = H_t^X$  for all  $t \in I$ . Similarly, for an E-fibration embedding  $g$ , there exists a

map  $F'' : Z \rightarrow (Y_{V'})^I$  such that  $F''_0 = g \circ (\pi_2 \circ h)$  and  $\Gamma_{V'} \circ F''_t = H_t^Y$  for all  $t \in I$ . Define a map  $F : Z \rightarrow (X_{U'} \times Y_{V'})^I$  by  $F_t = F'_t \times F''_t$  for all  $t \in I$ . We observe that

$$\begin{aligned} F_0 = F'_0 \times F''_0 &= [f \circ (\pi_1 \circ h)] \times [g \circ (\pi_2 \circ h)] = (f \times g) \circ [(\pi_1 \times \pi_2) \circ h] \\ &= (f \times g) \circ h. \end{aligned}$$

and

$$\begin{aligned} (\Gamma_{U'} \times \Gamma_{V'}) \circ F_t &= (\Gamma_{U'} \times \Gamma_{V'}) \circ (F'_t \times F''_t) = (\Gamma_{U'} \circ F'_t) \times (\Gamma_{V'} \circ F''_t) \\ &= H_t^X \times H_t^Y = (\pi_1 \circ H_t) \times (\pi_2 \circ H_t) \\ &= (\pi_1 \times \pi_2) \circ H_t = H_t \end{aligned}$$

for all  $t \in I$ . Hence, the proof is completed.  $\square$

**Theorem 2.9.** Let  $f : X_U \rightarrow X_{U'}$  be an E-fibration embedding from orbifold chart  $(X_U, G_U, \Gamma_U)$  into  $(X_{U'}, G_{U'}, \Gamma_{U'})$  and  $S$  be any subspace of  $X$ . Then, the E-triple

$$(\Gamma_{U_s} | \Gamma_U^{-1}(S), \Gamma_{U'_s} | \Gamma_{U'}^{-1}(S), S_{f_s})$$

has an E-fibration property where  $\Gamma_{U_s}$ ,  $\Gamma_{U'_s}$  and  $f_s$  are the restriction maps of  $\Gamma_U$ ,  $\Gamma_{U'}$  and  $f$  on  $\Gamma_U^{-1}(S)$ ,  $\Gamma_{U'}^{-1}(S)$  and  $\Gamma_U^{-1}(S)$ , respectively.

*Proof.* Let  $Z$  be any space and let  $h : Z \rightarrow \Gamma_U^{-1}(S)$  and  $H : Z \rightarrow S^I$  be any two maps with  $H_0 = \Gamma_{U_s} \circ h$ . Let  $j_u : \Gamma_U^{-1}(S) \rightarrow X_U$ ,  $j_{u'} : \Gamma_{U'}^{-1}(S) \rightarrow X_{U'}$  and  $j : S \rightarrow X$  be inclusion maps. Define a map  $H' : Z \rightarrow X^I$  by  $H'_t = j \circ H_t$  for all  $t \in I$ . Since  $f$  is an E-fibration embedding and

$$H'_0 = j \circ H_0 = H_0 = \Gamma_{U_s} \circ h = (\Gamma_{U_s} \circ j_u) \circ h = \Gamma_{U_s} \circ (j_u \circ h),$$

then there exists map  $F : Z \rightarrow (X_{U'})^I$  such that  $F_0 = f \circ (j_u \circ h)$  and  $\Gamma_{U'} \circ F_t = H'_t$  for all  $t \in I$ . Since

$$\Gamma_{U'}[F_t(z)] = H'_t = j \circ H_t = H_t \in S$$

then  $F(z)(t) \in \Gamma_{U'}^{-1}(S)$  for all  $z \in Z$ ,  $t \in I$ . Hence,  $F$  is a homotopy from  $Z$  into  $(\Gamma_{U'}^{-1}(S))^I$  and it is easy to note that

$$F_0 = f \circ (j_u \circ h) = (f \circ j_u) \circ h = f_s \circ h$$

and

$$\Gamma_{U'_s} \circ F_t = (\Gamma_{U'} \circ j_{u'}) \circ F_t = \Gamma_{U'} \circ (j_{u'} \circ F_t) = \Gamma_{U'} \circ F_t = H_t$$

for all  $t \in I$ . Then, the E-triple  $(\Gamma_{U_s} | \Gamma_U^{-1}(S), \Gamma_{U'_s} | \Gamma_{U'}^{-1}(S), S_{f_s})$  has an E-fibration property.  $\square$

**Remark 2.10.** For any embedding  $f : X_U \rightarrow X_{U'}$  of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$  and for any map  $P : X' \rightarrow X$  of a space  $X'$  into a space  $X$ , define the maps  $P_1 : X'(U) \rightarrow X'$ ,  $P'_1 : X'(U') \rightarrow X'$  and  $P_f : X'(U) \rightarrow X'(U')$  by

$$P_1(x', r) = x', \quad P'_1(x', r') = x' \quad \text{and} \quad P_f(x', r) = (x', f(r)),$$

respectively, for all  $(x', r) \in X'(U)$  and  $(x', r') \in X'(U')$ , where

$$X'(U) = \{(x', r) \in X' \times X_U : \Gamma_U(r) = P(x')\}$$

and

$$X'(U') = \{(x', r') \in X' \times X_{U'} : \Gamma_{U'}(r') = P(x')\}.$$

It is easy to note that for all  $(x', r) \in X'(U)$  we have that

$$(P'_1 \circ P_f)(x', r) = x' = P_1(x', r).$$

That is,  $(P_1|X'(U), P'_1|X'(U'), X'_{P_f})$  is an E-triple which is called the *E-pullback* of an embedding  $f$  by a map  $P$ .

**Theorem 2.11.** For any embedding E-fibration  $f : X_U \rightarrow X_{U'}$  of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$ , the E-pullback of  $f$  by any map  $P : X' \rightarrow X$  has an E-fibration property.

*Proof.* Let  $Z$  be any space. Let  $h' : Z \rightarrow X'(U)$  and  $H' : Z \rightarrow X'^I$  be two maps with  $H'_0 = P'_1 \circ h'$ . Define a map  $h : Z \rightarrow X_U$  by  $h(z) = \pi_2(h'(z))$  and a map  $H : Z \rightarrow X^I$  by  $H(z) = P \circ H'(z)$  for all  $z \in Z$ . Thus,

$$\begin{aligned} H(z)(0) &= (P \circ H'(z))(0) = P(H'(z)(0)) = P[(P'_1 \circ h')(z)] = P(P'_1(h'(z))) \\ &= P(\pi_1(h'(z))) = \Gamma_U(\pi_2(h'(z))) = \Gamma_U(h(z)) \end{aligned}$$

for all  $z \in Z$ . That is,  $H_0 = \Gamma_U \circ h$ , where  $\pi_1$  and  $\pi_2$  are the usual first and second projections. Since  $f$  is an E-fibration, then there exists a map  $F : Z \rightarrow X_{U'}^I$  such that  $F_0 = f \circ h$  and  $\Gamma_{U'} \circ F_t = H_t$  for all  $t \in I$ . Define a map  $F' : Z \rightarrow X'(U')$  by  $F'(z)(t) = [H'(z)(t), F(z)(t)]$  for all  $t \in I, z \in Z$ . Note  $P'_1 \circ F' = H'$  and

$$\begin{aligned} F'(z)(0) &= [H'(z)(0), F(z)(0)] = [P'_1(h'(z)), f(h(z))] = [\pi_1(h'(z)), f(\pi_2(h'(z)))] \\ &= P_f[\pi_1(h'(z)), \pi_2(h'(z))] = P_f(h'(z)) = (P_f \circ h')(z) \end{aligned}$$

for all  $z \in Z$ . Thus,  $F'_0 = P_f \circ h'$ . Hence, the E-pullback of  $f$  by a map  $P : X' \rightarrow X$  has an E-fibration property.  $\square$

### 3 E-lifting function

In this section, we introduce E-lifting function and study some of its properties.

**Definition 3.1.** Let  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  be an E-triple and let

$$\Delta\Theta_1 = \{(\Theta(x_2), \alpha) \in X_1 \times X^I : \Theta_2(x_2) = \alpha(0) \text{ for some } x_2 \in X_2\}.$$

The map  $\lambda_\Theta : \Delta\Theta_1 \rightarrow X_1^I$  is called an *E-lifting function* of an E-triple  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  if it satisfies the following:

1.  $\lambda_\Theta[\Theta(x_2), \alpha](0) = \Theta(x_2)$  for all  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$ ;

2.  $[\Theta_1 \circ \lambda_\Theta(\Theta(x_2), \alpha)](t) = \alpha(t)$  for all  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$  and  $t \in I$ .

Now, we discuss the following theorems related to E-lifting function.

**Theorem 3.2.** The E-triple  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has E-fibration property. if and only if it has an E-lifting function.

*Proof.* Let the E-triple  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has E-lifting function  $\lambda_\Theta$  and  $Z$  be any space. Let  $h : Z \rightarrow X_2$  be any map and  $H : Z \rightarrow X^I$  be any map such that  $\Theta_2 \circ h = H_0$ . For  $z \in Z$ , consider a path  $\alpha_z : I \rightarrow X$  defined by  $\alpha_z(t) = H(z, t)$  for all  $t \in I$ . Define a map  $F : Z \rightarrow (X_1)^I$  by

$$F(z)(t) = \lambda_\Theta[(\Theta \circ h)(z), \alpha_z](t)$$

for all  $z \in Z, t \in I$ . Then, we observe that

$$F(z)(0) = \lambda_\Theta[(\Theta \circ h)(z), \alpha_z](0) = (\Theta \circ h)(z)$$

and

$$\begin{aligned} (\Theta_1 \circ F)(z)(t) &= \Theta_1\{\lambda_\Theta[(\Theta \circ h)(z), \alpha_z](t)\} \\ &= \alpha_z(t) = H(z)(t) \end{aligned}$$

for all  $z \in Z, t \in I$ . Thus,  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has E-fibration property.

Conversely, let  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has E-fibration property. Let  $h : \Delta\Theta_1 \rightarrow X_2$  be a map and  $H : \Delta\Theta_1 \rightarrow X^I$  be a map defined by  $h(\Theta(x_2), \alpha) = x_2$  for all  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$  and  $H(\Theta(x_2), \alpha)(t) = \alpha(t)$  for all  $t \in I$  and  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$ . Note that

$$\begin{aligned} H(\Theta(x_2), \alpha)(0) &= \alpha(0) = \Theta_2(x_2) \\ &= (\Theta_2 \circ h)(\Theta(x_2), \alpha) \end{aligned}$$

for all  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$ . Thus,  $H_0 = \Theta_2 \circ h$ . Since  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has E-fibration property, then there exists a map  $F : \Delta\Theta_1 \rightarrow X_1^I$  such that  $\Theta_1 \circ F_t = H_t$  and  $\Theta \circ h = F_0$  for all  $t \in I$ . Now define a map  $\lambda_\Theta : \Delta\Theta_1 \rightarrow X_1^I$  by

$$\lambda_\Theta[\Theta(x_2), \alpha](t) = F(\Theta(x_2), \alpha)(t)$$

for all  $t \in I$  and  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$ . Note that

$$\begin{aligned} \lambda_\Theta[\Theta(x_2), \alpha](0) &= F(\Theta(x_2), \alpha)(0) \\ &= (\Theta \circ h)(\Theta(x_2), \alpha) = \Theta(x_2) \end{aligned}$$

for all  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$  and

$$\begin{aligned} \Theta_1 \circ \lambda_\Theta[\Theta(x_2), \alpha](t) &= (\Theta_1 \circ F)[\Theta(x_2), \alpha](t) \\ &= H(\Theta(x_2), \alpha)(t) = \alpha(t) \end{aligned}$$

for all  $(\Theta(x_2), \alpha) \in \Delta\Theta_1$  and  $t \in I$ . Hence,  $\lambda_\Theta$  is an E-lifting function of  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$ .  $\square$

We say that an E-triple  $(\Theta_2|X_2, \Theta_1|X_1, X_\Theta)$  has an *E-regular fibration property* if it has an *E-regular lifting function*  $\lambda_\Theta$ , i.e.  $\lambda_\Theta[\Theta(x_2), \Theta_1 \circ \widetilde{\Theta(x_2)}] = \widetilde{\Theta(x_2)}$  for all  $\Theta(x_2) \in X_1$ .

We say that an embedding map  $f : X_U \rightarrow X_{U'}$  is an *E-regular fibration embedding* if its E-triple  $(\Gamma_U|X_U, \Gamma_{U'}|X_{U'}, X_f)$  has E-regular lifting function.

**Example 3.3.** In Example 2.3, we can define the map  $\lambda_{f \times id_X} : \Delta\pi_2 \rightarrow (X_{U'} \times X)^I$  by

$$\lambda_{f \times id_X}[(f \times id_X)(r, x), \alpha](t) = (f(r), \alpha(t))$$

for all  $t \in I, ((f \times id_X)(r, x), \alpha) \in \Delta\pi_2$ . Note that for  $((f \times id_X)(r, x), \alpha) \in \Delta\pi_2$  and  $t \in I$ ,

$$\lambda_{f \times id_X}[(f \times id_X)(r, x), \alpha](0) = (f(r), x) = (f \times id_X)(r, x)$$

and

$$\pi_2 \circ \lambda_{f \times id_X}[(f \times id_X)(r, x), \alpha](t) = \pi_2(f(r), \alpha(t)) = \alpha(t).$$

Hence,  $\lambda_{f \times id_X}$  is an E- lifting function of E-triple  $(\pi_2|X_U \times X, \pi_2|X_{U'} \times X, X_{f \times id_X})$ . Also, we observe that for  $f \times id_X(r, x) \in X_{U'} \times X$ ,

$$\begin{aligned} \lambda_{f \times id_X}[(f \times id_X)(r, x), \pi_2 \circ \widetilde{f \times id_X}(r, x)](t) &= [f(r), (\pi_2 \circ \widetilde{f \times id_X}(r, x))(t)] \\ &= (f(r), x) = (f \times id_X)(r, x) \\ &= f \times \widetilde{id_X}(r, x)(t). \end{aligned}$$

Hence,  $\lambda_{f \times id_X}$  is an E-regular lifting function.

## 4 Preserving projection property

In this section, we show preserving projection property for E-lifting function.

**Theorem 4.1.** Let  $f : X_U \rightarrow X_{U'}$  be any E-regular fibration embedding of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . Let  $\lambda' : [f(X_U)]^I \rightarrow X_{U'}^I$  be a map defined by  $\lambda'(\beta) = \lambda_f(\beta(0), \Gamma_{U'} \circ \beta)$  for all  $\beta \in [f(X_U)]^I$ . If  $f$  is an injective, then  $\lambda' \simeq inclusion : [f(X_U)]^I \subset X_{U'}^I$ , *preserving projection*. This means that there exists a map  $H : [f(X_U)]^I \times I \rightarrow X_{U'}^I$  between two maps  $\lambda'$  and *inclusion* :  $[f(X_U)]^I \subset X_{U'}^I$ , such that

$$[\Gamma_{U'} \circ H(\beta, s)](t) = \Gamma_{U'}(\beta(t))$$

for all  $\beta \in [f(X_U)]^I, s, t \in I$ .

*Proof.* Since  $f$  is an injective, then for  $\beta \in [f(X_U)]^I$  and  $s \in I$ , there exists exactly one  $r$  such that  $f(r) = \beta(s)$ . Hence,  $\lambda'$  is well-defined and the maps in this proof will be well-defined. For  $\beta \in [f(X_U)]^I$  and  $s \in I$ , we can define a path  $\beta_s \in [f(X_U)]^I$  by

$$\beta_s(t) = \begin{cases} \beta(t) & \text{for } 0 \leq t \leq s, \\ \beta(s) & \text{for } s \leq t \leq 1. \end{cases}$$



For  $\alpha = \Gamma_{U'} \circ \beta$  and  $s \in I$ , we can define the path  $\alpha^{1-s} \in X^I$  by

$$\alpha^{1-s}(t) = \begin{cases} \alpha(s+t) & \text{for } 0 \leq t \leq 1-s, \\ \alpha(1) & \text{for } 1-s \leq t \leq 1. \end{cases}$$

Define a map  $H : [f(X_U)]^I \times I \rightarrow X_{U'}^I$  by

$$H(\beta, s)(t) = \begin{cases} \beta_s(t) & \text{for } 0 \leq t \leq s, \\ \lambda_f(\beta(s), \alpha^{1-s})(t-s) & \text{for } s \leq t \leq 1, \end{cases}$$

for all  $s \in I, \beta \in [f(X_U)]^I$ . By the E-regularity of  $f$ , we observe that

$$\begin{aligned} H(\beta, 0)(t) &= \lambda_f(\beta(0), \alpha^1)(t) = \lambda_f(\beta(0), \alpha)(t) \\ &= \lambda_f(\beta(0), \Gamma_{U'} \circ \beta)(t) = \lambda'(\beta)(t), \end{aligned}$$

and  $H(\beta, 1)(t) = \beta(t)$  for  $\beta \in [f(X_U)]^I, t \in I$ . Hence,

$$H(\beta, 0) = \lambda'(\beta) \quad \text{and} \quad H(\beta, 1) = \beta$$

for all  $\beta \in [f(X_U)]^I$ . Thus,  $\lambda' \simeq \text{inclusion} : [f(X_U)]^I \subset X_{U'}^I$ . Also,

$$\begin{aligned} [\Gamma_{U'} \circ H(\beta, s)](t) &= \begin{cases} \Gamma_{U'}(\beta_s(t)) & \text{for } 0 \leq t \leq s, \\ \Gamma_{U'}[\lambda_f(\beta(s), \alpha^{1-s})(t-s)] & \text{for } s \leq t \leq 1, \end{cases} \\ &= \begin{cases} \Gamma_{U'}(\beta(t)) & \text{for } 0 \leq t \leq s, \\ \alpha^{1-s}(t-s) & \text{for } 0 \leq t-s \leq 1-s, \end{cases} \\ &= \begin{cases} \Gamma_{U'}(\beta(t)) & \text{for } 0 \leq t \leq s, \\ \alpha(s+t-s) & \text{for } 0 \leq t-s \leq 1-s, \end{cases} \\ &= \begin{cases} \Gamma_{U'}(\beta(t)) & \text{for } 0 \leq t \leq s, \\ \alpha(t) & \text{for } s \leq t \leq 1, \end{cases} \\ &= \begin{cases} \Gamma_{U'}(\beta(t)) & \text{for } 0 \leq t \leq s, \\ \Gamma_{U'}(\beta(t)) & \text{for } s \leq t \leq 1, \end{cases} \\ &= \Gamma_{U'}(\beta(t)). \end{aligned}$$

Hence,  $\lambda' \simeq \text{inclusion} : [f(X_U)]^I \subset X_{U'}^I$  preserves projection.  $\square$

**Corollary 4.2.** Let  $f : X_U \rightarrow X_{U'}$  be any E-regular fibration embedding of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . Let  $\lambda' : X_{U'}^I \rightarrow X_{U'}^I$  be a map defined by

$$\lambda'(\beta) = \lambda_f(\beta(0), \Gamma_{U'} \circ \beta)$$

for all  $\beta \in X_{U'}^I$ . If  $f$  is a homeomorphism, then  $\lambda' \simeq \text{id}_{X_{U'}^I}$  preserves projection.

*Proof.* Since  $f$  is surjective, thus we can put  $[f(X_U)]^I = X_{U'}^I$ . Hence, the proof can be obtained easily.  $\square$

**Corollary 4.3.** In the Theorem 4.1, let  $f$  be a homeomorphism and for a path  $\beta \in X_{U'}^I$ , let  $g_\beta$  be a path in  $X_{U'}^I$  defined by  $g_\beta(s) = H(\beta, s)(1)$  for all  $s \in I$ . Let  $N : X_{U'}^I \rightarrow X_{U'}^I$  be a map defined by  $N(\beta) = \lambda_f(\beta(0), \Gamma_{U'} \circ \beta) \star g_\beta$  for all  $\beta \in X_{U'}^I$ . Then,  $N \simeq \text{id}_{X_{U'}^I}$  keeping end points fixed.

*Proof.* For a path  $\beta$  in  $X_{U'}^I$ , and  $s \in I$ , we can define a path  $\beta_s$  in  $X_{U'}^I$ , by

$$\beta_s(t) = \beta(s + (1 - s)t)$$

for all  $t \in I$ . Define the map  $G : X_{U'}^I \times I \rightarrow X_{U'}^I$ , by

$$G(\beta, s) = H(\beta, s) \circ (g_\beta)_s$$

for all  $s \in I, \beta \in X_{U'}^I$ . Hence, we observe that

$$\begin{aligned} G(\beta, 0) &= H(\beta, 0) \circ (g_\beta)_0 = \lambda_f(\beta(0), \Gamma_{U'} \circ \beta) \circ g_\beta \\ &= N(\beta) \end{aligned}$$

and

$$G(\beta, 1) = H(\beta, 1) \circ [g_\beta]_1 = \beta \circ \widetilde{\beta(1)} \simeq id_{X_{U'}^I}$$

for all  $\beta \in X_{U'}^I$ . Hence,  $N \simeq id_{X_{U'}^I}$ . Also we observe that

$$\begin{aligned} G(\beta, s)(0) &= [H(\beta, s) \circ (g_\beta)_s](0) \\ &= H(\beta, s)(0) = \beta_s(0) = \beta(0), \end{aligned}$$

and

$$G(\beta, s)(1) = [H(\beta, 0) \circ (g_\beta)_s](1) = (g_\beta)_s(1) = \beta(1).$$

for all  $\beta \in X_{U'}^I$ , and  $s \in I$ . Hence,  $N \simeq id_{X_{U'}^I}$ , keeping end points fixed.  $\square$

**Corollary 4.4.** Let  $f : X_U \rightarrow X_{U'}$  be an E-regular fibration embedding of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . If  $f$  is a homeomorphism, then  $X_{U'}^I$ , and  $\Delta\Gamma_{U'}$  are of the same map type.

*Proof.* We show that the E-lifting function  $\lambda_f : \Delta\Gamma_{U'} \rightarrow X_{U'}^I$ , is a map equivalence. We can define the map  $K : X_{U'}^I \rightarrow \Delta\Gamma_{U'}$  by  $K(\beta) = (\beta(0), \Gamma_{U'} \circ \beta)$  for all  $\beta \in X_{U'}^I$ . Then, we observe that from Corollary 4.3,  $\lambda' = \lambda_f \circ K \simeq id_{X_{U'}^I}$ . Also, we observe that for  $(f(r), \alpha) \in \Delta\Gamma_{U'}$ ,

$$\begin{aligned} (K \circ \lambda_f)(f(r), \alpha) &= K(\lambda_f(f(r), \alpha)) \\ &= (\lambda_f(f(r), \alpha)(0), \Gamma_{U'}[\lambda_f(f(r), \alpha)]) \\ &= (f(r), \alpha) = id_{\Delta\Gamma_{U'}}(f(r), \alpha). \end{aligned}$$

Hence,  $K \circ \lambda_f = id_{\Delta\Gamma_{U'}}$ . Then, we get that  $\lambda_f : \Delta\Gamma_{U'} \rightarrow X_{U'}^I$  is a map equivalence.  $\square$

**Theorem 4.5.** Let  $f : X_U \rightarrow X_{U'}$  be an E-regular fibration embedding of orbifold chart  $(X_U, G_U, \Gamma_U)$  into orbifold chart  $(X_{U'}, G_{U'}, \Gamma_{U'})$ . Let  $\lambda', \widehat{f} : X_U^I \rightarrow X_{U'}^I$ , be two maps defined by

$$\lambda'(\beta) = \lambda_f[f(\beta(0)), \Gamma_U \circ \beta] \text{ and } \widehat{f}(\beta) = f \circ \beta$$

for all  $\beta \in X_U^I$ . Then,  $\lambda' \simeq \widehat{f}$  preserves projection.

*Proof.* For  $\beta \in X_U^I$  and  $s \in I$ , we can define a path  $\beta_s \in X_U^I$  by

$$\beta_s(t) = \begin{cases} \beta(t) & \text{for } 0 \leq t \leq s, \\ \beta(s) & \text{for } s \leq t \leq 1, \end{cases}$$

For  $\alpha = \Gamma_U \circ \beta$  and  $s \in I$ , we can define the path  $\alpha^{1-s} \in X$  by

$$\alpha^{1-s}(t) = \begin{cases} \alpha(s+t) & \text{for } 0 \leq t \leq 1-s, \\ \alpha(1) & \text{for } 1-s \leq t \leq 1, \end{cases}$$

Define a map  $H : X_U^I \times I \rightarrow X_{U'}^I$  by

$$H(\beta, s)(t) = \begin{cases} f[\beta_s(t)] & \text{for } 0 \leq t \leq s, \\ \lambda_f[f(\beta(s)), \alpha^{1-s}](t-s) & \text{for } s \leq t \leq 1, \end{cases}$$

for all  $s \in I, \beta \in X_U^I$ . By the E-regularity of  $f$ , we get that

$$\begin{aligned} H(\beta, 0)(t) &= \lambda_f[f(\beta(0)), \alpha^1](t) = \lambda_f[f(\beta(0)), \alpha](t) \\ &= \lambda_f[f(\beta(0)), \Gamma_U \circ \beta](t) = \lambda'(\beta)(t), \end{aligned}$$

and  $H(\beta, 1)(t) = (f \circ \beta)(t)$  for all  $\beta \in X_U^I, t \in I$ . Hence,

$$H(\beta, 0) = \lambda'(\beta) \quad \text{and} \quad H(\beta, 1) = f \circ \beta = \widehat{f}(\beta),$$

for all  $\beta \in X_U^I$ , that is,  $\lambda' \simeq \widehat{f}$ . Also

$$\begin{aligned} [\Gamma_{U'} \circ H(\beta, s)](t) &= \begin{cases} \Gamma_{U'}\{f[\beta_s(t)]\} & \text{for } 0 \leq t \leq s, \\ \Gamma_{U'}\{\lambda_f[f(\beta(s)), \alpha^{1-s}](t-s)\} & \text{for } s \leq t \leq 1, \end{cases} \\ &= \begin{cases} \Gamma_{U'}\{f[\beta(t)]\} & \text{for } 0 \leq t \leq s, \\ \alpha^{1-s}(t-s) & \text{for } 0 \leq t-s \leq 1-s, \end{cases} \\ &= \begin{cases} \Gamma_{U'}\{f[\beta(t)]\} & \text{for } 0 \leq t \leq s, \\ \alpha(s+t-s) & \text{for } 0 \leq t-s \leq 1-s, \end{cases} \\ &= \begin{cases} \Gamma_{U'}\{f[\beta(t)]\} & \text{for } 0 \leq t \leq s, \\ \alpha(t) & \text{for } s \leq t \leq 1, \end{cases} \\ &= \begin{cases} \Gamma_{U'}\{f[\beta(t)]\} & \text{for } 0 \leq t \leq s, \\ (\Gamma_U \circ \beta)(t) & \text{for } s \leq t \leq 1, \end{cases} \\ &= \begin{cases} \Gamma_{U'}\{f[\beta(t)]\} & \text{for } 0 \leq t \leq s, \\ \Gamma_{U'}\{f[\beta(t)]\} & \text{for } s \leq t \leq 1, \end{cases} \\ &= \Gamma_{U'}[\widehat{f}(\beta)(t)] \end{aligned}$$

or all  $\beta \in X_U^I, s, t \in I$ . Therefore,  $\lambda' \simeq \widehat{f}$  preserving projection.  $\square$

## 5 Conclusion

In this paper, we introduced E-fibration embedding map in an orbifold chart. Later we established some fundamental properties of E-fibration such as the restriction, product, etc.

We also studied the relationship of E-fibration with Hurewicz fibration. Furthermore, we introduced the notion of E-lifting functions and studied preserving projection property of these lifting functions. Ideas of homotopy played a crucial role in this paper. Thus, we hope that this paper will open some new dimensions in research related to orbifold and Hurewicz fibration.

## Acknowledgement

The authors would like to thank the Deanship of Scientific Research at Umm Al-Qura University for supporting this work by Grant Code : 22UQU4330052DSR02

## References

- [1] A. Adem, J. Leida, and Y. Ruan, *Orbifolds and Stringy Topology*, Cambridge Tracts in Mathematics, (2007).
- [2] N. Brodsky, A. Chigogidze, E.V. Scepin, Sections of Serre fibrations with 2-manifold fibers, *Top. Appl.*, 155, (2008), 773-782.
- [3] B. Fantechi, L. Gottsche, Orbifold cohomology for global quotients, *Duke Math. Jour.*, 2 (2003), 197-227.
- [4] W. Hurewicz, On the concept of fiber space, *Proc. Nat. Acad. Sci. USA*, 14 (1955), 956-961.
- [5] C. Vafa and E. Witten, On orbifolds with discrete torsion, *Jour. Geom. Phys.*, 15 (1995), 189-214.
- [6] A. Hatcher, *Algebraic Topology*, Cambridge University Press, Cambridge, (2002).
- [7] A. Saif and H. A. Othman, Retractions In Homotopy Theory For Finite Topological Semigroups, *Tbilisi Math. Jour.*, 1(14) 2021, pp. 219-232.
- [8] I. Satake, On a Generalization of the Notion of Manifold, *Proceedings of the National Academy of Science U.S.A.*, 42 (1956), 359 - 363.
- [9] I. Satake, The Gauss-Bonnet theorem for V-manifolds, *Journal of the Mathematical Society of Japan*, 9 (1957), 464 - 492.
- [10] E.H. Spanier, *Algebraic Topology*, McGraw-Hill, New York, (1966).