

On Galois groups of linearized polynomials related to the special linear group of prime degree

Rod Gow and Gary McGuire

School of Mathematics and Statistics
University College Dublin
Ireland

Abstract

Let F be any field of prime characteristic p and let q be a power of p . We assume that F contains the finite field of order q . A q -polynomial L over F is an element of the polynomial ring $F[x]$ with the property that all exponents are a power of q . We assume that the coefficient of the x term of L is nonzero. We investigate the Galois group G of L over F , under the assumption that $L(x)/x$ is irreducible in $F[x]$. It is well known that if L has q -degree n , its roots are an n -dimensional vector space over the field of order q and G acts linearly on this space.

Our main theorem is the following. We consider a monic q -polynomial L whose q -degree is a prime, r , say, with $L(x)/x$ irreducible over F . Assuming that the coefficient of the x term of L is $(-1)^r$, we show that the Galois group of L over F is the special linear group $SL(r, q)$ of degree r over the field of order q . We also prove a projective version for the projective special linear group $PSL(r, q)$.

Keywords Galois group, linearized polynomial, special linear group

1 Introduction

Let p be a prime number, and let q be a positive power of p . Let \mathbb{F}_q denote the finite field with q elements. Let F be a field of characteristic p , and assume that F contains \mathbb{F}_q . A q -linearized polynomial over F is a polynomial of the form

$$L = a_0x + a_1x^q + a_2x^{q^2} + \cdots + a_nx^{q^n} \in F[x]. \quad (1)$$

If $a_n \neq 0$ we say that n is the q -degree of f . We will usually say q -polynomial instead of q -linearized polynomial.

The set of roots of a q -polynomial L forms an \mathbb{F}_q -vector space, which is contained in a splitting field of L . We make this statement more precise in the following lemma.

Lemma 1. *Let F be a field of prime characteristic p that contains \mathbb{F}_q . Let L be a q -polynomial of q -degree n in $F[x]$, with*

$$L = a_0x + a_1x^q + a_2x^{q^2} + \cdots + a_nx^{q^n}.$$

Let E be a splitting field for L over F and let V be the set of roots of L in E . Let G be the Galois group of E over F . Suppose that $a_0 \neq 0$. Then V is an \mathbb{F}_q -vector space of dimension n and G is naturally a subgroup of $GL(n, q)$.

Proof. The derivative $L'(x) = a_0$ and is thus a nonzero constant under the hypothesis above. It follows that L has no repeated roots and thus $|V| = q^n$. Let α and β be elements of V . Then $L(\alpha) = L(\beta) = 0$. Since

$$(\alpha + \beta)^{q^i} = \alpha^{q^i} + \beta^{q^i}$$

for all $i \geq 0$, it is clear that $L(\alpha + \beta) = 0$, and hence $\alpha + \beta \in V$.

Let λ be an element of \mathbb{F}_q . Since $\lambda^q = \lambda$, we have $(\lambda\alpha)^{q^i} = \lambda\alpha^{q^i}$. It follows that $L(\lambda\alpha) = \lambda L(\alpha) = 0$ and hence $\lambda\alpha \in V$. These arguments show that V is a vector space over \mathbb{F}_q . Furthermore, since $|V| = q^n$, V has dimension n over \mathbb{F}_q .

Let σ be an element of G . By definition of Galois group action, G permutes the roots of L and hence maps V into itself. In addition, since G is a group of field automorphisms of E , we have

$$\sigma(\alpha + \beta) = \sigma(\alpha) + \sigma(\beta)$$

for all α and β in V .

Now let λ be an element of \mathbb{F}_q . Since \mathbb{F}_q is assumed to be a subfield of F and G fixes F elementwise, we have $\sigma(\lambda) = \lambda$. Then, again by definition of Galois group action,

$$\sigma(\lambda\alpha) = \sigma(\lambda)\sigma(\alpha) = \lambda\sigma(\alpha).$$

This shows that the action of G on V is \mathbb{F}_q -linear and thus G may be considered to be a subgroup of $GL(n, q)$. \square

We call V the (\mathbb{F}_q -vector) space of roots of L .

The next result, a trivial consequence of elementary Galois theory, is the key idea in this paper.

Lemma 2. *Let F be a field of prime characteristic p that contains \mathbb{F}_q . Let L be a q -polynomial in $F[x]$, such that $L(x)/x$ is irreducible over F . Let E be a splitting field for L over F and let V be the space of roots of L in E . Let G be the Galois group of E over F . Then G acts transitively on the nonzero elements of V .*

Proof. G permutes the roots of $L(x)/x$, which are the nonzero elements of V . Since $L(x)/x$ is irreducible, the action of G on the roots of $L(x)/x$ is transitive. \square

It follows from these two lemmas that our Galois groups are subgroups of $GL(n, q)$ that act transitively on the nonzero vectors of the underlying n -dimensional space. There are a number of papers in the literature on this topic, as we will discuss in Sections 3 and 4. Before that, we present some elementary calculations in Section 2.

We remark that $SL(n, q)$ has been realized as a Galois group over $\mathbb{F}_q(t)$ by several authors, however our main result (Theorem 13) has a more general ground field.

2 Calculations with the determinant

Let F be a field of characteristic p and L be a monic q -polynomial of q -degree n in $F[x]$, with nonzero coefficient of x . Let $\alpha_1, \dots, \alpha_n$ be an \mathbb{F}_q -basis for the space of

roots of L in a splitting field E of L over F . We define the $(n+1) \times (n+1)$ matrix A by

$$A := \begin{bmatrix} \alpha_1 & \alpha_1^q & \alpha_1^{q^2} & \cdots & \alpha_1^{q^n} \\ \alpha_2 & \alpha_2^q & \alpha_2^{q^2} & \cdots & \alpha_2^{q^n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_n & \alpha_n^q & \alpha_n^{q^2} & \cdots & \alpha_n^{q^n} \\ x & x^q & x^{q^2} & \cdots & x^{q^n} \end{bmatrix}.$$

It is clear that $\det A$ is a q -polynomial with coefficients in E , and we claim that its roots are the same as those of L . For if we set $x = \alpha_i$, the determinant of the resulting matrix is zero. Thus the α_i are roots of $\det A$, and since $\det A$ is a q -polynomial, each element of the space of roots of L is a root of $\det A$. Thus $\det A$ has q^n different roots in L . As $\det A$ has q -degree at most n , we deduce that $\det A$ has q -degree exactly n and hence the coefficient of x^{q^n} is nonzero.

Let D denote the $n \times n$ matrix obtained by omitting the last row and last column of A . It is clear by expansion of $\det A$ that $\det D$ is the coefficient of x^{q^n} in $\det A$. Therefore $\det D$ is a nonzero element of E . Let $\delta = \det D$.

Lemma 3. *Continuing the notation above,*

1. $\delta^{q-1} \in F$, and
2. $F(\delta)$ is a normal extension of F .

Proof. Since $\det A$ and L have the same q -degree and same roots, and since L is assumed to be monic, and δ is the leading coefficient of $\det A$, we have

$$L = \frac{1}{\delta} \det A.$$

Expansion of $\det A$ shows that the coefficient of x in $\det A$ is $(-1)^n \delta^q$. It follows that the coefficient of x in L is $(-1)^n \delta^{q-1}$. An immediate consequence is that $\delta^{q-1} \in F$.

It follows that δ is a root in E of the polynomial $x^{q-1} - \delta^{q-1}$, which has coefficients in F . Since the roots of $x^{q-1} - \delta^{q-1}$ are the elements $\omega\delta$, where ω ranges over the nonzero elements of \mathbb{F}_q , E contains the splitting field $F(\delta)$ of $x^{q-1} - \delta^{q-1}$ over F . Thus $F(\delta)$ is a normal subfield of E . \square

Let G be the Galois group of E over F . Since $F(\delta)$ is a normal extension of F , it is mapped into itself by G . The next lemma tells us how G acts on $F(\delta)$.

Lemma 4. *Continuing the notation above, let σ be an element of G . Then*

$$\sigma(\delta) = (\det \sigma) \delta. \quad (2)$$

Proof. Since the roots α_i of L , $1 \leq i \leq n$, are an \mathbb{F}_q -basis of the space of roots of F , the action of σ on E is determined by its matrix with respect to this basis. We thus set S_σ to be the matrix of σ with respect to the α_i basis. A straightforward calculation using the fact that S_σ has entries in \mathbb{F}_q shows that we have the matrix equation

$$S_\sigma D = \sigma(D),$$

where $\sigma(D)$ is obtained from D by applying σ to the entries of D . We now take determinants on each side to obtain (2). \square

Corollary 5. *Let L be a monic q -polynomial of q -degree n in $F[x]$ and let E be its splitting field over F . Let G be the Galois group of L , considered as a subgroup of $GL(n, q)$. Then $G \leq SL(n, q)$ if and only if $\delta \in F$.*

Proof. If $G \leq SL(n, q)$ then $\sigma(\delta) = \delta$ for all $\sigma \in G$ by Lemma 4, and conversely. \square

We can now obtain a nice consequence of this analysis.

Corollary 6. *Let L be a monic q -polynomial of q -degree n in $F[x]$ and let E be its splitting field over F . Let the coefficient of x in L be $(-1)^n$. Let G be the Galois group of L , considered as a subgroup of $GL(n, q)$. Then $\det \sigma = 1$ for all $\sigma \in G$ and thus G is a subgroup of $SL(n, q)$.*

Proof. With the notation as before, $(-1)^n \delta^{q-1}$ is the coefficient of x in L and hence the hypotheses imply $\delta^{q-1} = 1$. This implies that $\delta \in \mathbb{F}_q$. Since we are assuming that \mathbb{F}_q is contained in F , $\delta \in F$, and so each element σ of G fixes δ . Equation (2) implies that $\det \sigma = 1$, as required. \square

We note that the same conclusion holds if the coefficient of x is $(-1)^n \mu^{q-1}$, where μ is any nonzero element of F .

3 Transitive actions on vector spaces

The results of the previous sections lead immediately to the following conclusion concerning a special type of Galois group action.

Theorem 7. *Let L be a monic q -polynomial of q -degree n in $F[x]$. Suppose that $L(x)/x$ is irreducible over F and the coefficient of x in L is $(-1)^n$. Then the Galois group G of L over F is isomorphic to a subgroup of the special linear group $SL(n, q)$ that acts transitively on the nonzero vectors of \mathbb{F}_q^n .*

We turn to consideration of a group G satisfying the conclusions of the theorem above when n is a prime, r , say. The main result of this paper is that if r is an odd prime and $q \neq 2$, then G must be equal to $SL(r, q)$.

We next provide some background detail. Let q be a power of p . The field \mathbb{F}_{q^n} is an n -dimensional vector space over \mathbb{F}_q , which we shall identify with the standard vector space \mathbb{F}_q^n . The multiplicative group $\mathbb{F}_{q^n}^*$ of \mathbb{F}_{q^n} , which is cyclic of order $q^n - 1$, acts on \mathbb{F}_q^n , and the set $\mathbb{F}_q^n \setminus 0$ of nonzero vectors is a single orbit under the group action. The Galois group of \mathbb{F}_{q^n} over \mathbb{F}_q , which is cyclic of order n and generated by the Frobenius q -th power element, also acts \mathbb{F}_q -linearly on \mathbb{F}_q^n and it normalizes the action of $\mathbb{F}_{q^n}^*$.

The resulting group generated by $\mathbb{F}_{q^n}^*$ and the Frobenius element acts on \mathbb{F}_q^n . We call this group the semilinear group of degree 1 and denote it by $\Gamma L(1, q^n)$. Thus $\Gamma L(1, q^n)$ is a subgroup of $GL(n, q)$ of order $n(q^n - 1)$. $\Gamma L(1, q^n)$ is the semidirect product of a normal cyclic subgroup, Z , say, of order $q^n - 1$ with a cyclic group of order n . The cyclic subgroup Z is a Singer cycle in $GL(n, q)$. It is known that the normalizer of Z in $GL(n, q)$ is $\Gamma L(1, q^n)$. Furthermore, the determinant homomorphism from $\Gamma L(1, q^n)$ into \mathbb{F}_q^* is surjective (see, for example, Satz 7.3, p.187, of [6]). Thus, if we set $\Gamma L_1(1, q^n) = \Gamma L(1, q^n) \cap SL(n, q)$, $\Gamma L_1(1, q^n)$ has order $n(q^n - 1)/(q - 1)$.

We study the action of $\Gamma L_1(1, q^n)$ in the cases q even and q odd separately.

3.1 q odd

The following lemma shows that $\Gamma L_1(1, q^n)$ is not transitive.

Lemma 8. *Let n and q both be odd. Then $\Gamma L_1(1, q^n) = \Gamma L(1, q^n) \cap SL(n, q)$ does not act transitively on $\mathbb{F}_q^n \setminus 0$.*

Proof. Suppose that $\Gamma L_1(1, q^n)$ acts transitively on $\mathbb{F}_q^n \setminus 0$. It follows then that $q^n - 1$ divides $n(q^n - 1)/(q - 1)$ and hence $q - 1$ divides n . This is impossible as $q - 1$ is even and n odd under the stated hypothesis. \square

3.2 q even

We need an improvement of Lemma 8 to deal with the case when q is a power of 2. We restrict ourselves to prime dimension, as this is the focus of our paper.

Lemma 9. *Let r be an odd prime and let q be a power of 2. Suppose that $\Gamma L_1(1, q^r) = \Gamma L(1, q^r) \cap SL(r, q)$ acts transitively on $\mathbb{F}_q^r \setminus 0$. Then one of the following holds:*

1. *The power q of 2 is 2.*
2. *The power q of 2 satisfies $q - 1 = r$. Furthermore, $|\Gamma L_1(1, q^r)| = q^r - 1$ and the action of $\Gamma L_1(1, q^r)$ on $\mathbb{F}_q^r \setminus 0$ is regular.*

Proof. Suppose that $\Gamma L_1(1, q^r)$ acts transitively on $\mathbb{F}_q^r \setminus 0$. Then by the proof of Lemma 8, $q - 1$ divides r . Thus since r is a prime, either $q - 1 = 1$, implying that $q = 2$, or $q - 1 = r$. Now, suppose that $q - 1 = r$. Then we have $|\Gamma L_1(1, q^r)| = q^r - 1$. Furthermore, since we are assuming that the action of $\Gamma L_1(1, q^r)$ is transitive, this action must be regular, given that $|\mathbb{F}_q^r \setminus 0| = q^r - 1$. \square

We note here that when $q = 2$, $\Gamma L_1(1, 2^n)$ acts transitively on $\mathbb{F}_q^n \setminus 0$ for all positive integers n . This is because $GL(n, 2) = SL(n, 2)$ and thus $\Gamma L_1(1, 2^n) = \Gamma L(1, 2^n)$. We want to show that the first case of Lemma 9 is the only case that can occur. This requires a more careful examination of the action of the Frobenius element on the field.

Lemma 10. *Let r be an odd prime and let q be a power of 2. Then $\Gamma L_1(1, q^r) = \Gamma L(1, q^r) \cap SL(r, q)$ does not act transitively on $\mathbb{F}_q^r \setminus 0$ when $q > 2$.*

Proof. Let α be a generator of $\mathbb{F}_{q^r}^*$ and define $\tau : \mathbb{F}_{q^r} \rightarrow \mathbb{F}_{q^r}$ by $\tau(z) = \alpha z$ for all $\alpha \in \mathbb{F}_{q^r}$. Then τ generates a Singer cycle Z , say, in $GL(r, q)$. Let σ be the Frobenius automorphism of \mathbb{F}_{q^r} defined by $\sigma(z) = z^q$ for all $z \in \mathbb{F}_{q^r}$. Then we find that $\sigma\tau\sigma^{-1} = \tau^q$, so that σ normalizes Z . Thus Z and σ generate a copy of $\Gamma L(1, q^r)$ in $GL(r, q)$.

The normal basis theorem for finite fields implies that \mathbb{F}_{q^r} has an \mathbb{F}_q -basis consisting of elements $\beta, \sigma(\beta), \dots, \sigma^{r-1}(\beta)$ for some element β of \mathbb{F}_{q^r} . Clearly, σ permutes the basis vectors as an r -cycle. It follows that $\det \sigma = 1$, since we are working in characteristic 2.

Now since σ normalizes Z , it also normalizes $Z_1 = Z \cap SL(r, q)$. Thus since σ is in $SL(r, q)$, the group generated by Z_1 and σ is a copy of $\Gamma L_1(1, q^r)$ in $SL(r, q)$. However, σ fixes the nonzero element 1 of \mathbb{F}_{q^r} , and hence an element of $\mathbb{F}_q^* \setminus 0$, whereas if $\Gamma L_1(1, q^r)$ acts transitively on $\mathbb{F}_q^* \setminus 0$, it acts regularly by Lemma 9. We deduce that it can only happen that the action of $\Gamma L_1(1, q^r)$ on $\mathbb{F}_q^* \setminus 0$ is not transitive. \square

Remark: The following observation is not relevant to the main theorem of this paper, but is perhaps of interest in as much as it illustrates the importance of various hypotheses that we have introduced. We consider the subgroup $\Gamma L_1(1, q^n)$ of $SL(n, q)$ and ask if there are any instances (other than when $q = 2$) when it acts transitively on $\mathbb{F}_q^n \setminus 0$. It is clear that for fixed n , only finitely many possibilities exist for q , since $q - 1$ must divide n if we have a transitive action. If we take say $q = 3$, we find that $\Gamma L_1(1, 3^n)$ does act transitively on $\mathbb{F}_3^n \setminus 0$ provided that $n \equiv 2 \pmod{4}$. In such circumstances, $\Gamma L_1(1, 3^n)$ has order $3^n - 1$ and its action is regular. In these examples, the Sylow 2-subgroup of $\Gamma L_1(1, 3^n)$ is quaternion of order 8 and thus $\Gamma L_1(1, 3^n)$ is not cyclic.

4 Proof of Main Theorem

We now present the classification of subgroups of $SL(r, q)$ that act transitively on $\mathbb{F}_q^r \setminus \{0\}$, which follows from results in the literature.

Theorem 11. *Let $q > 2$ be a prime power, let r be an odd prime, and let G be a subgroup of $SL(r, q)$ that acts transitively on the nonzero vectors of \mathbb{F}_q^r . Then either $G = SL(r, q)$ or $G \leq \Gamma L(1, q^r)$.*

A proof of Theorem 11 can be found in chapter 3 of the book [7] by Liebeck et al. The proof uses theorems of Hering [4], [5]. We note that Hering's work uses the classification of finite simple groups. We also note that there is a presentation given by Sambale, [8], Theorem 15.1, p.197, who provides some corrections and explanations to previously published statements, but this discussion deals only with q equal to a prime. The statement of Theorem 11 for q odd can also be derived from the statement of Proposition 3.2 in Ferraguti-Micheli [3].

Theorem 12. *Let r be an odd prime and let $q > 2$ be a power of a prime. Then the only subgroup of $SL(r, q)$ that acts transitively on $\mathbb{F}_q^r \setminus 0$ is $SL(r, q)$ itself.*

Proof. Let G be a subgroup of $SL(r, q)$ that enjoys the stated transitivity property. By Theorem 11, either $G = SL(r, q)$ or $G \leq \Gamma L(1, q^r)$. By Lemma 8 when q is odd, and Lemma 10 when q is even, $G \leq \Gamma L(1, q^r)$ is impossible. \square

We can now deduce the promised theorem relating to the Galois group of an irreducible q -polynomial whose q -degree is a prime.

Theorem 13. *Let F be a field of characteristic p that contains \mathbb{F}_q , where q is a power of p . Let L be a monic q -polynomial of q -degree r in $F[x]$, where r is an odd prime. Suppose that the coefficient of x in L is -1 and $L(x)/x$ is irreducible over F . Then the Galois group of L over F is isomorphic to the special linear group $SL(r, q)$, except possibly when $q = 2$.*

(More generally, the same conclusion holds if the coefficient of x in L is $-\mu^{p-1}$, where μ is any nonzero element of F .)

Theorem 13 is immediate from Theorem 7 and Theorem 12.

Example: We consider an example to illustrate Theorem 13. Let E be any field of characteristic p that contains \mathbb{F}_q . Let F be the function field $E(t)$, where t is an indeterminate over E . Let r be an odd prime and let $f(x)$ be a monic q -polynomial in $E[x]$ of q -degree r , with the coefficient of x in f equal to -1 . Let $g(x)$ also be a nonzero q -polynomial in $E[x]$ whose q -degree is less than r , with the coefficient of x in g equal to 0. Suppose also that $f(x)/x$ and $g(x)/x$ are relatively prime. Then the polynomial $f(x) + tg(x)$ in $F[x]$ is a monic q -polynomial of q -degree r , the coefficient of x is -1 , and $(f(x) + tg(x))/x$ is irreducible in $F[x]$. It follows that this polynomial has Galois group $SL(r, q)$ over F .

We remark that the hypotheses of Theorem 13 can never be satisfied when F is a finite field, as is easy to show directly.

5 Projective Polynomials

There is another Galois group related to the polynomial L described above, which we introduce now. Let $M(x) = L(x)/x$. Then it is easy to see that $M(x)$ is a monic polynomial in x^{q-1} , say $M(x) = P(x^{q-1})$. The polynomial P is monic of degree $(q^r - 1)/(q - 1)$ and is called the projective polynomial associated to L . Since we are assuming that M is irreducible, P is also irreducible over F .

Corollary 14. *Let F be a field of characteristic p that contains \mathbb{F}_q , where q is a power of p . Let L be a monic q -polynomial of q -degree r in $F[x]$, where r is an odd prime. Suppose that $L(x)/x$ is irreducible over F and the coefficient of x in L is -1 . Let $P(x)$ be the projective polynomial of degree $(q^r - 1)/(q - 1)$ associated to L . Then the Galois group of P over F is isomorphic to the projective special linear group $PSL(r, q)$.*

Proof. Given that $P(x^{q-1}) = L(x)/x$, it follows that the Galois group of P over F is a homomorphic image of that of L , since the splitting field of L over F contains that of P . Now we have shown that the Galois group of L over F is isomorphic to $SL(r, q)$. We consider two cases. Suppose that r does not divide $q - 1$. Then $SL(r, q)$ is a nonabelian finite simple group and $PSL(r, q)$ is isomorphic to $SL(r, q)$. The corollary is immediate in this case.

Suppose next that r divides $q - 1$. Then \mathbb{F}_q^* contains an element ω , say, of order r and similarly $SL(r, q)$ contains a central subgroup, Z , say, of order r generated by the scalar matrix ωI . The structure theory of special linear groups implies that Z is the unique proper normal subgroup of $SL(r, q)$. This follows since $PSL(r, q) = SL(r, q)/Z$ is a simple group (see, for example, Hauptsatz 6.13, p.182, of [6]). Since ωI acts trivially on the $(q - 1)$ -powers of the roots of L , and hence trivially on the roots of P , the homomorphism from the Galois group of L onto the Galois group of P contains Z in its kernel. The kernel is then precisely Z and hence the image is $PSL(r, q)$. \square

6 Polynomials of q -degree 2

We conclude the paper by considering q -polynomials of q -degree 2. As we mentioned earlier, we can employ a theorem of Dickson to prove reasonably complete results.

Theorem 15. *Suppose that F contains the field \mathbb{F}_q . Suppose also that the polynomial $M = x^{q^2-1} + bx^{q-1} + 1$ in $F[x]$ is irreducible in $F[x]$. Then the Galois group G of the polynomial over F is isomorphic to $SL(2, q)$, provided that q is not equal to any of 2, 3, 5, 7, 9 or 11.*

Proof. We note that xM is a q -polynomial of q -degree 2. It follows from Corollary 6 that G is a subgroup $SL(2, q)$. Furthermore, as we are assuming that M is irreducible, it follows that $q^2 - 1$ divides $|G|$.

We divide the rest of the proof into two cases. Suppose that q is a power of 2. Then $SL(2, q)$ is identical with $PSL(2, q)$ and G is thus a subgroup of $PSL(2, q)$ of order divisible by $q^2 - 1$. Thus since $PSL(2, q)$ has order $q(q^2 - 1)$, G has index a divisor of q . Theorem 262, p.286, of Dickson [2] shows that $PSL(2, q)$ has no subgroup of index less than $q + 1$ provided q is different from 2.

Suppose next that q is odd. Then $SL(2, q)$ contains a unique element of order 2, namely $-I$, and since $|G|$ is even, G contains $-I$. Let $\pi(G)$ be the image of G in $PSL(2, q)$. We have $|\pi(G)| = |G|/2$. A straightforward argument shows that the index of $\pi(G)$ in $PSL(2, q)$ divides q , and we again appeal to Dickson's theorem, [2], to show that no such subgroup exists when q is different from 3, 5, 7, 9 or 11 (but such subgroups do exist in the exceptional cases). \square

We remark that the case that $q = 2$ is certainly exceptional in the circumstances of Theorem 15. For if we take the polynomial $x^3 + x + 1$ in $\mathbb{F}_2[x]$, it is irreducible and thus its Galois group over \mathbb{F}_2 has order 3, not 6, the order of $SL(2, 2)$.

The following result is related to a result of Serre, found in an appendix to [1], p.131. This paper of Abhyankar [1] contains a substantial investigation of Galois groups in nonzero characteristic, as do subsequent papers by the author.

Corollary 16. *Suppose that F contains the field \mathbb{F}_q and the polynomial $x^{q^2-1} + bx^{q-1} + 1$ in $F[x]$ is irreducible in $F[x]$. Then the Galois group of the polynomial*

$x^{q+1} + bx + 1$ over F is isomorphic to $PSL(2, q)$ provided that q is not equal to any of 2, 3, 5, 7, 9 or 11.

We omit the proof, as it follows from the proof of Theorem 15, using the argument of Corollary 14.

References

- [1] Abhyankar, S. S. Galois theory on the line in nonzero characteristic. Bull. Amer. Math. Soc. **27** (1992), 68-133.
- [2] Dickson, L. E. Linear groups, with an exposition of the Galois field theory. B. G. Teubner, Leipzig, 1901.
- [3] Ferraguti, A., Micheli, G., Exceptional scatteredness in prime degree, J. Alg. **565** (2021) 691-701. <https://doi.org/10.1016/j.jalgebra.2020.09.034>
- [4] Hering, C. Transitive linear groups and linear groups which contain irreducible subgroups of prime order. Geometriae Dedicata **2** (1974), 425-460.
- [5] Hering, C. Transitive linear groups and linear groups which contain irreducible subgroups of prime order II. J. Algebra **93** (1985), 151-164.
- [6] Huppert, B. Endliche Gruppen I. Grundlehren der Mathematischen Wissenschaften, Vol. 134. Springer-Verlag, Berlin, 1967.
- [7] Liebeck, Martin W., Praeger, C., Saxl, J., Regular Subgroups of Primitive Permutation Groups, American Mathematical Society, 2010.
- [8] Sambale, B. Blocks of finite groups and their invariants. Lecture Notes in Mathematics, Vol. 2127. Springer, 2014.