

**Weighted Grand Lebesgue Spaces norm estimation  
for Hardy - Sobolev - Poincare - Wirtinger operators.**

**M.R.Formica, E.Ostrovsky, and L.Sirota.**

Università degli Studi di Napoli Parthenope, via Generale Parisi 13, Palazzo Pacanowsky, 80132, Napoli, Italy.

e-mail: mara.formica@uniparthenope.it

Department of Mathematics and Statistics, Bar-Ilan University, 59200, Ramat Gan, Israel.

e-mail: eugostrovsky@list.ru

Department of Mathematics and Statistics, Bar-Ilan University,  
59200, Ramat Gan, Israel.

e-mail: sirota3@bezeqint.net

*Abstract.*

We extend the classical Hardy - Sobolev - Poincare - Wirtinger inequalities from the ordinary Lebesgue - Riesz spaces into the Grand Lebesgue ones, with exact constants evaluation.

*Key words and phrases.* Hardy - Sobolev - Poincare - Wirtinger inequalities, domain, distance from the boundary, upper and lower estimates and limits, Lebesgue - Riesz, dilation method, Sobolev's and Grand Lebesgue Space norm and spaces, vector, dilatation and parameter of dilatation, Talenty's method, gradient, convex and property set, operators, examples.

## **1 Definitions. Notations. Statement of problem. Previous results.**

Let  $D$  be certain proper own:  $D \neq R^n$  sub - domain of the whole space  $R^n$ , equipped with ordinary Euclidean norm  $|x|$ ,  $x \in R^n$ , for instance open, connected,

convex, having non - trivial interior and Lipschitz boundary  $\partial D$ . The distance between arbitrary vector  $x \in R^n$  and boundary  $\partial D$  will be denoted as follows

$$d(x) = d(x, \partial D) = \inf_{y \in \partial D} |x - y|.$$

Let also  $u = u(x)$ ,  $x \in D$  be arbitrary valued function belonging to the Sobolev's space  $W^1(p)$  :

$$\|u\|_{W_p^1} \stackrel{def}{=} \|\nabla u\|_p^{1/p} + \|u\|_p < \infty,$$

where the usually Lebesgue - Riesz norm  $\|u\|_p$  has the form

$$\|u\|_p := \left[ \int_D |u(x)|^p dx \right]^{1/p}, 1 \leq p < \infty;$$

$\nabla(u)$  denotes the gradient in an distributional sense of the function  $u$ .

The following (and close) key *weight* Hardy - Sobolev - Poincare - Wirtinger (briefly, HSPW) inequality

$$\int_D \frac{|\nabla u|^p dx}{d^\alpha(x, \partial D)} \geq \left( \frac{\alpha + p - n}{p} \right)^p \cdot \int_D \frac{|u|^p dx}{d^{p+\alpha}(x, \partial D)}, p > n - \alpha, \alpha = \text{const} < n \quad (1)$$

belongs to many authors, see e.g. [1], [2], [3], [4], [18], [19], [27]. See also [12], [20], [21], [26], [28], [29] etc.

It may be rewritten as follows. Introduce the following (Borelian) measure  $\mu_\alpha = \mu_{\alpha, D}$  on the measurable subsets of the source set  $D$

$$\mu_{\alpha, D}(dx) \stackrel{def}{=} \frac{dx}{d^\alpha(x, \partial D)} \quad (2)$$

and put

$$K(p) = K(p; \alpha, n) := \frac{p}{p + \alpha - n}, p > n - \alpha. \quad (3)$$

Define also the (linear) operator defined on all the functions having as a domain of definition the set  $D \setminus \partial D$  :

$$T[u](x) \stackrel{def}{=} \frac{u(x)}{d(x, \partial D)}. \quad (4)$$

Then when  $p > n - \alpha$

$$\|T[u]\|_{L_p(D, \mu_\alpha)} \leq K(p) \cdot \|\nabla u\|_{L_p(D, \mu_\alpha)}. \quad (5)$$

Herewith the "constant"  $K = K(p)$  is the best possible for arbitrary proper domain  $D$  :

$$\sup_{u: \|\nabla u\|_{L_p(D, \mu_\alpha)}=1} \|T[u]\|_{L_p(D, \mu_\alpha)} = K(p), p > n - \alpha. \quad (6)$$

Our claim in this short preprint is to extrapolate the last estimate onto the so-called Grand Lebesgue Spaces instead the classical Lebesgue - Riesz ones, builded on the our set  $D$  equipped with the measure  $\mu_{\alpha,D}$ .

We intent herewith to calculate the exact value of correspondent embedding constants.

The particular case of these statement was considered in [26]; see also [23] - [25].

Let us recall here for readers convenience some known definitions and facts from the theory of Grand Lebesgue Spaces (GLS) adapted exclusively to offered article.

Let the number  $b$  be constants such that  $n - \alpha < b \leq \infty$ ; and let  $\psi = \psi(p) = \psi[b](p)$ ,  $p \in (n - \alpha, b)$ , be numerical valued strictly positive function not necessary to be finite in every point:

$$\inf_{p \in (n - \alpha, b)} \psi[b](p) > 0. \quad (7)$$

The set of all such a functions  $\psi(\cdot)$  will be denoted by  $\Psi(b) = \{ \psi = \psi(\cdot) \}$ .

For instance

$$\psi_m(p) := p^{1/m}, \quad m = \text{const} > 0,$$

or more generally

$$\psi_{m,\beta,L}(p) := p^{1/m} \ln^\beta(p + 1) L(\ln(p + 1)), \quad \beta = \text{const}, \quad (8)$$

where  $L(\cdot)$  is positive continuous slowly varying as  $p \rightarrow \infty$  function.

### Definition 1.1.

By definition, the (Banach) Grand Lebesgue Space (GLS)  $G\psi = G\psi[b]$ , consists on all the real (or complex) numerical valued measurable functions  $f : D \rightarrow R$  defined on the whole our space  $\Omega$  and having a finite norm

$$\| f \|_{G\psi} = \| f \|_{G\psi[b]} \stackrel{\text{def}}{=} \sup_{p \in (n - \alpha, b)} \left[ \frac{\| f \|_{p,D,\mu_\alpha}}{\psi(p)} \right]. \quad (9)$$

The function  $\psi = \psi(p) = \psi[b](p)$  is named as the *generating function* for this space.

If for instance

$$\psi(p) = \psi^{(r)}(p) = 1, \quad p = r; \quad \psi^{(r)}(p) = +\infty, \quad p \neq r,$$

where  $r = \text{const} \in [1, \infty)$ ,  $C/\infty := 0$ ,  $C \in R$ , (an extremal case), then the correspondent  $G\psi^{(r)}(p)$  space coincides with the classical Lebesgue - Riesz space  $L_r = L_r(\Omega, \mathbf{P})$ .

The finiteness of some GLS  $G\psi$  norm for the function  $f : D \rightarrow R$  is closely related with its tail function

$$T_f(u) := \mu_\alpha\{x : |f(x)| > u\}, \quad u \geq 0.$$

The GLS spaces are also closely related with the suitable exponential Orlicz ones, builded on the our measurable space  $(D, \mu_\alpha)$ .

See the detail investigation of these spaces in the works [5], [6] - [7], [9], [10], [8], [11], [15] - [17], [22], chapters 1,2.

## 2 Main result.

**Theorem 2.1.** Suppose that for certain function  $u = u(x)$ ,  $x \in D$  its gradient belongs to some Grand Lebesgue Space  $G\psi[b]$  :

$$\|\nabla u\|_{G\psi[b]} < \infty. \quad (10)$$

Introduce an auxiliary such a function

$$\psi_K(p) := K(p) \cdot \psi(p), \quad n - \alpha < p < b.$$

Our proposition:

$$\|T[u]\|_{G\psi_K} \leq 1 \times \|\nabla u\|_{G\psi}, \quad (11)$$

where the constant "1" in (11) is the best possible.

**Remark 2.1.** The generating function  $\psi[b](\cdot)$  in (10) may be choosed by the natural way:

$$\psi[b](\cdot) := \|\nabla u\|_p, \quad n - \alpha < p < b,$$

of course, if there exists such a value  $b > n - \alpha$ .

**Proof of upper bound** is very simple and is alike to one in the articles [24], [25]; as well as the proof of the lower bound. Suppose

$$\|\nabla u\|_{G\psi} < \infty;$$

one can assume without loss of generality

$$\|\nabla u\|_{G\psi} = 1.$$

It follows from the direct definition of the norm in GLS

$$\forall p \in (n - \alpha, b) \Rightarrow \|\nabla u\|_p \leq \psi(p).$$

We apply the inequality (5)

$$\|T[u]\|_{L_p(D, \mu_\alpha)} \leq K(p) \cdot \psi(p) = \psi_K(p),$$

and on the other words

$$\|T[u]\|_{G\psi_K} \leq 1 = \|\nabla u\|_{G\psi}.$$

**The lower bound** in (11) follows immediately from one of the results of the article [25]; see also [24], taking into account the *exactness* of the value  $K = K(p)$ , see (6).

Q.E.D.

### 3 Necessary conditions for these estimations.

We will ground in this section that the "configuration" given by the estimate (1) is essentially non - improvable. Namely, suppose that there exists non - trivial constants  $a, h; G(D, a, h)$  such that for *arbitrary* proper non - trivial sub - domain  $D \subset R^n$  and for *arbitrary* function  $u = u(x)$ ,  $x \in D$  belonging to the space  $C_\infty^0(D)$  there holds the estimate

$$L[u] := L[u](a) \stackrel{def}{=} \left[ \int_D \frac{|u(x)|^q}{d^a(x, \partial D)} dx \right]^{1/q} \leq \quad (12)$$

$$R[u] := R[u](h) \stackrel{def}{=} G(D, a, h) \left[ \int_D \frac{|\nabla u|^p}{d^h(x, \partial D)} dx \right]^{1/p} \quad (13)$$

for certain fixed values  $p, q \in [1, \infty)$ .

**Theorem 3.1.** Suppose that the relations (12) and (13) holds true for arbitrary proper domain  $D$  as well as for certain non - zero function  $u = u(x)$ ,  $x \in D$  from the set  $C_\infty^0(D)$ . Then

$$\frac{a - n}{q} = 1 + \frac{h - n}{p}. \quad (14)$$

**Proof.** We will apply the so - called dilation method, belonging at first, perhaps, to G.Talenty, see the article [29], in which was considered the particular case  $a = h = 0$ .

Let us choose the domain  $D$  on the form

$$D := R^d \otimes R_+^r, \quad n = \dim D = d + r, \quad r, d > 0.$$

Let's agree to write instead of  $x$  the vector  $(x, y)$ ,  $x \in R^d$ ,  $y \in R_+^r$ . Then

$$L^q[u] = \int_{R^d} \int_{R_+^r} \frac{|u(x, y)|^q dx dy}{|y|^a},$$

$$R^p[u] = \int_{R^d} \int_{R_+^r} \frac{|\nabla(x, y)|^p dx dy}{|y|^h}.$$

One can choose the function  $u = u(x, y)$  from the set  $C_\infty^0(D)$  such that  $L(u) > 0$ ,  $R[u] > 0$ ; therefore  $0 < L[u] \leq CR[u] < \infty$ .

Let also  $\lambda \in (0, \infty)$  be a parameter of dilatation:

$$V_\lambda[u](x, y) \stackrel{def}{=} u(\lambda x, \lambda y).$$

Obviously,  $V_\lambda[u](\cdot, \cdot) \in C_\infty^0(D)$ . Therefore, it satisfies the inequality (12) - (13):

$$L_a[V_\lambda u] \leq CR_h[V_\lambda u]. \quad (15)$$

But

$$\{L_a[V_\lambda u]\}^q = \lambda^{-d-r+a} L^q[u];$$

$$L_a[V_\lambda u] = \lambda^{(a-d-r)/q} L_a[u]$$

and analogously

$$R_h[V_\lambda u] = \lambda^{(p-d-r+h)/p} R_h[u],$$

and we get to the inequality

$$C_1 \lambda^{(a-d-r)/q} \leq C_2 \lambda^{(p-d-r+h)/p}$$

for arbitrary positive values  $\lambda$ . Following,

$$\frac{a-d-r}{q} = \frac{p-d-r+h}{p},$$

which is equal to (14).

Of course, in the case when  $a \cdot h = 0$  or  $p = q$  we get to the classical Sobolev's inequality, see e.g. [29].

## 4 Concluding remarks.

*Open question:* Suppose that there holds (14) and that the domain  $D$  is proper. Assume also the function  $u = u(x)$  belongs to the set  $C_\infty^0(D)$ . Will be the inequality (12) - (13) true, of course, in general case, i.e. when  $p \neq q$ ?

*Acknowledgement.* The first author has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and by Università degli Studi di Napoli Parthenope through the project “sostegno alla Ricerca individuale” (triennio 2015 - 2017).

## References

- [1] **F.G.Avkhadiev.** *Hardy type inequalities in higher dimensions with explicit estimate of constants.* Lobachevskii J. Math., **21**, (2006), 3 - 31.
- [2] **F. G. Avkhadiev.** *Families of domains with best possible Hardy constant.* Russian Mathematics, (Iz. VUZ), **57**, (2013), 49 - 52.
- [3] **G.Barbatis, S.Filippas and A.Tertikas.** *A unified approach to improved  $L_p$  Hardy inequalities with best constants.* Trans. Amer. Math. Soc., **356**, (2004), 2169 - 2196.
- [4] **B.Devyver and Y.Pinchover.** *Optimal  $L_p$  Hardy - type inequalities.* Ann. Inst. H. Poincaré. Anal. Non Linéaire 33 (2016), 93 - 118.
- [5] **V. Ermakov, and E. I. Ostrovsky.** *Continuity Conditions, Exponential Estimates, and the Central Limit Theorem for Random Fields.* Moscow, VINITI, (1986), (in Russian).
- [6] **A.Fiorenza.** *Duality and reflexivity in grand Lebesgue spaces.* Collectanea Mathematica, (electronic version), 51, 2, (2000), 131 - 148.
- [7] **A. Fiorenza and G.E.Karadzhov.** *Grand and small Lebesgue spaces and their analogs.* 2005, Consiglio Nazionale Delle Ricerche, Istituto per le Applicazioni del Calcolo Mauro Picone, Sezione di Napoli, Rapporto tecnico n., 272/03.
- [8] **A. Fiorenza, M. R. Formica and A. Gogatishvili.** *On grand and small Lebesgue and Sobolev spaces and some applications to PDE's.* Differ. Equ. Appl. **10** (2018), no. 1, 21–46.
- [9] **A. Fiorenza, M. R. Formica, A. Gogatishvili, T. Kopaliani and J. M. Rakotoson.** *Characterization of interpolation between grand, small or classical Lebesgue spaces.* Preprint arXiv:1709.05892, Nonlinear Anal., to appear.
- [10] **A. Fiorenza, M. R. Formica and J. M. Rakotoson.** *Pointwise estimates for  $G\Gamma$ -functions and applications.* Differential Integral Equations **30** (2017), no. 11-12, 809–824.
- [11] **M. R. Formica and R. Giova.** *Boyd indices in generalized grand Lebesgue spaces and applications.* Mediterr. J. Math. **12** (2015), no. 3, 987–995.

- [12] **G.H.Hardy, J.E.Littlewood and G.Polya.** *Inequalities.* Cambridge, (1952).
- [13] **T.Iwaniec and C. Sbordone.** *On the integrability of the Jacobian under minimal hypotheses.* Arch. Rat. Mech. Anal., 119, (1992), 129 - 143.
- [14] **T.Iwaniec, P.Koskela and J.Onninen.** *Mapping of finite distortion: Monotonicity and Continuity.* Invent. Math., 144, (2001), 507 - 531.
- [15] **Kozachenko Yu. V., Ostrovsky E.I.** (1985). *The Banach Spaces of random Variables of subgaussian Type.* Theory of Probab. and Math. Stat., (in Russian). Kiev, KSU, **32**, 43-57.
- [16] **Yu. V. Kozachenko, E. I. Ostrovsky and L. Sirota.** *Relations between exponential tails, moments and moment generating functions for random variables and vectors.* arXiv:1701.01901v1 [math.FA], 8 Jan 2017.
- [17] **Yu. V. Kozachenko, E. I. Ostrovsky and L. Sirota.** *Equivalence between tails, Grand Lebesgue Spaces and Orlicz norms for random variables without Cramer's condition.* arXiv:1710.05260v1 [math.PR], 15 Oct 2017.
- [18] **M. Marcus, V. J. Mizel and Y. Pinchover.** *On the best constant for Hardy's inequality in  $R^n$ .* Trans. Amer. Math. Soc., **350**, (1998), 3237 - 3255.
- [19] **T.Matskewich and P.E.Sobolevskii.** *The best possible constant in a generalized Hardy's inequality for convex domains in  $R^n$ .* Nonlinear Anal., **28**, (1997), 1601 - 1610.
- [20] **V.G. Maz'ya.** *Sobolev Spaces.* Springer Verlag, Berlin - New York, 1985.
- [21] **D.S.Mitrinovic, J.E.Pecaric and A.M.Fink.** *Inequalities Involving Functiona and Their Integrals and Derivatives.* Kluvner Academic Verlag, (1996), Dorderecht, Boston, London.
- [22] **Ostrovsky E.I.** (1999). *Exponential estimations for Random Fields and its applications,* (in Russian). Moscow-Obninsk, OINPE.
- [23] **Ostrovsky E., Sirota L.** *Grand Lebesgue Spaces norm estimates for multivariate functional operations.* arXiv:1805.01982v1 [math.FA] 5 May 2018
- [24] **Ostrovsky E., Sirota L., and Rogover E.** *Integral Operators in bilateral Grand Lebesgue Spaces.* arXiv:0912.2538v1 [math.FA] 13 Dec 2009.
- [25] **Ostrovsky E.** *Boundedness of Operators in Bilateral Grand Lebesgue Spaces with Exact and Weakly Exact Constant Calculation.* arXiv:1104.2963v1[math.FA] 15 Apr 201134.



- [26] **Ostrovsky E., Sirota L., and Rogover E.** *Poincare inequalities in bilateral Grand Lebesgue Spaces.*  
arXiv:0908.0546v1 [math.FA] 4 Aug 2009
- [27] **Yeguda Pinchover and Divya Goel.** *On weighted  $L_p$  - Hardy inequality on domains in  $R^n$ .*  
arXiv:2012.12860v1 [math.AP] 23 Dec 2020
- [28] **Sobolev, S.L.** (1963). *Some applications of functional analysis in mathematical physics.* Amer. Math. Soc..
- [29] **G.Talenty.** *Best constant in Sobolev inequality.* Instit. Mat. Univ. Firenze, **22**, (1974-1975), p. 1 - 32.
- [30] **A.Wannebo.** *Hardy Inequalities.* Proc. Amer. Math. Soc., **109**. (1990), 85 - 95.