

# The flat phase of quenched disordered membranes at three-loop order

S. Metayer<sup>1,\*</sup> and D. Mouhanna<sup>2,†</sup>

<sup>1</sup>*Sorbonne Université, CNRS, Laboratoire de Physique Théorique et Hautes Energies, LPTHE, 75005 Paris, France*

<sup>2</sup>*Sorbonne Université, CNRS, Laboratoire de Physique Théorique de la Matière Condensée, LPTMC, 75005 Paris, France*

We study quenched disordered polymerized membranes in their flat phase by means of a three-loop perturbative analysis performed in dimension  $D = 4 - \epsilon$ . We derive the renormalization group equations at this order and solve them up to order  $\epsilon^3$ . Our results confirm those obtained by Coquand *et al.* within a nonperturbative approach [Phys. Rev. E 97, 030102 (2018)] predicting a finite-temperature, finite-disorder *wrinkling* transition and those obtained by Coquand and Mouhanna within a recent two-loop order approach [Phys. Rev. E 103, 031001 (2021)], while correcting some of the results obtained in this last reference. We compute the anomalous dimensions that characterize the scaling behaviour at the various fixed points of the renormalization group flow diagram. They appear to be in strong agreement with those predicted within the nonperturbative context.

*Introduction.* Understanding the effects of quenched disorder in the flat phase of polymerized membranes has become a major challenge with, as a priority target, unveiling new phenomena in the physics of graphene and graphene-like materials [1–5] going from mechanical ones – e.g. a paradoxical enhancement of elasticity modulus with the density of defects [6] – to electronic ones – e.g. the possibility to open a tunable band gap [7–9]. However, the interest for quenched disorder in membranes has a longer history and goes back to the early experiments of Sackman *et al.* [10] followed by those of Mutz *et al.* [11] and Chaieb *et al.* [12–15] on partially polymerized lipid membranes. These authors have shown that, upon cooling below the melting temperature, these systems undergo a phase transition from a smooth structure at low-disorder or, equivalently, high polymerization to a wrinkled structure at high disorder, or low polymerization.

These investigations have stimulated an important theoretical work aiming to identify precisely the nature of this weakly polymerized, wrinkled, phase that has been conjectured to coincide with a *glassy* phase, a phase mainly controlled by *disorder* fluctuations. Nelson and Radzihovsky [16, 17], using a *one-loop* perturbative approach in the vicinity of the upper critical dimension  $D = 4$  have shown the irrelevance of a disorder acting only on the internal metric of the membrane. They have shown that the renormalization group (RG) flow was driven toward the disorder-free fixed point, called  $P_4$ , identified by Aronovitz and Lubensky [18] in their early RG approach of pure membranes. This result has then been confirmed by Radzihovsky and Le Doussal [19, 20] in the context of a leading order self-consistent screening approximation (SCSA). Morse *et al.* [21, 22] have then extended the one-loop study of Nelson and Radzihovsky by adding a curvature disorder to the metric one. They have confirmed the irrelevance of the disorder below  $D = 4$  and discovered a new, vanishing temperature,

fixed point, called  $P_5$ . This fixed point has been identified, within the one-loop computation of Morse *et al.*, as being stable with respect to the disorders but unstable in the direction associated with the temperature, making the fixed point  $P_5$  non-pertinent in the prospect of a glassy phase. These works have then been followed by a new approach relying on the use of the so-called nonperturbative renormalization group (NPRG) by Coquand *et al.* [23] on the metric/curvature-disordered model initially considered by Morse *et al.* [21, 22]. This approach is based on the use of an exact equation that controls the RG flow of an effective action  $\Gamma_k$  where  $k$  is a running scale. The quantity  $\Gamma_k$  is truncated in powers of the field and field-derivatives while keeping nonpolynomial contributions coming from the usual perturbative parameters: coupling constants, temperature,  $1/N - N$  being the number of components of the order parameter – and so on (see [24–29] for the use of this technique in the context of disorder-free membranes). A striking result obtained by means of this approach [23] is the discovery of a finite-temperature, finite-disorder, fixed point  $P_c$ , *unstable* with respect to the temperature, making the vanishing temperature fixed point  $P_5$  *fully attractive* at sufficiently low temperatures. This approach has thus confirmed theoretically, for the first time, the possibility of a whole glassy phase at low temperatures in quenched disordered membranes. Moreover, the various scaling laws observed by Chaieb *et al.* [12–15] in their investigations of partially polymerized lipid membranes have been qualitatively and quantitatively explained [30] on the basis of the analysis performed in [23].

Although convincing the NPRG approach of Coquand *et al.* [23] has happened to be at odds with the results obtained from the SCSA approach of Le Doussal and Radzihovsky [20] in which the fixed point  $P_c$  is missing. This is notably for this reason that, very recently, a *two-loop order* perturbative approach in the vicinity of  $D = 4$  has been performed by Coquand and Mouhanna [31], following the early one-loop order computation of Morse *et al.* [21, 22] and the two-loop order ones performed on disorder-free membranes by Mauri and Katnelson [32] and Coquand *et al.* [33]. This approach

\*Electronic address: smetayer@lpthe.jussieu.fr

†Electronic address: mouhanna@lptmc.jussieu.fr

has confirmed the existence of a fixed point  $P_c$  associated with a phase transition between a high-temperature phase [41] controlled by the disorder-free fixed point  $P_4$  and a low-temperature phase controlled by the vanishing-temperature, infinite-disorder, fixed point  $P_5$ . However, this approach has encountered difficulties taking the form of an indetermination of the coordinates of the various fixed points  $P_5$  and  $P_c$  as well as of the corresponding anomalous dimensions.

In order to clarify this situation we investigate here the flat phase of quenched disorder membranes by means of a *three-loop* order computation in the vicinity of  $D = 4$ , following the very recent approach performed in the pure case by Metayer *et al.* [34]; see also [35] for a four-loop computation. We show that the indetermination discussed above was associated to an incorrect expres-

sion for the RG function of the curvature disorder. We provide here the correct expressions of the RG functions at three-loop order and determine all physical quantities up to order  $\epsilon^3$  without ambiguity. Our results confirm, for the first time within the perturbative context, the existence of a new fixed point  $P_c$  in the flat phase of quenched polymerized membranes, even if it is found to be marginally – to order  $O(\epsilon^2)$  – stable in contradiction with the result of the NPRG approach. However, the anomalous dimensions computed at the various fixed points  $P_5$  and  $P_c$  are in strong agreement with those predicted within the NPRG approach.

*The action.* The action relevant to study the flat phase of quenched disordered  $D$ -dimensional membranes embedded in a  $d$ -dimensional euclidean space, is given by [21, 22]:

$$S = \int d^D x \left\{ \frac{\tilde{\kappa}^{\alpha\beta}}{2} \Delta \mathbf{h}^\alpha(\mathbf{x}) \Delta \mathbf{h}^\beta(\mathbf{x}) + \frac{\tilde{\lambda}^{\alpha\beta}}{2} u_{ii}^\alpha(\mathbf{x}) u_{jj}^\beta(\mathbf{x}) + \tilde{\mu}_{\alpha\beta} u_{ij}^\alpha(\mathbf{x}) u_{ij}^\beta(\mathbf{x}) \right\}. \quad (1)$$

In this expression the field  $\mathbf{h}^\alpha(\mathbf{x})$  describes the  $d - D$  flexural modes that parameterize, at each point  $\mathbf{x}$  of the membrane, the height, transverse, fluctuations with respect to a fully flat configuration while  $u_{ij}^\alpha(\mathbf{x})$  is the strain tensor given by

$$u_{ij}^\alpha \simeq \frac{1}{2} [\partial_i u_j^\alpha + \partial_i u_j^\alpha + \partial_i \mathbf{h}^\alpha \cdot \partial_j \mathbf{h}^\alpha]$$

where  $\mathbf{u}^\alpha$  represent  $D$  longitudinal – phonon – modes describing the elastic, transversal, fluctuations with respect to the flat configuration. Note that in all the expressions above the Greek indices are associated with the  $n$  replica that are used to performed the average over the disorder, see [21–23, 31]. The coupling constants entering in Eq.(1),  $\tilde{\kappa}^{\alpha\beta} = \tilde{\kappa} \delta^{\alpha\beta} - \tilde{\Delta}_\kappa J^{\alpha\beta}$ ,  $\tilde{\lambda}^{\alpha\beta} = \tilde{\lambda} \delta^{\alpha\beta} - \tilde{\Delta}_\lambda J^{\alpha\beta}$  and  $\tilde{\mu}^{\alpha\beta} = \tilde{\mu} \delta^{\alpha\beta} - \tilde{\Delta}_\mu J^{\alpha\beta}$ , where  $J^{\alpha\beta} \equiv 1, \forall \alpha, \beta$ , encode the bending rigidity  $\tilde{\kappa}$ , the elastic coupling constants – Lamé coefficients –  $\tilde{\lambda}$  and  $\tilde{\mu}$  and, finally, the disorder variances  $\tilde{\Delta}_\kappa$ , associated to the curvature disorder and  $(\tilde{\Delta}_\lambda, \tilde{\Delta}_\mu)$  associated to the metric disorder. As usual, see [21–23, 31], the temperature  $T$  has been re-absorbed in the definition of the coupling constants:  $\{\tilde{g}\} \equiv \{\kappa/T, \lambda/T, \mu/T, \Delta_\lambda/T^2, \Delta_\mu/T^2, \Delta_\kappa/T^2\}$ . Note that stability considerations imply that  $\mu, \lambda + 2\mu/D, \Delta_\kappa, \Delta_\mu$  and  $\Delta_\lambda + 2\Delta_\mu/D$  should be all positive.

Finally we define, as in [21–23, 31], the relevant correlation functions. Writing  $\delta \mathbf{h}(\mathbf{q}) = \mathbf{h}(\mathbf{q}) - \langle \mathbf{h}(\mathbf{q}) \rangle$  one has, denoting by  $[\dots]$  an average over a Gaussian disorder and by  $\langle \dots \rangle$  a thermal average:

$$G_{h_i h_j}(\mathbf{q}) = [\langle h_i(\mathbf{q}) h_j(-\mathbf{q}) \rangle] = T \chi_{h_i h_j}(\mathbf{q}) + C_{h_i h_j}(\mathbf{q})$$

where  $T \chi_{h_i h_j}(\mathbf{q}) = [\langle \delta h_i(\mathbf{q}) \delta h_j(-\mathbf{q}) \rangle]$  and  $C_{h_i h_j}(\mathbf{q}) =$

$[\langle h_i(\mathbf{q}) \rangle \langle h_j(-\mathbf{q}) \rangle]$  that behave, at low momenta, as:

$$\chi_{h_i h_j}(\mathbf{q}) \sim q^{-(4-\eta)}, \quad C_{h_i h_j}(\mathbf{q}) \sim q^{-(4-\eta')}$$

and are controlled by the exponents  $\eta$  and  $\eta'$  associated to thermal and disorder fluctuations, respectively. One defines analogous correlation functions for the phonon fields  $\mathbf{u}$  with critical exponents  $\eta_u$  and  $\eta'_u$  that are related to the previous ones by Ward identities [21–23, 31]:  $\eta_u + 2\eta = 4 - D$  and  $\eta'_u + 2\eta' = 4 - D$ . Finally, a relevant quantity obtained from  $\eta$  and  $\eta'$  is the exponent  $\phi$  given by [21, 22]  $\phi = \eta' - \eta$  that determines which kind of fluctuations dominates at a given fixed point:  $\phi > 0$ , respectively  $\phi < 0$ , corresponds to a fixed point where thermal, respectively disorder, fluctuations dominate. For  $\phi = 0$  both kinds of fluctuations coexist; the corresponding fixed point is then said to be marginal.

*The renormalization group equations and fixed points.* We have derived the three-loop order RG equations for model (1) within the modified minimal subtraction scheme and in the massless case, following the procedure used in [34]. In particular, we have made intensive use of techniques of massless Feynman diagram calculations, see, e.g., the review [36], and have recourse to Qgraf [37] for the generation of the diagrams, Mathematica to perform the numerator algebra and LiteRed [38, 39] to reduce the loop integrals to a finite set of master integrals. As in the disorder-free case [34] we had to evaluate 51 diagrams; here their evaluation is complicated by the heavy algebra associated to the replica structure. Computation details will be given in a forthcoming publication [40] while, in this paper, we focus on the analysis of the RG functions. The latter are too long

to be displayed in the main text; they are given in App.A up to two-loop order while the full three-loop expressions are given in a supplementary materials [42]. They are expressed in terms of dimensionless renormalized quantities:  $\bar{g} = k^{D-4} Z^2 Z_g^{-1} \tilde{\kappa}^{-2} \tilde{g}$  for  $g \in \{\mu, \lambda, \Delta_\mu, \Delta_\lambda\}$  and  $\bar{\Delta}_\kappa = Z Z_{\Delta_\kappa}^{-1} \tilde{\kappa}^{-1} \tilde{\Delta}_\kappa$  where  $Z_\alpha$  is the coupling constant renormalization for any coupling  $\alpha$  and  $Z$  is the field renormalization defined through  $\mathbf{h} = Z^{1/2} \tilde{\kappa}^{-1/2} \mathbf{h}_R$  and  $\mathbf{u} = Z \tilde{\kappa}^{-1} \mathbf{u}_R$ ,  $\mathbf{u}_R$  and  $\mathbf{h}_R$  being the renormalized fields. Note that with these definitions the scaling of the renormalized coupling constants with the temperature is given by  $\bar{\lambda} \sim \bar{\mu} \sim T$  (that can be considered as measures of the temperature  $T$ ),  $\bar{\Delta}_\kappa \sim 1/T$ , while the other coupling constants are temperature-independent. The running anomalous dimension  $\eta_t$  and the running exponent  $\phi_t$  are given by  $\eta_t = -\partial_t \ln Z$  and  $\phi_t = \eta'_t - \eta_t = \partial_t \ln \bar{\Delta}_\kappa$  with  $t = \ln k$ ,  $k$  being a renormalization momentum scale.

We discuss the fixed points of the RG equations and the corresponding field anomalous dimensions, whose expressions are explicitly provided at three-loop order in App.C. We first recall the one-loop order result [21, 22], then we provide our two-loop order results, while correcting those of Ref.[31] and, finally, our three-loop order ones.

*One-loop order.* At one-loop order one has, in  $D < 4$ , besides the Gaussian fixed point, two non-trivial fixed point [43] located on the hypersurfaces  $\bar{\lambda}/\bar{\mu} = \bar{\Delta}_\lambda/\bar{\Delta}_\mu = -1/3$ :

1) the disorder-free fixed point,  $P_4$ , for which  $\bar{\mu} = 96\pi^2 \epsilon / (24 + d_c)$ ,  $\bar{\Delta}_\mu = \bar{\Delta}_\kappa = 0$  and  $\eta = \eta'/2 = \phi = 12 \epsilon / (24 + d_c)$ . This attractive fixed point controls the long distance behaviour of both disorder-free and disordered membranes – see below.

2) The vanishing temperature, infinite disorder, fixed point,  $P_5$ , for which  $\bar{\mu} = 0$  and  $\bar{\Delta}_\kappa = \infty$ . Due to the infinite value of the curvature disorder at this fixed point one has to consider a special set of coupling constants  $\{\bar{g}\} \equiv \{\bar{\mu}, \bar{g}_\mu, \bar{g}_\lambda, \bar{\Delta}_\mu, \bar{\Delta}_\lambda\}$  involving  $\bar{g}_\mu = \bar{\mu} \bar{\Delta}_\kappa$  and  $\bar{g}_\lambda = \bar{\lambda} \bar{\Delta}_\kappa$  that stay finite at  $P_5$ . At this fixed point one has:  $\bar{\Delta}_\mu = 24\pi^2 \epsilon / d_{c6}$ ,  $\bar{g}_\mu = 48\pi^2 \epsilon / d_{c6}$  and  $\eta = \eta' = 3 \epsilon / d_{c6}$  with  $d_{c6} = d_c + 6$ . At this order  $\phi = 0$ ;  $P_5$  is thus marginal. An analysis of the nonlinearities of the RG flow shows that  $P_5$  is marginally unstable [21, 22].

*Two-loop order.* At two-loop order [31] one recovers the disorder-free fixed point  $P_4$  whose coordinates and anomalous dimension are given in [33]. We focus here on the non-trivial fixed points with non-vanishing disorder. To identify both the vanishing temperature fixed point  $P_5$  and the putative fixed point  $P_c$  obtained by Coquand *et al.* [23] one has again to recourse to the set of coupling constants  $\{\bar{g}\}$  that we expand in powers of  $\epsilon$  up to order 2,  $\bar{g} = \sum_{i=1}^2 \mathcal{C}_{\bar{g}}(i) \epsilon^i$ , where the  $\mathcal{C}_{\bar{g}}^{(1)}$ 's are the coordinates of  $P_5$  at one-loop order, given above. One finds, contrary to [31], that the coordinates of  $P_5$  at two-loop order are completely determined:  $\bar{\mu} = 0$ ,  $\bar{g}_\mu = 48\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$ ,  $\bar{g}_\lambda = -16\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$ ,  $\bar{\Delta}_\mu = 24\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$  and

$\bar{\Delta}_\lambda = -8\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$ . Their full expressions up to order  $\epsilon^3$  as well as the corresponding anomalous dimensions  $\eta_5$  and  $\eta'_5$  are given in Table V of App.C.

Note that, at this order,  $P_5$  no longer belongs to the hypersurfaces  $\bar{\lambda}/\bar{\mu} = \bar{g}_\lambda/\bar{g}_\mu = \bar{\Delta}_\lambda/\bar{\Delta}_\mu = -1/3$  by a distance of  $O(\epsilon)$ . This seems to be a general feature of the perturbative approach of flat phase of both disorder-free and disordered membranes beyond one-loop order. Let us give here the numerical values of the anomalous dimensions in the physical  $d_c = 1$  case :

$$\begin{aligned} \eta_5 &= 0.42857 \epsilon - 0.03779 \epsilon^2 + O(\epsilon^3) \\ \eta'_5 &= 0.42857 \epsilon - 0.03341 \epsilon^2 + O(\epsilon^3) \end{aligned} \quad (2)$$

and  $\phi_5 = 0.00437 \epsilon^2$ . An important fact is that one finds  $P_5$  to be *unstable*, in agreement with the positive value of  $\phi_5$  but in disagreement with the NPRG approach. We shall come back on this fact below.

As one of the main results of our computation at two-loop order is the identification of a finite temperature, finite disorder, “ $P_c$ -like”, fixed point whose coordinates  $\bar{\mu} = -8\pi^2 \epsilon^2 / d_{c6}^2 + O(\epsilon^3)$ ,  $\bar{g}_\mu = 48\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$ ,  $\bar{g}_\lambda = -16\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$ ,  $\bar{\Delta}_\mu = 24\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$  and  $\bar{\Delta}_\lambda = -8\pi^2 \epsilon / d_{c6} + O(\epsilon^2)$  differ, at leading order, with those of  $P_5$ , only by the (non-vanishing) value of  $\bar{\mu}$ . As found within the NPRG approach of Coquand *et al.* [33] this fixed point emerges from  $P_5$  as the dimension  $D$  is lowered from the upper critical dimension  $D = 4$ . The full expressions of the coordinates as well as the corresponding anomalous dimensions are given in the Table VI of App.C. Very surprisingly  $\bar{\mu}$  is found to be *negative* at leading order in  $\epsilon$ , which violates the (mean-field) condition of positivity of  $\bar{\mu}$  required by stability considerations. Moreover, and in agreement with the unstable character of  $P_5$ , the fixed point  $P_c$  is found to be stable, a result also at odds with those obtained within the NPRG approach. All these facts should be nevertheless considered with great care. Indeed, one knows since the two-loop perturbative approach to disorder-free membranes by Coquand *et al.* [33] that some physical fixed points, well defined at one-loop order, are ejected from the (mean-field) region of stability or from the hypersurface  $\bar{\lambda}/\bar{\mu} = -1/3$ , at two-loop order. However the physical quantities – as opposed to fixed points coordinates – are trustable and in agreement with those obtained from nonperturbative (SCSA and NPRG) approaches. One gets, for the anomalous dimensions at  $P_c$  for  $d_c = 1$ :

$$\eta_c = \eta'_c = 0.42857 \epsilon - 0.03695 \epsilon^2 + O(\epsilon^3) \quad (3)$$

and  $\phi_c = 0$  as  $\partial_t \ln \bar{\Delta}_\kappa = \eta' - \eta$  vanishes at this fixed point.

One immediately sees on the series (2) and (3) a strong decrease of their numerical coefficients with the order of the expansion. The same observation was done in the context of disorder-free membranes at two-[33] and three-loop [34] orders. This is both an indicator of the (apparent) convergence of the series and of a fruitful comparison

with the nonperturbative approaches. This is done below with the three-loop order results.

*Three-loop order.* At three-loop order the picture obtained at two-loop order is not fundamentally changed. One gets the disorder-free fixed point  $P_4$  and two non-trivial fixed points: the vanishing temperature, infinite disorder fixed point  $P_5$  and the finite disorder, finite temperature, fixed point  $P_c$  whose coordinates and anomalous dimensions for any  $d_c$  are given in Table V and VI of App.C. Again it is more appealing to consider the anomalous dimensions in the physical  $d_c = 1$  case. One has:

$$\begin{aligned}\eta_5 &= 0.42857 \epsilon - 0.03779 \epsilon^2 - 0.01205 \epsilon^3 + O(\epsilon^4) \\ \eta'_5 &= 0.42857 \epsilon - 0.03341 \epsilon^2 - 0.00964 \epsilon^3 + O(\epsilon^4).\end{aligned}\quad (4)$$

One immediately notice that the coefficients associated to the third order are still strongly decreasing with respect to those of second order. This was also observed in the disorder-free case [34]. The same phenomenon occurs at  $P_c$ :

$$\eta_c = \eta'_c = 0.42857 \epsilon - 0.03695 \epsilon^2 - 0.01191 \epsilon^3 + O(\epsilon^4) \quad (5)$$

*Comparison with NPRG approach.* It is very instructive to compare our results with those obtained from the NPRG approach and reexpanded in powers of  $\epsilon$ . The results for  $\eta_5$ ,  $\eta'_5$  and  $\eta_c$  for any  $d_c$  are given in Table IV of App.B and, again, we consider them for  $d_c = 1$ . For convenience all – perturbative as well as nonperturbative – results have been gathered in Table I and II. One first notice the structural identity between the series coming from the two approaches. Secondly, one see that the numerical agreement is very good both at two- and three-loop orders, almost up to three digits at the fixed point  $P_c$ . This fact advocates for – but does not ensure – the identification of the fixed point  $P_c$  found in this work with that obtained within the nonperturbative context.

Approach	$P_5$
Three-loop	$\eta_5 = 0.42857 \epsilon - 0.03779 \epsilon^2 - 0.01205 \epsilon^3$
	$\eta'_5 = 0.42857 \epsilon - 0.03341 \epsilon^2 - 0.00964 \epsilon^3$
NPRG	$\eta_5 = 0.42857 \epsilon - 0.03532 \epsilon^2 - 0.01293 \epsilon^3$
	$\eta'_5 = 0.42857 \epsilon - 0.03999 \epsilon^2 - 0.01636 \epsilon^3$

TABLE I: Anomalous dimensions  $\eta_5$  and  $\eta'_5$  at order  $\epsilon^3$  at the fixed point  $P_5$  from the three-loop order approach (this work) and from the NPRG approach [23].

Finally, on the basis of the fast decreasing character of the series giving the various anomalous dimensions one can compute, from the expressions given in Table I and II, successive approximations of  $\eta_5$  and  $\eta_c$  in  $D = 2$ , *i.e.* taking  $\epsilon = 2$ , without having recourse to resummation

Approach	$P_c$
Three-loop	$\eta_c = 0.42857 \epsilon - 0.03695 \epsilon^2 - 0.01191 \epsilon^3$
NPRG	$\eta_c = 0.42857 \epsilon - 0.03621 \epsilon^2 - 0.01318 \epsilon^3$

TABLE II: Anomalous dimension  $\eta_c = \eta'_c$  at order  $\epsilon^3$  at the fixed point  $P_c$  from the three-loop order approach (this work) and from the NPRG approach [23].

techniques that are worthless in the case of fast decreasing series. The results, displayed in Table III, show that the exponents  $\eta_5$  and  $\eta_c$  obtained in the perturbative context get closer and closer to the values obtained by means of the NPRG approach. This is only indicative but provide a further sign that the perturbative and non-perturbative techniques may describe the same physical situation.

Approach	$P_5$	$P_c$
Three-loop $\epsilon$	$\eta_5 = 0.85714$	$\eta_c = 0.85714$
Three-loop $\epsilon^2$	$\eta_5 = 0.70600$	$\eta_c = 0.70933$
Three-loop $\epsilon^3$	$\eta_5 = 0.60962$	$\eta_c = 0.61402$
NPRG	$\eta_5 = 0.449$	$\eta_c = 0.492$

TABLE III: Approximations in  $D = 2$  of the critical exponents  $\eta_5$  and  $\eta_c$  at order  $\epsilon$ ,  $\epsilon^2$  and  $\epsilon^3$  from the perturbative approach (this work) and from the NPRG approach [23].

*Conclusion.* We have analyzed quenched disordered membranes by means of a three-loop order perturbative approach. We have derived the RG functions for the various coupling constants and determined the fixed points relevant to the long-distance physics of membranes at two and three-loop orders. Our findings is that it clearly exists a new finite temperature, finite disorder, fixed point  $P_c$  in the RG flow diagram. The proximity between the values for the anomalous dimensions obtained within the perturbative and nonperturbative approaches strongly suggests that, despite different characteristics, the fixed point  $P_c$  identified in the former approach coincides with that discovered within the latter one. This would then be a confirmation of the existence of a finite disorder, finite temperature transition occurring in the phase transition of quenched disordered membranes as well as that of the existence of a low-temperature glassy phase in these systems. Obviously some questions remains open, in particular those of the position and stability of the fixed point found within the perturbative context. Recent studies of the flat phase of disorder-free membranes [33, 34] have clearly established that perturbative computations performed in the vicinity of  $D = 4$ , beyond one-loop order, lead to artefacts on non-universal quantities such as fixed point coordinates. Computations performed at fixed di-

mensions could be of strong help in this context.

### Acknowledgments

We wish to greatly thank O. Coquand and S. Teber for discussions.

- 
- [1] K. S. Novoselov and A. K. Geim and S. V. Morozov and D. Jiang and Y. Zhang and S. V. Dubonos and I. V. Grigorjeva and A. A. Firsov, *Science* **306**, 666 (2004).
- [2] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, *Rev. Mod. Phys.* **81**, 109 (2009).
- [3] M. I. Katsnelson, *Graphene: Carbon in Two Dimensions* (Cambridge University Press, Cambridge, U.K., 2012).
- [4] B. Amorim, A. Cortijo, F. de Juan, A. G. Grushin, F. Guinea, A. Gutiérrez-Rubio, H. Ochoa, V. Parente, R. Roldán, P. San-José, J. Schiefele, M. Sturla, and M. A. H. Vozmediano, *Phys. Rep.* **617**, 1 (2016).
- [5] D. Akinwande, C. J. Brennan, J. Scott Bunch, P. Egberts, J. R. Felts, H. Gao, R. Huang, J.S. Kim, T. Li, Y. Li, K. M. Liechti, N. Lu, H.S. Park, E. J. Reed, P. Wang, B.I. Yakobson T. Zhang, Y.W. Zhang, Y. Zhou and Y. Zhu, *Extreme Mech. Lett.* **13**, 42 (2017).
- [6] G. López-Polín, C. Gómez-Navarro, V. Parente, F. Guinea, M. I. Katsnelson, F. Pérez-Murano, and J. Gómez-Herrero, *Nature Physics* **11**, 26 (2015).
- [7] L. Liu, M. Qing, Y. Wang, and S. Chen, *J. Mater. Sci. Technol.* **31**, 599 (2015).
- [8] G. Yang, L. Li, W. B. Lee and M.C. Ng, *Sci. Technol. Adv. Mater.* **19**, 613 (2018).
- [9] B. Ni, T. Zhang, J. Li, X. Li and H. Gao, *Handbook of Graphene: Physics, Chemistry, and Biology* (John Wiley and Sons, 2019), vol. 2.
- [10] E. Sackman, P. Eggl, C. Fahn, H. Bader, H. Ringdorf and M. Schollmeier, *Ber. Bunsenges. Phys. Chem.* **89**, 1198 (1985).
- [11] D. B. M. Mutz and M. J. Brienne, *Phys. Rev. Lett.* **67**, 923 (1991).
- [12] S. Chaieb, V.K. Natrajan, and A. A. El-rahman, *Phys. Rev. Lett.* **96**, 078101 (2006).
- [13] S. Chaieb, S. Málková, and J. Lal, *J. Theor. Biol.* **251**, 60 (2008).
- [14] S. Chaieb, *Scientific Reports* **4**, 3699 (2013).
- [15] S. Chaieb, *Scientific Reports* **4**, 7347 (2014).
- [16] D. R. Nelson and L. Radzihovsky, *Europhys. Lett.* **16**, 79 (1991).
- [17] L. Radzihovsky and D. R. Nelson, *Phys. Rev. A* **44**, 3525 (1991).
- [18] J. A. Aronovitz and T. C. Lubensky, *Phys. Rev. Lett.* **60**, 2634 (1988).
- [19] L. Radzihovsky and P. Le Doussal, *J. Phys. I France* **2**, 599 (1992).
- [20] P. Le Doussal and L. Radzihovsky, *Ann. Phys. (N.Y.)* **392**, 340 (2018).
- [21] D. C. Morse, T. C. Lubensky and G. S. Grest, *Phys. Rev. A* **45**, R2151 (1992).
- [22] D. C. Morse and T. C. Lubensky, *Phys. Rev. A* **46**, 1751 (1992).
- [23] O. Coquand, K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev E* **97**, 030102(R) (2018).
- [24] J.-P. Kownacki and D. Mouhanna, *Phys. Rev. E* **79**, 040101(R) (2009).
- [25] F. L. Braghin and N. Hasselmann, *Phys. Rev. B* **82**, 035407 (2010).
- [26] K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev. Lett.* **106**, 128102 (2011).
- [27] N. Hasselmann and F. L. Braghin, *Phys. Rev. E* **83**, 031137 (2011).
- [28] K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev. E* **89**, 042101 (2014).
- [29] O. Coquand and D. Mouhanna, *Phys. Rev. E* **94**, 032125 (2016).
- [30] O. Coquand, K. Essafi, J.-P. Kownacki, and D. Mouhanna, *Phys. Rev. E* **101**, 042602 (2020).
- [31] O. Coquand and D. Mouhanna, *Phys. Rev. E* **103**, 031001 (2021).
- [32] A. Mauri and M.I. Katsnelson, *Nucl. Phys. B* **956**, 115040 (2020).
- [33] O. Coquand, D. Mouhanna, and S. Teber, *Phys. Rev. E* **101**, 062104 (2020).
- [34] S. Metayer, D. Mouhanna, and S. Teber, *Phys. Rev. E* **105**, L012603 (2022).
- [35] A. Pikelner (2021), arXiv:2112.07340.
- [36] A. V. Kotikov and S. Teber, *Phys. Part. Nucl.* **50**, 1 (2019), 1805.05109.
- [37] P. Nogueira, *J. Comput. Phys.* **105**, 279 (1993).
- [38] R. N. Lee (2012), arXiv:1212.2685.
- [39] R. N. Lee, *J. Phys. Conf. Ser.* **523**, 012059 (2014).
- [40] S. Metayer, Unpublished (2022).
- [41] This phase should not be confused with the high-temperature, crumpled, phase of membranes.
- [42] The supplementary material file “SM.m”, containing the full 3-loop RG expressions, is available along with the  $\LaTeX$  files.
- [43] One forgets unstable fixed points with vanishing disorder.
-

**Appendix A: Renormalization group equations at two-loop order.**

One gives here the RG equations at two-loop order for the – dimensionless – coupling constants (forgetting their overlining) entering in action Eq.(1). The three-loop order contributions are too long to be displayed on paper; they are given in the supplementary material [42]. For the sake of brevity, we use the notations  $\eta_{ut} = \epsilon - 2\eta_t$ ,  $a = \lambda + 2\mu$ ,  $\Delta_a = \Delta_\lambda + 2\Delta_\mu$ ,  $b_n = 1 + n\Delta_\kappa$ ,  $\Theta_1 = a\mu(a - \mu)$  and  $\Theta_2 = a^2\Delta_\mu + \mu^2(a + \Delta_a) - a\mu(a + 2\Delta_\mu)$ , leading:

$$\begin{aligned}
\partial_t \mu &= -\mu \eta_{ut} + \frac{d_c \mu^2 b_2}{6(4\pi)^2} + \frac{d_c \mu^2}{216a^2(4\pi)^4} \left[ 686\Delta_\kappa^2 \Theta_1 - 227b_4 \Theta_2 \right], \\
\partial_t \lambda &= -\lambda \eta_{ut} + \frac{d_c b_2}{6(4\pi)^2} \left[ 6a^2 - 18a\mu + 13\mu^2 \right] - \frac{d_c}{216a^2(4\pi)^4} \left[ 6d_c b_2^2 a^2 \mu (3a - 5\mu)^2 \right. \\
&\quad \left. - b_4 \Theta_2 (378a^2 - 1674a\mu + 1819\mu^2) + 2\Delta_\kappa^2 \Theta_1 (972a^2 - 3726a\mu + 3641\mu^2) \right], \\
\partial_t \Delta_\mu &= -\Delta_\mu \eta_{ut} + \frac{d_c \mu}{6(4\pi)^2} \left[ 2b_2 \Delta_\mu - \Delta_\kappa^2 \mu \right] + \frac{d_c \mu}{108a^2(4\pi)^4} \left[ 2\Delta_\kappa^2 \Theta_1 (58\mu b_{-2} + 343\Delta_\mu) + \Theta_2 (343\mu \Delta_\kappa^2 - 227b_4 \Delta_\mu) \right], \\
\partial_t \Delta_\lambda &= -\Delta_\lambda \eta_{ut} - \frac{d_c}{6(4\pi)^2} \left[ \Delta_\kappa^2 (6a^2 - 18a\mu + 13\mu^2) - 2b_2 (3a(2\Delta_a - 3\Delta_\mu) - \mu(9\Delta_a - 13\Delta_\mu)) \right] \\
&\quad + \frac{d_c}{108a^2(4\pi)^4} \left[ \Delta_\kappa^2 \left( 6d_c a^2 b_2 \mu (3a - 5\mu)^2 + a^2 \mu^2 (4698\Delta_a - 22101\Delta_\mu + 3493\mu) - 108a^3 \mu (18\Delta_a - 87\Delta_\mu + 19\mu) \right. \right. \\
&\quad \left. \left. - 54a^4 (18\Delta_\mu - 7\mu) + a\mu^3 (14564\Delta_\mu - 1819\mu) - 3641\mu^4 \Delta_a \right) - 9d_c a^2 b_2^2 (3a - 5\mu) (a\Delta_\mu + 2\Delta_a \mu - 5\Delta_\mu \mu) \right. \\
&\quad \left. + 4\Delta_\kappa^3 \Theta_1 (297a^2 - 1026a\mu + 911\mu^2) + b_4 \Theta_2 (378a\Delta_a - 837a\Delta_\mu - 837\Delta_a \mu + 1819\mu\Delta_\mu) \right], \\
\partial_t \Delta_\kappa &= 2\Delta_\kappa \eta_t - \frac{5\Theta_1 b_1 \Delta_\kappa}{a^2(4\pi)^2} + \frac{\Delta_\kappa}{72a^4(4\pi)^2} \left[ a^2 \mu^2 (b_3 + 2\Delta_\kappa^2) \left( a^2 (39d_c + 340) - 1220a\mu + 5(212 - 15d_c)\mu^2 \right) \right. \\
&\quad \left. - 20 \left( a^2 \mu^2 \Delta_\mu (45a\Delta_\kappa + 122\Delta_a + 424\Delta_\mu) - a^3 \mu \Delta_\mu (15a\Delta_\kappa + 244\Delta_\mu) + 34a^4 \Delta_\mu^2 \right. \right. \\
&\quad \left. \left. - a\mu^3 (15a\Delta_\kappa (\Delta_a + 2\Delta_\mu) + 424\Delta_a \Delta_\mu) + \mu^4 \Delta_a (15a\Delta_\kappa + 106\Delta_a) \right) \right], \\
\eta_t &= \frac{5}{a^2(4\pi)^2} \left[ \Delta_\kappa \Theta_1 - \Theta_2 \right] + \frac{1}{72a^4(4\pi)^2} \left[ a^2 \left( d_c (\mu^2 (39a^2 (\Delta_\kappa b_{-3} + 1) + 75\mu^2 (3\Delta_\kappa^2 - b_1))) \right. \right. \\
&\quad \left. \left. + \mu^2 \Delta_\kappa (20(17a^2 - 61a\mu + 53\mu^2) - 90\Delta_\kappa (13a^2 - 44a\mu + 37\mu^2)) + 6(\Delta_\mu - \mu) (d_c a b_2 \mu (13a - 25\mu) \right. \right. \\
&\quad \left. \left. + 30a(10\mu^2 \Delta_\kappa - a(\Delta_\mu - \mu b_{-6})) + 150\Theta_2 \right) + 10a\mu \Theta_2 (244a\Delta_\kappa + 15d_c b_2 \mu - 424\mu \Delta_\kappa) - 1060\Theta_2^2 \right].
\end{aligned}$$

**Appendix B: Anomalous dimensions at  $P_5$  and  $P_c$  at order  $\epsilon^3$  from NPRG approach.**

$\eta_5$	$\frac{3}{d_{c6}} \epsilon - \frac{d_c(4896 + 1734d_c + 155d_c^2)}{80d_{c6}^4} \epsilon^2 + \frac{d_c(5375d_c^5 - 5178d_c^4 - 1711125d_c^3 - 18385218d_c^2 - 75013452d_c - 109355832)}{19200d_{c6}^7} \epsilon^3$
$\eta'_5$	$\frac{3}{d_{c6}} \epsilon - \frac{d_c(5544 + 1962d_c + 175d_c^2)}{80d_{c6}^4} \epsilon^2 + \frac{d_c(6475d_c^5 - 7914d_c^4 - 2132397d_c^3 - 23019066d_c^2 - 94606380d_c - 138990168)}{19200d_{c6}^7} \epsilon^3$
$\eta_c$	$\frac{3}{d_{c6}} \epsilon - \frac{d_c(2556 + 425d_c)}{240d_{c6}^3} \epsilon^2 + \frac{d_c(129925d_c^3 - 894738d_c^2 - 24905043d_c - 89157186)}{518400d_{c6}^5} \epsilon^3$

TABLE IV: Anomalous dimensions  $\eta_5$ ,  $\eta'_5$  and  $\eta_c$  at order  $\epsilon^3$  obtained from the NPRG approach of Coquand *et al.* [23].

Appendix C: Fixed points coordinates at order  $\epsilon^3$  from three-loop approach.

$\mu_5$	0
$g_{\mu_5}$	$\frac{48\pi^2}{d_{c6}} \epsilon - \frac{4\pi^2(52 d_c^2 + 573 d_c + 486)}{5 d_{c6}^4} \epsilon^2 - \frac{\pi^2}{500 d_{c6}^7} \left( 31780 d_c^5 - d_c^4(4130687 - 3558816 \zeta_3) - 6 d_c^3(7627163 - 6225984 \zeta_3) - 12 d_c^2(1161793 - 1236384 \zeta_3) + 72 d_c(13807837 - 10855296 \zeta_3) + 864(2512621 - 2045088 \zeta_3) \right) \epsilon^3$
$g_{\lambda_5}$	$-\frac{1}{3} \frac{48\pi^2}{d_{c6}} \epsilon + \frac{4\pi^2(44 d_c^2 + 511 d_c + 1122)}{5 d_{c6}^4} \epsilon^2 + \frac{\pi^2}{1500 d_{c6}^7} \left( 26980 d_c^5 - d_c^4(3968047 - 3468096 \zeta_3) - 26 d_c^3(1541893 - 1353024 \zeta_3) + 12 d_c^2(3653327 - 396576 \zeta_3) + 72 d_c(17311117 - 11943936 \zeta_3) + 864(2987981 - 2181168 \zeta_3) \right) \epsilon^3$
$\Delta_{\mu_5}$	$\frac{24\pi^2}{d_{c6}} \epsilon - \frac{6\pi^2(14 d_c^2 + 121 d_c - 138)}{5 d_{c6}^4} \epsilon^2 - \frac{\pi^2}{1500 d_{c6}^7} \left( 3195 d_c^5 - 20 d_c^4(259615 - 192456 \zeta_3) - 12 d_c^3(3948749 - 2848932 \zeta_3) + 1440 d_c^2(33334 - 34263 \zeta_3) + 1296 d_c(907519 - 739692 \zeta_3) + 23328(78457 - 67716 \zeta_3) \right) \epsilon^3$
$\Delta_{\lambda_5}$	$-\frac{1}{3} \frac{24\pi^2}{d_{c6}} \epsilon - \frac{2\pi^2(6 d_c^2 + 119 d_c + 858)}{5 d_{c6}^4} \epsilon^2 - \frac{\pi^2}{1500 d_{c6}^7} \left( 10135 d_c^5 + 10 d_c^4(259801 - 174312 \zeta_3) + 4 d_c^3(8149309 - 5609412 \zeta_3) + 240 d_c^2(515581 - 345546 \zeta_3) + 144 d_c(1020203 - 541404 \zeta_3) + 3456(53663 - 20169 \zeta_3) \right) \epsilon^3$
$\eta_5$	$\frac{3}{d_{c6}} \epsilon - \frac{d_c(2802 + d_c(767 + 60 d_c))}{40 d_{c6}^4} \epsilon^2 - \frac{d_c}{24000 d_{c6}^7} \left( 16000 d_c^5 + 5 d_c^4(309539 - 152928 \zeta_3) + 4 d_c^3(1333339 + 261468 \zeta_3) - 480 d_c^2(263197 - 267543 \zeta_3) - 288 d_c(2968601 - 2664738 \zeta_3) - 432(3021431 - 2774088 \zeta_3) \right) \epsilon^3$
$\eta'_5$	$\frac{3}{d_{c6}} \epsilon - \frac{d_c(2442 + d_c(707 + 60 d_c))}{40 d_{c6}^4} \epsilon^2 - \frac{d_c}{12000 d_{c6}^7} \left( 8000 d_c^5 + d_c^4(1377941 - 787968 \zeta_3) + d_c^3(10751771 - 4582008 \zeta_3) - 6 d_c^2(8052667 - 9990216 \zeta_3) - 36 d_c(16694513 - 14815224 \zeta_3) - 216(5856221 - 4970808 \zeta_3) \right) \epsilon^3$
$\phi_5$	$\frac{3 d_c}{2 d_{c6}^3} \epsilon^2 - \frac{d_c}{8000 d_{c6}^6} \left( 3 d_c^3(134243 - 90144 \zeta_3) + 8 d_c^2(371711 - 222588 \zeta_3) - 252 d_c(31513 - 31104 \zeta_3) - 720(94493 - 73224 \zeta_3) \right) \epsilon^3$

TABLE V: Coordinates of the fixed point  $P_5$  and corresponding anomalous dimensions at three-loop order.

$\mu_c$	$-\frac{8\pi^2}{d_{c6}^2} \epsilon^2 + \frac{\pi^2}{1500 d_{c6}^4} \left( 9 d_c^2(44081 - 30048 \zeta_3) + 2 d_c(276557 - 79056 \zeta_3) - 120(92233 - 73224 \zeta_3) \right) \epsilon^3$
$g_{\mu_c}$	$\frac{48\pi^2}{d_{c6}} \epsilon - \frac{4\pi^2(47 d_c + 21)}{5 d_{c6}^3} \epsilon^2 - \frac{\pi^2}{3000 d_{c6}^5} \left( 3 d_c^3(195803 - 90144 \zeta_3) - 8 d_c^2(942497 - 1026756 \zeta_3) - 2196 d_c(12421 - 12528 \zeta_3) + 144(1591741 - 1312848 \zeta_3) \right) \epsilon^3$
$g_{\lambda_c}$	$-\frac{1}{3} \frac{48\pi^2}{d_{c6}} \epsilon + \frac{4\pi^2(127 d_c + 501)}{15 d_{c6}^3} \epsilon^2 + \frac{\pi^2}{9000 d_{c6}^5} \left( 3 d_c^3(186203 - 90144 \zeta_3) - 8 d_c^2(773567 - 958716 \zeta_3) - 684 d_c(9179 - 30672 \zeta_3) + 144(2085601 - 1448928 \zeta_3) \right) \epsilon^3$
$\Delta_{\mu_c}$	$\frac{24\pi^2}{d_{c6}} \epsilon - \frac{2\pi^2(32 d_c - 69)}{5 d_{c6}^3} \epsilon^2 - \frac{\pi^2}{3000 d_{c6}^5} \left( 9 d_c^3(44791 - 30048 \zeta_3) - 16 d_c^2(174209 - 167022 \zeta_3) - 1944 d_c(7109 - 5732 \zeta_3) + 144(706763 - 609444 \zeta_3) \right) \epsilon^3$
$\Delta_{\lambda_c}$	$-\frac{1}{3} \frac{24\pi^2}{d_{c6}} \epsilon - \frac{2\pi^2(28 d_c + 429)}{15 d_{c6}^3} \epsilon^2 + \frac{\pi^2}{3000 d_{c6}^5} \left( d_c^3(111973 - 90144 \zeta_3) - 4 d_c^2(598867 - 452736 \zeta_3) - 8 d_c(2499059 - 1844532 \zeta_3) - 96(108751 - 40338 \zeta_3) \right) \epsilon^3$
$\eta_c = \eta'_c$	$\frac{3}{d_{c6}} \epsilon - \frac{3 d_c(20 d_c + 149)}{40 d_{c6}^3} \epsilon^2 - \frac{d_c}{24000 d_{c6}^5} \left( 16000 d_c^3 + d_c^2(550237 - 223776 \zeta_3) - d_c(5283790 - 5670000 \zeta_3) - (47601132 - 42075936 \zeta_3) \right) \epsilon^3$

TABLE VI: Coordinates of the fixed point  $P_c$  and corresponding anomalous dimensions at three-loop order.