

# CORRESPONDENCE BETWEEN FACTORABILITY AND NORMALISATION IN MONOIDS

ALEN ĐURIĆ

**ABSTRACT.** The purpose of this note is to investigate the relation between two notions concerning monoids: factorability structure, introduced to simplify the bar complex; and quadratic normalisation, introduced to generalise quadratic rewriting systems and normalisations arising from Garside families. We characterise factorable monoids in the axiomatic setting of quadratic normalisation as monoids admitting quadratic normalisation satisfying a condition stronger than the class  $(5, 4)$  yet weaker than the class  $(4, 4)$ . We also prove an equivalence between the two assumptions known to ensure the convergence of the rewriting system associated to factorability.

## CONTENTS

Introduction	2
Factorability structure	2
Quadratic normalisation	2
Contributions	3
Acknowledgments	4
1. Preliminaries	4
1.1. Normal forms and rewriting systems	4
1.2. Divisibility in monoids	5
2. Factorability structure	5
2.1. Notion of factorability structure	5
2.2. Local factorability structure	7
2.3. Deviation from the original definition	8
2.4. Rewriting system associated to factorability	9
3. Quadratic normalisation	10
3.1. Normalisation and normal form	10
3.2. Quadratic normalisation	11
3.3. Quadratic normalisation of class $(4, 3)$	13
3.4. Rewriting system associated to quadratic normalisation	14
3.5. Left-weighted normalisation	15
4. Correspondence between factorability and normalisation	15
4.1. Characterisation of factorable monoids	15

*Date:* June 3, 2022.

*2020 Mathematics Subject Classification.* 20M05, 68Q42.

*Key words and phrases.* monoid, normal form, factorability, normalisation, rewriting system, termination, convergence.



This program has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 754362.

4.2. Factorability in relation to quadratic normalisation of class $(4, 3)$	19
References	23

INTRODUCTION

This text investigates combinatorial properties of a certain class of monoids, seen from two different viewpoints, with a goal of unifying the two. The main result answers the question, explicitly mentioned in [6] and [7], of determining the relation between these two approaches.

**Factorability structure.** The notion of factorability structure on monoids and categories is an extension by Wang [17] and Hess [11] of the definition of factorability structure on groups introduced by Bödiger [3] and Visy [16]. Their motivation was to abstract the structure, discovered in symmetric groups, which ensures the existence of a normal form admitting some remarkable properties. In particular, this normal form allows a reduction of the bar complex to a complex having considerably fewer cells, well-adapted for computing the homology of the algebraic structure in question.

This reduction is achieved using the theory of collapsing schemes, introduced in a topological flavour by Brown [2], elaborated for the algebraic setting by Cohen [5], and rediscovered under the name of discrete Morse theory by Forman [8]. The idea is to establish, for every  $n \geq 0$ , a bijection (called collapsing scheme in [2], and Morse matching in discrete Morse theory) between a class of  $n$ -cells called redundant and a class of  $(n + 1)$ -cells called collapsible in such a way that collapsing of paired cells preserves the homotopy type.

The idea of factorability for a given monoid  $M$  and its generating set  $S$  is to determine a convenient way to split off a generator from an element of the monoid. This is achieved in the notion of factorability structure consisting of a factorisation map  $\eta = (\eta', \bar{\eta}) : M \rightarrow M^2$  subject to several axioms ensuring, in particular, a compatibility with the multiplication in the monoid. In a manner of speaking,  $\eta$  acts on an element  $f$  of  $M$  by splitting off a generator  $\eta'(f)$ .

To every factorability structure, we can associate a confluent rewriting system that is not necessarily terminating. However, termination is obtained under the additional assumption that, for all  $s \in S$  and  $f \in M$ , the following equalities hold:

$$(0.1) \quad \eta'(sf) = \eta'(s \cdot \eta'(f)), \quad \bar{\eta}(sf) = \bar{\eta}(s \cdot \eta'(f)) \cdot \bar{\eta}(f).$$

Basic definitions concerning rewriting systems are recollected in Section 1. We recall basic notions and some results concerning factorability in Section 2.

**Quadratic normalisation.** Assume that a monoid is generated by a set  $S$ . By a normalisation, we mean a syntactic transformation of an arbitrary word over  $S$  to a “canonical” one, called normal. Quadratic normalisations in monoids, introduced by Dehornoy and Guiraud [7] (influenced by Krammer [13]), generalise, under the same axiomatic setting, two well-known classes of normalisations: those arising from quadratic rewriting systems, as studied in [10] for Artin-Tits monoids and in [1] and [4] for plactic monoids; and those arising from Garside families, as investigated in [6], resulting from successive generalisations of the greedy normal form in braid monoids, originating in the work of Garside [9]. Quadratic normalisations admit the

following locality properties: a word is normal if, and only if, its length-two factors are normal; and the procedure of transforming a word into a normal form consists of a finite sequence of rewriting steps, each of which transforms a length-two factor.

The notion of class of a quadratic normalisation is defined in order to measure the complexity of normalising length-three words. The class  $(m, n)$  means that every length-three word admits at most  $m$  rewriting steps starting from the left, and at most  $n$  rewriting steps starting from the right.

The class  $(4, 3)$  is explored in great detail in [7] as it exhibits exceptionally favourable computational properties. In particular, the rewriting system associated to a quadratic normalisation of class  $(4, 3)$  is always convergent.

Quadratic normalisations are recalled in Section 3.

**Contributions.** We establish a correspondence between factorability structure and quadratic normalisation in monoids, in spite of unrelated origins and motivations for these two notions. By a correspondence, we mean maps in both directions between appropriate subclasses, which are inverse to each other, up to technicalities, and moreover this bijection is compatible with associated rewriting systems.

Since the rewriting system associated to a factorability structure is not necessarily terminating, whereas the rewriting system associated to a quadratic normalisation of class  $(4, 3)$  is always terminating, it is known that a quadratic normalisation corresponding to a factorability structure is not necessarily of class  $(4, 3)$ .

We show that a quadratic normalisation corresponding to a factorability structure is always of class  $(5, 4)$ , and not smaller in general. We give a necessary and sufficient condition for a quadratic normalisation of class  $(5, 4)$  to correspond to a factorability structure. Thereby, we characterise factorable monoids in terms of quadratic normalisation, thus adding another important family of monoids to those unified under the axiomatic framework of quadratic normalisation.

In particular, a quadratic normalisation of class  $(4, 3)$  always yields a factorability structure, but not vice versa. However, the converse does hold under the stronger condition described as follows. Consider the map  $F := \eta \circ \mu$  from the set  $M^2$  to itself, with  $\mu : M^2 \rightarrow M$  denoting the multiplication in  $M$ . Denote by  $F_1$  (resp.  $F_2$ ) application of  $F$  to the first two elements (resp. the second and the third element) of a triple in  $M^3$ . In general, the factorability structure ensures the equality

$$F_1 F_2 F_1 F_2 (r, s, t) = F_2 F_1 F_2 (r, s, t)$$

whenever  $F_2 F_1 F_2 (r, s, t)$  contains no 1. The stronger condition is that this equality holds for every  $(r, s, t) \in S^3$ .

Furthermore, we show that this stronger condition is equivalent to the aforementioned additional assumption (0.1) which is known to grant termination of a rewriting system associated to factorability structure.

One of the benefits of the established correspondence between factorability structure and quadratic normalisation is that it provides a way of importing the results concerning homology, derived from the former (see e.g. [16], [17], [12]) to the framework of the latter, with the hope of generalising them to higher classes. This is going to be our next step in the current direction of research.

To simplify the presentation, we are considering exclusively monoids, but results stated here (recalled ones as well as new ones) mostly extend to categories, seen as monoids with partial multiplication. As a reminder that a monoid is thought of as

a monoid of endomorphisms (of an object), we tend to use letters  $f$ ,  $g$  and  $h$  for elements of a monoid.

**Acknowledgments.** I warmly thank Viktoriya Ozornova for supervising this work, for careful proofreading and numerous remarks which highly improved the quality of the present text.

## 1. PRELIMINARIES

**1.1. Normal forms and rewriting systems.** If  $S$  is a set,  $S^*$  denotes the free monoid over  $S$ . Elements of  $S$  and  $S^*$  are called  $S$ -letters and  $S$ -words, respectively. So, an  $S$ -word is a tuple<sup>1</sup> of  $S$ -letters, e.g.  $(s_1, \dots, s_p)$ . The prefix  $S$ - is left out if it is clear from the context. The product in  $S^*$  of two words  $u$  and  $v$  is denoted by  $u|v$ . A letter  $s$  is customarily identified with the single-letter word  $(s)$ . Accordingly, a word  $(s_1, \dots, s_p)$  can be written as the product (in  $S^*$ ) of its letters:  $s_1|\dots|s_p$ .

A monoid  $M$  is said to be generated by a set  $S$ , often written as  $(M, S)$ , if  $M$  is a homomorphic image of the free monoid  $S^*$ . Such a homomorphism is called an evaluation map and denoted  $\text{ev} : S^* \rightarrow M$ . A **normal form** for  $M$  with respect to  $S$  is a set-theoretic section, denoted by  $\text{nf}$ , of the evaluation map. To rephrase it, a normal form maps elements of  $M$  to their distinguished representative words of  $S^*$ .

We write  $|w|$  for the length of  $w$  in  $S^*$ . A normal form  $\text{nf}$  for a monoid  $M$  with respect to a generating set  $S$  is called **geodesic** if, for every  $f$  in  $M$ , the inequality  $|\text{nf}(f)| \leq |w|$  holds for every  $S$ -word  $w$  representing  $f$ , i.e. such that  $\text{ev}(w) = f$ .

A (word) **rewriting system** is a pair  $(S, R)$  consisting of a set  $S$  and a binary relation  $R$  on  $S^*$ , whose elements  $(u, v)$  are called **rewriting rules**, and also written  $u \rightarrow v$  to stress the fact that they are directed. Seeing relations between words, not as equalities but as rewriting rules, is a key concept of rewriting theory.

For a rewriting rule  $(u, v)$ , and for  $w$  and  $w'$  in  $S^*$ , a pair  $(w|u|w', w|v|w')$  is called a **rewriting step**. For  $u$  and  $v$  in  $S^*$ , we say that  $u$  **rewrites** to  $v$  if there is a finite composable sequence of rewriting steps with source  $u$  and target  $v$ . A word  $u$  is called **irreducible** with respect to  $R$  if there is no rewriting step whose source is  $u$ ; otherwise,  $u$  is reducible.

A rewriting system  $(S, R)$  is called:

- **confluent** if any two rewriting sequences starting with the same word can be completed in such a way that they eventually reach a common result;
- **normalising** if every  $u$  in  $S^*$  rewrites to at least one irreducible word;
- **terminating** if it admits no infinite rewriting sequence;
- **convergent** if it is both confluent and terminating;
- **reduced** (or minimal) if for every rewriting rule  $u \rightarrow v$ , the word  $v$  is irreducible with respect to  $R$ , and the word  $u$  is irreducible with respect to  $R \setminus \{(u, v)\}$ ;
- **strongly reduced** if it is reduced and, in addition, every element of  $S$  is irreducible;
- **quadratic** if the source and target of each element of  $R$  are of length two.

The monoid **presented by** a rewriting system  $(S, R)$  is the quotient  $M$  of the free monoid  $S^*$  by the congruence relation generated by  $R$ . If  $(S, R)$  is confluent, the irreducible word to which a word  $u$  rewrites, if it exists, is denoted by  $\hat{u}$ . If  $(S, R)$

<sup>1</sup>In the broader context of categories, words are generalised to paths of composable morphisms.

is convergent, the map  $M \rightarrow S^*$  defined by  $g \mapsto \widehat{u}$ , with  $\text{ev}(u) = g$ , is the normal form arising from the rewriting system  $(S, R)$ .

*Remark 1.1.1.* If  $S$  is a subset of  $M$ , then elements of  $S$  can be regarded in two ways: as length-one words in  $S^*$ , and as elements of  $M$ . When a rewriting system presenting  $M$  with respect to  $S$  is strongly reduced, this makes no essential difference, so elements of  $S$  are denoted in the same way, regardless of the viewpoint, relying on the context to provide the proper interpretation. In particular, one can say that  $S$  contains (or that it does not contain)  $1$ , the identity element of  $M$ . For clarity, we will emphasise such situation by calling  $S$  a generating subset (i.e. not just a generating set) of  $M$ . This is the case in the context of factorability (see [12], [11], [15]), but not in the context of normalisation in general (see [7]). However, when we characterise factorable monoids in terms of quadratic normalisation (Subsection 4.1), the corresponding normalisations can share this custom.

We adopt the following convention, for technical reasons.

**Convention 1.1.2.** *Whenever  $S$  denotes a generating set (resp. subset) of a monoid  $M$  other than  $S^*$ , it is implicitly understood that  $S$  contains exactly one letter, denoted by  $e$  (resp.  $1$ ), that is evaluated to the identity element of  $M$ . In case we want to exclude this letter from  $S$ , we write  $S_+$  instead of  $S$ .*

**1.2. Divisibility in monoids.** A monoid  $M$  is said to be **left-cancellative** (resp. **right-cancellative**) if, for every  $f, g$  and  $g'$  in  $M$ , the equality  $fg = fg'$  (resp.  $gf = g'f$ ) implies the equality  $g = g'$ .

An element  $f$  of a monoid  $M$  is called a **left divisor** of  $g \in M$ , and  $g$  is called a **right multiple** of  $f$ , denoted by  $f \preceq g$ , if there is an element  $f' \in M$  such that  $ff' = g$ . If  $M$  is left-cancellative, then the element  $f'$  is uniquely defined and called **the right complement of  $f$  in  $g$** .

## 2. FACTORABILITY STRUCTURE

This section recalls the notion of factorability structure. For elaboration, we refer the reader to [12]. Subsection 2.1 recollects the basic terminology. In Subsection 2.2, we recall an alternative approach to factorability through the notion of local factorability. We redefine certain notions in order to overcome the issues arising from the original definition, which we point out in Subsection 2.3. Subsection 2.4 reports on the rewriting system associated to a factorability structure.

**2.1. Notion of factorability structure.** Let  $M$  be a monoid having a generating subset  $S$ . For an element  $f$  of  $M$ , we write  $|f|$  for the minimum  $S$ -length of an  $S$ -word representing  $f$ . A pair  $(f, g)$  in  $M^2$  is called **geodesic** if  $|fg| = |f| + |g|$ .

**Convention 2.1.1.** *We adopt the convention that elements of a tuple are indexed starting from the leftmost one, as in  $(x_1, \dots, x_n)$ , thereby not following the convention used in [12] where elements are indexed starting from the rightmost one. The purpose is to make the notions which concern factorability more easily comparable (in Section 4) with those which concern normalisation.*

A **factorisation map** is a map  $\eta = (\eta', \overline{\eta}) : M \rightarrow M^2$  satisfying the following conditions:

- for  $f \in M \setminus \{1\}$ , the element  $\eta'(f) \in S_+$  is a left divisor of  $f$ , and the element  $\overline{\eta}(f)$  is a right complement of  $\eta'(f)$  in  $f$ ;

- $(\eta'(f), \bar{\eta}(f))$  is geodesic;
- $\eta$  maps  $1 \in M$  to  $\eta(1) = (1, 1)$ .

Whenever confusion is unlikely,  $\eta'(f)$  and  $\bar{\eta}(f)$  are abbreviated to  $f'$  and  $\bar{f}$ , respectively.

**Example 2.1.2.** Assume that  $M$  is a free commutative monoid over a non-empty finite totally ordered set  $A$ . Define  $\eta = (\eta', \bar{\eta}) : M \rightarrow M^2$  by setting  $\eta'(f)$  to be the least left divisor of  $f$ , lying in  $A$ . If there is no such divisor, then set  $\eta'(f) = 1$ . Note that this is well-defined since the left cancellation property of  $M$  implies uniqueness of right complements.

**Notation 2.1.3.** Assume that  $A$  is a set, and  $F$  is a map from  $A^k$  to  $A^l$ . Let  $A^*$  denote the set of all  $A$ -words, i.e. tuples of elements of  $A$ . Then the (partial) map  $F_i : A^* \rightarrow A^*$  consists of applying  $F$  to  $k$  consecutive elements starting from position  $i$ , i.e. to the elements at positions  $i, i+1, \dots, i+k-1$ .

**Example 2.1.4.** For the sake of illustration, take  $(A, <)$  to be a totally ordered set wherein  $A = \{a, b, c, d\}$ , and  $<$  is the alphabetical order  $a < b < c < d$ . We write  $<^*$  for the lexicographic extension of  $<$  to  $A^*$ . Let  $F : A^2 \rightarrow A^2$  map every length-two word  $w$  to the  $<^*$ -minimal word obtained by simply permuting letters of  $w$  if needed. Then, we have:

$$c|b|a \xrightarrow{F_2} c|a|b \xrightarrow{F_1} a|c|b \xrightarrow{F_3} a|b|c.$$

The multiplication in  $M$  is denoted by  $\mu : M^2 \rightarrow M$ , and  $\mu(f, g)$  is often abbreviated to  $f \cdot g$  or  $fg$ .

**Definition 2.1.5** ([12, Definition 2.1]). Assume that  $M$  is a monoid having a generating subset  $S$ . A **factorability structure** on  $(M, S)$  is a factorisation map  $\eta : M \rightarrow M^2$  such that, denoting the map  $\eta \circ \mu : M^2 \rightarrow M^2$  by  $F$ , for every triple in  $M^3$ , the three maps

$$F_1 F_2 F_1 F_2, \quad F_2 F_1 F_2, \quad F_2 F_1 F_2 F_1$$

coincide or each map reduces the sum of the lengths of the elements of the triple. If  $\eta : M \rightarrow M^2$  is a factorability structure on  $(M, S)$ , we call the triple  $(M, S, \eta)$  a **factorable monoid**.

Assume that  $(M, S, \eta)$  is a factorable monoid. A tuple  $x$  in  $M^n$  is called **stable** at the  $i$ th position if  $F_i(x) = x$ . A tuple is **everywhere stable** if it is stable at the  $i$ th position for every  $i$  in  $\{1, \dots, n-1\}$ .

To a factorable monoid  $(M, S, \eta)$ , one assigns the map  $\text{nf}_\eta : M \rightarrow S_+^*$  defined as

$$f \mapsto \eta_{|f|-1} \cdots \eta_2 \eta_1(f).$$

**Example 2.1.6.** The monoid  $(M, S)$  in Example 2.1.4 is factorable with a factorability structure splitting off the least letter. For  $f = bacabc$ , we get

$$\text{nf}_\eta(f) = \eta_5 \cdots \eta_2 \eta_1(f) = (a, a, b, b, c, c).$$

It turns out ([12, Section 3]) that  $\text{nf}_\eta$  is a normal form, called the **normal form with respect to the factorability structure**  $\eta$ , or the  $\eta$ -normal form, for short. It admits the following locality property.

**Lemma 2.1.7** ([12, Lemma 3.2]). *Let  $(M, S, \eta)$  be a factorable monoid. Then for every  $f \in M$ , the  $\eta$ -normal form is everywhere stable.*

**2.2. Local factorability structure.** There is an alternative definition of factorability, due to Moritz Rodenhausen. In order to resolve an issue detected in the original definition (to be addressed in the next subsection), we introduce the following notation.

**Notation 2.2.1.** Let  $A$  be a set. For a map  $\varphi : A^2 \rightarrow A^2$ , we define the composite map  $\varphi \xrightarrow{\circ} : A^* \rightarrow A^*$  as

$$w \mapsto \varphi_{|w|-1} \cdots \varphi_1(w).$$

Note that the last  $\varphi$  in the above composition is applied to the rightmost length-two factor of the argument  $w$ . In particular, if  $w$  has length one, then  $\varphi \xrightarrow{\circ}(w) = w$ .

**Definition 2.2.2.** Assume that  $A$  is a set containing 1, and  $\varphi : A^2 \rightarrow A^2$  is a map. The **normalisation map associated to the map  $\varphi$**  is the map  $N_\varphi : A^* \rightarrow A^*$  defined as follows:

- (1)  $N_\varphi(()) = ()$ ;
- (2)  $N_\varphi$  of a word containing 1 equals  $N_\varphi$  of the same word with 1 removed;
- (3) and, denoting  $\varphi \xrightarrow{\circ}(s_1 | N_\varphi(s_2, \dots, s_n))$  by  $u$ ,

$$N_\varphi(s_1, \dots, s_n) := \begin{cases} u, & \text{if } u \text{ contains no } 1, \\ N_\varphi(u), & \text{otherwise.} \end{cases}$$

*Remark 2.2.3.* Note that, by recursion, computation of  $N_\varphi(s_1, \dots, s_n)$  terminates and its length is bounded by  $n$ . Namely, all the recursive calls for  $\varphi$  are made on words of smaller length: directly in (2) and in the first case of (3); and indirectly in the second case of (3) which calls for (2). The length of  $N_\varphi(s_1, \dots, s_n)$  is less than or equal to the length of  $\varphi \xrightarrow{\circ}(s_1 | N_\varphi(s_2, \dots, s_n))$ , which is less than or equal to the length of  $N_\varphi(s_2, \dots, s_n)$  plus 1.

We state anew the definition of local factorability structure on  $(M, S)$ , using Definition 2.2.2.

**Definition 2.2.4.** Let  $M$  be a monoid generated by its subset  $S$ . A **local factorability structure** is a map  $\varphi : S^2 \rightarrow S^2$  satisfying the following properties:

- (1)  $M$  admits the presentation

$$\langle S_+ \mid \{(s, t) = \varphi(s, t) \mid s, t \in S_+\} \rangle;$$

- (2)  $\varphi$  is idempotent;
- (3)  $\varphi(1, s) = (s, 1)$  for every  $s \in S_+$ ;
- (4) for every  $(r, s, t) \in S_+^3$ , the equality

$$\varphi_1 \varphi_2 \varphi_1 \varphi_2(r, s, t) = \varphi_2 \varphi_1 \varphi_2(r, s, t)$$

- holds or  $\varphi_2 \varphi_1 \varphi_2(r, s, t)$  contains 1;
- (5) the normalisation map associated to  $\varphi$  satisfies  $N_\varphi(r, s, t) = N_\varphi(\varphi_1(r, s, t))$  for all  $(r, s, t) \in S_+^3$ .

By definition,  $N_\varphi(w)$  contains no 1 for any  $w$  in  $A^*$ . If we add to  $N_\varphi(w)$  a string of 1's on the right, the result is called an **extended form** of  $N_\varphi(w)$ . We recall a result to be used in Section 4.

**Lemma 2.2.5** ([15, Lemma 2.3.5]). *Let  $A$  be a set containing 1, and  $\varphi : A^2 \rightarrow A^2$  a map formally satisfying the second, the third and the fourth condition of Definition 2.2.4, having  $S$  replaced with  $A$ . Furthermore, let  $N_\varphi : A^* \rightarrow A^*$  be the normalisation map associated to the map  $\varphi$ . Then  $\varphi_1\varphi_2\varphi_1\varphi_2(r, s, t)$  is an extended form of  $N_\varphi(r, s, t)$  for all  $(r, s, t)$  in  $A^3$ .*

A factorability structure is equivalent to a local factorability structure, in the following sense.

**Theorem 2.2.6** ([12, Theorem 3.4]).

- (1) *If  $(M, S, \eta)$  is a factorable monoid, then the restriction of the map  $\eta\mu$  to  $S^2$  defines a local factorability structure on  $M$ .*
- (2) *Conversely, one can construct a factorability structure out of a local factorability structure by setting:  $\eta(1) = (1, 1)$ ; and*

$$\eta(f) = (s_1, \text{ev}(s_2, \dots, s_n)) \text{ for } N_\varphi(w) = (s_1, s_2, \dots, s_n),$$
*with  $w$  being any  $S$ -word representing  $f$ .*
- (3) *These constructions are inverse to each other.*
- (4) *By this correspondence, for  $f$  in  $M$ , the  $\eta$ -normal form  $\text{nf}_\eta(f)$  equals  $N_\varphi(w)$ , where  $w$  is any element of  $S^*$  representing  $f$ .*

The proof of Theorem 2.2.6 can be found in [15, Section 2.3].

*Remark 2.2.7.* Here are some observations about local factorability structures that will be used implicitly from now on.

- The property (3) implies  $N_\varphi(s) = s$  for every  $s$  in  $S_+$ .
- For every  $(s, t) \in S^2$ , the first element of  $\varphi(s, t)$  cannot be equal to 1 unless the second element is equal to 1. Namely, assume the opposite:  $\varphi(s, t) = (1, t')$  for  $t' \neq 1$ . Then the idempotency of  $\varphi$  gives  $\varphi(1, t') = (1, t')$  which contradicts the property (3).
- Note that, by Theorem 2.2.6, the equality  $\varphi(s, t) = (1, 1)$  holds if, and only if,  $st = 1$  in  $M$ .

**2.3. Deviation from the original definition.** As hinted by Convention 2.1.1, we have reformulated the original definition of a factorisation map which separates a right divisor, to make it separate a left divisor, instead. We have also modified the definition of local factorability structure.

Here we are going to recall the original definition of a local factorability structure, and explain the motivation for our modification thereof. For simplicity, we still assume Convention 1.1.2, so we do not actually copy the original verbatim, but we do preserve its essence (as well as the convention of indexing from the right).

**Definition** ([12, Definition 3.3]). Let  $M$  be a monoid having a generating subset  $S$ . A **local factorability structure** on  $(M, S)$  is a map  $\varphi : S^2 \rightarrow S^2$  satisfying the following properties:

- (1)  $M$  admits the presentation

$$\langle S_+ \mid \{(t, s) = \varphi(t, s) \mid s, t \in S_+\} \rangle;$$

- (2)  $\varphi$  is idempotent;
- (3)  $\varphi(s, 1) = (1, s)$  for every  $s \in S_+$ ;
- (4) for every  $(t, s, r) \in S_+^3$ , applying any  $\varphi_i$  to the triple  $\varphi_2\varphi_1\varphi_2(t, s, r)$  leaves it unchanged, or  $\varphi_2\varphi_1\varphi_2(t, s, r)$  contains 1;

(5)  $\text{NF}(t, s, r) = \text{NF}(\varphi_1(t, s, r))$  for all  $(t, s, r) \in S_+^3$ .

Here, the map  $\text{NF} : S^* \rightarrow S^*$  is defined inductively on the length of the word as follows:

- $\text{NF}(\epsilon) = \epsilon$ ;
- $\text{NF}(s) = s$  for all  $s \in S_+$ ;
- $\text{NF}$  of a word containing 1 equals  $\text{NF}$  of the same word with 1 removed;
- and

$$\text{NF}(s_n, \dots, s_1) := \begin{cases} \varphi_{n-1} \cdots \varphi_1(\text{NF}(s_n, \dots, s_2), s_1), & \text{if it contains no 1;} \\ \text{NF}(\varphi_{n-1} \cdots \varphi_1(\text{NF}(s_n, \dots, s_2), s_1)), & \text{otherwise.} \end{cases}$$

*Remark 2.3.1.* Observe that there is a flaw in the above definition. Since  $\varphi$  is defined on the domain  $S^2$ , it is not clear what should one do when one gets the expression of the form  $\varphi(s)$ , where  $s$  is a single element of  $S_+$ . However, such an expression may indeed occur in the definition of  $\text{NF}$ , namely, if  $\varphi$  is applied just after  $\text{NF}$  has eliminated 1 and thus shortened the word in question. For example, take  $s_2 = 1$ . Then

$$\text{NF}(s_2, s_1) = \varphi_1(\text{NF}(s_2), s_1) = \varphi_1(s_1)$$

which is not defined. We have resolved this issue by introducing Notation 2.2.1.

*Remark 2.3.2.* Note that in [12] and [15], one calls  $\text{NF}(w) : S^* \rightarrow S^*$  the normal form of  $w$ . In the present text (following [7]), however, normal forms  $\text{nf} : M \rightarrow S^*$  have elements of  $M$ , not of  $S^*$ , as their arguments (recall Subsection 1.1). Accordingly, we do not call the map  $N_\varphi$ , which is our analogue of  $\text{NF}$ , a normal form. Instead, we have called such a syntactic transformation (of arbitrary words to normal ones) a normalisation map, in Definition 2.2.2.

**2.4. Rewriting system associated to factorability.** A rewriting system is associated to a factorable monoid in a canonical way, as follows.

**Lemma 2.4.1** ([12, Lemma 5.1]). *Assume that  $(M, S, \eta)$  is a factorable monoid. Let  $R$  contain the rewriting rules of the form*

$$(s, t) \rightarrow \eta\mu(s, t)$$

*for all  $s$  and  $t$  in  $S_+$  such that  $(s, t)$  is not stable. Then  $(S_+, R)$  is a confluent, strongly reduced rewriting system presenting  $M$ . Here, if  $\bar{\eta}(st) = 1$ , we interpret the rewriting rule as  $(s, t) \rightarrow st$ .*

*Remark 2.4.2.* Observe that, strictly speaking, the presentation obtained by turning rewriting rules into equations in Lemma 2.4.1 is not the same as the one given by the property (1) of Definition 2.2.4. Namely, if  $st$  is in  $S_+$  (resp. equal to 1), then the former contains the relation  $(s, t) = st$  (resp.  $(s, t) = 1$ ), whereas the latter has  $(s, t) = (st, 1)$  (resp.  $(s, t) = (1, 1)$ ). However, one can obtain the former from the latter simply by removing extra 1's, and vice versa.

The associated rewriting system in Lemma 2.4.1 is not necessarily terminating, even if  $S$  is finite and  $M$  is left-cancellative. We refer the reader to [12, Appendix] for an example of a factorable monoid such that its associated rewriting system is not terminating. The following result gives sufficient conditions for the rewriting system associated to a factorable monoid to be terminating.

**Theorem 2.4.3** ([12, Theorem 7.3]). *Assume that  $(M, S, \eta)$  is a factorable monoid meeting the additional assumption that the equalities*

$$\eta'(sf) = \eta'(s \cdot \eta'(f)), \quad \bar{\eta}(sf) = \bar{\eta}(s \cdot \eta'(f)) \cdot \bar{\eta}(f)$$

*hold for all  $s \in S_+$  and  $f \in M$ . Then the associated rewriting system is terminating.*

### 3. QUADRATIC NORMALISATION

This section briefly recalls quadratic normalisation. After presenting basic notions concerning normalisation in Subsection 3.1, we recollect the notion of quadratic normalisation in Subsection 3.2, then we focus on a particular class of quadratic normalisation in Subsection 3.3. Subsection 3.4 reports on the rewriting system associated to a quadratic normalisation. Subsection 3.5 recalls the notion of left-weighted normalisation. For technical elaboration, see [7].

**3.1. Normalisation and normal form.** Having already hinted in the Introduction that a normalisation is a syntactic transformation of an arbitrary word to a normal one, here we recall a formal definition.

**Definition 3.1.1.** A **normalisation** is a pair  $(A, N)$ , where  $A$  is a set and  $N : A^* \rightarrow A^*$  is

- (1) a length-preserving map,
- (2) whose restriction to  $A$  is the identity map,
- (3) and which satisfies  $N(u|v|w) = N(u|N(v)|w)$  for all  $A$ -words  $u, v, w$ .

The map  $N$  is called a **normalisation map**. A word  $w$  such that  $N(w) = w$  is called  **$N$ -normal**. A **normalisation for a monoid  $M$**  is a normalisation  $(S_+, N)$  such that  $M$  admits the presentation

$$\langle S_+ | \{w = N(w) | w \in S_+^*\} \rangle.$$

*Remark 3.1.2.* We use a letter  $A$  in Definition 3.1.1 to point out that normalisation itself is a purely syntactic notion, as opposed to the notion of normalisation for a monoid. Recall that the subscript of  $S_+$  (or absence thereof) is explained by Convention 1.1.2.

**Example 3.1.3** ([7, Example 2.2]). Assume that  $(S_+, <)$  is a linearly ordered set. Let  $<^*$  denote the lexicographic extension of  $<$  to  $S_+^*$ . For an  $S_+$ -word  $w$ , define  $N(w)$  to be the  $<^*$ -minimal word obtained by permuting letters in  $w$ . Then  $(S_+, N)$  is a normalisation for the free commutative monoid over  $S_+$ .

Note that the definition of normalisation for a monoid  $M$  implies that there is a nontrivial homomorphism (or grading)  $d : M \rightarrow (\mathbb{N}, 1)$  such that  $d(s) = 1$  for all  $s$  in  $S_+$ . Namely, take  $d(f)$  to be the common length of all the  $S_+$ -words representing  $f$  (well-defined by the property (1) of normalisation). Such monoids  $(M, S_+)$  are called **graded**. In other words,  $M$  is graded with respect to a generating set  $S_+$  if all  $S_+$ -words representing the same element of  $M$  are equal in length.

*Remark 3.1.4* ([7, Proposition 2.6]). Observe that, in a graded monoid, a normalisation and a normal form are just different aspects of looking at the same notion. Indeed, given a normalisation  $(S_+, N)$  for a monoid  $M$ , a normal form  $\text{nf}$  for  $M$  with respect to  $S_+$  is obtained by setting  $\text{nf}(f) = N(w)$ , where  $w$  is any  $S_+$ -word representing  $f$ . Conversely, if  $\text{nf}$  is a normal form for a graded monoid  $M$  with

respect to a generating set  $S_+$ , one obtains a corresponding normalisation map by setting  $N(w) = \text{nf}(\text{ev}(w))$ . These two correspondences are inverse to each other.

If a monoid  $(M, S_+)$  is not graded, i.e. if an element of  $M$  may be represented by  $S_+$ -words of different lengths, then a new letter representing 1 can be introduced to formally preserve length. For a normalisation  $(S, N)$ , we say that an element  $e$  of  $S$  is  **$N$ -neutral** if the equalities

$$(3.1) \quad N(w|e) = N(e|w) = N(w)|e$$

hold for every  $S$ -word  $w$ . We say that  $(S, N)$  is a **normalisation mod  $e$  for a monoid  $M$**  if  $e$  is an  $N$ -neutral element of  $S$  and  $M$  admits the presentation

$$(3.2) \quad \langle S \mid \{w = N(w) \mid w \in S^*\} \cup \{e = 1\} \rangle.$$

Note that there can be at most one  $N$ -neutral element (hence the restriction to exactly one element representing 1 in Convention 1.1.2). We write  $\pi_e$  for the canonical projection from  $S^*$  onto  $S_+^*$ , which removes all the occurrences of  $e$ . This extends the equivalence between normalisation and normal form to general monoids (i.e. not necessarily graded), as follows.

**Proposition 3.1.5** ([7, Proposition 2.9]).

- (1) If  $(S, N)$  is a normalisation mod  $e$  for a monoid  $M$ , a geodesic normal form  $\text{nf}$  for  $M$  with respect to  $S$  is obtained by  $\text{nf}(f) = \pi_e(N(w))$ , where  $w$  is any  $S$ -word representing  $f$ .
- (2) Conversely, let  $\text{nf}$  be a geodesic normal form for a monoid  $M$  with respect to a generating set  $S_+$ . Write  $\text{ev}^e$  for the extension of the evaluation map  $\text{ev} : S_+^* \rightarrow M$  to  $S^*$  by putting  $\text{ev}^e(e) = 1$ . Then a normalisation  $(S, N)$  mod  $e$  for  $M$  is provided by the map

$$N(w) = \text{nf}(\text{ev}^e(w))|e^m,$$

with  $m$  denoting the number of letters  $e$  to be added in order to formally preserve length, namely  $m = |w| - |\text{nf}(\text{ev}^e(w))|$ .

- (3) These two correspondences are inverse to each other.

*Remark 3.1.6* ([7, Remark 2.10]). If a monoid  $(M, S_+)$  is graded and  $\text{nf}$  is a normal form on  $(M, S_+)$ , then there are two normalisations arising from  $\text{nf}$ , provided by Remark 3.1.4 and Proposition 3.1.5(2), respectively. To make a clear distinction, let us temporarily write  $N_+$  for the normalisation map of the former. The relation between normalisations  $(S_+, N_+)$  and  $(S, N)$  is given by the equality  $\pi_e(N(w)) = N_+(\pi_e(w))$  for all  $w$  in  $S^*$ . Therefore, it suffices to formally consider normalisations mod  $e$  for monoids.

**3.2. Quadratic normalisation.** We write  $\overline{N}$  for the restriction of  $N$  to the set of  $S$ -words of length-two and remind the reader of Notation 2.1.3 which we now extend.

**Notation.** For  $u = (i_1, \dots, i_n)$  in  $\mathbb{N}^n$ , the composite map  $F_{i_n} \circ \dots \circ F_{i_1}$  (note that  $F_{i_1}$  is applied first) is denoted by  $F_u$ , often omitting commas in  $u$  when  $i_k$  is a single-digit number for every  $k \in \{1, \dots, n\}$ .

**Definition 3.2.1.** A normalisation  $(A, N)$  is **quadratic** if the following two requirements are met.

- (1) An  $A$ -word  $w$  is  $N$ -normal if, and only if, every length-two factor of  $w$  is  $N$ -normal.
- (2) For every  $A$ -word  $w$ , there exists a finite sequence of positions  $u$ , depending on  $w$ , such that  $N(w) = \overline{N}_u(w)$ .

**Example 3.2.2.** Normalisation given in Example 3.1.3 is quadratic. Indeed, a word is  $<^*$ -minimal if, and only if, its every length-two factor is  $<^*$ -minimal; and each word can be ordered by switching pairs of adjacent letters that are not ordered as expected.

An advantage of quadratic normalisation is that it is completely determined by the restriction  $\overline{N}$ .

**Proposition 3.2.3** ([7, Proposition 3.6]).

- (1) If  $(S_+, N)$  is a quadratic normalisation for a monoid  $M$ , then  $\overline{N}$  is idempotent and  $M$  admits the presentation

$$\langle S_+ \mid \{s|t = \overline{N}(s|t) \mid s, t \in S_+\} \rangle.$$

- (2) If  $(S, N)$  is a quadratic normalisation, then an element  $e$  of  $S$  is  $N$ -neutral if, and only if, the following condition holds for every  $s$  in  $S$ :

$$\overline{N}(s|e) = \overline{N}(e|s) = \overline{N}(s) | e.$$

- (3) If  $(S, N)$  is a quadratic normalisation mod  $e$  for a monoid  $M$ , then  $\overline{N}$  is idempotent and  $M$  admits the presentation

$$\langle S_+ \mid \{s|t = \pi_e(\overline{N}(s|t)) \mid s, t \in S_+\} \rangle.$$

If  $(A, N)$  is a quadratic normalisation and  $w$  is an  $A$ -word, then  $N(w)$  is computed by sequentially applying  $\overline{N}$  to length-two factors at various positions. For length-three words, there are only two such positions. Since  $\overline{N}$  is idempotent, it suffices to consider alternating sequences of positions. For  $m \geq 0$ , we write  $12[m]$  (resp.  $21[m]$ ) for the alternating sequence  $121\dots$  (resp.  $212\dots$ ) of length  $m$ . This motivates the notion of class, which measures the complexity of normalising length-three words. A quadratic normalisation  $(A, N)$  is of **left-class**  $m$  (respectively, **right-class**  $n$ ) if the equality  $N(w) = \overline{N}_{12[m]}(w)$  (respectively,  $N(w) = \overline{N}_{21[n]}(w)$ ) holds for every  $A$ -word of length three. A quadratic normalisation  $(A, N)$  is of **class**  $(m, n)$  if it is of left-class  $m$  and of right-class  $n$ .

**Example 3.2.4.** Consider the quadratic normalisation given in Example 3.1.3. If  $S_+$  has only one element, then there is only one  $S_+$ -word of length three and it is  $N$ -normal. So,  $(S_+, N)$  is of class  $(0, 0)$  in this case. Now, assume that  $S_+$  has at least two elements. Then for all  $w$  in  $S_+^3$ , one finds that the words  $\overline{N}_{121}(w)$  and  $\overline{N}_{212}(w)$  are  $N$ -normal. Namely, assuming that the word  $r|s|t$  is normal, compute  $\overline{N}_{121}$  and  $\overline{N}_{212}$  of  $t|s|r$  which is the worst-case scenario:

$$t|s|r \xrightarrow{\overline{N}_1} s|t|r \xrightarrow{\overline{N}_2} s|r|t \xrightarrow{\overline{N}_1} r|s|t, \quad t|s|r \xrightarrow{\overline{N}_2} t|r|s \xrightarrow{\overline{N}_1} r|t|s \xrightarrow{\overline{N}_2} r|s|t.$$

So,  $(S_+, N)$  is of class  $(3, 3)$ . A smaller class cannot be obtained in general, as witnessed by  $\overline{N}_{12}(s|s|r) = s|r|s$  and  $\overline{N}_{21}(s|r|r) = r|s|r$ , with  $r$  and  $s$  being any two distinct elements of  $S_+$ .

A class of a quadratic normalisation  $(A, N)$  can be characterised by relations involving only the restriction  $\overline{N}$ , as follows.

**Proposition 3.2.5** ([7, Proposition 3.14]). *A quadratic normalisation  $(A, N)$  is*

(1) *of left-class  $m$  if, and only if*

$$\overline{N}_{12[m]} = \overline{N}_{12[m+1]} = \overline{N}_{21[m+1]},$$

(2) *of right-class  $n$  if, and only if*

$$\overline{N}_{21[n]} = \overline{N}_{21[n+1]} = \overline{N}_{12[n+1]},$$

(3) *of class  $(m, m)$  if, and only if*

$$\overline{N}_{12[m]} = \overline{N}_{21[m]},$$

where all the equalities are understood to hold on  $A^3$ .

The **minimal left-class** of  $(A, N)$  is the smallest natural number  $m$  such that  $(A, N)$  is of left-class  $m$  if such  $m$  exists, and  $\infty$  otherwise. The minimal right-class  $n$  of  $(A, N)$  is defined analogously. Then the minimal class of  $(A, N)$  is the pair  $(m, n)$ .

**Lemma 3.2.6** ([7, Lemma 3.13]). *The minimal class of quadratic normalisation is either of the form  $(m, n)$  with  $|m - n| \leq 1$ , or  $(\infty, \infty)$ .*

**3.3. Quadratic normalisation of class  $(4, 3)$ .** The class  $(4, 3)$  has particularly nice computational properties (see [7, Section 4]), not shared by the higher classes (see [7, Example 3.23]), thanks to a diagrammatic tool called the domino rule. Let  $A$  be a set and  $F : A^2 \rightarrow A^2$  a map. We say that the **domino rule** is valid for  $F$  if for all  $s_1, s_2, s'_1, s'_2, t_0, t_1, t_2$  in  $A$  such that  $F(t_0|s_1) = s'_1|t_1$  and  $F(t_1|s_2) = s'_2|t_2$ , the following implication holds: if  $s_1|s_2$  and  $s'_1|t_1$  and  $s'_2|t_2$  are fixed points of  $F$ , then so is  $s'_1|s'_2$ . The domino rule corresponds to the following commutative diagram.

$$(3.3) \quad \begin{array}{ccccc} \bullet & \xrightarrow{s'_1} & \bullet & \xrightarrow{s'_2} & \bullet \\ \downarrow t_0 & & \downarrow t_1 & & \downarrow t_2 \\ \bullet & \xrightarrow{s_1} & \bullet & \xrightarrow{s_2} & \bullet \end{array}$$

(Note: In the original image, there are arcs indicating fixed points. A dashed arc connects the top two dots, labeled  $s'_1|s'_2$ . Solid arcs connect the bottom two dots, labeled  $s_1|s_2$ . Solid arcs also connect the middle dot to the top and bottom dots, labeled  $s'_1|t_1$  and  $s'_2|t_2$  respectively.)

Arcs denote fixed points for  $F$ : the solid ones are assumptions, and the dashed one is the expected conclusion.

**Proposition 3.3.1** ([7, Lemma 4.2]). *A quadratic normalisation  $(A, N)$  is of class  $(4, 3)$  if, and only if, the domino rule is valid for  $\overline{N}$ .*

The domino rule allows one to devise a simple universal recipe for computing the normalisation map. We refer the reader to [7, Proposition 4.4] for details, while here we only recall the key step.

**Lemma 3.3.2** ([7, Lemma 4.5]). *Let  $(A, N)$  be a quadratic normalisation. If  $(A, N)$  is of class  $(4, 3)$ , then for every  $t$  in  $A$  and every  $N$ -normal  $A$ -word  $s_1|\cdots|s_q$ , we have*

$$N(t|s_1|\cdots|s_q) = \overline{N}_{1|2|\cdots|q-1|q}(t|s_1|\cdots|s_q).$$

*Remark 3.3.3.* Note, in particular, that Lemma 3.3.2 implies that the leftmost letter of  $N(t|s_1|\cdots|s_q)$  does not depend on  $s_2, \dots, s_q$ , but only on  $t$  and  $s_1$ . We will use this observation in Subsection 4.2.

By Proposition 3.2.5, given a quadratic normalisation  $(A, N)$  of class  $(4, 3)$ , the map  $\overline{N}$  satisfies

$$(3.4) \quad \overline{N}_{1212} = \overline{N}_{212} = \overline{N}_{2121}.$$

One of the major results of [7] is that every idempotent map satisfying (3.4) arises from such a normalisation.

**Proposition 3.3.4** ([7, Proposition 4.7]). *If  $A$  is a set and  $F : A^2 \rightarrow A^2$  is an idempotent map satisfying*

$$F_{1212} = F_{212} = F_{2121},$$

*then there is a quadratic normalisation  $(A, N)$  of class  $(4, 3)$  such that  $\overline{N} = F$ .*

Thus, quadratic normalisations of class  $(4, 3)$  are fully axiomatised.

**3.4. Rewriting system associated to quadratic normalisation.** There is a simple correspondence between quadratic normalisations and quadratic rewriting systems.

**Proposition 3.4.1** ([7, Proposition 3.7]).

- (1) *If  $(S_+, N)$  is a quadratic for a monoid  $M$ , we obtain a quadratic, reduced, normalising and confluent rewriting system  $(S_+, R)$  presenting  $M$ , by defining  $R$  as the set of rewriting rules of the form*

$$s|t \rightarrow \overline{N}(s|t)$$

*for all  $s$  and  $t$  in  $S_+$  such that  $s|t$  is not  $N$ -normal.*

- (2) *Conversely, if  $(S_+, R)$  is a quadratic, reduced, normalising and confluent rewriting system presenting a monoid  $M$ , then  $(S_+, N)$  is a quadratic normalisation for  $M$ , where*

$$N(u) = \widehat{u}$$

*for every  $S$ -word  $u$ .*

- (3) *These two correspondences are inverse to each other.*

Proposition 3.4.1(1) can be adapted to a case when there is an  $N$ -neutral element. In fact, an  $N$ -neutral element does not affect termination of the associated rewriting system.

**Proposition 3.4.2** ([7, Proposition 3.9]).

- (1) *If  $(S, N)$  is a quadratic normalisation mod  $e$  for a monoid  $M$ , then we obtain a reduced, normalising and confluent rewriting system  $(S_+, R)$  presenting  $M$ , by defining  $R$  as the set of rewriting rules of the form*

$$s|t \rightarrow \pi_e(\overline{N}(s|t))$$

*for all  $s$  and  $t$  in  $S_+$  such that  $s|t$  is not  $N$ -normal.*

- (2) *If the rewriting system in Proposition 3.4.1(1) terminates, then so does the one in (1).*

The rewriting system associated to a quadratic normalisation in Proposition 3.4.1 need not be terminating (see [7, Proposition 5.7]). However, it is terminating for quadratic normalisations of class  $(4, 3)$ .

**Proposition 3.4.3** ([7, Proposition 5.4]). *If  $(A, N)$  is a quadratic normalisation of class  $(4, 3)$ , then the associated rewriting system is convergent. More precisely, every rewriting sequence from an element of  $A^p$  has length at most  $2^p - p - 1$ .*

**3.5. Left-weighted normalisation.** Before closing this section, let us recall a notion to be used in the next section. A normalisation  $(S, N)$  for a monoid  $M$  is called **left-weighted** if, for all  $s, t, s', t'$  in  $S$ , the equality  $s'|t' = \overline{N}(s|t)$  implies  $s \preceq s'$  in  $M$ .


**Proposition 3.5.1** ([7, Proposition 6.10]). *Assume that  $S$  is a generating subset of a left-cancellative monoid  $M$  containing no nontrivial invertible element. Assume that  $(S, N)$  is a quadratic normalisation mod 1 for  $M$ . Then the following are equivalent.*

- For every element  $g$  of  $M$ , the  $N$ -normal word  $s_1| \cdots |s_q$  representing  $g$  enjoys the following property for all  $i < q$ ,

$$(3.5) \quad \forall t \in S, \forall f \in M, (t \preceq fs_i s_{i+1} \implies t \preceq fs_i).$$

- The normalisation  $(S, N)$  is of class  $(4, 3)$  and is left-weighted.

The property (3.5) is called being  **$S$ -greedy** (at the  $i$ th position). It can be expressed diagrammatically. Commutativity of the following diagram without the dashed arrow implies the existence of a dashed arrow making the square (and thus also the triangle) commute.



The arc joining  $s_i$  and  $s_{i+1}$  denotes greediness. This notion is extensively studied in Garside theory. To an interested reader, we recommend the monograph [6] on the subject.

#### 4. CORRESPONDENCE BETWEEN FACTORABILITY AND NORMALISATION

In this section, we establish a correspondence between factorability structures and quadratic normalisations. Subsection 4.1 gives a characterisation of factorable monoids in terms of quadratic normalisation. Subsection 4.2 shows that, although the class of a quadratic normalisation corresponding to factorable monoid is not of class  $(4, 3)$  in general, it is so if a defining condition of local factorability structure is strengthened. Finally, we show that this strengthened definition implies the additional assumption which is introduced in [12] to obtain termination of the associated rewriting system.

**4.1. Characterisation of factorable monoids.** In this subsection, we give a necessary and sufficient condition, in terms of a quadratic normalisation, for a monoid to be factorable. We achieve this through a syntactic correspondence between a local factorability structure and the restriction of a normalisation map to length-two words. The property (4) of Definition 2.2.4 is going to be essential in deriving our main result so, for convenience, we are going to express it compactly using the domino rule.

**Definition 4.1.1.** Let  $(A, N)$  be a quadratic normalisation having the  $N$ -neutral element  $e$ . We say that the **weak domino rule** is valid for  $\overline{N}$  if the domino rule fails for  $\overline{N}$  only when the upper-right part of the diagram (3.3) contains  $e$ .

Now, we can state our main result.

**Theorem 4.1.2.** *A monoid  $(M, S)$  is factorable if, and only if, it admits a quadratic normalisation  $(N, S)$  mod  $e$  such that the weak domino rule is valid for  $\overline{N}$ .*

The rest of this subsection presents a rather straightforward proof, relying on the relation between factorability structure and local factorability structure, as stated in Theorem 2.2.6. First we verify that a factorability structure yields a quadratic normalisation, rather canonically. Assume that  $(M, S, \eta)$  is a factorable monoid. Since  $N_\varphi$  does not enjoy the length-preserving property in general, it does not make a suitable candidate for a normalisation map in the sense of Subsection 3.1. To fix this, we introduce the following notation.

**Notation.** *Assume that  $(M, S, \eta)$  is a factorable monoid. Denote by  $N'_\varphi$  the point-wise length-preserving extended form of  $N_\varphi$ , defined as follows:*

$$w \mapsto N'_\varphi(w) |1^m, \text{ with } m = |w| - |N_\varphi(w)|.$$

**Lemma 4.1.3.** *Assume that  $(M, S, \eta)$  is a factorable monoid. Then  $(S, N'_\varphi)$  is a quadratic normalisation.*

*Proof.* Let us check that  $N'_\varphi$  enjoys the three defining properties of a normalisation (Definition 3.1.1). Properties (1) and (2) are satisfied by construction. Indeed,  $N'_\varphi$  is length-preserving by definition and  $N'_\varphi(s) = s$  for  $s \in S$ . Lemma 2.1.7 implies property (3), namely  $N'_\varphi(u|v|w) = N'_\varphi(u|N'_\varphi(v)|w)$ .

We verify that the obtained normalisation  $(S, N'_\varphi)$  is quadratic. The property (3) of  $N_\varphi$  (Definition 2.2.2) implies the property (2) of Definition (3.2.1) of quadratic normalisation and, consequently, also the right-to-left implication of the property (1) of quadratic normalisation. The other direction of the property (1) of quadratic normalisation follows from Lemma 2.1.7.  $\square$

We say that  $(S, N'_\varphi)$  is the **quadratic normalisation corresponding to a factorability structure  $\eta$** .

*Remark 4.1.4.* A note on formality is in order before we continue. As announced by Remark 1.1.1, we have tried to respect the two original conventions: of taking a generating set  $S$  to be a subset of a factorable monoid; and of distinguishing  $S$  from its image under an evaluation map arising from a normalisation. Now, however, that we consider those quadratic normalisations  $(S, N)$  which correspond to a factorability structure, we interchangeably write  $e$  and  $1$  for the  $N$ -neutral  $S$ -letter, depending on whether we take the viewpoint of normalisation or factorability.

*Remark 4.1.5.* Note that, by construction, the restriction  $\overline{N'_\varphi}$  of  $N'_\varphi$  to length-two words is identically equal to the local factorability structure of  $(M, S, \eta)$ . We often use this fact implicitly.

Having obtained a quadratic normalisation, we want to find out its class. To that end, let us reinforce Lemma 2.2.5.

**Lemma 4.1.6.** *Assume that  $(M, S, \eta)$  is a factorable monoid. Then the quadratic normalisation corresponding to  $\eta$  is of class  $(5, 4)$ .*

*Proof.* Let  $(a, b, c) \in S^3$ . Denote by  $\varphi$  the local factorability structure corresponding to  $\eta$  in the sense of Theorem 2.2.6. If  $c = 1$ , then  $\varphi_1(a, b, c)$  equals  $\varphi(a, b) |1$  which is everywhere stable. If  $b = 1$ , then  $\varphi_{21}(a, b, c)$  equals  $\varphi(a, c) |1$  which is

everywhere stable. If  $a = 1$ , then  $\varphi_{121}(a, b, c)$  equals  $\varphi_{212}(a, b, c)$  which equals  $\varphi(b, c) | 1$  which is, again, everywhere stable.

If  $(a, b, c)$  contains no 1, then by Lemma 2.2.5, we get that  $N'_\varphi(a, b, c)$  equals  $\varphi_{2121}(a, b, c)$ .

We conclude that quadratic normalisation  $(N'_\varphi, S)$  is of right-class 4. Then Lemma 3.2.6 grants the left-class 5.  $\square$

The following example demonstrates that the minimal right-class of a quadratic normalisation corresponding to a factorability structure is not smaller than 4 in general.

**Example 4.1.7** ([11, Example 2.1.13]). Consider the monoid  $(\mathbb{Z}, +)$  with respect to the generating set  $S = \{-1, +1\}$ . The factorisation map is defined by  $g \mapsto (\text{sgn}(g), g - \text{sgn}(g))$ , where  $\text{sgn} : \mathbb{Z} \rightarrow \{-1, 0, +1\}$  denotes the sign function. One can check that this is a factorable monoid (technically speaking, this is a factorable group by [16, Example 3.2.2], and a weakly factorable monoid and hence, by [11, Proposition 2.1.28], it is also a factorable monoid). Note that  $\varphi_{212}(1, -1, -1) = (0, -1, 0)$  whereas  $\varphi_{2121}(1, -1, -1) = (-1, 0, 0)$ .

Our next example shows that the minimal left-class of a quadratic normalisation corresponding to a factorability structure is not smaller than 5 in general.

**Example 4.1.8.** Consider the factorable monoid  $M$  given in [12, Proposition .7]. Computing  $\varphi_{1212}(c_1, b_2, a_2)$  gives us

$$(c_1, b_1, a_1) \xrightarrow{\varphi_1^1} (c_1, b_1, a_1) \xrightarrow{\varphi_2^2} (c_1, b_2, a_2) \xrightarrow{\varphi_1^1} (c_2, b_3, a_2) \xrightarrow{\varphi_2^2} (c_2, e_2, 1),$$

whereas

$$\varphi_{12121}(c_1, b_2, a_2) = \varphi_1(c_2, e_2, 1) = (g_2, f_2, 1).$$

Hence the minimal left-class of the corresponding quadratic normalisation is at least 5.


The above computation also gives  $\varphi_{212}(c_1, b_1, a_1) = (c_2, e_2, 1)$ , whereas

$$\varphi_{2121}(c_1, b_1, a_1) = \varphi_1(c_2, e_2, 1) = (g_2, f_2, 1).$$

Thus, the minimal right-class is at least 4, as expected according to Lemma 3.2.6.

The previous two examples witness that the estimate of class in Lemma 4.1.6 is as good as one can hope for. On the other hand, observe that not every quadratic normalisation of class  $(5, 4)$  is corresponding to a factorability structure, as demonstrated by the following example (adapted from [7, Example 3.15]).

**Example 4.1.9.** Let  $A = \{a, b_1, b_2, b_3, b_4, b_5\}$  and let  $R$  consist of the rules  $ab_i \rightarrow ab_{i+1}$  for  $i < 5$  even and  $b_i a \rightarrow b_{i+1} a$  for  $i < 5$  odd. The rewriting system  $(A, R)$  is clearly quadratic and reduced. Observe that it is also terminating because each rewriting rule only increases the subscript of a letter  $b$  in a word. Furthermore, it is confluent, as illustrated by the following diagram (it suffices to investigate so-called critical pairs, see e.g. [14, Section 1]).



Denote by  $(A, N)$  the quadratic normalisation associated to  $(A, R)$  by Proposition 3.4.1. Let us determine the minimal class of  $(A, N)$ . If a length-three word does not begin and end with the letter  $a$ , it is  $N$ -normal or becomes  $N$ -normal in a single step. On the other hand, normalising  $a|b_1|a$  looks like:

$$a|b_1|a \begin{pmatrix} \overline{N}_1 \\ \mapsto \end{pmatrix} a|b_1|a \begin{pmatrix} \overline{N}_2 \\ \mapsto \end{pmatrix} a|b_2|a \begin{pmatrix} \overline{N}_1 \\ \mapsto \end{pmatrix} a|b_3|a \begin{pmatrix} \overline{N}_2 \\ \mapsto \end{pmatrix} a|b_4|a \begin{pmatrix} \overline{N}_1 \\ \mapsto \end{pmatrix} a|b_5|a.$$

So, the minimal class of  $(A, N)$  is  $(5, 4)$ .

However, the normalisation  $(A, N)$  is not corresponding to any factorability structure. To see this, observe that  $\overline{N}_{212}(a, b_1, a) = (a, b_4, a)$  whereas  $\overline{N}_{2121}(a, b_1, a) = (a, b_5, a)$ . Thus,  $\overline{N}$  fails to admit the property (4) of Definition 2.2.4. Therefore, it does not correspond to a factorability structure, by Remark 4.1.5.

The natural question then is: among quadratic normalisations of class  $(5, 4)$ , what distinguishes those which correspond to a factorability structure? Example 4.1.9 suggests a candidate (by what it is lacking): simply impose the weak domino rule upon quadratic normalisation of class  $(5, 4)$ . Before testing sufficiency of these two properties combined, let us observe that they are not independent from each other.

**Lemma 4.1.10.** *Assume that  $(A, N)$  is a quadratic normalisation having the  $N$ -neutral element  $e$ , such that the weak domino rule is valid for  $\overline{N}$ . Then  $(A, N)$  is of class  $(5, 4)$ .*

*Proof.* Let  $(r, s, t)$  be an  $A$ -word. If  $\overline{N}_{2121}(r, s, t) = \overline{N}_{212}(r, s, t)$ , then  $N(r, s, t) = \overline{N}_{212}(r, s, t)$ ; so it takes at most three steps starting with  $\overline{N}_2$  to normalise  $(r, s, t)$ .

Otherwise,  $\overline{N}_{212}(r, s, t)$  contains  $e$  by assumption. Denote  $(a, b, c) := \overline{N}_{212}(r, s, t)$ .

*Case 1.* If  $e$  occurs exactly once in the triple  $(a, b, c)$ , it cannot be at the leftmost position after  $\overline{N}_1$ . Then it has to be  $c = e$  after the last  $\overline{N}_2$ . In that case,

$$(4.1) \quad N(a, b, c) = \overline{N}_1(a, b, c) = \overline{N}_{2121}(r, s, t).$$

*Case 2.* If  $e$  occurs exactly twice in the triple  $(a, b, c)$ , then either  $a \neq e$  or  $b \neq e$ . In any case, (4.1) holds.

*Case 3.* If  $e$  occurs three times in the triple  $(a, b, c)$ , then clearly (4.1) holds.

Therefore,  $(A, N)$  is of right class 4, hence of class  $(5, 4)$ , by Lemma 3.2.6.  $\square$

The following proposition shows that adding the weak domino rule to the properties of quadratic normalisation (and thus granting class  $(5, 4)$  by Lemma 4.1.10), suffices to yield factorability.

**Proposition 4.1.11.** *Assume that  $(S, N)$  is a quadratic normalisation mod  $e$  for  $M$ . Assume additionally that the weak domino rule is valid for  $\overline{N}$ . Then  $\overline{N}$  is a local factorability structure.*

*Proof.* The following list shows that the map  $\overline{N}$  enjoys all the properties of Definition 2.2.4.

(1) By Proposition 3.2.3(3),  $M$  admits the presentation

$$\langle S_+ \mid \{s|t = \pi_e(\overline{N}(s|t)) \mid s, t \in S_+\} \rangle.$$

Although this presentation is not the same as the one in Definition 2.2.4, the two are equivalent (see Remark 2.4.2).

- (2) By the property (3) of Definition 3.1.1 of normalisation,  $\overline{N}$  is idempotent.
- (3) By Proposition 3.2.3(2), the equality  $\overline{N}(e|s) = s|e$  holds for all  $s \in S$ .
- (4) This property is explicitly assumed.
- (5) By Lemma 4.1.10, the normalisation  $(S, N)$  is of class (5, 4). By the definition of right-class 4, the  $N$ -normal word  $N(r, s, t)$  equals  $\overline{N}_{2121}(r, s, t)$  which further equals  $\overline{N}_{12121}(r, s, t)$  by the definition of left-class 5. Hence,  $N(r, s, t)$  equals  $N\overline{N}_1(r, s, t)$  for all  $(r, s, t)$  in  $S_+^3$ .

□

We have, thus, proved one direction of Theorem 4.1.2. The other one follows from Remark 4.1.5, as the restriction to length-two words of a quadratic normalisation corresponding to a factorability structure enjoys all the properties of the corresponding local factorability structure and, in particular, the weak domino rule is valid. Thereby, we have completed the proof of Theorem 4.1.2.

The following corollary is an immediate consequence.

**Corollary 4.1.12.**

- (1) *Associating a factorability structure to a quadratic normalisation such that the weak domino rule is valid, and associating a quadratic normalisation (the weak domino rule is valid automatically) to a factorability structure, as given above, are inverse transformations.*
- (2) *Normal forms with respect to a factorability structure and to the corresponding quadratic normalisation are the same.*
- (3) *The rewriting systems associated to a factorability structure and to the corresponding quadratic normalisation are equivalent, the only difference being dummy letters to preserve length in the latter.*

**4.2. Factorability in relation to quadratic normalisation of class (4, 3).**

Having established a general correspondence between a factorability structure and a quadratic normalisation in the previous subsection, we are now going to further elaborate these links in the case when the quadratic normalisation involved is of class (4, 3). First we emphasise a particular, yet important, consequence of Proposition 4.1.11.

**Corollary 4.2.1.** *Assume that  $(S, N)$  is a quadratic normalisation of class (4, 3) mod  $e$  for  $M$ . Then  $(M, S, \eta_N)$  is a factorable monoid.*

Although the converse does not hold in general, it does so in the case of graded monoids, as defined in Subsection 3.1.

**Lemma 4.2.2.** *Assume that  $(M, S_+, \eta)$  is a graded factorable monoid. Then  $(S_+, N'_\varphi)$  is a quadratic normalisation of class (4, 3).*

*Proof.* In addition to the conclusion of Lemma 4.1.3, observe that the property of being of right-class 3 follows from the property (4) of Definition 2.2.4 and the assumption that  $M$  is graded. Then the property of being of left-class 4 follows from Lemma 3.2.6. □

Let us point out precisely which defining properties of quadratic normalisation of class (4, 3) mod 1 for  $M$  do not necessarily arise from a factorability structure on  $M$ . In other words, we are looking for a (not too restrictive) stronger property that

would complement a factorability structure to a quadratic normalisation of class (4, 3). The property (4) of Definition 2.2.4 grants the equality

$$(4.2) \quad \varphi_{2121}(r, s, t) = \varphi_{212}(r, s, t)$$

only for every  $(r, s, t)$  in  $S_+^3$  such that  $\varphi_{212}(r, s, t)$  contains no 1. On the other hand, the right-class 3 (i.e. the domino rule) requires the equality (4.2) to hold for every  $(r, s, t) \in S^3$ , regardless of whether  $\varphi_{212}(r, s, t)$  contains 1 or not. Note that Definition 2.2.4 already implies the equality (4.2) in case that any component of the argument equals 1, that is for all  $(r, s, t)$  in  $S^3 \setminus S_+^3$ . Therefore, in order for a factorability structure to induce a quadratic normalisation of class (4, 3), it suffices to strengthen the condition (4) of Definition 2.2.4, as follows.

**Proposition 4.2.3.** *Assume that  $(M, S, \eta)$  is a factorable monoid. Assume additionally that the equality (4.2) holds for every  $(r, s, t) \in S_+^3$ . Then  $(S, N'_\varphi)$  is a quadratic normalisation of class (4, 3) mod 1 for  $M$ .*

We have shown that a quadratic normalisation of class (4, 3) yields a factorability structure (Corollary 4.2.1), but not vice versa (Examples 4.1.7 and 4.1.8, or Theorem 4.1.2). However, a factorability structure does yield a quadratic normalisation of class (4, 3) under a stronger condition on local factorability (Proposition 4.2.3). Therefore, under the same condition, the rewriting system associated to a factorable monoid is terminating, by Proposition 3.4.3 and Corollary 4.1.12(3).

From another point of view, recall that Theorem 2.4.3 ensures termination of the rewriting system associated to a factorable monoid, under an additional assumption on the factorisation map. It is then natural to ask ourselves if there is a relation between the strengthened condition of Proposition 4.2.3 and the additional assumption of Theorem 2.4.3, which are both known to ensure termination of the associated rewriting system.

In the rest of the present subsection, we investigate the relation between these two optional properties of a factorable monoid  $(M, S, \eta)$ : for all  $(r, s, t) \in S_+^3$ , the condition

$$(4.3) \quad (\eta\mu)_{2121}(r, s, t) = (\eta\mu)_{212}(r, s, t),$$

and, for all  $s \in S_+$  and  $f \in M$ , the assumption

$$(4.4) \quad (sf)' = (sf')', \quad \overline{sf} = \overline{sf'} \cdot \overline{f}.$$

*Remark 4.2.4.* Note that (4.4) trivially holds in case  $f \in S$  or in case that  $(s, f)$  is stable.

It is already known that (4.4) implies (4.3), as follows.

**Lemma 4.2.5** ([12, Lemma 7.1]). *Let  $(M, S, \eta)$  be a factorable monoid satisfying the additional assumption (4.4). Then the maps  $(\eta\mu)_{212}$  and  $(\eta\mu)_{2121}$  coincide on  $M^3$ .*

**Corollary 4.2.6.** *Let  $(M, S, \eta)$  be a factorable monoid satisfying the additional assumption (4.4). Then the condition (4.3) holds.*

For the reader's convenience, we adapt the proof of Lemma 4.2.5 here, to prove Corollary 4.2.6.

*Proof.* Multiply (4.4) on the left by  $\eta_2$ , and on the right by  $\mu_2$  to get

$$\eta_2\eta_1\mu_1\mu_2 = \eta_2\mu_2\eta_1\mu_1\eta_2\mu_2 : S \times M \times M \rightarrow M \times M \times M$$

which implies

$$\eta_2\eta_1\mu_1\mu_2 = \varphi_2\varphi_1\varphi_2 : S^3 \rightarrow S^3.$$

Note that the composition  $\eta_2\eta_1$  produces the  $\eta$ -normal form by definition, so  $\varphi_2\varphi_1\varphi_2$  is everywhere stable by Lemma 2.1.7. Thus, the property (4.3) holds.  $\square$

In the opposite direction, a partial result is already known. Namely, (4.3) implies (4.4) under certain (quite restrictive) additional requirements imposed on both the monoid and the normalisation. The next lemma is a straightforward adaptation of [6, Proposition IV.1.49]. It is also hinted by the proof of [15, Corollary 7.4.5].

*Remark 4.2.7.* Relying on Theorem 4.1.2 and, in particular, Corollary 4.1.12(2), we are going to abuse terminology by saying “the normal form” without specifying whether it arises from factorability or normalisation.

**Lemma 4.2.8.** *Let  $(S, N)$  be a left-weighted quadratic normalisation of class (4, 3) mod 1 for a left-cancellative monoid  $M$  containing no nontrivial invertible element. Then  $(M, S, \eta_N)$  satisfies the condition (4.4).*

*Proof.* We need to show that the equalities  $(sf)' = (sf')'$  and  $\overline{sf} = \overline{sf'} \cdot \overline{f}$  hold for any  $s \in S_+$  and  $f \in M$ .

By the definition of factorability structure, we have  $(sf')' \preceq sf'$ , and  $f' \preceq f$  hence also  $sf' \preceq sf$ . The transitivity gives  $(sf')' \preceq sf'$ , and the assumption that  $(S, N)$  is left-weighted then yields  $(sf')' \preceq (sf)'$ .

By Corollary 4.1.12(2), the normal form of  $f$  has the form  $f'|s_2| \cdots |s_p$ . Hence,  $(sf)' \preceq sf = sf's_2 \cdots s_p$ . Since  $(sf)'$  lies in  $S$  and  $f'|s_2| \cdots |s_p$  is normal, we obtain  $(sf)' \preceq sf'$  by the property (3.5) in Proposition 3.5.1.

Since  $M$  is a left-cancellative monoid containing no nontrivial invertible element, we conclude that  $(sf)' = (sf')'$ . Then  $\overline{sf} = \overline{sf'} \cdot \overline{f}$  follows from the left cancellation property.  $\square$

We want to find out whether the condition (4.3) implies the assumption (4.4) in general (i.e. without all the additional requirements of the previous lemma) or the latter is strictly stronger than the former. We start by considering length-two words.

**Lemma 4.2.9.** *Let  $(M, S, \eta)$  be a factorable monoid such that the condition (4.3) holds. Then the assumption (4.4) is valid for  $|f| = 2$ .*

*Proof.* The idea is to equate two different expressions of the normal form of  $sf$ , in order to obtain (4.4). Fix arbitrary  $s \in S_+$  and  $f \in M$  such that  $|f| = 2$ .

First compute the  $\eta$ -normal form of  $sf$ , by definition:

$$sf \xrightarrow{\eta_1} ((sf)', \overline{sf}) \xrightarrow{\eta_2} ((sf)', \overline{sf'}, \overline{sf}).$$

Note that  $|f| = 2$  implies  $\overline{f} \in S_+$ . Hence we have  $(s, f', \overline{f}) \in S_+^3$ . So we have another length-three  $S_+$ -word evaluating to  $sf$ . By the condition (4.3), the word  $(\eta\mu)_{212}(s, f', \overline{f})$  is everywhere stable, hence also normal by Lemma 4.1.3 and the property (1) of Definition 3.2.1 of quadratic normalisation. Hence,  $(\eta\mu)_{212}(s, f', \overline{f})$

is the  $N'_\varphi$ -normal form of the evaluation of  $(s, f', \overline{f})$ , i.e. of  $sf \in M$ . Since computing  $(\eta\mu)_{212}(s, f', \overline{f})$  depends on whether any 1 occurs in the process, we consider the following cases.

*Case 1.* If  $sf' \notin S$ , then

$$(s, f', \overline{f}) \xrightarrow{(\eta\mu)_2} (s, f', \overline{f}) \xrightarrow{(\eta\mu)_1} ((sf')', \overline{sf'}, \overline{f}) \xrightarrow{(\eta\mu)_2} ((sf')', (\overline{sf'} \cdot \overline{f})', \overline{sf' \cdot f}).$$

Comparing the result to  $\text{nf}_\eta(sf) = ((sf)', \overline{sf'}, \overline{sf})$  yields (4.4), by Corollary 4.1.12(2). Indeed, equating the first components of the obtained normal forms gives  $(sf)'\overline{sf} = (sf')'\overline{sf}$ . Equating the second and the third components gives  $\overline{sf} = \overline{sf'} \cdot \overline{f}$ .

*Case 2.* If  $sf' \in S_+$ , then

$$(4.5) \quad (s, f', \overline{f}) \xrightarrow{(\eta\mu)_2} (s, f', \overline{f}) \xrightarrow{(\eta\mu)_1} ((sf')', 1, \overline{f}) \xrightarrow{(\eta\mu)_2} ((sf')', \overline{f}, 1).$$

Again, equating the first components of the result and  $\text{nf}_\eta(sf)$  gives  $(sf)'\overline{sf} = (sf')'\overline{sf}$ . Equating the second and the third components now gives  $\overline{sf} = \overline{f}$  but note that, in the present case, this equality is equivalent to  $\overline{sf} = \overline{sf'} \cdot \overline{f}$ .

Finally, observe that the case  $sf' = 1$  cannot occur. Namely, if  $sf'$  were equal to 1, then applying  $(\eta\mu)_1$  after (4.5) would yield  $(\overline{f}, 1, 1)$  since  $\overline{f}$  is in  $S_+$ , and that would contradict the condition (4.3).  $\square$

Lemma 4.2.9 suggests itself as the base case for an induction, which we achieve in Proposition 4.2.10. First we introduce some notation, to simplify its proof.

**Notation.** If  $(s_1, s_2, \dots, s_n)$  is the normal form of  $f$  in  $M$ , denote  $\underline{f} := s_1 s_2 \cdots s_{n-1}$ .

**Proposition 4.2.10.** *Let  $(M, S, \eta)$  be a factorable monoid such that the condition (4.3) holds. Then the assumption (4.4) holds in  $M$ .*

*Proof.* Fix an arbitrary  $s$  in  $S$  and  $f$  in  $M$  such that  $|f| = n$  is strictly greater than 1. Observe that Remark 3.3.3 (on Lemma 3.3.2) implies the equality  $(sf)'\overline{sf} = (sf')'\overline{sf}$ .

We need to show that the equality  $\overline{sf} = \overline{sf'} \cdot \overline{f}$  also holds. We prove this using induction on  $n$ . For  $n = 2$ , the assumption (4.4) is granted by Lemma 4.2.9.

Assume that the assumption (4.4) is valid for every  $g \in M$  satisfying  $2 \leq |g| < n$ . Since  $|\underline{f}| = n - 1$ , we have  $\overline{\underline{f}} = \overline{sf'} \cdot \overline{f}$  by the induction hypothesis. Denote the normal form of  $f$  by  $(r_1, r_2, \dots, r_n)$ . Multiplying the equality  $\overline{\underline{f}} = \overline{sf'} \cdot \overline{f}$  by  $r_n$  on the right yields  $\overline{\underline{f}} \cdot r_n = \overline{sf'} \cdot \overline{f}$ , hence it suffices to show that the equality  $\overline{\underline{f}} \cdot r_n = \overline{sf}$  holds.


Let us compute the normal forms of  $\overline{\underline{f}}$  and  $\overline{sf}$  using Lemma 3.3.2. Denoting  $t_i|s_i := \overline{N}(s_{i-1}|r_i)$  for  $i \in \{2, \dots, n-1\}$ , we obtain

$$\text{nf}(\overline{\underline{f}}) = t_2|t_3|\cdots|t_{n-1}|s_{n-1}$$

and

$$\text{nf}(\overline{sf}) = t_2|t_3|\cdots|t_n|s_n,$$

as displayed by the following diagram, with arcs having the same meaning as in the diagram (3.3).



Therefore,  $\overline{sf} = t_2 t_3 \cdots t_{n-1} s_{n-1}$ . Multiplying by  $r_n$  on the right yields

$$\overline{sf} \cdot r_n = t_2 t_3 \cdots t_{n-1} s_{n-1} r_n = t_2 t_3 \cdots t_{n-1} t_n = \overline{sf}.$$

By induction, we conclude that (4.4) holds.  $\square$

The results of the present subsection enable the following characterisation. Note that the implications (1)  $\implies$  (2)  $\implies$  (3) were already known.

**Proposition 4.2.11.** *Assume that  $(M, S, \eta)$  is a factorable monoid. Then the following properties are equivalent.*

- (1) *For every  $s \in S$  and  $f \in M$ , the equalities  $(sf)' = (sf)'$  and  $\overline{sf} = \overline{sf'} \cdot \overline{f}$  hold.*
- (2) *For every  $(f, g, h) \in M^3$ , the equality  $(\eta\mu)_{2121}(f, g, h) = (\eta\mu)_{212}(f, g, h)$  holds.*
- (3) *For every  $(r, s, t) \in S^3$ , the equality  $(\eta\mu)_{2121}(r, s, t) = (\eta\mu)_{212}(r, s, t)$  holds.*
- (4) *Normalisation  $(S, N'_\varphi)$  is a quadratic normalisation of class (4, 3) mod 1 for  $M$ . In other words, the domino rule is valid for  $\overline{N'_\varphi}$ .*

## REFERENCES

- [1] Leonid A. Bokut, Yuqun Chen, Weiping Chen and Jing Li, *New approaches to plactic monoid via Gröbner–Shirshov bases*, J. Algebra 423 (2015) 301–317.
- [2] Kenneth S. Brown, *The geometry of rewriting systems: a proof of the Anick–Groves–Squier theorem*, Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989), vol. 23, Math. Sci. Res. Inst. Publ. New York: Springer, 1992, pp. 137–163.
- [3] Carl-Friedrich Bödigheimer, Balázs Visy, *Factorable groups and their homology*, Mathematisches Forschungsinstitut Oberwolfach, Report No. 32/2010 DOI: 10.4171/OWR/2010/32 Cohomology of Finite Groups: Interactions and Applications.
- [4] Alan J. Cain, Robert D. Gray, António Malheiro, *Finite Gröbner–Shirshov bases for Plactic algebras and biautomatic structures for Plactic monoids*, J. Algebra 423 (2015) 37–52.
- [5] Daniel E. Cohen. *String rewriting and homology of monoids*. Math. Structures Comput. Sci., 7(3):207–240, 1997.
- [6] Patrick Dehornoy, with François Digne, Eddy Godelle, Daan Krammer, Jean Michel, *Foundations of Garside theory*, vol. 22, EMS Tracts in Mathematics, European Mathematical Society (EMS), Zürich, 2015, pp. xviii+691.
- [7] Patrick Dehornoy and Yves Guiraud, *Quadratic normalization in monoids*, Internat. J. Algebra Comput. 26.5 (2016), pp. 935–972.
- [8] Robin Forman. *Morse theory for cell complexes*, Adv. Math., 134(1):90–145, 1998.
- [9] Frank Garside, *The braid group and other groups*, Quart. J. Math. Oxford Ser. (2) 20 (1969), 235–254.
- [10] Stéphane Gaussent, Yves Guiraud, and Philippe Malbos, *Coherent presentations of Artin monoids*, Compos. Math. 151 (5 2015), pp. 957–998.
- [11] Alexander Hess, *Factorable monoids: Resolutions and homology via discrete Morse theory*, <http://hss.ulb.uni-bonn.de/2012/2932/2932.pdf>, 2012, PhD thesis.
- [12] Alexander Hess and Viktoriya Ozornova, *Factorability, string rewriting and discrete Morse theory*, arXiv:1412.3025, 2014.

- [13] Daan Krammer, *An asymmetric generalisation of Artin monoids*, Groups Complex. Cryptol. 5.2 (2013), pp. 141–167.
- [14] Yves Lafont and Alain Prouté, *Church-Rosser property and homology of monoids*, Math. Structures Comput. Sci. 1.3 (1991), pp. 297–326.
- [15] Viktoriya Ozornova, *Factorability, discrete Morse theory, and a reformulation of  $K(\pi, 1)$ -conjecture*, <http://hss.ulb.uni-bonn.de/2013/3117/3117.pdf>, 2013, PhD thesis.
- [16] Balázs Visy, *Factorable groups and their homology*, <http://hss.ulb.uni-bonn.de/2011/2559/2559.pdf>, 2011, PhD thesis.
- [17] Rui Wang, *Homology computations for mapping class groups, in particular for  $\Gamma_{3,1}^0$* , <http://hss.ulb.uni-bonn.de/2011/2610/2610.pdf>, 2011, PhD thesis.

(Alen Đurić) UNIVERSITÉ PARIS CITÉ, CNRS, INRIA, IRIF, F-75013, PARIS, FRANCE  
Email address: [alen.djuric@protonmail.com](mailto:alen.djuric@protonmail.com)