

---

# UNIQUENESS OF ORDERS AND PARAMETERS IN MULTI-TERM TIME-FRACTIONAL DIFFUSION EQUATIONS BY SHORT-TIME BEHAVIOR

---

**Yikan LIU**

Research Center of Mathematics for Social Creativity, Research Institute for Electronic Science,  
Hokkaido University, N12W7, Kita-Ward, Sapporo 060-0812, Japan.  
ykliu@es.hokudai.ac.jp

**Masahiro YAMAMOTO**

Graduate School of Mathematical Sciences, The University of Tokyo  
3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan;  
Honorary Member of Academy of Romanian Scientists  
Ilfov, nr. 3, Bucuresti, Romania;  
Correspondence member of Accademia Peloritana dei Pericolanti, Palazzo Università  
Piazza S. Pugliatti 1 98122 Messina, Italy.  
myama@next.odn.ne.jp

January 2, 2023

## ABSTRACT

As the most significant difference from parabolic equations, long-time or short-time behavior of solutions to time-fractional evolution equations is dominated by the fractional orders, whose unique determination has been frequently investigated in literature. Unlike all the existing results, in this article we prove the uniqueness of orders and parameters (up to a multiplier for the latter) only by principal terms of asymptotic expansions of solutions near  $t = 0$  at a single spatial point. Moreover, we discover special conditions on unknown initial values or source terms for the coincidence of observation data. As a byproduct, we can even conclude the uniqueness for initial values or source terms by the same data. The proof relies on the asymptotic expansion after taking the Laplace transform and the completeness of generalized eigenfunctions.

**Keywords** Multi-term time-fractional diffusion equation · Parameter inverse problem · Uniqueness · Short-time asymptotic behavior

**MR(2010) Subject Classification** 35R30 · 35R11 · 58J99

## 1 Introduction

Let  $T \in \mathbb{R}_+ := (0, \infty)$  and  $\Omega \subset \mathbb{R}^d$  ( $d \in \mathbb{N} = \{1, 2, \dots\}$ ) be a bounded domain with smooth boundary  $\partial\Omega$ . For a constant  $m \in \mathbb{N}$ , let  $\alpha_j, q_j$  ( $j = 1, \dots, m$ ) be positive constants such that  $1 > \alpha_1 > \alpha_2 > \dots > \alpha_m > 0$ . We define an elliptic operator  $L : \mathcal{D}(L) := H^2(\Omega) \cap H_0^1(\Omega) \rightarrow L^2(\Omega)$  by

$$Lh(\mathbf{x}) := -\operatorname{div}(\mathbf{a}(\mathbf{x})\nabla h(\mathbf{x})) + \mathbf{b}(\mathbf{x}) \cdot \nabla h(\mathbf{x}) + c(\mathbf{x})h(\mathbf{x}), \quad \mathbf{x} \in \Omega,$$

where  $\cdot$  and  $\nabla = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d})$  refer to the inner product in  $\mathbb{R}^d$  and the gradient in  $\mathbf{x}$ , respectively. We emphasize that  $L$  is not necessarily symmetric. Here

$$\mathbf{a} = (a_{ij})_{1 \leq i, j \leq d} \in C^1(\overline{\Omega}; \mathbb{R}^{d \times d}), \quad \mathbf{b} = (b_1, \dots, b_d) \in C^1(\overline{\Omega}; \mathbb{R}^d), \quad c \in C(\overline{\Omega}) \quad (1.1)$$

are real-valued  $\mathbf{x}$ -dependent matrix-, vector- and scalar-valued functions, respectively. In particular, we assume that  $\mathbf{a}$  is symmetric and strictly positive-definite on  $\overline{\Omega}$ , i.e., there exists a constant  $\delta > 0$  such that

$$a_{ij}(\mathbf{x}) = a_{ji}(\mathbf{x}), \quad \mathbf{a}(\mathbf{x})\boldsymbol{\xi} \cdot \boldsymbol{\xi} \geq \delta|\boldsymbol{\xi}|^2, \quad \forall i, j = 1, \dots, d, \quad \forall \mathbf{x} \in \overline{\Omega}, \quad \forall \boldsymbol{\xi} \in \mathbb{R}^d, \quad (1.2)$$

where  $|\boldsymbol{\xi}|^2 = \boldsymbol{\xi} \cdot \boldsymbol{\xi}$ .

In this article, we are concerned with the following initial-boundary value problem for a multi-term time-fractional diffusion equation

$$\begin{cases} \left( \sum_{j=1}^m q_j \partial_t^{\alpha_j} + L \right) u(\mathbf{x}, t) = \rho(t)f(\mathbf{x}), & (\mathbf{x}, t) \in \Omega \times (0, T), \\ u(\mathbf{x}, 0) = a(\mathbf{x}), & \mathbf{x} \in \Omega, \\ u(\mathbf{x}, t) = 0, & (\mathbf{x}, t) \in \partial\Omega \times (0, T). \end{cases} \quad (1.3)$$

Here  $\partial_t^{\alpha_j}$  denotes the  $\alpha_j$ -th order Caputo derivative in time, which is traditionally defined by (e.g. Podlubny [28])

$$\partial_t^{\alpha_j} h(t) := \frac{1}{\Gamma(1 - \alpha_j)} \int_0^t \frac{h'(s)}{(t - s)^{\alpha_j}} ds, \quad h \in C^1[0, \infty).$$

Here  $\Gamma(\cdot)$  stands for the Gamma function. Throughout this article, we basically assume  $\rho \in L^1(0, T)$  and for convenience, we understand  $\rho \in L^1(\mathbb{R}_+)$  by identifying it with its zero extension in  $(T, \infty)$ .

The governing equation in (1.3) is called a single-term time-fractional diffusion equation if  $m = 1$ , while it is called a multi-term one if  $m \geq 2$ . In the last decades, they have gathered increasing population as typical nonlocal models for anomalous diffusion in heterogeneous media or fractals (e.g. [4, 8]). Especially, linear theories for the case of  $m = 1$  have been established in recent years. Among many references, we refer only to [7, 16, 29] and see the references therein. Compared with classical parabolic equations, single-term time-fractional diffusion equations inherit the same time-analyticity and the maximum principle and see e.g. [27, 29]. Meanwhile, they also share quantitative similarities in the smoothing effect and the unique continuation property (e.g. [10, 29]). In contrast, they show essential difference from their integer counterpart in the asymptotic behavior as  $t \rightarrow \infty$ , which indicates the rate  $t^{-\alpha}$  (e.g. [29]).

Though less studied, it reveals that the case of  $m \geq 2$  in (1.3) also behaves similarly to that of  $m = 1$ . We refer to [20] for the well-posedness and the limited smoothing effect of solutions, and [25, 26] for weak and strong positivity properties, respectively. In [5, 17], the parameters  $q_j$  in (1.3) are generalized to be  $\mathbf{x}$ -dependent at the cost of slightly weaker solution regularity. Remarkably, the short- and long-time asymptotic behavior of solutions depend on the highest order  $\alpha_1$  and the lowest one  $\alpha_m$ , respectively (e.g. [17, 20]).

Since the orders of fractional derivatives are closely related to the heterogeneity and related physical properties of media, the determination of  $\alpha_j$  in (1.3) turns out to be significant in both theory and practice. For the single-term case in one-dimensional case, Cheng et al. [6] first proved the uniqueness of  $\alpha_1$  by the data  $u|_{\{\mathbf{x}_0\} \times (0, T)}$  with  $\mathbf{x}_0 \in \partial\Omega$ . Employing the above mentioned asymptotic behavior, Hatano et al. [9] deduced inversion formulae

$$\alpha_1 = - \lim_{t \rightarrow \infty} \frac{t \partial_t u(\mathbf{x}_0, t)}{u(\mathbf{x}_0, t)} = \lim_{t \rightarrow 0} \frac{t \partial_t u(\mathbf{x}_0, t)}{u(\mathbf{x}_0, t) - a(\mathbf{x}_0)}$$

with  $\mathbf{x}_0 \in \Omega$ . For the multi-term case, Li and Yamamoto [22] showed the unique determination of  $m, \alpha_j, q_j$  with constants  $q_j > 0$  by  $u|_{\{\mathbf{x}_0\} \times (0, T)}$  with  $\mathbf{x}_0 \in \Omega$ . For  $\mathbf{x}$ -dependent  $q_j \geq 0$ , Li et al. [19] proved the unique determination of  $m, \alpha_j, q_j$  and the zeroth-order coefficient  $c$  using the Dirichlet-to-Neumann map, and Li et al. [17] uniquely identified  $\alpha_j$  by  $u|_{\{\mathbf{x}_0\} \times (0, T)}$  with  $\mathbf{x}_0 \in \Omega$ . We refrain from providing a complete list, and refer to Li et al. [21] as a survey covering most literature on parameter inverse problems for (1.3) before 2019. We notice that due to the dimension of observation data, most literature dealt with the simultaneous identification of orders and unknown sources or coefficients in (1.3) (see [15, 23, 24, 31, 35, 36]). Only [18] obtained the stability in determining a single order by the data

$u(\mathbf{x}_0, t_0)$ . As recent progress, we refer to [3, 33] for the unique determination of time and space fractional orders, and Yamamoto [34] for treating a non-symmetric  $L$ . Very recently, Jin and Kian [11, 12] uniquely recovered  $m, \alpha_j, q_j$  in (1.3) in an unknown medium, that is,  $L$  is unknown.

Among the above literature on parameter inverse problems for (1.3), the majority used the observation of  $u$  at  $\{\mathbf{x}_0\} \times (0, T)$ . However, due to technical difficulties, almost all publications assume the symmetry of  $L$ , that is,  $\mathbf{b} \equiv \mathbf{0}$ , and only uniqueness was proved with data of solution such as  $u(\mathbf{x}_0, t)$  at a monitoring point  $\mathbf{x}_0 \in \Omega$  over a time interval. On the other hand, in the existing publications, since the observation data are a function of  $t$  and there are only finite unknowns, such an inverse problem is overdetermined, which indicates the possibility to reduce the accuracy of data or determine another unknown ingredient in (1.3) simultaneously. Moreover, since the orders  $\alpha_j$  greatly influence short-time behavior of the solution, we can expect that principal terms of asymptotic expansions of solutions near  $t = 0$  can characterize such orders, and we are motivated to establish the uniqueness for the following inverse problem.

**Problem 1.1** (Parameter inverse problem). *Let  $\mathbf{x}_0 \in \Omega$  and sufficiently small  $\tau > 0$  be fixed arbitrarily, and let  $u$  satisfy (1.3) with  $f \equiv 0$  or  $a \equiv 0$  in  $\Omega$ . Determine  $m$  and  $\alpha_j, q_j$  for  $j = 1, \dots, m$  simultaneously by asymptotic behavior of  $u(\mathbf{x}_0, t)$  for  $0 < t < \tau$ .*

In the formulation of Problem 1.1, as data we adopt the short-time behavior near  $t = 0$ , in other words, principal terms of the asymptotic expansion in  $t$  of the solution as  $t \downarrow 0$ , which is formulated by (3.2) later. We emphasize that we do not require any non-asymptotically observed quantities of the solution for the uniqueness of Problem 1.1, which is the main novelty of our article.

Moreover, the initial value  $a$  and the spatial component  $f$  of the source term in (1.3) are also assumed to be unknown. Finally, we mention that the elliptic part  $L$  can be non-symmetric, which seems also seldom handled in references.

Only by the information of principal terms of asymptotic expansions of solutions near  $t = 0$ , we establish the uniqueness in determining  $m$  and  $\alpha_j$  in (1.3) simultaneously in the respective cases  $f \equiv 0$  and  $a \equiv 0$ . Meanwhile, it reveals that  $q_j$  can be uniquely identified up to a common multiplier depending on the local information of the unknown initial value  $a$  or source term  $f$ , and the possible non-uniqueness is illustrated by some examples. Our uniqueness result can be regarded as the strong distinguishability effect of the fractional orders to the solution of (1.3).

On this direction, it can be proved that unknown initial values or source terms should satisfy rather special relations in order to produce coinciding asymptotics. As a corollary, in a special case we can even conclude the uniqueness of the initial value  $a$  or the spatial component  $f$  of the source term by data  $u(\mathbf{x}_0, \cdot)$ . It is surprising because the one-dimensional data can uniquely determine a  $d$ -dimensional unknown. This is indeed an underdetermined problem and the data are sometimes called sparse data in literature (e.g. [23]).

The remainder of this paper is organized as follows. In Section 2, we make preparations to state and prove the main results, especially the generalized eigenvalue theory of  $L$ . Next, we collect the main results concerning Problem 1.1 along with illustrative examples and remarks in Section 3. Then Section 4 is devoted to the proofs of main results, followed by conclusions provided in Section 5.

## 2 Preliminaries

Throughout this article, by  $C > 0$  we denote generic constants which may change line by line. For  $h \in L^1(\mathbb{R}_+)$ , we denote its Laplace transform by

$$\widehat{h}(p) := \int_{\mathbb{R}_+} e^{-pt} h(t) dt, \quad p > 0.$$

For  $h \in L^1(\mathbb{R}_+)$  and a constant  $\mu > -1$ , by the notation  $h(t) \sim t^\mu$  for small  $t > 0$ , we mean that there exists a constant  $C' \neq 0$  such that  $\lim_{t \rightarrow 0+} t^{-\mu} h(t) = C'$ . In this case, we know  $h(t) = O(t^\mu)$  but  $h(t) \neq o(t^{\mu'})$  for any  $\mu' > \mu$  a.e. as  $t \rightarrow 0+$ . We define  $\sim$  for a.e. large  $t > 0$  in the same manner.

In the sequel, the inner product of  $L^2(\Omega)$  is denoted by  $(\cdot, \cdot)$ . By  $\sigma(L) = \{\lambda_n\}_{n \in \mathbb{N}} \subset \mathbb{C}$  we denote the set of all distinct eigenvalues of  $L$ . Throughout this article, for our arguments we assume

$$0 \notin \sigma(L). \tag{2.1}$$

For the sake of self-containedness, we collect necessary ingredients concerning the eigenvalue theory of  $L$ , which can be found e.g. in Kato [14]. For each  $n \in \mathbb{N}$ , we take a circle  $\gamma_n$  centered at  $\lambda_n$  with sufficiently small radius such that  $\gamma_n$  does not enclose  $\lambda_{n'}$  with any  $n' \neq n$ . Defining an operator

$$P_n := -\frac{1}{2\pi\sqrt{-1}} \int_{\gamma_n} (L - z)^{-1} dz,$$

we know that

$$P_n P_{n'} = \begin{cases} P_n, & n' = n, \\ 0, & n' \neq n. \end{cases}$$

We call  $P_n$  the eigenprojection for  $\lambda_n$  and any  $\varphi \in P_n L^2(\Omega) \setminus \{0\}$  a generalized eigenfunction of  $\lambda_n$ . Meanwhile, the space  $P_n L^2(\Omega)$  is finite dimensional and we set  $d_n := \dim P_n L^2(\Omega) < \infty$ . We note that if  $L$  is symmetric, then there holds for all  $n \in \mathbb{N}$  that

$$\lambda_n \in \mathbb{R}, \quad P_n L^2(\Omega) = \ker(L - \lambda_n), \quad d_n = \dim \ker(L - \lambda_n). \quad (2.2)$$

Next, we introduce the eigennilpotent of  $\lambda_n$  as

$$D_n := (L - \lambda_n)P_n.$$

Then we can verify that  $P_n \varphi = \varphi$  for  $\varphi \in P_n L^2(\Omega)$  and

$$P_n L^2(\Omega) \subset \mathcal{D}(L), \quad L P_n L^2(\Omega) \subset P_n L^2(\Omega), \quad D_n P_n L^2(\Omega) \subset P_n L^2(\Omega), \quad D_n^{d_n} = (L - \lambda_n)^{d_n} P_n = 0.$$

Using  $P_n$  and  $D_n$ , we have the following Laurant expansion for the resolvent:

$$(L - z)^{-1} P_n = \frac{P_n}{\lambda_n - z} + \sum_{k=1}^{d_n-1} \frac{(-1)^k D_n^k}{(\lambda_n - z)^{k+1}}, \quad z \notin \sigma(L). \quad (2.3)$$

On the other hand, we recall the following resolvent estimate (see e.g. Tanabe [32]): there exist constants  $z_0 > 0$  and  $C > 0$  such that

$$\|(L + z)^{-1} h\|_{L^2(\Omega)} \leq \frac{C}{|z|} \|h\|_{L^2(\Omega)}, \quad \forall z > z_0, \quad \forall h \in L^2(\Omega). \quad (2.4)$$

Finally, we introduce the fractional power  $L^\gamma$  of  $L$  along with its domain  $\mathcal{D}(L^\gamma)$  for  $\gamma > 0$ . First for  $0 < \gamma < 1$ , the fractional operator  $L^{-\gamma}$  is defined by

$$L^{-\gamma} h := \frac{\sin \pi \gamma}{\pi} \int_{\mathbb{R}_+} z^{-\gamma} (L + z)^{-1} h dz, \quad h \in L^2(\Omega).$$

Then there holds  $L^{-\gamma} P_n L^2(\Omega) \subset P_n L^2(\Omega)$  and hence  $L^\gamma P_n L^2(\Omega) \subset P_n L^2(\Omega)$  for all  $\gamma \geq 0$ . Then for  $0 < \gamma < 1$ , we define  $L^\gamma$  as the inverse of  $L^{-\gamma}$ . For  $\gamma \in \mathbb{R}_+ \setminus \mathbb{N}$ , we define  $L^\gamma = L^{\gamma - \lfloor \gamma \rfloor} \circ L^{\lfloor \gamma \rfloor}$ , where  $\lfloor \gamma \rfloor := \max\{n \in \mathbb{Z} \mid n \leq \gamma\}$  stands for the flooring function and  $\circ$  denotes the composite. Then we set  $\mathcal{D}(L^\gamma) := \{h \in L^2(\Omega) \mid L^\gamma h \in L^2(\Omega)\}$  for  $\gamma > 0$ .

Now we recall the well-posedness results concerning the forward problems (1.3).

**Lemma 2.1.** *Assume (1.1)–(1.2) and (2.1). Let  $u$  satisfy (1.3) with  $f \equiv 0$  in  $\Omega$  and  $a \in \mathcal{D}(L^\gamma)$  with  $\gamma \geq 0$ . Then for any fixed constant  $\eta \in [0, 1)$ , the problem (1.3) admits a unique solution  $u \in C([0, \infty); \mathcal{D}(L^\gamma)) \cap C(\mathbb{R}_+; \mathcal{D}(L^{\gamma+\eta}))$ . Moreover, there exists a constant  $C > 0$  depending on  $\alpha_j, q_j$  ( $j = 1, \dots, m$ ),  $\Omega, L, \gamma$  such that*

$$\|u(\cdot, t)\|_{\mathcal{D}(L^{\gamma+\eta})} \leq C t^{-\alpha_1 \eta} e^{Ct} \|a\|_{\mathcal{D}(L^\gamma)}, \quad \forall t > 0. \quad (2.5)$$

The above lemma asserts that the solution norm  $\|u(\cdot, t)\|_{\mathcal{D}(L^{\gamma+\eta})}$  increases at most exponentially as  $t \rightarrow \infty$ , which guarantees the existence of its Laplace transform  $\hat{u}(\cdot, p)$  for large  $p > 0$ . Lemma 2.1 slightly improves Li, Huang and Yamamoto [17, Theorem 2.3] in the sense of more general choice of  $a$  and the key estimate (2.5). Especially, in [17, Theorem 2.3] the estimate (2.5) only held for  $t \in (0, T)$  with fixed  $T > 0$ . However, a close scrutiny into the proof reveals that  $T$  can be chosen arbitrarily. On the other hand, it was proved in [17, 20] that if  $\mathbf{b} \equiv \mathbf{0}$  and  $c \geq 0$  on  $\overline{\Omega}$ , then the solution  $u(\cdot, t)$  obeys the decay rate  $t^{-\alpha_m}$  as  $t \rightarrow \infty$ . Nevertheless, instead of the sharp decay estimate, here (2.5) is enough for applying the Laplace transform, which plays an essential role in the proof (see Section 4).

### 3 Main Results and Examples

Now we are well prepared to state the main results of this article. To this end, we introduce another initial-boundary value problem

$$\begin{cases} \left( \sum_{j=1}^{m'} r_j \partial_t^{\beta_j} + L \right) v(\mathbf{x}, t) = \rho(t)g(\mathbf{x}), & (\mathbf{x}, t) \in \Omega \times (0, T), \\ v(\mathbf{x}, 0) = b(\mathbf{x}), & \mathbf{x} \in \Omega, \\ v(\mathbf{x}, t) = 0, & (\mathbf{x}, t) \in \partial\Omega \times (0, T), \end{cases} \quad (3.1)$$

where  $m' \in \mathbb{N}$  is a constant and  $\beta_j, r_j$  ( $j = 1, \dots, m'$ ) are positive constants satisfying  $1 > \beta_1 > \beta_2 > \dots > \beta_{m'} > 0$ . We notice that (1.3) and (3.1) share the same elliptic part  $L$  and the temporal component  $\rho$  of the source term. Nevertheless, their initial values  $a, b$  and spatial components  $f, g$  of source terms differ from each other, which are basically assumed also to be unknown.

Let us state our first main result on the unique determination of orders and parameters with short-time asymptotic information for homogeneous problems, i.e.,  $f = g \equiv 0$  in (1.3) and (3.1).

**Theorem 3.1.** *Assume (1.1)–(1.2) and (2.1). Let  $u, v$  satisfy (1.3) and (3.1) respectively with  $f = g \equiv 0$  in  $\Omega$ , where  $a, b \in \mathcal{D}(L^\gamma)$  with  $\gamma > 1 + d/4$ . Choose  $\mathbf{x}_0 \in \Omega$  arbitrarily such that  $La(\mathbf{x}_0) \neq 0, Lb(\mathbf{x}_0) \neq 0$  and denote  $\kappa := La(\mathbf{x}_0)/Lb(\mathbf{x}_0)$ .*

(a) *If there exist constants  $C > 0$  and  $\nu > \min\{\alpha_1, \beta_1\}$  such that*

$$|u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)| \leq C t^\nu, \quad 0 \leq t \leq \tau, \quad (3.2)$$

where  $\tau > 0$  is some constant, then

$$\alpha_1 = \beta_1, \quad \frac{q_1}{r_1} = \kappa. \quad (3.3)$$

(b) *If further  $a, b \in \mathcal{D}(L^{1+\gamma})$  and (3.2) is satisfied with  $\nu > 2 \min\{\alpha_1, \beta_1\}$ , then (3.2) implies*

$$m = m', \quad \alpha_j = \beta_j, \quad \frac{q_j}{r_j} = \kappa, \quad j = 1, \dots, m. \quad (3.4)$$

In (3.2), we remark that  $\tau > 0$  can be arbitrarily small, and in view of (3.6), this assumption means that some principal parts of the asymptotic expansions of  $u(\mathbf{x}_0, t)$  and  $v(\mathbf{x}_0, t)$  coincide.

**Remark 3.2.** (a) According to the Sobolev embedding theorem (see Adams [1]), we see that  $a, b \in \mathcal{D}(L^\gamma)$  with  $\gamma > 1 + d/4$  implies  $a, b \in C^2(\bar{\Omega})$  and thus  $La(\mathbf{x}_0), Lb(\mathbf{x}_0)$  are well-defined. On the other hand, taking  $t = 0$  in (3.2), we automatically obtain

$$a(\mathbf{x}_0) = b(\mathbf{x}_0). \quad (3.5)$$

(b) Unlike most of the existing results on the uniqueness for inverse problems, we only need the short-time asymptotic behavior (3.2) instead of their coincidence. Moreover, we even do not need  $a = b$  as was assumed in most literature. Therefore, we minimize the assumptions on initial values to  $La(\mathbf{x}_0) \neq 0, Lb(\mathbf{x}_0) \neq 0$  and (3.5).

(c) The choices of the order  $\nu$  in (3.2) in Theorem 3.1(i)–(ii) turn out to be the minimum necessary condition used in the proof. If there is an a priori upper bound  $\bar{\alpha}$  of  $\min\{\alpha_1, \beta_1\}$ , then safer requirements of  $\nu$  would be  $\nu > \bar{\alpha}$  in (i) and  $\nu > 2\bar{\alpha}$  in (ii). If there is no a priori information on  $\alpha_1, \beta_1$ , then we should require  $\nu \geq 1$  in (i) and  $\nu \geq 2$  in (ii).

(d) Indeed, by the arguments in [20], one can prove

$$|u(\mathbf{x}_0, t) - a(\mathbf{x}_0)| = O(t^{\alpha_1}), \quad |v(\mathbf{x}_0, t) - b(\mathbf{x}_0)| = O(t^{\beta_1}) \quad \text{as } t \rightarrow 0,$$

which, together with (3.5), implies

$$|u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)| \leq |u(\mathbf{x}_0, t) - a(\mathbf{x}_0)| + |v(\mathbf{x}_0, t) - b(\mathbf{x}_0)| = O(t^{\min\{\alpha_1, \beta_1\}}) \quad \text{as } t \rightarrow 0. \quad (3.6)$$

However, (3.2) does not follow automatically from (3.6).

Since the key assumption (3.2) is satisfied if

$$\exists \tau > 0 \quad \text{such that } u(\mathbf{x}_0, t) = v(\mathbf{x}_0, t), \quad 0 \leq t \leq \tau, \quad (3.7)$$

Theorem 3.1 definitely covers the uniqueness with usual exact data. On the contrary, by an argument of the reverse proposition, we can immediately obtain the following distinguishability property: if there exists a constant  $\nu < \min\{\alpha_1, \beta_1\}$  such that

$$|u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)| \geq C t^\nu \quad \text{near } t = 0,$$

then there should be  $\alpha_1 \neq \beta_1$ .

In general, one can at most identify the parameters  $q_j$  up to a multiplier  $\kappa$ , but cannot further assert  $q_j = r_j$  ( $j = 1, \dots, m$ ) from (3.2). This is demonstrated by the following counterexample.

**Example 3.3.** In (1.3) and (3.1), let us simply take

$$\Omega = (0, \pi), \quad f = g \equiv 0 \text{ in } \Omega, \quad m = m' = 1, \quad \alpha_1 = \beta_1, \quad q_1 = 4, \quad r_1 = 1, \quad L = -\partial_x^2.$$

We pick any  $x_0 \in (0, \pi) \setminus \{\pi/2\}$  and select

$$a(x) = \frac{\sin 2x}{2 \cos x_0}, \quad b(x) = \sin x.$$

Then it is readily seen that

$$a(x_0) = b(x_0) = \sin x_0, \quad \frac{q_1}{r_1} = 4 = \frac{a''(x_0)}{b''(x_0)}.$$

However, employing the Mittag-Leffler function

$$E_{\alpha_1, 1}(z) := \sum_{\ell=0}^{\infty} \frac{z^\ell}{\Gamma(\alpha_1 \ell + 1)},$$

we can easily obtain the explicit solutions

$$u(x, t) = E_{\alpha_1, 1}(-t^{\alpha_1}) \frac{\sin 2x}{2 \cos x_0}, \quad v(x, t) = E_{\alpha_1, 1}(-t^{\alpha_1}) \sin x$$

and hence  $u(x_0, t) = E_{\alpha_1, 1}(-t^{\alpha_1}) \sin x_0 = v(x_0, t)$ . This means that even the exact data for all  $t \geq 0$  fails to guarantee  $q_1 = r_1$ .

Similarly to Theorem 3.1, we can obtain the same uniqueness with (3.2) for inhomogeneous problems, i.e.,  $a = b \equiv 0$  in (1.3) and (3.1).

**Theorem 3.4.** Assume (1.1)–(1.2) and (2.1). Let  $u, v$  satisfy (1.3) and (3.1) respectively with  $a = b \equiv 0$  in  $\Omega$ , where  $f, g \in \mathcal{D}(L^\gamma)$  with  $\gamma > d/4$  and there exists a constant  $\mu > -1$  such that

$$\rho(t) \sim t^\mu \quad \text{as } t \downarrow 0. \quad (3.8)$$

Fix sufficiently small  $\tau > 0$ , pick  $\mathbf{x}_0 \in \Omega$  such that  $f(\mathbf{x}_0) \neq 0, g(\mathbf{x}_0) \neq 0$  and denote  $\kappa := f(\mathbf{x}_0)/g(\mathbf{x}_0)$ .

- (a) If there exist constants  $C > 0$  and  $\nu > \min\{\alpha_1, \beta_1\} + \mu$  such that (3.2) is satisfied, then (3.3) holds.
- (b) If further  $f, g \in \mathcal{D}(L^{1+\gamma})$  and (3.2) is satisfied with  $\nu > 2 \min\{\alpha_1, \beta_1\} + \mu$ , then (3.2) implies (3.4).

The assumption (3.8) describes the short-time asymptotic behavior of the temporal component  $\rho$  of the source terms in (1.3) and (3.1), which turns out to be essential in proving Theorem 3.4. Indeed, since the information (3.2) is concerned only with small  $t > 0$ , we need more localized assumption on  $\rho$  than  $\rho \in L^1(\mathbb{R}_+)$ . On the other hand, the solution behavior near  $t = 0$  is also influenced by  $\rho$ . Therefore, compared with Theorem 3.1, the difference  $\nu - \mu$  of orders instead of  $\nu$  itself appears in the conditions guaranteeing the uniqueness in Theorem 3.4. We notice that (3.8) indicates  $\rho \not\equiv 0$  near  $t = 0$  and is not restrictive because it covers a wide choice of  $\rho$  behaving asymptotically like a power function for small  $t > 0$ . However, it does exclude e.g.

$$\rho(t) = \begin{cases} 0, & t = 0, \\ \exp(-1/t), & 0 < t < T, \end{cases}$$

which satisfies  $\rho^{(i)}(0) = 0$  for all  $i = 0, 1, \dots$ . With such functions, it is usually rather difficult to obtain the uniqueness or the stability for inverse problems. On the contrary, the ill-posedness is reduced if there exists  $i = 0, 1, \dots$  such that  $\rho^{(i)}(0) \neq 0$ , which falls within the framework of (3.8).

Similarly to Example 3.3, one can easily construct a counterexample against  $q_j = r_j$  ( $j = 1, \dots, m$ ) in the framework of Theorem 3.4. Owing to multiple unknown parameters, even a usual type of data (3.7) does not yield the uniqueness in determining coefficients  $q_j, r_j$ .

Finally, we discuss the uniqueness and the non-uniqueness by (3.7).

**Theorem 3.5.** *Assume that the elliptic operator  $L$  is symmetric, i.e.,  $\mathbf{b} \equiv \mathbf{0}$  on  $\overline{\Omega}$ . Let*

$$\ker(L - \lambda_n) = \text{span}\{\varphi_{n,k} \mid k = 1, \dots, d_n\},$$

where  $\varphi_{n,k}$  ( $k = 1, \dots, d_n$ ) are orthonormal eigenfunctions associated with  $\lambda_n$ . Define a countable set  $\Sigma := \{\lambda_n/\lambda_{n'} \mid n, n' \in \mathbb{N}\} \subset \mathbb{R}$ .

(a) Let  $u, v$  satisfy (1.3) and (3.1) respectively with  $f = g \equiv 0$  in  $\Omega$ . Under the same assumptions in Theorem 3.1, (3.7) holds if and only if

$$m = m', \quad \alpha_j = \beta_j \quad \frac{q_j}{r_j} = \frac{La(\mathbf{x}_0)}{Lb(\mathbf{x}_0)} =: \kappa \in \Sigma, \quad j = 1, \dots, m \quad (3.9)$$

and there exist nonempty sets  $M_\kappa, M'_\kappa \subset \mathbb{N}$  satisfying

$$\{\lambda_n \in \sigma(L) \mid n \in M_\kappa\} = \{\kappa\lambda_n \mid \lambda_n \in \sigma(L), n \in M'_\kappa\} \quad (3.10)$$

and a bijection  $\theta_\kappa : M_\kappa \rightarrow M'_\kappa$  such that

$$\begin{cases} P_n a(\mathbf{x}_0) = P_{\theta_\kappa(n)} b(\mathbf{x}_0), & P_n La(\mathbf{x}_0) = \kappa P_{\theta_\kappa(n)} Lb(\mathbf{x}_0), & n \in M_\kappa, \\ P_n a(\mathbf{x}_0) = P_n La(\mathbf{x}_0) = 0, & & n \notin M_\kappa, \\ P_n b(\mathbf{x}_0) = P_n Lb(\mathbf{x}_0) = 0, & & n \notin M'_\kappa. \end{cases} \quad (3.11)$$

(b) Let  $u, v$  satisfy (1.3) and (3.1) respectively with  $a = b \equiv 0$  in  $\Omega$ . Under the same assumptions in Theorem 3.4, (3.7) holds if and only if

$$m = m', \quad \alpha_j = \beta_j, \quad \frac{q_j}{r_j} = \frac{f(\mathbf{x}_0)}{g(\mathbf{x}_0)} =: \kappa \in \Sigma, \quad j = 1, \dots, m \quad (3.12)$$

and there exist the same sets  $M_\kappa, M'_\kappa \subset \mathbb{N}$  and bijection  $\theta_\kappa : M_\kappa \rightarrow M'_\kappa$  as that in (1) such that

$$\begin{cases} P_n f(\mathbf{x}_0) = \kappa P_{\theta_\kappa(n)} g(\mathbf{x}_0), & n \in M_\kappa, \\ P_n f(\mathbf{x}_0) = 0, & n \notin M_\kappa, \\ P_n g(\mathbf{x}_0) = 0, & n \notin M'_\kappa. \end{cases} \quad (3.13)$$

**Remark 3.6.** Theorem 3.5 characterizes the coincidence (3.7) of observation data via initial values  $a, b$  or source terms  $f, g$ . From the definition of  $\Sigma$ , it is obvious that  $1 \in \Sigma$ . We discuss the cases of  $\kappa = 1$ ,  $\kappa \notin \Sigma$  and  $\kappa \in \Sigma \setminus \{1\}$  separately.

(a) If  $\kappa = 1$ , i.e.,  $La(\mathbf{x}_0) = Lb(\mathbf{x}_0)$  or  $f(\mathbf{x}_0) = g(\mathbf{x}_0)$ , then  $M_1 = M'_1 = \mathbb{N}$  and  $\theta_1$  is the identity operator. In this case, it is readily seen that (3.11) and (3.13) are equivalent to

$$\begin{aligned} P_n a(\mathbf{x}_0) &= P_n b(\mathbf{x}_0), & P_n La(\mathbf{x}_0) &= P_n Lb(\mathbf{x}_0), & \forall n \in \mathbb{N}, \\ P_n f(\mathbf{x}_0) &= P_n g(\mathbf{x}_0), & & & \forall n \in \mathbb{N}, \end{aligned} \quad (3.14)$$

respectively.

(b) If  $\kappa \notin \Sigma$ , then Theorem 3.5 asserts that the observation data  $u(\mathbf{x}_0, t)$  and  $v(\mathbf{x}_0, t)$  ( $0 \leq t \leq \tau$ ) cannot coincide.

(c) If  $\kappa \in \Sigma \setminus \{1\}$ , then it is indicated in (3.11) and (3.13) that the initial values  $a, b$  or the source terms  $f, g$  should satisfy a rather special relation. To illustrate this point, we provide the following example.

**Example 3.7.** Similarly to Example 3.3, in (1.3) and (3.1) we take

$$\Omega = (0, \pi), \quad f = g \equiv 0 \text{ in } \Omega, \quad \kappa = \frac{La(x_0)}{Lb(x_0)} = 4, \quad L = -\partial_x^2.$$

Then it is readily seen that

$$\lambda_n = n^2, \quad \Sigma = \{(n/n')^2 \mid n, n' \in \mathbb{N}\} = \mathbb{Q}^2 \setminus \{0\}, \quad P_n f = (f, \varphi_n) \varphi_n, \quad f \in L^2(0, \pi),$$

where  $\varphi_n(x) := \sqrt{2/\pi} \sin nx$  ( $n \in \mathbb{N}$ ). Further, by the definitions of  $M_\kappa, M'_\kappa$  and  $\theta_\kappa$ , it is straightforward to verify that  $M_4 = 2\mathbb{N}, M'_4 = \mathbb{N}$  and  $\theta_4(n) = n/2$ . Therefore, the relation (3.11) can be rephrased as

$$(a, \varphi_{2n}) \varphi_{2n}(x_0) = (b, \varphi_n) \varphi(x_0), \quad (a, \varphi_{2n-1}) \varphi_{2n-1}(x_0) = 0, \quad n \in \mathbb{N}.$$

Especially, if  $x_0 \neq \pi\mathbb{Q}$ , i.e.,  $x_0$  is not a zero of  $\varphi_n$  ( $\forall n \in \mathbb{N}$ ), then  $a$  should take the special form of

$$a(x) = \sum_{n=1}^{\infty} \frac{(b, \varphi_n)}{\varphi_{2n}(x_0)} \varphi_{2n}(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=1}^{\infty} \frac{(b, \varphi_n)}{\cos nx_0} \sin 2nx,$$

which is a generalization of Example 3.3.

On the same direction of Theorem 3.5 and Example 3.7, it turns out that in a special case, we can even obtain the uniqueness of  $a, b$  or  $f, g$ .

**Corollary 3.8.** *In addition to the same assumptions in Theorem 3.5, we assume that all the eigenvalues of  $L$  are simple and*

$$\mathbf{x}_0 \notin \omega := \bigcup_{n=1}^{\infty} \{\mathbf{x} \in \Omega \mid \varphi_n(\mathbf{x}) = 0\} \subset \Omega, \quad (3.15)$$

where  $\varphi_n$  is an eigenfunction associated with  $\lambda_n$ . Let  $\Sigma, M_\kappa, M'_\kappa, \theta_\kappa$  be the same as that in Theorem 3.5.

(a) Let  $u, v$  satisfy (1.3) and (3.1) respectively with  $f = g \equiv 0$  in  $\Omega$ . Then (3.7) holds if and only if (3.9) holds and

$$a = \sum_{n \in M_\kappa} (a, \varphi_n) \varphi_n, \quad b = \sum_{n \in M'_\kappa} (b, \varphi_n) \varphi_n \quad (3.16)$$

$$\text{with } (a, \varphi_n) \varphi_n(\mathbf{x}_0) = (b, \varphi_{\theta_\kappa(n)}) \varphi_{\theta_\kappa(n)}(\mathbf{x}_0), \quad n \in M_\kappa.$$

Especially if  $\kappa = 1$ , i.e.  $La(\mathbf{x}_0) = Lb(\mathbf{x}_0)$ , then (3.16) is equivalent to  $a \equiv b$  in  $\Omega$ .

(b) Let  $u, v$  satisfy (1.3) and (3.1) respectively with  $a = b \equiv 0$  in  $\Omega$ . Then (3.7) holds if and only if (3.12) holds and

$$f = \sum_{n \in M_\kappa} (f, \varphi_n) \varphi_n, \quad g = \sum_{n \in M'_\kappa} (g, \varphi_n) \varphi_n \quad (3.17)$$

$$\text{with } (f, \varphi_n) \varphi_n(\mathbf{x}_0) = \kappa (g, \varphi_{\theta_\kappa(n)}) \varphi_{\theta_\kappa(n)}(\mathbf{x}_0), \quad n \in M_\kappa.$$

Especially if  $\kappa = 1$ , i.e.  $f(\mathbf{x}_0) = g(\mathbf{x}_0)$ , then (3.17) is equivalent to  $f \equiv g$  in  $\Omega$ .

The assumption (3.15) is a rank condition, which is widely used for observability of partial differential equations (e.g. [30]). In some special cases such as that in Example 3.7, the Lebesgue measure of  $\omega$  defined in (3.15) is zero. Therefore, the uniqueness claimed in Corollary 3.8 is generic, which is rather surprising because the single point observation can even almost determine the initial value or the spatial component of the source term uniquely.

## 4 Proofs of Main Results

For the proof of Theorem 3.1, we need the completeness of all the generalized eigenfunctions of  $L$ .

**Lemma 4.1.** (a) *For any  $h \in L^2(\Omega)$ , there exists a sequence  $\{h_N\}_{N \in \mathbb{N}} \subset L^2(\Omega)$  such that*

$$h_N \in \sum_{n=1}^N P_n L^2(\Omega), \quad \lim_{N \rightarrow \infty} \|h - h_N\|_{L^2(\Omega)} = 0.$$



(b) For any  $h \in \mathcal{D}(L^\gamma)$  with  $\gamma \geq 0$ , there exists a sequence  $\{h_N\}_{N \in \mathbb{N}} \subset \mathcal{D}(L^\gamma)$  such that

$$h_N \in \sum_{n=1}^N P_n L^2(\Omega), \quad \lim_{N \rightarrow \infty} \|L^\gamma(h - h_N)\|_{L^2(\Omega)} = 0.$$

*Proof.* (a) It follows immediately from the density of the linear subspace spanned by all generalized eigenfunctions of  $L$  in  $L^2(\Omega)$  (see Agmon [2]).

(b) Since  $h \in \mathcal{D}(L^\gamma)$  implies  $L^\gamma h \in L^2(\Omega)$ , we can apply (i) to choose a sequence  $\{\psi_N\}_{N \in \mathbb{N}} \subset L^2(\Omega)$  such that

$$\psi_N \in \sum_{n=1}^N P_n L^2(\Omega), \quad \lim_{N \rightarrow \infty} \|\psi_N - L^\gamma h\|_{L^2(\Omega)} = 0.$$

Owing to (2.1), we see that  $L^{-\gamma} : L^2(\Omega) \rightarrow L^2(\Omega)$  exists and is bounded, indicating

$$\lim_{N \rightarrow \infty} \|L^\gamma(L^{-\gamma}\psi_N) - L^\gamma h\|_{L^2(\Omega)} = 0.$$

By the definition of  $L^{-\gamma}$ , we see that  $h_N := L^{-\gamma}\psi_N \in \sum_{n=1}^N P_n L^2(\Omega) \subset \mathcal{D}(L^\gamma)$  and thus

$$\lim_{N \rightarrow \infty} \|L^\gamma h_N - L^\gamma h\|_{L^2(\Omega)} = 0.$$

The proof of Lemma 4.1 is complete. □

*Proof of Theorem 3.1.* Let  $f = g \equiv 0$  in  $\Omega$  in (1.3) and (3.1). We divide the proof into 4 steps.

**Step 1** To begin with, we make some overall preparations. Since the assumption  $\gamma - 1 > d/4$  implies the Sobolev embedding  $\mathcal{D}(L^{\gamma-1}) \subset C(\bar{\Omega})$ , it follows from Lemma 2.1 that  $u(\mathbf{x}_0, t)$  is well-defined for  $t > 0$  and fulfills the estimate

$$|u(\mathbf{x}_0, t)| \leq \|u(\cdot, t)\|_{C(\bar{\Omega})} \leq C \|u(\cdot, t)\|_{\mathcal{D}(L^{\gamma-1})} \leq C e^{Ct} \|a\|_{\mathcal{D}(L^{\gamma-1})}, \quad t > 0.$$

The similar estimate also holds for  $v(\mathbf{x}_0, t)$ , which, together with the key assumption (3.2), indicates

$$|u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)| \leq \begin{cases} C t^\nu, & 0 \leq t \leq \tau, \\ C e^{Ct}, & t > \tau. \end{cases}$$

Then for large  $p > 0$ , we estimate the Laplace transform of  $u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)$  from above as

$$\begin{aligned} \widehat{u}(\mathbf{x}_0, p) - \widehat{v}(\mathbf{x}_0, p) &\leq \int_{\mathbb{R}_+} e^{-pt} |u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)| dt \leq C \int_0^\tau e^{-pt} t^\nu dt + C \int_\tau^\infty e^{-(p-C)t} dt \\ &\leq C \int_{\mathbb{R}_+} e^{-pt} t^\nu dt + \frac{C e^{-(p-C)\tau}}{p-C} = \frac{C \Gamma(\nu+1)}{p^{\nu+1}} + o(p^{-\nu-1}) \leq C p^{-\nu-1}. \end{aligned}$$

We can similarly obtain the lower estimate  $\widehat{u}(\mathbf{x}_0, p) - \widehat{v}(\mathbf{x}_0, p) \geq -C p^{-\nu-1}$  for large  $p > 0$  and hence

$$\widehat{u}(\mathbf{x}_0, p) - \widehat{v}(\mathbf{x}_0, p) = O(p^{-\nu-1}) \quad \text{for large } p > 0. \quad (4.1)$$

Now we shall represent  $\widehat{u}(\mathbf{x}_0, p)$  and  $\widehat{v}(\mathbf{x}_0, p)$  explicitly for large  $p > 0$ . Performing the Laplace transform in (1.3) and using the formula

$$\widehat{\partial_t^\alpha h}(p) = p^\alpha \widehat{h}(p) - p^{\alpha-1} h(0),$$

we derive the boundary value problem for an elliptic equation for  $\widehat{u}(\cdot, p)$ :

$$\begin{cases} \left( \sum_{j=1}^m q_j p^{\alpha_j} + L \right) \widehat{u}(\cdot, p) = a \sum_{j=1}^m q_j p^{\alpha_j-1} & \text{in } \Omega, \\ \widehat{u}(\cdot, p) = 0 & \text{on } \partial\Omega. \end{cases}$$

Transforming (3.1) in the same manner and taking the inverses, we obtain

$$\widehat{u}(\cdot, p) = \frac{1}{p} \sum_{j=1}^m q_j p^{\alpha_j} \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} a, \quad (4.2)$$

$$\widehat{v}(\cdot, p) = \frac{1}{p} \sum_{j=1}^{m'} r_j p^{\beta_j} \left( L + \sum_{j=1}^{m'} r_j p^{\beta_j} \right)^{-1} b. \quad (4.3)$$

According to Lemma 4.1, there exist  $a_N = \sum_{n=1}^N a_{N,n}$  and  $b_N = \sum_{n=1}^N b_{N,n}$  ( $N \in \mathbb{N}$ ) such that

$$a_{N,n}, b_{N,n} \in P_n L^2(\Omega), \quad \lim_{N \rightarrow \infty} \|L^\gamma(a - a_N)\|_{L^2(\Omega)} = \lim_{N \rightarrow \infty} \|L^\gamma(b - b_N)\|_{L^2(\Omega)} = 0. \quad (4.4)$$

Especially, since  $\gamma > 1 + d/4$ , the Sobolev embedding implies

$$\lim_{N \rightarrow \infty} La_N(\mathbf{x}_0) = La(\mathbf{x}_0), \quad \lim_{N \rightarrow \infty} Lb_N(\mathbf{x}_0) = Lb(\mathbf{x}_0). \quad (4.5)$$

Then for any fixed  $N \in \mathbb{N}$ , we decompose  $a = a_N + (a - a_N)$  and employ the expansion (2.3) to rewrite (4.2) as

$$\begin{aligned} p\widehat{u}(\cdot, p) &= \sum_{j=1}^m q_j p^{\alpha_j} \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} \left\{ \sum_{n=1}^N a_{N,n} + (a - a_N) \right\} \\ &= \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} P_n a_{N,n} \\ &\quad + \left\{ \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right) - L \right\} \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} (a - a_N) \\ &= \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \sum_{k=0}^{d_n-1} \frac{(-1)^k D_n^k a_{N,n}}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}} + (a - a_N) - L \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} (a - a_N) \\ &= \sum_{n=1}^N \left( 1 - \frac{\lambda_n}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} \right) a_{N,n} + \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k D_n^k a_{N,n}}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}} \\ &\quad + a - a_N - \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} L(a - a_N) \\ &= a - \sum_{n=1}^N \frac{\lambda_n a_{N,n}}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} + \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k D_n^k a_{N,n}}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}} \\ &\quad - \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} L(a - a_N). \end{aligned} \quad (4.6)$$

Rewriting (4.3) in the same manner, we obtain

$$\begin{aligned} p\widehat{v}(\cdot, p) &= b - \sum_{n=1}^N \frac{\lambda_n b_{N,n}}{\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j}} + \sum_{j=1}^{m'} r_j p^{\beta_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k D_n^k b_{N,n}}{(\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j})^{k+1}} \\ &\quad - \left( L + \sum_{j=1}^{m'} r_j p^{\beta_j} \right)^{-1} L(b - b_N). \end{aligned} \quad (4.7)$$

Taking  $\mathbf{x} = \mathbf{x}_0$  in (4.6)–(4.7) and substituting them into (4.1), we notice (3.5) to arrive at

$$\sum_{i=1}^3 (I_N^i(p) - J_N^i(p)) = O(p^{-\nu}) \quad \text{for large } p > 0, \quad (4.8)$$

where

$$\begin{aligned} I_N^1(p) &:= \sum_{n=1}^N \frac{\lambda_n a_{N,n}(\mathbf{x}_0)}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}}, & J_N^1(p) &:= \sum_{n=1}^N \frac{\lambda_n b_{N,n}(\mathbf{x}_0)}{\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j}}, \\ I_N^2(p) &:= \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^{k+1} (D_n^k a_{N,n})(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}}, & J_N^2(p) &:= \sum_{j=1}^{m'} r_j p^{\beta_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^{k+1} (D_n^k b_{N,n})(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j})^{k+1}}, \\ I_N^3(p) &:= \left( \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} L(a - a_N) \right) (\mathbf{x}_0), & J_N^3(p) &:= \left( \left( L + \sum_{j=1}^{m'} r_j p^{\beta_j} \right)^{-1} L(b - b_N) \right) (\mathbf{x}_0). \end{aligned}$$

Here for  $I_N^3(p)$ , it follows from the Sobolev embedding and the estimate (2.4) that

$$\begin{aligned} |I_N^3(p)| &\leq \left\| \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} L(a - a_N) \right\|_{C(\bar{\Omega})} \leq C \left\| \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} L(a - a_N) \right\|_{\mathcal{D}(L^{\gamma-1})} \\ &\leq C \left\| \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} L^\gamma(a - a_N) \right\|_{L^2(\Omega)} \leq \frac{C}{\sum_{j=1}^m q_j p^{\alpha_j}} \|L^\gamma(a - a_N)\|_{L^2(\Omega)} \\ &\leq C p^{-\alpha_1} \|a - a_N\|_{\mathcal{D}(L^\gamma)}. \end{aligned} \quad (4.9)$$

Similarly, for  $J_N^3(p)$  we have

$$|J_N^3(p)| \leq C p^{-\beta_1} \|b - b_N\|_{\mathcal{D}(L^\gamma)}. \quad (4.10)$$

Finally, we note that the regularity of  $a, b$  guarantees the uniform boundedness of

$$\sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0), \quad \sum_{n=1}^N \sum_{k=1}^{d_n-1} (-1)^{k+1} (D_n^k a_{N,n})(\mathbf{x}_0), \quad \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0), \quad \sum_{n=1}^N \sum_{k=1}^{d_n-1} (-1)^{k+1} (D_n^k b_{N,n})(\mathbf{x}_0)$$

for all  $N \in \mathbb{N}$ . Especially, by the identity  $LP_n = \lambda_n P_n + D_n$  we know that

$$\begin{aligned} \sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0) + \sum_{n=1}^N D_n a_{N,n}(\mathbf{x}_0) &= \sum_{n=1}^N L a_{N,n}(\mathbf{x}_0) = L a_N(\mathbf{x}_0), \\ \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) + \sum_{n=1}^N D_n b_{N,n}(\mathbf{x}_0) &= \sum_{n=1}^N L b_{N,n}(\mathbf{x}_0) = L b_N(\mathbf{x}_0). \end{aligned} \quad (4.11)$$

**Step 2** Now we are well prepared to prove part (i). We starting with showing  $\alpha_1 = \beta_1$  by contradiction.

First let us assume that  $\alpha_1 < \beta_1$ . Our strategy is first multiplying both sides of (4.8) by  $p^{\alpha_1}$ , then passing  $N \rightarrow \infty$  and finally passing  $p \rightarrow \infty$  to deduce a contradiction. To this end, we calculate  $p^{\alpha_1} I_N^1(p)$ ,  $p^{\alpha_1} J_N^1(p)$ ,  $p^{\alpha_1} I_N^2(p)$  and  $p^{\alpha_1} J_N^2(p)$  as

$$p^{\alpha_1} I_N^1(p) = \sum_{n=1}^N \frac{\lambda_n a_{N,n}(\mathbf{x}_0)}{\lambda_n p^{-\alpha_1} + \sum_{j=1}^m q_j p^{\alpha_j - \alpha_1}} = \frac{1 + o(1)}{q_1} \sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0), \quad (4.12)$$

$$p^{\alpha_1} J_N^1(p) = \sum_{n=1}^N \frac{\lambda_n b_{N,n}(\mathbf{x}_0)}{\lambda_n p^{-\alpha_1} + \sum_{j=1}^{m'} r_j p^{\beta_j - \alpha_1}} = \frac{p^{\alpha_1 - \beta_1} (1 + o(1))}{r_1} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) = O(p^{\alpha_1 - \beta_1}), \quad (4.13)$$

$$\begin{aligned}
 p^{\alpha_1} I_N^2(p) &= \sum_{j=1}^m q_j p^{\alpha_j - \alpha_1} \sum_{n=1}^N \left\{ \frac{D_n a_{N,n}(\mathbf{x}_0)}{(\lambda_n p^{-\alpha_1} + \sum_{j=1}^m q_j p^{\alpha_j - \alpha_1})^2} + O(p^{-\alpha_1}) \right\} \\
 &= \frac{1 + o(1)}{q_1} \sum_{n=1}^N D_n a_{N,n}(\mathbf{x}_0) + O(p^{-\alpha_1}), \tag{4.14}
 \end{aligned}$$

$$p^{\alpha_1} J_N^2(p) = O(p^{\alpha_1 + \beta_1}) \sum_{n=1}^N O(p^{-2\beta_1}) \sum_{k=1}^{d_n-1} (-1)^{k+1} (D_n^k b_{N,n})(\mathbf{x}_0) = O(p^{\alpha_1 - \beta_1}) \tag{4.15}$$

for large  $p > 0$ . Substituting (4.12)–(4.15) and (4.9)–(4.10) into (4.8), we obtain

$$\begin{aligned}
 O(p^{-\nu + \alpha_1}) &= p^{\alpha_1} \sum_{i=1}^3 (I_N^i(p) - J_N^i(p)) \\
 &= \frac{1 + o(1)}{q_1} \left\{ \sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0) + \sum_{n=1}^N D_n a_{N,n}(\mathbf{x}_0) \right\} + O(p^{\alpha_1 - \beta_1}) + O(p^{-\alpha_1}) \\
 &\quad + O(1) \|a - a_N\|_{\mathcal{D}(L^\gamma)} + O(p^{\alpha_1 - \beta_1}) \|b - b_N\|_{\mathcal{D}(L^\gamma)} \\
 &= \frac{1 + o(1)}{q_1} L a_N(\mathbf{x}_0) + O(1) \|a - a_N\|_{\mathcal{D}(L^\gamma)} + o(1) \quad \text{for large } p > 0,
 \end{aligned}$$

where we used (4.11) and the uniform boundedness of  $\|b - b_N\|_{\mathcal{D}(L^\gamma)}$  with respect to  $N$ . By the assumption  $\nu > \min\{\alpha_1, \beta_1\} = \alpha_1$ , the above equation becomes

$$\frac{1 + o(1)}{q_1} L a_N(\mathbf{x}_0) + O(1) \|a - a_N\|_{\mathcal{D}(L^\gamma)} = o(1) \quad \text{for large } p > 0. \tag{4.16}$$

Then we pass  $N \rightarrow \infty$  in (4.16) and employ (4.4)–(4.5) to deduce

$$\frac{1 + o(1)}{q_1} L a(\mathbf{x}_0) = o(1) \quad \text{for large } p > 0.$$

Consequently, passing  $p \rightarrow \infty$  yields  $L a(\mathbf{x}_0) = 0$ , which contradicts with the assumption  $L a(\mathbf{x}_0) \neq 0$ .

In a completely parallel manner, assuming  $\alpha_1 > \beta_1$  results in a contradiction with the assumption  $L b(\mathbf{x}_0) \neq 0$ , indicating the only possibility  $\alpha_1 = \beta_1$ .

Now it suffices to verify  $q_1/r_1 = L a(\mathbf{x}_0)/L b(\mathbf{x}_0)$ . Similarly to the treatment for (4.12) and (4.14), we substitute  $\alpha_1 = \beta_1$  into  $p^{\alpha_1} J_N^1(p)$  and  $p^{\alpha_1} J_N^2(p)$  to see that now (4.13) and (4.15) become

$$p^{\alpha_1} J_N^1(p) = \frac{1 + o(1)}{r_1} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0), \quad p^{\alpha_1} J_N^2(p) = \frac{1 + o(1)}{r_1} \sum_{n=1}^N D_n b_{N,n}(\mathbf{x}_0) + O(p^{-\alpha_1}),$$

respectively. Substituting the above equalities, (4.12), (4.14), (4.9), (4.10) into (4.8) and repeating the same calculation as before, we arrive at

$$(1 + o(1)) \left( \frac{L a_N(\mathbf{x}_0)}{q_1} - \frac{L b_N(\mathbf{x}_0)}{r_1} \right) + O(1) (\|a - a_N\|_{\mathcal{D}(L^\gamma)} + \|b - b_N\|_{\mathcal{D}(L^\gamma)}) = o(1)$$

for large  $p > 0$ . Again passing first  $N \rightarrow \infty$  and then  $p \rightarrow \infty$ , we conclude

$$\frac{L a(\mathbf{x}_0)}{q_1} - \frac{L b(\mathbf{x}_0)}{r_1} = 0$$

or equivalently  $q_1/r_1 = L a(\mathbf{x}_0)/L b(\mathbf{x}_0)$ . This completes the proof of part (i).

**Step 3** From now on we proceed to the proof of part (ii). For later convenience, we set  $\kappa := L a(\mathbf{x}_0)/L b(\mathbf{x}_0)$ . Notice that owing to the further assumption  $a, b \in \mathcal{D}(L^{1+\gamma})$ , the sequences

$$\sum_{n=1}^N \lambda_n^2 a_{N,n}(\mathbf{x}_0), \quad \sum_{n=1}^N \lambda_n^2 b_{N,n}(\mathbf{x}_0), \quad \sum_{n=1}^N \lambda_n (D_n a_{N,n})(\mathbf{x}_0), \quad \sum_{n=1}^N \lambda_n (D_n b_{N,n})(\mathbf{x}_0)$$

are also uniformly bounded for all  $N \in \mathbb{N}$ .

In this step, we shall show by induction that for  $j = 1, 2, \dots, \min\{m, m'\}$ , there holds

$$\alpha_j = \beta_j, \quad \frac{q_j}{r_j} = \kappa. \quad (4.17)$$

Since the assumption of  $\nu$  in part (ii) is stronger than that in part (i), the case of  $j = 1$  was already shown in Step 2.

Now let us assume that for some  $\ell = 2, \dots, \min\{m, m'\}$ , (4.17) holds for  $j = 1, \dots, \ell - 1$ . Then our aim is to show that (4.17) also holds for  $j = \ell$ . Notice that now the terms  $J_N^1(p)$  and  $J_N^2(p)$  are written as

$$\begin{aligned} J_N^1(p) &= \sum_{n=1}^N \frac{\lambda_n b_{N,n}(\mathbf{x}_0)}{\lambda_n + \kappa^{-1} \sum_{j=1}^{\ell-1} q_j p^{\alpha_j} + \sum_{j=\ell}^{m'} r_j p^{\beta_j}}, \\ J_N^2(p) &= \left( \kappa^{-1} \sum_{j=1}^{\ell-1} q_j p^{\alpha_j} + \sum_{j=\ell}^{m'} r_j p^{\beta_j} \right) \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^{k+1} (D_n^k b_{N,n})(\mathbf{x}_0)}{(\lambda_n + \kappa^{-1} \sum_{j=1}^{\ell-1} q_j p^{\alpha_j} + \sum_{j=\ell}^{m'} r_j p^{\beta_j})^{k+1}}. \end{aligned} \quad (4.18)$$

Similarly to Step 2, we start with showing  $\alpha_\ell = \beta_\ell$  by contradiction.

Again let us first assume that  $\alpha_\ell < \beta_\ell$ . Now our strategy is first multiplying both sides of (4.8) by  $p^{2\alpha_1 - \beta_\ell}$ , then passing  $N \rightarrow \infty$  and finally passing  $p \rightarrow \infty$  to deduce a contradiction.

We first calculate  $p^{2\alpha_1 - \beta_\ell} (I_N^1(p) - J_N^1(p))$ . To this end, we rewrite the denominator in  $I_N^1(p)$  as

$$\begin{aligned} \frac{1}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} &= \frac{1}{\lambda_n + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j}} \left( 1 + \frac{\sum_{j=\ell}^m q_j p^{\alpha_j}}{\lambda_n + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j}} \right)^{-1} \\ &= \frac{1}{\lambda_n + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j}} - \frac{\sum_{j=\ell}^m q_j p^{\alpha_j}}{(\lambda_n + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j})^2} + O\left( \frac{(\sum_{j=\ell}^m q_j p^{\alpha_j})^2}{(\lambda_n + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j})^3} \right) \\ &= \frac{p^{-\alpha_1}}{\lambda_n p^{-\alpha_1} + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} - \frac{p^{\alpha_\ell - 2\alpha_1} (q_\ell + o(1))}{(q_1 + o(1))^2} + O(p^{2\alpha_\ell - 3\alpha_1}) \quad \text{for large } p > 0. \end{aligned} \quad (4.19)$$

For the denominator of the first term on the right-hand side, we further expand

$$\begin{aligned} \frac{1}{\lambda_n p^{-\alpha_1} + \sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} &= \frac{1}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} \left( 1 + \frac{\lambda_n p^{-\alpha_1}}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} \right)^{-1} \\ &= \frac{1}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} + O\left( \frac{\lambda_n p^{-\alpha_1}}{(\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1})^2} \right) \\ &= \frac{1}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} + \lambda_n O(p^{-\alpha_1}) \quad \text{for large } p > 0. \end{aligned}$$

Plugging the above expansion into (4.19), we obtain

$$\frac{1}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} = \frac{p^{-\alpha_1}}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} - \frac{p^{\alpha_\ell - 2\alpha_1} (q_\ell + o(1))}{(q_1 + o(1))^2} + \lambda_n O(p^{-2\alpha_1}) + O(p^{2\alpha_\ell - 3\alpha_1}) \quad (4.20)$$

for large  $p > 0$ . Treating the denominator in  $J_N^1(p)$  similarly and substituting them into  $p^{2\alpha_1 - \beta_\ell} (I_N^1(p) - J_N^1(p))$ , we deduce

$$\begin{aligned} p^{2\alpha_1 - \beta_\ell} (I_N^1(p) - J_N^1(p)) &= p^{\alpha_1 - \beta_\ell} \sum_{n=1}^N \lambda_n \left( \frac{a_{N,n}(\mathbf{x}_0)}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} - \frac{b_{N,n}(\mathbf{x}_0)}{\kappa^{-1} \sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} \right) \\ &\quad - p^{\alpha_\ell - \beta_\ell} (q_\ell + o(1)) \sum_{n=1}^N \frac{\lambda_n a_{N,n}(\mathbf{x}_0)}{(q_1 + o(1))^2} + (r_\ell + o(1)) \sum_{n=1}^N \frac{\lambda_n b_{N,n}(\mathbf{x}_0)}{(r_1 + o(1))^2} \end{aligned}$$

$$\begin{aligned}
 & + O(p^{-\beta_\ell}) \sum_{n=1}^N \lambda_n^2 (a_{N,n} - b_{N,n})(\mathbf{x}_0) + O(p^{2\alpha_\ell - \alpha_1 - \beta_\ell}) \sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0) \\
 & + O(p^{\beta_\ell - \alpha_1}) \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) \\
 = & \frac{p^{\alpha_1 - \beta_\ell} (1 + o(1))}{q_1} \sum_{n=1}^N \lambda_n (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) + \frac{r_\ell + o(1)}{r_1^2} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) \\
 & + o(1) \quad \text{for large } p > 0, \tag{4.21}
 \end{aligned}$$

where we utilized the assumption  $\alpha_1 < \beta_1$  and the uniform boundedness of all the involved sequences with respect to  $N$ .

Next we calculate  $p^{2\alpha_1 - \beta_\ell} (I_N^2(p) - J_N^2(p))$ . We further decompose  $I_N^2(p)$  as

$$\begin{aligned}
 I_N^2(p) & = \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \left\{ \frac{D_n a_{N,n}(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^2} + \sum_{k=2}^{d_n-1} \frac{(-1)^{k+1} (D_n a_{N,n})(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}} \right\} \\
 & = \sum_{n=1}^N \left\{ \sum_{j=1}^m q_j p^{\alpha_j} \frac{D_n a_{N,n}(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^2} + O(p^{-2\alpha_1}) \sum_{k=2}^{d_n-1} (-1)^{k+1} (D_n a_{N,n})(\mathbf{x}_0) \right\} \\
 & = \sum_{n=1}^N \left\{ \frac{D_n a_{N,n}(\mathbf{x}_0)}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} - \frac{\lambda_n (D_n a_{N,n})(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^2} \right\} + O(p^{-2\alpha_1}) \\
 & = \sum_{n=1}^N \frac{D_n a_{N,n}(\mathbf{x}_0)}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} + O(p^{-2\alpha_1}) \quad \text{for large } p > 0.
 \end{aligned}$$

Plugging the expansion (4.20) into the above equality and treating  $J_N^2(p)$  in the same manner, we deduce

$$\begin{aligned}
 p^{2\alpha_1 - \beta_\ell} (I_N^2(p) - J_N^2(p)) & = p^{\alpha_1 - \beta_\ell} \sum_{n=1}^N \left( \frac{D_n a_{N,n}(\mathbf{x}_0)}{\sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} - \frac{D_n b_{N,n}(\mathbf{x}_0)}{\kappa^{-1} \sum_{j=1}^{\ell-1} q_j p^{\alpha_j - \alpha_1}} \right) \\
 & \quad - p^{\alpha_\ell - \beta_\ell} (q_\ell + o(1)) \sum_{n=1}^N \frac{D_n a_{N,n}(\mathbf{x}_0)}{(q_1 + o(1))^2} + (r_\ell + o(1)) \sum_{n=1}^N \frac{D_n b_{N,n}(\mathbf{x}_0)}{(r_1 + o(1))^2} \\
 & \quad + O(p^{-\beta_\ell}) \sum_{n=1}^N \lambda_n D_n (a_{N,n} - b_{N,n})(\mathbf{x}_0) + O(p^{2\alpha_\ell - \alpha_1 - \beta_\ell}) \sum_{n=1}^N D_n a_{N,n}(\mathbf{x}_0) \\
 & \quad + O(p^{\beta_\ell - \alpha_1}) \sum_{n=1}^N D_n b_{N,n}(\mathbf{x}_0) \\
 = & \frac{p^{\alpha_1 - \beta_\ell} (1 + o(1))}{q_1} \sum_{n=1}^N D_n (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) + \frac{r_\ell + o(1)}{r_1^2} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) \\
 & + o(1) \quad \text{for large } p > 0. \tag{4.22}
 \end{aligned}$$

Collecting (4.21), (4.22), (4.9), (4.10) and using (4.11) again, we obtain

$$\begin{aligned}
 O(p^{-\nu + 2\alpha_1 - \beta_\ell}) & = p^{2\alpha_1 - \beta_\ell} \sum_{i=1}^3 (I_N^i(p) - J_N^i(p)) \\
 & = \frac{p^{\alpha_1 - \beta_\ell} (1 + o(1))}{q_1} \sum_{n=1}^N (\lambda_n + D_n) (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) + \frac{r_\ell + o(1)}{r_1^2} \sum_{n=1}^N (\lambda_n + D_n) b_{N,n}(\mathbf{x}_0) \\
 & \quad + o(1) + O(p^{\alpha_1 - \beta_\ell}) (\|a - a_N\|_{\mathcal{D}(L^\gamma)} + \|b - b_N\|_{\mathcal{D}(L^\gamma)})
 \end{aligned}$$

$$\begin{aligned}
 &= O(p^{\alpha_1 - \beta_\ell}) \left( \sum_{n=1}^N L(a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) + \|a - a_N\|_{\mathcal{D}(L\gamma)} + \|b - b_N\|_{\mathcal{D}(L\gamma)} \right) \\
 &\quad + \frac{r_\ell + o(1)}{r_1^2} \sum_{n=1}^N Lb_{N,n}(\mathbf{x}_0) + o(1) \\
 &= O(p^{\alpha_1 - \beta_\ell}) (L(a_N - \kappa b_N)(\mathbf{x}_0) + \|a - a_N\|_{\mathcal{D}(L\gamma)} + \|b - b_N\|_{\mathcal{D}(L\gamma)}) \\
 &\quad + \frac{r_\ell + o(1)}{r_1^2} Lb_N(\mathbf{x}_0) + o(1) \quad \text{for large } p > 0.
 \end{aligned}$$

By the assumption  $\nu > 2 \min\{\alpha_1, \beta_1\} = 2\alpha_1$ , we know  $-\nu + 2\alpha_1 - \beta_\ell < 0$  and thus

$$O(p^{\alpha_1 - \beta_\ell}) (L(a_N - \kappa b_N)(\mathbf{x}_0) + \|a - a_N\|_{\mathcal{D}(L\gamma)} + \|b - b_N\|_{\mathcal{D}(L\gamma)}) + \frac{r_\ell + o(1)}{r_1^2} Lb_N(\mathbf{x}_0) = o(1) \quad (4.23)$$

for large  $p > 0$ . By the definition of  $\kappa$ , we pass  $N \rightarrow \infty$  in (4.23) and employ (4.4)–(4.5) again to deduce

$$\frac{r_\ell + o(1)}{r_1^2} Lb(\mathbf{x}_0) = o(1) \quad \text{for large } p > 0.$$

Consequently, passing  $p \rightarrow \infty$  yields  $Lb(\mathbf{x}_0) = 0$ , which contradicts the assumption  $Lb(\mathbf{x}_0) \neq 0$ . In an identically parallel manner, we can exclude the possibility of  $\alpha_\ell > \beta_\ell$  and conclude  $\alpha_\ell = \beta_\ell$ .

Now it suffices to verify  $q_\ell/r_\ell = \kappa$ . Similarly to the argument at the end of Step 2, it follows immediately from  $\alpha_\ell = \beta_\ell$  that now (4.21) and (4.22) become

$$\begin{aligned}
 p^{2\alpha_1 - \beta_\ell} (I_N^1(p) - J_N^1(p)) &= \frac{p^{\alpha_1 - \beta_\ell} (1 + o(1))}{q_1} \sum_{n=1}^N \lambda_n (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) - \frac{q_\ell + o(1)}{q_1^2} \sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0) \\
 &\quad + \frac{r_\ell + o(1)}{r_1^2} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) + o(1), \\
 p^{2\alpha_1 - \beta_\ell} (I_N^2(p) - J_N^2(p)) &= \frac{p^{\alpha_1 - \beta_\ell} (1 + o(1))}{q_1} \sum_{n=1}^N D_n (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) - \frac{q_\ell + o(1)}{q_1^2} \sum_{n=1}^N \lambda_n a_{N,n}(\mathbf{x}_0) \\
 &\quad + \frac{r_\ell + o(1)}{r_1^2} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) + o(1).
 \end{aligned}$$

Repeating the same calculation as before, we arrive at

$$\begin{aligned}
 o(1) &= O(p^{\alpha_1 - \alpha_\ell}) (L(a_N - \kappa b_N)(\mathbf{x}_0) + \|a - a_N\|_{\mathcal{D}(L\gamma)} + \|b - b_N\|_{\mathcal{D}(L\gamma)}) \\
 &\quad - \frac{q_\ell + o(1)}{q_1^2} La_N(\mathbf{x}_0) + \frac{r_\ell + o(1)}{r_1^2} La_N(\mathbf{x}_0) \quad \text{for large } p > 0.
 \end{aligned}$$

Again passing first  $N \rightarrow \infty$  and then  $p \rightarrow \infty$ , we conclude

$$-\frac{q_\ell}{q_1^2} La(\mathbf{x}_0) + \frac{r_\ell}{r_1^2} Lb(\mathbf{x}_0) = 0$$

or equivalently  $q_\ell/r_\ell = \kappa$ . This completes the proof of (4.17) for  $j = \ell$ . By the inductive assumption, eventually (4.17) holds for all  $j = 1, 2, \dots, \min\{m, m'\}$ .

**Step 4** At last, it remains to show  $m = m'$  again by contradiction, that is, we assume  $m < m'$  without loss of generality. As those in the previous steps, our strategy is again multiplying both sides of (4.8) by  $p^{2\alpha_1 - \beta_{m+1}}$  to deduce a contradiction.

Notice that now  $J_N^1(p)$  and  $J_N^2(p)$  are represented by (4.18) with  $\ell = m + 1$ . Treating the denominators in  $I_N^1(p)$  and  $J_N^1(p)$  similarly as that in (4.20) yields

$$\frac{1}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} = \frac{p^{-\alpha_1}}{\sum_{j=1}^m q_j p^{\alpha_j - \alpha_1}} + \lambda_n O(p^{-2\alpha_1}),$$

$$\frac{1}{\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j}} = \frac{\kappa p^{-\alpha_1}}{\sum_{j=1}^m q_j p^{\alpha_j - \alpha_1}} - \frac{p^{2\alpha_1 - \beta_{m+1}}(r_{m+1} + o(1))}{(r_1 + o(1))^2} + \lambda_n O(p^{-2\alpha_1}) + O(p^{2\beta_{m+1} - 3\alpha_1})$$

for large  $p > 0$ . Then

$$\begin{aligned} p^{2\alpha_1 - \beta_{m+1}}(I_N^1(p) - J_N^1(p)) &= \frac{p^{\alpha_1 - \beta_{m+1}}(1 + o(1))}{q_1} \sum_{n=1}^N \lambda_n (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) \\ &\quad + \frac{r_{m+1} + o(1)}{r_1^2} \sum_{n=1}^N \lambda_n b_{N,n}(\mathbf{x}_0) + o(1) \quad \text{for large } p. \end{aligned} \quad (4.24)$$

Treating  $I_N^2(p)$  and  $J_N^2(p)$  similarly to that in Step 3 and expanding the involved denominators as above, we have

$$\begin{aligned} p^{2\alpha_1 - \beta_{m+1}}(I_N^2(p) - J_N^2(p)) &= \frac{p^{\alpha_1 - \beta_{m+1}}(1 + o(1))}{q_1} \sum_{n=1}^N D_n (a_{N,n} - \kappa b_{N,n})(\mathbf{x}_0) \\ &\quad + \frac{r_{m+1} + o(1)}{r_1^2} \sum_{n=1}^N D_n b_{N,n}(\mathbf{x}_0) + o(1) \quad \text{for large } p. \end{aligned} \quad (4.25)$$

Collecting (4.24), (4.25), (4.9), (4.10) and repeating the same calculation used in Step 3, we reach (4.23) with  $\ell = m + 1$ . Performing the same limiting process as before, we conclude again the contradiction  $Lb(\mathbf{x}_0) = 0$ . This completes the proof of  $m = m'$  and thus finalizes the proof of part (ii).  $\square$

Now we turn to the proof of Theorem 3.4, which relies on the following key lemma concerning the temporal component  $\rho(t)$  of the inhomogeneous terms in (1.3) and (3.1).

**Lemma 4.2.** *Let  $\rho \in L^1(\mathbb{R}_+)$  satisfy (3.8). Then its Laplace transform  $\widehat{\rho}(p)$  satisfies  $\widehat{\rho}(p) \sim p^{-\mu-1}$  for large  $p > 0$ .*

*Proof.* According to (3.8), there exist a small constant  $\varepsilon > 0$  and constants  $C_1, C_2 \in \mathbb{R}$  satisfying  $C_1 < C_2$  and  $C_1 C_2 > 0$  such that

$$C_1 t^\mu \leq \rho(t) \leq C_2 t^\mu \quad \text{a.e. } t \in (0, \varepsilon). \quad (4.26)$$

By the definition of the Laplace transform, we decompose  $\widehat{\rho}(p)$  as

$$\widehat{\rho}(p) = \left( \int_0^\varepsilon + \int_\varepsilon^\infty \right) e^{-pt} \rho(t) dt =: I_1(p) + I_2(p),$$

where

$$|I_2(p)| \leq \int_\varepsilon^\infty e^{-pt} |\rho(t)| dt \leq e^{-\varepsilon p} \|\rho\|_{L^1(\mathbb{R}_+)} = o(p^{-\mu-1}) \quad \text{for large } p > 0.$$

For  $I_1(p)$ , it follows from (4.26) and

$$\int_0^\varepsilon e^{-pt} t^\mu dt = \frac{1}{p^{\mu+1}} \int_0^{\varepsilon p} e^{-s} s^\mu ds \sim p^{-\mu-1} \quad \text{for large } p > 0$$

that  $I_1(p) \sim p^{-\mu-1}$  for large  $p > 0$ , which completes the proof.  $\square$

*Proof of Theorem 3.4.* Let  $a = b \equiv 0$  in  $\Omega$  in (1.3) and (3.1). Similarly to Step 1 in the proof of Theorem 3.1, it is not difficult to conclude that the difference between  $\widehat{u}(\mathbf{x}_0, p)$  and  $\widehat{v}(\mathbf{x}_0, p)$  still satisfies (4.1) for large  $p > 0$ .

Next, we take the Laplace transform in (1.3) and (3.1) and substitute  $\mathbf{x} = \mathbf{x}_0$  to deduce

$$\widehat{u}(\mathbf{x}_0, p) - \widehat{v}(\mathbf{x}_0, p) = \widehat{\rho}(p) (w(\mathbf{x}_0, p) - y(\mathbf{x}_0, p)) \quad \text{for large } p > 0,$$

where

$$w(\cdot, p) := \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} f, \quad y(\cdot, p) := \left( L + \sum_{j=1}^{m'} r_j p^{\beta_j} \right)^{-1} g. \quad (4.27)$$



Then we employ (4.1) and Lemma 4.2 to obtain

$$w(\mathbf{x}_0, p) - y(\mathbf{x}_0, p) = O(p^{\mu-\nu}) \quad \text{for large } p > 0. \quad (4.28)$$

Arguing similarly as before, we utilize Lemma 4.1 to pick two sequences  $f_N = \sum_{n=1}^N f_{N,n}$  and  $g_N = \sum_{n=1}^N g_{N,n}$  ( $N \in \mathbb{N}$ ) such that

$$f_{N,n}, g_{N,n} \in P_n L^2(\Omega), \quad \lim_{N \rightarrow \infty} \|L^\gamma(f - f_N)\|_{L^2(\Omega)} = \lim_{N \rightarrow \infty} \|L^\gamma(g - g_N)\|_{L^2(\Omega)} = 0.$$

Then for any fixed  $N \in \mathbb{N}$ , we decompose  $f = f_N + (f - f_N)$  and employ the expansion (2.3) to rewrite (4.27) as

$$\begin{aligned} w(\cdot, p) &= \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} \left\{ \sum_{n=1}^N f_{N,n} + (f - f_N) \right\} \\ &= \sum_{n=1}^N \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} P_n f_{N,n} + \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} (f - f_N) \\ &= \sum_{n=1}^N \frac{f_{N,n}}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}} + \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k D_n^k f_{N,n}}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}} + \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} (f - f_N) \end{aligned}$$

and similarly

$$y(\cdot, p) = \sum_{n=1}^N \frac{g_{N,n}}{\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j}} + \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k D_n^k g_{N,n}}{(\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j})^{k+1}} + \left( L + \sum_{j=1}^{m'} r_j p^{\beta_j} \right)^{-1} (g - g_N).$$

Then taking  $\mathbf{x} = \mathbf{x}_0$  in the above expressions and substituting them into (4.28) yield

$$\sum_{i=1}^3 (I_N^i(p) - J_N^i(p)) = O(p^{\mu-\nu}) \quad \text{for large } p > 0,$$

where

$$\begin{aligned} I_N^1(p) &:= \sum_{n=1}^N \frac{f_{N,n}(\mathbf{x}_0)}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}}, & J_N^1(p) &:= \sum_{n=1}^N \frac{g_{N,n}(\mathbf{x}_0)}{\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j}}, \\ I_N^2(p) &:= \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k (D_n^k f_{N,n})(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j})^{k+1}}, & J_N^2(p) &:= \sum_{j=1}^{m'} r_j p^{\beta_j} \sum_{n=1}^N \sum_{k=1}^{d_n-1} \frac{(-1)^k (D_n^k g_{N,n})(\mathbf{x}_0)}{(\lambda_n + \sum_{j=1}^{m'} r_j p^{\beta_j})^{k+1}}, \\ I_N^3(p) &:= \left( \left( L + \sum_{j=1}^m q_j p^{\alpha_j} \right)^{-1} (f - f_N) \right) (\mathbf{x}_0), & J_N^3(p) &:= \left( \left( L + \sum_{j=1}^{m'} r_j p^{\beta_j} \right)^{-1} (g - g_N) \right) (\mathbf{x}_0). \end{aligned}$$

The remaining part of the proof turns out to be essentially the same as that of Theorem 3.1 and we refrain from repeating the details.  $\square$

*Proof of Theorem 3.5.* We only provide the proof for part (1) because that of part (2) is mostly parallel. Then throughout this proof, we set  $f = g \equiv 0$  in  $\Omega$  in (1.3) and (3.1). Notice that thanks to the symmetry of  $L$ , we have (2.2) and

$$h = \sum_{n=1}^{\infty} P_n h, \quad \forall h \in L^2(\Omega), \quad (L - z)^{-1} P_n = \frac{P_n}{\lambda - z}, \quad z \notin \sigma(L). \quad (4.29)$$

First we assume (3.7) and show the “only if” part. According to Theorem 3.1, we have already reached (3.4). Together with (4.29), we can rewrite (4.2) and (4.3) as

$$\begin{aligned}\widehat{u}(\cdot, p) &= \frac{1}{p} \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^{\infty} \frac{P_n a}{\lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}}, \\ \widehat{v}(\cdot, p) &= \frac{1}{p} \sum_{j=1}^m q_j p^{\alpha_j} \sum_{n=1}^{\infty} \frac{P_n b}{\kappa \lambda_n + \sum_{j=1}^m q_j p^{\alpha_j}}\end{aligned}\quad \text{for large } p > 0. \quad (4.30)$$

On the other hand, owing to the time-analyticity of the solutions to (1.3) and (3.1), we see that (3.7) implies  $u(\mathbf{x}_0, \cdot) = v(\mathbf{x}_0, \cdot)$  in  $\mathbb{R}_+$ . Therefore, taking the Laplace transform results in  $\widehat{u}(\mathbf{x}_0, p) = \widehat{v}(\mathbf{x}_0, p)$  for large  $p > 0$  and hence

$$z \sum_{n=1}^{\infty} \frac{P_n a(\mathbf{x}_0)}{\lambda_n + z} = z \sum_{n=1}^{\infty} \frac{P_n b(\mathbf{x}_0)}{\kappa \lambda_n + z} \quad \text{for large } z, \quad z := \sum_{j=1}^m q_j p^{\alpha_j}.$$

Then we can analytically extend both sides of the above equality in  $z \in \mathbb{C}$  to obtain

$$\sum_{n=1}^{\infty} \frac{P_n a(\mathbf{x}_0)}{\lambda_n + z} = \sum_{n=1}^{\infty} \frac{P_n b(\mathbf{x}_0)}{\kappa \lambda_n + z}, \quad \sum_{n=1}^{\infty} \frac{P_n L a(\mathbf{x}_0)}{\lambda_n + z} = \sum_{n=1}^{\infty} \frac{\kappa P_n L b(\mathbf{x}_0)}{\kappa \lambda_n + z}, \quad z \in \mathbb{C} \setminus \Lambda, \quad (4.31)$$

where

$$\Lambda := (-\sigma(L)) \cup (-\kappa \sigma(L)), \quad \eta \sigma(L) := \{\eta \lambda_n \mid \lambda_n \in \sigma(L)\}, \quad \eta \in \mathbb{R}.$$

Here we used (3.5) and the fact that  $\lambda_n L = L P_n$ .

Now we concentrate on  $\kappa$  and consider the cases of  $\kappa = 1$  and  $\kappa \neq 1$  separately.

**Case 1** If  $\kappa = 1$ , then  $\Lambda = -\sigma(L)$  and for each  $n \in \mathbb{N}$ , we can take a sufficiently small circle  $\gamma_n$  enclosing  $-\lambda_n$  and excluding  $-\sigma(L) \setminus \{-\lambda_n\}$ . Integrating (4.31) on  $\gamma_n$  with respect to  $z$ , we can employ Cauchy’s integral theorem to conclude (3.14). In view of Remark 3.6, this completes the proof of (3.11) for  $\kappa = 1$ .

**Case 2** Now let us assume  $\kappa \neq 1$ . To obtain (3.9), it suffices to show  $\kappa \in \Sigma$  by contradiction. If  $\kappa \notin \Sigma$ , then it follows from the definition of  $\Sigma$  that  $\sigma(L) \cap \kappa \sigma(L) = \emptyset$ . Thus for each  $n \in \mathbb{N}$ , we can take the same circle  $\gamma_n$  as that in Case 1 enclosing  $-\lambda_n$  and excluding  $\Lambda \setminus \{-\lambda_n\}$ . Then integrating (4.31) on  $\gamma_n$  again yields  $P_n L a(\mathbf{x}_0) = 0$  ( $\forall n \in \mathbb{N}$ ) and hence

$$L a(\mathbf{x}_0) = \sum_{n=1}^{\infty} P_n L a(\mathbf{x}_0) = 0,$$

which contradicts the assumption  $L a(\mathbf{x}_0) \neq 0$ . This verifies  $\kappa \in \Sigma$  and also (3.9).

Now we turn to demonstrating (3.11) for  $\kappa \neq 1$ . Again by the definition of  $\Sigma$ , we know that  $\sigma(L) \cap \kappa \sigma(L) \neq \emptyset$  and the index sets  $M_\kappa, M'_\kappa$  defined by (3.10) are obviously nonempty. Meanwhile, by the strict monotonicity ordering of  $\sigma(L)$ , it is readily seen that for each  $n \in M_\kappa$ , there exists a unique  $n' \in M'_\kappa$  such that  $\lambda_n = \kappa \lambda_{n'}$  and vice versa. Then the bijection  $\theta_\kappa : M_\kappa \rightarrow M'_\kappa$  can be defined by  $\theta_\kappa(n) = n'$ .

Next, for each  $n \notin M_\kappa$ , we have  $\lambda_n \notin \kappa \sigma(L)$ . Then we can take the same circle  $\gamma_n$  as before and integrating (4.31) on  $\gamma_n$  to conclude

$$P_n a(\mathbf{x}_0) = P_n L a(\mathbf{x}_0) = 0, \quad \forall n \notin M_\kappa.$$

Similarly, we have

$$P_n b(\mathbf{x}_0) = P_n L b(\mathbf{x}_0) = 0, \quad \forall n \notin M'_\kappa.$$

Finally, for each  $n \in M_\kappa$ , we can take a circle  $\gamma_n$  enclosing  $\lambda_n = \kappa \lambda_{\theta_\kappa(n)}$  and excluding  $\Lambda \setminus \{-\lambda_n\}$ . Then again integrating (4.31) on  $\gamma_n$  gives

$$P_n a(\mathbf{x}_0) = P_{\theta_\kappa(n)} b(\mathbf{x}_0), \quad P_n L a(\mathbf{x}_0) = \kappa P_{\theta_\kappa(n)} L b(\mathbf{x}_0), \quad \forall n \in M_\kappa.$$

Collecting the above three cases completes the proof of (3.11) for  $\kappa \neq 1$ .

Now it remains to show the “if” part. Taking  $\boldsymbol{x} = \boldsymbol{x}_0$  in (4.30) and employing conditions (3.10)–(3.11), we obtain

$$\begin{aligned}\widehat{u}(\boldsymbol{x}_0, p) &= \frac{z}{p} \sum_{n=1}^{\infty} \frac{P_n a(\boldsymbol{x}_0)}{\lambda_n + z} = \frac{z}{p} \sum_{n \in M_\kappa} \frac{P_n a(\boldsymbol{x}_0)}{\lambda_n + z} = \frac{z}{p} \sum_{n \in M_\kappa} \frac{P_{\theta_\kappa(n)} b(\boldsymbol{x}_0)}{\lambda_n + z} \\ &= \frac{z}{p} \sum_{n \in M'_\kappa} \frac{P_n b(\boldsymbol{x}_0)}{\kappa \lambda_n + z} = \frac{z}{p} \sum_{n=1}^{\infty} \frac{P_n b(\boldsymbol{x}_0)}{\kappa \lambda_n + z} = \widehat{v}(\boldsymbol{x}_0, p) \quad \text{for large } p,\end{aligned}$$

where we abbreviated  $z := \sum_{j=1}^m q_j p^{\alpha_j}$ . Then the coincidence of  $u$  and  $v$  at  $\{\boldsymbol{x}_0\} \times \mathbb{R}_+$  follows immediately from the uniqueness of the Laplace transform.  $\square$

*Proof of Corollary 3.8.* Similarly to the proof of Theorem 3.5, we only deal with part (1) and omit the proof of part (2). Then it suffices to show the equivalence of (3.11) and (3.16).

By the assumption (3.15) on the choice of  $\boldsymbol{x}_0$ , there holds  $\varphi_n(\boldsymbol{x}_0) \neq 0$  for all  $n \in \mathbb{N}$ . On the other hand, since all eigenvalues of  $L$  are assumed to be simple, it reveals that

$$P_n h = (h, \varphi_n) \varphi_n, \quad P_n L h = \lambda_n (h, \varphi_n) \varphi_n, \quad \forall h \in \mathcal{D}(L).$$

Therefore, noticing that  $\lambda_n = \kappa \lambda_{\theta_\kappa(n)}$  for  $n \in M_\kappa$ , we immediately see that (3.11) is equivalent to

$$\begin{cases} (a, \varphi_n) \varphi_n(\boldsymbol{x}_0) = (b, \varphi_{\theta_\kappa(n)}) \varphi_{\theta_\kappa(n)}(\boldsymbol{x}_0), & n \in M_\kappa, \\ (a, \varphi_n) = 0, & n \notin M_\kappa, \\ (b, \varphi_n) = 0, & n \notin M'_\kappa. \end{cases}$$

Substituting the above relation into the Fourier expansions of  $a, b$ , we immediately obtain (3.16). The opposite side can also be verified similarly.

Especially if  $\kappa = 1$ , it turns out that (3.11) simply equals

$$(a, \varphi_n) = (b, \varphi_n), \quad \forall n \in \mathbb{N}$$

or equivalently  $a = b$  in  $\Omega$ . The proof of Corollary 3.8 is completed.  $\square$

## 5 Concluding Remarks

As information of solutions for our inverse problem, we assume only (3.2), which means the coincidence of principal terms of asymptotic expansions of solutions as  $t \downarrow 0$ . On the other hand, after taking Laplace transform of the solutions with respect to  $t$ , we only take advantage of the asymptotic behavior of  $\widehat{u}(\boldsymbol{x}_0, p) - \widehat{v}(\boldsymbol{x}_0, p)$  for large  $p > 0$ . This reflects the dominating effect of fractional orders and corresponding parameters to the solution. Simultaneously, it also suggests that the asymptotic behavior of solutions possesses abundant information about the equations.

The starting point of the proofs relies on the Laplace transforms of solutions, whose existence was guaranteed by the key estimate (2.5) in Lemma 2.1. Nevertheless, especially for  $\sigma(L) \subset \mathbb{R}_+$ , the estimate (2.5) is not at all sharp for large  $t$ . The sharp decay rate with a non-symmetric  $L$  keeps open even in the single-term case of (1.3), which deserves further investigation.

By reviewing the proof, we immediately see that our arguments also work for multi-term time-fractional wave equations, that is, (1.3) and (3.1) with

$$\begin{aligned}2 > \alpha_1 > \alpha_2 > \cdots > \alpha_m > 0, \quad 2 > \beta_1 > \beta_2 > \cdots > \beta_{m'} > 0, \\ \partial_t u = 0 \quad \text{if } \alpha_1 > 1, \quad \partial_t v = 0 \quad \text{if } \beta_1 > 1 \quad \text{in } \Omega \times \{0\}.\end{aligned}$$

However, since the forward theory for multi-term time-fractional wave equations (especially with a non-symmetric elliptic part) is not well established for the moment, we postpone this generalization as a future work. Meanwhile, other possible future topics include the case of  $\boldsymbol{x}$ -dependent parameters  $q_j(\boldsymbol{x}), r_j(\boldsymbol{x})$ .

Recently researches of wave equations with time-fractional damping terms, are developed. Such fractional systems and inverse problems are important also from physical points of view, and we can refer to Kaltenbacher and Rundell [13]. We can expect that our arguments may be available, but this should be a future work.

Our main purpose is the determination of scalar parameters  $m, \alpha_j, q_j$  in a multi-term time-fractional diffusion equation by some minimal information of solutions, as which we adopt the asymptotic behavior near  $t = 0$ . In general, it is difficult to prove the uniqueness in determining spatially varying initial data or source terms by such sparse data.

We can discuss weakened information of (3.2):

There exist a sequence  $\{t_m\}_{m=1}^{\infty} \subset \mathbb{R}_+$  and a constant  $C > 0$  such that

$$\lim_{m \rightarrow \infty} t_m = 0 \quad \text{and} \quad |u(\mathbf{x}_0, t_m) - v(\mathbf{x}_0, t_m)| \leq C t_m^{\tilde{\nu}}, \quad m \in \mathbb{N}, \quad (5.1)$$

where  $\tilde{\nu} > \min\{\alpha_1, \beta_1\}$ . From a practical viewpoint,  $\{t_m\}$  is a set of sampling times at a spatial monitoring point  $\mathbf{x}_0 \in \Omega$ . We can expect that (5.1) is equivalent to (3.2), so that replacing (3.2) by the weakened condition (5.1), we may obtain the same conclusion as in Theorem 3.1.

The proof in the general case is postponed to a future work, and here we sketch in the case  $m = 1$  with smooth  $a, b \in \mathcal{D}(L^\gamma)$  where  $\gamma > 0$  is sufficiently large. We assume that  $(La)(\mathbf{x}_0) \neq 0, (Lb)(\mathbf{x}_0) \neq 0$  and we write  $\alpha = \alpha_1, \beta = \beta_1$  for simplicity. Then (5.1) holds with some  $\tilde{\nu} > \min\{\alpha, \beta\}$  if and only if (3.2) holds with some  $\nu > \min\{\alpha, \beta\}$ .

*Proof.* It is trivial that (3.2) implies (5.1). Assuming (5.1), we see that

$$\begin{aligned} u(\mathbf{x}_0, t) &= \sum_{n=1}^{\infty} E_{\alpha,1} \left( -\frac{\lambda_n}{q_1} t^\alpha \right) (P_n a)(\mathbf{x}_0), \\ v(\mathbf{x}_0, t) &= \sum_{n=1}^{\infty} E_{\beta,1} \left( -\frac{\lambda_n}{r_1} t^\alpha \right) (P_n b)(\mathbf{x}_0), \end{aligned} \quad t > 0.$$

For  $a, b \in \mathcal{D}(L^\gamma)$  with large  $\gamma > 0$ , we can calculate

$$\begin{aligned} u(\mathbf{x}_0, t) &= \sum_{n=1}^{\infty} E_{\alpha,1} \left( -\frac{\lambda_n}{q_1} t^\alpha \right) (P_n a)(\mathbf{x}_0) \\ &= \sum_{n=1}^{\infty} \left\{ \left( 1 - \frac{\lambda_n t^\alpha}{q_1 \Gamma(\alpha + 1)} \right) P_n a(\mathbf{x}_0) + \sum_{k=2}^{\infty} \frac{(-\lambda_n t^\alpha)^k}{\Gamma(\alpha k + 1)} P_n a(\mathbf{x}_0) \right\} \\ &= \sum_{n=1}^{\infty} (P_n a)(\mathbf{x}_0) - \frac{t^\alpha}{q_1 \Gamma(\alpha + 1)} \sum_{n=1}^{\infty} \lambda_n (P_n a)(\mathbf{x}_0) \\ &\quad + \sum_{n=1}^{\infty} \sum_{j=0}^{\infty} \frac{1}{\Gamma(\alpha j + 2\alpha + 1)} \left( -\frac{\lambda_n}{q_1} t^\alpha \right)^2 \left( -\frac{\lambda_n}{q_1} t^\alpha \right)^j (P_n a)(\mathbf{x}_0) \\ &= a(\mathbf{x}_0) - \frac{t^\alpha}{q_1 \Gamma(\alpha + 1)} (La)(\mathbf{x}_0) + \frac{t^{2\alpha}}{q_1^2} \sum_{n=1}^{\infty} E_{\alpha,2\alpha+1} \left( -\frac{\lambda_n}{q_1} t^\alpha \right) P_n (L^2 a)(\mathbf{x}_0). \end{aligned}$$

Since  $a$  is sufficiently smooth, we can verify by the estimate of  $E_{\alpha,2\alpha+1}$  (e.g., [28, Theorem 1.6 (p.35)]) that

$$\sum_{n=1}^{\infty} \left| E_{\alpha,2\alpha+1} \left( -\frac{\lambda_n}{q_1} t^\alpha \right) \right| |P_n (L^2 a)(\mathbf{x}_0)| \leq \sum_{n=1}^{\infty} \frac{C}{1 + \frac{\lambda_n}{q_1} t^\alpha} |P_n (L^2 a)(\mathbf{x}_0)| < \infty.$$

Hence, we obtain

$$u(\mathbf{x}_0, t) = a(\mathbf{x}_0) - \frac{t^\alpha}{q_1 \Gamma(\alpha + 1)} (La)(\mathbf{x}_0) + O(t^{2\alpha}) \quad \text{as } t \downarrow 0. \quad (5.2)$$

Similarly, we can prove

$$v(\mathbf{x}_0, t) = b(\mathbf{x}_0) - \frac{t^\beta}{r_1 \Gamma(\beta + 1)} (Lb)(\mathbf{x}_0) + O(t^{2\beta}) \quad \text{as } t \downarrow 0. \quad (5.3)$$

Therefore, (5.1) yields

$$\left| a(\mathbf{x}_0) - b(\mathbf{x}_0) - \frac{(La)(\mathbf{x}_0)}{q_1\Gamma(\alpha+1)}t_m^\alpha + \frac{(Lb)(\mathbf{x}_0)}{r_1\Gamma(\beta+1)}t_m^\beta + O(t_m^{2\alpha}) + O(t_m^{2\beta}) \right| \leq C t_m^{\tilde{\nu}}, \quad m \in \mathbb{N}.$$

Here  $\tilde{\nu} > \min\{\alpha, \beta\}$ . Letting  $m \rightarrow \infty$ , we see (3.5). Assuming  $\alpha \leq \beta$  without loss of generality, we see  $\tilde{\nu} > \alpha$ . Dividing the above inequality by  $t_m^\alpha$ , we have

$$\left| -\frac{(La)(\mathbf{x}_0)}{q_1\Gamma(\alpha+1)} + \frac{(Lb)(\mathbf{x}_0)}{r_1\Gamma(\beta+1)}t_m^{\beta-\alpha} + O(t_m^\alpha) + O(t_m^{2\beta-\alpha}) \right| \leq C t_m^{\tilde{\nu}-\alpha}, \quad m \in \mathbb{N}. \quad (5.4)$$

Letting  $m \rightarrow \infty$ , since  $O(t_m^{2\beta-\alpha}) = o(1)$  and  $t_m^{\tilde{\nu}-\alpha} = o(1)$  as  $m \rightarrow \infty$ , we reach

$$\frac{(La)(\mathbf{x}_0)}{q_1\Gamma(\alpha+1)} = \lim_{m \rightarrow \infty} t_m^{\beta-\alpha} \left( \frac{(Lb)(\mathbf{x}_0)}{r_1\Gamma(\beta+1)} \right).$$

By the assumption  $(La)(\mathbf{x}_0) \neq 0$  and  $(Lb)(\mathbf{x}_0) \neq 0$ , we obtain

$$\lim_{m \rightarrow \infty} t_m^{\beta-\alpha} \left( \frac{(Lb)(\mathbf{x}_0)}{r_1\Gamma(\beta+1)} \right) \neq 0,$$

that is,  $\lim_{m \rightarrow \infty} t_m^{\beta-\alpha} \neq 0$ , which implies  $\beta = \alpha$ . Letting  $m \rightarrow \infty$  in (5.4) with  $\alpha = \beta$ , we see that

$$\frac{(Lb)(\mathbf{x}_0)}{r_1\Gamma(\beta+1)} = \frac{(La)(\mathbf{x}_0)}{q_1\Gamma(\alpha+1)}.$$

Therefore, by (5.2)–(5.4), we obtain

$$|u(\mathbf{x}_0, t) - v(\mathbf{x}_0, t)| = O(t^{2\alpha}) \quad \text{as } t \downarrow 0.$$

Hence, (3.2) is satisfied with  $\nu := 2\alpha > \min\{\alpha, \beta\}$ . Thus (3.2) and (5.1) provide equivalent information.  $\square$

Although the single point observation is sufficient for the uniqueness of Problem 1.1 theoretically, it seems meaningful to develop numerical methods with multiple observation points in order to improve the accuracy for the reconstruction. If the corresponding rank condition (see [30]) is also satisfied in this case, it is even possible to uniquely identify the initial value  $a$  or the spatial component  $f$  of the source term in (1.3), whose numerical reconstruction may also be interesting.

Finally, for Problem 1.1, we can adopt information of the asymptotic data as  $t \rightarrow \infty$ , but we postpone details to another future work.

## Acknowledgments

Y.Liu is supported by Grant-in-Aid for Early-Career Scientists 20K14355 and 22K13954, Japan Society for the Promotion of Science (JSPS). M. Yamamoto is supported by Grant-in-Aid for Scientific Research (A) 20H00117 and Grant-in-Aid for Challenging Research (Pioneering) 21K18142, JSPS.

## References

- [1] R. A. Adams, Sobolev Spaces, Academic Press, New York, 1975.
- [2] S. Agmon, Lectures on Elliptic Boundary Value Problems, AMS, Providence, 1965.
- [3] R. Ashurov and S. Umarov, Determination of the order of fractional derivative for subdiffusion equations, *Fract. Calc. Appl. Anal.*, **23** (2020), 1647–1662.

- 
- [4] M. T. Barlow and E. A. Perkins, Brownian motion on the Sierpinski gasket, *Probab. Theory Related Fields*, **79** (1988), 543–623.
- [5] S. Beckers and M. Yamamoto, Regularity and unique existence of solution to linear diffusion equation with multiple time-fractional derivatives, in: Bredies, K. et al. (eds.) *Control and Optimization with PDE Constraints*, Springer, Basel, 2013, 45–55.
- [6] J. Cheng, J. Nakagawa, M. Yamamoto and T. Yamazaki, Uniqueness in an inverse problem for a one dimensional fractional diffusion equation, *Inverse Problems*, **25** (2009), 115002.
- [7] R. Gorenflo, Y. Luchko and M. Yamamoto, Time-fractional diffusion equation in the fractional Sobolev spaces, *Fract. Calc. Appl. Anal.*, **18** (2015), 799–820.
- [8] Y. Hatano and N. Hatano, Dispersive transport of ions in column experiments: an explanation of long-tailed profiles, *Water Resour. Res.*, **34** (1998), 1027–1033.
- [9] Y. Hatano, J. Nakagawa, S. Wang, and M. Yamamoto, Determination of order in fractional diffusion equation, *J. Math-for-Ind.*, **5A** (2013), 51–57.
- [10] D. Jiang, Z. Li, Y. Liu and M. Yamamoto, Weak unique continuation property and a related inverse source problem for time-fractional diffusion-advection equations, *Inverse Problems*, **33** (2017), 055013.
- [11] B. Jin and Y. Kian, Recovery of the order of derivation for fractional diffusion equations in an unknown medium, *SIAM J. Appl. Math.*, **82** (2022), 1045–1067.
- [12] B. Jin and Y. Kian, Recovering multiple fractional orders in time-fractional diffusion in an unknown medium, *Proc. R. Soc. A*, **477** (2021), 20210468.
- [13] B. Kaltenbacher and W. Rundell, Determining damping terms in fractional wave equations, *Inverse Problems*, **38** (2022), 075004.
- [14] T. Kato, *Perturbation Theory for Linear Operators* (Second Edition), Springer-Verlag, Berlin, 1976.
- [15] Y. Kian, Z. Li, Y. Liu and M. Yamamoto, The uniqueness of inverse problems for a fractional equation with a single measurement, *Math. Ann.*, **380** (2021), 1465–1495.
- [16] A. Kubica, K. Ryszewska and M. Yamamoto, *Time-Fractional Differential Equations: A Theoretical Introduction*, Springer, Singapore, 2020.
- [17] Z. Li, X. Huang and M. Yamamoto, Initial-boundary value problems for multi-term time-fractional diffusion equations with  $x$ -dependent coefficients, *Evol. Equ. Control Theory*, **9** (2020), 153–179.
- [18] Z. Li, X. Huang and M. Yamamoto, A stability result for the determination of order in time-fractional diffusion equations, *J. Inverse Ill-Posed Probl.*, **28** (2020), 379–388.
- [19] Z. Li, O. Y. Imanuvilov and M. Yamamoto, Uniqueness in inverse boundary value problems for fractional diffusion equations, *Inverse Problems*, **32** (2016), 015004.
- [20] Z. Li, Y. Liu and M. Yamamoto, Initial-boundary value problems for multi-term time-fractional diffusion equations with positive constant coefficients, *Appl. Math. Comput.*, **257** (2015), 381–397.
- [21] Z. Li, Y. Liu and M. Yamamoto, Inverse problems of determining parameters of the fractional partial differential equations, in: Kochubei, A., Luchko, Y. (eds.) *Handbook of Fractional Calculus with Applications, Volume 2: Fractional Differential Equations*, De Gruyter, Berlin, 2019, 431–442.
- [22] Z. Li and M. Yamamoto, Uniqueness for inverse problems of determining orders of multi-term time-fractional derivatives of diffusion equation, *Appl. Anal.*, **94** (2015), 570–579.
- [23] Z. Li and Z. Zhang, Unique determination of fractional order and source term in a fractional diffusion equation from sparse boundary data, *Inverse Problems*, **36** (2020), 115013.

- [24] K. Liao and T. Wei, Identifying a fractional order and a space source term in a time-fractional diffusion-wave equation simultaneously, *Inverse Problems*, **35** (2019), 115002.
- [25] Y. Liu, Strong maximum principle for multi-term time-fractional diffusion equations and its application to an inverse source problem, *Comput. Math. Appl.*, **73** (2017), 96–108.
- [26] Y. Luchko, Initial-boundary-value problems for the generalized multi-term time-fractional diffusion equation, *J. Math. Anal. Appl.*, **374** (2011), 538–548.
- [27] Y. Luchko and M. Yamamoto, Maximum principle for the time-fractional PDEs, in: Kochubei, A., Luchko, Y. (eds.) *Handbook of Fractional Calculus with Applications, Volume 2: Fractional Differential Equations*, De Gruyter, Berlin, 2019, 299–326.
- [28] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [29] K. Sakamoto and M. Yamamoto, Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems, *J. Math. Anal. Appl.*, **382** (2011), 426–447.
- [30] Y. Sakawa, Observability and related problems for partial differential equations of parabolic type, *SIAM J. Control*, **13**, 1975, 14–27.
- [31] L. L. Sun, Y. S. Li and Y. Zhang, Simultaneous inversion of the potential term and the fractional orders in a multi-term time-fractional diffusion equation, *Inverse Problems*, **37** (2021), 055007.
- [32] H. Tanabe, *Equations of Evolution*, Pitman, London, 1975.
- [33] M. Yamamoto, Uniqueness in determining the orders of time and spatial fractional derivatives, preprint, arXiv:2006.15046.
- [34] M. Yamamoto, Uniqueness for inverse problem of determining fractional orders for time-fractional advection-diffusion equations, *Math. Control Relat. Fields*, accepted, doi: 10.3934/mcrf.2022017.
- [35] M. Yamamoto, Uniqueness in determining fractional orders of derivatives and initial values, *Inverse Problems*, **37** (2021), 095006.
- [36] X. B. Yan, Y. X. Zhang and T. Wei, Identify the fractional order and diffusion coefficient in a fractional diffusion wave equation, *J. Comput. Appl. Math.*, **393** (2021), 113497.