

The Order of the Unitary Subgroups of Group Algebras

Zsolt Adam Balogh

*Department of Mathematical Sciences
United Arab Emirates University, Al Ain,
United Arab Emirates, P.O.Box: 15551*

Abstract

Let FG be the group algebra of a finite p -group G over a finite field F of positive characteristic p . Let \ast be an involution of the algebra FG which is a linear extension of an anti-automorphism of the group G to FG . If p is an odd prime, then the order of the \ast -unitary subgroup of FG is established. For the case $p = 2$ we generalize a result obtained for finite abelian 2-groups. It is proved that the order of the \ast -unitary subgroup of FG of a non-abelian 2-group is always divisible by a number which depends only on the size of F , the order of G and the number of elements of order two in G . Moreover, we show that the order of the \ast -unitary subgroup of FG determines the order of the finite p -group G .

Keywords: group algebras; unit group of group algebras; unitary subgroups

1. Introduction and main results

Let FG be the group algebra of a finite p -group G over a finite field F of positive characteristic p . Let

$$V(FG) = \left\{ x = \sum_{g \in G} \alpha_g g \in FG \mid \chi(x) = \sum_{g \in G} \alpha_g = 1 \right\}$$

be the group of normalized units of FG , where $\chi(x)$ is the augmentation map (see [5, Chapters 2-3, p. 194-196]). In this case, the order of the group $V(FG)$ is equal to $|F|^{|G|-1}$, so the order of $V(FG)$ can be very large even for

Email address: baloghza@gmail.com (Zsolt Adam Balogh)

a small group G . Note that, studying the structure of the group $V(FG)$ is a rather difficult task (for more details see the survey [5]).

Let \otimes be an involution of the algebra FG . We say that the involution \otimes arises from the group G , if \otimes is a linear extension of an anti-automorphism of G to FG . An example for such kind of involution is the canonical involution that is the linear extension of the anti-automorphism of G which sends each element of G to its inverse. This involution is usually denoted by $*$.

An element $u \in V(FG)$ is called \otimes -unitary, if $u^\otimes = u^{-1}$ with respect to the involution \otimes of FG . The set $V_\otimes(FG)$ of all \otimes -unitary units forms a subgroup of $V(FG)$ which is called \otimes -unitary subgroup. Interest in the structure of unitary subgroups arose in algebraic topology and unitary K -theory (see Novikov's papers [17] and Bovdi's paper [8]). Let L be a finite Galois extension of F with Galois group G , where F is a finite field of characteristic two. Serre [18] identified an interesting relation between the self-dual normal basis of L over F and the $*$ -unitary subgroup of FG . This relationship also makes the study of the unitary subgroups timely.

The unitary subgroups have been proven to be very useful subgroups in several studies (see [2, 3, 4, 6, 7, 10, 11, 13, 14, 15] and [16]). However, we know very little about their structure, as even finding their order is a challenging problem. The first results in this area were published in the 1980's. For finite abelian p -groups G the order of $V_*(FG)$ was given in [9].

Proposition 1. ([9, Theorem 2]) *Let G be a finite abelian 2-group. If F is a finite field of characteristic 2, then the order $|V_*(FG)|$ is divisible by $|F|^{\frac{1}{2}(|G|+|G\{2\}|)-1}$, that is,*

$$|V_*(FG)| = \Theta \cdot |F|^{\frac{1}{2}(|G|+|G\{2\}|)-1},$$

where $\Theta = |G^2\{2\}|$ and $|S|$ denotes the size of a finite set S .

It follows that the number Θ does not depend on the size of the field F . The following breakthrough result was proved for certain non-abelian 2-groups by Bovdi and Roza.

Proposition 2. ([12, Corollary 2]) *If $|F| = 2^m \geq 2$, then:*

$$|V_*(FG)| = \Theta \cdot |F|^{\frac{1}{2}(|G|+|G\{2\}|)-1},$$

where

- (i) $\Theta = 1$ if G is a dihedral 2-group;
- (ii) $\Theta = 4$ if G is a generalized quaternion 2-group.

In [1], the value of the number Θ was given for all non-abelian groups of order 2^4 . Although, for these groups, the number Θ is not equal to $|G^2\{2\}|$ it does not depend on the field F . Wang and Liu [19] evaluated Θ in the case when G is a non-abelian 2-group given by a central extension of the form

$$1 \longrightarrow C_{2^m} \longrightarrow G \longrightarrow C_2 \times \cdots \times C_2 \longrightarrow 1,$$

in which $m \geq 1$ and $|G'| = 2$. At present, the question of whether the quotient Θ depends on the field F is still open.

Our main results are the following.

Theorem 1. *Let G be a finite p -group, where p is an odd prime and let F be a finite field of characteristic p . If \otimes is an involution of FG that arises from the group G , then*

$$|V_{\otimes}(FG)| = |F|^{\frac{1}{2}(|G| - |G_{\otimes}|)},$$

where $G_{\otimes} = \{g \mid g = g^{\otimes}\}$.

Let $\xi(G)$ denote the center of the group G and $\xi(G)\{2\}$ denote the set of elements of order two in $\xi(G)$.

Theorem 2. *Let G be a finite 2-group. If F is a finite field of characteristic two, then*

$$|V_*(FG)| = \Theta \cdot |F|^{\frac{1}{2}(|G| + |G\{2\}|) - 1}$$

for some integer Θ . Moreover, if the set $T_c = \{g \in G \mid g^2 = c\}$ is commutative for some $c \in \xi(G)\{2\}$, then Θ does not depend on the field F .

By combining Theorem 1 and Theorem 2, we have the following.

Corollary 1. *Let G be a finite p -group. If F is a finite field of characteristic p , then the order of the $*$ -unitary subgroup of FG determines the order of G .*

2. Proofs

Let G be a finite p -group, let F be a finite field of $\text{char}(F) = p > 2$ and let \otimes be an involution of FG which arises from G . An element $x \in FG$ is called skew-symmetric under the involution \otimes if $x^\otimes = -x$. Let FG_\otimes^- denote the set of all skew-symmetric elements of FG .

Proof of Theorem 1. Let $z \in FG$ such that $1 + z$ is invertible. Clearly, $1 - z$ and $1 + z$ commute, therefore $1 - z$ and $(1 + z)^{-1}$ also commute.

Let $Q = \{x \in FG \mid 1 + x \text{ is invertible in } FG\}$. Let us define the map $f : Q \rightarrow FG$ by

$$f(x) = (1 - x)(1 + x)^{-1}.$$

If $y \in FG_\otimes^-$, then $\chi(y) = 0$, so

$$\chi(1 + y) = \chi(1 - y) = \chi(1 + y^\otimes) = 1$$

and $1 + y, 1 - y, 1 + y^\otimes$ are normalized units. Hence

$$\begin{aligned} f(y)f(y)^\otimes &= (1 - y)(1 + y)^{-1}(1 + y^\otimes)^{-1}(1 - y^\otimes) \\ &= (1 - y)(1 + y)^{-1}(1 - y)^{-1}(1 + y) \\ &= (1 + y)^{-1}(1 - y)(1 - y)^{-1}(1 + y) \\ &= 1. \end{aligned}$$

Consequently, $f(y) \in V_\otimes(FG)$ and $f : FG_\otimes^- \rightarrow V_\otimes(FG)$ is a surjection.

Let $x \in V_\otimes(FG)$. Evidently, $1 + x, 1 + x^\otimes$ and $1 + x^{-1}$ are invertible, because $\chi(1 + x) = \chi(1 + x^\otimes) = \chi(1 + x^{-1}) = 2$ is invertible in F . Therefore $V_\otimes(FG)$ is a subset of Q . Let y denote the element $f(x)$. Then

$$\begin{aligned} y^\otimes &= f(x)^\otimes = (1 + x^\otimes)^{-1}(1 - x^\otimes) = (1 + x^{-1})^{-1}(1 - x^{-1}) \\ &= (x^{-1}(x + 1))^{-1}x^{-1}(x - 1) = -(1 + x)^{-1}xx^{-1}(1 - x) \\ &= -y. \end{aligned}$$

Therefore $f : V_\otimes(FG) \rightarrow FG_\otimes^-$ is a surjection.

Similar computation shows that

$$\begin{aligned} f(f(x)) &= (1 - (1 - x)(1 + x)^{-1})(1 + (1 - x)(1 + x)^{-1})^{-1} \\ &= ((1 + x) - (1 - x))(1 + x)^{-1} \left(((1 + x) + (1 - x))(1 + x)^{-1} \right)^{-1} \\ &= ((1 + x) - (1 - x))(1 + x)^{-1}(1 + x) \left((1 + x) + (1 - x) \right)^{-1} \\ &= ((1 + x) - (1 - x)) \left((1 + x) + (1 - x) \right)^{-1} \\ &= x \end{aligned}$$

for every $x \in V_{\otimes}(FG)$, so f is a bijection between FG_{\otimes}^- and $V_{\otimes}(FG)$.

Since FG_{\otimes}^- is a linear space over F with basis $\{g - g^{\otimes} \mid g \in G \setminus G_{\otimes}\}$,

$$|V_{\otimes}(FG)| = |FG_{\otimes}^-| = |F|^{\frac{1}{2}(|G| - |G_{\otimes}|)}.$$

□

Let H be a normal subgroup of G and let $I(H) := \langle 1 + h \mid h \in H \rangle_{FG}$ be an ideal of FG generated by the set $\{1 + h \mid h \in H\}$. Clearly,

$$FG/I(H) \cong FG/\ker(\Psi) \cong F[G/H],$$

where $\Psi : FG \rightarrow FG/I(H)$ is the natural homomorphism.

Let us denote by $V_*(F\overline{G})$ the $*$ -unitary subgroup of the factor algebra $FG/I(H)$, where $\overline{G} = G/H$. It is easy to check that the set

$$N_{\Psi}^* = \{x \in V(FG) \mid \Psi(x) \in V_*(F\overline{G})\}$$

forms a subgroup in $V(FG)$. Let $I(H)^+ = \{1 + x \mid x \in I(H)\}$. The subgroup $I(H)^+$ is normal in $V(FG)$ and $S_H = \{xx^* \mid x \in N_{\Psi}^*\}$ is a subset of $I(H)^+$, because $xx^* \in 1 + \ker(\Psi) = I(H)^+$ for all $x \in N_{\Psi}^*$.

First, we need the following.

Lemma 1. *Let H be a normal subgroup of a finite 2-group G . Set $\overline{G} = G/H$. If $|F| = 2^m \geq 2$, then*

$$|V_*(FG)| = |F|^{|\overline{G}|} \cdot \frac{|V_*(F\overline{G})|}{|S_H|}. \quad (1)$$

Proof. Let $\Phi : V(FG) \rightarrow V(FG)$ be a map such that $\Phi(x) = xx^*$ for every $x \in V(FG)$. The sets $\Phi(x)$ and $\Phi(y)$ coincide if and only if $y \in x \cdot V_*(FG)$. Indeed, if $y \in x \cdot V_*(FG)$, then $y = xv$ for some $v \in V_*(FG)$. Therefore

$$\Phi(y) = yy^* = xv(xv)^* = xvv^*x^* = xx^* = \Phi(x).$$

Assume that $\Phi(x) = \Phi(y)$ for some $x, y \in V(FG)$. Then $xx^* = yy^*$, or equivalently, $y^{-1}x = y^*(x^*)^{-1}$. Therefore

$$(x^{-1}y)^{-1} = y^{-1}x = y^*(x^*)^{-1} = (x^{-1}y)^*$$

which confirms that $x^{-1}y \in V_*(FG)$.

Since $[N_{\Psi}^* : V_*(FG)] = |S_H|$,

$$|V_*(FG)| = \frac{|N_{\Psi}^*|}{|S_H|} = |I(H)^+| \cdot \frac{|V_*(F\overline{G})|}{|S_H|}.$$

We should note that $V_*(FG)$ is usually not a normal subgroup of N_{Ψ}^* .

The ideal $I(H)$ can be considered as a vector space over F with the following basis $\{u(1+h) \mid u \in T(G/H), h \in H\}$, where $T(G/H)$ is a complete set of left coset representatives of H in G . Consequently,

$$|I(H)^+| = |I(H)| = |F|^{\frac{|G|}{|H|}}.$$

□

Proof of Theorem 2. Let G be a 2-group of order 2^n and let H be a subgroup of G generated by a central element c of order two. Evidently, the set $S_H = \{xx^* \mid x \in N_{\Psi}^*\}$ is a subset of $I(H)^+ \cap V_*(FG)$ and every $y \in S_H$ is a *-symmetric element. Moreover, the support of y does not contain elements of order two by [1, Lemma 2.5]. Thus, every $y \in S_H$ can be written as

$$y = 1 + \sum_{g \in G \setminus (G\{2\} \cup T_c)} \alpha_g(g + g^{-1})\widehat{H} + \sum_{g \in T_c} \beta_g g \widehat{H}, \quad (\alpha_g, \beta_g \in F)$$

where $T_c = \{g \in G \mid g^2 = c\}$ and $\widehat{H} = 1 + c$. This yields that

$$|S_H| \leq |F|^{\frac{1}{4}(|G| - |G\{2\}| - |T_c|) + \frac{1}{2}|T_c|} = |F|^{\frac{1}{4}(|G| - |G\{2\}| + |T_c|)}. \quad (2)$$

Let us prove that if T_c is a commutative set, then

$$|F|^{\frac{1}{4}(|G| - |G\{2\}| + |T_c|)} \cdot 2^{-\frac{1}{2}|T_c|} \leq |S_H|. \quad (3)$$

Let N_1 be a group generated by the elements $1 + \alpha_g(g + g^{-1})\widehat{H}$, in which $g^2 \notin H$ and $\alpha_g \in F$. Evidently, N_1 is an elementary abelian subgroup of $I(H)^+$. Since $g^2 \notin H$ (equivalently $g \in G \setminus (G\{2\} \cup T_c)$),

$$1 + \alpha_g(g + g^{-1})\widehat{H} = 1 + \alpha_g g^{-1}(1 + g^2)\widehat{H} \neq 1$$

and

$$1 + \alpha_g(g + g^{-1})\widehat{H} = (1 + \alpha_g g \widehat{H})(1 + \alpha_g g \widehat{H})^* \in S_H.$$

If $z \in N_1$, then

$$z = \prod_{g^2 \notin H} (1 + \alpha_g(g + g^{-1})\widehat{H}) = \prod_{g^2 \notin H} (1 + \alpha_g g \widehat{H})(1 + \alpha_g g \widehat{H})^*,$$

so $(1 + \alpha_g g \widehat{H}) \in I(H)^+$ and

$$z = \left(\prod_{g^2 \notin H} (1 + \alpha_g g \widehat{H}) \right) \cdot \left(\prod_{g^2 \notin H} (1 + \alpha_g g \widehat{H}) \right)^* \in S_H.$$

Thus N_1 is a subgroup in the set S_H and $|N_1| = |F|^{\frac{1}{4}(|G| - |G\{2\}| - |T_c|)}$.

The map $\tau : F \rightarrow F$ defined by $\tau(\alpha) = \alpha + \alpha^2$ ($\alpha \in F$) is a homomorphism on the additive group of the field F with kernel $\ker(\tau) = \{0, 1\}$ (see [1, Lemma 10]). Therefore $|\text{im}(\tau)| = \frac{|F|}{2}$.

Suppose that T_c is a commutative set. Let N_2 be a group generated by the elements $1 + \alpha_g g \widehat{H}$, in which $g \in T_c$ and $\alpha_g \in F$. Since

$$(1 + \omega g + \omega g^2)(1 + \omega g + \omega g^2)^* = 1 + (\omega + \omega^2)g \widehat{H}$$

we have $1 + \alpha_g g \widehat{H} \in S_H$ for every $\alpha_g \in \text{im}(\tau)$. The group N_2 , being T_c commutative, is a subgroup in S_H and

$$|N_2| = |\text{im}(\tau)|^{\frac{1}{2}|T_c|} = |F|^{\frac{1}{2}|T_c|} \cdot 2^{-\frac{1}{2}|T_c|}.$$

Let $x \in N_1$ and $y \in N_2$. There exist $x_1 \in I(H)^+$ and $y_1 \in N_{\Psi}^*$ such that $x_1 x_1^* = x$ and $y_1 y_1^* = y$. Since $I(H)^+$ is an elementary 2-group, the element y commutes with x_1 and

$$yx = yx_1 x_1^* = x_1 y x_1^* = x_1 y_1 y_1^* x_1^* = (x_1 y_1)(x_1 y_1)^* \in S_H.$$

Therefore $N_1 \times N_2$ is a subgroup in S_H and

$$|N_1 \times N_2| = |F|^{\frac{1}{4}(|G| - |G\{2\}| - |T_c|) + \frac{1}{2}|T_c|} \cdot 2^{-\frac{1}{2}|T_c|}.$$

Consequently,

$$|F|^{\frac{1}{4}(|G| - |G\{2\}| + |T_c|)} \cdot 2^{-\frac{1}{2}|T_c|} \leq |S_H|.$$

Now, we are ready to prove the theorem. If $n = 3$, then the theorem is true by Propositions 1 and 2. Suppose that $n > 3$. In the factor group $\overline{G} = G/H$ the element \overline{g} has order two if and only if either $g \in G\{2\}$ or $g \in T_c$. Therefore, $|\overline{G}\{2\}| = \frac{|G\{2\}| + |T_c|}{2}$. According to the inductive hypothesis

$$|V_*(F\overline{G})| = 2^s \cdot |F|^{\frac{1}{2}(|\overline{G}| + |\overline{G}\{2\}|) - 1} = 2^s \cdot |F|^{\frac{1}{4}(|G| + |G\{2\}| + |T_c|) - 1}$$

for some $s \geq 0$. Using Lemma 1 and equation (2) we obtain that

$$\begin{aligned} |V_*(FG)| &= |F|^{\frac{|G|}{2}} \cdot \frac{|V_*(F\bar{G})|}{|S_H|} \\ &\geq |F|^{\frac{|G|}{2}} \cdot \frac{2^s \cdot |F|^{\frac{1}{4}(|G|+|G\{2\}|+|T_c|)-1}}{|F|^{\frac{1}{4}(|G|-|G\{2\}|+|T_c|)}} \\ &= 2^s \cdot |F|^{\frac{1}{2}(|G|+|G\{2\}|-1)}. \end{aligned}$$

Therefore

$$2^s \cdot |F|^{\frac{1}{2}(|G|+|G\{2\}|-1)} \leq |V_*(FG)| \quad (4)$$

and $|V_*(FG)|$ is divisible by $|F|^{\frac{1}{2}(|G|+|G\{2\}|-1)}$.

Similarly, using Lemma 1 and (3), we obtain that

$$\begin{aligned} |V_*(FG)| &= |F|^{\frac{|G|}{2}} \cdot \frac{|V_*(F\bar{G})|}{|S_H|} \\ &\leq |F|^{\frac{|G|}{2}} \cdot \frac{2^s \cdot |F|^{\frac{1}{4}(|G|+|G\{2\}|+|T_c|)-1}}{|F|^{\frac{1}{4}(|G|-|G\{2\}|+|T_c|), 2^{-\frac{1}{2}|T_c|}}} \\ &= 2^{s+\frac{1}{2}|T_c|} \cdot |F|^{\frac{1}{2}(|G|+|G\{2\}|-1)}. \end{aligned}$$

The size of the set T_c does not depend on the field F . Since 2^s does not depend on the field F by the inductive hypothesis, the proof is complete. \square

We should remark that 2^s in the inequality (4) is usually not inherited via the factorization. For example, for the semidihedral group D_{16}^- of order 16 we have that

$$|V_*(FD_{16}^-)| = 2 \cdot |F|^{\frac{1}{2}(|D_{16}^-|+|D_{16}^-\{2\}|-1)}$$

by [1, Lemma 3.4]. However, $D_8 \cong D_{16}^-/H$, where H is the center of D_{16}^- and D_8 is the dihedral group of order 8 and $|V_*(FD_8)| = |F|^{\frac{1}{2}(|D_8|+|D_8\{2\}|-1)}$ by Proposition 2.

Proof of Corollary 1. If $p = 2$, then Theorem 2 implies that

$$|F|^{\frac{|G|}{2}-1} \leq |F|^{\frac{1}{2}(|G|+|G\{2\}|-1)} \leq |V_*(FG)| \leq |F|^{|G|-1}.$$

If p is an odd prime, then $|V_*(FG)| = |F|^{\frac{1}{2}(|G|-1)}$ by Theorem 1. Hence, $|F|^{\frac{|G|}{2}-1} \leq |V_*(FG)| \leq |F|^{|G|-1}$, which confirms that the order of $V_*(FG)$ determines the order of G for every finite p -groups. \square

References

- [1] Z. Balogh. On unitary subgroups of group algebras. *Int. Electron. J. Algebra*, 29:187–198, 2021.
- [2] Z. Balogh. Unitary units of the group algebra of modular groups. *J. Algebra Appl.*, 21(2):Paper No. 2250027, 7, 2022.
- [3] Z. Balogh, L. Creedon, and J. Gildea. Involutions and unitary subgroups in group algebras. *Acta Sci. Math. (Szeged)*, 79(3-4):391–400, 2013.
- [4] Z. Balogh and V. Laver. Unitary subgroups of commutative group algebras of characteristic 2. *Ukrain. Mat. Zh.*, 72(6):751–757, 2020.
- [5] A. Bovdi. The group of units of a group algebra of characteristic p . *Publ. Math. Debrecen*, 52(1-2):193–244, 1998.
- [6] A. Bovdi and L. Erdei. Unitary units in modular group algebras of groups of order 16. *Technical Reports, Universitas Debrecen, Dept. of Math., L. Kossuth Univ.*, 4(157):1–16, 1996.
- [7] A. Bovdi and L. Erdei. Unitary units in modular group algebras of 2-groups. *Comm. Algebra*, 28(2):625–630, 2000.
- [8] A. A. Bovdi. Unitarity of the multiplicative group of an integral group ring. *Mat. Sb. (N.S.)*, 119(161)(3):387–400, 448, 1982.
- [9] A. A. Bovdi and A. A. Sakach. Unitary subgroup of the multiplicative group of a modular group algebra of a finite abelian p -group. *Mat. Zametki*, 45:6:23–29, 1989.
- [10] A. A. Bovdi and A. Szakács. A basis for the unitary subgroup of the group of units in a finite commutative group algebra. *Publ. Math. Debrecen*, 46(1-2):97–120, 1995.
- [11] V. Bovdi and L. G. Kovács. Unitary units in modular group algebras. *Manuscripta Math.*, 84(1):57–72, 1994.
- [12] V. Bovdi and A. L. Rosa. On the order of the unitary subgroup of a modular group algebra. *Comm. Algebra*, 28(4):1897–1905, 2000.

- [13] V. Bovdi and M. Salim. On the unit group of a commutative group ring. *Acta Sci. Math. (Szeged)*, 80(3-4):433–445, 2014.
- [14] V. A. Bovdi and A. N. Grishkov. Unitary and symmetric units of a commutative group algebra. *Proc. Edinb. Math. Soc. (2)*, 62(3):641–654, 2019.
- [15] L. Creedon and J. Gildea. Unitary units of the group algebra $\mathbb{F}_{2^k}Q_8$. *Internat. J. Algebra Comput.*, 19(2):283–286, 2009.
- [16] L. Creedon and J. Gildea. The structure of the unit group of the group algebra $\mathbb{F}_{2^k}D_8$. *Canad. Math. Bull.*, 54(2):237–243, 2011.
- [17] S. P. Novikov. Algebraic construction and properties of Hermitian analogs of K -theory over rings with involution from the viewpoint of Hamiltonian formalism. Applications to differential topology and the theory of characteristic classes. I. II. *Izv. Akad. Nauk SSSR Ser. Mat.*, 34:253–288; *ibid.* 34 (1970), 475–500, 1970.
- [18] J.-P. Serre. Bases normales autoduales et groupes unitaires en caractéristique 2. *Transform. Groups*, 19(2):643–698, 2014.
- [19] Y. Wang and H. Liu. The unitary subgroups of group algebras of a class of finite p -groups. *Journal of Algebra and Its Applications*, 2021.