

# On Hamiltonian-Connected and Mycielski graphs

Ashok Kumar Das\* and Indrajit Paul

Department of Pure Mathematics

University of Calcutta

35, Ballygunge Circular Road

Kolkata-700019

Email Address - ashokdas.cu@gmail.com & paulindrajit199822@gmail.com

February 7, 2023

## Abstract

A graph  $G$  is Hamiltonian-connected if there exists a Hamiltonian path between any two vertices of  $G$ . It is known that if  $G$  is 2-connected then the graph  $G^2$  is Hamiltonian-connected. In this paper we prove that the square of every self-complementary graph of order greater than 4 is Hamiltonian-connected. If  $G$  is a  $k$ -critical graph, then we prove that the Mycielski graph  $\mu(G)$  is  $(k+1)$ -critical graph. Jarnicki et al. [9] proved that for every Hamiltonian graph of odd order, the Mycielski graph  $\mu(G)$  of  $G$  is Hamiltonian-connected. They also pose a conjecture that if  $G$  is Hamiltonian-connected and not  $K_2$  then  $\mu(G)$  is Hamiltonian-connected. In this paper we also prove this conjecture.

Keywords: *Hamiltonian-connected graphs, self-complementary graphs, Mycielski graphs.*

## 1. INTRODUCTION

A graph  $G$  is Hamiltonian if it has a cycle containing all the vertices of  $G$ . Hamiltonian graphs have been extensively studied by several researchers. Ronald J. Gould has written a survey paper [8] on Hamiltonian graphs. But unfortunately, no easy testable characterization is known for Hamiltonian graphs. Bondy and Chvátal [1] proved that a graph is Hamiltonian if and only if its closure  $Cl(G)$  is Hamiltonian. *Closure* of  $G$  is formed by recursively joining two non-adjacent vertices of  $G$  whose degree sum is at least  $n$ , where  $n$  is the number of vertices of  $G$ . But this condition for Hamiltonicity doesn't help much as we are required to test another graph to be Hamiltonian. However, it provides a method for proving a sufficient condition for Hamiltonicity. A condition that forces  $Cl(G)$  to be Hamiltonian also forces  $G$  to be Hamiltonian. For example, if  $Cl(G)$  is complete then  $G$  is also Hamiltonian. Chvátal [4] used this method to provide one of the strongest sufficient conditions for a graph to be Hamiltonian based on the degrees of the vertices of the graph.

---

\*Corresponding author

A path in  $G$  is *Hamiltonian path* if it contains all the vertices of the graph  $G$ . A graph  $G$  is *Hamiltonian-connected* if for any two vertices  $u, v \in V(G)$ , there exists a  $u$ - $v$  Hamiltonian path. Obviously, a Hamiltonian-connected graph is Hamiltonian. However, the converse is not true.

Let  $G = (V, E)$  be a connected graph and  $u, v$  are two distinct vertices of  $G$ . Then *distance* between  $u$  and  $v$ , denoted by  $d(u, v)$ , is the length of the shortest path between  $u$  and  $v$ . *Diameter* of  $G$ , denoted by  $diam(G)$  is the largest distance between the pair of vertices in  $G$ . The square of a graph  $G$ , denoted by  $G^2$ , is a graph with vertex set  $V$  and  $uv$  is an edge of  $G^2$  if and only if  $d(u, v) \leq 2$  in  $G$ . Similarly, the cube of  $G$ , denoted by  $G^3$ , is the graph with vertex set  $V$  and  $uv$  is an edge of  $G^3$  if and only if  $d(u, v) \leq 3$ . In this paper we give few more results on Hamiltonian-Connected graphs and Mycielski's graphs.

## 2. Self-complementary Graphs and Hamiltonian Connectedness

A graph is *self-complementary* if the graph is isomorphic to its complement. A graph  $G$  is  $k$ -*connected* ( $k \geq 1$ ) if removal of vertices, (fewer than  $k$ ) keeps the graph connected. If removal of a single vertex  $v$  of  $G$  disconnects the graph, then  $v$  is a *cut vertex* of  $G$ .

As mentioned in the introduction the following theorem, due to Chvátal [4], is a sufficient condition for a graph to be Hamiltonian based on the degrees of the vertices of the graph.

**Theorem 1** ([4]). *Let  $G$  be a graph of order  $n > 3$ , the degrees  $d_i$  of whose vertices satisfy  $d_1 \leq d_2 \leq \dots \leq d_n$ . If for any  $k < \frac{n}{2}$ , we have  $d_k \geq k + 1$  or  $d_{n-k} \geq n - k$ , then  $G$  is Hamiltonian.*

It is easy to observe that a graph  $G$  has a Hamiltonian path if and only if the graph  $G \vee K_1$  has a Hamiltonian cycle, where  $G \vee K_1$  is the graph obtained from  $G$  by joining a new vertex  $w$  to each vertex of  $G$ . Based on this observation, the next theorem analogous to the Theorem 1, gives a sufficient condition for the existence of a Hamiltonian path in  $G$ .

**Theorem 2** ([12]). *Let  $G$  be a graph of order  $n \geq 2$ , the degree  $d_i$  of whose vertices satisfy  $d_1 \leq d_2 \leq \dots \leq d_n$ . If for any  $k < \frac{n+1}{2}$ , we have  $d_k \geq k$  or  $d_{n+1-k} \geq n - k$ , then  $G$  has a Hamiltonian path.*

Next, we state the theorem due to O.Ore, which provides a sufficient condition for a graph to be Hamiltonian-connected.

**Theorem 3** ([10]). *Let  $G$  be a graph of order  $n \geq 3$ . If for any two non-adjacent vertices  $u$  and  $v$  of  $G$ ,  $d(u) + d(v) \geq n + 1$ , then  $G$  is Hamiltonian-connected.*

It can be easily checked that the square of a tree of order at least 4 is not necessarily Hamiltonian. But M.Sekanina [11] and others proved that for any connected graph  $G$ , the *cube* of  $G$  is Hamiltonian-connected. Although Nash-Williams and M.D. Plummer conjectured that for any 2-*connected* graph  $G$ , the graph  $G^2$  is Hamiltonian. Fleischner [6] verified this conjecture. The work of Fleischner was strengthened by Chartrand, Hobbs, Jung, Kapoor and Nash-Williams[2] showing that the square of a 2-*connected* graph is Hamiltonian-connected.

**Theorem 4** ([2]). *Let  $G$  be a 2-connected graph of order at least 3. Then  $G^2$  is Hamiltonian-connected.*

A self complementary graph of order  $n \geq 5$ , may not be 2-connected. For example, the following self complementary graph is not 2-connected.




Figure 1: Self-complementary graph on 5 vertices which is not 2-connected.

In this context we prove the following result.

**Theorem 5.** *If  $G$  is self-complementary graph of order  $n > 3$ , then  $G$  has a Hamiltonian path. Moreover, if  $n \geq 5$ , then  $G^2$  is Hamiltonian connected.*

*Proof.* The first part of the theorem is known [3]. But, for the sake of completeness we give here a new proof. Let the degree sequence of  $G$  be  $d_1, d_2, \dots, d_k, \dots, d_n$  satisfying the condition  $d_1 \leq d_2 \leq \dots \leq d_k \leq \dots \leq d_n$ , where  $d_i$  is the degree of the vertex  $v_i$ . Then the degrees of  $\overline{G}$  satisfy the condition

$$n - 1 - d_n \leq n - 1 - d_{n-1} \leq \dots \leq n - 1 - d_k \leq \dots \leq n - 1 - d_2 \leq n - 1 - d_1 \quad (1)$$

Now, for any  $k < \frac{n+1}{2}$ , let  $d_k \leq k - 1$ . Then

$$n - 1 - d_k \geq n - k \quad (2)$$

Again Eq. (1) can be written as

$$n - 1 - d_{(n+1)-1} \leq n - 1 - d_{(n+1)-2} \leq \dots \leq n - 1 - d_{(n+1)-k} \leq \dots \leq n - 1 - d_{(n+1)-(n-1)} \leq n - 1 - d_{(n+1)-n}$$

Because  $G$  is self-complementary,  $d_k = n - 1 - d_{n+1-k}$ . So  $d_{n+1-k} = n - 1 - d_k$ . Therefore from Eq. (2),  $d_{n+1-k} \geq n - k$ . Therefore from Theorem 2,  $G$  has a Hamiltonian path.

We can also conclude that  $G$  and hence  $\overline{G}$  are connected.

To prove the second part, first we shall state the following known result.

**Lemma 6.** [7] *If  $G$  is a self-complementary graph of order at least 5, then  $\text{diam}(G) = 2$  or 3.*

Now we shall prove the second part of the Theorem 5.

Let  $G$  be a self complementary graph of order  $n \geq 5$ . We shall prove that  $G^2$  is Hamiltonian-connected.

First, suppose  $\text{diam}(G) = 2$ . Then  $G^2$  is complete and hence  $G^2$  is Hamiltonian-connected. Next, suppose  $\text{diam}(G) = 3$ . If  $G$  is 2-connected then by Theorem 4,  $G^2$  is Hamiltonian-connected. So, we assume  $G$  has a cut-vertex  $v$  and  $G_1, G_2, \dots, G_k$  are the components of  $G \setminus \{v\}$ . Let  $G_i$  and  $G_j$  be any two components of  $G \setminus \{v\}$ . Then every vertex of  $G_i$  is adjacent to every vertex of  $G_j$  in  $\overline{G}$ . Thus we conclude that  $\overline{G}^2 \setminus \{v\} = K_{n-1}$ . Now  $d_G(v) \geq 2$ , then if  $d_{\overline{G}}(v) \geq 3$ , we consider two non-adjacent vertices of  $\overline{G}^2$ . One of these two

vertices must be  $v$  and, say, another vertex is  $u$ , then  $d_{\overline{G}^2}(u) = n - 2$ . Since  $d_{\overline{G}}(v) \geq 3$  implies  $d_{\overline{G}^2}(v) \geq 3$ , we have for non-adjacent vertices  $u$  and  $v$ ,  $d_{\overline{G}^2}(u) + d_{\overline{G}^2}(v) \geq n - 2 + 3 = n + 1$ . Therefore by [Theorem 3](#),  $\overline{G}^2$  is Hamiltonian-connected. Now, we examine the other possibilities.

**Case 1.** Let  $d_{\overline{G}}(v) = 1$ . Thus  $d_G(v) = n - 2$  and there exists only one vertex  $x$  such that  $vx \in E(\overline{G})$ . Since  $\text{diam}(\overline{G}) = 3$ , every vertex in  $\overline{G}$  is at a distance at most 2 from  $x$ . Now, if  $d_{\overline{G}}(x) \geq 3$  then  $d_{\overline{G}^2}(v) \geq 3$ . Then, as before  $\overline{G}^2$  is Hamiltonian-connected. Therefore assume  $d_{\overline{G}}(x) = 2$ , where  $N_{\overline{G}}(x) = \{v, z\}$  (Figure 2). Since  $\text{diam}(\overline{G}) = 3$ , every vertex of  $V(G) \setminus \{v\}$  is adjacent to  $z$  in  $\overline{G}$ . Thus  $d_{\overline{G}}(z) = n - 2$  and no other vertex in  $\overline{G}$  is of degree  $n - 2$ . Since  $G$  is a self complementary graph and  $z$  is a cut vertex of  $\overline{G}$ , as before  $G^2 \setminus \{z\} = K_{n-1}$ . Again, all the vertices except  $x$ , that are adjacent to  $z$  in  $\overline{G}$  are also adjacent to  $v$  in  $G$ . This implies  $d_{G^2}(z) = n - 2 \geq 3$ , as  $n \geq 5$ . So in  $G^2$ ,  $d_{G^2}(z) + d_{G^2}(u) \geq n - 2 + 3 = n + 1$ , where  $u$  is any other vertex of  $G$  other than  $z$ . Hence  $G^2$  is Hamiltonian-connected.




Figure 2: A possible figure in  $\overline{G}$ , where dotted edges represent the edges of  $G$ .

**Case 2.** Let  $d_{\overline{G}}(v) = 2$ . Since  $v$  is a cut vertex of  $G$ ,  $v$  is not a cut vertex of  $\overline{G}$ . Now, assume  $N_{\overline{G}}(v) = \{a, b\}$ . Since  $v$  is not a cut vertex of  $\overline{G}$ , at least one vertex of  $a$  and  $b$ , say  $a$ , has a neighbour other than  $b$  and  $v$  in  $\overline{G}$ . This implies  $d_{\overline{G}^2}(v) \geq 3$ . Again, as before  $\overline{G}^2 \setminus v = K_{n-1}$ . So for any two vertices  $x, y$  of  $G$  other than  $v$ , we have  $d(x) + d(y) = 2n - 4$  in  $K_{n-1}$ . As  $n \geq 5$ ,  $2n - 4 \geq n + 1$ , i.e.  $d_{\overline{G}^2}(x) + d_{\overline{G}^2}(y) \geq n + 1$ . Finally,  $d_{\overline{G}^2}(v) + d_{\overline{G}^2}(x) \geq n - 2 + 3 = n + 1$ , where  $x \in V - \{v\}$ . Hence  $\overline{G}^2$  is Hamiltonian-connected and accordingly  $G^2$  is Hamiltonian-connected. This completes the proof of the [Theorem 5](#). ■

### 3. Mycielski's Graphs

Let  $G = (V, E)$  be a connected graph with vertex set  $V = \{v_1, v_2, \dots, v_n\}$ . The Mycielski's graph  $\mu(G)$  of  $G$  is a graph with the vertex set  $X \cup Y \cup \{z\}$ , where  $X = \{x_1, x_2, \dots, x_n\}$ ,

$Y = \{y_1, y_2, \dots, y_n\}$ , and the vertices of  $X$  induce  $G$ . The new edges in  $\mu(G)$  are  $zy_i$  for all  $i$  also  $x_i y_j$  is an edge of  $\mu(G)$  if  $x_i x_j$  is an edge of  $G$ . For example,  $\mu(K_2) = C_5$  and  $\mu(C_5)$  is the Grötzsch graph (Figure 3).




Figure 3: The Grötzsch Graph.

Thus starting with  $M_2 = K_2$ , the Mycielski's graph  $M_n$  are formed by iterating  $\mu$  successively  $(n - 2)$  times on  $K_2$ . Thus  $M_3 = \mu(K_2)$ ,  $M_4 = \mu(M_3)$  and in general  $M_{k+1} = \mu(M_k)$ . Mycielski's graphs are of interest because if  $G$  is triangle-free then  $\mu(G)$  is also triangle-free and if the chromatic number  $\chi(G)$  of  $G$  is  $k$ , then  $\chi(\mu(G)) = k + 1$ .

Now,  $M_3$  is  $C_5$ ; also it can be observed that  $M_4$ , the Grötzsch graph is Hamiltonian. Fisher et al. [5] proved that if  $G$  is Hamiltonian then  $\mu(G)$  is also Hamiltonian. Jarnicki et al. [9] have extended this result and proved that  $\mu(G)$  is Hamiltonian-connected if  $G$  is Hamiltonian and of odd order. In this paper we prove that each Mycielski's graph is a  $k$ -critical graph. Finally, we shall prove a conjecture posed by Jarnicki et al. [9] regarding Hamiltonian-connectedness of Mycielski's graphs. First, we shall prove the following proposition.

**Proposition 7.** *If  $P_n$  is a path on  $n$  vertices, then  $\mu(P_n)$  has a Hamiltonian path.*

*Proof.* Let  $x_1 x_2 \dots x_{n-1} x_n$  be the path  $P_n$ . First, suppose  $n$  is odd, then  $\mu(P_n)$  has a Hamiltonian path,  $x_1 y_2 x_3 y_4 \dots x_{n-2} y_{n-1} x_n x_{n-1} y_n z y_1 x_2 y_3 x_4 \dots x_{n-3} y_{n-2}$ . Next, if  $n$  is even then  $x_1 y_2 x_3 y_4 \dots x_{n-1} y_n z y_1 x_2 y_3 x_4 \dots y_{n-1} x_n$  is a Hamiltonian path in  $\mu(P_n)$ . ■

A graph  $G$  is  **$k$ -critical graph** if  $\chi(G) = k$  but  $\chi(H) < \chi(G)$  for every proper subgraph  $H$  of  $G$ . Now we prove the following propositions.

**Proposition 8.** *Let  $G$  be a  $k$ -chromatic graph, where  $k \geq 2$ . Then for every vertex  $v$  of  $G$ ,  $\chi(G - v) = k$  or  $k - 1$ .*

*Proof.* On the contrary, Let  $\chi(G - v) = p \leq k - 2$ . Now, if  $v$  is adjacent to the vertices in  $G - v$  which receives all the  $p$  colors. Then we can color the vertex  $v$  by  $(p + 1)$ th color. Which implies that  $\chi(G) \leq k - 1$ . Thus we arrive at a contradiction. Hence,  $\chi(G - v) = k$  or  $k - 1$ . ■

**Proposition 9.** Let  $G$  be a  $k$ -chromatic graph, where  $k \geq 2$ . Then for every edge  $e$  of  $G$ ,  $\chi(G - e) = k$  or  $k - 1$ .

*Proof.* On the contrary, assume  $\chi(G - e) = p \leq k - 2$  where  $e = x_i x_j$ . Now in  $G - e$  if both  $x_i$  and  $x_j$  receive the  $p$ th color then recolor  $x_i$  or  $x_j$  by  $(p + 1)$ th color. Which is a contradiction. Therefore  $\chi(G - e) = k$  or  $k - 1$ . ■

From the above two propositions we conclude that if  $G$  is  $k$ -critical graph, then for every vertex  $v$ ,  $\chi(G - v) = k - 1$  and for every edge  $e$ ,  $\chi((G - e)) = k - 1$ . Now, we prove the following theorem.

**Theorem 10.** Every Mycielski graph  $M_k$  is a  $k$ -critical graph for  $k \geq 2$ .

*Proof.* To prove this theorem, we use mathematical induction on  $k$ . We can easily see that  $M_2 = K_2$  and  $M_3 = C_3$  are respectively 2-critical and 3-critical graphs. Now, we assume that  $M_p$  is a  $p$ -critical graph for  $p > 3$  and we shall prove that  $M_{p+1}$  is  $(p + 1)$ -critical graph. Again  $\chi(M_p) = p$  and suppose  $V(M_p) = \{x_1, x_2, \dots, x_n\}$ . The vertex set  $V(M_{p+1})$ , of  $M_{p+1}$  is  $\{x_1, x_2, \dots, x_n\} \cup \{y_1, y_2, \dots, y_n\} \cup \{z\}$ . We know that  $\chi(M_{p+1}) = p + 1$  and we shall prove that  $\chi(M_{p+1} - v) = p$  for  $v \in V(M_{p+1})$ . We consider the following cases.

**Case 1.** Let  $v = z$ . Since  $\chi(M_p) = p$ , from the construction of  $M_{p+1}$ , it follows that  $\chi(M_{p+1} - z) = p$ .

**Case 2.** Let  $v = x_i \in V(M_p)$ . Since  $M_p$  is  $p$ -critical,  $\chi(M_p - x_i) = p - 1$ . Suppose  $1, 2, 3, \dots, (p - 1)$  colors are needed to color the vertices  $x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ .

Again, each vertex  $y_i$  where  $(1 \leq i \leq n)$  is adjacent to some vertices of the set  $\{x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n\}$ . So we can use  $p$ -th color to color the vertices  $\{y_1, y_2, \dots, y_n\}$ . Then we can use any of the colors  $1, 2, 3, \dots, (p - 1)$  to color the vertex  $z$ . Thus  $\chi(M_{p+1} - x_i) = p$ .

**Case 3.** Let  $v = y_i$ . Now  $x_i$  and  $y_i$  are non adjacent and  $x_i$  and  $y_i$  have same adjacency in  $V(M_{p+1}) \setminus \{z\}$  for each  $i$  ( $1 \leq i \leq n$ ). Since  $M_p$  is  $p$ -critical,  $\chi(M_p - x_i) = p - 1$  and we recolor the vertices  $x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n$  by  $1, 2, 3, \dots, (p - 1)$  colors. We can color the vertices  $y_j$  and  $x_j$  by same color for every  $j \neq i$ . Then we color the vertices  $x_i$  and  $z$  by  $p$ -th color. Thus  $\chi(M_{p+1} - y_i) = p$ .

Next, we consider the edge deletion cases.

**Case 4.** Let  $e = x_i x_j$ . Since  $M_p$  is  $p$ -critical,  $\chi(M_p - e) = p - 1$ . Again, existence of the edge  $x_i x_j$  implies the existence of the edges  $x_i y_j$  and  $x_j y_i$  in  $M_{p+1}$ . Now the vertices of  $M_p - e$  can be colored by  $(p - 1)$  numbers of colors, say  $1, 2, \dots, (p - 1)$ ; where  $x_i$  and  $x_j$  both receive the same color. Now in  $M_{p+1}$  the vertices  $y_1, y_2, \dots, y_i, \dots, y_j, \dots, y_n$  are colored by  $1, 2, \dots, p - 1, p$  colors (i.e.  $p$  number of colors). Next, we delete the edges  $x_i y_j$  and  $x_j y_i$  from  $M_{p+1} - e$ . Now, since in  $M_{p+1} - \{x_i y_j, x_j y_i, x_i x_j\}$  the vertices  $x_k$  and  $y_k$  ( $1 \leq k \leq n$ ) have the same neighbours in the vertex set of  $M_p - x_i x_j$ , the vertices  $y_1, y_2, \dots, y_n$  now can be colored by  $1, 2, \dots, (p - 1)$  colors. Next, we recolor the vertices  $x_i$  and  $x_j$  in  $M_p - e$  by the  $p$ th color and insert the edges  $x_i y_j$  and  $x_j y_i$  to obtain the graph  $M_{p+1} - e$ . Then the  $p$ th color can be used to color  $z$ . This proves that  $\chi(M_{p+1} - e) = p$ .

**Case 5.** Let  $e = x_i y_j$ . Now, color the vertices  $x_1, x_2, \dots, x_i, \dots, x_{j-1}, x_{j+1}, \dots, x_n$  of  $M_{p+1}$  by  $(p - 1)$  colors as above and use the  $p$ th color to color the vertex  $x_j$ . Again in  $M_{p+1} - e$ , the edge  $x_j y_i$  exists and the vertex  $x_j$  is colored by  $p$ th color. So the vertices  $y_1, y_2, \dots, y_n$  can be colored by  $1, 2, \dots, (p - 1)$  colors in  $M_{p+1} - e$  as in the case 4. Thus we can use the  $p$ th color to color the vertex  $z$ . Hence  $\chi(M_{p+1} - e) = p$ .

**Case 6.** Let  $e = y_i z$ . Recolor the vertices of  $M_p$  by  $1, 2, \dots, p$  colors, where  $x_i$  is colored by  $p$ th color as  $M_p - x_i x_j$  is colored by  $1, 2, \dots, (p - 1)$  colors. In  $M_{p+1}$  the vertices  $x_k$  and  $y_k (1 \leq k \leq n)$  receive the same color as their adjacencies are same in  $M_p$ . Thus the vertex  $y_i$  can also be colored by  $p$ th color. So in  $M_{p+1} - e$ , the vertex  $z$  can be colored by  $p$ th color. Hence  $\chi(M_{p+1} - e) = p$ .

Hence  $M_{p+1}$  is  $(p + 1)$ -critical graph. This completes the proof of this theorem.  $\blacksquare$

From the proof of Theorem 10, it follows that the above result can be generalized for arbitrary  $k$ -critical graph  $G$ .

**Corollary 11.** *If  $G$  is any  $k$ -critical graph, then  $\mu(G)$  is  $(k + 1)$ -critical graph.*

We have mentioned earlier that Jarnickie et al. [9] proved that if  $G$  is Hamiltonian and of odd order then  $\mu(G)$  is Hamiltonian-connected. They have also proved that this result does not hold if the order of  $G$  is even. In this context they have conjectured the following, which we shall prove in the next theorem.

**Theorem 12.** *If the graph  $G$  is Hamiltonian-connected and not  $K_2$ , then  $\mu(G)$  is also Hamiltonian-connected.*

To prove the above theorem, first we prove the following lemmas.

**Lemma 13.** *If  $G$  is a Hamiltonian-connected graph, then  $G$  must contains odd cycle.*

*Proof.* If  $G$  is bipartite then  $G$  is not Hamiltonian-connected. Thus  $G$  must contains an odd cycle.

**Lemma 14.** *If  $G$  is a Hamiltonian-connected graph of order  $\geq 4$ , then degree of  $v$ ,  $d(v) \geq 3$  for all  $v \in V(G)$ .*

*Proof.* Let  $n$  be the order of  $G$ . Suppose  $v_1 v_2 v_3 \dots v_n v_1$  is a Hamiltonian cycle in  $G$ . Next, assume  $d(v_2) = 2$ . Then we do not have any  $v_1$ - $v_3$  Hamiltonian path in  $G$ . Which contradicts that  $G$  is Hamiltonian-connected and complete the proof of the lemma.  $\blacksquare$

Now we prove the Theorem 12.

*Proof.* Let  $G = (V, E)$  be a Hamiltonian-connected graph, where  $V = \{v_1, v_2, v_3, \dots, v_{n-1}, v_n\}$ . Since Hamiltonian-connected graph must be Hamiltonian, so if  $n$  is odd then by Theorem 5.1 of [9],  $\mu(G)$  is Hamiltonian-connected. Thus we suppose that  $n$  is even.

Now, we assume  $\mu(G)$  has vertices  $X \cup Y \cup \{z\}$  where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $Y = \{y_1, y_2, \dots, y_n\}$ . Also  $G[X]$ , the subgraph induced by the vertex set  $X$  is  $G$  and  $x_1, x_2, x_3, \dots, x_{n-1}, x_n, x_1$  is a Hamiltonian cycle. Also,  $\mu(G)$  has extra edges  $z y_i$  for all  $i$ , and  $y_i x_j$  and  $y_j x_i$  are edges of  $\mu(G)$  if  $x_i x_j \in E(G)$ . To prove that  $\mu(G)$  is Hamiltonian-connected, we have to show that for any two vertices  $u, v \in V(\mu(G))$  there exists a  $u$ - $v$  Hamiltonian path in  $\mu(G)$ .

Now, without loss of generality we consider the following cases.

**Case 1.** Let  $x_1, x_k \in V(G)$ . Since  $G$  is Hamiltonian-connected suppose we have the following  $x_1$ - $x_k$  Hamiltonian path:  $x_1 x_j x_l \dots x_i x_m x_k$  in  $G$ . Then we construct the following  $x_1$ - $x_k$

Hamiltonian path  $P_1$  in  $\mu(G)$  where

$$P_1 : x_1 y_j x_l \cdots y_i x_m y_k z y_1 x_j y_l \cdots x_i y_m x_k$$

**Case 2.** Let  $x_1, y_p \in V(\mu(G))$ . Suppose we have the following  $x_1$ - $x_p$  Hamiltonian path  $P$  in  $G$ , where

$$P : x_1 x_i x_j x_k x_r x_s \dots x_t x_l x_m x_p$$

Now, we construct the following Hamiltonian  $x_1$ - $y_p$  path  $P_1$  in  $\mu(G)$  where

$$P_1 : x_1 y_i x_j y_k \dots x_t y_l x_m x_p y_m x_l y_t \dots x_k y_j x_i y_1 z y_p$$

**Case 3.** Let  $x_1, z \in V(\mu(G))$ . Since  $G$  is Hamiltonian-connected, assume  $P'$  be the  $x_1$ - $x_n$  Hamiltonian path in  $G$ , with  $x_1 x_n \in E(G)$ , where  $P' : x_1 x_2 \dots x_i x_r \dots x_q x_s \dots x_n$ . Now,  $n$  is even and  $G$  contains an odd cycle, also  $d(v) \geq 3$  for all  $v \in V(G)$ , there must exist two even (or odd) indexed vertices of  $P'$  that are adjacent. If  $x_r$  and  $x_s$  are two vertices of odd index, then we relabel the vertices of the Hamiltonian cycle  $x_1 x_2 \dots x_i x_r \dots x_q x_s \dots x_n x_1$  so that  $x_r$  and  $x_s$  are of even index. Thus without loss of generality, let  $x_r$  and  $x_s$  are two such vertices of even index. Next, as  $G$  is Hamiltonian-connected, there exists a  $x_i$ - $x_r$  Hamiltonian path  $P''$  in  $G$ , where  $P'' : x_i x_j \dots x_s \dots x_t x_r$ , and  $x_i x_r$  and  $x_s x_r \in E(G)$ .

Now, we relabel the vertices of  $P''$  to get  $P$ , where  $P : x_1 x_2 x_3 \dots x_{p-1} x_p x_{p+1} \dots x_{n-1} x_n$  and  $x_1 x_n, x_p x_n \in E(G)$  (i.e. we relabel  $x_i$  by  $x_1$ ,  $x_j$  by  $x_2, \dots$ ,  $x_s$  by  $x_p$ ,  $x_r$  by  $x_n$  etc.).

Next, we construct the  $x_1$ - $z$  Hamiltonian path  $P_1$  in  $\mu(G)$  as follows (Fig. 4).

$$P_1 : x_1 y_2 x_3 y_4 x_5 \dots y_{p-2} x_{p-1} y_p x_{p+1} y_{p+2} \dots x_{n-3} y_{n-2} x_{n-1} y_n x_p y_{p-1} x_{p-2} \dots y_5 x_4 y_3 x_2 y_1 x_n y_{n-1} x_{n-2} \\ y_{n-3} \dots x_{p+2} y_{p+1} z$$






Figure 4: The  $x_1$ - $z$  path  $P_1$ .

**Case 4.** Let  $x_1, y_1 \in V(\mu(G))$ . Now we have the following  $x_1$ - $x_n$  Hamiltonian path  $P$  in  $G$  where  $P : x_1x_2x_3x_4 \dots x_{r-1}x_rx_{r+1} \dots x_{n-2}x_{n-1}x_n$ . Again, as in the case 3, suppose  $x_1x_n$  and  $x_rx_n \in E(G)$  and  $x_r$  is a vertex of even index. Then we construct the following  $x_1$ - $y_1$  Hamiltonian path  $P_1$  in  $\mu(G)$ , where

$$P_1 : x_1x_2y_3x_4y_5 \dots y_{r-1}x_r y_n x_{n-1}y_{n-2} \dots y_{r+2}x_{r+1}y_r x_{r-1} \dots y_4x_3y_2z y_{r+1}x_{r+2} \dots x_{n-2}y_{n-1}x_n y_1$$

**Case 5.** Let  $y_1, z \in V(\mu(G))$  also  $y_1z$  is an edge of  $\mu(G)$ . Let  $x_1x_l \in E(G)$ , then since  $G$  is Hamiltonian-connected we have a  $x_1$ - $x_l$  Hamiltonian path  $P$  in  $G$  where

$$P : x_1 x_i x_j x_m \dots x_k x_p x_l$$

Thus in  $\mu(G)$ , we have the following  $y_1$ - $z$  Hamiltonian path  $P_1$  where

$$P_1 : y_1 x_i y_j x_m \dots x_k y_p x_l x_1 y_i x_j \dots y_k x_p y_l z$$

**Case 6.** Let  $y_1, y_2 \in V(\mu(G))$  where  $x_1x_2 \in E(G)$ . Now, let  $x_1x_l \in E(G)$  and assume the following  $x_1$ - $x_l$  Hamiltonian path  $P$  in  $G$ , where  $P : x_1 x_2 x_i x_j \dots x_k x_m x_l$ . Then we construct the following  $y_1$ - $y_2$  Hamiltonian path  $P_1$  in  $\mu(G)$ , where

$$P_1 : y_1 x_2 x_i y_j \dots y_k x_m y_l x_1 x_l y_m x_k \dots x_j y_i z y_2$$

**Case 6.1.** Let  $y_1, y_p \in V(\mu(G))$ , where  $p$  is even and  $x_1x_p \notin E(G)$ . Also, assume  $x_1x_n \in E(G)$ . Since  $G$  is Hamiltonian-connected,  $P$  be the  $x_1$ - $x_n$  Hamiltonian path, where  $P :$

$x_1 x_2 x_3 x_4 x_5 \dots x_{p-1} x_p x_{p+1} \dots x_{n-2} x_{n-1} x_n$ . Next, we construct the  $y_1$ - $y_p$  Hamiltonian path  $P_1$  in  $\mu(G)$  as follows ( Figure 5).

$$P_1 : y_1 x_2 y_3 x_4 y_5 \dots y_{p-1} x_p x_{p+1} y_{p+2} \dots y_{n-2} x_{n-1} y_n x_1 x_n y_{n-1} x_{n-2}$$

$$\dots x_{p+2} y_{p+1} z y_2 x_3 y_4 \dots x_{p-1} y_p.$$




Figure 5: The  $y_1$ - $y_p$  path  $P_1$  ( $p$  is even).

**Case 6.2.** Let  $y_1, y_p \in V(\mu(G))$  where  $p$  is odd and  $x_1 x_p \notin E(G)$ . Now, assume  $x_1$ - $x_n$  Hamiltonian path  $P$  in  $G$ , where  $P : x_1 x_2 x_3 x_4 \dots x_{p-1} x_p x_{p+1} \dots x_{n-2} x_{n-1} x_n$  and  $x_1 x_n \in E(G)$ . Next, we construct the  $y_1$ - $y_p$  Hamiltonian path  $P_1$  in  $\mu(G)$  as follows ( Figure 6).

$$P_1 : y_1 x_2 y_3 x_4 \dots y_{p-2} x_{p-1} x_p y_{p+1} x_{p+2} \dots y_{n-2} x_{n-1} y_n z y_{p-1} x_{p-2} y_{p-3} \dots x_5 y_4 x_3 y_2 x_1 x_n$$

$$y_{n-1} x_{n-2} \dots x_{p+3} y_{p+2} x_{p+1} y_p.$$




Figure 6: The  $y_1$ - $y_p$  path  $P_1$  ( $p$  is odd).

This completes the proof of Theorem 12. ■

## References

- [1] J.A. Bondy and V. Chvátal, A method in graph theory. *Discrete Math* 15(1976), 111–136.
- [2] G. Chartrand, A.M. Hobbs, H.A. Jung, S.F. Kapoor and C.St.J.A. Nash-Williams, The square of a block is Hamiltonian-connected. *J. Combin. Theory Ser. B* 16(1974), 290–292.
- [3] G.Chartrand, L.Lesniak and P.Zang, Graphs and Digraphs, *CRC Press* 2016.
- [4] V. Chvátal, On Hamiltonian ideas. *J. Combin. Theory Ser. B* (1972), 163–168.
- [5] D.C. Fisher, P.A. McKenna, and E.D. Boyer, Hamiltoniancity, diameter, domination, packing, and biclique partition of Mycielski’s graphs. *Disc. Appl. Math.* 84(1998), 93–105.

- [6] H. Fleischner, The square of every two-connected graph is Hamiltonian. *J. Combin. Theory Ser. B* 16(1974), 29–34.
- [7] T. Gangopadhyay and S.P. Rao Hebbare, r-Partite Self-Complementary Graphs-Diameters. *Discrete Mathematics* 32(1980) 245-255.
- [8] R.J. Gould, Advances on Hamiltonian Problem - A survey, 2002, Emory University, Atlanta GA30322.
- [9] W. Jarnicki, W. Myrvold, P. Saltzman, and S. Wagon, Properties, proved and conjectured, of Keller, Mycielski, and queen graphs. *Ars Mathematica Contemporanea* 13(2017), 427–460.
- [10] O. Ore, Hamilton Connected graphs. *J. Math Pures Appl.* 42(1963), 21–27.
- [11] M. Sekanina, On an ordering of the set of vertices of a connected graph. *Spisy Prirod Fak. Univ. Brno.* (1960), 137–141.
- [12] D.B. West, Introduction to graph theory (Second Editon). PHI Learning Pvt. Limited, New Delhi (2012).