

Turaev–Viro TQFT and the Rank versus Genus Conjecture

QING LAN

June 7, 2022

Abstract

This paper presents a way to estimate the Heegaard genus of a 3-manifold using the Turaev–Viro state sum TQFT. The Turaev–Viro state sum TQFT is derived from the modular category associated to the quantum group $U_q(\mathfrak{sl}_2)$, which is unitary for some q by Wenzl. Hence by Turaev and Virelizier the corresponding TQFT is unitary. We modify a proof by Garoufalidis to give a lower bound of the Heegaard genus using a unitary TQFT, and then use the software *Regina* to provide some known counterexamples to the rank versus genus conjecture.

1 Overview

For a closed orientable 3-manifold M , the *Heegaard genus* $g(M)$ of M , is the minimal genus over all Heegaard splittings of M . The *rank* $r(M)$ of M is the rank of the fundamental group $\pi_1(M)$. Hence $r(M) \leq g(M)$.

Conjecture 1. $r(M) = g(M)$?

This conjecture was proposed by Waldhausen [Wal78]. In 1984, Boileau–Zieschang [BZ84] discovered the first counterexamples among Seifert manifolds. Later, for any n , graph manifolds of genus $4n$ whose fundamental group is $3n$ -generated are constructed by Schultens and Weidmann in [SW07]. In 2009, Namazi and Souto [NS09] showed that rank equals genus if the gluing map of a Heegaard splitting is a high power of a generic pseudo-Anosov map. In 2013, Tao Li showed in [Li13] that there are counterexamples among hyperbolic manifolds, using topological argument.

In this thesis, we apply the Turaev–Viro TQFT to the rank versus genus conjecture. We obtain a lower bound of the Heegaard genus of a 3-manifold, and provide some known counterexamples to the rank versus genus conjecture.

1.1 Background

Inspired by Witten, Reshetikhin and Turaev [RT91] constructed quantum invariants using representations of a quantum group $U_q(\mathfrak{sl}_2(\mathbb{C}))$. This invariant is part of a topological quantum field theory (TQFT). This was generalized to simple Lie algebras of type A, B, C and D in [TW93]. More generally, the TQFT can be constructed using a general modular category, as is explained in [Tur92]

or [Tur10]. There is another approach via Kauffman bracket, see [BHMV92], [BHMV95]. The book [Lus10] is an introduction to quantum groups.

In [TV92], Turaev and Viro constructed a TQFT using quantum $6j$ -symbols associated to $U_q(\mathfrak{sl}_2(\mathbb{C}))$. More generally, the construction can be performed using a unimodal category, see [Tur10], or using a finite semisimple spherical category of non-zero dimension, see [BW96]. This is called the Turaev–Viro–Barrett–Westbury invariant in the book by Turaev and Virelizier [TV17]. In 2018, Qingtao Chen and Tian Yang [CY18] defined the invariant for a manifold with boundary and proposed volume conjectures for the Reshetikhin–Turaev and the Turaev–Viro invariants.

The process of constructing a modular category from $U_q(\mathfrak{sl}_2(\mathbb{C}))$ using the notion of tilting modules is explained in [BK01]. In [BW96], Barrett and Westbury used the paper by Andersen and Paradowski [HAP95], and showed that the quantum \mathfrak{sl}_2 can be used to produce a state sum TQFT. (This general procedure is needed in the proof of unitarity.)

Wenzl [Wen98] showed that, if \mathfrak{g} is a simple Lie algebra and $q = e^{\frac{\pi i}{l d}}$ where $d = m \in \{1, 2, 3\}$ is the ratio of the square lengths of a long and a short root and $l = \varkappa$ is larger than the dual Coxeter number, then the category for $U_q \mathfrak{g}$ is unitary. (It is shown in [KJ96] that the category is Hermitian.) Note that the ratio is 1 for Lie type A . The dual Coxeter number of A_n is $n + 1$, and \mathfrak{sl}_2 corresponds to A_1 . There is a survey [Row06] on the general case.

It is shown in [TV17], Appendix G, that a unitary fusion category \mathcal{C} gives rise to a unitary state sum TQFT.

1.2 A Lower Bound of the Heegaard Genus

In [Gar98], Theorem 2.2 gives a lower bound of Heegaard genus using a unitary TQFT. **This theorem is not correct in general.** Note that Theorem 2.2 uses Lemma 2.1, and the author claims that Lemma 2.1 follows from a paper by Witten [Wit89]. In [Wit89], there is an implicit condition on the TQFT, namely the space associated to S^2 must be one-dimensional. This condition is not mentioned in [Gar98].

We modify this proof in [Gar98] to give the bound.

Parallel to Lemma 2.1 in [Gar98], our unitary TQFT Z must have the following three additional properties:

- (1) If M, N are closed 3-manifolds, then $Z(M \# N)Z(S^3) = Z(M)Z(N)$.
- (2) $Z(S^2 \times S^1) = 1$.
- (3) $0 < Z(S^3) < 1$.

Assuming the space associated to S^2 is one-dimensional, property (1) follows from a cut-and-paste argument (see Lemma 10.2 in [TV17]). This is true at least for the state sum TQFT. Lemma 13.6 in [TV17] states that, over any commutative ring \mathbb{k} , the \mathbb{k} -module $|S^2|_{\mathcal{C}}$ is isomorphic to \mathbb{k} . (Here \mathcal{C} is a spherical fusion \mathbb{k} -category.)

As for property (2), section 13.1.3 in [TV17] shows that $|S^1 \times S^2|_{\mathcal{C}} = 1$.

For property (3), $|S^3|_{\mathcal{C}} = (\dim(\mathcal{C}))^{-1}$. If \mathcal{C} is spherical, then $\dim(\mathcal{C}) = \sum_{i \in I} (\dim(i))^2$, where I is a (finite) representative set of simple objects of \mathcal{C} . Note that the quantum dimension is not necessarily an integer. Here, we

consider the original Turaev–Viro TQFT [TV92], at $q = e^{\pi i/r}$, $r \geq 3$. Then

$$|S^3| = -\frac{\left(e^{\frac{i\pi}{r}} - e^{-\frac{i\pi}{r}}\right)^2}{2r} = \frac{2 \sin^2\left(\frac{\pi}{r}\right)}{r} \in (0, 1).$$

Combining the results, we obtain a lower bound of the Heegaard genus. It is worth noting that Corollary 11.7 in [TV13] gives the same bound.

We use the software *Regina* to search for counterexamples. The confirmed ones have appeared in [BZ84].

Here is the structure of the thesis. Chapter 2 includes necessary definitions and the construction of the category needed, described in [BK01]. In Chapter 3 we construct the state sum TQFT and sketch a proof of unitarity, described in [TV17] and [TV13]. In Chapter 4 we present the estimation and the computer search.

2 Modular Category Associated to $U_q(\mathfrak{sl}_2)$

2.1 Definitions

Our definitions are taken from [TV17], [TV13], [Row06], and [BK01].

A *monoidal category* is a category with a tensor product \otimes and an identity object $\mathbb{1}$ satisfying axioms that guarantee that the tensor product is associative (up to specified isomorphism) and that $\mathbb{1} \otimes X \cong X \otimes \mathbb{1} \cong X$ for any object X .

Let $\mathcal{C} = (\mathcal{C}, \otimes, \mathbb{1})$ and $\mathcal{D} = (\mathcal{D}, \otimes', \mathbb{1}')$ be monoidal categories. A *monoidal functor* from \mathcal{C} to \mathcal{D} is a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ endowed with a morphism $F_0 : \mathbb{1}' \rightarrow F(\mathbb{1})$ in \mathcal{D} and with a natural transformation

$$F_2 = \{F_2(X, Y) : F(X) \otimes' F(Y) \rightarrow F(X \otimes Y)\}_{X, Y \in \text{Ob}(\mathcal{C})}$$

satisfying some compatibility conditions. The monoidal functor is *strong* if F_0 and $F_2(X, Y)$ are isomorphisms for all X, Y .

Let \mathbb{k} be a commutative ring. A \mathbb{k} -*category* \mathcal{C} is a category such that all Hom sets are left \mathbb{k} -modules and the composition of morphisms is \mathbb{k} -bilinear. An object D of \mathcal{C} is a *direct sum* of objects X_1, \dots, X_n if there is a family of morphisms $p_i : D \rightarrow X_i, q_i : X_i \rightarrow D$, such that

$$\text{id}_D = \sum q_i p_i, p_i q_j = \delta_{ij} \text{id}_{X_i}, \forall i, j.$$

An object X of \mathcal{C} is *simple* if $\text{End}_{\mathcal{C}}(X)$ is isomorphic to \mathbb{k} as a \mathbb{k} -module.

A *monoidal \mathbb{k} -category* is a \mathbb{k} -category which is monoidal and such that monoidal product of morphisms is \mathbb{k} -bilinear. It is *semisimple* if any object is isomorphic to a direct sum of simple ones.

A monoidal category has (*left*) *duality* if for any object X there is an object X^* (the *left dual* of X) and

$$b_X = \text{coev}_X : \mathbb{1} \rightarrow X \otimes X^*, d_X = \text{ev}_X : X^* \otimes X \rightarrow \mathbb{1}$$

such that

$$(\text{id}_X \otimes d_X)(b_X \otimes \text{id}_X) = \text{id}_X, (d_X \otimes \text{id}_{X^*})(\text{id}_{X^*} \otimes b_X) = \text{id}_{X^*}.$$

A *left rigid category* is a monoidal category with a left duality. A *rigid category* is a monoidal category which is left rigid and right rigid. Here a right duality is similarly defined using

$$\widetilde{\text{ev}}_X : X \otimes X^* \rightarrow \mathbb{1}, \quad \widetilde{\text{coev}}_X : \mathbb{1} \rightarrow X^* \otimes X.$$

For $f \in \text{Hom}(X, Y)$, let

$$f^* = (d_Y \otimes \text{id}_{X^*}) (\text{id}_{Y^*} \otimes f \otimes \text{id}_{X^*}) (\text{id}_{Y^*} \otimes b_X).$$

A *braiding* in a monoidal category is a collection of natural isomorphisms

$$c_{X,Y} : X \otimes Y \cong Y \otimes X$$

such that

$$c_{X,Y \otimes Z} = (\text{id}_Y \otimes c_{X,Z}) (c_{X,Y} \otimes \text{id}_Z), \quad c_{X \otimes Y, Z} = (c_{X,Z} \otimes \text{id}_Y) (\text{id}_X \otimes c_{Y,Z}).$$

A braiding is *symmetric* if for all X, Y , $c_{Y,X} c_{X,Y} = \text{id}$. In this case the category is *symmetric*. A *braided functor* between braided categories (\mathcal{C}, c) and (\mathcal{C}', c') is a monoidal functor $F : \mathcal{C} \rightarrow \mathcal{C}'$ such that for all $X, Y \in \text{Ob}(\mathcal{C})$,

$$F_2(Y, X) c'_{F(X), F(Y)} = F(c_{X,Y}) F_2(X, Y).$$

A *symmetric functor* is a braided monoidal functor between symmetric categories.

A *twist* in a braided monoidal category is a natural family of isomorphisms $\theta_X : X \cong X$, such that

$$\theta_{X \otimes Y} = c_{Y,X} c_{X,Y} (\theta_X \otimes \theta_Y).$$

A *ribbon category* is a braided monoidal category with a twist and a left duality such that $\theta_{X^*} = (\theta_X)^*$.

A *pivotal category* \mathcal{C} is a rigid category with distinguished duality such that the induced left and right dual functors coincide as monoidal functors. The *left trace* of an endomorphism f of an object X is

$$\text{tr}_l(f) = \text{ev}_X (\text{id}_{X^*} \otimes f) \widetilde{\text{coev}}_X \in \text{End}_{\mathcal{C}}(\mathbb{1}).$$

The *right trace* of f is

$$\text{tr}_r(f) = \widetilde{\text{ev}}_X (f \otimes \text{id}_{X^*}) \text{coev}_X \in \text{End}_{\mathcal{C}}(\mathbb{1}).$$

The *left dimension* of an object X is $\dim_l(X) = \text{tr}_l(\text{id}_X)$. The *right dimension* is $\dim_r(X) = \text{tr}_r(\text{id}_X)$.

A *spherical category* is a pivotal category whose left and right traces coincide. All ribbon categories are spherical.

A *pre-fusion \mathbb{k} -category* is a monoidal \mathbb{k} -category \mathcal{C} such that there is a set I of simple objects of \mathcal{C} such that $\mathbb{1} \in I$, $\text{Hom}_{\mathcal{C}}(i, j) = 0$ for any distinct $i, j \in I$, and every object of \mathcal{C} is a direct sum of a finite family of elements of I .

A *fusion \mathbb{k} -category* is a rigid pre-fusion \mathbb{k} -category which has only a finite number of isomorphism classes of simple objects.

The *dimension* of a pivotal fusion \mathbb{k} -category \mathcal{C} is defined by

$$\dim(\mathcal{C}) = \sum_{i \in I} \dim_l(i) \dim_r(i) \in \mathbb{k}.$$

Let \mathcal{C} be a ribbon fusion \mathbb{k} -category. For any $i, j \in I$, set

$$S_{i,j} = \text{tr}(c_{j,i}c_{i,j}) \in \text{End}_{\mathcal{C}}(\mathbb{1}) = \mathbb{k}.$$

The category \mathcal{C} is said to be *modular* if this S -matrix is invertible over \mathbb{k} . The modularity of \mathcal{C} implies that $\dim(\mathcal{C})$ is invertible in \mathbb{k} .

A *Hermitian fusion category* is a spherical fusion category \mathcal{C} over \mathbb{C} endowed with antilinear homomorphisms

$$\{f \in \text{Hom}_{\mathcal{C}}(X, Y) \mapsto \bar{f} \in \text{Hom}_{\mathcal{C}}(Y, X)\}_{X, Y \in \text{Ob}(\mathcal{C})}$$

such that $\bar{\bar{f}} = f, \overline{gf} = \bar{f}\bar{g}, \overline{f \otimes g} = \bar{f} \otimes \bar{g}, \overline{\text{ev}_X} = \widetilde{\text{coev}_X}, \overline{\text{coev}_X} = \widetilde{\text{ev}_X}$.

A *unitary fusion category* is a Hermitian fusion category \mathcal{C} such that $\text{tr}(f\bar{f}) > 0$ for any non-zero morphism f in \mathcal{C} . The dimension of a unitary fusion category is a positive real number.

2.2 Representations of $U_q(\mathfrak{sl}_2)$

We will focus on the case where $\mathfrak{g} = \mathfrak{sl}_2, q = e^{\pi i/r}, r \geq 3$. We follow [BK01].

Definition 1. A *Hopf algebra* A over a field \mathbb{k} is a bialgebra with an algebra anti-isomorphism $\gamma : A \rightarrow A$, called the antipode, satisfying $\mu(\text{id} \otimes \gamma)\Delta = \epsilon = \mu(\gamma \otimes \text{id})\Delta$.

The category of finite-dimensional representations of A is a rigid monoidal category.

Let \mathfrak{g} be a finite-dimensional simple Lie algebra over \mathbb{C} , and \mathfrak{h} be its Cartan subalgebra. Let $\Delta \subset \mathfrak{h}^*$ be the root system, $\Pi = \{\alpha_1, \dots, \alpha_r\} \subset \Delta$ be the set of simple roots, $h_i = \alpha_i^\vee \in \mathfrak{h}$ be the coroots, and $A = (a_{ij})$ be the Cartan matrix, $a_{ij} = (\alpha_i^\vee, \alpha_j)$.

Let $P \subset \mathfrak{h}^*$ be the weight lattice, $Q \subset \mathfrak{h}^*$ be the root lattice, and $Q^\vee \subset \mathfrak{h}$ be the dual root lattice (coroot lattice). Let $\langle \langle \cdot, \cdot \rangle \rangle$ be an invariant bilinear form on \mathfrak{g} normalized by $\langle \langle \alpha, \alpha \rangle \rangle = 2$ for short roots α . Then $d_i := \langle \langle \alpha_i, \alpha_i \rangle \rangle / 2 \in \mathbb{Z}_+$ for all $i = 1, \dots, r$. Let \mathbb{C}_q be the field $\mathbb{C}(q^{1/|P/Q|})$ where q is a formal variable.

Definition 2. The quantum group $U_q(\mathfrak{g})$ is the associative algebra over \mathbb{C}_q with generators $e_i, f_i (i = 1, \dots, r), q^h (h \in Q^\vee)$ and relations

$$q^{h'} q^{h''} = q^{h'+h''}, q^0 = 1, \quad h', h'' \in Q^\vee,$$

$$q^h e_i q^{-h} = q^{\langle h, \alpha_i \rangle} e_i, q^h f_i q^{-h} = q^{-\langle h, \alpha_i \rangle} f_i, [e_i, f_j] = \delta_{ij} \frac{q^{d_i h_i} - q^{-d_i h_i}}{q^{d_i} - q^{-d_i}},$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_i e_i^{1-a_{ij}-k} e_j e_i^k = 0, \quad i \neq j,$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_i f_i^{1-a_{ij}-k} f_j f_i^k = 0, \quad i \neq j,$$

where

$$\begin{bmatrix} n \\ k \end{bmatrix}_i := \frac{[n]_i!}{[k]_i! [n-k]_i!}, \quad [n]_i! := [1]_i [2]_i \cdots [n]_i, \quad [n]_i := \frac{q^{d_i n} - q^{-d_i n}}{q^{d_i} - q^{-d_i}}.$$

This is a Hopf algebra with

$$\begin{aligned} \Delta(q^h) &= q^h \otimes q^h, \\ \Delta(e_i) &= e_i \otimes q^{d_i h_i} + 1 \otimes e_i, \\ \Delta(f_i) &= f_i \otimes 1 + q^{-d_i h_i} \otimes f_i, \\ \epsilon(q^h) &= 1, \quad \epsilon(e_i) = \epsilon(f_i) = 0, \\ \gamma(q^h) &= q^{-h}, \quad \gamma(e_i) = -e_i q^{-d_i h_i}, \quad \gamma(f_i) = -q^{d_i h_i} f_i. \end{aligned}$$

Let $\mathcal{A} = \mathbb{Z}[q^{\pm 1/|P/Q|}]$ and let $U_q(\mathfrak{g})_{\mathbb{Z}}$ be the \mathcal{A} -subalgebra of $U_q(\mathfrak{g})$ generated by the elements

$$e_i^{(n)} = \frac{e_i^n}{[n]_i!}, \quad f_i^{(n)} = \frac{f_i^n}{[n]_i!}, \quad q^h, \quad n = 1, 2, \dots, i = 1, \dots, r, h \in Q^{\vee}.$$

Fix $\varkappa \in \mathbb{C}^{\times}$ and consider \mathbb{C} as an \mathcal{A} -module via the homomorphism

$$\mathcal{A} \rightarrow \mathbb{C}, \quad q^a \mapsto e^{\frac{a \pi i}{m \varkappa}}, \quad m := \max d_i.$$

Set

$$U_q(\mathfrak{g})|_{q=e^{\frac{\pi i}{m \varkappa}}} := U_q(\mathfrak{g})_{\mathbb{Z}} \otimes_{\mathcal{A}} \mathbb{C}.$$

We abbreviate $U_q(\mathfrak{g})|_{q=e^{\frac{\pi i}{m \varkappa}}}$ to $U_q(\mathfrak{g})$.

The category $\mathcal{C}(\mathfrak{g}, \varkappa)$ of finite-dimensional representations of $U_q(\mathfrak{g})$ over \mathbb{C} possessing a weight decomposition

$$V = \bigoplus_{\lambda \in P} V^{\lambda}, \quad q^h|_{V^{\lambda}} = q^{(h, \lambda)} \text{id}_{V^{\lambda}},$$

$$e_i^{(n)}(V^{\lambda}) \subset V^{\lambda + n \alpha_i}, \quad f_i^{(n)}(V^{\lambda}) \subset V^{\lambda - n \alpha_i},$$

is a ribbon category.

The Weyl modules of $U_q(\mathfrak{sl}_2)$ are $V_n = \bigoplus_{i=0}^n \mathbb{C} v_i$, $n \in \mathbb{Z}_+$, where v_0 is the highest weight vector and $v_i = f^{(i)} v_0$. The action is

$$q^h v_i = q^{n-2i} v_i, \quad e v_i = [n-i+1] v_{i-1}, \quad f v_i = [i+1] v_{i+1}.$$

The module V_n is irreducible for $n < \varkappa$, and $\dim_q V_n = [n+1] = 0$ if and only if \varkappa divides $n+1$. Hence for $0 \leq n \leq \varkappa - 2$, V_n is irreducible and $\dim_q V_n \neq 0$. Correspondingly, let $C := \{0, \frac{1}{2}, \dots, \frac{\varkappa-2}{2}\}$ be the set of weights, where $\mathfrak{g} = \mathfrak{sl}_2$, $q = e^{\pi i / \varkappa}$, $\varkappa \geq 3$. From now on we index the Weyl modules by their weights in $\frac{1}{2}\mathbb{Z}$.

A module T over $U_q(\mathfrak{sl}_2)$ is called *tilting* if both T and T^* have composition series with factors Weyl modules. Let \mathcal{T} be the full subcategory of $\mathcal{C}(\mathfrak{sl}_2, \varkappa)$ consisting of all tilting modules. If $\lambda \in C$ then the module V_{λ} is tilting. It is shown in [HAP95] that the category of tilting modules \mathcal{T} is closed under $*$, \oplus , \otimes and direct summands. It is a ribbon category.

A tilting module T is called *negligible* if $\text{tr}_q f = 0$ for any $f \in \text{End} T$. A morphism $f : T_1 \rightarrow T_2$ is called *negligible* if $\text{tr}_q(fg) = 0$ for all $g : T_2 \rightarrow T_1$.

Let $\mathcal{C}^{\text{int}} := \mathcal{C}^{\text{int}}(\mathfrak{sl}_2, \varkappa)$ be the category with objects tilting modules and morphisms

$$\text{Hom}_{\mathcal{C}^{\text{int}}}(V, W) = \text{Hom}_{\mathcal{T}}(V, W)/\text{negligible morphisms}.$$

Then \mathcal{C}^{int} is a ribbon category. Any object V in \mathcal{C}^{int} is isomorphic to $\bigoplus_{\lambda \in C} n_{\lambda} V_{\lambda}$. \mathcal{C}^{int} is a semisimple abelian category. Indeed, \mathcal{C}^{int} is a modular tensor category with simple objects $V_{\lambda} (\lambda \in C)$.

2.3 Unitarity of the category

For a survey on the general case, see [Row06]. We will focus on the case where $\mathfrak{g} = \mathfrak{sl}_2$, $q = e^{\pi i/r}$, $r \geq 3$.

The invertibility of S -matrix is shown in [TW93]. Their work is based on the notion of quasimodular Hopf algebra, which is a weakened version of modular Hopf algebra introduced in the paper [RT91].

In [KJ96], Hermitian structure on the category is defined in Section 4. Note that the notation in [KJ96] is different. In [Wen98], positivity is proved by Wenzl. Xu [Xu98] independently showed some of the cases covered by Wenzl.

Theorem 2 ([Wen98], [Xu98]). *The categories $\mathcal{C}^{\text{int}}(\mathfrak{sl}_2, \varkappa)$ are unitary when $q = e^{\pi i/r}$.*

3 Turaev–Viro TQFT

3.1 TQFT

Manifolds considered are assumed to be oriented.

An n -dimensional cobordism is a quadruple $(M, \Sigma_0, \Sigma_1, h)$, where M is a compact oriented n -manifold, Σ_0, Σ_1 are closed oriented $(n-1)$ -manifolds, and $h : (-\Sigma_0) \sqcup \Sigma_1 \rightarrow \partial M$ is an orientation preserving homeomorphism. Two cobordisms $(M, \Sigma_0, \Sigma_1, h)$ and $(M', \Sigma_0, \Sigma_1, h')$ between Σ_0 and Σ_1 are *homeomorphic* if there is an orientation preserving homeomorphism $\alpha : M \rightarrow M'$ such that $h' = \alpha h$.

Let Cob_3 be the category with objects closed oriented 2-manifolds and morphisms cobordisms between them considered up to homeomorphisms. It is a symmetric monoidal category, where the monoidal product is the ordered disjoint union of manifolds and cobordisms. Each closed oriented 3-manifold represents a morphism $\emptyset \rightarrow \emptyset$.

Let $\text{Mod}_{\mathbb{k}}$ be the symmetric monoidal category of \mathbb{k} -modules.

Definition 3. A 3-dimensional *Topological Quantum Field Theory* (TQFT) is a symmetric strong monoidal functor

$$Z : \text{Cob}_3 \rightarrow \text{Mod}_{\mathbb{k}}.$$

A *conjugation* in a monoidal category \mathcal{C} is a family of maps

$$\{f \in \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \bar{f} \in \text{Hom}_{\mathcal{C}}(Y, X)\}_{X, Y \in \text{Ob}(\mathcal{C})}$$

compatible with the associativity and unitality constraints of \mathcal{C} , such that for all morphisms f, g in \mathcal{C} and all $X \in \text{Ob}(\mathcal{C})$

$$\overline{gf} = \bar{f}\bar{g}, \quad \overline{\text{id}_X} = \text{id}_X, \quad \bar{\bar{f}} = f, \quad \overline{f \otimes g} = \bar{f} \otimes \bar{g}.$$

The category Cob_3 has a conjugation

$$(M, \Sigma_0, \Sigma_1, h : (-\Sigma_0) \sqcup \Sigma_1 \rightarrow \partial M) \\ \mapsto (-M, \Sigma_1, \Sigma_0, hP : (-\Sigma_1) \sqcup \Sigma_0 \rightarrow -\partial M = \partial(-M)),$$

where P is the orientation reversing permutation homeomorphism

$$(-\Sigma_1) \sqcup \Sigma_0 \simeq (-\Sigma_0) \sqcup \Sigma_1.$$

Let Hilb be the category of finite-dimensional Hilbert spaces over \mathbb{C} and \mathbb{C} -linear homomorphisms between them. This is a symmetric ribbon \mathbb{C} -category. The conjugation is given by assigning to every morphism its hermitian adjoint.

Definition 4. A *unitary 3-dimensional TQFT* is a symmetric strong monoidal functor

$$Z : \text{Cob}_3 \rightarrow \text{Hilb}$$

commuting with the conjugations in Cob_3 and in Hilb .

Note that there is a forgetful functor $\text{Hilb} \rightarrow \text{Mod}_{\mathbb{C}}$. Composing such a functor Z with the forgetful functor, we obtain a 3-dimensional TQFT. We call Z a *unitary lift* of the latter TQFT.

The category Cob_3 has a subcategory isomorphic to the category of closed oriented surfaces and isotopy classes of orientation-preserving homeomorphisms. Any such homeomorphism $f : \Sigma \rightarrow \Sigma'$ determines a morphism

$$(C = \Sigma' \times [0, 1], \Sigma, \Sigma', h : (-\Sigma) \sqcup \Sigma' \simeq \partial C)$$

in Cob_3 , where $h(x) = (f(x), 0)$ for $x \in \Sigma$ and $h(x') = (x', 1)$ for $x' \in \Sigma'$. Restricting a TQFT $Z : \text{Cob}_3 \rightarrow \text{Mod}_{\mathbb{k}}$ to this subcategory, we obtain the action of homeomorphisms, and in particular the action of mapping class groups. We denote the induced map by $f_{\#} : Z(\Sigma) \rightarrow Z(\Sigma')$. It follows that $(f^{-1})_{\#} = (f_{\#})^{-1}$.

If, in addition, Z is unitary, then

Proposition 3. $\langle f_{\#}x, f_{\#}y \rangle = \langle x, y \rangle$.

Proof. The adjoint of $f_{\#}$ is the same as the inverse of $f_{\#}$. Indeed there is a homeomorphism from the conjugation of $(C = \Sigma' \times [0, 1], \Sigma, \Sigma', h : (-\Sigma) \sqcup \Sigma' \simeq \partial C)$ to the cobordism associated to f^{-1} . \square

Theorem 10.3.1 in [Tur10] is a similar result.

3.2 Construction of Turaev–Viro state sum TQFT

We sketch a construction following [TV17] and [TV13].

Fix a spherical fusion \mathbb{k} -category \mathcal{C} of invertible dimension. Fix a (finite) representative set I of simple objects of \mathcal{C} .

3.2.1 Colored graphs in surfaces

A *graph* G in an oriented surface Σ is a finite graph without isolated vertices embedded in Σ . A graph G is \mathcal{C} -*colored*, if each edge of G is oriented and endowed with an object of \mathcal{C} called the *color* of the edge.

For a vertex v of a \mathcal{C} -colored graph G , the set E_v of half-edges of G incident to v with cyclic order induced by the opposite orientation of Σ , and the color $c = c_v : E_v \rightarrow \text{Ob}(\mathcal{C})$, the orientation $\epsilon = \epsilon_v : E_v \rightarrow \{+, -\}$, together form a *cyclic \mathcal{C} -set*. The sign is $+$ if $e \in E_v$ is oriented towards v .

Starting from $e = e_1 \in E_v$, let

$$H_e = \text{Hom}_{\mathcal{C}} \left(\mathbb{1}, c(e_1)^{\epsilon(e_1)} \otimes \cdots \otimes c(e_n)^{\epsilon(e_n)} \right),$$

where $X^+ = X$, $X^- = X^*$.

This module can be made into the *symmetrized multiplicity module* $H(E_v)$, independent of the starting point chosen.

Set $H_v(G) = H(E_v)$ and

$$H(G) = \otimes_v H_v(G).$$

Orient the plane \mathbb{R}^2 counterclockwise. Let G be a \mathcal{C} -colored graph in \mathbb{R}^2 . Pick any $\alpha_v \in H_v(G)$ and replace v by a box colored with the corresponding element in the Hom set. This transforms G into a planar diagram which determines, by the graphical calculus, an element $\mathbb{F}_{\mathcal{C}}(G) (\otimes_v \alpha_v)$ of $\text{End}_{\mathcal{C}}(\mathbb{1}) = \mathbb{k}$. See [Tur10], [TV17] or [TV13] for an introduction to the graphical calculus. This procedure defines a vector

$$\mathbb{F}_{\mathcal{C}}(G) \in H(G)^* = \text{Hom}_{\mathbb{k}}(H(G), \mathbb{k}).$$

As \mathcal{C} is spherical, the invariant $\mathbb{F}_{\mathcal{C}}$ generalizes to graphs in the 2-sphere.

3.2.2 State sums on skeletons of 3-manifolds

A *2-polyhedron* is a compact topological space that can be triangulated using a finite number of simplices of dimension ≤ 2 so that all 0-simplices and 1-simplices are faces of 2-simplices. For a 2-polyhedron P , let $\text{Int}(P)$ be the subspace consisting of all points having a neighborhood homeomorphic to \mathbb{R}^2 .

A *stratification* of a 2-polyhedron P is a graph G embedded in P so that $P \setminus \text{Int}(P) \subset G$. The vertices and edges of $P^{(1)} := G$ are called respectively the vertices and edges of P .

Cutting a stratified 2-polyhedron P along the graph $P^{(1)} \subset P$, we obtain a compact surface \tilde{P} with interior $P \setminus P^{(1)}$. The (connected) components of \tilde{P} are called the *regions* of P . We let $\text{Reg}(P)$ be the (finite) set of all regions of P .

A *branch* of a stratified 2-polyhedron P at a vertex x is a germ at x of an adjacent region. A *branch* of P at an edge e is a germ at e of an adjacent region. The set of branches of P at e is denoted P_e . The number of elements of P_e is called the *valence* of e .

The edges of P of valence 1 and their vertices form a graph called the *boundary* of P and denoted ∂P . We say that P is *orientable* (resp. *oriented*) if all regions of P are orientable (resp. oriented).

A *skeleton* of a closed 3-manifold M is an oriented stratified 2-polyhedron $P \subset M$ such that $\partial P = \emptyset$ and $M \setminus P$ is a disjoint union of open 3-balls.

Any vertex x of a skeleton $P \subset M$ has a closed ball neighborhood $B_x \subset M$ such that $\Gamma_x = P \cap \partial B_x$ is a finite non-empty graph and $P \cap B_x$ is the cone over Γ_x . The pair $(\partial B_x, \Gamma_x)$ is the *link* of x in (M, P) . If M is oriented, then we endow ∂B_x with orientation induced by that of M restricted to $M \setminus \text{Int}(B_x)$.

Let M be a closed oriented 3-manifold. Fix a skeleton P of M . For a map $c : \text{Reg}(P) \rightarrow I$ and an oriented edge e of P , we define a \mathbb{k} -module

$$H_c(e) = H(P_e),$$

where P_e is the cyclic \mathcal{C} -set of branches of P at e . If e^{op} is the same edge with opposite orientation, then there is a contraction

$$*_e : H_c(e)^* \otimes H_c(e^{\text{op}})^* \rightarrow \mathbb{k}.$$

The link of a vertex $x \in P$ determines a \mathcal{C} -colored graph Γ_x in $\partial B_x \simeq S^2$. Hence there is a tensor $\mathbb{F}_{\mathcal{C}}(\Gamma_x) \in H_c(\Gamma_x)^*$. Note that $H_c(\Gamma_x) = \otimes_e H_c(e)$, where e runs over all edges of P incident to x and oriented away from x . Hence $\otimes_x \mathbb{F}_{\mathcal{C}}(\Gamma_x) \in \otimes_e H_c(e)^*$, where e runs over all oriented edges of P . Set $*_P = \otimes_e *_e : \otimes_e H_c(e)^* \rightarrow \mathbb{k}$.

Define

$$|M|_{\mathcal{C}} = (\dim(\mathcal{C}))^{-|P|} \sum_c \left(\prod_{r \in \text{Reg}(P)} (\dim c(r))^{\chi(r)} \right) *_P (\otimes_x \mathbb{F}_{\mathcal{C}}(\Gamma_x)) \in \mathbb{k},$$

where $|P|$ is the number of components of $M \setminus P$, c runs over all maps $\text{Reg}(P) \rightarrow I$, and $\chi(r)$ is the Euler characteristic of r .

Theorem 4 ([TV13], [TV17]). *The invariant $|M|_{\mathcal{C}}$ is independent of P chosen.*

3.2.3 The state sum TQFT

Let M be a compact 3-manifold with boundary. Let G be an oriented graph in ∂M such that all vertices of G have valence ≥ 2 . A *skeleton* of the pair (M, G) is an oriented stratified 2-polyhedron $P \subset M$ satisfying certain requirements including $P \cap \partial M = \partial P = G$.

For any compact oriented 3-manifold M and any I -colored graph G in ∂M , we define a topological invariant $|M, G| \in \mathbb{k}$ as follows. Pick a skeleton $P \subset M$ of the pair (M, G) . Pick a map $c : \text{Reg}(P) \rightarrow I$ extending the coloring of G .

Let E_0 be the set of oriented edges of P with both endpoints in $\text{Int}(M)$, and let E_{∂} be the set of edges of P with exactly one endpoint in ∂M oriented towards this endpoint. We have

$$\otimes_{e \in E_{\partial}} H_c(e)^* = \otimes_v H_v(G^{\text{op}}; -\partial M)^* = H(G^{\text{op}}; -\partial M)^*,$$

where G^{op} in $-\Sigma$ is obtained by reversing orientation in all edges of G and in Σ .

There is a contraction homomorphism

$$*_P : \otimes_{e \in E_0 \cup E_{\partial}} H_c(e)^* \rightarrow \otimes_{e \in E_{\partial}} H_c(e)^* = H(G^{\text{op}}; -\partial M)^*.$$

The tensor product $\otimes_x \mathbb{F}_{\mathcal{C}}(\Gamma_x)$ over all vertices x of P lying in $\text{Int}(M)$ is a vector in $\otimes_{e \in E_0 \cup E_{\partial}} H_c(e)^*$.

Set

$$|M, G| = (\dim(\mathcal{C}))^{-|P|} \sum_c \left(\prod_{r \in \text{Reg}(P)} (\dim c(r))^{\chi(r)} \right) *_P (\otimes_x \mathbb{F}_{\mathcal{C}}(\Gamma_x)),$$

where $|P|$ is the number of components of $M \setminus P$, c runs over all maps $\text{Reg}(P) \rightarrow I$ extending the coloring of G .

Theorem 5 ([TV13], [TV17]). $|M, G| \in H(G^{\text{op}}; -\partial M)^*$ does not depend on the choice of P .

Let Σ_0, Σ_1 be closed oriented surfaces, and let $f : \Sigma_0 \rightarrow \Sigma_1$ be a morphism in Cob_3 represented by a pair (M, h) . For I -colored graphs $G_0 \subset \Sigma_0$ and $G_1 \subset \Sigma_1$,

$$|M, h(G_0^{\text{op}} \cup G_1)|_{\mathcal{C}} \in H(h(G_0^{\text{op}} \cup G_1); \partial M) \simeq \text{Hom}_{\mathbb{k}}(H(G_0), H(G_1)).$$

Set

$$|f, G_0, G_1| = \frac{(\dim(\mathcal{C}))^{|\Sigma_1 \setminus G_1|}}{\dim(G_1)} |M, h(G_0^{\text{op}} \cup G_1)|_{\mathcal{C}} : H(G_0) \rightarrow H(G_1),$$

where $|\Sigma_1 \setminus G_1|$ is the number of components of $\Sigma_1 \setminus G_1$ and $\dim(G_1)$ is the product over all edges of G_1 of the dimensions of their colors.

A *skeleton* of a closed surface Σ is an oriented graph G embedded in Σ such that all vertices of G have valence ≥ 2 and all components of $\Sigma \setminus G$ are open disks. We let $\text{col}(G)$ be the set of all maps from the set of edges of G to I .

For a closed oriented surface Σ and a skeleton $G \subset \Sigma$, consider the \mathbb{k} -module

$$|\Sigma, G|^{\circ} = \bigoplus_{c \in \text{col}(G)} H((G, c); \Sigma).$$

For a morphism $f : \Sigma_0 \rightarrow \Sigma_1$ in Cob_3 and skeletons $G_0 \subset \Sigma_0$ and $G_1 \subset \Sigma_1$, consider the \mathbb{k} -linear homomorphism

$$|f, G_0, G_1|^{\circ} : |\Sigma_0, G_0|^{\circ} \rightarrow |\Sigma_1, G_1|^{\circ}$$

whose restriction to every summand $H((G_0, c_0); \Sigma_0)$ of $|\Sigma_0, G_0|^{\circ}$ is equal to $\sum_{c_1 \in \text{col}(G_1)} |f, (G_0, c_0), (G_1, c_1)|$.

For any skeletons G, G' of a closed oriented surface Σ , set

$$p(G, G') = |\text{id}_{\Sigma}, G, G'|^{\circ} : |\Sigma, G|^{\circ} \rightarrow |\Sigma, G'|^{\circ},$$

$$|\Sigma, G| = \text{Im}(p(G, G')).$$

The \mathbb{k} -linear homomorphism $p(G, G')$ restricts to an isomorphism $|\Sigma, G| \rightarrow |\Sigma, G'|$ and the family $(\{|\Sigma, G|\}_G, \{p(G, G')\}_{G, G'})$ is a projective system. The projective limit

$$|\Sigma| = \varprojlim |\Sigma, G|$$

is a \mathbb{k} -module depending only on Σ .

We associate with each morphism $f : \Sigma_0 \rightarrow \Sigma_1$ in Cob_3 a homomorphism $|f| : |\Sigma_0| \rightarrow |\Sigma_1|$. For each skeleton G of Σ , we have a \mathbb{k} -linear cone isomorphism $\tau_G : |\Sigma| \rightarrow |\Sigma, G|$. Pick any skeletons $G_0 \subset \Sigma_0$ and $G_1 \subset \Sigma_1$. Set

$$|f| = \tau_{G_1}^{-1} \circ |f, G_0, G_1|^\circ \circ \tau_{G_0} : |\Sigma_0| \rightarrow |\Sigma_1|.$$

This homomorphism does not depend on the choice of G_0 and G_1 .

Theorem 6 ([TV13], [TV17]). *The functor $|\cdot| = |\cdot|_{\mathcal{C}}$ is a 3-dimensional TQFT.*

3.3 Unitarity

Let \mathbb{k} be an algebraically closed field. Let \mathcal{C} be a unitary fusion category (over \mathbb{C}). Then $\dim(\mathcal{C}) \neq 0$.

Theorem 7 (Theorem G.1 in [TV17], Corollary 11.6 in [TV13]). *The state sum TQFT $|\cdot|_{\mathcal{C}}$ has a unitary lift.*

The proof is involved. Recall the construction of the Reshetikhin–Turaev extended TQFT $\tau_{\mathcal{B}}$ associated to a modular category \mathcal{B} over \mathbb{k} equipped with a distinguished square root of $\dim(\mathcal{B})$, described in detail in [Tur10]. In this case unitarity is proved in [Tur10], Chapter IV, section 11.

The center $\mathcal{Z}(\mathcal{C})$ of \mathcal{C} has a subcategory the unitary center $\mathcal{Z}^u(\mathcal{C})$ of \mathcal{C} , such that the inclusion is an equivalence of categories. Theorem 17.1 in [TV17] implies that

$$|\cdot|_{\mathcal{C}} \simeq \tau_{\mathcal{Z}(\mathcal{C})} \simeq \tau_{\mathcal{Z}^u(\mathcal{C})}.$$

Since $\mathcal{Z}^u(\mathcal{C})$ is an anomaly-free unitary modular category, these TQFTs are unitary.

In this section we introduce the idea of the proof, following [TV13].

3.3.1 Extending to graph TQFTs

The goal of this subsection is Theorem 11.1 and Theorem 11.2 in [TV13]:

Theorem 8 ([TV13]). *Let \mathcal{C} be a spherical fusion category over an algebraically closed field \mathbb{k} such that $\dim \mathcal{C} \neq 0$. Then $|M|_{\mathcal{C}} = \tau_{\mathcal{Z}(\mathcal{C})}(M)$ for any closed oriented 3-manifold M . Moreover the TQFTs $|\cdot|_{\mathcal{C}}$ and $\tau_{\mathcal{Z}(\mathcal{C})}$ are isomorphic.*

Let \mathcal{C} be a monoidal category. A *half braiding* is a pair (A, σ) , where $A \in \text{Ob}(\mathcal{C})$ and $\sigma = \{\sigma_X : A \otimes X \rightarrow X \otimes A\}_{X \in \text{Ob}(\mathcal{C})}$ is a natural isomorphism such that $\sigma_{X \otimes Y} = (\text{id}_X \otimes \sigma_Y)(\sigma_X \otimes \text{id}_Y)$ for all $X, Y \in \text{Ob}(\mathcal{C})$.

The *center* of \mathcal{C} is the braided category $\mathcal{Z}(\mathcal{C})$ whose objects are half braidings. A morphism $(A, \sigma) \rightarrow (A', \sigma')$ in $\mathcal{Z}(\mathcal{C})$ is a morphism $f : A \rightarrow A'$ such that $(\text{id}_X \otimes f)\sigma_X = \sigma'_X(f \otimes \text{id}_X)$ for all $X \in \text{Ob}(\mathcal{C})$. There is a forgetful functor.

The following theorem is Theorem 1.2 and Proposition 5.18 in [Mü03].

Theorem 9 ([Mü03]). *Let \mathcal{C} be a spherical fusion \mathbb{k} -category such that $\dim \mathcal{C} \neq 0$. Then $\mathcal{Z}(\mathcal{C})$ is an anomaly free modular category with $\Delta_+ = \Delta_- = \dim \mathcal{C}$, $\dim \mathcal{Z}(\mathcal{C}) = \Delta_+ \Delta_- = (\dim \mathcal{C})^2$.*

Hence it makes sense to talk about the Reshetikhin–Turaev extended TQFT.

It is necessary to extend the definition of a TQFT. For any category \mathcal{B} , we define a category $\mathcal{L}_{\mathcal{B}}$ of 3-cobordisms with \mathcal{B} -colored framed oriented links inside. The links may be empty so that the category Cob_3 is a subcategory of $\mathcal{L}_{\mathcal{B}}$.

By a *link TQFT* we mean a symmetric monoidal functor $Z : \mathcal{L}_{\mathcal{B}} \rightarrow \text{Mod}_{\mathbb{k}}$.

For a category \mathcal{B} , we define a category $\mathcal{G}_{\mathcal{B}}$ of 3-cobordisms with \mathcal{B} -colored ribbon graphs inside, see [Tur10]. Here we consider only ribbon graphs disjoint from the bases of cobordisms.

By a *graph TQFT* we mean a symmetric monoidal functor $\mathcal{G}_{\mathcal{B}} \rightarrow \text{Mod}_{\mathbb{k}}$.

The TQFT $\tau_{\mathcal{Z}(\mathcal{C})}$ extends to a graph TQFT still denoted $\tau_{\mathcal{Z}(\mathcal{C})}$, see [Tur10]. This TQFT is *non-degenerate*, i.e. the vector space $\tau_{\mathcal{Z}(\mathcal{C})}(\Sigma)$ is generated by the vectors $\tau_{\mathcal{Z}(\mathcal{C})}(M, \emptyset, \Sigma)(1)$, where M runs over all compact oriented 3-manifolds with $\mathcal{Z}(\mathcal{C})$ -colored ribbon graphs inside and with $\partial M = \Sigma$.

The TQFT $|\cdot|_{\mathcal{C}}$ also can be extended to a graph TQFT. We discuss the extension to a link TQFT, which is similar.

Let M be a compact oriented 3-manifold, and P be a skeleton. A link in M can be presented by possibly intersecting circles in P in general position, called the *link diagram*. The underlying 4-valent graph may be added to the 1-skeleton of P . An *enriched link diagram* in P is a link diagram with half integers added to record the framing. Two enriched link diagrams in P represent isotopic framed links if and only if these diagrams may be related by a finite sequence of certain moves and ambient isotopies in P .

Consider a pair (M, L) , where M is a compact oriented 3-manifold (with empty graph $G = \emptyset$ in the boundary) and $L = L_1 \sqcup \cdots \sqcup L_N \subset \text{Int}(M)$ is an oriented framed link whose all components are non-trivial and colored by simple objects $J_1 \dots J_N$ of $\mathcal{Z}(\mathcal{C})$. We define $|M, L|_{\mathcal{C}} \in \mathbb{k}$ as follows. Fix a (finite) representative set I of simple objects of \mathcal{C} . Pick a special skeleton (see [TV13]) P of M and an oriented enriched link diagram d in P representing L . After adding d to $P^{(1)}$, we obtain regions.

Pick distinguished square root $\nu_{J_q} \in \mathbb{k}$ of the twist scalar v_{J_q} of J_q . Let $n_q \in \frac{1}{2}\mathbb{Z}$ be the pre-twist of the loop of d representing L_q . Set

$$|M, L|_{\mathcal{C}} = (\dim(\mathcal{C}))^{-|P|} \prod_{q=1}^N \nu_{J_q}^{2n_q} \sum_c \left(\prod_{r \in \text{Reg}(d)} (\dim c(r))^{x(r)} \right) *_P (\otimes_x |x|_{\mathcal{C}}).$$

This is a topological invariant of the pair (M, L) . It extends to the case where links may have trivial components.

It is shown in Theorem 1.2 in [Mü03] that here $\mathcal{Z}(\mathcal{C})$ is semisimple with finitely many simple objects. The invariant extends by linearity to arbitrary colors of the components of L .

We extend the TQFT $|\cdot|_{\mathcal{C}} : \text{Cob}_3 \rightarrow \text{Mod}_{\mathbb{k}}$ to a link TQFT $\mathcal{L}_{\mathcal{Z}(\mathcal{C})} \rightarrow \text{Mod}_{\mathbb{k}}$, and moreover a graph TQFT.

3.3.2 $|M, R|_{\mathcal{C}} = \tau_{\mathcal{Z}(\mathcal{C})}(M, R)$

For a closed (connected) oriented 3-manifold M with $\mathcal{Z}(\mathcal{C})$ -colored ribbon graph R without free ends inside, we claim

Lemma 10. $|M, R|_{\mathcal{C}} = \tau_{\mathcal{Z}(\mathcal{C})}(M, R)$.

Present M by surgery on S^3 along a framed oriented link L . Pick a closed regular neighborhood U of L , and set $E = S^3 \setminus \text{Int}(U)$. We consider $R \subset E$.

Pick a homeomorphism

$$f_L : (S^1 \times S^1)^{\sqcup n} \longrightarrow \partial U = -\partial E$$

which carries the q -th copy of $S^1 \times \{\text{pt}\}$ onto a positively oriented meridian of L_q and carries the q -th copy of $\{\text{pt}\} \times S^1$ onto a positively oriented longitude of L_q determined by the framing.

Define $V = -(S^1 \times D^2)^{\sqcup n}$, $W = (D^2 \times S^1)^{\sqcup n}$. Using f_L , the gluing of W to E yields S^3 , while the gluing of V to E yields M . The *torus vector* of a TQFT Z is the vector $w = w(Z) = Z(V)(1_{\mathbb{k}}) \in Z(\partial V) = Z(S^1 \times S^1)$ where $n = 1$.

There are morphisms

$$Z(f_L) = (f_L)_\# : Z\left((S^1 \times S^1)^{\sqcup n}\right) \rightarrow Z(-\partial E),$$

$$Z(E, R) : Z(-\partial E) \rightarrow \mathbb{k}.$$

Define Z_R^L to be the composition. Hence

$$Z(M, R) = Z_R^L(w^{\otimes n}).$$

We fix a representative set \mathcal{J} of simple objects of $\mathcal{Z}(\mathcal{C})$.

For $X \in \text{Ob}(\mathcal{Z}(\mathcal{C}))$, let

$$K_X = \{0\} \times S^1 \subset \text{Int}(D^2 \times S^1)$$

be the $\mathcal{Z}(\mathcal{C})$ -colored knot with orientation induced by that of S^1 , constant framing, and color X . Set

$$[X] = Z(D^2 \times S^1, K_X)(1_{\mathbb{k}}) \in Z(S^1 \times S^1).$$

Indeed,

Lemma 11. *The vectors $\{[j]\}_{j \in \mathcal{J}}$ form a basis.*

One can explicitly expand w in terms of this basis. To calculate $|M, R|_{\mathcal{C}}$, we need $Z_R^L(w^{\otimes n})$, and it suffices to calculate

$$\begin{aligned} Z_R^L([j_1] \otimes_{\mathbb{k}} \cdots \otimes_{\mathbb{k}} [j_n]) &= Z(S^3, T_{j_1, \dots, j_n}) = (\dim(\mathcal{C}))^{-1} \langle T_{j_1, \dots, j_n} \rangle_{\mathcal{Z}(\mathcal{C})} \\ &\in \text{End}_{\mathcal{Z}(\mathcal{C})}(\mathbb{1}_{\mathcal{Z}(\mathcal{C})}) = \text{End}_{\mathcal{C}}(\mathbb{1}) = \mathbb{k}, \end{aligned}$$

where $j_1, \dots, j_n \in \mathcal{J}$, T_{j_1, \dots, j_n} is the union of R with the link L whose components L_1, \dots, L_n are colored with j_1, \dots, j_n , and $\langle \cdot \rangle_{\mathcal{Z}(\mathcal{C})}$ is the graphical calculus, see Theorem 16.1 in [TV17].

Now Lemma 10 follows from a complete expansion.

3.3.3 Proof of Theorem 8

We claim that there is a monoidal isomorphism of the functors $\tau_{\mathcal{Z}(\mathcal{C})}$ and $|\cdot|_{\mathcal{C}}$ from $\mathcal{G}_{\mathcal{Z}(\mathcal{C})}$ to $\text{Mod}_{\mathbb{k}}$.

Indeed there is a general criterion establishing isomorphism of two (generalized) TQFTs, see [Tur10], Chapter III, Section 3. If

- (1) at least one of the TQFTs is non-degenerate,
- (2) the values of these TQFTs on cobordisms with empty bases are equal, and
- (3) the vector spaces associated with any object have equal dimensions, then these TQFTs are isomorphic.

The graph TQFT $\tau_{\mathcal{Z}(\mathcal{C})}$ is non-degenerate. That $|M, K|_{\mathcal{C}} = \tau_{\mathcal{Z}(\mathcal{C})}(M, K)$ for any $\mathcal{Z}(\mathcal{C})$ -colored ribbon graph K in a closed oriented 3-manifold M is proven. The equality of dimensions is provided by the following lemma. This completes the proof of the theorem.

The following lemma is Lemma 11.3 in [TV13].

Lemma 12 ([TV13]). *The vector spaces $|\Sigma|_{\mathcal{C}}$ and $\tau_{\mathcal{Z}(\mathcal{C})}(\Sigma)$ associated with any closed connected oriented surface Σ have equal dimensions.*

Its proof involves more tools like coends and Hopf monads, see [TV13].

3.3.4 Proof of unitarity

A half braiding (A, σ) of \mathcal{C} is *unitary* if $\overline{\sigma_X} = \sigma_X^{-1}$ for all $X \in \text{Ob}(\mathcal{C})$. The *unitary center* $\mathcal{Z}^u(\mathcal{C})$ of \mathcal{C} is the full subcategory of $\mathcal{Z}(\mathcal{C})$ formed by the unitary half braidings.

The inclusion $\mathcal{Z}^u(\mathcal{C}) \subset \mathcal{Z}(\mathcal{C})$ is a braided equivalence, see Theorem 6.4 in [Mü03]. Therefore $\mathcal{Z}^u(\mathcal{C})$ is also modular. The induced conjugation makes $\mathcal{Z}^u(\mathcal{C})$ a unitary modular category, and hence the TQFT $\tau_{\mathcal{Z}^u(\mathcal{C})}$ is unitary.

Now there are isomorphisms of TQFTs

$$\tau_{\mathcal{Z}^u(\mathcal{C})} \simeq \tau_{\mathcal{Z}(\mathcal{C})} \simeq |\cdot|_{\mathcal{C}}.$$

The first one is induced by the braided equivalence $\mathcal{Z}^u(\mathcal{C}) \simeq \mathcal{Z}(\mathcal{C})$. The unitary structure of $\tau_{\mathcal{Z}^u(\mathcal{C})}$ is transported to $|\cdot|_{\mathcal{C}}$ via the isomorphism.

By Corollary 11.6, Theorem 11.2, and Theorem 11.5 of [Tur10],

Corollary 13 ([TV13]). *If \mathcal{C} is a unitary fusion category, then*

$$||M|_{\mathcal{C}}| \leq (\dim(\mathcal{C}))^{g(M)-1}$$

for any closed oriented 3-manifold M , where $g(M)$ is the Heegaard genus of M .

In the next chapter we give another proof of a similar theorem.

4 Counterexamples to Rank versus Genus Conjecture

4.1 Estimating the Heegaard genus

In this section we estimate the Heegaard genus, using arguments similar to Theorem 2.2 in [Gar98]. This theorem is not correct in general. Note that Theorem 2.2 uses Lemma 2.1, and the author claims that Lemma 2.1 follows

from a paper by Witten [Wit89]. In [Wit89], there is an implicit condition on the TQFT, namely the space associated to S^2 must be one-dimensional. This condition is not mentioned in [Gar98].

However, the estimate is valid in our special case.

Lemma 14 (Lemma 13.6 in [TV17]). *Over any commutative ring \mathbb{k} , the \mathbb{k} -module $|S^2|_{\mathcal{C}}$ is isomorphic to \mathbb{k} .*

The proof is just a direct calculation using definitions.

Lemma 15 (Lemma 10.2 in [TV17]). *Let $Z : \text{Cob}_n \rightarrow \text{Mod}_{\mathbb{k}}$ be an n -dimensional TQFT such that $Z(S^{n-1}) \simeq \mathbb{k}$. Then, for any connected sum $M = M_0 \# M_1$, we have*

$$Z(S^n) Z(M, -\partial M_0, \partial M_1) = Z(M_1, \emptyset, \partial M_1) \circ Z(M_0, -\partial M_0, \emptyset).$$

In particular, if $\partial M_0 = \partial M_1 = \emptyset$, then $Z(S^n) Z(M) = Z(M_0) Z(M_1)$.

Sketch of proof. For $i = 0, 1$, pick a ball $B_i \subset \text{Int}(M_i)$ and set $N_i = M_i \setminus \text{Int}(B_i)$. Let

$$b_0 : S^{n-1} \rightarrow \emptyset, \quad b_1 : \emptyset \rightarrow S^{n-1}, \quad n_0 : -\partial M_0 \rightarrow S^{n-1}, \quad n_1 : S^{n-1} \rightarrow \partial M_1$$

be the corresponding morphisms.

Pick a \mathbb{k} -linear isomorphism $z : Z(S^{n-1}) \rightarrow Z(\emptyset)$. Then $Z(b_0) = \lambda_0 z$ and $Z(b_1) = \lambda_1 z^{-1}$ for some $\lambda_0, \lambda_1 \in \mathbb{k}$. It follows that

$$Z(S^n) \text{id}_{Z(\emptyset)} = Z(S^n, \emptyset, \emptyset) = Z(b_0 \circ b_1) = Z(b_0) Z(b_1) = \lambda_0 \lambda_1 \text{id}_{Z(\emptyset)}.$$

Hence

$$\begin{aligned} & Z(S^n) Z(M, -\partial M_0, \partial M_1) \\ &= \lambda_0 \lambda_1 Z(n_1 \circ n_0) \\ &= \lambda_0 \lambda_1 Z(n_1) \circ Z(n_0) \\ &= Z(n_1) \circ (\lambda_1 z^{-1}) \circ (\lambda_0 z) \circ Z(n_0) \\ &= Z(n_1 \circ b_1) \circ Z(b_0 \circ n_0) \\ &= Z(M_1, \emptyset, \partial M_1) \circ Z(M_0, -\partial M_0, \emptyset). \end{aligned}$$

□

A direct calculation in section 13.1.3 in [TV17] shows that

Lemma 16 ([TV17]). $|S^1 \times S^2|_{\mathcal{C}} = 1$, and $|S^3|_{\mathcal{C}} = (\dim(\mathcal{C}))^{-1}$.

Here, we consider the original Turaev–Viro TQFT [TV92], at $q = e^{\pi i/r}$, $r \geq 3$. Then

$$|S^3| = -\frac{\left(e^{\frac{i\pi}{r}} - e^{-\frac{i\pi}{r}}\right)^2}{2r} = \frac{2 \sin^2\left(\frac{\pi}{r}\right)}{r} \in (0, 1).$$

Lemma 17. *Let M, N be compact oriented 3-manifolds and $f : \partial M \rightarrow \partial N$ be an orientation-preserving homeomorphism. Let Z be a unitary TQFT, and write $f_{\#} : Z(\partial M) \rightarrow Z(\partial N)$ for the isomorphism induced by f . Then*

$$Z(M \cup_f -N) = \langle f_{\#} Z(M), Z(N) \rangle_{Z(\partial N)} \in \mathbb{C}.$$

Proof. In Cob_3 there are morphisms

$$-N : \partial N \rightarrow \emptyset, \quad N : \emptyset \rightarrow \partial N.$$

Hence

$$Z(-N) : Z(\partial N) \rightarrow Z(\emptyset) = \mathbb{C}, \quad Z(N) : \mathbb{C} \rightarrow Z(\partial N).$$

Note that N and $-N$ are conjugate. By the definition of unitary TQFT,

$$\begin{aligned} & \langle f_{\#} Z(M), Z(N) \rangle_{Z(\partial N)} \\ &= \langle f_{\#} Z(M) 1, Z(N) 1 \rangle_{Z(\partial N)} \\ &= \langle Z(-N) f_{\#} Z(M) 1, 1 \rangle \\ &= Z(M \cup_f -N). \end{aligned}$$

□

In the case where we glue two components Σ_1, Σ_2 of the boundary of one manifold M , it is equivalent to glue $\Sigma_1 \times I$ to M , and the lemma applies as well.

Theorem 18. *Let Z be the original Turaev–Viro TQFT [TV92], at $q = e^{\pi i/r}$, $r \geq 3$. Let M be a closed oriented 3-manifold. Then*

$$|Z(M)| \leq Z(S^3)^{-g(M)+1},$$

where $g(M)$ is the Heegaard genus of M . Thus

$$g(M) - 1 \geq -\frac{\log |Z(M)|}{\log Z(S^3)}.$$

Proof. Let $M = H \cup_f (-H)$ be a Heegaard splitting of M , where H is a handlebody of genus g , and $f : \partial H \rightarrow \partial H$ is an orientation-preserving homeomorphism. Let $u := Z(H) \in Z(\partial H)$. Then

$$\begin{aligned} |Z(M)| &= |\langle u, f_{\#}(u) \rangle| \\ &\leq \sqrt{\langle u, u \rangle \langle f_{\#}(u), f_{\#}(u) \rangle} \\ &= \langle u, u \rangle \\ &= Z(\#_{i=1}^g S^2 \times S^1) \\ &= Z(S^2 \times S^1)^g Z(S^3)^{-g+1} \\ &= Z(S^3)^{-g+1}. \end{aligned}$$

□

4.2 Searching for counterexamples

First we claim that

Proposition 19. *There are no counterexamples to the rank versus genus conjecture among manifolds with Heegaard genus 0, 1, 2.*

Symbol	Base orbifold
S2	2-sphere
RP2/n2	Real projective plane
T	Torus
KB/n2	Klein bottle
D	Disc with regular boundary
M/n2	Möbius band with regular boundary
A	Annulus with two regular boundaries

Proof. If $\pi_1(M)$ is a finite cyclic group, by elliptisation theorem M is elliptic. The only elliptic 3-manifolds with finite cyclic fundamental group are the lens spaces (and S^3).

If $\pi_1(M) = \mathbb{Z}$, M is irreducible and does not contain incompressible torus and hence is geometric. M has $S^2 \times \mathbb{R}$ geometry, and is hence either $S^1 \times S^2$ or $\mathbb{R}P^2 \tilde{\times} S^1$. $\pi_1(S^1 \times S^2) = \mathbb{Z}$, while $\pi_1(\mathbb{R}P^2 \tilde{\times} S^1)$ is not \mathbb{Z} . Thus $M = S^1 \times S^2$. \square

For the theorems used in the proof, see [Mar16]. Manifolds we are looking for have Heegaard genus at least 3.

The software *Regina* [BBP+21] is used in concrete calculation. Given a triangulation, *Regina* computes the state sum invariant. The function `turaevViroApprox()` in *Regina* computes the given Turaev–Viro state sum invariant of a 3-manifold using a fast but inexact floating-point approximation. The function `turaevViro()` in *Regina* computes the given Turaev–Viro state sum invariant of a 3-manifold using exact arithmetic.

We use the census of all minimal triangulations of all closed prime orientable 3-manifolds that can be built from ≤ 11 tetrahedra, as tabulated using *Regina* [Bur11].

We use the original Turaev–Viro TQFT [TV92], at $q = e^{\pi i/r}$, $r = 5$.

4.2.1 Notation

Notation used in naming 3-manifolds is described in *Regina*.

The name $T \times I / [a, b - c, d]$ denotes a torus bundle over the circle. This torus bundle is expressed as the product of the torus and the interval, with the two torus boundaries identified according to the monodromy $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

A Seifert fibred space is denoted by SFS [B: $(p_1, q_1) (p_2, q_2) \dots (p_k, q_k)$], where B is the base orbifold and each (p_i, q_i) describes an exceptional fibre.

In the special case where there are no exceptional fibres at all, we simply write the space as $B \times S^1$ (if there are no fibre-reversing twists) or $B \tilde{\times} S^1$ (if there are).

The name SFS [$B_1: \dots$] U/m SFS [$B_2: \dots$], $m = [a, b - c, d]$ denotes a graph manifold formed from a pair of Seifert fibred spaces $\mathcal{M}_1, \mathcal{M}_2$. Let ϕ_i and ω_i be curves on the boundary of \mathcal{M}_i representing the fibres and base orbifold, $i = 1, 2$. We identify both boundaries so that

$$\begin{bmatrix} \phi_2 \\ \omega_2 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \times \begin{bmatrix} \phi_1 \\ \omega_1 \end{bmatrix}.$$

3-piece graph manifolds are defined similarly.

4.2.2 Data for $r = 5$

Here is a table of manifolds with Turaev–Viro invariant ≥ 7.235 . Their H_1 are shown. Here \star means the Heegaard genus is larger than the rank of H_1 .

SFS [S2: (2,1) (2,1) (2,1) (2,-1)] : #1: 13.105572809000083
 2 $Z_{.2} + Z_{.4}$
 T x S1 : #1: 15.999999999999984
 3 Z
 KB/n2 x~ S1 : #1: 15.999999999999988
 Z + 2 $Z_{.2}$
 SFS [S2: (2,1) (2,1) (2,1) (2,1)] : #1: 14.894427190999902
 2 $Z_{.2} + Z_{.8}$
 SFS [S2: (2,1) (2,1) (2,1) (3,-2)] : #1: 7.999999999999994
 $Z_{.2} + Z_{.10}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (3,-1)] : #1: 7.447213595499956
 $Z_{.2} + Z_{.14}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (2,3)] : #1: 14.894427190999908
 2 $Z_{.2} + Z_{.12}$
 SFS [S2: (2,1) (2,1) (2,1) (3,2)] : #1: 7.4472135954999485
 $Z_{.2} + Z_{.26}$ \star True by [BZ84].
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], $m = [-1,3 \text{ --- } 0,1]$: #1:
 7.788854381999807
 2 $Z_{.4}$ \star
 SFS [S2: (2,1) (2,1) (2,1) (2,5)] : #1: 13.105572809000096
 2 $Z_{.2} + Z_{.16}$
 SFS [S2: (2,1) (2,1) (2,1) (3,4)] : #1: 7.44721359549994
 $Z_{.2} + Z_{.34}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (5,-9)] : #1: 9.919349550499504
 $Z_{.2} + Z_{.6}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (5,-4)] : #1: 9.919349550499517
 $Z_{.2} + Z_{.14}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (5,-1)] : #1: 9.919349550499518
 $Z_{.2} + Z_{.26}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (7,-12)] : #1: 7.4472135954999406
 $Z_{.2} + Z_{.6}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (7,-4)] : #1: 7.447213595499941
 $Z_{.2} + Z_{.26}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (7,-3)] : #1: 7.999999999999984
 $Z_{.2} + Z_{.30}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (7,-2)] : #1: 7.447213595499941
 $Z_{.2} + Z_{.34}$ \star True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (2,-3)] : #1: 74.2111456179999 Note!
 4 $Z_{.2}$
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (2,-1)] : #1: 77.78885438199974 Note!
 3 $Z_{.2} + Z_{.6}$
 SFS [RP2/n2: (2,1) (2,1) (2,-1)] : #1: 18.89442719099984
 2 $Z_{.2} + Z_{.8}$
 SFS [RP2/n2: (2,1) (2,1) (2,1)] : #1: 22.894427190999874

$2 Z_2 + Z_8$
 SFS [T: (2,3)] : #1: 9.236067977499754
 $2 Z + Z_3$
 SFS [KB/n2: (2,3)] : #1: 9.236067977499763
 $Z + Z_8$ ★ False.
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], $m = [-1,4 \text{ --- } 0,1]$: #1:
 8.422291236000305
 $2 Z_2 + Z_8$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [0,1 \text{ --- } 1,0]$: #1:
 8.161300899000896
 $2 Z_2 + Z_4$
 SFS [S2: (2,1) (2,1) (2,1) (2,7)] : #1: 15.999999999999968
 $2 Z_2 + Z_{20}$
 SFS [S2: (2,1) (2,1) (2,1) (3,7)] : #1: 7.447213595499941
 $Z_2 + Z_{46}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (3,8)] : #1: 7.999999999999971
 $Z_2 + Z_{50}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (5,1)] : #1: 9.919349550499517
 $Z_2 + Z_{34}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (5,4)] : #1: 9.919349550499517
 $Z_2 + Z_{46}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (7,2)] : #1: 7.999999999999983
 $Z_2 + Z_{50}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (7,3)] : #1: 7.447213595499948
 $Z_2 + Z_{54}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (13,-21)] : #1: 7.447213595499955
 $Z_2 + Z_6$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (13,-8)] : #1: 7.447213595499948
 $Z_2 + Z_{46}$ ★ True by [BZ84].
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (2,1)] : #1: 71.99999999999923 Note!
 $3 Z_2 + Z_{10}$
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (3,-5)] : #1: 37.105572809000008
 $2 Z_2 + Z_4$
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (3,-2)] : #1: 37.105572809000008
 $2 Z_2 + Z_{16}$
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (3,-4)] : #1: 38.894427190999856
 $2 Z_2 + Z_8$
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (3,-1)] : #1: 36.00000000000012
 $2 Z_2 + Z_{20}$
 SFS [RP2/n2: (2,1) (2,1) (2,3)] : #1: 16.42229123600026
 $2 Z_2 + Z_8$
 SFS [RP2/n2: (2,1) (2,1) (3,-2)] : #1: 9.447213595499948
 $Z_4 + Z_{12}$ ★ False.
 SFS [RP2/n2: (2,1) (2,1) (3,1)] : #1: 9.447213595499948
 $Z_4 + Z_{12}$ ★ False.
 SFS [RP2/n2: (2,1) (2,1) (3,-1)] : #1: 11.447213595499925
 $Z_4 + Z_{12}$ ★ False.
 SFS [RP2/n2: (2,1) (2,1) (3,2)] : #1: 8.211145618000165
 $Z_4 + Z_{12}$ ★ False.
 SFS [T: (1,5)] : #1: 7.999999999999971

$2 Z + Z_5$
 SFS [T: (3,4)] : #1: 9.527864045000376
 $2 Z + Z_4$
 SFS [KB/n2: (1,5)] : #1: 7.999999999999987
 $Z + Z_4$ ★ False.
 SFS [KB/n2: (3,4)] : #1: 9.52786404500038
 $Z + Z_2 + Z_6$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [1,4 — 0,1] : #1: 9.105572809000032
 $2 Z_2 + Z_{16}$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [4,3 — 1,1] : #1: 7.2360679774997605
 $Z_4 + Z_{20}$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-1,5 — 0,1] : #1:
 7.44721359549993
 $Z_4 + Z_{12}$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [5,3 — 2,1] : #1: 7.236067977499749
 $Z_4 + Z_{20}$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-5,7 — -3,4] : #1:
 7.788854381999794
 $2 Z_4$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-5,8 — -3,5] : #1:
 15.577708763999555
 $2 Z_2 + Z_4$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-5,8 — -2,3] : #1:
 10.89442719099983
 $2 Z_2 + Z_8$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (3,1) (3,2)], m = [-1,3 — 0,1] : #1:
 7.788854381999803
 $Z_2 + Z_{18}$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (3,2) (3,2)], m = [-1,3 — -1,2] : #1:
 7.236067977499761
 $Z_2 + Z_{30}$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [1,1 — 0,1] : #1:
 12.1613008990009
 $2 Z_2 + Z_{16}$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [1,1 — 1,0] : #1:
 16.161300899000842
 $2 Z_2 + Z_{16}$
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [1,1 — 1,2] : #1:
 15.05572809000077
 $Z + 2 Z_2$
 SFS [D: (2,1) (3,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [0,1 — 1,0] : #1:
 11.447213595499912
 $Z_2 + Z_6$ ★
 SFS [D: (2,1) (3,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [0,1 — 1,1] : #1:
 9.447213595499933
 $Z_2 + Z_{14}$ ★
 SFS [D: (2,1) (3,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [0,1 — -1,1] : #1:
 9.44721359549994
 $Z_2 + Z_{34}$ ★

SFS [D: (2,1) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [0,1 — 1,0] : #1:
8.080650449500448
Z_2 + Z_18 ★

SFS [D: (2,1) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [0,1 — 1,1] : #1:
7.527864045000408
Z_2 + Z_10 ★

SFS [S2: (2,1) (2,1) (2,1) (2,9)] : #1: 13.105572809000092
2 Z_2 + Z_24

SFS [S2: (2,1) (2,1) (2,1) (5,6)] : #1: 9.919349550499511
Z_2 + Z_54 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (5,9)] : #1: 9.919349550499511
Z_2 + Z_66 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (7,-13)] : #1: 7.999999999999955
Z_2 + Z_10 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (7,11)] : #1: 7.447213595499942
Z_2 + Z_86 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (7,12)] : #1: 7.999999999999976
Z_2 + Z_90 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (13,-23)] : #1: 7.447213595499916
Z_2 + Z_14 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (13,-22)] : #1: 7.999999999999953
Z_2 + Z_10 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (13,8)] : #1: 7.999999999999988
Z_2 + Z_110 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (13,-3)] : #1: 7.447213595499927
Z_2 + Z_66 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (15,-26)] : #1: 9.919349550499476
Z_2 + Z_14 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (15,-11)] : #1: 9.919349550499486
Z_2 + Z_46 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (15,-4)] : #1: 9.919349550499486
Z_2 + Z_74 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (17,-29)] : #1: 7.447213595499921
Z_2 + Z_14 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (17,-12)] : #1: 7.447213595499927
Z_2 + Z_54 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (17,-27)] : #1: 7.4472135954999255
Z_2 + Z_6 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (17,-7)] : #1: 7.447213595499917
Z_2 + Z_74 ★ True by [BZ84].

SFS [S2: (2,1) (2,1) (2,1) (18,-31)] : #1: 14.894427190999846
2 Z_2 + Z_8

SFS [S2: (2,1) (2,1) (2,1) (18,-13)] : #1: 14.894427190999856
2 Z_2 + Z_28

SFS [S2: (2,1) (2,1) (2,1) (18,-29)] : #1: 13.105572809000005
2 Z_2 + Z_4

SFS [S2: (2,1) (2,1) (2,1) (18,-11)] : #1: 14.894427190999844
2 Z_2 + Z_32

SFS [S2: (2,1) (2,1) (2,1) (18,-7)] : #1: 15.999999999999904
2 Z_2 + Z_40

SFS [S2: (2,1) (2,1) (2,1) (18,-5)] : #1: 13.10557280900005
 2 Z_2 + Z_44
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (2,3)] : #1: 77.788854382 [Note!](#)
 3 Z_2 + Z_14
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (3,1)] : #1: 38.89442719099981
 2 Z_2 + Z_28
 SFS [S2: (2,1) (2,1) (2,1) (2,1) (3,2)] : #1: 38.89442719099981
 2 Z_2 + Z_32
 SFS [S2: (2,1) (2,1) (2,1) (3,1) (3,-5)] : #1: 19.447213595499942
 Z_2 + Z_6 **★False.**
 SFS [S2: (2,1) (2,1) (2,1) (3,1) (3,-2)] : #1: 18.552786404500043
 Z_2 + Z_42 **★False.**
 SFS [S2: (2,1) (2,1) (2,1) (3,1) (3,-4)] : #1: 18.55278640450001
 Z_2 + Z_18 **★False.**
 SFS [S2: (2,1) (2,1) (2,1) (3,1) (3,-1)] : #1: 19.4472135955
 Z_2 + Z_54 **★False.**
 SFS [S2: (2,1) (2,1) (2,1) (3,2) (3,-4)] : #1: 17.999999999999954
 Z_2 + Z_30 **★False.**
 SFS [S2: (2,1) (2,1) (2,1) (3,2) (3,-1)] : #1: 19.447213595500003
 Z_2 + Z_66 **★False.**
 SFS [RP2/n2: (2,1) (2,1) (2,5)] : #1: 22.894427191000126
 2 Z_2 + Z_8
 SFS [RP2/n2: (2,1) (2,1) (3,4)] : #1: 11.447213595499914
 Z_4 + Z_12 **★False.**
 SFS [RP2/n2: (2,1) (2,1) (3,5)] : #1: 11.447213595499916
 Z_4 + Z_12 **★False.**
 SFS [T: (2,7)] : #1: 9.236067977499735
 2 Z + Z_7
 SFS [T: (3,7)] : #1: 9.236067977499753
 2 Z + Z_7
 SFS [T: (3,8)] : #1: 18.472135954999473
 2 Z + Z_8
 SFS [T: (4,5)] : #1: 7.9999999999999645
 2 Z + Z_5
 SFS [KB/n2: (2,7)] : #1: 9.236067977499738
 Z + Z_8 **★False.**
 SFS [KB/n2: (3,7)] : #1: 9.23606797749975
 Z + Z_12 **★False.**
 SFS [KB/n2: (3,8)] : #1: 18.472135954999477
 Z + Z_2 + Z_6
 SFS [KB/n2: (4,5)] : #1: 7.999999999999974
 Z + Z_16 **★False.**
 T x I / [13,8 — 8,5] : #1: 10.472135954999501
 Z + 2 Z_4
 T x I / [-13,-8 — -8,-5] : #1: 10.472135954999501
 Z + Z_2 + Z_10
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [1,4 — 0,-1] : #1:
 11.577708763999615
 2 Z_2 + Z_24

SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-3,-4 — 1,1] : #1:
 9.105572809000037
 2 Z_2 + Z_36
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-3,-4 — 2,3] : #1:
 8.422291236000294
 2 Z_2 + Z_48
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [1,5 — 0,1] : #1: 7.999999999999984
 Z_4 + Z_20 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [4,5 — 1,1] : #1: 7.447213595499944
 Z_4 + Z_28 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [4,5 — 3,4] : #1: 7.447213595499939
 Z_4 + Z_8 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [5,7 — 3,4] : #1: 7.2360679774997685
 Z_4 + Z_20
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [3,8 — 2,5] : #1: 11.57770876399964
 2 Z_2 + Z_16
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [5,8 — 2,3] : #1: 10.894427190999833
 2 Z_2 + Z_32
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [5,8 — 3,5] : #1: 14.472135954999498
 2 Z_2 + Z_20
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-7,12 — -3,5] : #1:
 10.894427190999867
 2 Z_2 + Z_12
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-5,12 — -2,5] : #1:
 15.577708763999548
 2 Z_2 + Z_16
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1)], m = [-5,13 — -2,5] : #1:
 7.236067977499755
 Z_4 + Z_20
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (3,1)], m = [-5,8 — -2,3] : #1:
 7.788854381999799
 Z_2 + Z_18 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (3,2)], m = [-5,8 — -2,3] : #1:
 7.236067977499767
 Z_2 + Z_30
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [1,1 — -1,0] : #1:
 17.950155281000672
 2 Z_2 + Z_32
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [1,1 — 0,-1] : #1:
 11.478019326001158
 2 Z_2 + Z_32
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [2,1 — 1,0] : #1:
 17.95015528100069
 2 Z_2 + Z_28
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [2,1 — 1,1] : #1:
 15.05572809000081
 2 Z_2 + Z_20
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [-1,3 — 0,1] : #1:
 34.04984471899907
 2 Z_2 + Z_16

SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [2,1 — 3,1] : #1:
 12.1613008990009
 2 Z_2 + Z_4
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [-2,3 — -1,1] : #1:
 27.83869910099898
 2 Z_2 + Z_12
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [-2,3 — -1,2] : #1:
 28.521980673998794
 2 Z_2 + Z_4
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [-1,3 — -1,2] : #1:
 36.52198067399852
 2 Z_2 + Z_16
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [-2,3 — -3,4] : #1:
 22.049844718999115
 2 Z_2 + Z_4
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (2,1)], m = [-1,3 — -1,4] : #1:
 32.944271909999095
 Z + 2 Z_2
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (3,1)], m = [1,1 — 0,1] : #1:
 7.527864045000402
 Z_4 + Z_20 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (3,1)], m = [1,1 — 1,0] : #1:
 7.527864045000403
 Z_4 + Z_20 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (3,1)], m = [1,1 — 1,2] : #1:
 8.080650449500455
 2 Z_4 ★
 SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (2,1) (3,2)], m = [1,1 — 0,1] : #1:
 8.975077640500356
 Z_4 + Z_28 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [-2,-1 — 1,1] : #1:
 8.552786404500011
 Z_4 + Z_24 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [-1,-1 — 2,1] : #1:
 8.552786404500011
 2 Z_8 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [-1,3 — 0,1] : #1:
 7.788854381999817
 2 Z_8 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [1,-3 — 0,1] : #1:
 11.788854381999764
 2 Z_8 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [-2,3 — 1,-2] : #1:
 9.91934955049948
 Z_4 + Z_8 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [-2,3 — 1,-1] : #1:
 10.683281572999679
 Z_4 + Z_8 ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (2,1)], m = [-1,3 — 1,-2] : #1:
 9.024922359499575

$2 Z_8$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (3,1)], $m = [1,1 \text{ --- } 0,-1]$: #1:
 10.633436854000454
 $2 Z_2 + Z_{24}$
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (3,1)], $m = [1,-2 \text{ --- } 0,1]$: #1:
 10.683281572999697
 $Z_4 + Z_{12}$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (3,1)], $m = [-1,2 \text{ --- } 1,-1]$: #1:
 10.683281572999693
 $Z_4 + Z_{12}$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (3,2)], $m = [0,1 \text{ --- } 1,-2]$: #1:
 8.422291236000298
 $2 Z_2 + Z_{12}$
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (3,2)], $m = [-1,2 \text{ --- } 1,-1]$: #1:
 9.919349550499494
 $Z_4 + Z_{12}$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [M/n2: (5,3)], $m = [0,1 \text{ --- } 1,-1]$: #1:
 9.527864045000399
 $2 Z_2 + Z_{20}$
 SFS [D: (2,1) (3,1)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [0,-1 \text{ --- } 1,0]$: #1:
 11.447213595499914
 $Z_2 + Z_{54}$ ★
 SFS [D: (2,1) (3,1)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [0,1 \text{ --- } 1,-1]$: #1:
 8.211145618000156
 $Z_2 + Z_{26}$ ★
 SFS [D: (2,1) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [0,-1 \text{ --- } 1,0]$: #1:
 8.975077640500354
 $Z_2 + Z_{66}$ ★
 SFS [D: (2,1) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [1,1 \text{ --- } 0,1]$: #1:
 7.527864045000412
 $Z_2 + Z_{50}$ ★
 SFS [D: (2,1) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [1,1 \text{ --- } 1,0]$: #1:
 8.975077640500354
 $Z_2 + Z_{54}$ ★
 SFS [D: (3,1) (3,1)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [-1,1 \text{ --- } 1,0]$: #1:
 8.975077640500354
 $Z_6 + Z_{18}$ ★
 SFS [D: (3,2) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [-1,1 \text{ --- } 0,1]$: #1:
 7.527864045000394
 $Z_2 + Z_{30}$ ★
 SFS [D: (3,2) (3,2)] U/m SFS [D: (2,1) (2,1) (2,1)], $m = [-1,1 \text{ --- } 1,0]$: #1:
 8.080650449500455
 $2 Z_6$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [A: (4,1)] U/n SFS [D: (2,1) (2,1)], $m = [0,1 \text{ --- } 1,0]$, $n = [1,1 \text{ --- } 1,0]$: #1: 7.447213595499939
 $Z_4 + Z_{12}$ ★
 SFS [D: (2,1) (2,1)] U/m SFS [A: (4,3)] U/n SFS [D: (2,1) (2,1)], $m = [0,1 \text{ --- } 1,0]$, $n = [1,1 \text{ --- } 1,0]$: #1: 7.788854381999809
 $2 Z_4$ ★

SFS [D: (2,1) (2,1)] U/m SFS [A: (5,2)] U/n SFS [D: (2,1) (2,1)], m = [0,1 — 1,0], n = [1,1 — 1,0] : #1: 8.21114561800015
Z_4 + Z_12 ★

SFS [D: (2,1) (2,1)] U/m SFS [A: (5,3)] U/n SFS [D: (2,1) (2,1)], m = [0,1 — 1,0], n = [0,1 — 1,0] : #1: 8.21114561800015
Z_4 + Z_28 ★

SFS [D: (2,1) (2,1)] U/m SFS [A: (5,3)] U/n SFS [D: (2,1) (2,1)], m = [0,1 — 1,0], n = [1,1 — 1,0] : #1: 16.42229123600025
2 Z_2 + Z_8

SFS [D: (2,1) (2,1)] U/m SFS [A: (5,3)] U/n SFS [D: (2,1) (2,1)], m = [0,-1 — 1,0], n = [0,1 — 1,0] : #1: 8.21114561800015
Z_4 + Z_12 ★

4.2.3 Analysis

Here ★ means the Heegaard genus is larger than the rank of H_1 . These are potential counterexamples.

Proposition 20. (Proposition 1.5 in [BZ84])

$$G = \langle s_1, \dots, s_{2q}, f \mid s_1^2 f, \dots, s_{2q-1}^2 f, s_{2q}^{2\lambda+1} f^\beta, s_1 \dots s_{2q} f^e \rangle$$

with $q \geq 2$, $\lambda > 0$ and β arbitrary, admits a presentation with $2q - 2$ generators and $2q - 2$ defining relators.

Usually the fundamental group considered is a quotient of G . We write **True** by [BZ84]. if the manifold is a counterexample by this proposition.

Observe that among the 92 potential counterexamples, 35 of them are marked as above. By Theorem 3.1 in [BZ84], they are counterexamples. Also, in the list, any manifold that can be judged by Theorem 3.1 are marked.

Note that the Seifert Euler number in [BZ84] differs from the one defined in [Mar16] by a sign.

Theorem 2.1 and Theorem 1.1 in [BZ84] also help, and 17 manifolds are marked with **False**.

4.2.4 Other remarks

In [Wei03] the writer describes a family of 3-dimensional graph manifolds consisting of two Seifert fibre spaces whose base orbifolds are the Möbius band and a disk and which have one, respectively two, exceptional fibres. The considered spaces have 2-generated fundamental group but are not of Heegaard genus 2. It is not difficult to see that these manifolds are of genus 3. The precise description of those manifolds is given in Theorem 1 there.

Observe that as manifolds, $\text{SFS [D: (2,1) (2,-1)]} \cong$ the circle bundle over the Möbius strip with total space orientable. (See Proposition 10.3.33 in [Mar16].) As Seifert fibrations, $\text{SFS [D: (2,k) (2,j)]} \cong \text{SFS [D: (2,1) (2,-1)]}$ for k, j odd, and they need to be considered. (See Proposition 10.3.11 in [Mar16].) The TQFT method only tells us that among the 92 selected manifolds, genus ≥ 3 . What we can detect is the case genus ≥ 3 and rank = 2.

Accordingly, the author thinks that the following manifold might be another counterexample:

SFS [D: (2,1) (2,1)] U/m SFS [D: (2,1) (3,1)], $m = [-5,8 \text{ --- } -2,3]$: #1:
7.788854381999799
Z_2 + Z_18

The case $r = 4$ has been considered by the author and nothing interesting was found in the database.

There is also a census of smallest known closed hyperbolic 3-manifolds (where “smallest” refers to volume), as tabulated by Hodgson and Weeks [HW94]. The cases $r = 3, 4, 5, 6$ have been considered by the author and nothing interesting was found in this database.

Acknowledgements

I would like to thank Professor Nicolai Reshetikhin for guiding me to quantum topology, which is the key tool in this thesis, and is in itself interesting and useful. Also I would like to thank Professor Yi Liu for introducing me to the conjecture, and Professor Xiaoming Du for discussions before the use of TQFT. I thank Professor Yin Tian for telling me there is something called TQFT.

I thank Professor Qingtao Chen, Tian Yang and Tao Li for related talks, and Professor Yang Su for inviting me to give a talk at Academy of Mathematics and Systems Science, CAS. I thank Xiang Liu for his idea after the talk, and Professor Yinghua Ai, Si Li, Zhongzi Wang and Fan Ye for discussions.

I thank teachers in Yau Mathematical Sciences Center and Department of Mathematical Sciences. Finally I would like to thank my family, especially Rui Ling, for companion.

About Regina

Regina is a software for low-dimensional topology, with a focus on triangulations, knots and links, normal surfaces, and angle structures.

Any database used here can be downloaded from:

<https://regina-normal.github.io/data.html>.

References

- [BBP⁺21] Benjamin A. Burton, Ryan Budney, William Pettersson, et al., *Regina: Software for low-dimensional topology*, <http://regina-normal.github.io/>, 1999–2021. [18](#)
- [BHMV92] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Three-manifold invariants derived from the Kauffman bracket*, *Topology* **31** (1992), no. 4, 685–699. [2](#)
- [BHMV95] ———, *Topological quantum field theories derived from the Kauffman bracket*, *Topology* **34** (1995), no. 4, 883–927. [2](#)
- [BK01] Bojko Bakalov and A.A. Kirillov, *Lectures on tensor categories and modular functors*, Amer. Math. Soc. Univ. Lect. Ser., vol. 21, 01 2001. [2](#), [3](#), [5](#)

- [Bur11] Benjamin A. Burton, *Detecting genus in vertex links for the fast enumeration of 3-manifold triangulations*, Proceedings of the 36th international symposium on Symbolic and algebraic computation - ISSAC '11, ACM Press, 2011. [18](#)
- [BW96] John Barrett and Bruce Westbury, *Invariants of piecewise-linear 3-manifolds*, Transactions of the American Mathematical Society **348** (1996), no. 10, 3997–4022. [2](#)
- [BZ84] Michel Boileau and Heiner Zieschang, *Heegaard genus of closed orientable Seifert 3-manifolds*, Inventiones mathematicae **76** (1984), no. 3, 455–468. [1](#), [3](#), [19](#), [20](#), [22](#), [27](#)
- [CY18] Qingtao Chen and Tian Yang, *Volume conjectures for the Reshetikhin–Turaev and the Turaev–Viro invariants*, Quantum Topology (2018). [2](#)
- [Gar98] Stavros Garoufalidis, *Applications of quantum invariants in low dimensional topology*, Topology **37** (1998), no. 1, 219–224. [2](#), [15](#), [16](#)
- [HAP95] Henning Haahr Andersen and Jan Paradowski, *Fusion categories arising from semisimple lie algebras*, Communications in Mathematical Physics **169** (1995), no. 3, 563–588. [2](#), [6](#)
- [HW94] Craig D. Hodgson and Jeffrey R. Weeks, *Symmetries, isometries and length spectra of closed hyperbolic three-manifolds*, Experimental Mathematics **3** (1994), no. 4, 261–274. [28](#)
- [KJ96] Alexander Kirillov Jr, *On an inner product in modular tensor categories*, Journal of the American Mathematical Society **9** (1996), no. 4, 1135–1169. [2](#), [7](#)
- [Li13] Tao Li, *Rank and genus of 3-manifolds*, Journal of the American Mathematical Society **26** (2013), no. 3, 777–829. [1](#)
- [Lus10] George Lusztig, *Introduction to quantum groups*, Springer Science & Business Media, 2010. [2](#)
- [Mar16] Bruno Martelli, *An introduction to geometric topology*, 2016. [18](#), [27](#)
- [Mü03] Michael Müger, *From subfactors to categories and topology II: The quantum double of tensor categories and subfactors*, Journal of Pure and Applied Algebra **180** (2003), no. 1, 159–219. [12](#), [13](#), [15](#)
- [NS09] Hossein Namazi and Juan Souto, *Heegaard splittings and pseudo-Anosov maps*, Geometric and Functional Analysis **19** (2009), no. 4, 1195–1228. [1](#)
- [Row06] Eric C. Rowell, *From quantum groups to unitary modular tensor categories*, 2006. [2](#), [3](#), [7](#)
- [RT91] Nicolai Reshetikhin and Vladimir Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Inventiones mathematicae **103** (1991), no. 1, 547–597. [1](#), [7](#)

- [SW07] Jennifer Schultens and Richard Weidmann, *On the geometric and the algebraic rank of graph manifolds*, Pacific Journal of Mathematics **231** (2007), 481–510. [1](#)
- [Tur92] Vladimir Turaev, *Modular categories and 3-manifold invariants*, International Journal of Modern Physics B **06** (1992), no. 11n12, 1807–1824. [1](#)
- [Tur10] ———, *Quantum invariants of knots and 3-manifolds*, De Gruyter, 2010. [2](#), [8](#), [9](#), [12](#), [13](#), [15](#)
- [TV92] V.G. Turaev and O.Y. Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, Topology **31** (1992), no. 4, 865–902. [2](#), [3](#), [16](#), [17](#), [18](#)
- [TV13] Vladimir Turaev and Alexis Virelizier, *On two approaches to 3-dimensional TQFTs*, 2013. [3](#), [8](#), [9](#), [10](#), [11](#), [12](#), [13](#), [15](#)
- [TV17] ———, *Monoidal categories and topological field theory*, vol. 322, Springer, 2017. [2](#), [3](#), [8](#), [9](#), [10](#), [11](#), [12](#), [14](#), [16](#)
- [TW93] Vladimir Turaev and Hans Wenzl, *Quantum invariants of 3-manifolds associated with classical simple Lie algebras*, Internat. J. Math **4** (1993), no. 2, 323–358. [1](#), [7](#)
- [Wal78] Friedhelm Waldhausen, *Some problems on 3-manifolds*, Proc. Symp. Pure Math, vol. 32, 1978, pp. 313–322. [1](#)
- [Wei03] Richard Weidmann, *Some 3-manifolds with 2-generated fundamental group*, Archiv der Mathematik **81** (2003), no. 5, 589–595. [27](#)
- [Wen98] Hans Wenzl, *C^* tensor categories from quantum groups*, Journal of the American Mathematical Society **11** (1998), no. 2, 261–282. [2](#), [7](#)
- [Wit89] Edward Witten, *Quantum field theory and the Jones polynomial*, Communications in Mathematical Physics **121** (1989), no. 3, 351–399. [2](#), [16](#)
- [Xu98] Feng Xu, *Standard λ -lattices from quantum groups*, Inventiones mathematicae **134** (1998), no. 3, 455–487. [7](#)