

**ELEMENTARY CONSTRUCTION
OF A MINIMAL FREE RESOLUTIONS
OF THE SPECHT IDEAL OF SHAPE $(n - d, d)$**

KOSUKE SHIBATA AND KOHJI YANAGAWA

ABSTRACT. Let K be a field with $\text{char}(K) = 0$. For a partition λ of $n \in \mathbb{N}$, let I_λ^{Sp} be the ideal of $R = K[x_1, \dots, x_n]$ generated by all Specht polynomials of shape λ . These ideals have been studied from several points of view (and under several names). Using advanced tools of the representation theory, Berkesch Zamaere et al. [1] constructed a minimal free resolution of $I_{(n-d,d)}^{\text{Sp}}$ except differential maps. The present paper constructs the differential maps, and also gives an elementary proof of the result of [1].

1. INTRODUCTION

Throughout this paper, K is a field with $\text{char}(K) = 0$. For a positive integer n , a *partition* of n is a sequence $\lambda = (\lambda_1, \dots, \lambda_l)$ of integers with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_l \geq 1$ and $\sum_{i=1}^l \lambda_i = n$. The notation $\lambda \vdash n$ means that λ is a partition of n . A partition λ is represented by its Young diagram. The *Young tableau* of shape λ is a bijective filling of the boxes of the Young diagram of λ by the integers in $\{1, \dots, n\}$. Let $\text{Tab}(\lambda)$ be the set of tableaux of shape λ . For a tableau $T \in \text{Tab}(\lambda)$, its *Specht polynomial* $f_T \in R := K[x_1, \dots, x_n]$ is the product of all $x_i - x_j$ such that i and j are in the same column of T and j is in a lower position than i . For example, a partition $(4, 2, 1)$ is represented as $\begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & & \\ \square & & & \end{array}$, and

$$T = \begin{array}{cccc} \boxed{4} & \boxed{2} & \boxed{1} & \boxed{7} \\ \boxed{3} & \boxed{5} & & \\ \boxed{6} & & & \end{array}$$

is a tableau of shape $(4, 2, 1)$. In this case, its Specht polynomial is $f_T = (x_4 - x_3)(x_3 - x_6)(x_4 - x_6)(x_2 - x_5)$.

The *vector space* spanned by $\{f_T \mid T \in \text{Tab}(\lambda)\}$ has the natural \mathfrak{S}_n -module structure, where \mathfrak{S}_n is the n -th symmetric group. In fact, this is isomorphic to the *Specht module* V_λ , which is crucial in the representation theory of \mathfrak{S}_n . The *ideal*

$$I_\lambda^{\text{Sp}} = (f_T \mid T \in \text{Tab}(\lambda)) \subset R$$

2020 *Mathematics Subject Classification.* Primary 13F99.

Key words and phrases. Specht polynomial, Specht ideal, subspace arrangement, Cohen–Macaulay ring.

The author is partially supported by JSPS Grant-in-Aid for Scientific Research (C) 19K03456.

is called the *Specht ideal* of shape λ . These ideals have been studied from several points of view (and under several names and characterizations), so the present authors had sometimes “re-proved” existing results in their preceding papers ([6, 10]). Anyway, the Cohen-Macaulay Specht ideals are classified as follows. (Assuming the results in [3], the proof in [10] can be largely simplified. See also [4] for the characteristic free results.)

Theorem 1.1 (c.f. [10, Proposition 2.8 and Corollary 4.4]). *R/I_λ^{Sp} is Cohen-Macaulay if and only if one of the following conditions holds.*

$$(1) \lambda = (n - d, 1, \dots, 1), \quad (2) \lambda = (n - d, d), \quad (3) \lambda = (d, d, 1).$$

The minimal free resolution of I_λ^{Sp} in the case (1) is treated in the joint paper [9] with J. Watanabe. In the cases (2) and (3), C. Berkesch Zamaere et al. [1] determined the \mathfrak{S}_n -module structure of $\text{Tor}_i^R(K, R/I_\lambda^{\text{Sp}})$, especially they gave the graded betti numbers

$$\beta_{i,j}(R/I_\lambda^{\text{Sp}}) := \dim_K[\text{Tor}_i^R(K, R/I_\lambda^{\text{Sp}})]_j.$$

A crucial point of their result is that

$$(*) \beta_{i,j}(I_{(n-d,d)}^{\text{Sp}}) \neq 0 \text{ implies } j = d + i \text{ (resp. } j = d + i + 1) \text{ if } i \leq n - 2d \text{ (resp. } i > n - 2d).$$

In the previous paper [7], the authors constructed minimal free resolutions of $I_{(n-2,2)}^{\text{Sp}}$ and $I_{(d,d,1)}^{\text{Sp}}$ explicitly using Specht modules. In these cases (and the case $I_{(n-1,1)}^{\text{Sp}}$), the fact (*) is easy, and this is a key point of [7]. Extending this result, the present paper constructs a minimal free resolution of $I_{(n-d,d)}^{\text{Sp}}$ for general d .

Contrary to [1], the present paper gives the differential maps of our resolution explicitly. Moreover, while [1] uses highly advanced tools of the representation theory (rational Cherednik algebras, Jack polynomials, etc), we only use the basic theory of Specht modules. Of course, our proof is self-contained, in other words, we give a new and elementary proof of (*).

Here we do not give the precise construction of our resolutions, but introduce an example. Our minimal free resolution $\mathcal{F}_\bullet^{(4,3)}$ of $I_{(4,3)}^{\text{Sp}}$ is as follows.

$$(1.1) \quad 0 \longrightarrow V_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \otimes_K R(-7) \xrightarrow{\partial_3} V_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \otimes_K R(-6) \xrightarrow{\partial_2} V_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \otimes_K R(-4) \\ \xrightarrow{\partial_1} V_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \otimes_K R(-3) \xrightarrow{\partial_0} I_{(4,3)}^{\text{Sp}} \longrightarrow 0.$$

Here the degree of an element in V_λ is 0, so we have $[V_\lambda \otimes R(-i)]_j \cong (R_{j-i})^{\oplus \dim V_\lambda}$. The differential maps of $\mathcal{F}_\bullet^{(4,3)}$ are described in Examples 3.2 below.

The composition of this paper is as follows. §2 collects basic definitions and preliminary results. In §3, we give the construction of our resolution $\mathcal{F}_\bullet^{(n-d,d)}$, and §4 mainly treats the proof of the statement (*), from which we can easily show that $\mathcal{F}_\bullet^{(n-d,d)}$ is a minimal free resolution. So if one can assume (*), which is given by [1], one can skip the most part of this section. Our proof is an induction on n using the branching rule of Specht modules.

2. PRELIMINARIES

In this section, we briefly explain Specht modules and related notions. See [5, Chapter 2] for details.

For a partition $\lambda = (\lambda_1, \dots, \lambda_l)$, we sometimes use “exponential notation”. For example, $(4, 3^2, 2, 1^3)$ means $(4, 3, 3, 2, 1, 1, 1)$. For convenience, a partition like $(4, 3)$ sometimes denoted by $(4, 3, 1^0)$. If $\lambda = (\lambda_1, \dots, \lambda_l)$, then $\text{Tab}(\lambda)$ can be simply written as $\text{Tab}(\lambda_1, \dots, \lambda_l)$. Given any set A , let $S(A)$ be the set of all permutations of A . If a tableau $T \in \text{Tab}(\lambda)$ with $\lambda \vdash n$ has columns $C_1, \dots, C_{\lambda_1} \subset [n] := \{1, 2, \dots, n\}$, then we call $C(T) := S(C_1) \times \dots \times S(C_{\lambda_1})$ the *column-stabilizer* of T . For $T, T' \in \text{Tab}(\lambda)$, T and T' are *row equivalent*, if the corresponding rows of T and T' contain the same elements. For $T \in \text{Tab}(\lambda)$, the *tabloid* $\{T\}$ of T is defined by

$$\{T\} := \{T' \in \text{Tab}(\lambda) \mid T \text{ and } T' \text{ are row equivalent}\},$$

and the *polytabloid* of T is defined by

$$e(T) := \sum_{\sigma \in C(T)} \text{sgn}(\sigma) \sigma\{T\}.$$

The vector space

$$V_\lambda := \sum_{T \in \text{Tab}(\lambda)} Ke(T)$$

becomes an \mathfrak{S}_n -module in the natural way, and it is called the *Specht module* of λ . The Specht modules V_λ are irreducible, and $\{V_\lambda \mid \lambda \vdash n\}$ forms a complete list of the irreducible representations of \mathfrak{S}_n (recall that $\text{char}(K) = 0$ now).

In the previous section, we defined the Specht polynomial $f_T \in R = K[x_1, \dots, x_n]$. Since \mathfrak{S}_n acts on R , the vector subspace

$$\sum_{T \in \text{Tab}(\lambda)} Kf_T$$

is also an \mathfrak{S}_n -module. Moreover, the map

$$(2.1) \quad V_\lambda \longrightarrow \sum_{T \in \text{Tab}(\lambda)} Kf_T, \quad e(T) \longmapsto f_T.$$

is well-defined, and gives an isomorphism as \mathfrak{S}_n -modules.

The set $\{e(T) \mid T \in \text{Tab}(\lambda)\}$ is linearly dependent. For linear relations among them, a *Garnir relation* is fundamental. See [5, §2.6]. We say a tableau T is *standard*, if all columns (resp. rows) are increasing from top to bottom (resp. from left to right). Let $\text{SYT}(\lambda)$ be the set of standard Young tableaux of shape λ . It is a classical result that $\{e(T) \mid T \in \text{SYT}(\lambda)\}$ is a basis of V_λ (c.f. [5, Theorem 2.6.5]).

Lemma 2.1. *We have that $\beta_{i,j}(I_{(n-d,d)}^{\text{Sp}}) \neq 0$ implies $j = d + i, d + i + 1$, and*

$$\beta_{i,d+i}(I_{(n-d,d)}^{\text{Sp}}) - \beta_{i-1,d+i}(I_{(n-d,d)}^{\text{Sp}}) = \begin{cases} \dim_K V_{(n-d-i,d,1^i)} & (i \leq n - 2d), \\ 0 & (i = n - 2d + 1), \\ -\dim_K V_{(d-1,n-d-i+1,1^i)} & (i > n - 2d + 1), \end{cases}$$

Proof. In [1] (see also [6]), the Hilbert series of $R/I_{(n-d,d)}^{\text{Sp}}$ is given by

$$H(R/I_{(n-d,d)}^{\text{Sp}}, t) = \frac{1 + h_1 t + h_2 t^2 + \cdots + h_d t^d}{(1-t)^d}$$

with

$$h_i = \begin{cases} \binom{n-d+i-1}{i} & \text{if } 1 \leq i \leq d-1, \\ \binom{n-1}{d-2} & \text{if } i = d. \end{cases}$$

Since $R/I_{(n-d,d)}^{\text{Sp}}$ is Cohen–Macaulay, the above Hilbert series states that the regularity of $I_{(n-d,d)}^{\text{Sp}}$ is $d+1$. Since $I_{(n-d,d)}^{\text{Sp}}$ is generated by degree d elements, we have the first assertion of the lemma.

Since $\dim_K V_\lambda$ can be computed by the hook formula ([5, Theorem 3.10.2]), the second assertion follows from the first one and the direct computations as in [7, Lemma 4.1]. \square

Remark 2.2. The above computation of the Hilbert series is also treated in the authors' previous paper [6]. While [6] gives extra results (e.g. the characteristic free-ness of the Hilbert series and the “branching rule” for Specht ideals), the computation in [1] is much more direct and quicker than [7].

3. CONSTRUCTION OF THE FREE RESOLUTION

For the Specht ideal $I_{(n-d,d)}^{\text{Sp}}$, we define the chain complex

$$\mathcal{F}_\bullet^{(n-d,d)} : 0 \longrightarrow F_{n-d-1} \xrightarrow{\partial_{n-d-1}} F_{n-d-2} \xrightarrow{\partial_{n-d-2}} \cdots \xrightarrow{\partial_2} F_1 \xrightarrow{\partial_1} F_0 \longrightarrow 0$$

of graded free R -modules as follows (if there is no danger of confusion, we simply denote $\mathcal{F}_i^{(n-d,d)}$ as F_i). Here

$$F_i = V_{(n-d-i, d, 1^i)} \otimes_K R(-d-i)$$

for $0 \leq i \leq n-2d$, and

$$F_i = V_{(d-1, n-d-i, 1^{i+1})} \otimes_K R(-d-i-1)$$

for $n-2d+1 \leq i \leq n-d-1$.

Remark 3.1. The second assertion of Lemma 2.1 states that

$$(3.1) \quad \beta_{i,d+i}(I_{(n-d,d)}^{\text{Sp}}) - \beta_{i-1,d+i}(I_{(n-d,d)}^{\text{Sp}}) = \begin{cases} \text{rank } \mathcal{F}_i^{(n-d,d)} & (i \leq n-2d), \\ 0 & (i = n-2d+1), \\ -\text{rank } \mathcal{F}_{i-1}^{(n-d,d)} & (i > n-2d+1), \end{cases}$$

where $\text{rank } F$ denotes the rank of a free module F .

First, we define the differential map $\partial_i : F_i \rightarrow F_{i-1}$ for $1 \leq i \leq n - 2d$. For a tableau

$$T := \begin{array}{|c|c|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_d & \cdots & b_{n-d-i} \\ \hline a_2 & c_2 & \cdots & c_d & & \\ \hline a_3 & & & & & \\ \hline \vdots & & & & & \\ \hline a_{i+2} & & & & & \\ \hline \end{array} \in \text{Tab}(n-d-i, d, 1^i)$$

and an integer j with $1 \leq j \leq i + 2$, set

$$T^j := \begin{array}{|c|c|c|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_d & \cdots & b_{n-d-i} & a_j \\ \hline a_2 & c_2 & \cdots & c_d & & & \\ \hline a_3 & & & & & & \\ \hline \vdots & & & & & & \\ \hline a_{j-1} & & & & & & \\ \hline a_{j+1} & & & & & & \\ \hline \vdots & & & & & & \\ \hline a_{i+2} & & & & & & \\ \hline \end{array} \in \text{Tab}(n-d-i+1, d, 1^{i-1}).$$

Then, for $1 \leq i \leq n - 2d$, we set

$$\partial_i(e(T) \otimes 1) := \sum_{j=1}^{i+2} (-1)^{j-1} e(T^j) \otimes x_{a_j} \in V_{(n-d-i+1, d, 1^{i-1})} \otimes_K R(-d-i+1) = F_{i-1}.$$

Recall that $e(\sigma T) = \text{sgn}(\sigma)e(T)$ for $\sigma \in C(T)$. It is easy to check that the maps ∂_i defined above satisfies $\partial_i(e(\sigma T) \otimes 1) = \text{sgn}(\sigma) \partial_i(e(T) \otimes 1)$ for all $\sigma \in C(T)$. The same is true for ∂_i for $i > n - 2d$ defined below. However, this is not enough. Since

$\{e(T) \mid T \in \text{Tab}(\lambda)\}$ is linearly dependent, the well-definedness of ∂_i is non-trivial, and we will prove this Proposition 3.4 below.

To define ∂_{n-2d+1} , we need further preparation. For a tableau

$$(3.2) \quad T := \begin{array}{|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{d-1} \\ \hline a_2 & c_2 & \cdots & c_{d-1} \\ \hline a_3 & & & \\ \hline \vdots & & & \\ \hline a_{n-2d+4} & & & \\ \hline \end{array} \in \text{Tab}((d-1)^2, 1^{n-2d+2})$$

and integers j, k with $1 \leq j < k \leq n - 2d + 4$, set

$$T_{j,k} := \begin{array}{|c|c|c|c|c|} \hline \vdots & b_2 & \cdots & b_{d-1} & a_j \\ \hline \vdots & c_2 & \cdots & c_{d-1} & a_k \\ \hline \vdots & & & & \\ \hline \end{array} \in \text{Tab}(d^2, 1^{n-2d}),$$

where the first column is the “transpose” of

$$\begin{array}{|c|c|c|c|c|c|c|c|c|c|} \hline a_1 & a_2 & \cdots & a_{j-1} & a_{j+1} & \cdots & a_{k-1} & a_{k+1} & \cdots & a_{n-2d+4} \\ \hline \end{array}.$$

Then we set

$$\begin{aligned} \partial_{n-2d+1}(e(T) \otimes 1) &:= \sum_{1 \leq j < k \leq n-2d+4} (-1)^{j+k-1} e(T_{j,k}) \otimes x_{a_j} x_{a_k} \\ &\in V_{(d^2, 1^{n-2d})} \otimes_K R(-n+d) = F_{n-2d}. \end{aligned}$$

Finally, we define ∂_i for $n-2d+2 \leq i \leq n-d-1$, but this case is a bit more complicated. For a tableau

$$T := \begin{array}{|c|c|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{n-d-i} & \cdots & b_{d-1} \\ \hline a_2 & c_2 & \cdots & c_{n-d-i} & & \\ \hline a_3 & & & & & \\ \hline \vdots & & & & & \\ \hline a_{i+3} & & & & & \\ \hline \end{array}$$

in $\text{Tab}(d-1, n-d-i, 1^{i+1})$ and an integer j with $1 \leq j \leq i+3$, set

$$T_j := \begin{array}{|c|c|c|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{n-d-i} & b_{n-d-i+1} & \cdots & b_{d-1} \\ \hline a_2 & c_2 & \cdots & c_{n-d-i} & a_j & & \\ \hline \vdots & & & & & & \\ \hline a_{j-1} & & & & & & \\ \hline a_{j+1} & & & & & & \\ \hline \vdots & & & & & & \\ \hline a_{i+3} & & & & & & \\ \hline \end{array}$$

in $\text{Tab}(d-1, n-d-i+1, 1^i)$. Then we set

(3.3)

$$\partial_i(e(T) \otimes 1) := \sum_{j=1}^{i+2} \sum_{\sigma \in H} (-1)^{j-1} e(\sigma(T_j)) \otimes x_{a_j} \in V_{(d-1, d-i+2, 1^{i-1})} \otimes_K R(-d-i) = F_{i-1},$$

where H is the set of the transpositions $(b_{n-d-i+1} b_{n-d-i+k})$ for $k \geq 2$.

Example 3.2. Recall that the “form” of $\mathcal{F}_\bullet^{(4,3)}$ was given in Introduction. The differential maps are as follows.

$$\begin{aligned} \partial_3(e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline 4 & \\ \hline 5 & \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array}) \otimes 1) &= e(\begin{array}{|c|c|} \hline 3 & 2 \\ \hline 4 & 1 \\ \hline 5 & \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array}) \otimes x_1 - e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 4 & 3 \\ \hline 5 & \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array}) \otimes x_3 + e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array}) \otimes x_4 \\ &- e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array}) \otimes x_5 + e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 6 \\ \hline 4 & \\ \hline 5 & \\ \hline 7 & \\ \hline \end{array}) \otimes x_6 - e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 7 \\ \hline 4 & \\ \hline 5 & \\ \hline 6 & \\ \hline \end{array}) \otimes x_7 \end{aligned}$$

(in this case, $H = \{\varepsilon\}$),

$$\begin{aligned} \partial_2(e(\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array}) \otimes 1) &= e(\begin{array}{|c|c|c|} \hline 5 & 2 & 1 \\ \hline 6 & 4 & 3 \\ \hline 7 & & \\ \hline \end{array}) \otimes x_1 x_3 - e(\begin{array}{|c|c|c|} \hline 3 & 2 & 1 \\ \hline 6 & 4 & 5 \\ \hline 7 & & \\ \hline \end{array}) \otimes x_1 x_5 \\ &\vdots \\ &\vdots \\ &- e(\begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & 7 \\ \hline 6 & & \\ \hline \end{array}) \otimes x_5 x_7 + e(\begin{array}{|c|c|c|} \hline 1 & 2 & 6 \\ \hline 3 & 4 & 7 \\ \hline 5 & & \\ \hline \end{array}) \otimes x_6 x_7, \end{aligned}$$

and

$$\begin{aligned} \partial_1(e(\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & 6 \\ \hline 7 & & \\ \hline \end{array}) \otimes 1) &= e(\begin{array}{|c|c|c|c|} \hline 4 & 2 & 3 & 1 \\ \hline 7 & 5 & 6 & \\ \hline \end{array}) \otimes x_1 - e(\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 7 & 5 & 6 & \\ \hline \end{array}) \otimes x_4 \\ &+ e(\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 7 \\ \hline 4 & 5 & 6 & \\ \hline \end{array}) \otimes x_7. \end{aligned}$$

Remark 3.3. (1) The complex $\mathcal{F}_\bullet^{(n-2,2)}$ has been constructed in [7], but the convention is not the same. The complex in [7] is a free resolution of the residue ring $R/I_{(n-2,2)}^{\text{Sp}}$, while the present version is a free resolution of the ideal $I_{(n-2,2)}^{\text{Sp}}$. Similarly, [7] gave a minimal free resolution $\mathcal{F}_\bullet^{(d,d,1)}$ of the residue ring $R/I_{(d,d,1)}^{\text{Sp}}$.

(2) The differential maps ∂_i of $\mathcal{F}_\bullet^{(n-d,d)}$ for $1 \leq i \leq n-2d$ are essentially same as the differential maps of $\mathcal{F}_\bullet^{(n-2,2)}$ except the last one, which is the same type as ∂_{n-2d+1} of $\mathcal{F}_\bullet^{(n-d,d)}$. Similarly, the differential maps ∂_i of $\mathcal{F}_\bullet^{(n-d,d)}$ for $n-2d+2 \leq i \leq n-d-1$ are essentially same as those of $\mathcal{F}_\bullet^{(d,d,1)}$.

Proposition 3.4. *The maps ∂_i are well-defined for all i .*

Proof. The well-definedness of ∂_i for $i \leq n-2d$ (resp. $i \geq n-2d+2$) can be shown by the same way as [7, Theorem 4.2] for $\mathcal{F}_\bullet^{(d-2,2)}$ (resp. [7, Theorem 6.2] for

$\mathcal{F}_{\bullet}^{(d,d,1)}$). So only ∂_{n-2d+1} remains. To prove this, we use the following linear maps

$$\varphi : V_{((d-1)^2, 1^{n-2d+2})} \longrightarrow V_{(d, d-1, 1^{n-2d+1})} \otimes_K R_1$$

$$\psi : V_{(d, d-1, 1^{n-2d+1})} \longrightarrow V_{(d^2, 1^{n-2d})} \otimes_K R_1$$

defined by

$$\varphi(e(T)) := \sum_{i \geq 1} (-1)^{i-1} e(T^i) \otimes x_{a_i} \in V_{(d, d-1, 1^{n-2d+1})} \otimes_K R_1$$

for

$$T = \begin{array}{|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{d-1} \\ \hline a_2 & c_2 & \cdots & c_{d-1} \\ \hline a_3 & & & \\ \hline \vdots & & & \\ \hline \end{array} \in \text{Tab}((d-1)^2, 1^{n-2d+2}),$$

and

$$\psi(e(T')) := \sum_{j \geq 1} (-1)^{j-1} e(T'_j) \otimes x_{a_j} \in V_{(d^2, 1^{n-2d})} \otimes_K R_1$$

for

$$T' = \begin{array}{|c|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{d-1} & b_d \\ \hline a_2 & c_2 & \cdots & c_{d-1} & \\ \hline a_3 & & & & \\ \hline \vdots & & & & \\ \hline \end{array} \in \text{Tab}(d, d-1, 1^{n-2d+1}).$$

The well-definedness of φ (resp. ψ) is the same thing as that of ∂_i for $i \leq n-2d$ (resp. $i \geq n-2d+2$). Finally, consider the linear map $\mu : R_1 \otimes_K R_1 \rightarrow R_2$ defined by

$$\mu(x_j \otimes x_i) = \frac{1}{2} x_i x_j.$$

Identifying the degree $(n-d+2)$ -part of $F_{n-2d+1} = V_{((d-1)^2, 1^{n-2d+2})} \otimes_K R(-n+d-2)$ with $V_{((d-1)^2, 1^{n-2d+2})}$, the composition

$$\begin{array}{ccc} V_{((d-1)^2, 1^{n-2d+2})} & \xrightarrow{\varphi} & V_{(d, d-1, 1^{n-2d+1})} \otimes_K R_1 \\ & & \xrightarrow{\psi \otimes \text{Id}} V_{(d^2, 1^{n-2d})} \otimes_K R_1 \otimes_K R_1 \\ & & \xrightarrow{\text{Id} \otimes \mu} V_{(d^2, 1^{n-2d})} \otimes_K R_2 \end{array}$$

coincides with the degree $(n-d+2)$ -part $[\partial_{n-2d+1}]_{n-d+2}$ of ∂_{n-2d+1} . Here, for example, we will compute the $-\otimes x_{a_1} \otimes x_{a_2}$ and $-\otimes x_{a_2} \otimes x_{a_1}$ parts of $(\psi \otimes \text{Id}) \circ \varphi(e(T))$

for the above T .

$$\begin{aligned}
& (\psi \otimes \text{Id}) \circ \varphi(e(T)) \\
= & (\psi \otimes \text{Id}) \left(e \left(\begin{array}{|c|c|c|c|} \hline a_2 & \cdots & b_{d-1} & a_1 \\ \hline a_3 & \cdots & c_{d-1} & \\ \hline \vdots & & & \\ \hline \end{array} \right) \otimes x_{a_1} - e \left(\begin{array}{|c|c|c|c|} \hline a_1 & \cdots & b_{d-1} & a_2 \\ \hline a_3 & \cdots & c_{d-1} & \\ \hline \vdots & & & \\ \hline \end{array} \right) \otimes x_{a_2} + \cdots \right) \\
= & e \left(\begin{array}{|c|c|c|c|} \hline a_3 & \cdots & b_{d-1} & a_1 \\ \hline a_4 & \cdots & c_{d-1} & a_2 \\ \hline \vdots & & & \\ \hline \end{array} \right) \otimes x_{a_2} \otimes x_{a_1} - e \left(\begin{array}{|c|c|c|c|} \hline a_3 & \cdots & b_{d-1} & a_2 \\ \hline a_4 & \cdots & c_{d-1} & a_1 \\ \hline \vdots & & & \\ \hline \end{array} \right) \otimes x_{a_1} \otimes x_{a_2} + \cdots \\
= & e \left(\begin{array}{|c|c|c|c|} \hline a_3 & \cdots & b_{d-1} & a_1 \\ \hline a_4 & \cdots & c_{d-1} & a_2 \\ \hline \vdots & & & \\ \hline \end{array} \right) \otimes x_{a_2} \otimes x_{a_1} + e \left(\begin{array}{|c|c|c|c|} \hline a_3 & \cdots & b_{d-1} & a_1 \\ \hline a_4 & \cdots & c_{d-1} & a_2 \\ \hline \vdots & & & \\ \hline \end{array} \right) \otimes x_{a_1} \otimes x_{a_2} + \cdots .
\end{aligned}$$

Anyway, we have shown that $[\partial_{n-2d+1}]_{n-d+2}$ is well-defined. Since ∂_{n-2d+1} is determined by $[\partial_{n-2d+1}]_{n-d+2}$, it is also well-defined. \square

Proposition 3.5. $\mathcal{F}_\bullet^{(n-d,d)}$ is actually a chain complex.

Proof. It suffices to show that $\partial_i \partial_{i+1}(e(T) \otimes 1) = 0$ for all i and T . The cases $i \neq n - 2d, n - 2d + 1$ are easy, and have been essentially shown in [7]. Consider the case $i = n - 2d + 1$. Take $e(T) \otimes 1 \in F_{n-2d+2}$, where

$$T = \begin{array}{|c|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{d-2} & b_{d-1} \\ \hline a_2 & c_2 & \cdots & c_{d-2} & \\ \hline a_3 & & & & \\ \hline \vdots & & & & \\ \hline \end{array} \in \text{Tab}(d-1, d-2, 1^{n-2d+3}).$$

Note that $\partial_{n-2d+1} \partial_{n-2d+2}(e(T) \otimes 1)$ can be decomposed as

$$\partial_{n-2d+1} \partial_{n-2d+2}(e(T) \otimes 1) = \sum_{i < j < k} \alpha_{i,j,k} \otimes x_i x_j x_k,$$

for $\alpha_{i,j,k} \in V_{(d^2, 1^{n-2d})}$, and

$$\begin{aligned}
\alpha_{i,j,k} &= (-1)^{i-1} (-1)^{(j-1)+(k-1)-1} e((T_i)_{j,k}) + (-1)^{j-1} (-1)^{i+(k-1)-1} e((T_j)_{i,k}) \\
&\quad + (-1)^{k-1} (-1)^{i+j-1} e((T_k)_{i,j}) \\
&= (-1)^{i+j+k} (e((T_i)_{j,k}) - e((T_j)_{i,k}) + e((T_k)_{i,j})).
\end{aligned}$$

The right most boxes of $(T_i)_{j,k}$, $(T_j)_{i,k}$ and $(T_k)_{i,j}$ are the following

$$(T_i)_{j,k} = \begin{array}{c} \cdots \\ \cdots \end{array} \begin{array}{|c|c|} \hline b_{d-1} & a_j \\ \hline a_i & a_k \\ \hline \end{array}, \quad (T_j)_{i,k} = \begin{array}{c} \cdots \\ \cdots \end{array} \begin{array}{|c|c|} \hline b_{d-1} & a_i \\ \hline a_j & a_k \\ \hline \end{array}, \quad (T_k)_{i,j} = \begin{array}{c} \cdots \\ \cdots \end{array} \begin{array}{|c|c|} \hline b_{d-1} & a_i \\ \hline a_k & a_j \\ \hline \end{array}$$

(these tableau are same in the remaining boxes). By a Garnir relation, we have $e((T_i)_{j,k}) - e((T_j)_{i,k}) + e((T_k)_{i,j}) = 0$, and hence $\partial_{n-2d+1}\partial_{n-2d+2}(e(T) \otimes 1) = 0$.

That $\partial_{n-2d}\partial_{n-2d+1}(e(T) \otimes 1) = 0$ can be shown in a similar way (in some sense, it has been shown in [7], since $\partial_{n-2d}\partial_{n-2d+1}$ is essentially the last of $\mathcal{F}_\bullet^{(n-2,2)}$). \square

For the complex $\mathcal{F}_\bullet^{(n-d,d)}$, the augmentation map $\partial_0 : F_0 \rightarrow I_{(n-d,d)}^{\text{Sp}}$ is defined by

$$\partial_0 : F_0 = V_{(n-d,d)} \otimes_K R(-d) \ni e(T) \otimes 1 \mapsto f_T \in I_{(n-d,d)}^{\text{Sp}}.$$

Clearly, ∂_0 is surjective and well-defined (recall the map (2.1)).

Lemma 3.6. *With the above notation, we have $\partial_0\partial_1 = 0$.*

Proof. The case $d = 2$ is shown in [7], and the proof there works for the general case. (Since $\mathcal{F}_\bullet^{(n-2,2)}$ in [7] is a resolution of $R/I_{(n-2,2)}^{\text{Sp}}$, our $\partial_0\partial_1 = 0$ corresponds to $\partial_1\partial_2 = 0$ there.) \square

We define the \mathfrak{S}_n -action on $\mathcal{F}_\bullet^{(n-d,d)}$. For $F_i = V_\lambda \otimes_K R(-j)$ (here λ is a suitable partition of n and j is a suitable integer) and $\sigma \in \mathfrak{S}_n$, set $\sigma(v \otimes f) := \sigma v \otimes \sigma f$. It is easy to check that ∂_i is an \mathfrak{S}_n -homomorphism for all i .

Lemma 3.7. *With the above notation, the degree j -part $[\partial_i]_j : [F_i]_j \rightarrow [F_{i-1}]_j$ is injective for all $i \geq 1$. Similarly, the degree d -part of the augmentation map $\partial_0 : F_0 \rightarrow I_{(n-d,d)}^{\text{Sp}}$ is injective (actually, bijective).*

The corresponding fact has been shown in [7], but we will explain it again.

Proof. Since $[F_i]_j \cong V_\lambda$ is irreducible as an \mathfrak{S}_n -module and $[\partial_i]_j$ is an \mathfrak{S}_n -homomorphism, it is either the zero map or an injection. However, it is clearly non-zero. \square

By Lemmas 3.6 and 3.7, we see that the former half $F_{n-2d} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow 0$ of $\mathcal{F}_\bullet^{(n-d,d)}$ is a subcomplex of a minimal free resolution P_\bullet of $I_{(n-d,d)}^{\text{Sp}}$ such that each F_i is a direct summand of P_i for each $0 \leq i \leq n-2d$.

4. THE PROOF OF THE MAIN THEOREM

The following is the main result of the present paper, and this section is devoted to its proof.

Theorem 4.1. *The complex $\mathcal{F}_\bullet^{(n-d,d)}$ is a minimal free resolution of $I_{(n-d,d)}^{\text{Sp}}$.*

The theorem is easy when $n = 2d$.

Proposition 4.2. *If $n = 2d$, Theorem 4.1 holds, that is, $\mathcal{F}_\bullet^{(d,d)}$ is a minimal free resolution of $I_{(d,d)}^{\text{Sp}}$.*

Proof. By the equation (3.1), we have $\beta_{1,d+1}(I_{(d,d)}^{\text{Sp}}) = \beta_{0,d+1}(I_{(d,d)}^{\text{Sp}}) = 0$, and hence $\beta_{i,d+i}(I_{(d,d)}^{\text{Sp}}) = 0$ for all $i \geq 1$. So, by the first statement of Lemma 2.1, $\beta_{i,j}(I_{(d,d)}^{\text{Sp}}) \neq 0$ for $i \geq 1$ implies $j = d + i + 1$. By Lemmas 3.6 and 3.7, $\mathcal{F}_{\bullet}^{(d,d)}$ is a subcomplex of a minimal free resolution P_{\bullet} of $I_{(d,d)}^{\text{Sp}}$ such that $\mathcal{F}_i^{(d,d)}$ is a direct summand of P_i for all i , but (3.1) implies that $P_i = \mathcal{F}_i^{(d,d)}$ for all $i \geq 0$, and hence $P_{\bullet} = \mathcal{F}_{\bullet}^{(d,d)}$. \square

In the general case, the proof of Theorem 4.1 is much more complicated, and preparation is required. For $n, d \in \mathbb{N}$ with $n \geq 2d$, we construct the chain complex $\mathcal{L}_{\bullet}^{n,d}$ of graded free R -modules as follows.

$$0 \longrightarrow \mathcal{L}_{n-2d}^{n,d} \longrightarrow \cdots \longrightarrow \mathcal{L}_1^{n,d} \longrightarrow \mathcal{L}_0^{n,d} \longrightarrow 0,$$

$$\mathcal{L}_i^{n,d} = V_{(n-d-i,d,1^i)} \otimes_K R(-d-i),$$

and the differential maps are given by the same way as the former half of $\mathcal{F}_{\bullet}^{(n-d,d)}$. Clearly, $\mathcal{L}_{\bullet}^{n,d}$ is the d -linear strand of $\mathcal{F}_{\bullet}^{(n-d,d)}$, and it is actually a chain complex. If $n = 2d$, then $\mathcal{L}_i^{n,d} = 0$ for $i \neq 0$, and Theorem 4.1 has been proved in this case (Proposition 4.2). So we mainly treat the case $n > 2d$. Since the case $n = 2d + 1$ is still somewhat special, we first assume that $n > 2d + 1$.

Proposition 4.3. *For $n, d \in \mathbb{N}$ with $n > 2d + 1$, the following are equivalent.*

- (1) *Theorem 4.1 holds in this case.*
- (2) *$H_i(\mathcal{L}_{\bullet}^{n,d}) = 0$ for $0 < i < n - 2d$.*

Proof. (1) \Rightarrow (2): Clear.

(2) \Rightarrow (1): First, we prove that $H_1(\mathcal{L}_{\bullet}^{n,d}) = 0$ (or even $[H_1(\mathcal{L}_{\bullet}^{n,d})]_{d+2} = 0$) implies $H_0(\mathcal{L}_{\bullet}^{n,d}) \cong I_{(n-d,d)}^{\text{Sp}}$. Note that $\mathcal{L}_{\bullet}^{n,d}$ is a subcomplex of a minimal free resolution P_{\bullet} of $I_{(n-d,d)}^{\text{Sp}}$ with $\mathcal{L}_0^{n,d} = P_0$. Hence if $\mathcal{L}_1^{n,d} = P_1$, then $H_0(\mathcal{L}_{\bullet}^{n,d}) = H_0(P_{\bullet}) \cong I_{(n-d,d)}^{\text{Sp}}$. By (3.1), we see that

$$\begin{aligned} \dim_K[\mathcal{L}_1^{n,d}]_{d+1} = \text{rank } \mathcal{F}_1^{(n-d,d)} &= \beta_{1,d+1}(I_{(n-d,d)}^{\text{Sp}}) - \beta_{0,d+1}(I_{(n-d,d)}^{\text{Sp}}) \\ &= \beta_{1,d+1}(I_{(n-d,d)}^{\text{Sp}}) = \dim_K[P_1]_{d+1}, \end{aligned}$$

and hence $[\mathcal{L}_1^{n,d}]_{d+1} = [P_1]_{d+1}$. By the first statement of Lemma 2.1, if $\mathcal{L}_1^{n,d} \neq P_1$ then $P_1 = \mathcal{L}_1^{n,d} \oplus R(-d-2)^m$ for some $m > 0$. By (3.1), we have $P_2 = \mathcal{L}_2^{n,d} \oplus R(-d-2)^m \oplus R(-d-3)^l$ (l can be 0). Consider the exact sequence

$$0 \longrightarrow \mathcal{L}_{\bullet}^{n,d} \longrightarrow P_{\bullet} \longrightarrow P_{\bullet}/\mathcal{L}_{\bullet}^{n,d} \longrightarrow 0.$$

Both $P_2/\mathcal{L}_2^{n,d}$ and $P_1/\mathcal{L}_1^{n,d}$ have the direct summand $R(-d-2)^m$. Since P_{\bullet} is minimal, we have $\text{Im } \partial_2^{P_{\bullet}} \subset \mathfrak{m}P_1$, and hence $\partial_2^{P_{\bullet}}$ induces the zero map between these two $R(-d-2)^m$. Hence $[H_2(P_{\bullet}/\mathcal{L}_{\bullet}^{n,d})]_{d+2} \neq 0$, but the above exact sequence induces $[H_1(\mathcal{L}_{\bullet}^{n,d})]_{d+2} = [H_2(P_{\bullet}/\mathcal{L}_{\bullet}^{n,d})]_{d+2} \neq 0$. This is a contradiction, and we have $m = 0$. Hence $H_0(\mathcal{L}_{\bullet}^{n,d}) \cong I_{(n-d,d)}^{\text{Sp}}$.

If (2) holds, $\mathcal{L}_{\bullet}^{n,d}$ coincides with the first $n - 2d$ steps of P_{\bullet} (since $H_{n-2d-1}(\mathcal{L}_{\bullet}^{n,d}) = 0$, the kernel of $\partial : \mathcal{L}_{n-2d-1}^{n,d} \longrightarrow \mathcal{L}_{n-2d-2}^{n,d}$ coincides with $\partial(\mathcal{L}_{n-2d}^{n,d})$). In particular, $\beta_{n-2d,n-d+1}(I_{(n-d,d)}^{\text{Sp}}) = 0$. By (3.1), we have $\beta_{n-2d+1,n-d+1}(I_{(n-d,d)}^{\text{Sp}}) = 0$, and hence

$\beta_{i,d+i}(I_{(n-d,d)}^{\text{Sp}}) = 0$ for all $i > n - 2d$. Summing up, $\beta_{i,j}(I_{(n-d,d)}^{\text{Sp}}) \neq 0$ and $i \leq n - 2d$ (resp. $i > n - 2d$) implies $j = d + i$ (resp. $j = d + i + 1$). The argument similar to the last step of the proof Proposition 4.2, we have $P_{\bullet} = \mathcal{F}_{\bullet}^{(n-d,d)}$. \square

The next lemma can be proved by a similar way to Proposition 4.3, and we leave the details to the reader.

Proposition 4.4. *For $n, d \in \mathbb{N}$ with $n = 2d + 1$, the following are equivalent.*

- (1) *Theorem 4.1 holds in this case.*
- (2) $[H_1(\mathcal{L}_{\bullet}^{n,d})]_{d+2} = 0$.

We regard \mathfrak{S}_{n-1} as the symmetric group on $\{1, \dots, n-1\}$. Regarding the \mathfrak{S}_n -module $V_{(n-d-i,d,1^i)}$ as an \mathfrak{S}_{n-1} -module in the natural way, we get the restriction $V_{(n-d-i,d,1^i)} \downarrow_{\mathfrak{S}_{n-1}}^{\mathfrak{S}_n}$. By the *branching rule* ([5, Theorem 2.8.3]), we have

$$V_{(n-d-i,d,1^i)} \downarrow_{\mathfrak{S}_{n-1}}^{\mathfrak{S}_n} \cong V_{(n-d-i-1,d,1^i)} \oplus V_{(n-d-i,d-1,1^i)} \oplus V_{(n-d-i,d,1^{i-1})}.$$

Here, if $(n-d-i-1, d, 1^i)$ is not a partition (i.e., if $n-d-i-1 < d$), we set $V_{(n-d-i-1,d,1^i)} = 0$. Similarly, if $i = 0$ then $V_{(n-d-i,d,1^{i-1})} = 0$.

It is easy to see that $\mathcal{F}_{\bullet}^{(n-1,1)}$ is a minimal free resolution of a complete intersection ideal $I_{(n-1,1)}^{\text{Sp}}$, and [7, Theorem 3.2] states that $\mathcal{F}_{\bullet}^{(n-2,2)}$ is a minimal free resolution of $I_{(n-2,2)}^{\text{Sp}}$. So we may assume that $d \geq 3$.

Recall that $\{e(T) \mid T \in \text{SYT}(n-d-i, d, 1^i)\}$ forms a basis of $V_{(n-d-i,d,1^i)}$. For $T \in \text{SYT}(n-d-i, d, 1^i)$, the box \boxed{n} is located in one of the following positions

- (i) the right end of the first row (if $n-d-i > d$),
- (ii) the right end of the second row,
- (iii) the bottom of the first column (if $i \geq 1$).

Removing the box \boxed{n} from T , we get the tableau T' . If \boxed{n} is in the position (i) (resp. (ii) or (iii)), then T' is a standard tableau of shape $(n-d-i-1, d, 1^i)$ (resp. $(n-d-i, d-1, 1^i)$ or $(n-d-i, d, 1^{i-1})$).

We consider the following subspaces of $V_i := V_{(n-d-i,d,1^i)}$.

$$U_i := \langle e(T) \mid T \in \text{SYT}(n-d-i, d, 1^i), \boxed{n} \text{ is in the position (i)} \rangle$$

$$W_i := \langle e(T) \mid T \in \text{SYT}(n-d-i, d, 1^i), \boxed{n} \text{ is in the positions (i) or (ii)} \rangle$$

Note that V_i/W_i has a basis

$$(4.1) \quad \{ \bar{e}(T) \mid T \in \text{SYT}(n-d-i, d, 1^i), \boxed{n} \text{ is in the position (iii)} \},$$

where $\bar{e}(T)$ denotes the natural image of $e(T)$. These vector spaces are closed under the \mathfrak{S}_{n-1} -action, and we have

$$U_i \cong V_{(n-d-i-1,d,1^i)}, \quad W_i/U_i \cong V_{(n-d-i,d-1,1^i)} \quad \text{and} \quad V_i/W_i \cong V_{(n-d-i,d,1^{i-1})}$$

as \mathfrak{S}_{n-1} -modules.

In the above notation, we have $\mathcal{L}_i^{n,d} = V_i \otimes_K R(-d-i)$. If $z \in U_i \otimes_K R(-d-i) \subset \mathcal{L}_i^{n,d}$, then it is easy to see that $\partial_i^{\mathcal{L}_i^{n,d}}(z) \in U_{i-1} \otimes_K R(-d-i+1) \subset \mathcal{L}_{i-1}^{n,d}$. Similarly, $z \in W_i \otimes_K R(-d-i)$ implies $\partial_i^{\mathcal{L}_i^{n,d}}(z) \in W_{i-1} \otimes_K R(-d-i+1)$. Hence, if we

set $A_i := U_i \otimes_K R(-d-i)$ and $B_i := W_i \otimes_K R(-d-i)$ for $i \geq 0$, A_\bullet and B_\bullet are subcomplexes of $\mathcal{L}_\bullet^{n,d}$. Clearly, these are complexes of \mathfrak{S}_{n-1} -modules.

Set $R' := K[x_1, \dots, x_{n-1}]$. Then $\mathcal{L}_\bullet^{n-1,d}$ and $\mathcal{L}_\bullet^{n-1,d-1}$ are chain complexes of graded free R' -modules.

Lemma 4.5. *With the above notation, we have the following.*

- (1) $A_\bullet \cong \mathcal{L}_\bullet^{n-1,d} \otimes_{R'} R$
- (2) B_\bullet/A_\bullet is isomorphic to a truncation of $\mathcal{L}_\bullet^{n-1,d-1}(-1) \otimes_{R'} R$, where (-1) denotes the degree shift for graded modules. More precisely, if we remove the “last term” $\mathcal{L}_{n-2d+1}^{n-1,d-1}(-1) \otimes_{R'} R$ from $\mathcal{L}_\bullet^{n-1,d-1}(-1) \otimes_{R'} R$, it is isomorphic to B_\bullet/A_\bullet .
- (3) $\mathcal{L}_\bullet^{n,d}/B_\bullet \cong \mathcal{L}_{\bullet-1}^{n-1,d}(-1) \otimes_{R'} R$.

Proof. (1) Easy.

(2) We have

$$\mathcal{L}_i^{n-1,d-1} \otimes_{R'} R \cong V_{(n-d-i,d-1,1^i)} \otimes_K R(-d-i+1)$$

for $0 \leq i \leq (n-1) - 2(d-1) = n - 2d + 1$. Similarly,

$$B_i/A_i \cong V_{(n-d-i,d-1,1^i)} \otimes_K R(-d-i)$$

for $0 \leq i \leq n - 2d$ (note that $\mathcal{L}_i^{n,d} = 0$ for $i > n - 2d$). The differential maps clearly coincide, and (2) holds.

(3) We have $\mathcal{L}_0^{n,d}/B_0 = 0$ and

$$\mathcal{L}_i^{n,d}/B_i \cong (V_i/W_i) \otimes_K R(-d-i) \cong V_{(n-d-i,d,1^{i-1})} \otimes_K R(-d-i),$$

for $1 \leq i \leq n - 2d$. Similarly,

$$\mathcal{L}_{i-1}^{n-1,d} \otimes_{R'} R \cong V_{(n-d-i,d,1^{i-1})} \otimes_K R(-d-i+1)$$

for $1 \leq i \leq n - 2d$. Let $\bar{e}(T)$ be an element of the basis (4.1) of V_i/W_i . If the first column of T has the entries $a_1, a_2, \dots, a_{i+2} (= n)$ from top to bottom, we have

$$\partial_i^{\mathcal{L}_\bullet^{n,d}}(e(T) \otimes 1) = \sum_{j=1}^{i+2} (-1)^{j-1} (e(T^j) \otimes x_{a_j}).$$

Since $e(T_{i+2}) \in W_{i-1}$ and $\bar{e}(T_{i+2}) = 0$, we have

$$\partial_i^{\mathcal{L}_\bullet^{n,d}/B_\bullet}(\bar{e}(T) \otimes 1) = \sum_{j=1}^{i+1} (-1)^{j-1} (\bar{e}(T^j) \otimes x_{a_j}).$$

This clearly coincides with $\partial_{i-1}^{\mathcal{L}_\bullet^{n-1,d}}$, and hence (3) holds. \square

The proof of Theorem 4.1. We use induction on n . Since the theorem has been shown for $d \leq 2$ in [7], the theorem holds for small n 's. We may also assume that $n > 2d$ by Proposition 4.2.

Clearly, $A_i = 0$ for $i \geq n - 2d$. By the induction hypothesis and Lemma 4.5, we have

- $\mathcal{F}_\bullet^{(n-d-1, d)}$ (resp. $\mathcal{F}_\bullet^{(n-d, d-1)}$) is a minimal free resolution of the Specht ideal $I_{(n-d-1, d)}^{\text{Sp}}$ (resp. $I_{(n-d, d-1)}^{\text{Sp}}$) of R' .
- A_\bullet is the first $n-2d-1$ steps of $\mathcal{F}_\bullet^{(n-d-1, d)} \otimes_{R'} R$. In particular, $H_i(A_\bullet) = 0$ for $0 < i < n-2d-1$.
- B_\bullet/A_\bullet is the first $n-2d$ steps of $\mathcal{F}_\bullet^{(n-d, d-1)}(-1) \otimes_{R'} R$. In particular, $H_i(B_\bullet/A_\bullet) = 0$ for $0 < i < n-2d$. (We have $H_i(\mathcal{L}_\bullet^{n-1, d-1}) = 0$ for $0 < i \leq n-2d$, but we have to exclude the case $i = n-2d$ by the truncation).

First, we will study $H_i(B_\bullet)$ using the exact sequence

$$(4.2) \quad 0 \longrightarrow A_\bullet \longrightarrow B_\bullet \longrightarrow B_\bullet/A_\bullet \longrightarrow 0.$$

Claim 1. If $n > 2d+1$, then $H_i(B_\bullet) = 0$ for $0 < i < n-2d$.

The proof of Claim 1. By (4.2) and the above observation, we have $H_i(B_\bullet) = 0$ for $0 < i < n-2d-1$, and it remains to prove $H_{n-2d-1}(B_\bullet) = 0$. Since $A_{n-2d} = 0$, we have the exact sequence

$$(4.3) \quad 0 \longrightarrow H_{n-2d}(B_\bullet) \xrightarrow{f} H_{n-2d}(B_\bullet/A_\bullet) \xrightarrow{g} H_{n-2d-1}(A_\bullet) \xrightarrow{h} H_{n-2d-1}(B_\bullet) \longrightarrow 0.$$

By the above observation,

$$H_{n-2d-1}(A_\bullet) = \text{Ker}(\partial_{n-2d-1}^{A_\bullet}) = \text{Ker}(\partial_{n-2d-1}^{\mathcal{F}_\bullet^{(n-d-1, d)}}) \otimes_{R'} R = \text{Im}(\partial_{n-2d}^{\mathcal{F}_\bullet^{(n-d-1, d)}}) \otimes_{R'} R.$$

Since $\mathcal{F}_{n-2d}^{(n-d-1, d)} \otimes_{R'} R \cong V_{((d-1)^2, 1^{n-2d+1})} \otimes_K R(-n+d-1)$, $H_{n-2d-1}(A_\bullet)$ is generated by its degree $(n-d+1)$ -part

$$[H_{n-2d-1}(A_\bullet)]_{n-d+1} \cong V_{((d-1)^2, 1^{n-2d+1})}.$$

On the other hand, since the ‘‘tail’’ of B_\bullet/A_\bullet is of the form

$$0 \longrightarrow \mathcal{L}_{n-2d}^{n-1, d-1}(-1) \otimes_{R'} R \longrightarrow \mathcal{L}_{n-2d-1}^{n-1, d-1}(-1) \otimes_{R'} R$$

by Lemma 4.5 (2) (while $\mathcal{L}_{n-2d+1}^{n-1, d-1} \neq 0$ and $H_{n-2d}(\mathcal{L}_\bullet^{n-1, d-1}) = 0$), we have

$$H_{n-2d}(B_\bullet/A_\bullet) \cong \text{Ker}(\partial_{n-2d}^{\mathcal{L}_\bullet^{n-1, d-1}})(-1) \otimes_{R'} R \cong \text{Im}(\partial_{n-2d+1}^{\mathcal{L}_\bullet^{n-1, d-1}})(-1) \otimes_{R'} R.$$

Since $\mathcal{L}_{n-2d+1}^{n-1, d-1} \otimes_{R'} R \cong V_{((d-1)^2, 1^{n-2d+1})} \otimes_K R(-n+d)$, we have

$$[H_{n-2d}(B_\bullet/A_\bullet)]_{n-d+1} \cong V_{((d-1)^2, 1^{n-2d+1})}.$$

Since, in the sequence (4.3),

$$\begin{array}{ccc} [g]_{n-d+1} : [H_{n-2d}(B_\bullet/A_\bullet)]_{n-d+1} & \longrightarrow & [H_{n-2d-1}(A_\bullet)]_{n-d+1} \\ & \cong \parallel & \cong \parallel \\ & V_{((d-1)^2, 1^{n-2d+1})} & V_{((d-1)^2, 1^{n-2d+1})} \end{array}$$

is an \mathfrak{S}_{n-1} -homomorphism, and $H_{n-2d-1}(A_\bullet)$ is generated by $[H_{n-2d-1}(A_\bullet)]_{n-d+1}$, we have

$$\begin{aligned} [g]_{n-d+1} \neq 0 &\implies [g]_{n-d+1} \text{ is bijective} \implies g \text{ is surjective} \\ &\implies h \text{ is injective} \implies H_{n-2d-1}(B_\bullet) = 0. \end{aligned}$$

So it suffices to show that $[g]_{n-d+1} \neq 0$.

Since $A_{n-2d} = B_{n-2d+1} = 0$, we have

$$H_{n-2d}(B_\bullet) = \text{Ker } \partial_{n-2d}^{B_\bullet} \quad \text{and} \quad H_{n-2d}(B_\bullet/A_\bullet) = (\partial_{n-2d}^{B_\bullet})^{-1}(A_{n-2d-1})$$

(both are submodules of B_{n-2d}), and f is the inclusion map.

As defined in the construction of the differential map of $\mathcal{F}_\bullet^{(n-d,d)}$, for a tableau

$$T = \begin{array}{|c|c|c|c|} \hline a_1 & b_2 & \cdots & b_{d-1} \\ \hline a_2 & c_2 & \cdots & c_{d-1} \\ \hline a_3 & & & \\ \hline \vdots & & & \\ \hline \end{array} \in \text{Tab}((d-1)^2, 1^{n-2d+1})$$

we have the tableau $T^i \in \text{Tab}(d, d-1, 1^{n-2d})$ for each $1 \leq i \leq n-2d+3$. Adding \boxed{n} to the right end of the second row of T^i , we get $\tilde{T}^i \in \text{Tab}(d^2, 1^{n-2d})$. Set

$$\alpha := \sum_{i \geq 1} (-1)^{i-1} e(\tilde{T}^i) \otimes x_{a_i} \in [W_{n-2d} \otimes_{R'} R(-n+d)]_{n-d+1} = [B_{n-2d}]_{n-d+1}.$$

We want to compute $\partial_{n-2d}^{B_\bullet}(\alpha)$. If $T', T'', T''' \in \text{Tab}(d+1, d, 1^{n-2d-1})$ are tableaux whose right most 3 entries are

$$T' = \begin{array}{|c|c|} \hline \cdots & a_i \quad a_j \\ \hline \cdots & n \\ \hline \end{array}, \quad T'' = \begin{array}{|c|c|} \hline \cdots & a_j \quad a_i \\ \hline \cdots & n \\ \hline \end{array} \quad \text{and} \quad T''' = \begin{array}{|c|c|} \hline \cdots & a_i \quad n \\ \hline \cdots & a_j \\ \hline \end{array},$$

and the rests are same, then $e(T') = e(T'') + e(T''')$ and $e(T''') \in A_{n-2d-1}$. Hence it is easy to see that $\partial_{n-2d}^{B_\bullet}(\alpha) \in A_{n-2d-1}$ and $\alpha \in H_{n-2d}(B_\bullet/A_\bullet)$. Moreover, the term

$$e \left(\begin{array}{|c|c|c|c|c|c|} \hline a_3 & b_2 & \cdots & b_{d-1} & a_1 & n \\ \hline a_4 & c_2 & \cdots & c_{d-1} & a_2 & \\ \hline \vdots & & & & & \\ \hline \end{array} \right) \otimes x_{a_1} x_{a_2}$$

survives in $\partial_{n-2d}^{B_\bullet}(\alpha)$. Hence $\partial_{n-2d}^{B_\bullet}(\alpha) \neq 0$, and it means that $\alpha \notin H_{n-2d}(B_\bullet) = \text{Im } f = \text{Ker } g$ and $g(\alpha) \neq 0$. We are done. \square

Claim 2. If $n = 2d + 1$, then $[H_1(B_\bullet)]_{d+2} = 0$.

The proof of Claim 2. By the same argument as in Claim 1, we have an injection $f : H_1(B_\bullet) \rightarrow H_1(B_\bullet/A_\bullet)$, and we see that $[H_1(B_\bullet/A_\bullet)]_{d+2} \cong V_{((d-1)^2, 1^2)}$. Since $V_{((d-1)^2, 1^2)}$ is a simple \mathfrak{S}_{n-1} -module, and f is an \mathfrak{S}_{n-1} -homomorphism, the degree $(d+2)$ -part $[f]_{d+2}$ is either a bijection or the zero map (then $[H_1(B_\bullet)]_{d+2} = 0$). So it suffices to show that $[H_1(B_\bullet)]_{d+2} \neq [H_1(B_\bullet/A_\bullet)]_{d+2}$.

As in Claim 1, we have $H_1(B_\bullet) = \text{Ker } \partial_1^{B_\bullet}$ and $H_1(B_\bullet/A_\bullet) = (\partial_1^{B_\bullet})^{-1}(A_0)$. So it suffices to show that there is some $\alpha \in [B_1]_{d+2}$ with $0 \neq \partial_1^{B_\bullet}(\alpha) \in A_0$. However, we can find such α in the same way as in Claim 1. \square

Let us return to the main line of the proof. The proof is divided into two cases. In both cases, we use the exact sequence

$$(4.4) \quad 0 \longrightarrow B_\bullet \longrightarrow \mathcal{L}_\bullet^{n,d} \longrightarrow \mathcal{L}_\bullet^{n,d}/B_\bullet \longrightarrow 0.$$

Case 1. First, we consider the case $n > 2d+1$. It suffices to prove that $H_i(\mathcal{L}_\bullet^{n,d}) = 0$ for $0 < i < n - 2d$ by Proposition 4.3. By Lemma 4.5 (3) and the induction hypothesis, we have $H_i(\mathcal{L}_\bullet^{n,d}/B_\bullet) = 0$ for $1 < i < n - 2d$. By (4.4), Claim 1 and the above observation, we have $H_i(\mathcal{L}_\bullet^{n,d}) = 0$ for $1 < i < n - 2d$, and it remains to show that $H_1(\mathcal{L}_\bullet^{n,d}) = 0$. By Lemma 4.5 (3) and the induction hypothesis, we have $H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet) \cong H_0(\mathcal{L}_\bullet^{n-1,d})(-1) \otimes_{R'} R \cong (I_{(n-d-1,d)}^{\text{Sp}})(-1) \otimes_{R'} R \cong (I_{(n-d-1,d)}^{\text{Sp}} R)(-1)$, where $I_{(n-d-1,d)}^{\text{Sp}} R$ is the extension of the ideal $I_{(n-d-1,d)}^{\text{Sp}} \subset R'$ to R . Hence we have the following exact sequence

$$0 \longrightarrow H_1(\mathcal{L}_\bullet^{n,d}) \xrightarrow{\varphi} (I_{(n-d-1,d)}^{\text{Sp}} R)(-1) \xrightarrow{\psi} H_0(B_\bullet) \longrightarrow H_0(\mathcal{L}_\bullet^{n,d}) \longrightarrow 0.$$

By Lemma 4.5 (1), (2) and the induction hypothesis, $H_0(A_\bullet) \cong I_{(n-d-1,d)}^{\text{Sp}} R$ and $H_0(B_\bullet/A_\bullet) \cong (I_{(n-d,d-1)}^{\text{Sp}} R)(-1)$. Since $H_1(B_\bullet/A_\bullet) = 0$ now, (4.2) yields

$$0 \longrightarrow I_{(n-d-1,d)}^{\text{Sp}} R \longrightarrow H_0(B_\bullet) \longrightarrow (I_{(n-d,d-1)}^{\text{Sp}} R)(-1) \longrightarrow 0,$$

and hence $\text{Ass}_R(H_0(B_\bullet)) = \{(0)\}$. Note that \mathfrak{S}_n acts on $H_1(\mathcal{L}_\bullet^{n,d})$, but does not act on $I_{(n-d-1,d)}^{\text{Sp}} R$. So φ cannot be an isomorphism. Since $I_{(n-d-1,d)}^{\text{Sp}} R \subset R$, $H_1(\mathcal{L}_\bullet^{n,d}) \neq 0$ implies that $\text{Im } \psi = \text{Coker } \varphi$ has an associated prime $\mathfrak{p} \neq (0)$. Since $\text{Im } \psi$ is a submodule of $H_0(B_\bullet)$, \mathfrak{p} is also an associated prime of $H_0(B_\bullet)$, and this is a contradiction. Hence we have $H_1(\mathcal{L}_\bullet^{n,d}) = 0$, and we are done.

Case 2. Finally, we consider the case $n = 2d + 1$. It suffices to show that $[H_1(\mathcal{L}_\bullet^{n,d})]_{d+2} = 0$ by Proposition 4.4. By Claim 2, the exact sequence (4.4) yields $\varphi : H_1(\mathcal{L}_\bullet^{n,d}) \rightarrow H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet)$ such that

$$[\varphi]_{d+2} : [H_1(\mathcal{L}_\bullet^{n,d})]_{d+2} \longrightarrow [H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet)]_{d+2}$$

is injective. Recall that $H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet) \cong H_0(\mathcal{L}_\bullet^{n-1,d})(-1) \otimes_{R'} R$. Now $\mathcal{F}_\bullet^{(n-d-1,d)}$ ($= \mathcal{F}_\bullet^{(d,d)}$) is a minimal free resolution of $I_{(d,d)}^{\text{Sp}} \subset R'$, $\mathcal{F}_0^{(d,d)} \cong \mathcal{L}_0^{n-1,d}$ and $\mathcal{L}_i^{n-1,d} = 0$ for $i > 0$, hence we have

$$H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet) \cong H_0(\mathcal{L}_\bullet^{n-1,d}) \otimes_{R'} R = \mathcal{L}_0^{n-1,d} \otimes_{R'} R = \mathcal{F}_0^{(d,d)} \otimes_{R'} R.$$

Since there is a graded submodule N of $\mathcal{F}_0^{(d,d)}$ such that $\mathcal{F}_0^{(d,d)}/N \cong I_{(d,d)}^{\text{Sp}}$ and N is generated by N_{d+2} , there is a submodule L ($\cong N(-1) \otimes_{R'} R$) of $H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet)$ such that L is generated by L_{d+3} and

$$H_1(\mathcal{L}_\bullet^{n,d}/B_\bullet)/L \cong I_{(d,d)}^{\text{Sp}}(-1) \otimes_{R'} R \cong (I_{(d,d)}^{\text{Sp}} R)(-1).$$

If $[H_1(\mathcal{L}_{\bullet}^{n,d})]_{d+2} \neq 0$, then we have a subideal $J \subset I_{(d,d)}^{\text{Sp}}R$ with

$$J(-1) \cong \text{Im } \varphi / (L \cap \text{Im } \varphi)$$

and $J_{d+1} \neq 0$. Since \mathfrak{S}_n acts on $J_{d+1} (\cong [\text{Im } \varphi]_{d+2} \cong [H_1(\mathcal{L}_{\bullet}^{n,d})]_{d+2})$, the ideal $J' := (J_{d+1})$ is a symmetric ideal. For a subsets $F \subset [n-1]$ (resp. $F \subset [n]$), set

$$P'_F := (x_i - x_j \mid i, j \in F) \subset R' \quad (\text{resp. } P_F := (x_i - x_j \mid i, j \in F) \subset R)$$

to be the prime ideal. Since

$$I_{(d,d)}^{\text{Sp}} = \bigcap_{\substack{F \subset [n-1] \\ \#F=d+1}} P'_F,$$

we have

$$I_{(d,d)}^{\text{Sp}}R = \bigcap_{\substack{F \subset [n-1] \\ \#F=d+1}} P_F.$$

Since $J' \subset I_{(d,d)}^{\text{Sp}}R$ is a symmetric ideal, we have

$$J' \subset \bigcap_{\substack{F \subset [n] \\ \#F=d+1}} P_F = I_{(d,d,1)}^{\text{Sp}}.$$

However, we have $J_{d+1} = J'_{d+1} \subset [I_{(d,d,1)}^{\text{Sp}}]_{d+1} = 0$. This is a contradiction. □

ACKNOWLEDGEMENTS

We are grateful to Professors Stephen Griffeth, Steven V Sam and Alexander Woo for giving us valuable information on the preceding works on Specht ideals, especially [1]. We also thank Professors Mitsuyasu Hashimoto and Satoshi Murai for stimulating discussion.

REFERENCES

- [1] C. Berkesch Zamaere, S. Griffeth, and S.V. Sam, Jack polynomials as fractional quantum Hall states and the Betti numbers of the $(k+1)$ -equals ideal, *Comm. Math. Phys.* **330** (2014), 415–434
- [2] D. Eisenbud, *Commutative Algebra with a view toward algebraic geometry*, Graduate texts in Mathematics 150, Springer-Verlag, 1995.
- [3] S.-Y.R. Li and W.C.W. Li, Independence numbers of graphs and generators of ideals, *Combinatorica* **1** (1981) 55–61.
- [4] C. McDaniel and J. Watanabe, Principal radical systems, Lefschetz properties and perfection of Specht Ideals of two-rowed partitions, to appear in *Nagoya Math. J.*, arXiv:2103.00759.
- [5] B.E. Sagan, *The Symmetric Group, Representations, Combinatorial Algorithms, and Symmetric Functions*, second edition, Graduate texts in Mathematics 203, Springer-Verlag, 2001.
- [6] K. Shibata and K. Yanagawa, Regularity of Cohen-Macaulay Specht ideals, *J. Algebra* **582** (2021), 73–87.
- [7] K. Shibata and K. Yanagawa, Elementary construction of minimal free resolutions of the Specht ideals of shapes $(n-2, 2)$ and $(d, d, 1)$, to appear in *J. Algebra its Appl.*, arXiv:2010.06522.

- [8] R. H. Villarreal, Monomial algebras, Second edition, Monographs and research notes in mathematics, CRC Press, Boca Raton FL, 2015.
- [9] J. Watanabe and K. Yanagawa, Vandermonde determinantal ideals, Math. Scand. **125** (2019), 179-184.
- [10] K. Yanagawa, When is a Specht ideal Cohen-Macaulay? J. Commut. Algebra **13** (2021), 589-608.

NATIONAL INSTITUTE OF TECHNOLOGY, YONAGO COLLEGE, YONAGO, TOTTORI, 683-8502, JAPAN

Email address: shibata@yonago.kosen-ac.jp

DEPARTMENT OF MATHEMATICS, KANSAI UNIVERSITY, SUITA, OSAKA 564-8680, JAPAN

Email address: yanagawa@kansai-u.ac.jp