

LOCAL COHOMOLOGY OF MODULE OF DIFFERENTIALS OF INTEGRAL EXTENSIONS II

S. P. DUTTA

ABSTRACT. In this note (R, m) denotes a complete regular local ring and B mostly denotes its absolute integral closure. The four objectives of this paper are the following: i) to determine the highest non-vanishing local cohomology of $\Omega_{B/R}$ in equicharacteristic 0, ii) to establish a connection between each of $\Omega_{B/R}$ and $\Omega_{B/V}$ and pull-back of $\Omega_{A/V}$ via a short exact sequence together with new observations on corresponding local cohomologies in mixed characteristic where V is the coefficient ring of R and A is its absolute integral closure, iii) to demonstrate that $\Omega_{B/R}$ can be mapped onto a cohomologically Cohen-Macaulay module and iv) to study torsion-free property for $\Omega_{C/V}$ and $\Omega_{C/k}$ along with their respective completions where C is an integral domain and a module finite extension of R . In this connection an extension of Suzuki's theorem on normality of complete intersections to the formal set-up in all characteristics is accomplished.

1. INTRODUCTION

This note is a sequel to our previous work in ([3]). Here (R, m) denotes a complete regular local ring of dimension n , B denotes its absolute integral closure unless otherwise stated. In ([3]) it was pointed out that $H_m^n(\Omega_{B/R}) = 0$ and in mixed characteristic $H_m^{n-1}(\Omega_{B/R}) \neq 0 \iff$ the direct summand property for integral extensions of regular local rings is valid and hence by André's work in ([1]) it follows that $H_m^{n-1}(\Omega_{B/R}) \neq 0$ (theorem 4.1, [3]). However, we were not sure that the above observation on non-vanishing of local cohomology ([6]) is valid in equicharacteristic 0 and we raised it as a question along with a couple of other questions in ([3]). In this paper we are mainly concerned with the following: i) to provide affirmative answers to some of the questions raised in ([3]) on highest non-vanishing local cohomology of $\Omega_{B/R}$ with respect to m and with respect to any ideal generated by a part of

AMS Subject Classification: 13D02, 13D22, 13D45, 13N05

Key words and phrases: module of differentials, integral extension, formal smoothness, local cohomology, absolute integral closure.

regular system of parameters of length $n - 1$ in equicharacteristic 0, ii) to establish a connection between each of $\Omega_{B/R}$ and $\Omega_{B/V}$ and pull back of $\Omega_{A/V}$ via a short exact sequence together with several new observations on corresponding local cohomologies in mixed characteristic where V is the coefficient ring of R and A is the absolute integral closure of V , iii) to demonstrate that in mixed characteristic $\Omega_{B/R}$ can be mapped onto cohomologically Cohen-Macaulay modules N such that for every $t > 0$, $(0 : p^t)N$ are finitely generated Cohen-Macaulay free $B/p^t B$ -modules of rank $n - 1$ and dimension $n - 1$ and iv) to study torsion free property for standard generators of $\Omega_{B/V}$, $\Omega_{B/k}$ and to study the same property for $\Omega_{C/V}$, $\Omega_{C/k}$ along with their respective completions where C is an integral domain and a module finite extension of R . In this connection an extension of Suzuki's theorem on normality of complete intersection equicharacteristic affine domains to the formal set-up in all characteristics is accomplished. Several new results are also discussed. A brief overview of most of the results of this paper is given below.

In section 2.1 first we derive the following **corollary** from theorem 3.5 in ([3]):

Let R be a normal domain containing a field k of characteristic 0 with a non-null derivation $D \in \text{Der}_k(R)$. Let C denote the integral closure of R in a finite extension of $Q(R)$ and $\tilde{C} = \{x \in Q(C) | \text{Tr}(xC) \subset R\}$ ([13]). Let $\tilde{D} : \Omega_{C/k} \rightarrow Q(C)$ denote the C -linear map induced by the extension of D to $Q(C)$. Then $\text{Im} \tilde{D} \subset \tilde{C}$.

We use the above result to prove the following theorem.

Theorem 2.2. Let (R, m) be a complete regular local ring of dimension n in equicharacteristic 0 and let B denote its absolute integral closure. Let x_2, \dots, x_n denote a part of a regular system of parameters of R and \underline{x} denote the corresponding ideal. Then $H_{\underline{x}}^{n-1}(\Omega_{B/R}) \neq 0$.

As a **corollary** (2.3) we derive that $H_m^{n-1}(\Omega_{B/R}) \neq 0$.

The above theorem settles affirmatively question 2a) in equicharacteristic 0 and the corollary settles affirmatively question 1) in (4.4, [3]).

Using the above corollary we are now able to prove the following result in mixed characteristic,

Proposition 2.5. The surjective map $H_m^{n-1}(\Omega_{B/R}) \rightarrow H_{\underline{x}}^{n-1}(\Omega_{B/R})$ is not an isomorphism (here p, x_2, \dots, x_n form a regular system of parameters of R).

In section 3 first we point out in lemma 3.1 that if $R_i = V[[X_1, \dots, X_i]]$, $i \leq n - 1$, $R = R_{n-1}$, $R_0 = V$, $B_i =$ absolute integral closure of R_i ,

$A = B_0$ and $B = B_{n-1}$, then each of $\Omega_{B_i/V}$ and Ω_{B_i/R_i} is a B_i -summand of $\Omega_{B/V}$ and $\Omega_{B/R}$ respectively. Similar statement is also valid in the equicharacteristic case.

For lemma 3.2 and corollaries of theorem 3.3 in 3.4, we assume that $[\mathbb{Q}(V) : \mathbb{Q}(\hat{\mathbb{Z}}_p\mathbb{Z})] < \infty$

Let \underline{X} denote the ideal generated by X_1, \dots, X_{n-1} in R .

Lemma 3.2 points out the following:

$$H_{\underline{X}}^{n-1}(\Omega_{A/V} \otimes B) \simeq H_m^{n-1}(\Omega_{A/V} \otimes B) \simeq H_m^n(B).$$

Thus the validity of the direct summand property for integral extensions ([7]) is equivalent to the non-null property for $H_{\underline{X}}^{n-1}(\Omega_{A/V} \otimes B)$.

In our next theorem we prove the following.

Theorem 3.3. The following sequence is exact:

$$0 \rightarrow \Omega_{A/V} \otimes B \rightarrow \Omega_{B/V} \rightarrow \Omega_{B/A} \rightarrow 0$$

In one of the two **corollaries** of this theorem we point out in 3.4 that

$$H_p^1(B) \text{ imbeds into both } \Omega_{B/V} \text{ and } \Omega_{B/R}.$$

This implies (Remark 3.4) that both $\Omega_{B/V}$ and $\Omega_{B/R}$ contain isomorphic copies of a Cohen-Macaulay torsion B -module. It follows from an exact sequence in proposition 2.5 or in theorem 4.1, ([3]) that both $\Omega_{B/V}$ and $\Omega_{B/k}$ also contain cohomologically Cohen-Macaulay flat B -modules. For the later observation no restriction on $Q(V)$ is necessary.

The final theorem of this section demonstrates the following.

Theorem 3.5. $\Omega_{B/R}$ can be mapped onto a B -module M such that i) M is cohomologically Cohen-Macaulay of cohomological dimension $n-1$, ii) for every $t > 0$, $(0 : p^t)M$ is finitely generated Cohen-Macaulay free $B/p^t B$ -module of rank $n-1$ and dimension $n-1$ and iii) $(0 : p^t)M$ is a summand of $(0 : p^t)\Omega_{B/R}$.

As a corollary we derive that, for every $t > 0$, $\Omega_{B/R}$ contains a finitely generated Cohen-Macaulay free $B/p^t B$ -module of rank $n-1$ and dimension $n-1$. Moreover, if $[\mathbb{Q}(V) : \mathbb{Q}(\hat{\mathbb{Z}}_p\mathbb{Z})] < \infty$, the corresponding rank of this submodule is n .

In section 4 we study torsion-free property for standard generators of each of $\Omega_{B/V}$, $\Omega_{B/k}$ and for each of $\Omega_{C/V}$, $\Omega_{C/k}$ where C is the integral closure of R in a finite extension of $Q(R)$. First we prove the following proposition.

Proposition 4.1. Let $b (\neq 0) \in B$; consider an element w of the form $cdb \in \Omega_{B/V}$, $c \in B$. If w is a torsion element in $\Omega_{B/V}$, then $b \in A$.

If R is equicharacteristic i.e. $R = k[[X_1, \dots, X_n]]$, then for every $b \in B - \bar{k}$, $db \in \Omega_{B/k}$ is torsion free over B where \bar{k} is the algebraic closure of k . Same arguments work in the affine case.

The above proposition induced me to inquire about conditions under which each of $\Omega_{B/V}$ and $\Omega_{B/k}$ would become torsion free as B -modules. This led me to Suzuki's theorem ([10]) (statement in (4.3)). Theorem 4.3 extends Suzuki's work on the connection between normality of complete intersection affine domains over a field k and torsion-freeness of the corresponding module of differentials to the formal set-up in all characteristics. To this end we first extend Grothendieck's result: Proposition 22.7.2 in ([5]) to mixed characteristic in the following way:

Theorem 4.2. Let V be a complete discrete valuation ring in mixed characteristic p with maximal ideal pV such that the residue field k of V is perfect. Let $R = V[[X_1, \dots, X_{n-1}]]$ and $C = R/I$ be an integral domain such that $p \neq 0$ in C . Let P be a prime ideal of R containing I and let $q = P/I$. The following conditions are equivalent.

- i) C_q is formally smooth over V for the q -adic topology.
- ii) There exists derivations $D_i, 1 \leq i \leq r, \in \text{Der}_V(R)$ and elements $\{f_i\}_{1 \leq i \leq r}$ of I such that $\{f_i\}_{1 \leq i \leq r}$ generate IR_P and $\det(D_i f_j) \notin P$.
- iii) C_q is an unramified regular local ring.

Next we prove the following extension of Suzuki's theorem.

Theorem 4.3. Let $C = R/I$ where (R, m) is either unramified or equicharacteristic complete regular local ring with coefficient ring V (field k) such that $V/pV(k)$ is perfect and I is a complete intersection prime ideal in R . Let $\widehat{\Omega}_{C/V}, \widehat{\Omega}_{C/k}$ denote the m -adic completions of $\Omega_{C/V}$ and $\Omega_{C/k}$ respectively. We have the following

- 1) Suppose that if $p \neq 0 \in C$, every prime ideal P of height 1 in C containing p is such that $p \notin P^2 C_P$. If C is normal then both $\widehat{\Omega}_{C/V}$ and $\widehat{\Omega}_{C/k}$ are torsion-free over C .

- 2) If $\widehat{\Omega}_{C/V}$ or $\widehat{\Omega}_{C/k}$ is torsion free then C is normal.

As a **corollary** we point out that, with R as above, if $C = R[[Y_1, \dots, Y_r]]/I$ is a module finite extension of R and is a complete intersection domain such that $Q(C)$ is a separable extension of $Q(R)$, then part 1) of the above theorem implies that $\Omega_{C/V}$ and $\Omega_{C/k}$ are torsion-free over C .

Our final **theorem** (4.5) points out the connection between derivations of R and the torsion sub-module T of $\widehat{\Omega}_{C/V}$, where C is as mentioned in the statement of theorem 4.2. We prove the following:

for any $\omega \neq 0$ in $\widehat{\Omega}_{C/V}$, $\omega \in T \iff \tilde{D}_i(\omega) = 0, \tilde{D}_i$ induced by the extension D_i of the standard derivation $\partial/\partial X_i : R \rightarrow R$ for $1 \leq i \leq n-1$ to $Q(C)$.

For similar observation in the equicharacteristic case see remark 4) in 4.7.

In **Corollary 3**, 4.6, we again point out that $\Omega_{B/R}$ has an image N such that N is cohomologically Cohen-Macaulay i.e. $H_m^i(N) = 0$ for $i < n - 1$, $H_m^{n-1}(N) \neq 0$ and $\Omega_{B/V}$ can be mapped on to a B -module W such that $H_m^i(W) = 0$ for $i \geq 0$. N shares the same properties with M as in theorem 3.5 and its corollary. Moreover, for any $x \neq 0 \in m$, $(0 : x)N$ can be imbedded into a cohomologically Cohen-Macaulay flat B/xB -module.

We provide construction of another such module in remark 5), 4.7.

We are thankful to B. Bhatt for his note (about a couple months back) through which we became aware of his awesome piece of work in [2]. We used some of his main results to draw our conclusions on Cohen-Macaulayness in several places within this paper.

Notations. Given an extension $A \rightarrow B$ of commutative rings, the corresponding module of differentials is denoted by $\Omega_{B/A}$ (instead of $\Omega_{B/A}^1$); d.v.r. stands for discrete valuation ring; for any integral domain (ring) C , $Q(C)$ denotes the field of fractions (quotient ring) of C ; for any local ring (R, m) , E denotes the injective hull of R/m over R , $E(k(P))$ denotes the injective hull of R/P over R and for any R -module M , M^\vee stands $\text{Hom}_R(M, E)$, M^* stands for $\text{Hom}_R(M, R)$. Unless stated otherwise, in a local ring by local cohomology of a module we mean the same with respect to the maximal ideal.

SECTION 2.

2.1. For the convenience of the reader we state theorem 3.5, ([3]) and add a sentence or two for indicating its proof.

Theorem. (Theorem 3.5, [3]) Let R be a normal domain of dimension n containing a field k of characteristic 0 with a non-null derivation $D \in \text{Der}_k(R)$ and let B be its integral closure in an algebraic extension F of K . Then $\text{Hom}_R(\Omega_{B/k}, R) \neq 0$. If (R, m) is a complete local normal domain of dimension n with maximal ideal m and a non-null derivation $D \in \text{Der}_k(R)$, then $H_m^n(\Omega_{B/k}) \neq 0$.

The proof of this theorem is obtained by proving the following assertion: Let K denote the field of fractions of R . Since characteristic of $k = 0$, D can be extended to a derivation D (same notation): $B \rightarrow F$ and hence to a derivation: $B_P \rightarrow F$ for any prime ideal P of R . Let $\text{Tr} : F \rightarrow K$ denote the trace map. We define an R -linear map $L : \Omega_{B/k} \rightarrow K$ in the following way:

For any $\omega \in \Omega_{B/k}$, $L(\omega) = \text{Tr}(\tilde{D}(\omega))$ where $\tilde{D} : \Omega_{B/k} \rightarrow F$ is the B -linear map induced by D ; the same prescription works for $\Omega_{B^P/k} \rightarrow F$ for any prime ideal P of R . Then $\text{im}L \subset R$.

See ([3]) for details.

Next we derive the following corollary.

Corollary. Let R be a normal domain containing a field k of characteristic 0 with a non-null derivation $D \in \text{Der}_k(R)$. Let C denote the integral closure of R in a finite extension $Q(C)$ of $Q(R)$. Let $\tilde{C} = \{x \in Q(C) | \text{Tr}(xC) \subset R\}$ ([13]). Then $\tilde{D}(\Omega_{C/k}) \subset \tilde{C}$, where the C -linear map $\tilde{D} : \Omega_{C/k} \rightarrow Q(C)$ is induced by the extension of D to $Q(C)$.

Proof. Recall in the proof of the above theorem we have shown that if $L = \text{Tr}_\bullet \tilde{D} : \Omega_{C/k} \rightarrow Q(R)$, then $\text{im}L \subset R$. Let $\lambda \in \text{im} \tilde{D}$, i.e. $\lambda = \tilde{D}(w)$ for some $w \in \Omega_{C/k}$. For any $b \in C$, $b\lambda = b\tilde{D}(w) = \tilde{D}(bw)$. Hence by above the theorem, it follows that $\text{Tr}(b\lambda) = \text{Tr}(\tilde{D}(bw)) \in R$; this implies that $\lambda \in \tilde{C}$. Thus the corollary follows.

2.2. In the following theorem and its corollary we are going to answer affirmatively a question that was raised in ([3]).

Theorem. Let $R = k[[X_1, \dots, X_n]]$, characteristic of $k = 0$. Let B be the integral closure of R in an algebraic extension of $Q(R)$ such that B contains an element y with the property that $y^2 = X_i$, for some i , $1 \leq i \leq n$. Let \underline{X} denote the ideal $(X_1, \dots, \hat{X}_i, \dots, X_n)$ ($\hat{} \equiv$ omitted). Then $H_{\underline{X}}^{n-1}(\Omega_{B/R}) \neq 0$.

Proof. Can assume $i = 1$. Let $K = Q(R)$ and $F = Q(B)$. Let $D : B \rightarrow F$ denote the extension of $\partial/\partial X_1 : R \rightarrow R$, $\partial/\partial X_1(X_i) = 0$ for $i \neq 1$ and $= 1$ for $i = 1$. Let $\tilde{D} : \Omega_{B/k} \rightarrow F$ denote the B -linear map induced by D . By hypothesis,

$$y^2 = X_1 \Rightarrow 2ydy = dX_1 \Rightarrow \tilde{D}(dy) = 1/2y \text{ and}$$

$$X_1 dy = 0 \in \Omega_{B/R} \dots \dots \dots (1).$$

To prove our assertion it will be enough to show that for $t > 0$, $dy(X_2, \dots, X_n)^{t-1} \notin (X_2^t, \dots, X_n^t)\Omega_{B/R}$. If possible, let

$$dy(X_2, \dots, X_n)^{t-1} = \sum_{i \geq 2} X_i^t w_i, w_i \in \Omega_{B/R}, 2 \leq i \leq n \dots \dots \dots (2)$$

Consider the short exact sequence (theorem 2.3, lemma 2.4, [3]).

$$0 \rightarrow \Omega_{R/k} \otimes B \rightarrow \Omega_{B/k} \xrightarrow{\eta} \Omega_{B/R} \rightarrow 0$$

Due to lemma in theorem 3.6, [3] we have a split exact sequence

$$0 \rightarrow \bigcap_{s>0} m^s \Omega_{R/k} \rightarrow \Omega_{R/k} \rightarrow \bigoplus_1^n R dX_i \rightarrow 0$$

Tensoring with B we obtain from above the following split exact sequence

$$0 \rightarrow \bigcap_{s>0} m^s \Omega_{R/k} \otimes B \rightarrow \Omega_{R/k} \otimes B \rightarrow \bigoplus_1^n B dX_i \rightarrow 0.$$

Let $\tilde{w}_i \in \Omega_{B/k}$ be such that $\eta(\tilde{w}_i) = w_i$. The following equality then follows from (2):

$$dy(X_2, \dots, X_n)^{t-1} = \sum_{i \geq 2} X_i^t \tilde{w}_i + \sum_{i \geq 1} b_i dx_i + \lambda, b_i \in B,$$

$$\lambda \in \bigcap_{s>0} m^s \Omega_{R/k} \otimes B \dots \dots \dots (3)$$

Note that $\tilde{D}(\lambda) = 0$. Applying $\tilde{D} : \Omega_{B/k} \rightarrow F$ to (3) we obtain:

$$\tilde{D}(dy)(X_2, \dots, X_n)^{t-1} = \sum X_i^t \tilde{D}(\tilde{w}_i) + b_1$$

From (1), it follows that

$$(X_2, \dots, X_n)^{t-1} = 2y \sum X_i^t \tilde{D}(\tilde{w}_i) + 2yb_1 \dots \dots \dots (4)$$

Let $S = k[[X_1, \dots, X_n, Y]]/(Y^2 - X_1) \simeq k[[X_2, \dots, X_n, Y]]$.

For any finite extension C of S contained in B such that $b_1 \in C$ and $\tilde{w}_i \in \Omega_{C/k}, 2 \leq i \leq n$, we have $\text{Tr}_{Q(C)/Q(R)}(\mu \tilde{D}(\tilde{w}_i) | \mu \in C) \subset R$ (theorem 2.1). By corollary in 2.1 we obtain $\tilde{D}(\tilde{w}_i) \subset \tilde{C}, 2 \leq i \leq n$.

Since $\text{Tr}_{Q(C)/Q(R)} = \text{Tr}_{Q(S)/Q(R)} \bullet \text{Tr}_{Q(C)/Q(S)}$, it follows by the above lemma that $\text{Tr}_{Q(C)/Q(S)}(\tilde{D}(\tilde{w}_i)) \subset \tilde{S} = R\{1/2y, y/2y\}, 2 \leq i \leq n$, where $\tilde{S} = \{z \in Q(S) | \text{Tr}(zS) \subset R\}$; \tilde{S} is a free R -module generated by $\{1/2y, y/2y\}$ ([9], [13]).

Applying $\text{Tr}_{Q(C)/Q(S)}$ to (4) we obtain (ignoring the units):

$$(X_2, \dots, X_n)^{t-1} = \sum X_i^t 2y \text{Tr}_{Q(C)/Q(S)}(\tilde{D}(\tilde{w}_i)) + 2y \text{Tr}_{Q(C)/Q(S)}(b_1).$$

It follows from above that

$$(X_2, \dots, X_n)^{t-1} = \sum_{i \geq 2} X_i^t s_i + 2yc_1, s_i, c_1 \in S$$

. This contradicts the monomial property over S ([6]). Hence $H_{\underline{X}}^{n-1}(\Omega_{B/R}) \neq 0$.

2.3. Corollary. Let (R, m) and B be as above. Then $H_m^{n-1}(\Omega_{B/R}) \neq 0$.

Proof. We continue with our notations from the above theorem. Let us recall that localization commutes with module of differentials even in non-noetherian situations. By corollary 2.1 in ([3]) we have $H_{\underline{X}}^{n-1}(\Omega_{B/R}[1/X_1]) = 0$. This implies, due to the above theorem and the following exact sequence

$$H_m^{n-1}(\Omega_{B/R}) \rightarrow H_{\underline{X}}^{n-1}(\Omega_{B/R}) \rightarrow H_{\underline{X}}^{n-1}(\Omega_{B/R}[1/X_1]) \rightarrow 0$$

that $H_m^{n-1}(\Omega_{B/R}) \neq 0$.

Remark. Since $H_m^n(\Omega_{B/R}) = 0$ (corollary. 2.1, [3]), the above corollary implies that the highest non-vanishing local cohomology of $\Omega_{B/R}$

occurs at the $(n-1)$ -th stage. A proof of this observation can also be obtained without applying corollary 2.1.

2.4. Due to corollary 2 lemma 2.1 in [3] we know if (R, m) is a complete local normal domain of dimension n of equicharacteristic 0 and B is an integral domain integral over R , then $H_m^n(\Omega_{B/R}) = 0$. In this perspective we have the following proposition.

Proposition. Let (R, m) be a complete regular local ring of dimension n of equicharacteristic 0 and B be an integral domain integral over R . Suppose B contains a square root of a regular parameter of R . Then for any system of parameters y_1, \dots, y_n of R , $\Omega_{B/R} \otimes B/yB = \Omega_{(B/yB)/(R/y)} \neq 0$; here y denotes the ideal generated by y_1, \dots, y_n .

Proof. Let $R = k[[X_1, \dots, X_n]]$ and let $\underline{X} = (X_1, \dots, X_n)$. It is enough to show that $\Omega_{(B/\underline{X}B)/k} \neq 0$. Let $y \in B$ be such that $y^2 = X_1$; then

$$2ydy = dX_1 \text{ i.e. } 2X_1dy = ydX_1 \text{ in } \Omega_{B/k}.$$

If possible let $\text{im } dy = 0$ in $\Omega_{(B/\underline{X}B)/k}$.

Consider the exact sequence (theorem 2.3, lemma 2.4, [3]):

$$0 \rightarrow \Omega_{R/k} \otimes B \rightarrow \Omega_{B/k} \rightarrow \Omega_{B/R} \rightarrow 0$$

Tensoring the sequence with $B/\underline{X}B$ we obtain the following exact sequence:

$$\rightarrow \Omega_{R/k} \otimes B/\underline{X}B \rightarrow \Omega_{B/k} \otimes B/\underline{X}B \rightarrow \Omega_{(B/\underline{X}B)/k} \rightarrow 0$$

$\text{im } dy = 0$ in $\Omega_{(B/\underline{X}B)/k} \Rightarrow$

$dy = \sum X_i w_i + \sum b_i dX_i + \lambda$ in $\Omega_{B/k}$, $b_i \in B$, $w_i \in \Omega_{B/k}$ for $1 \leq i \leq n$ and $\lambda \in \cap m^t \Omega_{R/k} \otimes B$.

Let $\tilde{D} : \Omega_{B/k} \rightarrow Q(B)$ denote the B -linear map induced by the extension of $\partial/\partial X_1 : R \rightarrow R$. Applying \tilde{D} to the above equation, we obtain

$$1/2y = \sum X_i D(w_i) + b_1.$$

This implies

$$1 - 2yb_1 = \sum X_i 2y \tilde{D}(w_i).$$

Since R is complete, B is local and $1 - 2yb_1$ is unit, say u in B . Thus

$$1 = \sum X_i 2iu^{-1} \tilde{D}(w_i).$$

Applying the trace map we obtain via theorem 2.1.

$$1 = \sum X_i r_i, r_i \in R \text{ — a contradiction.}$$

Remark. This result is valid even if R is not complete provided B is local.

2.5. Here we will use theorem 2.3 to prove an observation in the mixed characteristic.

Proposition 2.5. Let $R = V[[X_1, \dots, X_{n-1}]]$, $m = (p, X_1, \dots, X_{n-1})$ and (V, pV) is a complete discrete valuation ring. let B denote the absolute integral closure of R and let \underline{X} denote the ideal generated by X_1, \dots, X_{n-1} in R . Then the map $\alpha : H_{\underline{X}}^{n-2}(\Omega_{B/R}[1/p]) \rightarrow H_m^{n-1}(\Omega_{B/R})$ is non-null.

Proof. Let us recall that the above map is part of a long exact sequence:

$$H_{\underline{X}}^{n-2}(\Omega_{B/R}) \rightarrow H_{\underline{X}}^{n-2}(\Omega_{B/R}[1/p]) \xrightarrow{\alpha} H_m^{n-1}(\Omega_{B/R}) \rightarrow H_{\underline{X}}^{n-1}(\Omega_{B/R}) \dots \dots (5)$$

Let $y \in B$ be such that $y^p = X_1 \Rightarrow py^{p-1}dy = dX_1 \Rightarrow pX_1dy = ydX_1 \in \Omega_{B/V} \Rightarrow pX_1dy = 0 \in \Omega_{B/R}$.

Consider the element $\lambda = (dy(X_2, \dots, X_{n-1})^{t-1}, 0 \dots 0) \in \oplus_1^{n-1} \Omega_{B/R}[1/p]$.

Note that $X_1^t dy(X_2, \dots, X_{n-1})^{t-1} = X_1 dy(X_1, \dots, X_{n-1})^{t-1} = 0 \in \Omega_{B/R}[1/p]$. Thus λ represents a non-null element in $H^{n-1}(\underline{X}^t, \Omega_{B/R}[1/p])$ (proof of theorem 2.2). Since $\Omega_{B/R}(\Omega_{B/V}) = p\Omega_{B/R}(\Omega_{B/V})$ (theorem 4.1, [3]), we have an exact sequence

$$0 \rightarrow H_p^0(\Omega_{B/R}) \rightarrow \Omega_{B/R} \rightarrow \Omega_{B/R}[1/p] \rightarrow 0$$

. It follows that $H_{\underline{X}}^i(H_p^0(\Omega_{B/R})) \simeq H_m^i(\Omega_{B/R})$ for $i \geq 0$. Then $\alpha(\text{class of } \lambda) = \text{class of } X_1 dy(X_1, \dots, X_{n-1})^{t-1} \in H_{\underline{X}}^{n-1}(H_p^0(\Omega_{B/R}))$. So $\alpha(\text{class of } \lambda) = 0 \Rightarrow$:

$$X_1 dy(X_1, \dots, X_{n-1})^{t-1} = \sum_{i \geq 2} X_i^t w_i, \quad w_i \in (0 : p^t)\Omega_{B/R}, 2 \leq i \leq n-1 \dots (6)$$

We have an exact sequence (theorem 2.3, lemma 2.4, [3]):

$$0 \rightarrow \Omega_{R/V} \otimes B \rightarrow \Omega_{B/V} \rightarrow \Omega_{B/R} \rightarrow 0$$

Applying $\text{Hom}_B(B/p^t B, -)$ we obtain the following short exact sequence:

$$0 \rightarrow (0 : p^t)\Omega_{B/V} \rightarrow (0 : p^t)\Omega_{B/R} \xrightarrow{\beta} \Omega_{R/V} \otimes B/p^t B \rightarrow 0 \dots (7)$$

Applying β to (6) we obtain the following equality from equ (7):

$$p^{t-1} y dX_1(X_1, \dots, X_{n-1})^{t-1} = \sum_{i \geq 1} X_i^t \mu_i \in \Omega_{R/V} \otimes B/p^t B, \quad \mu_i \in \Omega_{R/V} \otimes B, 1 \leq i \leq n.$$

Hence,

$$p^{t-1} y dX_1(X_1, \dots, X_{n-1})^{t-1} = \sum_{i \geq 1} X_i^t \mu_i + p^t \nu, \quad \nu \in \Omega_{R/V} \otimes B \dots \dots (8)$$

Let D_1 be the extension of $\partial/\partial X_1 : R \rightarrow R$ to $B \rightarrow Q(B)$ and let $\tilde{D}_1 : \Omega_{B/k} \rightarrow Q(B)$ denote the corresponding B-linear map.

Applying \tilde{D}_1 to (8) we obtain

$p^{t-1}y(X_1, \dots, X_{n-1})^{t-1} = \Sigma X_i^t b_i + p^t c$ where $b_i, c \in B, 1 \leq i \leq n$.

Since $y^p = X_1$, the above equality contradicts the monomial property in B . Hence $\text{im}\alpha \neq 0$.

Remark. Since $H_{\underline{X}}^{n-1}(\Omega_{B/R}[1/p]) = 0$ (corollary 2.1, [3]), the above proposition shows that the surjective map $H_m^{n-1}(\Omega_{B/R}) \rightarrow H_{\underline{X}}^{n-1}(\Omega_{B/R})$ is not an isomorphism. Let us mention that we are still not able to prove that $H_{\underline{X}}^{n-1}(\Omega_{B/R}) \neq 0$ in mixed characteristic (raised as a question in [3]).

SECTION 3.

3.1. Lemma. Let $R_i = V[[X_1, \dots, X_i]]$, $1 \leq i \leq n-1$, $R_0 = V$ and $R_{n-1} = R$. Let B, B_i denote the absolute integral closure of R, R_i respectively for $0 \leq i \leq n-1$, $B_0 = A$. Then each of $\Omega_{B_i/V}$ and Ω_{B_i/R_i} is B_i -summand of $\Omega_{B/V}$ and $\Omega_{B/R}$ respectively for $0 \leq i < n-1$.

Similar result is also valid in equicharacteristic zero.

Proof. Let Q_i be a prime ideal of B lying over $(X_{i+1}, \dots, X_{n-1})$ in R . Since B is absolutely integrally closed, we have $Q_i = Q_i^2$, B/Q_i is the absolute integral closure of R_i and hence $B_i \simeq B/Q_i$. Thus $B_i \xrightarrow{J_i} B \xrightarrow{\tilde{\eta}_i} B/Q_i$ is an R_i -algebra isomorphism. We have the following exact sequences :

$$Q_i/Q_i^2 \rightarrow \Omega_{B/V} \otimes B/Q_i \rightarrow \Omega_{(B/Q_i)/V} \rightarrow 0$$

and

$$Q_i/Q_i^2 \rightarrow \Omega_{B/R} \otimes B/Q_i \rightarrow \Omega_{(B/Q_i)/R_i} \rightarrow 0.$$

it follows from above that

$$\Omega_{B/V} \otimes B/Q_i \simeq \Omega_{B_i/V}$$

and

$$\Omega_{B/R} \otimes B/Q_i \simeq \Omega_{B_i/R_i}.$$

Thus the composite maps:

$$\Omega_{B_i/V} \xrightarrow{\tilde{J}_i} \Omega_{B/V} \xrightarrow{\tilde{\eta}_i} \Omega_{(B/Q_i)/V}$$

and

$$\Omega_{B_i/R} \xrightarrow{\tilde{J}_i} \Omega_{B/R} \xrightarrow{\tilde{\eta}_i} \Omega_{(B/Q_i)/R_i}$$

are B_i -module isomorphism. Hence the result.

3.2. For the rest of section 3 we assume $R = V[[X_1, \dots, X_{n-1}]]$, V -a complete discrete valuation ring in mixed characteristic p such that V/pV is perfect; let A, B denote the absolute integral closures of V, R respectively. For lemma 3.2 and corollary 2 in 3.4 we assume that $[Q(V) : Q_p] < \infty, Q_p = Q(\hat{Z}_p Z)$.

Since $\Omega_{Q(A)/Q(V)} = 0$, $\Omega_{A/V}$ is a p -torsion module. In a more general setup Fontaine [4] showed that $\Omega_{A/V}$ is generated by $\{d\xi_t\}_{t \geq 1}$, where ξ_t is a p^t -th root of unity such that $\xi_{t+1}^p = \xi_t$ for $t > 0$. We have the following lemma.

Lemma. $\Omega_{A/V} \otimes B \simeq H_p^1(B)$ and $H_m^{n-1}(\Omega_{A/V} \otimes B) \simeq H_m^n(B)$.

Proof. Since $\Omega_{A/V}$ is a p -torsion module, $\Omega_{A/V} \simeq \varinjlim (0 : p^t)\Omega_{A/V}$. In ([12]) it was derived from Fontaine’s work that $(0 : p^t)\Omega_{A/V} \simeq A/p^t A$. We have the following commutative diagram:

$$\begin{array}{ccc} (0 : p^t)\Omega_{A/V} & \longrightarrow & A/p^t A \\ \downarrow & & \downarrow \\ (0 : p^{t+1})\Omega_{A/V} & \longrightarrow & A/p^{t+1} A \end{array}$$

Where the horizontal arrows are isomorphisms, the right vertical arrow is multiplication by p and the left vertical arrow is the inclusion map.

It follows from above that $\Omega_{A/V} \simeq \varinjlim A/p^t A = H_p^1(A)$.

Hence $\Omega_{A/V} \otimes B \simeq H_p^1(B)$.

Moreover $H_m^{n-1}(\Omega_{A/V} \otimes B) \simeq H_m^{n-1}(H_p^1(B)) \simeq H_m^n(B)$

Corollary. The validity of the monomial property for local rings is equivalent to the condition that $H_m^{n-1}(\Omega_{A/V} \otimes B) \neq 0$, i.e. $H_{\underline{X}}^{n-1}(\Omega_{A/V} \otimes B) \neq 0$

3.3. Our next observation is the following.

Theorem. Let $AR = A \otimes_V R$. With notations as above the following sequence is exact:

$$0 \rightarrow \Omega_{A/V} \otimes_A B \xrightarrow{\alpha} \Omega_{B/R} \rightarrow \Omega_{B/AR} \rightarrow 0$$

Proof. The following sequence is exact

$$\Omega_{A/V} \otimes_A B \xrightarrow{\alpha} \Omega_{B/R} \rightarrow \Omega_{B/AR} \rightarrow 0.$$

We need to show that α is injective.

We have $A = \varinjlim A_t$ where, for every t , A_t is a complete d.v.r. and a module finite extension of V ;

$A_t = V[Y]/(f_t(Y))$, $f_t(Y)$ is a monic irreducible polynomial in $V[Y]$ ([9]).

$\Omega_{A/V} = \varinjlim \Omega_{A_t/V}$. It is enough to show that

$\Omega_{A_t/V} \otimes B \rightarrow \Omega_{B/R}$ is injective for every t (9).

$B = \varinjlim C$, C is a module finite extension of R .

$$\Omega_{B/R} = \varinjlim \Omega_{C/R}. \quad A_t R = A_t \otimes_V R \simeq A_t[[X_1, \dots, X_{n-1}]] \simeq R[Y]/(f_t(Y));$$

$A_t R$ is a complete regular local ring.

$$\Omega_{A_t/V} \otimes_{A_t} B \simeq \Omega_{A_t/V} \otimes_{A_t} A_t R \otimes_{A_t R} B;$$

$$\Omega_{A_t/V} \otimes_{A_t} A_t R \simeq \Omega_{A_t/V} \otimes_V R \simeq \Omega_{A_t R/R}.$$

Write $f = f_t(Y)$, $f' = f'_t(Y)$ and $S = A_t R = R[Y]/(f)$. We need to show $\Omega_{S/R} \otimes_S B \rightarrow \Omega_{B/R}$ is injective.

It is enough to show that $\Omega_{S/R} \otimes_S C \rightarrow \Omega_{C/R}$ is injective where C is any normal domain module finite extension of S (hence a module finite extension of R).

$$\Omega_{S/R} = S/f'S; \text{ hence } \Omega_{S/R} \otimes_S C = C/f'C.$$

We have an exact sequence ([11]):

$$\Gamma_{C/S} \xrightarrow{\beta} \Omega_{S/R} \otimes_S C \xrightarrow{v} \Omega_{C/R} \rightarrow \Omega_{C/S} \rightarrow 0$$

we need to show v is injective i.e. to show that $\text{im}\beta = 0$.

Since S and C are complete local normal domains and C is a module finite extension of S , for every prime ideal P of height 1 in S , C_P is a Dedekind domain module finite extension of S_P . Hence, by lemma 2.2 in [3] $(\Gamma_{C/S})_P = \Gamma_{C_P/S_P} = 0$. This implies that $\text{im}\beta$ has at least codimension 1 in $\Omega_{S/R} \otimes_S C = C/f'C$. Since C is a complete local normal domain, $C/f'C$ can't have any sub-module of codimension ≥ 1 in $C/f'C$. Thus $\text{im}\beta = 0$ and the assertion follows.

3.4. Corollaries.

1. There exists a commutative diagram of short exact sequences

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \Omega_{A/V} \otimes B & \xlongequal{\quad} & \Omega_{A/V} \otimes B & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \Omega_{R/V} \otimes B & \longrightarrow & \Omega_{B/V} & \longrightarrow & \Omega_{B/R} \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Omega_{R/V} \otimes B & \longrightarrow & \Omega_{B/A} & \longrightarrow & \Omega_{B/AR} \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

Proof. The exactness of the middle row follows from theorem 2.3 and lemma 2.4 in [3]. Since $\Omega_{A/V} \otimes B$ is a p -torsion module and $\Omega_{R/V} \otimes B$ is B -flat and hence torsion free over B , the assertion follows. For local cohomological implications see Remarks 2 in (4.7).

2. There exist exact sequences:

$$0 \rightarrow H_p^1(B) \rightarrow H_p^0(\Omega_{B/V}) \rightarrow H_p^0(\Omega_{B/A}) \rightarrow 0$$

and

$$0 \rightarrow H_p^1(B) \rightarrow H_p^0(\Omega_{B/R}) \rightarrow H_p^0(\Omega_{B/AR}) \rightarrow 0$$

.

Proof. By the above corollary we have an exact sequence:

$$0 \rightarrow \Omega_{A/V} \otimes_A B \rightarrow \Omega_{B/V} \rightarrow \Omega_{B/A} \rightarrow 0.$$

This implies, due to lemma (3.2), and the fact that $\Omega_{A/V} = p\Omega_{A/V}$, the exactness of the first exact sequence.

The second exact sequence is also proved by similar arguments.

Remark. Corollary 2 above implies, due to corollary 5.10 [2], that both $\Omega_{B/V}$ and $\Omega_{B/R}$ contains a Cohen-Macaulay torsion B -module. From middle row of the diagram in corollary 1 above it follows, due to theorem 5.1 [2] and flatness of $\Omega_{R/V}$ over R , that $\Omega_{B/V}$ also contains a cohomologically Cohen-Macaulay flat B -module. For the second assertion no restriction on $Q(V)$ is necessary.

3.5. In our final theorem of this section we prove the following.

Theorem. With V, R, B as in 3.2, $\Omega_{B/R}$ can be mapped onto a B -module M such that i) M is cohomologically Cohen-Macaulay of cohomological dimension $n - 1$, ii) for every $t > 0$, $(0 : p^t)M$ is finitely generated Cohen-Macaulay free $B/p^t B$ -module of rank $n - 1$ and dimension $n - 1$ and iii) $(0 : p^t)M$ is a summand of $(0 : p^t)\Omega_{B/R}$.

Proof. We have a short exact sequence:

$$0 \rightarrow H_p^0(\Omega_{B/V}) \rightarrow \Omega_{B/V} \rightarrow \Omega_{B/V}[1/p] \rightarrow 0.$$

(the exactness follows from the fact that $\Omega_{B/V} \otimes B/pB = 0$ (theorem 4.1, [3]))

Let M be the cokernel of $(H_p^0(\Omega_{B/V}) \rightarrow \Omega_{B/R})$

Consider the following commutative diagram of short exact sequences:

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & H_p^0(\Omega_{B/V}) & \xlongequal{\quad} & H_p^0(\Omega_{B/V}) & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & \Omega_{R/V} \otimes B & \longrightarrow & \Omega_{B/V} & \longrightarrow & \Omega_{B/R} \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Omega_{R/V} \otimes B & \longrightarrow & \Omega_{B/V}[1/p] & \longrightarrow & M \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

The exactness of the left column follows from the fact that $\Omega_{R/V}$ being a flat R -module ([3]), $\Omega_{R/V} \otimes B$ is a flat B -module and hence torsion-free over B .

Since $H_m^i(\Omega_{B/V}[1/p]) = 0$ for $i \geq 0$ and $\Omega_{R/V}$ is R -flat (theorem 2.3, [3]), it follows from the last row of the above diagram that $H_m^i(M) \simeq \Omega_{R/V} \otimes H_m^{i+1}(B) \simeq \bigoplus_1^{n-1} H_m^{i+1}(B)$ (lemma, theorem 3.6, [3]) for $i \geq 0$(*)

in particular $H_m^{n-1}(M) \simeq \bigoplus_1^{n-1} H_m^n(B)$.

By theorem 5.1, [2] it follows that $H_m^i(B) = 0$ for $i < n$; hence $H_m^i(M) = 0$ for $i < n - 1$ and $H_m^{n-1}(M) \neq 0$ ([1], [7]). Thus M is cohomologically Cohen-Macaulay of cohomological dimension $n - 1$. From the last row of the above commutative diagram we obtain $(0 : p^t)M \simeq \Omega_{R/V} \otimes B/p^t B \simeq \bigoplus_1^{n-1} B/p^t B$ (lemma within theorem 3.6, [3]). Since, by corollary 5.10 [2], $B/p^t B$ is Cohen-Macaulay, our assertion follows.

Corollary. For every $t > 0$, $\Omega_{B/R}$ contains a finitely generated Cohen-Macaulay free $B/p^t B$ -module of rank $n - 1$ and dimension $n - 1$. If $[Q(V) : Q(\hat{Z}_{pZ})] < \infty$, then the corresponding rank is n .

Proof follows from: a) the above commutative diagram by applying $\text{Hom}(B/p^t B, -)$ to the last row and the last column of the above diagram, b) corollary 2, 3.4 and c) corollary 5.10, [2].

Remark. Since $H_m^{n-1}(\Omega_{B/R}) \rightarrow H_m^{n-1}(M) \rightarrow 0$ is exact, the observation in (*) also leads to a new proof of the fact (theorem 4.1, [3]) that

$H_m^{n-1}(\Omega_{B/R}) \neq 0$ if and only if the direct summand property is valid over R .

SECTION 4.

4.1. Our first proposition deals with torsion-free property of standard generators of $\Omega_{B/V}$.

Proposition. Let $R = V[[X_1, \dots, X_{n-1}]]$ or $k[[X_1, \dots, X_n]]$ or $k[X_1, \dots, X_n]$ where characteristic of k is 0. Let B denote an integral extension of R in an algebraic extension of $Q(R)$. If for some $c \in B, b \in B - V, cdb \in \Omega_{B/V}$ is a torsion element, then $b \in A \cap B$ where A is the absolute integral closure of V .

If R , as above, contains a field k of characteristic 0 then for $b \in B - \bar{k}, db$ is torsion-free, where \bar{k} is the algebraic closure of k in $Q(B)$.

Proof. Let $R = V[[X_1, \dots, X_{n-1}]]$. Let $b \in B$ be such that db is a non-null torsion element in $\Omega_{B/V}$. We have an exact sequence

$$0 \rightarrow \Omega_{R/V} \otimes B \xrightarrow{\beta} \Omega_{B/V} \rightarrow \Omega_{B/R} \rightarrow 0$$

Let $c' \in B - (0)$ be such that $c'db = 0$. Let $\tilde{c} = c'$; $\tilde{c}db = 0$. \tilde{c} satisfies an integral equation over R : $\tilde{c}^s + r'_1\tilde{c}^{s-1} + \dots + r'_s = 0$, $r'_i \in R, 1 \leq i \leq s$ and $r'_s \neq 0$. This implies that $r'_s db = 0$. We write r for r'_s i.e. $rdb = 0, r \neq 0 \in R$.

Since $Q(B)$ is separable algebraic over $Q(R)$, if $f(X)$ denotes the minimal polynomial for b over R then $f'(b) \neq 0$. Hence

$$f'(b)rdb = 0 \dots \dots \dots (10)$$

Let $f(X) = X^t + r_1X^{t-1} + \dots + r_t$. Then

$$f(b) = b^t + r_1b^{t-1} + \dots + r_t = 0 \dots \dots \dots (11)$$

Hence $f'(b)db + \sum d(r_i)b^{t-i} = 0$ in $\Omega_{B/V}$. This implies, due to (10)

$$r\sum d(r_i)b^{t-i} = 0 \dots \dots \dots (12)$$

Since $\Omega_{R/V}$ is flat over R and B is an integral domain, $\Omega_{R/V} \otimes_R B$ is torsion-free B -module. From the above exact sequence we obtain, by (12),

$$\sum d(r_i) \otimes b^{t-i} = 0 \text{ in } \Omega_{R/V} \otimes_R B, \text{ i.e. } \sum d(r_i)b^{t-i} = 0 \in \Omega_{B/V}.$$

$R[b]$ is a free R -module of rank t (due to (10)) with basis $1, b, \dots, b^{t-1}$. We have an isomorphism

$$\eta : \oplus_1^t R \rightarrow R[b], \eta(e_i) = b^{t-i} \dots \dots \dots (13)$$

Tensoring (13) with $\Omega_{R/V}$ we get an isomorphism,

$$\tilde{\eta} : \bigoplus_1^t \Omega_{R/V} \rightarrow \Omega_{R/V} \otimes_R R[b], \quad \tilde{\eta}(d(s_1), \dots, d(s_t)) = \Sigma d(s_i) \otimes b^{t-i}.$$

Since $\Omega_{R/V}$ is R -flat, $\alpha : \Omega_{R/V} \otimes_R R[b] \rightarrow \Omega_{R/V} \otimes_R B$ is injective.

Composing this with the injection $\beta : \Omega_{R/V} \otimes_R B \rightarrow \Omega_{B/V}$ we have an injective map $\beta \cdot \alpha : \Omega_{R/V} \otimes_R R[b] \rightarrow \Omega_{B/V}$.

Since $\beta \cdot \alpha(\Sigma d(r_i) \otimes b^{t-i} = \Sigma d(r_i)b^{t-i}) = 0$, and $\tilde{\eta}$ is an isomorphism we have $d(r_i) = 0, 1 \leq i \leq t$. This implies that $r_i \in V, 1 \leq i \leq t$.

Hence from (11) it follows that b is integral over V and thus $b \in A$.

If R is of equicharacteristic 0, the above arguments show that for every $b \in B - \bar{k}$, db is torsion-free over B .

4.2. As mentioned in the introduction proposition 4.1 led us to look for sufficient conditions for torsion-free property of the module of differentials as a whole. And in this respect we were guided by Suzuki's work ([10]). Our main goal in 4.2 and 4.3 is to prove a formal version in all characteristics of Suzuki's theorem on the connection between normality of complete intersection affine domains over a field k and torsion-freeness of the corresponding module of differentials. For this purpose we first extend Grothendieck's proposition 22.7.2 in ([5]) to mixed characteristic. Grothendieck's arguments work in this case with several minor modifications.

Theorem. Let V be a complete discrete valuation ring in mixed characteristic p with maximal ideal pV such that the residue field k of V is perfect. Let $R = V[[X_1, \dots, X_{n-1}]]$ and $C = R/I$ be an integral domain such that $p \neq 0$ in C . Let P be a prime ideal of R containing I and let $q = P/I$. The following conditions are equivalent.

- i) C_q is formally smooth over V for the q -adic topology.
- ii) There exists derivations $D_i, 1 \leq i \leq r, \in \text{Der}_V(R)$ and elements $\{f_i\}_{1 \leq i \leq r}$ of I such that $\{f_i\}_{1 \leq i \leq r}$ generate IR_P and $\det(D_i f_j) \notin P$.
- iii) C_q is an unramified regular local ring.

Proof. i) \iff iii). Since V is a d.v.r. and C_q is a domain, $V \rightarrow C_q$ is formally smooth if and only if $V/pV \rightarrow C_q/pC_q$ is formally smooth. Since V/pV is perfect this is equivalent to C_q/pC_q being a regular local ring, i.e. C_q is an unramified regular local ring.

The proof of i) \iff ii) becomes evident by following Grothendieck's arguments in his proof of 22.7.2 in ([5]) with relevant minor modifications needed for the mixed characteristic case. Grothendieck's assumption for ensuring separability is assured here by assuming that the residue field k of V is perfect.

First we recall two basic facts that will be needed in our proof.

a) If p is in J for any prime ideal J of R then $p \notin J^2R_J$ ([8]).

b) $\widehat{\Omega}_{R/V} \simeq \bigoplus_1^{n-1} RdX_i, 1 \leq i \leq n - 1$, where $\widehat{\Omega}_{R/V}$ is the m -adic completion of $\Omega_{R/V}, m =$ the maximal ideal of R ([3], [5]).

Let $L = k(p) = R_P/PR_P = C_q/qC_q$. By (22.6.2, (ii), [5]) the condition i) is equivalent to the condition

$d_0 : I/I^2 \otimes L \rightarrow \Omega_{R/V} \otimes L$ is injective, d_0 is induced by the standard differential map "d". Let us consider the composite map

$$\hat{d} : I/I^2 \otimes L \rightarrow \Omega_{R/V} \otimes L \rightarrow \widehat{\Omega}_{R/V} \otimes L \dots \dots \dots (14)$$

and show that condition ii) is equivalent to the assertion that \hat{d} is injective. Note that $I/I^2 \otimes_C L = IR_P/I^2R_P \otimes_{RP} L$. Condition ii) implies

that, if $F_i = \hat{d}(\bar{f}_i \otimes 1)$, where \bar{f}_i is the image of f_i in IR_P/I^2R_P , the matrix $(\langle F_i, D_j \otimes 1 \rangle)$ is invertible. This implies $\{F_i\}$ are linearly independent, and hence, so are $\{\bar{f}_i \otimes 1\}$ which generate $IR_P/I^2R_P \otimes_{RP} L$.

Thus \hat{d} is injective. Inversely suppose \hat{d} is injective and let $\{f_i \in I\}_{1 \leq i \leq r}$ be such that $\{\bar{f}_i \otimes 1\}$ form a basis of $IR_P/I^2R_P \otimes_{RP} L$. Then $\{F_i\}$ form

a basis of $\text{Im}(\hat{d})$. By fact b), $\widehat{\Omega}_{R/V}$ is a free R -module of rank $(n - 1)$ and the V -derivations of R generate its dual. The fact that $\{F_i\}$ are linearly independent implies the existence of V -derivations D_i of R into itself such that the matrix $(\langle F_i, D_j \otimes 1 \rangle)$ is invertible as stated in ii).

i) \Rightarrow ii). We need to show that i) implies that \hat{d} is injective.

Note that \hat{d} is the composite of

$I/I^2 \otimes L \xrightarrow{\alpha} P/(P^2 + pR) \otimes L \xrightarrow{\beta} \widehat{\Omega}_{R/V} \otimes L$, where β is induced by the standard differential map "d".

Since R_P is a regular local ring and IR_P, p form a part of a regular system of parameters of R_P , it follows that the natural injection α is injective. Hence it is enough to show that

$$P/(P^2 + pR) \otimes L \xrightarrow{\beta} \widehat{\Omega}_{R/V} \otimes L \dots \dots \dots (15)$$

is injective.

We have an exact sequence

$$P/(P^2 + pR) \rightarrow \Omega_{R/V} \otimes R/P \rightarrow \Omega_{(R/P)/V} \rightarrow 0.$$

Since (R, m) is a Zarishki ring, it follows from (20.7.20, [5]) that

$$P/(P^2 + pR) \rightarrow \widehat{\Omega}_{R/V} \otimes R/P \rightarrow \widehat{\Omega}_{(R/P)/V} \rightarrow 0$$

is exact. And hence

$$P/(P^2 + pR) \otimes L \xrightarrow{\beta} \widehat{\Omega}_{R/V} \otimes L \rightarrow \widehat{\Omega}_{(R/P)/V} \otimes L \rightarrow 0 \dots \dots (16)$$

is exact.

It follows from (theorem 3.6, [3]) that, since $p \in P$, $\text{rank}(\widehat{\Omega}_{(R/P)/V} \otimes L) = \dim R/P$ and $\widehat{\Omega}_{R/V} \otimes L$ is of rank $(n - 1)$. Moreover, since $\dim P/(P^2 + pR) \otimes L = \text{height } P - 1$, it follows from (16) that

$$0 \rightarrow P/(P^2 + pR) \otimes L \xrightarrow{\beta} \widehat{\Omega}_{R/V} \otimes L \rightarrow \widehat{\Omega}_{(R/P)/V} \otimes L \rightarrow 0$$

is exact.

Hence our proof is complete.

4.3. Next we extend Suzuki's theorem to the formal set-up ([10]).

Suzuki's theorem. Let R be a complete intersection locality over a field k . Then R is normal if and only if Ω_{R/k^*} is torsion-free, where $k^* = k$ if characteristic of $k = 0$ or a differential constant field of R if the characteristic is positive.

Our theorem is the following.

Theorem. Let $C = R/I$, where (R, m) is either unramified or equicharacteristic complete regular local ring with coefficient ring V (field k) such that $V/pV(k)$ is perfect and I is a complete intersection prime ideal of R . Let $\widehat{\Omega}_{C/V}$, $\widehat{\Omega}_{C/k}$ denote the m -adic completions of $\Omega_{C/V}$, $\Omega_{C/k}$ respectively. We have the following.

i) For mixed-characteristic assume that if $p \neq 0$ in C , then every prime ideal P of height 1 in C containing p is such that $p \notin P^2 C_P$. If C is normal then $\widehat{\Omega}_{C/V}$ and $\widehat{\Omega}_{C/k}$ are torsion-free over C .

ii) If $\widehat{\Omega}_{C/V}$ or $\widehat{\Omega}_{C/k}$ is torsion-free then C is normal.

Proof. If $p \in I$, our assertion reduces to the equicharacteristic case. We here prove the mixed characteristic case; the equicharacteristic case would follow by similar arguments. Our goal is to reduce this formal case to a situation where arguments similar to those given by Suzuki ([10]) would work. Let $R = V[[X_1, \dots, X_{n-1}]]$ and let $I = (f_1, \dots, f_r)$. In order to attain such reduction, for part i) our main tool is theorem 4.2, in particular the observation (14) in the proof; and for part ii) several observations from ([3]) and ([5]).

i) Let $d : R \rightarrow \Omega_{R/V}$ be the usual derivation. Let $g (\neq 0) \in R$; then $g = \lim g_t$ where g_t is the sum of terms of g of degree $\leq t$. We have $dg = \lim dg_t$; $dg_t = \sum_{i=1}^{n-1} (\partial g_t / \partial X_i) dX_i$, $\lim dg_t = \sum_{i=1}^{n-1} \lim (\partial g_t / \partial X_i) dX_i = \sum_{i=1}^{n-1} (\partial g / \partial X_i) dX_i$ (by continuity). Since $\Omega_{R/V}$ is not Hausdorff, $dg = \sum_{i=1}^{n-1} (\partial g / \partial X_i) dX_i + \mu$ where $\mu \in \cap m^t \Omega_{R/V}$. let us recall that the following

sequence is split exact (lemma in theorem 3.6, [3]):

$$0 \rightarrow \cap m^t \Omega_{R/V} \rightarrow \Omega_{R/V} \xrightarrow{\eta} \widehat{\Omega}_{R/V} (= \bigoplus_1^{n-1} R dX_i) \rightarrow 0$$

We also denote by d the composite of $R \rightarrow \Omega_{R/V} \rightarrow \widehat{\Omega}_{R/V}$; then, for any $f \in R, df = \Sigma(\partial f/\partial X_i)dX_i$. Since (R, m) is a Zariski ring, it can be checked easily that the exact sequence

$$I/I^2 \rightarrow \Omega_{R/V} \otimes C \xrightarrow{\alpha} \Omega_{C/V} \rightarrow 0$$

gives rise to the following exact sequence.

$$I/I^2 \xrightarrow{\bar{d}} \widehat{\Omega}_{R/V} \otimes C \xrightarrow{\hat{\alpha}} \widehat{\Omega}_{C/V} \rightarrow 0 \dots \dots \dots (17)$$

First we assume that C is normal. If possible, let $w \in \widehat{\Omega}_{C/V}$ and $c(\neq 0) \in C$ be such that $cw = 0$. Then it follows from (17) that

$$c \sum_{i=1}^{n-1} \lambda_i \bar{d}X_i = \sum_{j=1}^r \nu_j \bar{d}f_j \dots \dots \dots (18)$$

where $\hat{\alpha}(\Sigma \lambda_i \bar{d}X_i) = w, \bar{d} = d \otimes Id_C, \lambda_i, \nu_j \in C$; hence

$$c \Sigma \lambda_i \bar{d}X_i = \sum_{j=1}^d \nu_j \sum_{i=1}^{n-1} (\partial f_j / \partial X_i) \bar{d}X_i \dots \dots \dots (19)$$

Since $\bar{d}X_1, \dots, \bar{d}X_{n-1}$ are linearly independent in $\widehat{\Omega}_{R/V} \otimes C$, we have

$$\lambda_i = 1/c \sum_{j=1}^d \nu_j (\partial f_j / \partial X_i) \dots \dots \dots (20)$$

. We need to show that $\nu_j/c \in C, 1 \leq j \leq r$. Then from (18) it would follow that $w = 0 \in \widehat{\Omega}_{C/V}$.

Let P be a prime ideal of height 1 of C . Then C_P is a regular local ring. By our assumption C_P is formally smooth over V Let $L = k(P)$ -the field of fractions of C_P . By the proof of proposition 4.2, C_P is formally smooth over V if and only if

$$I/I^2 \otimes L \xrightarrow{\bar{d}_L} \widehat{\Omega}_{R/V} \otimes L$$

is injective, i.e.

$$I/I^2 \otimes L \xrightarrow{\bar{d}_L} \bigoplus_{i=1}^{n-1} L dX_i \dots \dots \dots (21)$$

is injective.

Due to equation (20), this implies that certain $r \times r$ minor of $(\partial f_j / \partial X_i)$ $1 \leq j \leq r, 1 \leq i \leq n - 1$ is invertible and hence $\nu_j/c \in C_P$ for $1 \leq j \leq$

d. Since C is normal, $C = \cap C_P$ and thus $\nu_j/c \in C$ for $1 \leq j \leq d$.

ii) R is geometrically regular over V . Hence $\Gamma_{R/V} = 0$ (theorem. 2.3 [3]). We have an exact sequence ([11])

$$0 \rightarrow \Gamma_{C/V} \rightarrow I/I^2 \xrightarrow{\bar{d}} \Omega_{R/V} \otimes C \rightarrow \Omega_{C/V} \rightarrow 0 \dots (22)$$

Since $R(C)$ is complete and I is a complete intersection ideal of codimension r , I/I^2 is a finitely generated free complete C -module of rank r . Hence, $\Gamma_{C/V}$ and $\text{im} \bar{d}$ are also finitely generated complete C -modules. taking the m-adic completion we obtain an exact sequence (20.7.20, [5])

$$0 \rightarrow N \rightarrow I/I^2 \xrightarrow{\bar{d}} \widehat{\Omega}_{R/V} \otimes C \rightarrow \widehat{\Omega}_{C/V} \rightarrow 0 \dots (23)$$

where $\Gamma_{C/V} \subset N$ (we are not changing the notation for \bar{d}).

Since the $\text{rank}_R \widehat{\Omega}_{R/V} = \dim R - 1$ and $\text{rank}_C \widehat{\Omega}_{C/V} = \dim C - 1$ (21.9.5 [5], theorem 3.6, [3]), via rank calculation it follows that $N = 0$ and hence $\Gamma_{C/V} = 0$. Thus

$$0 \rightarrow I/I^2 \xrightarrow{\bar{d}} \widehat{\Omega}_{R/V} \otimes C \rightarrow \widehat{\Omega}_{C/V} \rightarrow 0 \dots (24)$$

is exact.

Let us assume that C is not normal. Let \bar{C} denote the normalization of C in $Q(C)$ and let $J = (C : \bar{C}) = \text{ann}_C \bar{C}/C$. Let P be a prime ideal of height 1 in C containing J . Then C_P is not regular. By proposition 4.2, $\text{rank jac}(f_1, \dots, f_r) \text{ mod } P < r$. The arguments for the rest of the proof is similar to that given by Suzuki for his proof of the affine case. There exist b_1, \dots, b_r of C , at least one of which, say b_1 , is not in P such that $\sum_{i=1}^r b_i \bar{d}f_i = 0$ in $\widehat{\Omega}_{R/V} \otimes C/P$.

Let $\tilde{J} = J : P$. There exist an element $j \in \tilde{J}\bar{C}$ such that $j \notin C_P \dots \dots \dots (25)$

Let $j = c/g, c, g \in C$. We have
 $j \sum b_i \bar{d}f_i \in j P \widehat{\Omega}_{R/V} \otimes C \subset J \bar{C} \widehat{\Omega}_{R/V} \otimes C \subset \widehat{\Omega}_{R/V} \otimes C$

It follows from above and (24) that $j \sum b_i \bar{d}f_i = \sum c_i \bar{d}X_i, c_i \in C$
 i.e. $c \sum b_i \bar{d}f_i = g \sum c_i \bar{d}X_i$ in $\widehat{\Omega}_{R/V} \otimes C \dots \dots \dots (26)$

From (24) it follows that $\text{im}(g \sum c_i \bar{d}X_i) = 0$ in $\widehat{\Omega}_{C/V}$. If $\text{im}(\sum c_i \bar{d}X_i) \neq 0$ in $\widehat{\Omega}_{C/V}$, then this is a torsion element in $\widehat{\Omega}_{C/V}$. If $\text{im}(\sum c_i \bar{d}X_i) = 0$ in

$\widehat{\Omega}_{C/V}$, then it follows from (24) that there are r elements $s_1, \dots, s_r \in C$ such that $\Sigma c_i \bar{d}X_i = \Sigma s_i \bar{d}f_i$.

Hence it follows from (26) that $\Sigma(gs_i - cb_i)\bar{d}f_i = 0$.

Due to (24) $\bar{d}f_1, \dots, \bar{d}f_r$ are linearly independent in $\widehat{\Omega}_{R/V} \otimes C$. Hence $gs_i = cb_i, 1 \leq i \leq r$, i.e. $j = c/g = s_i/b_i, 1 \leq i \leq r$. Since $b_1 \notin P$, it follows that $j \in C_P$ - a contradiction due to (25).

Corollary. Let (R, m) be an unramified or an equicharacteristic complete regular local ring and let $C = R[[X_1, \dots, X_n]]/I$ be a module finite integral extension of R such that I is a complete intersection prime ideal. Suppose that a) $Q(C)/Q(R)$ is separable, b) $\dim R > 1$ if R is mixed characteristic, $\dim R > 0$ if R is equicharacteristic and c) in mixed-characteristic for any prime ideal P of height 1 in C containing p , $p \notin P^2C/P$. If C is normal, then $\Omega_{C/V}$ and $\Omega_{C/k}$ are torsion-free over C .

Proof. We have a commutative diagram of short exact sequences

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
& & \cap m^t \Omega_{R/V} \otimes C & \equiv & \cap m^t \Omega_{R/V} \otimes C & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \Omega_{R/V} \otimes C & \longrightarrow & \Omega_{C/V} & \longrightarrow & \Omega_{C/R} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \parallel \\
0 & \longrightarrow & \widehat{\Omega}_{R/V} \otimes C & \longrightarrow & \widehat{\Omega}_{C/V} & \longrightarrow & \Omega_{C/R} \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0 & &
\end{array} \tag{27}$$

$\Omega_{C/R}$, being a finitely generated C -module, is complete. Since $\Omega_{R/V}$ is a flat R -module and $0 \rightarrow \cap m^t \Omega_{R/V} \rightarrow \Omega_{R/V} \rightarrow \widehat{\Omega}_{R/V} \rightarrow 0$ splits, $\cap m^t \Omega_{R/V} \otimes C$ is a flat C -module. Since C is an integral domain the assertion follows immediately.

4.4. Remarks. 1. The above diagram shows that for any module-finite extension C of a power series ring R such that $Q(C)$ is separably algebraic over $Q(R)$, the torsion sub-modules of $\Omega_{C/V}$ and $\Omega_{C/k}$ imbed

into $\widehat{\Omega}_{C/V}$ and $\widehat{\Omega}_{C/k}$) respectively and the corresponding local cohomology modules of $\Omega_{C/V}$ and $\Omega_{C/k}$ are isomorphic to the same of $\widehat{\Omega}_{C/V}$ and $\widehat{\Omega}_{C/k}$ respectively.

2. With notations as above let B denote the absolute integral closure of $C(R)$. Let $\widetilde{\Omega}_{B/V} = \varinjlim \widehat{\Omega}_{C/V}$, $R \subset C \subset B, C$ as in Remark 1 above, in mixed characteristic/equicharacteristic 0. Then the torsion sub-module of $\Omega_{B/V}$ imbeds into $\widetilde{\Omega}_{B/V}$ and the corresponding local cohomology modules of $\Omega_{B/V}$ and $\widetilde{\Omega}_{B/V}$ are isomorphic. This follows from (27) by taking direct limit and the fact that $\cap m^t \Omega_{R/V}$ is R -flat and $\cap m^t \Omega_{R/V} \otimes R/pR = 0$

4.5. Our final theorem points out a connection between torsion sub-module of $\widehat{\Omega}_{C/V}$ and derivations $\partial/\partial X_i, 1 \leq i \leq n-1$, of R .

Theorem. Let C be a complete local domain in mixed characteristic and $R = V[[X_1, \dots, X_{n-1}]] \hookrightarrow C$ be such that C is a module finite extension of R . Let m denote the maximal ideal of R and let $\widehat{\Omega}_{C/V}$ denote the m -adic completion of $\Omega_{C/V}$. Let $T =$ the torsion sub-module of $\widehat{\Omega}_{C/V}$. Then, for any $\omega \in \widehat{\Omega}_{C/V}$, $\omega \in T \iff \widetilde{D}_i(\omega) = 0$ for $1 \leq i \leq n-1$, where $\widetilde{D}_i : \Omega_{C/V}$ or $\widehat{\Omega}_{C/V} \rightarrow Q(C)$ is a C -linear map induced by the extension of the derivation $\partial/\partial X_i : R \rightarrow R, \partial/\partial X_i(X_j) = \delta_{ij}$, to $D_i : C \rightarrow Q(C)$.

Proof. \implies : is obvious.

\impliedby : let \widehat{d} denote the composite of $C \xrightarrow{d} \Omega_{C/V} \rightarrow \widehat{\Omega}_{C/V} (R \xrightarrow{d} \Omega_{R/V} \rightarrow \widehat{\Omega}_{R/V})$. It follows from the middle column of (27) that $(\cap m^t \Omega_{R/V}) \otimes C \simeq \cap m^t \Omega_{C/V}$.

It has been pointed out in the beginning of proof of part i), theorem 4.3 that for $r \in R$, $dr = \Sigma(\partial r/\partial X_i)dX_i + \mu$ where $\mu \in \cap m^t \Omega_{R/V}$ and $\widehat{d}r = \Sigma(\partial r/\partial X_i)dX_i$.

Let $\alpha (\neq 0) \in C$. First we want to show that

$f'(\alpha)d\alpha = f'(\alpha)\Sigma D_j(\alpha)dX_j + \nu, \nu \in \cap m^t \Omega_{C/V}$, where $f(X) \in R[X]$ denotes the minimal polynomial for α .

Let $f(X) = X^h + r_1X^{h-1} + \dots + r_h, f(\alpha) = 0$.

Then $f'(\alpha)d\alpha + \sum_{i=1}^h d(r_i)\alpha^{h-i} = 0$.

For $1 \leq j \leq n-1$, $f'(\alpha)D_j(\alpha) + \sum_{i=1}^h D_j(r_i)\alpha^{h-i} = 0$.

Hence $f'(\alpha)d\alpha - \sum_{j=1}^{n-1} f'(\alpha)D_j(\alpha)dX_j$

$$= \sum_{i=1}^h [d(r_i) - \sum_{j=1}^{n-1} D_j(r_i)dX_j]\alpha^{h-i}$$

$$= \sum \nu_i \alpha^{h-i}, \nu_i = [d(r_i) - \sum_{j=1}^{n-1} D_j(r_i)dX_j] \in \cap m^t \Omega_{R/V}.$$

Hence $f'(\alpha)d\alpha - \sum_{j=1}^{n-1} f'(\alpha)D_j(\alpha)dX_j \in \cap m^t \Omega_{R/V} \otimes C = \cap m^t \Omega_{C/V} \dots (28)$

Now let $\omega \in \Omega_{C/V}$; can write $\omega = \sum_{i=1}^t \lambda_i d\alpha_i$, $\lambda_i, \alpha_i \in C$. Let $f_i(X) \in R[X]$ denote the minimal polynomial of α_i and let $f = \pi f'_i(\alpha_i)$. We have $\tilde{D}_j(\omega) = \sum_{i=1}^t \lambda_i D_j \alpha_i$.

It follows from (28) that

$$f\omega - \sum f \tilde{D}_j(\omega) dX_j \in \cap m^t \Omega_{C/V} \dots (29)$$

Hence it follows from (29) that if $\tilde{D}_j(\omega) = 0$ for $1 \leq j \leq n-1$, then $f\omega \in \cap m^t \Omega_{C/V} \implies f \operatorname{im} \omega = 0 \in \hat{\Omega}_{C/V}$, i.e. $\operatorname{im} \omega \in \hat{\Omega}_{C/V}$ is a torsion element (middle vertical exact sequence in (27)).

4.6. Corollaries. Notations are as in Remark 2, 4.4.

1. $\omega \in T(\tilde{\Omega}_{B/V}) \iff \tilde{D}_i(\omega) = 0, 1 \leq i \leq n-1$, where $T(\tilde{\Omega}_{B/V})$ denotes the torsion sub-module of $\tilde{\Omega}_{B/V}$ and \tilde{D}_i denotes the corresponding extension: $\Omega_{B/V}$ or $\tilde{\Omega}_{B/V} \rightarrow Q(B)$ obtained via the extension of the derivation $\partial/\partial X_i, 1 \leq i \leq n-1$, as mentioned earlier.

This occurs due to the observation that

$$0 \rightarrow \cap m^t \hat{\Omega}_{R/V} \otimes B \rightarrow \Omega_{B/V} \rightarrow \tilde{\Omega}_{B/V} \rightarrow 0,$$

obtained by taking direct limit over C of the middle vertical column in (27), is exact.

2. we have a commutative diagram of short exact sequences:

$$\begin{array}{ccccccc}
& & & 0 & & 0 & \\
& & & \downarrow & & \downarrow & \\
& & & T(\tilde{\Omega}_{B/V}) & \equiv & T(\tilde{\Omega}_{B/V}) & \\
& & & \downarrow & & \downarrow & \\
0 & \longrightarrow & \widehat{\Omega}_{R/V} \otimes B & \longrightarrow & \tilde{\Omega}_{B/V} & \longrightarrow & \Omega_{B/R} \longrightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow & \text{.....(30)} \\
0 & \longrightarrow & \widehat{\Omega}_{R/V} \otimes B & \longrightarrow & W & \longrightarrow & N \longrightarrow 0 \\
& & & & \downarrow & & \downarrow & \\
& & & & 0 & & 0 &
\end{array}$$

where W is a B -submodule of $\bigoplus_{i=1}^{n-1} Q(B)$ and the vertical map $\tilde{D} : \tilde{\Omega}_{B/V} \rightarrow W$ is given by $\tilde{D} = (\tilde{D}_1, \dots, \tilde{D}_{n-1})$; moreover, for $i \geq 0$, $H_m^i(W) = 0$ and $H_m^i(N) \simeq \bigoplus_{i=1}^{n-1} H_m^{i+1}(B)$.

The proof follows easily from the facts: i) $\tilde{\Omega}_{B/V} = \varinjlim \widehat{\Omega}_{C/V}$ and the bottom row of (30) is exact

ii) $W \xrightarrow{p} W$, multiplication by p , is an isomorphism, and iii) $\widehat{\Omega}_{R/V} \otimes B \simeq \bigoplus_1^{n-1} B$.

Note that W is torsion-free and $W/pW = 0$. Thus $W \xrightarrow{p} W$ is an isomorphism and hence $H_m^i(W) = 0$ for $i \geq 0$.

3. $\Omega_{B/R}$ has an image N such that i) N is cohomologically Cohen-Macaulay i.e. $H_m^i(N) = 0$ for $i < n - 1$, $H_m^{n-1}(N) \neq 0$, ii) $(0 : p^t)N$ is finitely generated Cohen-Macaulay free $B/p^t B$ -module of rank $n - 1$ and of dimension $n - 1$, iii) $(0 : p^t)N$ is a summand of $(0 : p^t)\Omega_{B/R}$ and iv) for any $x \neq 0 \in R$, $(0 : x)N$ can be imbedded into a cohomologically Cohen-Macaulay flat B/xB -module. Also $\Omega_{B/V}$ can be mapped onto a B -module W such that $H_m^i(W) = 0$ for $i \geq 0$.

Proof. Since $\Omega_{B/V} \rightarrow \tilde{\Omega}_{B/V}$ is onto, the second assertion is obvious from corollary 2, (diagram 30). The last row of (30) implies that $H_m^i(N) \simeq H_m^{i+1}(\bigoplus_1^{n-1} B)$, $(0 : p^t)N \simeq \bigoplus_1^{n-1} B/p^t B$ and the last column of (30) implies that $H_m^{n-1}(\Omega_{B/R}) \rightarrow H_m^{n-1}(N) \rightarrow 0$ is exact. Due to: a) theorem 5.1 and corollary 5.10 in ([2]), b) x is a non-zero divisor on W and c) $\Omega_{R/V}$ is R -flat, our assertion follows from the last row of (30).

4.7. Remarks. 1) Corollary 2 above provides a new proof of the fact that $H_m^{n-1}(\Omega_{B/R}) \neq 0$ (Theorem 4.1, [3]). Last row of (30) implies that $H_m^{n-1}(N) \simeq \bigoplus_1^{n-1} H_m^n(B)$ and last column of (30) implies that $H_m^{n-1}(\Omega_{B/R}) \rightarrow H_m^{n-1}(N)$ is onto. Hence, $H_m^{n-1}(\Omega_{B/R}) \neq 0$ ([1], [3]).

2) Using theorem 5.1 in ([2]) it follows from proposition 4.2 in ([3]) that $H_m^i(\Omega_{B/V}) \simeq H_m^i(\Omega_{B/R})$ for $i \leq n-2$, and from corollary 1 (3.4) that a) $H_m^i(\Omega_{B/A}) \simeq H_m^i(\Omega_{B/AR})$, for $i \leq n-2$; b) $H_m^i(\Omega_{B/V}) \simeq H_m^i(\Omega_{B/A})$, for $i \leq n-3$ and injectivity of maps at the next level. The crucial question is whether $H_m^{n-1}(\Omega_{A/V} \otimes B) \simeq H_m^n(B) \rightarrow H_m^{n-1}(\Omega_{B/V})$ is non-null.

3) B. Bhatt kindly shared with me his approach for providing another new proof of the fact that $H_m^{n-1}(\Omega_{B/R}) \neq 0$ (theorem 4.1, [3]). He used Cohen-Macaulayness of the m -adic completion \hat{B} along with facts from derived categories and perfectoids to accomplish his proof.

4) In equicharacteristic 0 same proof works for corresponding statements for theorem 4.5, corollary 1 and corollary 2 in 4.6 without the assertion on local cohomology modules. In positive characteristic the same arguments work for corresponding statement for theorem 4.5 with the assumption that $Q(C)$ is separably algebraic over $Q(R)$.

5) Let T denote the torsion sub-module of $\Omega_{B/V}$, let $U = \text{Coker}(T \rightarrow \Omega_{B/V})$ and $L = \text{Coker}(T \rightarrow \Omega_{B/R})$. Then, as in corollary 2 and corollary 3 in 4.6 it follows that $H_m^i(U) = 0$ for $i \geq 0$; $H_m^i(L) = 0$ for $i \neq n-1$ and $H_m^{n-1}(L) \simeq H_m^{n-1}(\bigoplus_1^{n-1} B)$ and for $t > 0$, $(0 : p^t)L \simeq \bigoplus_1^{n-1} B/p^t B$. The above isomorphisms are obtained from the exact sequence:

$$0 \rightarrow \Omega_{R/V} \otimes B \rightarrow U \rightarrow L \rightarrow 0.$$

Thus L shares the same properties with N in corollary 3, 4.6 above. Moreover, with N as in corollary 2 above, there exists an onto map $\eta : L \rightarrow N$ such that $H_m^i(\ker \eta) = 0$ for $i \geq 0$.

REFERENCES

- [1] Y. André, La conjecture de facteur direct, Publ. Math. IHES 127, 71-93 (2018).
- [2] B. Bhatt, Cohen-Macaulayness of absolute integral closures, arXiv.2008.08070 [math.AG].
- [3] S. P. Dutta, Local Cohomology of Module of Differentials of Integral Extensions, J. of Algebra 582(2021), 136-156.
- [4] J-M Fontaine, Formes Différentielles et modules de Tate des variétés abéliennes sur les corps locaux, Inventiones math. 65, 379-409 (1982).
- [5] A. Grothendieck & J. Dieudonné, EGA IV, Publ. Math. IHES 20 (1964).

- [6] R. Hartshorne, Local Cohomology, Lecture Notes in Math, 41 (1967), Springer-Verlag. Berlin. Heidelberg. NewYork.
- [7] M. Hochster, Canonical elements in local cohomology modules and the direct summand conjecture, J. of Algebra, vol. 84, No. 3, October (1983), 503-553.
- [8] M. Nagata, Local Rings, R. E. K. Publ. Company, Huntington, New York, 1975.
- [9] J. P. Serre, Local Fields, Grad. Text in math. Springer-Verlag, New York Heidelberg Berlin.
- [10] S. Suzuki, On torsion of the module of differentials of a locality which is a complete intersection, J. Math. Kyoto. Univ., 4-3(1965), 471-475.
- [11] R. G. Swan, Nèron-Popescu desingularization, Proc. Alg. and Geom (1995), 135-192.
- [12] T. Szamuely & Z. Zabradi, The p-adic Hodge decomposition according to Beilinson, arXiv: 1606.01921 v3 [mathNT], 17Jan. 2017.
- [13] O. Zariski & P. Samuel, Commutative Algebra, vol i, Springer Verlag, New York Heidelberg Berlin.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN,
1409 WEST GREEN STREET, URBANA, ILLINOIS 61801