

The Farahat-Higman Algebra of Centralizers of Symmetric Group Algebras

Samuel Creedon

Abstract

We define a generalisation of the Farahat-Higman algebra introduced in [FH59] which replaces the role of the center of the group algebra of the symmetric groups with certain centralizer algebras instead. We prove various structural properties of this new algebra and provide an isomorphism which connects it to the degenerate affine Hecke algebra and the algebra of symmetric functions.

1 Introduction

Let \mathfrak{S}_n denote the symmetric group on n letters. The center $Z(\mathbb{Z}\mathfrak{S}_n)$ has a basis consisting of conjugacy class sums. In [FH59, Theorem 2.2] the structure constants associated with this basis were shown to be polynomial in n . This allowed H. Farahat and G. Higman to construct an algebra, denoted here by FH_0 , by in some sense replacing the structure constants with their corresponding polynomials. In particular this algebra is given over the subring R of integer-valued polynomials in $\mathbb{Q}[t]$, and via “evaluation at n ” one has surjective ring homomorphisms $\text{pr}_{n,0} : \text{FH}_0 \rightarrow Z(\mathbb{Z}\mathfrak{S}_n)$ for each $n \geq 0$. The algebra FH_0 is determined by the centers $Z(\mathbb{Z}\mathfrak{S}_n)$ in the sense that a relation holds in FH_0 if and only if its image under each $\text{pr}_{n,0}$ holds. The motivation for the construction of such an algebra was to present an alternative proof of Nakayama’s Conjecture regarding the p -blocks of symmetric groups. Also, the main result of [FH59] was in establishing a set of generators for FH_0 , whose non-zero images generate the centers $Z(\mathbb{Z}\mathfrak{S}_n)$. It was shown in [Jucys74] that these images correspond to the elementary symmetric polynomials in the Jucys-Murphy elements, demonstrating that $Z(\mathbb{Z}\mathfrak{S}_n)$ is the subalgebra of symmetric polynomials in such elements.

In [Mac95, p.131-132] a certain associated graded algebra \mathbf{G} of FH_0 is considered. This algebra may be regarded as a \mathbb{Z} -algebra as the non-constant polynomials appearing in the product of elements of FH_0 get filtered out. The algebra \mathbf{G} was shown to be isomorphic to the algebra of symmetric functions, and it has been studied in relation to enumeration problems for permutation factorisation, see for example [GJ94]. The algebras FH_0 and \mathbf{G} have seen counterparts in the setting of the spin symmetric group algebras in [TW09], of wreath products of the symmetric groups with any finite group in [W03] and [Ryba21], and of the general linear group over finite fields in [WW19].

In this paper we construct generalisations of FH_0 by considering certain centralizer algebras of the symmetric group in place of the centers $Z(\mathbb{Z}\mathfrak{S}_n)$. For $n \geq m$ let $\text{Stab}_n(m)$ denote the subgroup of \mathfrak{S}_n of all permutations which fix each element $1, 2, \dots, m$. Then we consider the centralizer algebra $Z_{n,m}$ consisting of all elements of $\mathbb{Z}\mathfrak{S}_n$ which commute with $\text{Stab}_n(m)$. When $m = 0$ then $Z_{n,0} = Z(\mathbb{Z}\mathfrak{S}_n)$. The algebra $Z_{n,m}$ has its own counterparts to class sums which provide a basis, and we prove in *Theorem 3.0.3* that the associated structure constants are polynomial in n . These polynomials again belong to the subring R of integer-valued polynomials in $\mathbb{Q}[t]$. From that, we can define a new R -algebra FH_m which we refer to as the Farahat-Higman algebra associated to $Z_{n,m}$. There exist analogous surjective ring homomorphisms $\text{pr}_{n,m} : \text{FH}_m \rightarrow Z_{n,m}$ for any $n \geq m$, and the algebras FH_m are determined by the centralizer algebras $Z_{n,m}$ in a manner comparable to above. To the best of the author’s knowledge, this is the first time the techniques of Farahat and Higman have been employed on an algebra which is not the center of a group algebra, and the algebra FH_m is in general not commutative.

The remainder of the paper proves various structural properties of FH_m . We show that the symmetric group algebra $R\mathfrak{S}_m$ and the Farahat-Higman algebra FH_0 are subalgebras of FH_m . The indexing set for the basis elements of FH_m are called m -marked cycle shapes, and have only made brief appearances elsewhere (see for example [K05, p. 8]). We define a monoid structure on the set of m -marked cycle shapes, and use

it to prove a leading term result for the product of basis elements in FH_m . This result, along with other known results of FH_0 , allow us to prove the main result *Theorem 9.0.5* of the paper where we establish an isomorphism between the R -algebras FH_m and $R \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$ where \mathcal{H}_m is the degenerate affine Hecke algebra (viewed over the integers) and Sym is the \mathbb{Z} -algebra of symmetric functions. This isomorphism gives us a description of the center of FH_m and a new basis involving elements which mimic the Jucys-Murphy elements. The \mathbb{C} -algebra $\mathbb{C} \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$ has appeared in the context of the Heisenberg category of [Kho14], where it arises as an endomorphism algebra of certain objects. Also a closely related algebra is studied in [MO01] when investigating a centralizer construction of the degenerate affine Hecke algebra. In this sense the algebra FH_m gives an organic means of realising the degenerate affine Hecke algebra from a Farahat-Higman perspective. A conscious effort has been made to keep all the results of FH_m integral (i.e. over R as opposed to its field of fractions $\mathbb{Q}(t)$). The hope is that this will allow FH_m to remain open as a tool for investigating the modular representations of $Z_{n,m}$ in future work.

The structure of the paper is as follows: *Section 2* proves some technical results regarding the cardinalities of certain orbits. These results are used in *Section 3* to prove the polynomial property of the structure constants in the centralizer algebras $Z_{n,m}$. We also set up some notation including that of an m -marked cycle shape and its degree. In *Section 4* we define the R -algebras FH_m and prove that $\text{R}\mathfrak{S}_m$ and FH_0 are subalgebras. In *Section 5* we describe a monoid structure on the set of m -marked cycle shapes and describe a concrete criteria for a permutation to belong to the orbit associated to a given m -marked cycle shape. In *Section 6* we use this product to describe a leading term of a product of basis elements in FH_m . *Section 7* simply recalls the algebra of symmetric functions and *Section 8* recalls the Jucys-Murphy elements of $\mathbb{Z}\mathfrak{S}_n$ and their relationship to the Farahat-Higman algebra FH_0 . We then end with *Section 9* which connects FH_m with the degenerate affine Hecke algebra and the algebra of symmetric functions.

2 Polynomial Cardinality of m -Classes

Given a finite set $A \subset \mathbb{N}$ we write $\mathfrak{S}(A)$ to denote the group of permutations of A . For any $n \geq 0$ we write $[n] := \{1, \dots, n\}$ and $\mathfrak{S}_n := \mathfrak{S}([n])$ for the symmetric group on n letters, with convention that $[0] = \emptyset$ and \mathfrak{S}_0 the trivial group. We set $\mathfrak{S}_{\mathbb{N}} := \cup_{n \geq 1} \mathfrak{S}_n$, the group of permutations of \mathbb{N} with finite support. We denote the support of any $\pi \in \mathfrak{S}_{\mathbb{N}}$ by $\text{Sup}(\pi) = \{i \in \mathbb{N} \mid \pi(i) \neq i\}$ and set $\|\pi\| := |\text{Sup}(\pi)|$. For any $r \geq 1$ consider the r -fold direct product $\mathfrak{S}_{\mathbb{N}}^{\times r}$. For $\boldsymbol{\pi} = (\pi_i)_{i=1}^r \in \mathfrak{S}_{\mathbb{N}}^{\times r}$ let $\text{Sup}(\boldsymbol{\pi}) := \text{Sup}(\pi_1) \cup \dots \cup \text{Sup}(\pi_r)$ and set $\|\boldsymbol{\pi}\| := |\text{Sup}(\boldsymbol{\pi})|$. For any $m \geq 0$ let $\text{Stab}(m)$ denote the subgroup of $\mathfrak{S}_{\mathbb{N}}$ consisting of the permutations which fix each element of $[m]$, in particular $\text{Stab}(0) = \mathfrak{S}_{\mathbb{N}}$. The group $\text{Stab}(m)$ acts on the r -fold direct product $\mathfrak{S}_{\mathbb{N}}^{\times r}$ by component-wise conjugation. We refer to the respected orbits as the m -classes of $\mathfrak{S}_{\mathbb{N}}^{\times r}$. We denote the m -class of $\mathfrak{S}_{\mathbb{N}}^{\times r}$ containing $\boldsymbol{\pi} = (\pi_i)_{i=1}^r \in \mathfrak{S}_{\mathbb{N}}^{\times r}$ by

$$\text{CL}_{\mathbb{N},m}(\boldsymbol{\pi}) := \{(\sigma_i)_{i=1}^r \in \mathfrak{S}_{\mathbb{N}}^{\times r} \mid \sigma_i = \tau \pi_i \tau^{-1} \text{ for all } i \in [r] \text{ and some } \tau \in \text{Stab}(m)\}.$$

Example 2.0.1. Let $m = r = 2$ and consider $\boldsymbol{\pi} = ((1, 3)(2, 4), (3, 4, 5)) \in \mathfrak{S}_{\mathbb{N}}^{\times 2}$. Then the m -class containing $\boldsymbol{\pi}$ is given by $\text{CL}_{\mathbb{N},2}(\boldsymbol{\pi}) = \{((1, a)(2, b), (a, b, c)) \mid (a, b, c) \in (\mathbb{N} \setminus [2])^{13}\}$ where $(\mathbb{N} \setminus [2])^{13}$ is the subset of the 3-fold direct product of $\mathbb{N} \setminus [2]$ consisting of all tuples with pairwise distinct entries.

Given any $m \geq 0$ and $\boldsymbol{\pi} = (\pi_i)_{i=1}^r \in \mathfrak{S}_{\mathbb{N}}^{\times r}$ define

$$\text{Sup}_m(\boldsymbol{\pi}) := \text{Sup}(\boldsymbol{\pi}) \cap [m], \text{ and } \text{Sup}^m(\boldsymbol{\pi}) := \text{Sup}(\boldsymbol{\pi}) \setminus [m],$$

and write $\|\boldsymbol{\pi}\|_m$ and $\|\boldsymbol{\pi}\|^m$ for the cardinalities of $\text{Sup}_m(\boldsymbol{\pi})$ and $\text{Sup}^m(\boldsymbol{\pi})$ respectively. Let \mathbf{C} be an m -class of $\mathfrak{S}_{\mathbb{N}}^{\times r}$ and let $\boldsymbol{\pi} = (\pi_i)_{i=1}^r, \boldsymbol{\sigma} = (\sigma_i)_{i=1}^r \in \mathbf{C}$. Hence there exists some $\tau \in \text{Stab}(m)$ such that $\sigma_i = \tau \pi_i \tau^{-1}$ for each $i \in [r]$. Conjugating π_i by τ permutes the entries within the cycles of π_i according to τ , thus for each $i \in [r]$ the permutations π_i and σ_i must have the same cycle structure, and the relative positions of the elements of $[m]$ among their cycles must agree. Hence $\|\boldsymbol{\pi}\| = \|\boldsymbol{\sigma}\|$, $\|\boldsymbol{\pi}\|_m = \|\boldsymbol{\sigma}\|_m$, and $\|\boldsymbol{\pi}\|^m = \|\boldsymbol{\sigma}\|^m$, and so it makes sense to define $\|\mathbf{C}\| := \|\boldsymbol{\pi}\|$, $\|\mathbf{C}\|_m := \|\boldsymbol{\pi}\|_m$, and $\|\mathbf{C}\|^m := \|\boldsymbol{\pi}\|^m$ for any $\boldsymbol{\pi} \in \mathbf{C}$.

Let $n \geq m$, then we set $\text{Stab}_n(m) := \text{Stab}(m) \cap \mathfrak{S}_n$. Similar to the above situation, this group acts on the r -fold direct product $\mathfrak{S}_n^{\times r}$ by component-wise conjugation. We refer to the respected orbits as the m -classes of $\mathfrak{S}_n^{\times r}$. For any $\boldsymbol{\pi} = (\pi_i)_{i=1}^r \in \mathfrak{S}_n^{\times r}$ we denote the m -class of $\mathfrak{S}_n^{\times r}$ containing $\boldsymbol{\pi}$ by

$$\text{CL}_{n,m}(\boldsymbol{\pi}) := \{(\sigma_i)_{i=1}^r \in \mathfrak{S}_n^{\times r} \mid \sigma_i = \tau \pi_i \tau^{-1} \text{ for all } i \in [r] \text{ and some } \tau \in \text{Stab}_n(m)\}.$$

Again we can set $\|C\| := \|\pi\|$, $\|C\|_X := \|\pi\|_X$, and $\|C\|^X := \|\pi\|^X$ for any m -class C of $\mathfrak{S}_n^{\times r}$ and any $\pi \in C$. Given any m -class C of $\mathfrak{S}_n^{\times r}$ we write $C_n := C \cap \mathfrak{S}_n^{\times r}$.

Example 2.0.2. Continuing from *Example 2.0.1* let $C = \text{CL}_{\mathbb{N},2}(\pi)$. For any $n \geq 2$ one can deduce that

$$C_n = \begin{cases} \{((1, a)(2, b), (a, b, c)) \mid (a, b, c) \in ([n] \setminus [2])^{13}\}, & n \geq 5 \\ \emptyset, & 2 \leq n \leq 4 \end{cases}$$

When $n \geq 5$ the set C_n is an m -class of $\mathfrak{S}_n^{\times 2}$.

As the above example suggests, when C is an m -class of $\mathfrak{S}_n^{\times r}$ and $n \geq m$, then the set C_n is either empty or an m -class of $\mathfrak{S}_n^{\times r}$. The following proposition proves this and gives a criteria for when C_n is empty.

Proposition 2.0.3. *Let C be an m -class of $\mathfrak{S}_n^{\times r}$ and $n \geq m$. Then C_n is non-empty if and only if*

$$n \geq \|C\|^m + m. \quad (1)$$

In this case C_n is an m -class of $\mathfrak{S}_n^{\times r}$ and the m -classes of $\mathfrak{S}_n^{\times r}$ appear uniquely in this manner.

Proof. Fix $\pi = (\pi_i)_{i=1}^r \in C$. All elements of C are of the form $(\tau\pi_i\tau^{-1})_{i=1}^r$ for $\tau \in \text{Stab}(m)$. The set $\text{Sup}^m(\tau\pi\tau^{-1})$ is all elements of $\mathbb{N} \setminus [m]$ for which at least one $\tau\pi_i\tau^{-1}$ acts non-trivially. Thus $\tau\pi\tau^{-1} \in C_n$ if and only if $\text{Sup}^m(\tau\pi\tau^{-1}) \subset [n] \setminus [m]$. For this to be the case we must have

$$|\text{Sup}^m(\tau\pi\tau^{-1})| = \|C\|^m \leq |[n] \setminus [m]| = n - m.$$

Rearranging gives *Equation (1)*. So $C_n = \emptyset$ whenever $n < \|C\|^m + m$. Now assume *Equation (1)* holds. We have $\text{Sup}^m(\tau\pi\tau^{-1}) = \tau(\text{Sup}^m(\pi))$, the image of $\text{Sup}^m(\pi)$ under τ . Let $t : \text{Sup}^m(\pi) \rightarrow [n] \setminus [m]$ be an injective map, which exists since *Equation (1)* holds. Then fix a permutation $\tau_t \in \text{Stab}(m)$ such that $\tau(i) = t(i)$ for all $i \in \text{Sup}^m(\pi)$. Hence $\text{Sup}^m(\tau_t\pi\tau_t^{-1}) \subset [n] \setminus [m]$ and so C_n is non-empty as it contains $\sigma := \tau_t\pi\tau_t^{-1}$.

We now want to prove that C_n is an m -class of $\mathfrak{S}_n^{\times r}$. Since $\sigma := \tau_t\pi\tau_t^{-1} \in C_n$ and $C_n = C \cap \mathfrak{S}_n^{\times r}$, any element of C_n is of the form $\tau\sigma\tau^{-1}$ for $\tau \in \text{Stab}(m)$ such that $\text{Sup}^m(\tau\sigma\tau^{-1}) \subset [n] \setminus [m]$. If $\tau \in \text{Stab}_n(m)$ then clearly $\text{Sup}^m(\tau\sigma\tau^{-1}) \subset [n] \setminus [m]$, thus we have $\text{CL}_{n,m}(\sigma) \subset C_n$. Hence the equality $\text{CL}_{n,m}(\sigma) = C_n$ occurs if one can show that whenever $\tau\sigma\tau^{-1} \in C_n$ for some $\tau \in \text{Stab}(m)$ there exists a $\tau' \in \text{Stab}_n(m)$ such that $\tau\sigma\tau^{-1} = \tau'\sigma(\tau')^{-1}$. Given such a τ we must have that $\text{Sup}^m(\tau\sigma\tau^{-1}) = \tau(\text{Sup}^m(\sigma)) \subset [n] \setminus [m]$. Since we also have that $\text{Sup}^m(\sigma) \subset [n] \setminus [m]$, let τ' be any permutation of $[n] \setminus [m]$ with the property that $\tau'(i) := \tau(i)$ for all $i \in \text{Sup}^m(\sigma)$, then it is clear that $\tau\sigma\tau^{-1} = \tau'\sigma(\tau')^{-1}$ and so $\text{CL}_{n,m}(\sigma) = C_n$. Hence for any $\sigma \in \mathfrak{S}_n^{\times r}$ we have $(\text{CL}_{\mathbb{N},m}(\sigma))_n = \text{CL}_{n,m}(\sigma)$. Since orbits intersect trivially all such m -classes of $\mathfrak{S}_n^{\times r}$ appear as the intersection of a unique m -class of $\mathfrak{S}_n^{\times r}$ with $\mathfrak{S}_n^{\times r}$. \square

We now end this section by presenting a result describing the cardinality of the set C_n when C is an m -class of $\mathfrak{S}_n^{\times r}$ and $n \geq m$. Notably we show that the cardinality of such a set is polynomial in n .

Proposition 2.0.4. *Let C be an m -class of $\mathfrak{S}_n^{\times r}$. For $n \geq m$,*

$$|C_n| = \frac{1}{b(C)}(n-m)(n-m-1)\cdots(n-m-\|C\|^m+1)$$

where $b(C) \in \mathbb{N}$ is a constant depending only on the class C and not on n .

Proof. Assume $n \geq \|C\|^m + m$, so C_n is an m -class of $\mathfrak{S}_n^{\times r}$ by *Proposition 2.0.3*. Pick $\pi = (\pi_i)_{i=1}^r \in C_n$, then $C_n = \text{CL}_{n,m}(\pi)$ is the orbit of π in $\mathfrak{S}_n^{\times r}$ under the action of $\text{Stab}_n(m)$ by component-wise conjugation. Thus by the *Orbit-Stabilizer Theorem* we have

$$|C_n| = \frac{|\text{Stab}_n(m)|}{|\text{Stab}_{\text{Stab}_n(m)}(\pi)|}$$

where $\text{Stab}_{\text{Stab}_n(m)}(\pi) = \{\tau \in \text{Stab}_n(m) \mid \tau\pi\tau^{-1} = \pi\}$ is the subgroup of $\text{Stab}_n(m)$ whose elements fix π under component-wise conjugation. Consider $\text{Stab}_n([m] \cup \text{Sup}^m(\pi))$, the subgroup of \mathfrak{S}_n consisting of all

permutations which act trivially on $[m] \cup \text{Sup}^m(\pi)$. Naturally $\text{Stab}_n([m] \cup \text{Sup}^m(\pi)) \subset \text{Stab}_{\text{Stab}_n(m)}(\pi)$. Now let $\mathfrak{S}(\text{Sup}^m(\pi))$ denote the subgroup of \mathfrak{S}_n consisting of the permutations of $\text{Sup}^m(\pi)$. Then we have that $\mathfrak{S}(\text{Sup}^m(\pi)) \subset \text{Stab}_n(m)$, and hence $\text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi) = \{\tau \in \mathfrak{S}(\text{Sup}^m(\pi)) \mid \tau\pi\tau^{-1} = \pi\}$ is also a subgroup of $\text{Stab}_{\text{Stab}_n(m)}(\pi)$.

Claim: We have a group isomorphism

$$\text{Stab}_{\text{Stab}_n(m)}(\pi) \cong \text{Stab}_n([m] \cup \text{Sup}^m(\pi)) \times \text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi).$$

Note that the two subgroups $\text{Stab}_n([m] \cup \text{Sup}^m(\pi))$ and $\text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi)$ commute and have trivial intersection. So to prove this claim we only need to show that any permutation $\tau \in \text{Stab}_{\text{Stab}_n(m)}(\pi)$ can be expressed as $\tau = \tau_1\tau_2$ for some $\tau_1 \in \text{Stab}_n([m] \cup \text{Sup}^m(\pi))$ and $\tau_2 \in \text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi)$. Let $\tau \in \text{Stab}_{\text{Stab}_n(m)}(\pi)$, then by definition $\tau\pi_i\tau^{-1} = \pi_i$ for each $i \in [r]$. We seek to show that the sets $[n] \setminus \text{Sup}(\pi)$ and $\text{Sup}(\pi)$ are invariant under the action of τ . Suppose for contradiction this is not the case, hence there exists an $a \in [n] \setminus \text{Sup}(\pi)$ such that $\tau(a) = b \in \text{Sup}(\pi)$. Then for each $i \in [r]$,

$$\pi_i(b) = (\tau\pi_i\tau^{-1})(b) = (\tau\pi_i)(a) = \tau(a) = b.$$

Thus π_i fixes b for each $i \in [r]$, but this gives the desired contradiction since $b \in \text{Sup}(\pi)$. Thus the sets $[n] \setminus \text{Sup}(\pi)$ and $\text{Sup}(\pi)$ are invariant under the action of τ , which implies a decomposition $\tau = \tau_1\tau_2$ where τ_1 is a permutation of $[n] \setminus \text{Sup}(\pi)$ and τ_2 is a permutation of $\text{Sup}(\pi)$. Naturally τ_1 and τ_2 commute, and since τ fixes $[m]$, both τ_1 and τ_2 must also fix $[m]$. As such τ_1 is an element of $\text{Stab}_n([m] \cup \text{Sup}^m(\pi))$ as desired, and $\tau_2 \in \mathfrak{S}(\text{Sup}^m(\pi))$. Lastly note that for each $i \in [r]$,

$$\pi_i = \tau\pi_i\tau^{-1} = \tau_2\tau_1\pi_i\tau_1^{-1}\tau_2^{-1} = \tau_2\pi_i\tau_2^{-1}$$

since τ_1 commutes with both τ_2 and π_i . Thus $\tau_2\pi_i\tau_2^{-1} = \pi_i$ for each $i \in [r]$, implying that τ_2 belongs to $\text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi)$. Hence the claim holds.

Returning to the cardinality of C_n , recall that $|\text{Sup}^m(\pi)| = \|\mathbf{C}\|^m$. Also observe that the size of the group $\mathfrak{S}(\text{Sup}^m(\pi))$ depends only on the class \mathbf{C} , in particular it is independent of n . This implies the same for the cardinality of $\text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi)$, and so we write $b(\mathbf{C}) := |\text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi)|$. Hence we have that

$$\begin{aligned} |C_n| &= \frac{|\text{Stab}_n(m)|}{|\text{Stab}_n([m] \cup \text{Sup}^m(\pi))| |\text{Stab}_{\mathfrak{S}(\text{Sup}^m(\pi))}(\pi)|} \\ &= \frac{(n-m)!}{(n-m - \|\mathbf{C}\|^m)! b(\mathbf{C})} \\ &= \frac{1}{b(\mathbf{C})} (n-m)(n-m-1) \cdots (n-m - \|\mathbf{C}\|^m + 1). \end{aligned}$$

We assumed $n \geq \|\mathbf{C}\|^m + m$ so that C_n is an m -class of $\mathfrak{S}_n^{\times r}$. By *Proposition 2.0.3* the set C_n is empty when $m \leq n < \|\mathbf{C}\|^m + m$, and such values of n are precisely those which give zero in the above formula for $|C_n|$. Hence this formula holds for all $n \geq m$, completing the proof. \square

3 Centralizer Algebras and Polynomial Structure Constants

Let $m \geq 0$ and \mathbf{C} be an m -class of $\mathfrak{S}_{\mathbb{N}}$ and $\pi \in \mathbf{C}$. As mentioned before \mathbf{C} is completely determined by the cycle structure of π and the relative positions of the elements of $[m]$ among the cycles. If one was to take π and replace each of the elements in $\mathbb{N} \setminus [m]$ among the non-trivial cycles with a formal symbol, say $*$, then the resulting object would retain the defining characteristics of \mathbf{C} , and so could represent it.

Definition 3.0.1. For a finite set A , we call $(a_i)_{i=1}^l \in A^{\times l}$, for some $l \in \mathbb{N}$, a *cycle* if we only care about the order of the entries up to cyclic shifts, and say it has length l . For $*$ a formal symbol, we define an *m -marked cycle shape* to be a finite collection of cycles with entries in $[m] \cup \{*\}$ with the following properties:

- (1) The multiset of entries among the cycles equals $[m] \cup \{*\}^n$ for some $n \in \mathbb{Z}_{\geq 0}$, in particular each element of $[m]$ appears precisely once.

(2) Cycles containing only $*$ must be of length at least two.

We write an m -marked cycle shape as a formal product of its cycles by juxtaposition, where the order of the cycles is immaterial. We let $\Lambda(m)$ denote the set of all such m -marked cycle shapes.

Example 3.0.2. An example of a 6-marked cycle shape is

$$\lambda = (2, 6)(5)(1, *, *, 4, *) (3, *, *) (*, *) (*, *, *) \in \Lambda(6).$$

The multiset of entries among the cycles of λ is $[6] \cup \{*\}^{10}$. Cyclicly shifting the entries of any of the cycles or rearranging the cycles in any order will result in an alternative expression of λ .

The 0-marked cycle shapes only contain the symbol $*$, and we will refer to these as *cycle shapes*. Clearly one may identify the set $\Lambda(0)$ with the set of proper partitions (i.e. partitions with no parts equal to 1). We consider the empty set \emptyset an element of $\Lambda(0)$ consisting of no cycles. Also the subset of $\Lambda(m)$ consisting of the m -marked cycle shapes which contain no symbols $*$ may be identified with \mathfrak{S}_m , and so we write $\mathfrak{S}_m \subset \Lambda(m)$.

As discussed above the set of m -marked cycle shapes $\Lambda(m)$ provides a natural indexing set for the collection of m -classes of $\mathfrak{S}_{\mathbb{N}}$: For $\lambda \in \Lambda(m)$ let $\text{CL}_{\mathbb{N}, m}(\lambda)$ denote the corresponding m -class. The permutations of $\text{CL}_{\mathbb{N}, m}(\lambda)$ are those obtained from λ by replacing the symbols $*$ with distinct elements from $\mathbb{N} \setminus [m]$. For example, letting λ be the 6-marked cycle shape displayed in *Example 3.0.2*, we have that

$$\text{CL}_{\mathbb{N}, 6}(\lambda) = \left\{ (2, 6)(1, a_1, a_2, 4, a_3)(3, a_4, a_5)(a_6, a_7)(a_8, a_9, a_{10}) \mid (a_i)_{i=1}^{10} \in (\mathbb{N} \setminus [6])^{10} \right\}.$$

Recall the quantities $\|\text{C}\|$, $\|\text{C}\|^m$, and $\|\text{C}\|_m$ for any m -class C of $\mathfrak{S}_{\mathbb{N}}$. Then for $\lambda \in \Lambda(m)$ we write

$$\|\lambda\| := \|\text{CL}_{\mathbb{N}, m}(\lambda)\|, \quad \|\lambda\|^m := \|\text{CL}_{\mathbb{N}, m}(\lambda)\|^m, \quad \text{and} \quad \|\lambda\|_m := \|\text{CL}_{\mathbb{N}, m}(\lambda)\|_m.$$

The quantity $\|\lambda\|$ is the number of entries among the cycles of length at least two, $\|\lambda\|^m$ is the number of $*$ symbols appearing among the cycles, and $\|\lambda\|_m$ is the number of elements from $[m]$ which appear in cycles of length at least two. We define a map $\text{deg}_m : \Lambda(m) \rightarrow \mathbb{Z}_{\geq m}$ by setting

$$\text{deg}_m(\lambda) := \|\lambda\|^m + m.$$

We refer to $\text{deg}_m(\lambda)$ as the *degree* of λ . We have defined the degree map with the inequality of *Proposition 2.0.3* in mind, in particular for any $n \geq m$ we have that the set $\Lambda_{\leq n}(m) := \{\lambda \in \Lambda(m) \mid \text{deg}_m(\lambda) \leq n\}$ gives us an indexing set for the m -classes of \mathfrak{S}_n . Given $\lambda \in \Lambda_{\leq n}(m)$ we denote the corresponding m -class by $\text{CL}_{n, m}(\lambda) = (\text{CL}_{\mathbb{N}, m}(\lambda))_n$, which consists of all permutations of \mathfrak{S}_n one can obtain from λ by replacing the symbols $*$ with distinct elements from $[n] \setminus [m]$.

Consider the \mathbb{Z} -algebra $\mathbb{Z}\mathfrak{S}_n$. For any $n \geq m$ we define the *m -centralizer algebra* as

$$Z_{n, m} := \{z \in \mathbb{Z}\mathfrak{S}_n \mid \tau z = z \tau \text{ for all } \tau \in \text{Stab}_n(m)\}.$$

When $m = 0$ then $\text{Stab}_n(0) = \mathfrak{S}_n$ and $Z_{n, 0} = Z(\mathbb{Z}\mathfrak{S}_n)$. For any $\lambda \in \Lambda(m)$ we define the m -class sum by

$$K_n(\lambda) := \sum_{\pi \in \text{CL}_{n, m}(\lambda)} \pi.$$

Note by *Proposition 2.0.3* we have that $K_n(\lambda) \neq 0$ if and only if $\text{deg}_m(\lambda) \leq n$. Also one can deduce that the set of m -class sums $\{K_n(\lambda) \mid \lambda \in \Lambda_{\leq n}(m)\}$ provides a \mathbb{Z} -basis for the centralizer algebra $Z_{n, m}$. As such the product of any two such elements must decomposed into a linear combination of the same, that is for any $\lambda, \mu \in \Lambda_{\leq n}(m)$ we have that

$$K_n(\lambda)K_n(\mu) = \sum_{\nu \in \Lambda_{\leq n}(m)} a_{\lambda, \mu}^{\nu}(n)K_n(\nu)$$

with structure constants $a_{\lambda, \mu}^{\nu}(n) \in \mathbb{Z}_{\geq 0}$. We now show these structure constants are polynomial in n . Let t be a formal variable and R the subring of $\mathbb{Q}[t]$ of all polynomials $f(t)$ such that $f(n) \in \mathbb{Z}$ whenever $n \in \mathbb{Z}$.

Theorem 3.0.3. For each $\lambda, \mu, \nu \in \Lambda(m)$ there exists a unique polynomial $f_{\lambda, \mu}^\nu(t) \in R$ such that

$$K_n(\lambda)K_n(\mu) = \sum_{\nu \in \Lambda_{\leq n}(m)} f_{\lambda, \mu}^\nu(n)K_n(\nu)$$

in $Z_{n,m}$ for any $n \geq m$. We refer to the polynomials $f_{\lambda, \mu}^\nu(t)$ as the structure polynomials.

Proof. Fix $\lambda, \mu, \nu \in \Lambda(m)$. Consider the set of pairs

$$A = \{(\pi_1, \pi_2) \in \text{CL}_{\mathbb{N}, m}(\lambda) \times \text{CL}_{\mathbb{N}, m}(\mu) \mid \pi_1 \pi_2 \in \text{CL}_{\mathbb{N}, m}(\nu)\} \subset \mathfrak{S}_{\mathbb{N}} \times \mathfrak{S}_{\mathbb{N}}.$$

When $A = \emptyset$ we set $f_{\lambda, \mu}^\nu(t) := 0$. Assume $A \neq \emptyset$ and let $(\pi_1, \pi_2) \in A$. For any $\tau \in \text{Stab}(m)$ we have that $(\tau \pi_1 \tau^{-1})(\tau \pi_2 \tau^{-1}) = \tau(\pi_1 \pi_2) \tau^{-1}$ which belongs to $\text{CL}_{\mathbb{N}, m}(\nu)$ since $\pi_1 \pi_2$ does. So $(\tau \pi_1 \tau^{-1}, \tau \pi_2 \tau^{-1})$ belongs to A for any $\tau \in \text{Stab}(m)$. Thus for some indexing set I , the set A is the union of m -class $\mathbf{C}^{(i)}$ of $\mathfrak{S}_{\mathbb{N}} \times \mathfrak{S}_{\mathbb{N}}$ for each $i \in I$. For any $(\pi_1^{(i)}, \pi_2^{(i)}) \in \mathbf{C}^{(i)}$,

$$\|\mathbf{C}^{(i)}\|^m = |\text{Sup}^m(\pi_1^{(i)}) \cup \text{Sup}^m(\pi_2^{(i)})| \leq |\text{Sup}^m(\pi_1^{(i)})| + |\text{Sup}^m(\pi_2^{(i)})| = \|\lambda\|^m + \|\mu\|^m.$$

Thus by *Proposition 2.0.3* we have for any $n \geq \|\lambda\|^m + \|\mu\|^m + m$ that $\mathbf{C}_n^{(i)}$ is an m -class of $\mathfrak{S}_n \times \mathfrak{S}_n$ for each $i \in I$. This implies that I is finite. Also by *Proposition 2.0.4* we have for any $n \geq m$ that

$$|A \cap (\mathfrak{S}_n \times \mathfrak{S}_n)| = \sum_{i \in I} \frac{1}{b(\mathbf{C}^{(i)})} (n-m)(n-m-1) \cdots (n-m - \|\mathbf{C}^{(i)}\|^m + 1),$$

where $b(\mathbf{C}^{(i)})$ are constants independent of n . By definition of A , the multiplicity of $K_n(\nu)$ in the product $K_n(\lambda)K_n(\mu)$ is $|A \cap (\mathfrak{S}_n \times \mathfrak{S}_n)|$ divided by $|\text{CL}_{n,m}(\nu)|$. Again by *Proposition 2.0.4* we have that

$$|\text{CL}_{n,m}(\nu)| = \frac{1}{b(\nu)} (n-m)(n-m-1) \cdots (n-m - \|\nu\|^m + 1)$$

where $b(\nu) = b(\text{CL}_{\mathbb{N}, m}(\nu))$ is a constant independent of n . Now for any $(\pi_1^{(i)}, \pi_2^{(i)}) \in \mathbf{C}^{(i)}$,

$$\|\nu\|^m = |\text{Sup}^m(\pi_1^{(i)} \pi_2^{(i)})| \leq |\text{Sup}^m(\pi_1^{(i)}) \cup \text{Sup}^m(\pi_2^{(i)})| = \|\mathbf{C}^{(i)}\|^m.$$

Thus we have that $|A \cap (\mathfrak{S}_n \times \mathfrak{S}_n)|$ divided by $|\text{CL}_{n,m}(\nu)|$ is given by

$$b(\nu) \sum_{i \in I} \frac{1}{b(\mathbf{C}^{(i)})} (n-m - \|\nu\|^m)(n-m - \|\nu\|^m - 1) \cdots (n-m - \|\mathbf{C}^{(i)}\|^m + 1).$$

Hence let $f_{\lambda, \mu}^\nu(t)$ be the polynomial obtained from the above expression by replacing n with t . What remains to show is that $f_{\lambda, \mu}^\nu(t)$ belongs to R . It is known that a polynomial $f(t) \in \mathbb{Q}[t]$ of degree d belongs to R if and only if it is integer valued on $d+1$ consecutive integers. Well we have shown that each $f_{\lambda, \mu}^\nu(t)$ is integer valued on the infinite set $\{m, m+1, \dots\}$ since the evaluation at such an integer n gives the coefficient of the term $K_n(\nu)$ in the product $K_n(\lambda)K_n(\mu)$ in the \mathbb{Z} -algebra $Z_{n,m}$, hence we must have $f_{\lambda, \mu}^\nu(t) \in R$. \square

Specialising the above theorem to the case $m = 0$ recovers [FH59, Theorem 2.2]. It is worth remarking that in [FH59] they instead used *reduced partitions* as the indexing set for their classes as aposed to cycle shapes, but there is a natural correspondence between them.

4 The Farahat-Higman Algebras FH_m

Knowing that the structure constants of $Z_{n,m}$ are polynomial in n allows us to define a new R -algebra FH_m , which we refer to as the Farahat-Higman algebra associated to $Z_{n,m}$. This is constructed by replacing the structure constants $a_{\lambda, \mu}^\nu(n)$ with the structure polynomials $f_{\lambda, \mu}^\nu(t)$. We first define FH_m as a free R -module equipped with a product, and then shortly prove that such a product realises it as an R -algebra.

Definition 4.0.1. Let FH_m be the free R -module with basis $\{K(\lambda) \mid \lambda \in \Lambda(m)\}$. Equip this module with the product given by the R -linear extension of

$$K(\lambda)K(\mu) = \sum_{\nu \in \Lambda(m)} f_{\lambda,\mu}^\nu(t)K(\nu)$$

where $f_{\lambda,\mu}^\nu(t)$ are the structure polynomials given in *Theorem 3.0.3*.

Since $f_{\lambda,\mu}^\nu(n)$, for any $n \geq m$, gives the multiplicity of $K_n(\nu)$ in the product $K_n(\lambda)K_n(\mu)$, it is clear that $f_{\lambda,\mu}^\nu(t) = 0$ whenever $\|\nu\| > \|\lambda\| + \|\mu\|$. As such the product described above for FH_m is well-defined since only a finite number of terms will appear in the product of any two elements.

We say a *distributive ring* is an object satisfying all the axioms of a ring except possibly the associativity of the product and the existence of a multiplicative identity. In particular, a ring is a special case of a distributive ring. From definition we certainly have that FH_m is a distributive ring. We seek to show that the product given in *Definition 4.0.1* is associative and that FH_m possess a multiplicative identity. This will follow since FH_m is determined by the algebras $Z_{n,m}$ for all $n \geq m$, as we will now demonstrate.

By definition of the structure polynomials we must have a surjective homomorphism of distributive rings $\text{pr}_{n,m} : \text{FH}_m \rightarrow Z_{n,m}$ given by $\text{pr}_{n,m}(K(\lambda)) = K_n(\lambda)$ and $\text{pr}_{n,m}(f(t)) = f(n)$ for all $\lambda \in \Lambda(m)$ and $f(t) \in R$.

Lemma 4.0.2. *The intersection of the kernels of $\text{pr}_{n,m}$ for all $n \geq m$ is trivial, that is*

$$\bigcap_{n \geq m} \text{Ker}(\text{pr}_{n,m}) = \{0\}$$

Proof. Since $\{K_n(\lambda) \mid \lambda \in \Lambda_{\leq n}(m)\}$ forms a \mathbb{Z} -basis of $Z_{n,m}$, it is clear that the kernel of $\text{pr}_{n,m}$ is given by the sum $\text{Ker}(\text{pr}_{n,m}) = R^{(n)}\text{FH}_m + \text{Span}_R\{K(\lambda) \mid \text{deg}_m(\lambda) > n\}$, where $R^{(n)}$ is the subring of R given by all such polynomials with n as a root. Note for any $n_1, n_2 \geq m$ one has that

$$R^{(n_1)}\text{FH}_m \cap \text{Span}_R\{K(\lambda) \mid \text{deg}_m(\lambda) > n_2\} \subset R^{(n_1)}\text{FH}_m.$$

Therefore we have

$$\bigcap_{n \geq m} \text{Ker}(\text{pr}_{n,m}) \subset \left(\bigcap_{n \geq m} R^{(n)} \right) \text{FH}_m + \bigcap_{n \geq m} \text{Span}_R\{K(\lambda) \mid \text{deg}_m(\lambda) > n\} = 0,$$

since no element λ can have a degree greater than all $n \geq m$, and since the only polynomial with infinitely many roots is the trivial polynomial zero. □

Lemma 4.0.3. *For any $X, Y \in \text{FH}_m$, then $X = Y$ if and only if $\text{pr}_{n,m}(X) = \text{pr}_{n,m}(Y)$ for all $n \geq m$.*

Proof. The forward implication is immediate, while the reverse implication follows since it implies that $X - Y$ belongs to $\bigcap_{n \geq m} \text{Ker}(\text{pr}_{n,m}) = \{0\}$. □

Thus we can solve any product in FH_m by computing a corresponding one in $Z_{n,m}$ for arbitrary $n \geq m$.

Example 4.0.4. Consider the 2-marked cycle shapes $\lambda = (1, 2)(*, *)$ and $\mu = (1)(2)(*, *)$. We have that $\text{deg}_2(\lambda) = \text{deg}_2(\mu) = 4$, hence the class sums $K_n(\lambda)$ and $K_n(\mu)$ are non-zero if and only if $n \geq 4$. Let $n \geq 2$, then we have that

$$K_n(\lambda)K_n(\mu) = \left(\sum_{\{a,b\} \subseteq [n] \setminus [2]} (1, 2)(a, b) \right) \left(\sum_{\{c,d\} \subseteq [n] \setminus [2]} (c, d) \right) = \sum_{\{a,b\} \subseteq [n] \setminus [2]} \sum_{\{c,d\} \subseteq [n] \setminus [2]} (1, 2)(a, b)(c, d).$$

If the two cycles (a, b) and (c, d) are disjoint then the resulting permutation is simply $(a, b)(c, d)$, and there are two such ways to do so. If the two cycles share a single element, then the resulting permutation gives a 3-cycle (a, b, c) , and the number of ways to arrive at this 3-cycle from a product of two transpositions is

three. Lastly if the two cycles agree then the result is the identity, and there are as many ways to do this as there are two-element subsets of $[n] \setminus [2]$. Thus altogether we have that

$$K_n(\lambda)K_n(\mu) = 2K_n(\tau_1) + 3K_n(\tau_2) + \binom{n-2}{2}K_n(\tau_3)$$

where $\tau_1 = (1, 2)(*, *)(*, *)$, $\tau_2 = (1, 2)(*, *, *)$, and $\tau_3 = (1, 2)$. Note that

$$\binom{n-2}{2} = \frac{1}{2}(n-2)(n-3).$$

Thus $K_n(\lambda)K_n(\mu) = 2K_n(\tau_1) + 3K_n(\tau_2) + \frac{1}{2}(n-2)(n-3)K_n(\tau_3)$ for all $n \geq 2$, noting that when $n = 2, 3$ both sides of the equality are zero since $K_n(\tau) = 0$ for all $\tau \in \{\lambda, \mu, \tau_1, \tau_2\}$, and the polynomial in n which is the coefficient of $K_n(\tau_3)$ has both 2 and 3 as roots. Thus by *Lemma 4.0.3* we have that the relation

$$K(\lambda)K(\mu) = 2K(\tau_1) + 3K(\tau_2) + \frac{1}{2}(z-2)(z-3)K(\tau_3)$$

holds in FH_m , and so $f_{\lambda, \mu}^{\tau_1}(z) = 2$, $f_{\lambda, \mu}^{\tau_2}(z) = 3$, and $f_{\lambda, \mu}^{\tau_3}(z) = \frac{1}{2}(z-2)(z-3)$.

Proposition 4.0.5. *The distributive ring FH_m is an R -algebra.*

Proof. We need to prove that the product is associative and a multiplicative identity exists. For the identity, consider $K(\emptyset_m)$ where \emptyset_m is the m -marked cycle shape $(1)(2) \dots (m)$. The m -class $\text{CL}_{n,m}(\emptyset_m)$ contains only the identity, so $K_n(\emptyset_m)$ is the identity of $Z_{n,m}$ for any $n \geq m$. Thus $K(\emptyset_m)$ is the identity of FH_m by *Lemma 4.0.3*. For any $(X, Y, W) \in Z_{n,m}^{\times 3}$ let $[X, Y, W] := (XY)W - X(YW)$. Then $\text{pr}_{n,m}([X, Y, W]) = 0$ for all $n \geq m$ as $Z_{n,m}$ is associative. So $[X, Y, W] = 0$ by *Lemma 4.0.3*, showing that FH_m is associative. \square

We end this section by describing two natural R -subalgebras of FH_m .

Lemma 4.0.6. *We have an injective R -algebra homomorphism $R\mathfrak{S}_m \rightarrow \text{FH}_m$ defined by the R -linear extension of $\pi \mapsto K(\pi)$ for all $\pi \in \mathfrak{S}_m(\subset \Lambda(m))$.*

Proof. For $n \geq m$, $\text{CL}_{n,m}(\pi) = \{\pi\}$ for any $\pi \in \mathfrak{S}_m$. Hence $\text{pr}_{n,m}(K(\pi)) = K_n(\pi) = \pi$ showing we have a homomorphism by *Lemma 4.0.3*. Injectivity follows by construction of FH_m . \square

Consider the injective map $(-)_m : \Lambda(0) \rightarrow \Lambda(m)$ which sends a cycle shape $\lambda \in \Lambda(0)$ to the m -marked cycle shape λ_m which is obtained from λ by adjoining the trivial cycles $(1)(2) \dots (m)$.

Lemma 4.0.7. *We have an injective R -algebra homomorphism $\text{FH}_0 \rightarrow \text{FH}_m$ defined by the R -linear extension of $K(\lambda) \mapsto K(\lambda_m)$ for all $\lambda \in \Lambda(0)$.*

Proof. An application of *Lemma 4.0.3* shows that $f_{\lambda_m, \mu_m}^{\nu_m}(t) = 0$ for any $\lambda, \mu \in \Lambda(0)$ whenever $\nu \neq \gamma_m$ for $\gamma \in \Lambda(0)$. Thus it suffices to show that $f_{\lambda_m, \mu_m}^{\nu_m}(t) = f_{\lambda, \mu}^{\nu}(t)$ for any $\lambda, \mu, \nu \in \Lambda(0)$. Let FH_m^* denote the R -subalgebra of FH_m generated by $K(\lambda_m)$ for all $\lambda \in \Lambda(0)$. Clearly for any $n \geq 0$ we have that $\text{pr}_{n+m,m}(\text{FH}_m^*) = Z(\mathbb{Z}\mathfrak{S}([n+m] \setminus [m]))$. Similarly for any $n \geq 0$ we have that $\text{pr}_{n,0}(\text{FH}_0) = Z(\mathbb{Z}\mathfrak{S}_n)$. Clearly we have a \mathbb{Z} -algebra isomorphism $Z(\mathbb{Z}\mathfrak{S}([n+m] \setminus [m])) \cong Z(\mathbb{Z}\mathfrak{S}_n)$, which may be realised by the R -linear extension of conjugation $\sigma_n(-)\sigma_n^{-1} : Z(\mathbb{Z}\mathfrak{S}([n+m] \setminus [m])) \rightarrow Z(\mathbb{Z}\mathfrak{S}_n)$ where σ_n is any permutation of \mathfrak{S}_{n+m} such that $\sigma([n+m] \setminus [m]) = [n]$. By definition of $(-)_m$ we have that $K_n(\lambda) = \sigma_n K_{n+m}(\lambda_m) \sigma_n^{-1}$ for all $n \geq 0$ and $\lambda \in \Lambda(0)$. Applying *Lemma 4.0.3* we deduce that $f_{\lambda_m, \mu_m}^{\nu_m}(t) = f_{\lambda, \mu}^{\nu}(t)$ for any $\lambda, \mu, \nu \in \Lambda(0)$. Injectivity of $\text{FH}_0 \rightarrow \text{FH}_m$ follows since $\{K(\lambda_m) \mid \lambda \in \Lambda(0)\}$ is R -linearly independent by construction. \square

With this result in mind we will often identify FH_0 as the R -subalgebra of FH_m consisting of R -linear combination of basis elements $K(\lambda)$ where λ contains the trivial cycles $(1)(2) \dots (m)$, i.e. identifying FH_0 with the R -subalgebra FH_m^* described in the above proof. Since $\text{pr}_{n,0}(\text{FH}_0) = Z(\mathbb{Z}\mathfrak{S}_n)$, it is easy to deduce from *Lemma 4.0.3* that FH_0 is commutative, and hence a commutative R -subalgebra of FH_m .

5 Monoid of m -Marked Cycle Shapes

To show certain structural properties of the algebras FH_m it will prove helpful to equip the set of m -marked cycle shapes $\Lambda(m)$ with an associative product. To do such we will construct a monoid which we may identify with $\Lambda(m)$. This will also allow us to give a more concrete criteria for a permutation to belong to an m -class.

Let \mathcal{U}_m be the free commutative monoid on the set $\{u_1, \dots, u_m\}$. We have a natural left monoid action $\varphi : \mathfrak{S}_m \rightarrow \text{End}(\mathcal{U}_m)$ given by $\varphi(\pi)(u_i) = u_{\pi(i)}$ for all $\pi \in \mathfrak{S}_m$ and $i \in [m]$, where $\text{End}(\mathcal{U}_m)$ is the monoid of all monoid endomorphisms $\mathcal{U}_m \rightarrow \mathcal{U}_m$. Now let \mathcal{C} denote the free commutative monoid on the infinite set $\{c_1, c_2, \dots\}$. Then the monoid we are interested in is $(\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C}$, where \times_{φ} denotes the semidirect product with respect to the action φ . The underlying set is $\mathfrak{S}_m \times \mathcal{U}_m \times \mathcal{C}$ and the product is given by

$$(\pi, p, a)(\sigma, q, b) = (\pi\sigma, \varphi(\sigma)(p)q, ab)$$

for all $(\pi, p, a), (\sigma, q, b) \in \mathfrak{S}_m \times \mathcal{U}_m \times \mathcal{C}$. We abuse notation and write $\pi = (\pi, 1, 1)$, $p = (1, p, 1)$, and $a = (1, 1, a)$. For any set A we let $\mathbb{Z}_{\geq 0}^A$ denote the set of all functions $\mathbf{f} : A \rightarrow \mathbb{Z}_{\geq 0}$ with finite support, that is the subset of elements $a \in A$ such that $\mathbf{f}(a) \neq 0$ is finite. Then for any $\mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}$ and $\mathbf{l} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}}$ we let

$$u^{\mathbf{d}} := \prod_{i=1}^m u_i^{\mathbf{d}(i)} \quad \text{and} \quad c^{\mathbf{l}} := \prod_{i \in \mathbb{N}} c_i^{\mathbf{l}(i)}$$

which are well-defined by commutativity and since \mathbf{l} has finite support. Then as sets we have that

$$(\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C} = \left\{ \pi u^{\mathbf{d}} c^{\mathbf{l}} \mid \pi \in \mathfrak{S}_m, \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}, \mathbf{l} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}} \right\}.$$

Moreover, the product may be described by

$$(\pi u^{\mathbf{d}} c^{\mathbf{l}})(\sigma u^{\mathbf{e}} c^{\mathbf{k}}) = \pi \sigma u^{\sigma \circ \mathbf{d} + \mathbf{e}} c^{\mathbf{l} + \mathbf{k}} \quad (2)$$

where $\sigma \circ \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}$ is defined by $(\sigma \circ \mathbf{d})(i) = \mathbf{d}(\sigma^{-1}(i))$. In particular, the operator \circ realises the set $\mathbb{Z}_{\geq 0}^{[m]}$ as a left action set of \mathfrak{S}_m . The set $\Lambda(m)$ of m -marked cycle shapes is in a natural one-to-one correspondence with this monoid: Consider the map $(\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C} \rightarrow \Lambda(m)$ given by sending $\pi u^{\mathbf{d}} c^{\mathbf{l}}$ to the m -marked cycle shape consisting, for each $i \in \mathbb{N}$, $\mathbf{l}(i)$ many cycles of length $i + 1$ containing only the symbol $*$, and where the remaining cycles are constructed from those of π by adding $\mathbf{d}(i)$ symbols $*$ after the entry i , for each $i \in [m]$ (see examples below). This map is a bijection since there is a natural inverse to consider. With this correspondence in mind, we identify $\Lambda(m)$ with the monoid $(\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C}$.

Examples 5.0.1.

- (1) Let λ be the 6-marked cycle shape given in *Example 3.0.2*, then we have the identification

$$(1, 4)(2, 6)(3)(5)u_1^2 u_3^2 u_4^1 c_1^1 c_2^1 = (2, 6)(5)(1, *, *, *, 4, *) (3, *, *) (*, *) (*, *, *),$$

where we have added colours to aid in demonstrating the correspondence.

- (2) Consider the 4-marked cycle shapes written as elements of $(\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C}$ by $\lambda = (1, 2)(3, 4)u_1 u_2$, and $\mu = (1, 4)(2)(3)u_3^2 c_1$. Their product is given by

$$\lambda\mu = ((1, 2)(3, 4)u_1 u_2) ((1, 4)(2)(3)u_3^2 c_1) = (1, 3, 4, 2)u_2 u_3^2 u_4 c_1.$$

Thus when witting λ and μ as m -marked cycle shapes we have

$$\lambda\mu = ((1, *, 2, *) (3, 4)) ((1, 4)(2)(3, *, *) (*, *)) = (1, 3, *, *, 4, *, 2, *) (*, *).$$

Given any element $\lambda \in \Lambda(m)$, we will freely move between viewing it as an m -marked cycle shape or as an element of $(\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C}$, i.e. as an expression of the form $\pi u^{\mathbf{d}} c^{\mathbf{l}}$. As we know, a permutation σ belongs to the m -class $\text{CL}_{\mathbb{N}, m}(\lambda)$ if and only if σ may be obtained from λ by replacing the symbols $*$ with distinct elements from $\mathbb{N} \setminus [m]$. When expressing $\lambda = \pi u^{\mathbf{d}} c^{\mathbf{l}}$, then from the above discussion it is clear that we have the following equivalent criteria for when σ belongs to $\text{CL}_{\mathbb{N}, m}(\lambda)$ given in terms of π , \mathbf{d} , and \mathbf{l} .

Lemma 5.0.2. *We have $\sigma \in \text{CL}_{\mathbb{N},m}(\pi u^{\mathbf{d}} c^{\mathbf{l}})$ if and only if the following hold:*

- (i) *the number of cycles of σ of length $i + 1$ containing no elements of $[m]$ is $\mathbf{l}(i)$,*
- (ii) *$\sigma^{\mathbf{d}(i)+1}(i) = \pi(i)$ for each $i \in [m]$, and $\sigma^n(i) \notin [m]$ for any $1 \leq n \leq \mathbf{d}(i)$.*

□

Item (ii) above gives a concrete way of describing the relative positions of the elements of $[m]$ within a permutation σ , and is completely captured by π and \mathbf{d} . Recall the quantities $\|\lambda\|$, $\|\lambda\|_m$, and $\|\lambda\|^m$ for an m -marked cycle shape $\lambda \in \Lambda(m)$. Then when expressing $\lambda = \pi u^{\mathbf{d}} c^{\mathbf{l}}$, one can deduce that

$$\|\lambda\| = |\text{Sup}(\pi)| + \sum_{i \in [m]} \mathbf{d}(i) + \sum_{i \in \mathbb{N}} (i+1)\mathbf{l}(i), \quad \|\lambda\|^m = \sum_{i \in [m]} \mathbf{d}(i) + \sum_{i \in \mathbb{N}} (i+1)\mathbf{l}(i), \quad \|\lambda\|_m = |\text{Sup}(\pi)|.$$

Recall the degree function $\text{deg}_m : \Lambda(m) \rightarrow \mathbb{Z}_{\geq m}$ given by $\text{deg}_m(\lambda) = \|\lambda\|^m + m$. The codomain $\mathbb{Z}_{\geq m}$ is a monoid under the addition $a +_m b := a + b - m$ for all $a, b \in \mathbb{Z}_{\geq m}$ with m the identity. Viewing $\Lambda(m)$ as a monoid, then by consulting *Equation (2)* one can deduce that deg_m is a monoid homomorphism, and hence provides a grading for $\Lambda(m)$. Lastly, with the identification $\Lambda(m) = (\mathfrak{S}_m \times_{\varphi} \mathcal{U}_m) \times \mathcal{C}$ we may express the Farahat-Higman algebras by

$$\text{FH}_m = \text{Span}_R \{K(\pi u^{\mathbf{d}} c^{\mathbf{l}}) \mid \pi \in \mathfrak{S}_m, \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}, \mathbf{l} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}}\}.$$

Note that the R -subalgebra generated by the elements $K(c^{\mathbf{l}})$ for all $\mathbf{l} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}}$ is precisely the subspace FH_m^* described in the proof of *Lemma 4.0.7* consisting of all basis elements indexed by m -marked cycles shapes with the elements within $[m]$ appearing in trivial cycles. Therefore $\text{FH}_0 \cong \text{Span}_R \{K(c^{\mathbf{l}}) \mid \mathbf{l} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}}\} \subset \text{FH}_m$.

6 Leading Terms via Monoid Product

We may extend the degree function $\text{deg}_m : \Lambda(m) \rightarrow \mathbb{Z}_{\geq m}$ to one acting on FH_m by setting

$$\text{deg} \left(\sum_{\lambda \in \Lambda(m)} f_{\lambda}(t) K(\lambda) \right) = \max\{\text{deg}_m(\lambda) \mid f_{\lambda}(t) \neq 0\}.$$

In particular $\text{deg}_m(K(\lambda)) = \text{deg}_m(\lambda)$. We seek to show that the product of two basis elements of FH_m admits a unique leading term with regards to this degree function. To prove this we will use the following lemma.

Lemma 6.0.1. *For $\lambda, \mu \in \Lambda(m)$, let $g \in \text{CL}_{\mathbb{N},m}(\lambda)$ and $h \in \text{CL}_{\mathbb{N},m}(\mu)$. The equality*

$$\text{Sup}^m(g) \cap \text{Sup}^m(h) = \emptyset \tag{3}$$

holds if and only if $gh \in \text{CL}_{\mathbb{N},m}(\lambda\mu)$.

Proof. Suppose that $gh \in \text{CL}_{\mathbb{N},m}(\lambda\mu)$, then since deg_m is a monoid homomorphism

$$|\text{Sup}^m(gh)| = \|\lambda\mu\|^m = \|\lambda\|^m + \|\mu\|^m = |\text{Sup}^m(g)| + |\text{Sup}^m(h)|.$$

Since also $\text{Sup}^m(gh) \subset \text{Sup}^m(g) \cup \text{Sup}^m(h)$, we have $\text{Sup}^m(g) \cap \text{Sup}^m(h) = \emptyset$. This proves the “if” statement. For the “only if” let $\lambda = \pi u^{\mathbf{d}} c^{\mathbf{l}}$ and $\mu = \sigma u^{\mathbf{e}} c^{\mathbf{k}}$ for $\pi, \sigma \in \mathfrak{S}_m$, $\mathbf{d}, \mathbf{e} \in \mathbb{Z}_{\geq 0}^{[m]}$, and $\mathbf{l}, \mathbf{k} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}}$. By *Equation (2)*,

$$\lambda\mu = \pi\sigma u^{\sigma \circ \mathbf{d} + \mathbf{e}} c^{\mathbf{l} + \mathbf{k}}.$$

Hence the result follows if we show that gh satisfies items (i) and (ii) of *Lemma 5.0.2* with respect to $\pi\sigma u^{\sigma \circ \mathbf{d} + \mathbf{e}} c^{\mathbf{l} + \mathbf{k}}$. For item (i), since *Equation (3)* is upheld, it is clear that the number of cycles of gh of length $i + 1$ which contain no elements of $[m]$ is $\mathbf{l}(i) + \mathbf{k}(i)$, since this is the sum of such cycles of g and h . For item (ii), pick any $x \in [m]$ and set $y := \sigma(x)$ and $z := \pi(y)$. Since $h \in \text{CL}_{\mathbb{N},m}(\sigma u^{\mathbf{e}} c^{\mathbf{k}})$, item (ii) of

Lemma 5.0.2 tells us that $h : x \mapsto i_1 \mapsto i_2 \mapsto \cdots \mapsto i_{e(x)} \mapsto y$ where $\{i_1, i_2, \dots, i_{e(x)}\} \cap [m] = \emptyset$. Similarly since $g \in \mathbf{CL}_{\mathbb{N}, m}(\pi u^{\mathbf{d}} c^{\mathbf{l}})$, $g : y \mapsto j_1 \mapsto j_2 \mapsto \cdots \mapsto j_{\mathbf{d}(y)} \mapsto z$ where $\{j_1, j_2, \dots, j_{\mathbf{d}(y)}\} \cap [m] = \emptyset$. Since *Equation (3)* is upheld, we must have that

$$gh : x \mapsto i_1 \mapsto \cdots \mapsto i_{e(x)} \mapsto j_1 \mapsto \cdots \mapsto j_{\mathbf{d}(y)} \mapsto z.$$

Thus $(gh)^{\mathbf{d}(y)+e(x)+1}(x) = (\pi\sigma)(x)$ and $(gh)^n(x) \notin [m]$ for any $1 \leq n \leq \mathbf{d}(y) + e(x)$. One may also note that $\mathbf{d}(y) = \mathbf{d}(\sigma^{-1}(x)) = (\sigma \circ \mathbf{d})(x)$, and hence gh also upholds item (ii) of *Lemma 5.0.2*. \square

Proposition 6.0.2. *Let $\lambda = \pi u^{\mathbf{d}} c^{\mathbf{l}}, \mu = \sigma u^{\mathbf{e}} c^{\mathbf{k}} \in \Lambda(m)$. In \mathbf{FH}_m we have that*

$$K(\lambda)K(\mu) = c_{\lambda, \mu} K(\lambda\mu) + \sum_{\substack{\nu \in \Lambda(m) \\ \deg_m(\nu) < \deg_m(\lambda\mu)}} f_{\lambda, \mu}^{\nu}(t) K(\nu)$$

where $c_{\lambda, \mu} \in \mathbb{N}$ is a constant given by

$$c_{\lambda, \mu} = \prod_{i=1}^n \binom{(\mathbf{l} + \mathbf{k})(i)}{\mathbf{l}(i)},$$

Proof. Let $g \in \mathbf{CL}_{\mathbb{N}, m}(\lambda), h \in \mathbf{CL}_{\mathbb{N}, m}(\mu)$, then $\mathbf{Sup}^m(gh) \subset \mathbf{Sup}^m(g) \cup \mathbf{Sup}^m(h)$. Thus if $gh \in \mathbf{CL}_{\mathbb{N}, m}(\nu)$ for $\nu \in \Lambda(m)$ then $\|\nu\|^m \leq \|\lambda\|^m + \|\mu\|^m$. Hence $\deg_m(\nu) \leq \deg_m(\lambda) + \deg_m(\mu) = \deg_m(\lambda\mu)$. Thus we have

$$K(\lambda)K(\mu) = \sum_{\substack{\nu \in \Lambda(m) \\ \deg_m(\nu) \leq \deg_m(\lambda\mu)}} f_{\lambda, \mu}^{\nu}(t) K(\nu).$$

Suppose $gh \in \mathbf{CL}_{\mathbb{N}, m}(\nu)$ with $\deg_m(\nu) = \deg_m(\lambda\mu)$. This implies that $\|\nu\|^m = \|\lambda\mu\|^m$ and hence we must have $\mathbf{Sup}^m(g) \cap \mathbf{Sup}^m(h) = \emptyset$. Thus by *Lemma 6.0.1* we have that $\nu = \lambda\mu$. Therefore

$$K(\lambda)K(\mu) = f_{\lambda, \mu}^{\lambda\mu}(t) K(\lambda\mu) + \sum_{\substack{\nu \in \Lambda(m) \\ \deg_m(\nu) < \deg_m(\lambda\mu)}} f_{\lambda, \mu}^{\nu}(t) K(\nu).$$

It remains to show $f_{\lambda, \mu}^{\lambda\mu}(t) = c_{\lambda, \mu}$. Let $\omega \in \mathbf{CL}_{\mathbb{N}, m}(\lambda\mu)$ and $A_{\lambda, \mu}(\omega) := \{(g, h) \in \mathbf{CL}_{\mathbb{N}, m}(\lambda) \times \mathbf{CL}_{\mathbb{N}, m}(\mu) \mid gh = \omega\}$. By *Equation (2)* we have $\lambda\mu = \pi\sigma u^{\sigma \circ \mathbf{d} + \mathbf{e}} c^{\mathbf{l} + \mathbf{k}}$, and so by *Lemma 5.0.2*, for any $x \in [m]$, we have that $\omega : x \mapsto i_1 \mapsto \cdots \mapsto i_{(\sigma \circ \mathbf{d} + \mathbf{e})(x)} \mapsto (\pi\sigma)(x)$, where $\{i_1, \dots, i_{(\sigma \circ \mathbf{d} + \mathbf{e})(x)}\} \cap [m] = \emptyset$. Any pair $(g, h) \in A_{\lambda, \mu}(\omega)$ satisfies *Equation (3)*, and since their product gives ω we must have that

$$\begin{aligned} h : x \mapsto i_1 \mapsto \cdots \mapsto i_{e(x)} \mapsto \sigma(x), \\ g : \sigma(x) \mapsto i_{e(x)+1} \mapsto \cdots \mapsto i_{(\sigma \circ \mathbf{d} + \mathbf{e})(x)} \mapsto (\pi\sigma)(x). \end{aligned}$$

So if we construct a pair $(g, h) \in \mathbf{CL}_{\mathbb{N}, m}(\lambda) \times \mathbf{CL}_{\mathbb{N}, m}(\mu)$ such that $gh = \omega$, the cycles containing elements of $[m]$ in g and h are predetermined by ω . Hence we are just concerned with the cycles which contain no elements of $[m]$. In ω there are $(\mathbf{l} + \mathbf{k})(i)$ number of such cycles of length $i + 1$, while g and h contain $\mathbf{l}(i)$ and $\mathbf{k}(i)$ such cycles respectively. Thus to construct a pair $(g, h) \in A_{\lambda, \mu}(\omega)$, it is simply a matter of how one distributes the cycles containing no elements of $[m]$ of ω among either g or h . The binomial coefficient

$$\binom{(\mathbf{l} + \mathbf{k})(i)}{\mathbf{l}(i)}$$

counts the number of ways to allocate such cycles of length $i + 1$ of ω to the permutation g (with the remaining cycles allocated to h). Therefore

$$f_{\lambda, \mu}^{\lambda\mu}(t) = |A_{\lambda, \mu}(\omega)| = \prod_{i=1}^{\infty} \binom{(\mathbf{l} + \mathbf{k})(i)}{\mathbf{l}(i)},$$

where the product is only formally infinite since the functions $(\mathbf{l} + \mathbf{k})$, \mathbf{l} , and \mathbf{k} have finite support. \square

Thus the leading term in the product $K(\lambda)K(\mu)$ is $c_{\lambda,\mu}K(\lambda\mu)$. So the product of $\Lambda(m)$ is governing the leading terms. From the formula given above one can deduce that $c_{\lambda,\mu} = 1$ if and only if λ and μ share no cycles of the same length consisting of only the symbols $*$, i.e. whenever $\mathbf{l}(i) \neq 0$ then $\mathbf{k}(i) = 0$.

If one was to extend R to a ring which contains the inverses to any natural number, say its field of fractions $\mathbb{Q}(t)$, then the above leading term result could be applied to easily deduce generating sets for the $\mathbb{Q}(t)$ -algebra $\mathbb{Q}(t) \otimes \text{FH}_m$. For example, let $s_i = (i, i+1)$ for any $i \in [m]$ be the simple transposition in \mathfrak{S}_m , then by arguing by induction on the degree, *Proposition 6.0.2* can be employed to show that the set

$$\{K(s_i), K(u_i), K(c_j) \mid i \in [m], j \in \mathbb{N}\} \quad (4)$$

generates $\mathbb{Q}(t) \otimes \text{FH}_m$ as a $\mathbb{Q}(t)$ -algebra since $\{s_i, u_i, c_j \mid i \in [m], j \in \mathbb{N}\}$ generates $\Lambda(m) = (\mathfrak{S}_m \ltimes_{\varphi} \mathcal{U}_m) \times \mathcal{C}$.

7 Symmetric Functions

Consider the polynomial \mathbb{Z} -algebra $\mathbb{Z}[x_1, \dots, x_n]$ in n commuting variables. The group \mathfrak{S}_n acts on such polynomials by permuting the variables. A polynomial is *symmetric* if it is \mathfrak{S}_n -invariant. Denote the \mathbb{Z} -subalgebra of symmetric polynomials by Sym_n . We are interested in two types of symmetric polynomials, and to describe one type it will be helpful to recall the definition of a partition. We say a tuple $\alpha = (a_1, a_2, \dots, a_l)$ of positive integers is a partition whenever $a_1 \geq a_2 \geq \dots \geq a_l$. We say α has length l and size $a_1 + \dots + a_l$. Let $\mathcal{P}(k, l)$ be the set of partitions of size k and length l . When $k = l = 0$ let $\mathcal{P}(0, 0) = \{\emptyset\}$.

Definition 7.0.1. For any $k \geq 0$, the *elementary symmetric polynomials* are given by

$$e_k(x_1, \dots, x_n) := \sum_{i_1 < i_2 < \dots < i_k} x_{i_1} x_{i_2} \cdots x_{i_k}$$

Hence e_k is the sum of all monomials with k variables. In particular $e_0 = 1$ and $e_k = 0$ for $k > n$.

Definition 7.0.2. For $\alpha = (a_1, a_2, \dots, a_l) \in \mathcal{P}(k, l)$, the *monomial symmetric polynomials* are given by

$$m_{\alpha}(x_1, \dots, x_n) := \sum_{\{i_1, i_2, \dots, i_l\} \subset [n]} x_{i_1}^{a_1} x_{i_2}^{a_2} \cdots x_{i_l}^{a_l}$$

Hence m_{α} is the sum of all monomials whose exponents match the partition α up to rearrangement. In particular $m_{\emptyset} = 1$ and $m_{\alpha} = 0$ whenever $l > n$.

It is well known that Sym_n is generated (transcendentally) by e_1, e_2, \dots, e_n . That is we have a \mathbb{Z} -algebra isomorphism $\text{Sym}_n \cong \mathbb{Z}[e_1, \dots, e_n]$ with the latter being a free polynomial algebra in n commuting generators. One may show that $\{m_{\alpha} \mid \alpha \in \mathcal{P}(k, l), k \geq 0, n \geq l \geq 0\}$ forms a \mathbb{Z} -basis for Sym_n . Assign each variable x_i a degree of 1, then let Sym_n^k denote the degree k component of Sym_n . In particular $e_k, m_{\alpha} \in \text{Sym}_n^k$ for any $\alpha \in \mathcal{P}(k, l)$. For any $N > n$ we have surjective \mathbb{Z} -module homomorphisms $\rho_{N,n} : \text{Sym}_N^k \rightarrow \text{Sym}_n^k$ given by evaluating the variables x_{n+1}, \dots, x_N at zero. One can show that the collection of such morphisms defines an inverse system for the \mathbb{Z} -modules Sym_n^k (with fixed k), and so we have the inverse limit

$$\text{Sym}^k := \varprojlim \text{Sym}_n^k.$$

Any element of Sym^k is of the form (f_1, f_2, \dots) with $f_n \in \text{Sym}_n^k$ and where $\rho_{N,n}(f_N) = f_n$ for all $N > n$. One can note that $\rho_{N,n}(m_{\alpha}(x_1, \dots, x_N)) = m_{\alpha}(x_1, \dots, x_n)$ for any $N > n$ and $\alpha \in \mathcal{P}(k, l)$. Thus we write $m_{\alpha} = (m_{\alpha}(x_1), m_{\alpha}(x_1, x_2), \dots)$ and call such elements of Sym^k the *monomial symmetric functions*. The same can be said for e_k since $e_k = m_{(1^k)}$ where (1^k) is the partition of size k consisting of k parts equal to 1. The \mathbb{Z} -algebra of *symmetric functions* is given by

$$\text{Sym} := \bigoplus_{k \in \mathbb{Z}_{\geq 0}} \text{Sym}^k.$$

We have the isomorphism of \mathbb{Z} -algebras $\text{Sym} \cong \mathbb{Z}[e_1, e_2, \dots]$, and a \mathbb{Z} -basis $\{m_{\alpha} \mid \alpha \in \mathcal{P}(k, l), k, l \geq 0\}$.

8 Jucys-Murphy Elements and Generators of \mathbf{FH}_0

In this section we recall the Jucys-Murphy elements of the symmetric group algebras, and their connections to the Farahat-Higman algebras.

Definition 8.0.1. For each $1 \leq i \leq n$, the i -th *Jucys-Murphy element* L_i of $\mathbb{Z}\mathfrak{S}_n$ is defined by

$$L_i := \sum_{j < i} (j, i).$$

Note that $L_1 = 0$. The following relations regarding the Jucys-Murphy elements are well-known.

Lemma 8.0.2. *The following relations hold within $\mathbb{Z}\mathfrak{S}_n$:*

- (1) $L_i L_j = L_j L_i$ for all $i, j \in [n]$
- (2) $s_i L_j = L_j s_i$ for all $i \in [n-1]$ and $j \neq i, i+1$
- (3) $L_{i+1} = s_i L_i s_i + s_i$ for all $i \in [n-1]$

where $s_i = (i, i+1)$ is the simple transposition exchanging i and $i+1$ for all $i \in [n-1]$.

It was shown in [Murphy83, Theorem 1.9] that the center is precisely the collection of all symmetric polynomials in the Jucys-Murphy elements. Then from the last section we know that

$$Z(\mathbb{Z}\mathfrak{S}_n) = \langle e_1(L_1, \dots, L_n), \dots, e_n(L_1, \dots, L_n) \rangle. \quad (5)$$

We know the center $Z(\mathbb{Z}\mathfrak{S}_n)$ has a \mathbb{Z} -basis given by the set $\{K_n(\lambda) \mid \lambda \in \Lambda_{\leq n}(0)\}$. It is natural to ask how the elementary symmetric polynomials in the Jucys-Murphy elements decompose as a linear combination of class sums. This was answered in [Jucys74, Section 3] as we recall now. Let $\lambda \in \Lambda(0)$ contain l many cycles. We say the *reduced degree* of λ is $\text{rd}(\lambda) := \text{deg}_0(\lambda) - l$, i.e. the number of symbols $*$ present in λ minus the number of cycles. Then it was shown that

$$e_k(L_1, \dots, L_n) = \sum_{\substack{\lambda \in \Lambda_{\leq n}(0) \\ \text{rd}(\lambda) = k}} K_n(\lambda)$$

for any $k \in [n]$. Recall that $\text{pr}_{n,0}(\mathbf{FH}_0) = Z(\mathbb{Z}\mathfrak{S}_n)$, and from the above formula there is natural elements of \mathbf{FH}_0 to consider which project down to the elementary symmetric polynomials in the Jucys-Murphy elements.

Definition 8.0.3. For any $k \in \mathbb{Z}_{\geq 0}$ define

$$E_k := \sum_{\substack{\lambda \in \Lambda(0) \\ \text{rd}(\lambda) = k}} K(\lambda),$$

Note only finitely many elements of $\Lambda(0)$ have reduced degree k . Thus $\text{pr}_{n,0}(E_k) = e_k(L_1, \dots, L_n)$ for any $n \geq 0$ and when $k > n$ both $\text{pr}_{n,0}(E_k)$ and $e_k(L_1, \dots, L_n)$ are zero. The images of E_1, E_2, \dots, E_n under $\text{pr}_{n,0}$ thus give a generating set for $Z(\mathbb{Z}\mathfrak{S}_n)$. Relation-wise \mathbf{FH}_0 is determined by the centers $Z(\mathbb{Z}\mathfrak{S}_n)$, so one may expect the elements E_1, E_2, \dots to generate \mathbf{FH}_0 . This was proved in [FH59], which we record here.

Theorem 8.0.4. [Theorem 2.5 of [FH59]] *The set of elements $\{E_1, E_2, \dots\}$ generates \mathbf{FH}_0 as an R -algebra.*

Furthermore, it was proven in [Ryba21, Theorem 3.8] that there is an isomorphism $\mathbf{FH}_0 \cong R \otimes_{\mathbb{Z}} \text{Sym}$ of R -algebras which sends e_k to E_k . In the next section we will prove an analogous result for \mathbf{FH}_m in *Theorem 9.0.5*, and to do so it will be helpful to discuss elements of \mathbf{FH}_0 which are to the monomial symmetric polynomials m_α what the elements E_k are to the elementary symmetric polynomials e_k .

Lemma 8.0.5. *For any $k, l \geq 0$ and $\alpha \in \mathcal{P}(k, l)$, there exists an element $M_\alpha \in \mathbf{FH}_0$ such that*

$$\text{pr}_{n,0}(M_\alpha) = m_\alpha(L_1, \dots, L_n).$$

Proof. As $\text{Sym} = \mathbb{Z}[e_1, e_2, \dots]$ then m_α is a finite \mathbb{Z} -linear combination of monomials in e_1, e_2, \dots . Letting M_α be obtained from m_α by replacing e_k with E_k gives the element we are looking for. \square

We wish to say a little more about the elements M_α . For any $\alpha = (a_1, \dots, a_l) \in \mathcal{P}(k, l)$ let $\bar{\alpha} \in \Lambda(0)$ denote the cycle shape with l cycles of lengths $a_1 + 1, a_2 + 1, \dots, a_l + 1$. When $\alpha = \emptyset$ then $\bar{\alpha} = \emptyset$ also. We have that $\text{rd}(\bar{\alpha}) = k$, the size of α . Then in the proof of Theorem 1.9 of [Murphy83], see also [Ryba21, Proposition 3.11], the following proposition was shown. Let $\mathcal{P} := \cup_{k, l \geq 0} \mathcal{P}(k, l)$ and note that the map $\overline{(-)} : \mathcal{P} \rightarrow \Lambda(0)$ sending $\alpha \mapsto \bar{\alpha}$ is a bijection.

Proposition 8.0.6. *Let $\alpha \in \mathcal{P}$ such that $\text{deg}_0(\bar{\alpha}) \leq n$, then*

$$m_\alpha(L_1, \dots, L_n) = K_n(\bar{\alpha}) + \sum c_\mu(n) K_n(\mu),$$

where the sum runs over all $\mu \in \Lambda(0)$ such that $\text{rd}(\mu) < \text{rd}(\bar{\alpha})$ or that $\text{rd}(\mu) = \text{rd}(\bar{\alpha})$ and μ contains less cycles than $\bar{\alpha}$ (noting that only finitely many such μ exist), and where $c_\mu(n) \in \mathbb{Z}_{\geq 0}$.

Applying Lemma 4.0.3 allows us to deduce the following:

Lemma 8.0.7. *Let $\alpha \in \mathcal{P}$ then*

$$M_\alpha = K(\bar{\alpha}) + \sum c_\mu(t) K(\mu), \tag{6}$$

where the sum runs over all $\mu \in \Lambda(0)$ such that $\text{rd}(\mu) < \text{rd}(\bar{\alpha})$ or that $\text{rd}(\mu) = \text{rd}(\bar{\alpha})$ and μ contains less cycles than $\bar{\alpha}$ (noting that only finitely many such μ exist), and where $c_\mu(t) \in R$ are such that $c_\mu(n)$ are the coefficients of $K_n(\mu)$ in the above proposition.

We may now show that the elements M_α provide an R -basis for FH_0 .

Proposition 8.0.8. *The set $\{M_\alpha \mid \alpha \in \mathcal{P}\}$ gives an R -basis for FH_0 .*

Proof. By Equation (6) it is clear that $\{M_\alpha \mid \alpha \in \mathcal{P}\}$ is R -linearly independent, hence we only need to show that it spans FH_0 . Equip the R -basis $\{K(\lambda) \mid \lambda \in \Lambda(0)\}$ of FH_0 with the partial order $<$ defined by letting $K(\mu) < K(\lambda)$ if $\text{rd}(\mu) < \text{rd}(\lambda)$ or if $\text{rd}(\mu) = \text{rd}(\lambda)$ and μ contains less cycles than λ . For any $n \in \mathbb{Z}_{\geq 0}$ let $\text{FH}_0^{\leq n}$ denote the \mathbb{Z} -module spanned by the elements $K(\lambda)$ such that $\text{rd}(\lambda) \leq n$. We prove that $\{M_\alpha \mid \alpha \in \mathcal{P}\}$ spans $\text{FH}_0^{\leq n}$ by induction on n . For the base case we have $\text{FH}_0^{\leq 0} = \text{Span}_R\{K(\emptyset)\}$, and hence this case follows since $M_\emptyset = K(\emptyset) = K(\emptyset)$. Consider $\text{FH}_0^{\leq n}$ for some $n > 1$, and let

$$K = \sum_{\substack{\lambda \in \Lambda(0) \\ \text{rd}(\lambda) \leq n}} f_\lambda(t) K(\lambda)$$

be an element of $\text{FH}_0^{\leq n}$. Let $K(\lambda')$ be maximal in $\{K(\lambda) \mid f_\lambda(t) \neq 0\}$ with respect to $<$. Then let α be the unique partition of \mathcal{P} such that $\bar{\alpha} = \lambda'$, then $K - f_{\lambda'}(t)M_\alpha$ is an element of $\text{FH}_0^{\leq n}$ whose terms are strictly less than $K(\lambda')$ in the partial order or incomparable. Continuing this procedure of removing the maximal basis element will lead to an element of $\text{FH}_0^{\leq n-1}$, and by induction the resulting element will belong to the span of $\{M_\alpha \mid \alpha \in \mathcal{P}\}$. Since we got there by subtracting some R -linear combination of elements M_α , the starting element K must also belong to the span of $\{M_\alpha \mid \alpha \in \mathcal{P}\}$ which completes the proof by induction. The proposition follows since FH_0 is the union of $\text{FH}_0^{\leq n}$ for all $n \geq 0$. \square

9 An Isomorphism $\text{FH}_m \cong R \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$

In this section we recall the definition of the degenerate affine Hecke algebra \mathcal{H}_m , and some of its structural properties. We define some elements of FH_m which mimic the variables generators of \mathcal{H}_m , and end by proving the isomorphism in the section title.

Definition 9.0.1. The *degenerate affine Hecke Algebra* \mathcal{H}_m is the \mathbb{Z} -algebra presented with generating set $\{s_i, y_j \mid 1 \leq i \leq m-1, j \in [m]\}$ and defining relations

- (1i) $s_i^2 = 1$, for $i \in [m-1]$. (2i) $y_i y_j = y_j y_i$ for all $i, j \in [m]$.
(1ii) $s_i s_j = s_j s_i$, for $j \neq i-1, i+1$. (2ii) $s_i y_j = y_j s_i$ for all $j \neq i, i+1$.
(1iii) $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, for $i \in [m-2]$. (2iii) $y_{i+1} = s_i y_i s_i + s_i$ for all $i \in [m-1]$.

The elements s_i are the simple transpositions $(i, i+1)$ of \mathfrak{S}_m . The algebra \mathcal{H}_m has a basis of the form

$$\{\pi y_1^{\mathbf{d}(1)} \cdots y_m^{\mathbf{d}(m)} \mid \pi \in \mathfrak{S}_m, \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}\}. \quad (7)$$

For a proof see [K05, Theorem 3.2.2]. Hence one can deduce that the subalgebra generated by the elements s_i is a copy of $\mathbb{Z}\mathfrak{S}_m$, and similarly the subalgebra generated by the variable generators y_1, \dots, y_m is a copy of the polynomial \mathbb{Z} -algebra $\mathbb{Z}[y_1, \dots, y_m]$. Moreover, as a \mathbb{Z} -module we have $\mathcal{H}_m \cong \mathbb{Z}\mathfrak{S}_m \otimes \mathbb{Z}[y_1, \dots, y_m]$. By *Lemma 8.0.2*, we have a surjective \mathbb{Z} -algebra homomorphism $\mathcal{H}_m \rightarrow \mathbb{Z}\mathfrak{S}_m$ given by $s_i \mapsto s_i$ and $y_j \mapsto L_j$.

We now show that the algebra FH_m contains like-minded elements.

Definition 9.0.2. For any $i \in [m]$ define elements $Y_i \in \text{FH}_m$ by

$$Y_i := K(u_i) + \sum_{\substack{j \in [m] \\ j < i}} K((j, i)) = K(u_i) + L_i,$$

where we have identified $R\mathfrak{S}_m$ with $\{K(\pi) \mid \pi \in \mathfrak{S}_m \subset \Lambda(m)\}$.

Example 9.0.3. In FH_3 we have

$$Y_1 = K((* , 1)), \quad Y_2 = K((* , 2)) + K((1, 2)), \quad Y_3 = K((* , 3)) + K((1, 3)) + K((2, 3)),$$

where we have dropped the trivial cycles (1), (2), and (3) from each of the 3-marked cycle shapes appearing.

Since $\deg_m(u_i) = m+1$ we have that $\text{pr}_{m,m}(K(u_i)) = 0$, hence $\text{pr}_{m,m}(Y_i) = L_i$. However, one can deduce that when $n > 0$ the elements $\text{pr}_{n+m,m}(Y_i)$ do not get sent to the Jucys-Murphy elements L_i of $\mathbb{Z}\mathfrak{S}_{n+m}$ but rather $\sigma_n L_{i+m} \sigma_n^{-1}$ for any $\sigma_n \in \mathfrak{S}_{n+m}$ for which $\sigma_n([m]) = \{n+1, \dots, n+m\}$. Nonetheless they retain counterparts to the relations of *Lemma 8.0.2*:

Lemma 9.0.4. *The following relations hold within FH_m :*

- (1) $Y_i Y_j = Y_j Y_i$ for all $i, j \in [n]$
- (2) $K(s_i) Y_j = Y_j K(s_i)$ for all $i \in [n-1]$ and $j \neq i, i+1$
- (3) $Y_{i+1} = K(s_i) Y_i K(s_i) + K(s_i)$ for all $i \in [n-1]$

where $s_i = (i, i+1)$ is the simple transposition exchanging i and $i+1$ for all $i \in [n-1]$.

Proof. (1): For any $n \geq 0$ and $i, j \in [m]$ we have

$$\text{pr}_{n+m,m}(Y_i Y_j) = \sigma_n L_{i+m} L_{j+m} \sigma_n^{-1} = \sigma_n L_{j+m} L_{i+m} \sigma_n^{-1} = \text{pr}_{n+m,m}(Y_j Y_i),$$

since the Jucys-Murphy elements commute. So applying *Lemma 4.0.3* shows that $Y_i Y_j = Y_j Y_i$. (2): It follows as a straight forward application of *Lemma 4.0.3* that $K(s_i)$ commutes with $K(u_j)$ whenever $i \in [n-1]$ and $j \neq i, i+1$, and we already know that $K(s_i)$ commutes with L_j , so (2) follows. (3): Again by *Lemma 4.0.3* one can show that $K(s_i) K(u_i) K(s_i) = K(u_{i+1})$, and hence

$$K(s_i) Y_i K(s_i) = K(u_{i+1}) + s_i L_i s_i = K(u_i) + L_{i+1} - K(s_i) = Y_j - K(s_i),$$

where we used (3) of *Lemma 8.0.2*. Rearranging gives (3). □

Theorem 9.0.5. *We have an isomorphism of R -algebras $\phi : R \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym}) \rightarrow \text{FH}_m$ defined by $s_i \mapsto K(s_i)$, $y_j \mapsto Y_j$, and $e_k \mapsto E_k$ for each $i \in [m-1]$, $j \in [m]$, and $k \geq 0$.*

Proof. To prove ϕ is a homomorphism it suffices to show that the relations of *Definition 9.0.1* are respected, and that the elements $\phi(e_k)$ commute with $\text{Im}(\phi)$. Relations (1) of *Definition 9.0.1* are respected by *Lemma 4.0.6*, and relations (2) are upheld by *Lemma 9.0.4*. Also $\phi(e_k) = E_k \in \text{FH}_0 \subset \text{FH}_m$, hence the images $\phi(e_1), \phi(e_2), \dots$ commute with one another since FH_0 is commutative. Moreover, since they belong to FH_0 , their projections down to $Z_{n,m}$ by $\text{pr}_{n,m}$ consist of permutations which act trivially on $[m]$. Therefore $\phi(e_k)$ commutes with $\phi(s_i) = K(s_i)$ for any $k \geq 0$ and $i \in [m-1]$. Lastly we have that

$$[\phi(e_k), \phi(y_j)] = [\phi(e_k), Y_j] = [\phi(e_k), K(u_j)].$$

We have that $\text{pr}_{n,m}(K(u_j))$ is the sum of transpositions (a, j) for all $a \in [n] \setminus [m]$, which commutes with any permutation which fixes the elements $[m]$. In particular $\text{pr}_{n,m}(K(u_j))$ must commute with $\text{pr}_{n,m}(\phi(e_k))$, and so by *Lemma 4.0.3* we have that $K(u_j)$ commutes with $\phi(e_k)$. Hence $[\phi(e_k), \phi(y_j)] = 0$, and so $\phi(e_k)$ commutes with $\text{Im}(\phi)$. So ϕ is an R -algebra homomorphism. For surjectivity we show $K(\lambda) \in \text{Im}(\phi)$ for any $\lambda \in \Lambda(m)$. Write $\lambda = \pi u^{\mathbf{d}} c^{\mathbf{l}}$ and consider the element $K(c^{\mathbf{l}})$ which belongs to the R -subalgebra FH_0 of FH_m . By *Theorem 8.0.4* there exists $C \in \langle \phi(e_1), \phi(e_2), \dots \rangle$ such that $\phi(C) = K(c^{\mathbf{l}})$. Then by employing the leading term result of *Proposition 6.0.2*, we have that

$$\phi\left(\pi y_1^{\mathbf{d}^{(1)}} \dots y_m^{\mathbf{d}^{(m)}} C\right) = K(\pi) Y_1^{\mathbf{d}^{(1)}} \dots Y_m^{\mathbf{d}^{(m)}} K(c^{\mathbf{l}}) = K(\pi u^{\mathbf{d}} c^{\mathbf{l}}) + T$$

where T stands for an R -linear combination of terms $K(\mu)$ such that $\text{deg}_m(\mu) < \text{deg}_m(\lambda)$. Hence arguing by induction on the degree of λ shows surjectivity, noting that the base case is immediate since the basis elements of degree zero are precisely $K(\pi) = \phi(\pi)$ for some $\pi \in \mathfrak{S}_m \subset \Lambda(m)$. For injectivity, by *Equation (7)* and recalling that the monomial symmetric functions form a \mathbb{Z} -basis of Sym , we have that the set

$$\mathbf{B} := \left\{ \pi y_1^{\mathbf{d}^{(1)}} \dots y_m^{\mathbf{d}^{(m)}} \otimes m_\alpha \mid \pi \in \mathfrak{S}_m, \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}, \alpha \in \mathcal{P} \right\}$$

forms an R -basis of $R \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$. We seek to show that $\phi(\mathbf{B})$ is R -linearly independent. Equip the basis set $\{K(\pi u^{\mathbf{d}} c^{\mathbf{l}}) \mid \pi \in \mathfrak{S}_m, \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}, \mathbf{l} \in \mathbb{Z}_{\geq 0}^{\mathbb{N}}\}$ of FH_m with the partial order $<$ define by $K(\sigma u^e c^k) < K(\pi u^{\mathbf{d}} c^{\mathbf{l}})$ if (i) the degree of $\sigma u^e c^k$ is strictly less than that of $\pi u^{\mathbf{d}} c^{\mathbf{l}}$, (ii) their degrees agree but $\text{rd}(c^k) < \text{rd}(c^{\mathbf{l}})$, (iii) their degrees agree and $\text{rd}(c^k) = \text{rd}(c^{\mathbf{l}})$ but c^k contains less cycles than $c^{\mathbf{l}}$. Note *Proposition 6.0.2* tells us that the product of $K(\lambda)$ and $K(\mu)$ in FH_m results in $c_{\lambda, \mu} K(\lambda \mu)$ plus addition terms all lower in the order $<$ where $c_{\lambda, \mu} \in \mathbb{N}$. Recall that $c_{\lambda, \mu} = 1$ whenever λ and μ share no cycles of the same size which contain only the symbol $*$. From *Equation (6)* and *Proposition 6.0.2* we have that ϕ acts on a basis element of \mathbf{B} by

$$\phi\left(\pi y_1^{\mathbf{d}^{(1)}} \dots y_m^{\mathbf{d}^{(m)}} \otimes m_\alpha\right) = K(\pi) Y_1^{\mathbf{d}^{(1)}} \dots Y_m^{\mathbf{d}^{(m)}} M_\alpha = K(\pi u^{\mathbf{d}} \bar{\alpha}) + T$$

where T is an R -linear combination of basis elements $K(\lambda)$ of FH_m which are strictly less with respect to $<$. As such, in the image of any finite R -linear combination of elements of \mathbf{B} under ϕ , we may pick out a non-zero term which is uncomparable or strictly greater than any other term with respect to $<$, showing that $\phi(\mathbf{B})$ is R -linearly independent in FH_m , which shows ϕ is injective. \square

The \mathbb{C} -algebra $\mathbb{C} \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$ and close variations have made appearances within the literature. In the [MO01] they aimed to give a centralizer construction for the degenerate affine Hecke algebra in a manner comparable to how the Yangians arise from a projective limit of universal enveloping algebras of \mathfrak{gl}_n . No such projective system exists for the group algebras of the symmetric groups, but does for the larger semigroup of partial permutations. Working with these instead algebras $A_m := A_0 \otimes \tilde{\mathcal{H}}_m$ were constructed, where $\tilde{\mathcal{H}}_m$ is a degenerate affine counterpart to the semigroup algebra of partial permutations, and A_0 was shown to be isomorphic to the algebra of shifted symmetric functions. In our setting the lack of a projective system was sidestepped by employing the techniques of Farahat and Higman on the centralizer algebras $Z_{n,m}$, which allowed us to stay working with the symmetric group itself. The \mathbb{C} -algebra $\mathbb{C} \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$ also appears in the Heisenberg category Heis of M. Khovanov defined in [Kho14]. Such a category is a monoidal category generated by two objects \uparrow and \downarrow , and where the morphism spaces are defined diagrammatically. It was shown in [Kho14, Proposition 4] that the endomorphism algebra $\text{End}_{\text{Heis}}(\uparrow^{\otimes m})$ is isomorphic to $\mathbb{C} \otimes_{\mathbb{Z}} (\mathcal{H}_m \otimes \text{Sym})$.

We close this section with some consequences of the above theorem.

Corollary 9.0.6. *The algebra FH_m has a R -basis given by the set*

$$\left\{ K(\pi)Y_1^{\mathbf{d}(1)} \cdots Y_m^{\mathbf{d}(m)} M_\alpha \mid \pi \in \mathfrak{S}_m, \mathbf{d} \in \mathbb{Z}_{\geq 0}^{[m]}, \alpha \in \mathcal{P} \right\}.$$

Corollary 9.0.7. *The algebra FH_m is generated by $K(s_i)$, Y_j , and E_k for $i \in [m-1]$, $j \in [m]$, and $k \geq 0$.*

Corollary 9.0.8. *The center of FH_m is $\text{Sym}[Y_1, \dots, Y_m] \otimes \text{FH}_0$, which is generated by the elements E_k and the elementary symmetric polynomials $e_k(Y_1, \dots, Y_m)$ for all $k \geq 0$.*

Proof. By Theorem 9.0.5 it is clear that the result follows if $\text{Sym}[y_1, \dots, y_m]$ is the center of \mathcal{H}_m , which is shown in [K05, Theorem 3.3.1]. □

References

- [FH59] H. Farahat and G. Higman, *The centres of symmetric group rings*, Proc. Roy. Soc. (A) 250 (1959), 212–221. URL: <https://doi.org/10.1098/rspa.1959.0060>
- [GJ94] I. Goulden and D. Jackson, *Symmetric functions and Macdonald’s result for top connexion coefficients in the symmetric group*, J. Algebra 166 (1994), no. 2, 364–378. URL: <https://doi.org/10.1006/jabr.1994.1157>
- [Jucys74] A. Jucys, *Symmetric polynomials and the center of the symmetric group ring*, Rep. Mathematical Phys. 5 (1974), no. 1, 107–112. MR 0419576 (54 7597), URL: [https://doi.org/10.1016/0034-4877\(74\)90019-6](https://doi.org/10.1016/0034-4877(74)90019-6)
- [Kho14] M. Khovanov, *Heisenberg algebra and a graphical calculus*, Fundamenta Mathematicae 225 (2014), 169–210, URL: <https://arxiv.org/abs/1009.3295>
- [K05] A. Kleshchev, *Linear and projective representations of the symmetric group*, Cambridge Tract in Mathematics, series number 163, CUP, 2005.
- [Mac95] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford University Press, 1995.
- [MO01] A.I. Molev and G.I. Olshanski, *Degenerate affine Hecke algebras and centralizer construction for the symmetric groups*, J. Algebra 237 (1) (2001) 302–341.
- [Murphy83] G. E. Murphy, *The idempotents of the symmetric group and Nakayama’s conjecture*, J. Algebra 81 (1983), no. 1, 258–265. MR 696137 (84k:20007), URL: [https://doi.org/10.1016/0021-8693\(83\)90219-3](https://doi.org/10.1016/0021-8693(83)90219-3)
- [Ryba21] C. Ryba, *Stable Centres I: Wreath Products* URL: <https://doi.org/10.48550/arXiv.2107.03752>
- [TW09] J. Tysse and W. Wang, *The centers of spin symmetric group algebras and Catalan numbers* J Algebr Comb (2009) 29: 175–193 URL: <https://doi.org/10.1007/s10801-008-0128-1>
- [W03] W. Wang, *The Farahat-Higman ring of wreath products and Hilbert schemes* Adv. Math., 187 (2) (2004), pp. 417–446 URL: <https://doi.org/10.1016/j.aim.2003.09.003>
- [WW19] J. Wan and W. Wang, *Stability of the centers of group algebras of $GL_n(q)$* Adv. Math 349 (2019), pp. 749–780 URL: <https://doi.org/10.1016/j.aim.2019.04.026>