

Criteria for solubility and nilpotency of finite groups with automorphisms

Cristina Acciarri, Robert M. Guralnick, and Pavel Shumyatsky

ABSTRACT. Let G be a finite group admitting a coprime automorphism α . Let $J_G(\alpha)$ denote the set of all commutators $[x, \alpha]$, where x belongs to an α -invariant Sylow subgroup of G . We show that $[G, \alpha]$ is soluble or nilpotent if and only if any subgroup generated by a pair of elements of coprime orders from the set $J_G(\alpha)$ is soluble or nilpotent, respectively.

1. Introduction

In [4] Baumslag and Wiegold established the following sufficient condition for the nilpotency of a finite group G .

THEOREM 1.1. *Let G be a finite group in which $|ab| = |a||b|$, whenever the elements a, b have coprime orders. Then G is nilpotent.*

Here the symbol $|x|$ stands for the order of an element x in a group G . Obviously the condition above is also necessary for the nilpotency of G . We mention that there are several recent results related to the theorem of Baumslag and Wiegold (see for example [2, 3, 10, 12, 6]).

An automorphism α of a finite group G is said to be coprime if $(|G|, |\alpha|) = 1$. Following [1] denote by $I_G(\alpha)$ the set of commutators $g^{-1}g^\alpha$, where $g \in G$. Let $[G, \alpha]$ be the subgroup generated by $I_G(\alpha)$. In [1] the authors studied the impact of $I_G(\alpha)$ on the structure of $[G, \alpha]$.

Here we establish the following variation of the result of Baumslag and Wiegold.

2010 *Mathematics Subject Classification.* 20D45.

Key words and phrases. Finite groups, automorphisms.

The second author was partially supported by the NSF grant DMS-1901595 and a Simons Foundation Fellowship 609771. The third author was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Fundação de Apoio à Pesquisa do Distrito Federal (FAPDF), Brazil.

THEOREM 1.2. *Let G be a finite group admitting a coprime automorphism α . Then $[G, \alpha]$ is nilpotent if, and only if, $|xy| = |x||y|$ whenever x and y are elements of coprime prime power orders from $I_G(\alpha)$.*

At the start of this project, we did not know whether the hypothesis on the orders of elements in Theorem 1.2 is inherited by quotient groups. In order to overcome this issue we work instead with a somewhat different condition that behaves well with respect to forming quotients. Let $J_G(\alpha)$ denote the set of all commutators $[x, \alpha]$, where x belongs to an α -invariant Sylow subgroup of G . Observe that $J_G(\alpha)$ is a subset of $I_G(\alpha)$ and the elements of $J_G(\alpha)$ have prime power order. Moreover note that $J_G(\alpha)$ is a generating set for $[G, \alpha]$. Indeed this easily follows from [13, Lemma 2.4]. It turns out that properties of G are pretty much determined by those of subgroups generated by elements of coprime orders from $J_G(\alpha)$.

It is well known that if any pair of elements of a finite group generates a soluble (respectively nilpotent) subgroup, then the whole group is soluble (respectively nilpotent). One of the theorems established in [1] provides a variation of this for groups with automorphisms.

Let G be a finite group admitting a coprime automorphism α . If any pair of elements from $I_G(\alpha)$ generates a soluble subgroup, then $[G, \alpha]$ is soluble. If any pair of elements from $I_G(\alpha)$ generates a nilpotent subgroup, then $[G, \alpha]$ is nilpotent.

Here this will be extended as follows.

THEOREM 1.3. *Let G be a finite group admitting a coprime automorphism α . Then $[G, \alpha]$ is soluble if and only if any subgroup generated by a pair of elements of coprime orders from $J_G(\alpha)$ is soluble.*

The above theorem is used to establish the following related necessary and sufficient conditions for the nilpotency of $[G, \alpha]$.

THEOREM 1.4. *Let G be a finite group admitting a coprime automorphism α . Then the following statements are equivalent.*

- (i) *The subgroup $[G, \alpha]$ is nilpotent;*
- (ii) *Any subgroup generated by a pair of elements of coprime orders from $J_G(\alpha)$ is nilpotent;*
- (iii) *Any subgroup generated by a pair of elements of coprime orders from $J_G(\alpha)$ is abelian;*
- (iv) *If x and y are elements of coprime orders from $J_G(\alpha)$, then $|xy| = |x||y|$;*

- (v) If x and y are elements of coprime orders from $J_G(\alpha)$, then $\pi(xy) = \pi(x) \cup \pi(y)$.

Here, $\pi(g)$ denotes the set of prime divisors of the order of $g \in G$.

Note that Theorem 1.2 easily follows from Theorem 1.4. Furthermore, observe that the results provide criteria for solubility and nilpotency of finite groups admitting coprime groups of automorphisms: If G is acted on by a group A of coprime order, then $[G, A]$ is soluble or nilpotent if and only if any subgroup generated by a pair of elements of coprime orders from the set $J_G(A)$ is soluble or nilpotent, respectively. Here $J_G(A)$ stands for the set of all commutators $[x, a]$, where $a \in A$ and x belongs to an A -invariant Sylow subgroup of G . Indeed, this is straightforward using that $[G, A] = \prod_{a \in A} [G, a]$.

2. A condition for solubility of $[G, \alpha]$

All groups considered in this paper are finite. We start with a collection of well-known facts about coprime automorphisms of finite groups (see for example [8]).

LEMMA 2.1. *Let a group G admit a coprime automorphism α . The following conditions hold:*

- (i) $G = [G, \alpha]C_G(\alpha)$ and $|I_G(\alpha)| = [G : C_G(\alpha)]$;
- (ii) If N is any α -invariant normal subgroup of G , we have $C_{G/N}(\alpha) = C_G(\alpha)N/N$, $I_{G/N}(\alpha) = \{gN \mid g \in I_G(\alpha)\}$ and $J_{G/N}(\alpha) = \{gN \mid g \in J_G(\alpha)\}$;
- (iii) If N is any α -invariant normal subgroup of G such that $N = C_N(\alpha)$, then $[G, \alpha]$ centralizes N ;
- (iv) Any α -invariant p -subgroup of G is contained in an α -invariant Sylow p -subgroup.

The purpose in this section is to establish conditions that lead to the solubility of $[G, \alpha]$. Throughout, by a simple group we mean a nonabelian simple group. If a simple group G admits a coprime automorphism α of order e , then $G = L(q)$ is a group of Lie type and α is a field automorphism. Furthermore, $C_G(\alpha) = L(q_0)$ is a group of the same Lie type (and rank) defined over the subfield such that $q = q_0^e$ (see [9]).

In what follows, $q = p^s$ is a power of a prime p . For any positive integer n , we say that a prime r is a *primitive prime divisor* of $q^n - 1$ if r divides $q^n - 1$ and r does not divide $q^k - 1$ for any positive integer $k < n$. Primitive prime divisors of $q^n + 1$ are defined in a similar way. The following result of Zsigmondy ([18]) on the existence of primitive prime divisors will be used in the sequel.

THEOREM 2.2 ([18]). *Let $a > b > 0$, $\gcd(a, b) = 1$ and $n > 1$ be positive integers. Then*

- (i) $a^n - b^n$ has a prime divisor that does not divide $a^k - b^k$ for all positive integers $k < n$, unless $a = 2, b = 1$ and $n = 6$; or $a + b$ is a power of 2 and $n = 2$.
- (ii) $a^n + b^n$ has a prime divisor that does not divide $a^k + b^k$ for all positive integers $k < n$, with exception $2^3 + 1^3$.

Throughout, the term “semisimple group” means direct product of simple groups. We are now ready to deal with the proof of Theorem 1.3. For the reader’s convenience we restate it here.

THEOREM. *Let G be a group admitting a coprime automorphism α . Then $[G, \alpha]$ is soluble if and only if any subgroup generated by a pair of elements of coprime orders from $J_G(\alpha)$ is soluble.*

PROOF OF THEOREM 1.3. Suppose that this is false and let $G = [G, \alpha]$ be a counterexample of minimal order.

Note that without loss of generality we may assume that α has prime order, say e . Indeed, arguing by induction on the order of α assume that $[G, B]$ is soluble for any proper subgroup B of $\langle \alpha \rangle$. Since G is a counterexample of minimal order, it follows that G does not have soluble normal subgroups and so α must have prime order.

We wish to show that there are α -invariant subgroups P and Q of coprime prime power orders such that $[x, \alpha]$ and $[y, \alpha]$ generate a nonsoluble subgroup for some $x \in P$ and $y \in Q$.

Let M be a minimal α -invariant normal subgroup of G . Since G/M satisfies the hypothesis, by induction G/M is soluble. The subgroup M is either elementary abelian or semisimple. It is enough to consider the case where M is semisimple. Hence, we can assume that M is a direct product of isomorphic simple groups, say $M = S_1 \times \cdots \times S_k$, and α transitively permutes the simple factors. Moreover because of minimality $G = M$. Suppose first that $k \geq 2$.

It follows by [11, Theorem 1.2] that in S_1 there are two elements, say x and y , of coprime prime power orders generating a nonsoluble subgroup and such that $|xy|$ is coprime to both $|x|$ and $|y|$. Since α transitively permutes the simple factors of G , we deduce that both $\langle x, x^\alpha, \dots, x^{\alpha^i}, \dots \rangle$ and $\langle y, y^\alpha, \dots, y^{\alpha^i}, \dots \rangle$ are α -invariant abelian subgroups. Therefore both are contained in α -invariant Sylow subgroups. Write $[x, \alpha][y, \alpha] = x^{-1}x^\alpha y^{-1}y^\alpha = (yx)^{-1}(xy)^\alpha$ and observe that $|[x, \alpha][y, \alpha]|$ is coprime to both $|[x, \alpha]|$ and $|[y, \alpha]|$. The elements $[x, \alpha]$ and $[y, \alpha]$ generate a nonsoluble subgroup and we are done (an alternative proof of this case can be obtained using [7, Corollary F]).

Therefore we are reduced to the case where G is simple. It follows that $G = L(q)$ is a group of Lie type, say over the field of $q = p^s$ elements and α is a field automorphism of coprime order e . Moreover for any α -invariant subgroup H of G the subgroup $[H, \alpha]$ is soluble. Write $s = s_1 s_2$ where s_1 is a e -power and s_2 is coprime to e . Observe that α nontrivially acts on the subgroup $L(p^{s_1})$. Because of minimality we conclude that $q = p^{s_1}$. Thus, s is a e -power. Furthermore, the centralizer $C_G(\alpha)$ is the group of the same Lie type defined over the subfield of $q_0 = p^{s/e}$ elements.

Assume that G has rank at least 2. Note that α normalizes a Borel subgroup B and, by the structure of field automorphisms, α normalizes each parabolic subgroup of G containing B . We see that α normalizes (but does not centralize) a Levi subgroup L and so, by the minimality of G , the subgroup L should be soluble, a contradiction (since $q \geq 8$). This means that G must be of rank 1.

Hence suppose that G has (twisted) Lie rank 1. If $G = \text{PSU}_3(q)$ or ${}^2\text{G}_2(q)$, we observe that there is an α -invariant subgroup isomorphic to $\text{PSL}_2(q)$ which is not centralized by α and so we can reduce to the case $G = \text{PSL}_2(q)$. Thus, we only need to consider $G = \text{PSL}_2(q)$ with $q = p^s$ for s odd and $s \geq 5$ or $G = \text{Sz}(q)$ with $q = 2^s$ for odd $s > 1$.

If $G = \text{PSL}_2(q)$, take U to be an α -invariant Sylow p -subgroup and note that $[U, \alpha] \neq 1$. Note that any element in $[U, \alpha]$ is a commutator $[u, \alpha]$ with $u \in U$. Let now r be a primitive prime divisor of $q+1$, i.e. r does not divide $p^i + 1$ for $i < s$ (that always exists by Theorem 2.2), and let R be an α -invariant Sylow r -subgroup. Since r does not divide the order of $C_G(\alpha)$ (and of any subfield subgroup), we have $[R, \alpha] = R$. Let $1 \neq x \in [R, \alpha]$ and let $1 \neq y \in [U, \alpha]$. It follows that $G = \langle x, y \rangle$ since there is no proper subgroup of order divisible by pr (see [15, Theorem 6.25 in Chap. 3]), a contradiction.

It remains to consider the case $G = \text{Sz}(q)$, where $q = 2^s$ for odd $s > 1$. See [14] for properties of Suzuki groups. The order of G is $q^2(q-1)(q^2+1)$. Observe that the maximal subgroups of G are (up to conjugacy) a Borel subgroup of order $q^2(q-1)$, a dihedral subgroup of order $2(q-1)$, subfield subgroups, and two subgroups of the form $T.4$, where T is cyclic of order $q \pm l + 1$ with $l^2 = 2q$, i.e. of order $2^s \pm 2^{(s+1)/2} + 1$. Note that $(q+l+1)(q-l+1) = q^2 + 1$.

Let r be a primitive prime divisor of $q^2 + 1 = 2^{2s} + 1$. Let R be an α -invariant Sylow r -subgroup. Let t be a primitive prime divisor of $q - 1 = 2^s - 1$. Then t does not divide the order of any subfield subgroup and so also t does not divide $q^2 + 1$. By [14, Theorem 9], there is no proper subgroup of G whose order is divisible by rt . Let

S be an α -invariant Sylow t -subgroup. Neither of R and S intersects $C_G(\alpha)$, whence $[R, \alpha] = R$ and $[S, \alpha] = S$. Moreover any element in R or S is a commutator with α . It follows that G is generated by $[x, \alpha]$ and $[y, \alpha]$ with $x \in R$ and $y \in S$, a contradiction. This completes the proof. \square

As a by-product of the proof of Theorem 1.3 we obtain the following proposition.

PROPOSITION 2.3. *Let G be a finite group admitting a coprime automorphism α . Assume that $\pi(xy) = \pi(x) \cup \pi(y)$ whenever x and y are elements of coprime orders from $J_G(\alpha)$. Then $[G, \alpha]$ is soluble.*

The proof will be omitted since it is just an obvious modification of the argument used in Theorem 1.3. The proposition will be used in the proof of Theorem 1.4, which in particular shows that under the hypotheses of the proposition the subgroup $[G, \alpha]$ is nilpotent.

3. Criteria for nilpotency of $[G, \alpha]$

In this section we aim to establish Theorem 1.4 that gives necessary and sufficient conditions for the nilpotency of $[G, \alpha]$. For the reader's convenience we restate it here.

THEOREM. *Let G be a finite group admitting a coprime automorphism α . Then the following statements are equivalent.*

- (i) *The subgroup $[G, \alpha]$ is nilpotent;*
- (ii) *Any subgroup generated by a pair of elements of coprime orders from $J_G(\alpha)$ is nilpotent;*
- (iii) *Any subgroup generated by a pair of elements of coprime orders from $J_G(\alpha)$ is abelian;*
- (iv) *If x and y are elements of coprime orders from $J_G(\alpha)$, then $|xy| = |x||y|$;*
- (v) *If x and y are elements of coprime orders from $J_G(\alpha)$, then $\pi(xy) = \pi(x) \cup \pi(y)$.*

PROOF OF THEOREM 1.4. Note that obviously (i) implies (ii). Now if x and y are elements of coprime prime power orders from $J_G(\alpha)$ generating a nilpotent subgroup, then x and y commute each other and so (ii) implies (iii). If x and y are elements of coprime prime power orders from $J_G(\alpha)$ generating an abelian subgroup, then $|xy| = |x||y|$, and so (iii) implies (iv). Moreover it is immediate to see that (iv) implies (v).

We now prove the implication (v) \Rightarrow (i). Assume (v). In view of Proposition 2.3, the subgroup $[G, \alpha]$ is soluble. Suppose that the result is false and let $G = [G, \alpha]$ be a counterexample of minimal order. Let

M be a minimal α -invariant normal subgroup of G . Hence, M is an elementary abelian p -group, for some prime p , because G is soluble. Since G/M satisfies the hypothesis, by induction G/M is nilpotent. It follows that $G = MQ$, where Q is an α -invariant Sylow q -subgroup of G satisfying $Q = [Q, \alpha]$. Moreover, for any $x \in J_Q(\alpha)$ and $y \in J_M(\alpha)$, by hypothesis the order of xy is divisible by both p and q . Note that the element y is of the form $[m, \alpha]$, for suitable element $m \in M$ and so $y \in [M, \alpha]$. If $y \in [M, x]$, then xy is a q -element, contradicting the hypothesis. Hence we have $[M, x] \cap [M, \alpha] = 1$, whenever $x \in J_Q(\alpha)$.

Suppose that for a nontrivial element u there is $x \in J_Q(\alpha)$ such that $u \in C_M(x\alpha^{-1})$ while $u \notin C_M(\alpha)$. Then the element $[u, \alpha]$ is nontrivial and belongs to the intersection $[M, \alpha] \cap [M, x]$. This leads to a contradiction with the assumption above that $[M, x] \cap [M, \alpha] = 1$, whenever $x \in J_Q(\alpha)$.

Therefore

$$C_M(x\alpha^{-1}) \leq C_M(\alpha) \text{ whenever } x \in J_Q(\alpha). \quad (1)$$

Now, let g be an arbitrary element of Q and apply (1) with $x = [g, \alpha]$. Then $x\alpha^{-1} = \alpha^{-g}$ and we obtain that $C_M(\alpha)^g = C_M(\alpha)$. Since this holds for any $g \in Q$, we conclude that $C_M(\alpha)$ is normal in G . Lemma 2.1 (iii) now shows that $C_M(\alpha) \leq Z(G)$. Since G is a counterexample of minimal order, $G/Z(G)$ is nilpotent unless $Z(G) = 1$. Hence $Z(G) = 1$ and therefore $C_M(\alpha) = 1$. Thus, we have $M = [M, \alpha]$. Recall that $[M, x] \cap [M, \alpha] = 1$. It follows that $[M, x] = 1$ for any $x \in J_Q(\alpha)$ and so $M \leq Z(G)$, a contradiction. This proves that (v) implies (i) and thus the theorem is established. \square

References

- [1] C. Acciarri, R.M. Guralnick and P. Shumyatsky, *Coprime automorphisms of finite groups*, Trans. Amer. Math. Soc., 375:7 (2022), 4549–4565. <https://doi.org/10.1090/tran/8553>
- [2] R. Bastos and P. Shumyatsky, *A sufficient condition for nilpotency of the commutator subgroup*, Sib. Math. J. 57 (2016), 762–763.
- [3] R. Bastos, C. Monetta and P. Shumyatsky, *A criterion for metanilpotency of a finite group*, J. Group Theory 21 (2018), 713–718.
- [4] B. Baumslag and J. Wiegold, *A sufficient condition for nilpotency in a finite group*, arXiv:1411.2877v1 (2014).
- [5] R.W. Carter, *Simple Groups of Lie Type*, John Wiley & Sons, 1972.
- [6] J. Alves Da Silva and P. Shumyatsky, *On Nilpotency of Higher Commutator Subgroups of a Finite Soluble Group*, Arch. Math., 116:1 (2021), 1–6.
- [7] S. Dolfi, R.M. Guralnick, M. Herzog and C. Praeger, *A new solvability criterion for finite groups*, J. Lond. Math. Soc. (2) 85 (2012), 269–281.
- [8] D. Gorenstein, *Finite Groups*, Chelsea Publishing Company, New York, 1980.

- [9] D. Gorenstein, R. Lyons and R. Solomon, *The classification of the finite simple groups*. Number 3. Part I. Chapter A: Almost Simple K-groups, Mathematical Surveys and Monographs, vol. 40, AMS, Providence, RI, 1998.
- [10] R. M. Guralnick and A. Moretó, *Conjugacy classes, characters and products of elements*, Math. Nachr. 292 (2019), 1315–1320.
- [11] R. M. Guralnick and P. H. Tiep, *Lifting in Frattini covers and a characterization of finite solvable groups*, J. Reine Angew. Math. 708 (2015), 49–72.
- [12] A. Moretó and A. Sáez, *Prime divisors of orders of products*, Proc. Roy. Soc. Edinb. Sect. A 149 (2019), 1153–1162.
- [13] P. Shumyatsky, *Linear groups with almost right Engel elements*, Proc. Edinb. Math. Soc. (2) 62 (2019), no. 3, 789–797.
- [14] M. Suzuki, *On a Class of Doubly Transitive Groups*, Annals Math. 75 (1962), 105–145.
- [15] M. Suzuki, *Group Theory I*, Grundlehren der mathematischen Wissenschaften 247, Springer-Verlag, Berlin, 1982.
- [16] J. G. Thompson, *Nonsolvable finite groups all of whose local subgroups are solvable*, Bull. Amer. Math. Soc., 74 (1968), 383–437.
- [17] R. A. Wilson, *The Finite Simple Groups*, Graduate Texts in Mathematics, vol. 251. Springer-Verlag London, Ltd., London, 2009.
- [18] K. Zsigmondy, *Zur Theorie der Potenzreste*, Monatsh. für Math. u. Phys. 3 (1892), 265–284.

CRISTINA ACCIARRI: DIPARTIMENTO DI SCIENZE FISICHE, INFORMATICHE E MATEMATICHE, UNIVERSITÀ DEGLI STUDI DI MODENA E REGGIO EMILIA, VIA CAMPI 213/B, I-41125 MODENA, ITALY

Email address: `cristina.acciarri@unimore.it`

ROBERT M. GURALNICK: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA90089-2532, USA

Email address: `guralnic@usc.edu`

PAVEL SHUMYATSKY: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA, BRASILIA-DF, 70910-900 BRAZIL

Email address: `pavel@unb.br`