

# STABILITY OF A NEW GENERALIZED RECIPROCAL TYPE FUNCTIONAL EQUATION

I. SADANI

ABSTRACT. In this paper, we investigate the generalized Hyers-Ulam stability of the following reciprocal type functional equation

$$f(2x + y) + f(2x - y) = \frac{2f(x)f(y) \sum_{\substack{k=0 \\ k \text{ is even}}}^l 2^{l-k} \binom{l}{k} f(x)^{\frac{k}{l}} f(y)^{\frac{l-k}{l}}}{\left(4f(y)^{\frac{2}{l}} - f(x)^{\frac{2}{l}}\right)^l}$$

in non-zero real and non-Archimedean spaces.

## 1. INTRODUCTION

Since the birth of theory of stability, which was first proposed by Ulam [16] in the form of a question and after the first answer due to Hyers [7], many results have been obtained for many functional equations in different spaces. In this work we are interested in the case of reciprocal functional equations.

In 2010, Ravi and Senthil Kumar [9] studied the stability of the reciprocal type functional equation

$$(1.1) \quad f(x + y) = \frac{f(x)f(y)}{f(x) + f(y)},$$

where  $f : X \rightarrow \mathbb{R}$  is a mapping with  $X$  as the space of non-zero real numbers.

In 2014, Kim and Bodaghi [3] introduced and studied the stability of the quadratic reciprocal functional equation

$$(1.2) \quad f(2x + y) + f(2x - y) = \frac{2f(x)f(y)[4f(y) + f(x)]}{(4f(y) - f(x))^2}$$

In the same year, Ravi et al. [11] investigated the generalized Hyers-Ulam-Rassias stability of a reciprocal-quadratic functional equation of the form

$$(1.3) \quad r(x + 2y) + r(2x + y) = \frac{r(x)r(y)[5r(x) + 5r(y) + 8\sqrt{r(x)r(y)}]}{[2r(x) + 2r(y) + 5\sqrt{r(x)r(y)}]^2}$$

in intuitionistic fuzzy normed spaces.

In 2017, Kim et al. [15] introduced and studied the stability of the reciprocal-cubic functional equation

---

2010 *Mathematics Subject Classification.* 39B52, 39B72 .

*Key words and phrases.* Reciprocal Functional Equation, non-Archimedean space, Hyers-Ulam-Rassias stability.

$$(1.4) \quad c(2x+y) + c(x+2y) = \frac{9c(x)c(y)[c(x) + c(y) + 2c(x)^{\frac{1}{3}}c(y)^{\frac{1}{3}}(c(x)^{\frac{1}{3}} + c(y)^{\frac{1}{3}})]}{[2c(x)^{\frac{2}{3}} + 2c(y)^{\frac{2}{3}} + 5c(x)^{\frac{1}{3}}c(y)^{\frac{1}{3}}]^3}$$

and the reciprocal-quartic functional equation

$$(1.5) \quad q(2x+y) + q(2x-y) = \frac{2q(x)q(y)[q(x) + 16q(y) + 24\sqrt{q(x)q(y)}]}{[4\sqrt{q(y)} - \sqrt{q(x)}]^4}$$

in non-Archimedean fields.

Very recently, in [2], Kumar et al. focused on the stability of reciprocal-septic functional equation

$$(1.6) \quad S(2x+y) + S(2x-y) = \frac{2S(x)S(y)[128S(x) + 672S(x)^{\frac{2}{7}}S(y)^{\frac{5}{7}} + 280S(x)^{\frac{4}{7}}S(y)^{\frac{3}{7}} + 14S(x)^{\frac{6}{7}}S(y)^{\frac{1}{7}}]}{[4S(y)^{\frac{2}{7}} - S(x)^{\frac{2}{7}}]^7},$$

and the reciprocal-octic functional equation

$$(1.7) \quad \mathcal{O}(2x+y) + \mathcal{O}(2x-y) = \frac{2\mathcal{O}(x)\mathcal{O}(y)[\mathcal{O}(x) + 1792\mathcal{O}(x)^{\frac{1}{4}}\mathcal{O}(y)^{\frac{3}{4}} + 1120\mathcal{O}(x)^{\frac{1}{2}}\mathcal{O}(y)^{\frac{1}{2}} + 112\mathcal{O}(x)^{\frac{3}{4}}\mathcal{O}(y)^{\frac{1}{4}} + 256\mathcal{O}(y)]}{[4\mathcal{O}(y)^{\frac{1}{4}} - \mathcal{O}(x)^{\frac{1}{4}}]^8}.$$

The other results pertaining to the stability of different reciprocal-type functional equations can be found in [1, 12, 13, 14, 4, 10, 6, 5, 8].

In this paper, we introduce the following new reciprocal functional equation

$$(1.8) \quad f(2x+y) + f(2x-y) = \frac{2f(x)f(y) \sum_{\substack{k=0 \\ k \text{ is even}}}^l 2^{l-k} \binom{l}{k} f(x)^{\frac{k}{l}} f(y)^{\frac{l-k}{l}}}{(4f(y)^{\frac{2}{l}} - f(x)^{\frac{2}{l}})^l}$$

then, we investigate the general solution and its generalized Hyers-Ulam stability in the non-zero real numbers and in non-Archimedean fields.

## 2. GENERAL SOLUTION OF (1.8)

In this section, the solution of the functional equation (1.8) is given. Throughout this section, we assume that  $\mathbb{R}^*$  is the set of non-zero real numbers.

**Theorem 1.** *Let  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  be a continuous real-valued function of a two real variables satisfying the reciprocal type functional equation (1.8) and assume that  $f(x) \neq 0$  and  $4f(y)^{\frac{2}{l}} - f(x)^{\frac{2}{l}} \neq 0$  for all  $x, y \in \mathbb{R}^*$ . Then  $f$  is of the form*

$$f(x) = \frac{c}{x^l},$$

for all  $x \in \mathbb{R}^*$ , where  $c \neq 0$ .

*Proof.* Let  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  satisfies the functional equation (1.8). Replacing  $(x, y)$  by  $(x, x)$  in (1.8), we obtain

$$f(3x) + f(x) = \frac{2f(x) \sum_{\substack{k=0 \\ k \text{ is even}}}^l 2^{l-k} \binom{l}{k}}{3^l}$$

By induction, it is straightforward to show that

$$(2.1) \quad \sum_{\substack{k=0 \\ k \text{ is even}}}^l 2^{l-k} \binom{l}{k} = \frac{3^l + 1}{2}, l \geq 0.$$

Then,

$$f(3x) = \frac{1}{3} f(x)$$

for all  $x \in \mathbb{R}^*$ . Therefore  $f$  is a reciprocal function of the form  $\frac{c}{x}$ . This completes the proof.  $\square$

### 3. STABILITY OF (1.8) IN NON-ZERO REAL NUMBERS

For future use and convenience, we introduce the following difference operator  $\Lambda : \mathbb{R}^* \rightarrow \mathbb{R}$  as

$$\Lambda(x, y) = f(2x + y) + f(2x - y) - \frac{2f(x)f(y) \sum_{\substack{k=0 \\ k \text{ is even}}}^l 2^{l-k} \binom{l}{k} f(x)^{\frac{k}{l}} f(y)^{\frac{l-k}{l}}}{\left(4f(y)^{\frac{2}{l}} - f(x)^{\frac{2}{l}}\right)^l},$$

with  $l \geq 1$  and  $x, y \in \mathbb{R}^*$ .

**Theorem 2.** Let  $Q : \mathbb{R}^* \times \mathbb{R}^* \rightarrow \mathbb{R}$  a function satisfying

$$(3.1) \quad \sum_{s=0}^{\infty} \frac{1}{3^{ls}} Q\left(\frac{x}{3^{s+1}}, \frac{y}{3^{s+1}}\right) < \infty$$

for all  $x, y \in \mathbb{R}^*$ . If a function  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  satisfies functional inequality

$$(3.2) \quad |\Lambda(x, y)| \leq Q(x, y)$$

for all  $x, y \in \mathbb{R}^*$ , then there exists a unique reciprocal function  $g : \mathbb{R}^* \rightarrow \mathbb{R}$  which satisfies (1.8) and the inequality

$$(3.3) \quad |f(x) - g(x)| \leq \sum_{s=0}^{\infty} \frac{1}{3^{ls}} Q\left(\frac{x}{3^{s+1}}, \frac{x}{3^{s+1}}\right)$$

for all  $x \in \mathbb{R}^*$ .

*Proof.* We substitute  $(x, y)$  by  $(x, x)$  in (3.2) and using (2.1) we get

$$(3.4) \quad \left| f(3x) - \frac{f(x)}{3} \right| \leq Q(x, x)$$

for all  $x \in \mathbb{R}^*$ . Substituting  $x$  by  $\frac{x}{3}$  in (3.4), we obtain

$$(3.5) \quad \left| f(x) - \frac{1}{3} f\left(\frac{x}{3}\right) \right| \leq Q\left(\frac{x}{3}, \frac{x}{3}\right)$$

for all  $x \in \mathbb{R}^*$ . Now, by replacing  $x$  into  $\frac{x}{3}$  in (3.5), dividing by  $3^l$  and then adding the ensuing inequality with (3.5), we get

$$(3.6) \quad \left| f(x) - \frac{1}{3^{2l}} f\left(\frac{x}{3^2}\right) \right| \leq Q\left(\frac{x}{3}, \frac{x}{3}\right) + \frac{1}{3^l} Q\left(\frac{x}{3^2}, \frac{x}{3^2}\right)$$

for all  $x \in \mathbb{R}^*$ . In a similar way, proceeding further and applying induction procedure on a positive integer  $m$ , we get

$$(3.7) \quad \left| f(x) - \frac{1}{3^{ml}} f\left(\frac{x}{3^m}\right) \right| \leq \sum_{s=0}^{m-1} \frac{1}{3^{ls}} Q\left(\frac{x}{3^{s+1}}, \frac{x}{3^{s+1}}\right)$$

for all  $x \in \mathbb{R}^*$ .

Next, for any integers  $m', m$  with  $m' > m > 0$ , we obtain

$$(3.8) \quad \begin{aligned} \left| \frac{1}{3^{lm'}} f\left(\frac{x}{3^{m'}}\right) - \frac{1}{3^{lm}} f\left(\frac{x}{3^m}\right) \right| &= \left| \frac{1}{3^{lm'}} f\left(\frac{x}{3^{m'}}\right) - \frac{1}{3^{l(m'-1)}} f\left(\frac{x}{3^{l(m'-1)}}\right) + \dots \right. \\ &\quad \left. + \frac{1}{3^{l(m'-1)}} f\left(\frac{x}{3^{l(m'-1)}}\right) - \dots + \frac{1}{3^{l(m+1)}} f\left(\frac{x}{3^{m+1}}\right) - \frac{1}{3^{lm}} f\left(\frac{x}{3^m}\right) \right| \\ &\leq \frac{1}{3^{l(m'-1)}} Q\left(\frac{x}{3^{m'}}, \frac{x}{3^{m'}}\right) + \dots + \frac{1}{3^{lm}} Q\left(\frac{x}{3^{m+1}}, \frac{x}{3^{m+1}}\right) \\ &\leq \sum_{j=m}^{m'-1} \frac{1}{3^{lj}} Q\left(\frac{x}{3^{j+1}}, \frac{x}{3^{j+1}}\right) \end{aligned}$$

for all  $x \in \mathbb{R}^*$ . Taking  $m' \rightarrow \infty$  in (3.8) and applying (3.1), we observe that the sequence  $\{\frac{1}{3^{lm}} f(\frac{x}{3^m})\}$  is a Cauchy sequence for each  $x \in \mathbb{R}^*$ . Since  $\mathbb{R}$  is complete, we can define  $g : \mathbb{R}^* \rightarrow \mathbb{R}$  by  $g(x) = \lim_{m \rightarrow \infty} \frac{1}{3^{lm}} f(\frac{x}{3^m})$ . To prove that  $g$  satisfies (1.8), substituting  $(x, y)$  by  $(3^{-m}x, 3^{-m}y)$  in (3.2) and dividing by  $3^{lm}$ , we arrive

$$(3.9) \quad |3^{-lm} \Lambda(3^{-m}x, 3^{-m}y)| \leq 3^{-lm} Q(3^{-m}x, 3^{-m}y)$$

for all  $x, y \in \mathbb{R}^*$  and for all positive integer  $m$ . Letting  $m \rightarrow \infty$  in (3.9) and using (3.1), we see that  $g$  satisfies (1.8) for all  $x, y \in \mathbb{R}^*$ . Again, by taking  $m \rightarrow \infty$  in (3.7), we achieve (3.3). Now, it remains to show that  $g$  is unique. Let  $g' : \mathbb{R}^* \rightarrow \mathbb{R}$  be another reciprocal mapping which satisfies (1.8) and the inequality (3.3). Obviously, we have  $g'(3^{-m}x) = 3^{lm} g'(x)$ ,  $g(3^{-m}x) = 3^{lm} g(x)$  and using (3.3), we get

$$(3.10) \quad \begin{aligned} |g'(x) - g(x)| &= 3^{-lm} |g'(3^{-m}x) - g(3^{-m}x)| \\ &\leq 3^{-lm} (|g'(3^{-m}x) - f(3^{-m}x)| + |f(3^{-m}x) - g(3^{-m}x)|) \\ &\leq 2 \sum_{j=0}^{\infty} \frac{1}{3^{l(m+j)}} Q\left(\frac{x}{3^{m+j+1}}, \frac{x}{3^{m+j+1}}\right) \\ &\leq 2 \sum_{j=m}^{\infty} \frac{1}{3^{lj}} Q\left(\frac{x}{3^{j+1}}, \frac{x}{3^{j+1}}\right) \end{aligned}$$

for all  $x \in \mathbb{R}^*$ . Letting  $m \rightarrow \infty$  in (3.10), we find that  $g$  is unique. The proof of Theorem 2 is complete.  $\square$

The following corollaries are immediate consequences of Theorem 2.

**Corollary 1.** Let  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  be a mapping for which there exists  $\epsilon > 0$  such that

$$|\Lambda(x, y)| \leq \epsilon$$

holds for all  $x, y \in \mathbb{R}^*$ . Then,

$$g(x) = \lim_{m \rightarrow \infty} \frac{1}{3^{lm}} f\left(\frac{x}{3^m}\right)$$

for all  $x \in \mathbb{R}^*$ ,  $m \in \mathbb{N}$  and  $g : \mathbb{R}^* \rightarrow \mathbb{R}$  is the unique mapping satisfying the reciprocal functional equation (1.8) such that

$$|f(x) - g(x)| \leq \frac{3^l}{3^l - 1} \epsilon$$

for every  $x \in \mathbb{R}^*$ .

*Proof.* By taking  $Q(x, y) = \epsilon$  in Theorem 2 we arrive at the desired result.  $\square$

**Corollary 2.** Let  $\epsilon > 0$  and  $\alpha \neq -l$  be real numbers, and  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  be a mapping satisfying the functional inequality

$$|\Lambda(x, y)| \leq \epsilon(|x|^\alpha + |y|^\alpha)$$

for all  $x, y \in \mathbb{R}^*$ . Then, there exists a unique reciprocal mapping  $g : \mathbb{R}^* \rightarrow \mathbb{R}$  satisfying the functional equation (1.8) and

$$|f(x) - g(x)| \leq \frac{2 \cdot 3^l \epsilon}{3^{\alpha+l} - 1} |x|^\alpha$$

for all  $x \in \mathbb{R}^*$ .

*Proof.* By letting  $Q(x, y) = \epsilon(|x|^\alpha + |y|^\alpha)$  for all  $x, y \in \mathbb{R}^*$  in Theorem 2 we get the desired result.  $\square$

**Corollary 3.** Let  $\epsilon > 0$  and  $\alpha \neq -l$  be real numbers, and  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  be a mapping satisfying

$$|\Lambda(x, y)| \leq \epsilon(|x|^{\frac{\alpha}{2}} |y|^{\frac{\alpha}{2}} + |x|^\alpha + |y|^\alpha)$$

for all  $x, y \in \mathbb{R}^*$ . Then, there exists a unique reciprocal mapping  $g : \mathbb{R}^* \rightarrow \mathbb{R}$  satisfying the functional equation (1.8) and

$$|f(x) - g(x)| \leq \frac{\epsilon 3^{l+1}}{3^{\alpha+l} - 1} |x|^\alpha$$

for all  $x \in \mathbb{R}^*$ .

*Proof.* By taking  $Q(x, y) = \epsilon(|x|^{\frac{\alpha}{2}} |y|^{\frac{\alpha}{2}} + |x|^\alpha + |y|^\alpha)$  for all  $x, y \in \mathbb{R}^*$  in Theorem 2 we get the desired result.  $\square$

**Corollary 4.** Let  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  be a mapping and there exist  $p, q$  with  $p + q \neq -l$ . If there exists  $\epsilon \geq 0$  such that

$$|\Lambda(x, y)| \leq \epsilon |x|^p |y|^q$$

for all  $x, y \in \mathbb{R}^*$ , then there exists a unique reciprocal mapping  $g : \mathbb{R}^* \rightarrow \mathbb{R}$  satisfying the functional equation (1.8) and

$$|f(x) - g(x)| \leq \frac{3^l \epsilon}{3^{p+q+l} - 1} |x|^{p+q}$$

for all  $x \in \mathbb{R}^*$ .

*Proof.* Letting  $Q(x, y) = \epsilon|x|^p|y|^q$  for all  $x, y \in \mathbb{R}^*$  in Theorem 2, we obtain the required result.  $\square$

#### 4. STABILITY OF (1.8) IN NON-ARCHIMEDEAN FIELD

Throughout this section,  $\mathbb{A}$  and  $\mathbb{B}$  will denote a non-Archimedean field and a complete non-Archimedean field, respectively. Moreover, for a non-Archimedean field  $\mathbb{A}$ , we put  $\mathbb{A}^* = \mathbb{A} - \{0\}$ . We also assume that the reader is familiar with the basic properties of non-Archimedean fields.

**Theorem 3.** *Let  $G : \mathbb{A}^* \times \mathbb{A}^* \rightarrow [0, \infty[$  be a mapping such that*

$$(4.1) \quad \lim_{m \rightarrow \infty} \left| \frac{1}{3^l} \right|^m G \left( \frac{x}{3^{m+1}}, \frac{y}{3^{m+1}} \right) = 0$$

for all  $x, y \in \mathbb{A}^*$ . Suppose that  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  is a mapping satisfying the inequality

$$(4.2) \quad |\Lambda(x, y)| \leq G(x, y)$$

for all  $x, y \in \mathbb{A}^*$ . Then, there exists a unique reciprocal mapping  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  such that such that

$$(4.3) \quad |f(x) - g(x)| \leq \max \left\{ \left| \frac{1}{3^l} \right|^{k+1} G \left( \frac{x}{3^{k+1}}, \frac{x}{3^{k+1}} \right) : k \in \mathbb{N} \cup \{0\} \right\}$$

for all  $x \in \mathbb{A}^*$ .

*Proof.* Replacing  $(x, y)$  by  $(x, x)$  in (4.2), one finds

$$(4.4) \quad \left| f(x) - \frac{1}{3^l} f \left( \frac{x}{3} \right) \right| \leq |3^l| G \left( \frac{x}{3}, \frac{x}{3} \right)$$

for all  $x \in \mathbb{A}^*$ . Now, considering  $x$  as  $\frac{x}{3^m}$  in (4.4) and multiplying by  $\left| \frac{1}{3^l} \right|^m$ , we get

$$(4.5) \quad \left| \frac{1}{3^{lm}} f \left( \frac{x}{3^m} \right) - \frac{1}{3^{l(m+1)}} f \left( \frac{x}{3^{(m+1)}} \right) \right| \leq \left| \frac{1}{3^l} \right|^m G \left( \frac{x}{3^{m+1}}, \frac{x}{3^{m+1}} \right)$$

for all  $x \in \mathbb{A}^*$ . It is easy to obtain from the inequalities (4.1) and (4.5) that the sequence  $\left\{ \frac{1}{3^{lm}} f \left( \frac{x}{3^m} \right) \right\}$  is Cauchy. Since  $\mathbb{B}$  is complete, we can say that this sequence converges to a mapping  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  defined by

$$(4.6) \quad g(x) = \lim_{m \rightarrow \infty} \frac{1}{3^{lm}} f \left( \frac{x}{3^m} \right)$$

Besides, for each  $x \in \mathbb{A}^*$  and nonnegative integers  $m$ , we obtain

$$(4.7) \quad \begin{aligned} \left| \frac{1}{3^{lm}} f \left( \frac{x}{3^m} \right) - g(x) \right| &= \left| \sum_{k=0}^{m-1} \left[ \frac{1}{3^{l(k+1)}} f \left( \frac{x}{3^{(k+1)}} \right) - \frac{1}{3^{lm}} f \left( \frac{x}{3^m} \right) \right] \right| \\ &\leq \max \left\{ \left| \frac{1}{3^{l(k+1)}} \right| \left| f \left( \frac{x}{3^{(k+1)}} \right) - \frac{1}{3^{lm}} f \left( \frac{x}{3^m} \right) \right| : 0 \leq k < m \right\} \\ &\leq \max \left\{ \left| \frac{1}{3^l} \right|^m G \left( \frac{x}{3^{m+1}}, \frac{x}{3^{m+1}} \right) : 0 \leq k < m \right\}. \end{aligned}$$

Taking  $m \rightarrow \infty$  in the inequality (4.7) and using (4.6), we observe that the inequality (4.3) is valid. By the application of inequalities (4.1), (4.2) and (4.6), for all  $x, y \in \mathbb{A}^*$ , we arrive at

$$\begin{aligned} |\Lambda(x, y)| &= \lim_{m \rightarrow \infty} \left| \frac{1}{3^l} \right|^m \left| \Lambda \left( \frac{x}{3^m}, \frac{y}{3^m} \right) \right| \\ &\leq \lim_{m \rightarrow \infty} \left| \frac{1}{3^l} \right|^m G \left( \frac{x}{3^m}, \frac{y}{3^m} \right) \\ &= 0. \end{aligned}$$

Hence, the mapping  $g$  satisfies (4.1) and so it is reciprocal mapping. In order to confirm the uniqueness of  $g$ , let us presume  $g' : \mathbb{A}^* \rightarrow \mathbb{B}$  be another reciprocal mapping verifying (4.3). Then

$$\begin{aligned} |g(x) - g'(x)| &= \lim_{n \rightarrow \infty} \left| \frac{1}{3^l} \right|^n \left| g \left( \frac{x}{3^n} \right) - g' \left( \frac{x}{3^n} \right) \right| \\ &\leq \lim_{n \rightarrow \infty} \left| \frac{1}{3^l} \right|^n \max \left\{ \left| g \left( \frac{x}{3^n} \right) - f \left( \frac{x}{3^n} \right) \right|, \left| f \left( \frac{x}{3^n} \right) - g' \left( \frac{x}{3^n} \right) \right| \right\} \\ &\leq \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \max \left\{ \max \left\{ \left| \frac{1}{3^l} \right|^{k+n} G \left( \frac{x}{3^{k+n+1}}, \frac{x}{3^{k+n+1}} \right) \right. \right. \\ &\quad \left. \left. : n \leq k \leq m+n \right\} \right\} \\ &= 0 \end{aligned}$$

for all  $x \in \mathbb{A}^*$ . This implies that  $g$  is unique, which completes the proof.  $\square$

As a direct consequence of Theorem 3, we have the following corollaries.

**Corollary 5.** *Let  $\mu > 0$  be a constant. If  $f : \mathbb{A}^* \rightarrow \mathbb{B}$  satisfies*

$$|\Lambda(x, y)| \leq \mu$$

*for all  $x, y \in \mathbb{A}^*$ , then there exists a unique reciprocal mapping  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  satisfying (1.8) and*

$$|f(x) - g(x)| \leq \mu$$

*for all  $u \in \mathbb{A}^*$ .*

*Proof.* Taking  $G(x, y) = \mu$  in Theorem 3, we get the required result.  $\square$

**Corollary 6.** *Let  $\mu \geq 0$  and  $a \neq -l$ , be fixed constants. If  $f : \mathbb{A}^* \rightarrow \mathbb{B}$  satisfies*

$$|\Lambda(x, y)| \leq \mu(|x|^a + |y|^a)$$

*for all  $x, y \in \mathbb{A}^*$ , then there exists a unique reciprocal mapping  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  satisfying (1.8) and*

$$|f(x) - g(x)| \leq \begin{cases} \frac{2|\mu|}{|3|^a} |u|^a, & a > -l, \\ 2|\mu| |3|^l |u|^a, & a < -l, \end{cases}$$

*for all  $x \in \mathbb{A}^*$ .*

*Proof.* Considering  $G(x, y) = \mu(|x|^a + |y|^a)$  in Theorem 3, we obtain the desired result.  $\square$

**Corollary 7.** *Let  $f : \mathbb{A}^* \rightarrow \mathbb{B}$  be a mapping and let there exist real numbers  $p, q, a = p + q \neq -l$  and  $\mu \geq 0$  such that*

$$|\Lambda(x, y)| \leq \mu |u|^p |v|^q$$

for all  $x, y \in \mathbb{A}^*$ . Then, there exists a unique reciprocal mapping  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  satisfying (1.8) and

$$|f(x) - g(x)| \leq \begin{cases} \frac{\mu}{|3|^a} |u|^a, & a > -l, \\ \mu |3|^l |u|^a, & a < -l, \end{cases}$$

for all  $x \in \mathbb{A}^*$ .

*Proof.* Letting  $G(x, y) = \mu |x|^p |y|^q$ , for all  $x, y \in \mathbb{A}^*$  in Theorem 3, we acquire the requisite result.  $\square$

**Corollary 8.** *Let  $\mu \geq 0$  and  $a \neq -l$  be real numbers, and  $f : \mathbb{A}^* \rightarrow \mathbb{B}$  be a mapping satisfying the functional inequality*

$$|\Lambda(x, y)| \leq \mu (|x|^{\frac{a}{2}} |y|^{\frac{a}{2}} + |x|^a + |y|^a)$$

for all  $x, y \in \mathbb{A}^*$ . Then, there exists a unique reciprocal mapping  $g : \mathbb{A}^* \rightarrow \mathbb{B}$  satisfying (1.8) and

$$|f(x) - g(x)| \leq \begin{cases} \frac{|3|\mu}{|3|^a} |u|^a, & a > -l, \\ |3|\mu |3|^l |u|^a, & a < -l, \end{cases}$$

for all  $x \in \mathbb{A}^*$ .

*Proof.* Letting  $G(x, y) = \mu (|x|^{\frac{a}{2}} |y|^{\frac{a}{2}} + |x|^a + |y|^a)$  in Theorem 3, the result follows directly.  $\square$

*Proof.* Considering  $\mu(x, y) = \epsilon (|x|^a + |y|^a)$  in Theorem 3 the desired result follows directly.  $\square$

**Corollary 9.** *Let  $\alpha : [0, +\infty[ \rightarrow ]0, +\infty[$  be a function satisfying*

$$\alpha(|3|^{-1}t) \leq \alpha(|3|^{-1})\alpha(t) \quad (t \geq 0), \quad \left| \frac{1}{3^l} \right| \alpha(|3|^{-1}) < 1.$$

Let  $\delta > 0$  and  $f : \mathbb{A}^* \rightarrow \mathbb{B}$  be a mapping satisfying

$$(4.8) \quad |\Lambda(x, y)| \leq \delta (\alpha(|x|) + \alpha(|y|))$$

with  $x, y \in \mathbb{A}^*$ . Then there exists a unique reciprocal mapping  $h : \mathbb{A}^* \rightarrow \mathbb{B}$  such that

$$|f(x) - h(x)| \leq 2\delta \alpha \left( \left| \frac{x}{3} \right| \right) \quad (x \in \mathbb{A}^*).$$

*Proof.* Defining  $G : \mathbb{A}^* \times \mathbb{A}^* \rightarrow ]0, +\infty[$  by  $G(x, y) = \delta (\alpha(|x|) + \alpha(|y|))$ . On the one hand, we have



$$\begin{aligned} \lim_{m \rightarrow \infty} \left| \frac{1}{3^l} \right|^m G\left(\frac{x}{3^{m+1}}, \frac{y}{3^{m+1}}\right) &= \lim_{m \rightarrow \infty} \left| \frac{1}{3^l} \right|^m \delta\left(\alpha\left(\left|\frac{x}{3^{m+1}}\right|\right) + \alpha\left(\left|\frac{y}{3^{m+1}}\right|\right)\right) \\ &\leq \lim_{m \rightarrow \infty} \left| \frac{1}{3^l} \right|^m \alpha(|3|^{-1})^{m+1} G(x, y) \\ &= \lim_{m \rightarrow \infty} \left( \left| \frac{1}{3^l} \right| \alpha(|3|^{-1}) \right)^m \alpha(|3|^{-1}) G(x, y) \\ &= \lim_{m \rightarrow \infty} \left( \left| \frac{1}{3^l} \right| \alpha(|3|^{-1}) \right)^{m+1} \left| \frac{1}{3^l} \right|^{-1} G(x, y) \\ &= 0. \end{aligned}$$

for all  $x, y \in \mathbb{A}^*$ . On the other hand,

$$\lim_{m \rightarrow \infty} \max \left\{ \left| \frac{1}{3^l} \right|^k G\left(\frac{x}{3^{k+1}}, \frac{x}{3^{k+1}}\right); 0 \leq k < m \right\} = G\left(\frac{x}{3}, \frac{x}{3}\right) = 2\delta\alpha\left(\left|\frac{x}{3}\right|\right),$$

for all  $x \in \mathbb{A}^*$ , and

$$\begin{aligned} \lim_{j \rightarrow \infty} \lim_{m \rightarrow \infty} \max \left\{ \left| \frac{1}{3^l} \right|^k G\left(\frac{x}{3^{k+1}}, \frac{x}{3^{k+1}}\right), j \leq k < m + j \right\} &= \lim_{j \rightarrow \infty} \left| \frac{1}{3^l} \right|^j G\left(\frac{x}{3^{j+1}}, \frac{x}{3^{j+1}}\right) \\ &= 0. \end{aligned}$$

Applying Theorem 2, we get desired result. □

### 5. CONCLUSION

- (1) The advantage of this paper is that we do not need to prove the stability of quadratic (1.2), quartic (1.5), septic (1.6) and octic (1.7) reciprocal functional equations separately. Instead we can apply our main theorem to prove the stability of those functional equations simultaneously.
- (2) Similarly, we can give the generalized form of the functional equation (1.3) and (1.4) as follows

$$(5.1) \quad f(2x + y) + f(x + 2y) = \frac{f(x)f(y) \sum_{k=0}^n (2^{n-k} + 2^k) \binom{n}{k} f(x)^{\frac{k}{n}} f(y)^{\frac{n-k}{n}}}{\left(2f(x)^{\frac{2}{n}} + 5f(x)^{\frac{1}{n}} f(y)^{\frac{1}{n}} + 2f(y)^{\frac{2}{n}}\right)^n},$$

with  $n \geq 1$ . The reciprocal function  $f(x) = \frac{c}{x^n}$  is solution of (5.1). Then the generalized Ulam-Hyers stability can be obtained in the same way as above study or as [11] in various spaces.

### REFERENCES

- [1] A. Bodaghi, P. Narasimman, J.M. Rassias, K. Ravi, Ulam stability of the reciprocal functional equation in non-Archimedean fields, *Acta Math. Univ. Comen.* 85(1) (2016), 113-124.
- [2] A. Bodaghi, B. V. Senthil Kumar and A. B. Vakilabad, Various stabilities of reciprocal-septic and reciprocal-octic functional equations, *Asian-European Journal of Mathematics* 2150034 (19 pages) (2021).
- [3] A. Bodaghi, S. O. Kim, Approximation on the quadratic reciprocal functional equation, *J. Funct. Spaces Appl.*, (2014), Article ID 532463.
- [4] A. Bodaghi, Y. Ebrahimoost, On the stability of quadratic reciprocal functional equation in non-Archimedean fields, *Asian-Eur. J. Math.*, 9(1) (2016), Article ID 1650002.

- [5] A. Bodaghi, J. M. Rassias, C. Park, Fundamental stabilities of an alternative quadratic reciprocal functional equation in non-Archimedean fields, *Proc. Jangjeon Math. Soc.*, 18(3) (2015), 313-320.
- [6] B. V. Senthil Kumar, K. Ravi, J. M. Rassias, Solution and generalized Ulam-Hyers stability of a reciprocal type functional equation in non-Archimedean fields, *World Sci. News*, 31(2016), 71-81.
- [7] D. H. Hyers, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci., U.S.A.*, 27 (1941), 222-224.
- [8] I. Sadani, On the stability of  $f(2x+y) + f\left(\frac{x+y}{2}\right) = \frac{2f(x)f(y)}{f(x)+f(y)} + \frac{2f(x+y)f(y-x)}{3f(y-x)-f(x+y)}$ , *Tatra Mt. Math. Publ.*, 78 (2020), 1-10.
- [9] K. Ravi, B. V. Senthil Kumar, Ulam-Gavruta-Rassias stability of Rassias reciprocal functional equation, *Glob. J. Appl. Math. Math. Sci.*, 3 (2016), 57-79.
- [10] K. Ravi, J. M. Rassias and B. V. Senthil Kumar, Ulam stability of a generalized reciprocal type functional equation in non-Archimedean fields, *Arab. J. Math.* 4 (2015), 117-126.
- [11] K. Ravi, J. M. Rassias, B. V. Senthil Kumar, A. Bodaghi, Intuitionistic fuzzy stability of a reciprocal-quadratic functional equation, *Int. J. Appl. Sci. Math.*, 1(1) (2014), 9-14.
- [12] K. Ravi, J. M. Rassias, B.V. Senthil Kumar, Ulam stability of reciprocal difference and adjoint functional equations. *Aust. J. Math. Anal. Appl.*, 8(1) (2011), Article ID 13.
- [13] K. Ravi, J. M. Rassias, B. V. Senthil Kumar, Ulam stability of generalized reciprocal functional equation in several variables, *Int. J. Appl. Math. Stat.*, 19 (2010), Article ID D10.
- [14] K. Ravi, E. Thandapani, B. V. Senthil Kumar, Stability of reciprocal type functional equations, *Panam. Math. J.*, 21(1) (2011), 59-70.
- [15] S. O. Kim, B. V. Senthil Kumar, A. Bodaghi, Approximation on the reciprocal-cubic and reciprocal-quartic functional equations in non-Archimedean fields, *Adv. Differ. Equ.*, 77 (2017).
- [16] S. M. Ulam, *Problems in Modern Mathematics*, Science Editions, John Wiley and Sons, New York, NY, USA, (1964).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MOULoud MAMMERI, TIZI-OUZOU 15000, ALGERIA

Email address: sadani.idir@yahoo.fr