

States localized on a boundary of the time-dependent parity-breaking medium

Oleg O. Novikov^{*1} and Anna Zakharova^{†1,2}

¹Saint Petersburg State University, 7/9 Universitetskaya nab., St. Petersburg, Russia 199034

²Petersburg Nuclear Physics Institute named by B.P. Konstantinov of National Research Centre "Kurchatov Institute", Gatchina, 1, mkr. Orlova roshcha, 188300, Russia

September 8, 2022

Abstract

We consider the massive vector field propagating in the inhomogeneous parity-breaking medium, such as the dense hot hadronic matter with chiral imbalance. The transition between the regions with approximately constant values of the parity-breaking parameter allows for the states localized on such boundary to occur. The adiabatic change of the background introduces either decay or the amplification of the localized states.

1 Introduction

It is a well-known feature of the Standard model all the parities \mathcal{P} , \mathcal{C} , \mathcal{T} and \mathcal{CP} are non-conserved [1]. Many models of the new physics predict new sources of the \mathcal{CP} violation. This may affect the physics at the energies much lower than the masses of the new particles through the radiative loop corrections. Thus, the searches of the \mathcal{CP} -violating effects in the collider, atomic and molecular experiments [2–8] constitute an important strategy for the exploration of the physics beyond the Standard model.

One of the ways the \mathcal{CP} -symmetry may be violated is through the θ -term of the gluon field,

$$S = -\frac{\theta}{16\pi^2} \int d^4x \operatorname{Tr} [G_{\mu\nu} \tilde{G}^{\mu\nu}], \quad \tilde{G}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} G_{\alpha\beta} \quad (1)$$

When θ is constant this term happens to be a topological charge. Thus, it does not appear in the perturbative processes, however it changes the nonperturbative dynamics. One of the effects is

^{*}o.novikov@spbu.ru

[†]zakharova.annet@gmail.com

the appearance of the electric dipole moment (EDM) for the neutron [9]. Its measurement put very strong constraints on the value of θ which constitutes a so-called strong \mathcal{CP} -problem [10,11].

However, even if fundamentally θ is zero, the strong interaction dynamics may admit the local \mathcal{P} and \mathcal{CP} -violation [12–16]

Such an effect may occur when the large fluctuations of the gluon field produce metastable configurations with nonzero value of the axial topological charge that is tied through the anomaly to the quark axial charge. The emergence of the parity-violating regions may occur within the fireballs of very hot and very dense hadronic medium approaching a phase transition curve of the QCD phase diagram. Such conditions may be achieved in the heavy-ion collisions at NICA, STAR, PHOENIX, SPS and CBM experiments. The manifestations of the QCD local parity violation include the chiral magnetic effect in the peripheral collisions [17–24] and the excessive production of the dileptons in the central collisions [25–29]. The chiral imbalance environment may also produce exotic decays of mesons [30, 31].

The presence of the chiral imbalance background affects the propagation and interaction of the effective photon and meson particles. The finite temperature and axial chemical potential as well as the local nature of the fireball result in the Lorentz symmetry breaking. This is associated with the local violation of the \mathcal{CPT} invariance.

The fireball of chiral matter exists only for a short time. After the cooldown of the medium and disappearance of the local chiral imbalance the effective vacuum state transforms into a squeezed state of ordinary photons and mesons [32].

Similar situation may occur on different scales if the strong \mathcal{CP} -problem is solved with help of the axion particle. In that scenario, the constant θ is promoted to the condensate of the pseudoscalar axion field. The axion field must also have similar \mathcal{CP} -violating interaction with the electromagnetic field. Local variations of the condensate of the axion particles may play the role of the Dark matter. In the vicinity of the very dense stars the condensate may be especially high [33]. The passage of the photons through the spatial boundary of such an axion lump was investigated in [32]. Those results also give a rough idea on the similar passage of the vector mesons through the spatial boundary of the static hadronic medium.

However, the aforementioned papers considered only the very edge of the transition region where θ -parameter behaves linearly. This may be valid only for a very high energy particles that have short wavelengths, and, thus, insensitive to the large scale behavior. In contrast, particles with lower energies should be sensitive to the entire transition area. As will be demonstrated in this paper, the transition between two plateau regions may contain the localized states of the massive vector field. While the model we considered is simplified, this implies that the fireball of the chiral hadronic matter possesses the boundary currents. We also consider the influence of the temporal evolution of the medium which results in the amplification or the decay of the boundary states.

2 Vector field in the parity breaking medium

As in [32] we start with the massive electrodynamics model with local parity-violating term,

$$S = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{4} \theta(x) F_{\mu\nu} \tilde{F}^{\mu\nu} + \frac{1}{2} m^2 A^\mu A_\mu + A^\mu \partial_\mu B + \frac{1}{2} B^2 \right) \quad (2)$$

This model is related to the vector dominance model in the presence of the chiral imbalance as described in [34]. The auxiliary field B is introduced to employ the Stückelberg mechanism for self-consistency of the massive vector field theory [35,36]. The equations of motion for this field yield,

$$(\square + m^2)B = 0, \quad \partial_\mu A^\mu = B \quad (3)$$

where $\square = \partial_\mu \partial^\mu$. From that we can derive the condition,

$$(\square + m^2)(\partial_\mu A^\mu) = 0, \quad (4)$$

and omit B from our consideration. The equations of motion for A^μ take the form,

$$(\square + m^2)A^\mu = \epsilon^{\mu\nu\alpha\beta}(\partial_\nu \theta)(\partial_\alpha A_\beta) \quad (5)$$

We will assume that the boundary thickness of the chiral breaking matter region is much smaller than its size. Then we can approximate θ by a function only of t and z . In the components the equations take the form,

$$(\square + m^2)A^t = -(\partial_z \theta) \left[\partial_x A^y - \partial_y A^x \right], \quad (6)$$

$$(\square + m^2)A^z = (\partial_t \theta) \left[\partial_x A^y - \partial_y A^x \right], \quad (7)$$

$$(\square + m^2)A^x = (\partial_t \theta) \left[\partial_y A^z - \partial_z A^y \right] + (\partial_z \theta) \left[\partial_t A^y + \partial_y A^t \right], \quad (8)$$

$$(\square + m^2)A^y = -(\partial_t \theta) \left[\partial_x A^z - \partial_z A^x \right] - (\partial_z \theta) \left[\partial_t A^x + \partial_x A^t \right] \quad (9)$$

As the translational invariance in x and y directions is not broken it is useful to rewrite these equations in the momentum representation,

$$A^\mu(t, x, y, z) = \int d^2k A^\mu(t, z) \exp(ik_x x + ik_y y), \quad (10)$$

It is also convenient to use the circular components,

$$A_\pm = A^x \pm iA^y, \quad k_\pm = k_x \pm ik_y \quad (11)$$

Then the equations take the form,

$$(\square + m^2)A^t = -\frac{1}{2}(\partial_z \theta) \left[k_- A_+ - k_+ A_- \right], \quad (12)$$

$$(\square + m^2)A^z = \frac{1}{2}(\partial_t \theta) \left[k_- A_+ - k_+ A_- \right], \quad (13)$$

$$(\square + m^2)A_\pm = \pm k_\pm \left[(\partial_t \theta) A^z + (\partial_z \theta) A^t \right] \pm i \left[(\partial_t \theta) \partial_z A_\pm - (\partial_z \theta) \partial_t A_\pm \right] \quad (14)$$

For the simplicity we will consider the case of the momentum directed perpendicularly to the boundary i.e. $k_\pm = 0$ assuming that the results may serve as an approximation for sufficiently low momenta in the x and y directions. Then the A_\pm decouple both from the equations on A^t and A^z , and from the condition (4).

The (t, z) sector does not depend on θ and corresponds to the longitudinal polarization of the massive photon in the empty space. In contrast the equation for the transverse polarizations looks like,

$$(\partial_t^2 - \partial_z^2 + m^2)A_{\pm} = \pm i \left[(\partial_t \theta) \partial_z A_{\pm} - (\partial_z \theta) \partial_t A_{\pm} \right], \quad (15)$$

As A_- satisfies the same equation as A_+ but for the reversed time, we will consider only A_+ from now on.

3 Adiabatically changing parity-violating background

First, we will consider the equation (15) in the adiabatic regime, when θ is changing in time slowly compared to the frequency of the wave whereas its spatial variation is sufficiently high compared to the wavelentgth. Then we will assume that the solution behaves as,

$$A_+ = a(t, z) \exp \left(-i \int dt \omega(t) \right) \quad (16)$$

where the profile function a and the frequency $\omega(t)$ are slowly varying at the same rate as θ .

Then in the leading approximation the $i(\partial_t \theta)$ term may be neglected. In this approximation,

$$a \simeq a_0, \quad \omega \simeq \omega_0 \quad (17)$$

and the equation becomes,

$$-\partial_z^2 a_0 + \left[m^2 - \omega_0^2 + \omega_0 (\partial_z \theta) \right] a_0 = 0, \quad (18)$$

which looks like a Schrödinger equation though with the potential proportional to the spectral parameter ω_0 .

Consider the boundary between regions of approximately homogeneous θ . Let us consider the following shape of θ ,

$$\theta(t, z) = \bar{\theta}(t) + \Delta\theta(t) \cdot \tanh \left[\mu(t) (z - z_0(t)) \right], \quad \theta \xrightarrow{z \rightarrow -\infty} \bar{\theta} - \Delta\theta, \quad \theta \xrightarrow{z \rightarrow +\infty} \bar{\theta} + \Delta\theta, \quad (19)$$

where all the functions only slowly depend on time. For this shape of θ the potential becomes the Pöschl-Teller one [37],

$$\left[-\partial_z^2 + m^2 - \omega_0^2 + \frac{\omega_0 \mu \cdot \Delta\theta}{\cosh^2 \mu(z - z_0)} \right] a_0 = 0, \quad (20)$$

As at large z the potential becomes constant $-\varepsilon \equiv -(\omega_0^2 - m^2)$ the potential has the continuous spectrum of the wavelike solutions for $\varepsilon > 0$ i.e. $|\omega_0| > m$.

If $\varepsilon < 0$ i.e. in the band $|\omega_0| < m$ the bound states may exist. This is indeed true for the case of $\omega_0 \mu \cdot \Delta\theta < 0$ when the last term gives a potential well. Then, as well-known for the Pöschl-Teller potential the bound states correspond to,

$$\varepsilon_n = -\mu^2 (\lambda - n - 1)^2, \quad \lambda(\lambda - 1) = \left| \frac{\omega_0 \Delta\theta}{\mu} \right| \quad (21)$$

where n is integer from 0 to the largest integer $n_{max} \leq \lambda - 1$. As $\varepsilon = \omega_0^2 - m^2$ this turns into equation on λ ,

$$\lambda^4 - 2\lambda^3 + (1 + \Delta\theta^2)\lambda^2 - 2(n+1)\Delta\theta^2\lambda + \Delta\theta^2\left((n+1)^2 - \frac{m^2}{\mu^2}\right) = 0 \quad (22)$$

When this equation has a real solution $\lambda > 1$ the bound state exists. From the discussion above it is evident that $n_{max}(n_{max} + 1) \leq |m\Delta\theta/\mu|$.

The normalized $n = 0$ bound state takes the form,

$$a_{0,n=0} = \frac{C}{\cosh^{\lambda_0-1} \mu(z - z_0)}, \quad C^2 = \frac{\mu}{\sqrt{\pi}} \frac{\Gamma(\lambda_0 - \frac{1}{2})}{\Gamma(\lambda_0 - 1)}. \quad (23)$$




Figure 1: The spectrum of ω_0 for $m/\mu = 0.5$

The examples of the spectrum for (19) are presented on Fig. 1 and Fig. 2. One can see that the levels appear with growth of the $\Delta\theta$ in pairs of different ω_0 corresponding to the same n . The levels with different n may also intersect. Both of these phenomena must lead to the breakdown of the adiabatic approximation used in this paper.

4 Correction due to the $\partial_t\theta$ term

At small times we may approximate the influence of the $\partial_t\theta(t, z) \equiv v(t, z)$ term with the adiabatically changing perturbation that changes the frequencies and profile functions,

$$\omega = \omega_0 + \omega_1, \quad a = a_0 + a_1, \quad \omega_1, a_1 \sim |v| \quad (24)$$

The next order equation takes the form,

$$\left[\omega_0^2 - \partial_z^2 + m^2 + \omega(\partial_z\theta) \right] a_1 = iv(\partial_z a_0) - 2\omega_0\omega_1 a_0 - \omega_1(\partial_z\theta)a_0 \quad (25)$$




Figure 2: The spectrum of ω_0 for $m/\mu = 1.0$

Let a be one of the localized bound states we found in the preceding section. For simplicity we will choose it to be real (which can always be done in this dimensionality by the appropriate phase transformation). If we multiply this equation on a_0^* and integrate over z , l.h.s. vanishes thanks to the Hermiticity of the leading order operator and the equation (18). From that we derive,

$$\omega_1 = i \frac{\int_{-\infty}^{+\infty} dz v a_0 \partial_z a_0}{\int_{-\infty}^{+\infty} dz (2\omega_0 + \partial_z \theta) a_0^2} = -\frac{i}{2} \frac{\int_{-\infty}^{+\infty} dz (\partial_z v) a_0^2}{\int_{-\infty}^{+\infty} dz (2\omega_0 + \partial_z \theta) a_0^2} \quad (26)$$

From this it is evident that ω_1 is purely imaginary. This should not be a surprise as the perturbation is non-Hermitian,

$$(iv\partial_z)^\dagger = i\partial_z v = i(\partial_z v) + iv\partial_z \quad (27)$$

Therefore the $\partial_t \theta$ term leads to the decay or the amplification of the bound states that were stable in the leading approximation.

Suppose for simplicity that in (19) only $\bar{\theta}$ and $\Delta\theta$ depend on time. Then for the level $n = 0$ we get from (23),

$$\omega_{1,n=0} = -\frac{i}{2} \frac{\frac{\lambda-1}{2\lambda-1} \mu (\partial_t \Delta\theta)}{\omega_0 + \frac{\lambda-1}{2\lambda-1} \mu \Delta\theta} \quad (28)$$

5 Conclusions

We have demonstrated that the massive vector field on a parity-violating background may possess localized states on the boundary of the parity-violating medium. While greatly simplified, this model may imply that a similar statement is valid for the vector meson states in a chiral-breaking medium [34]. The time evolution of the background means that the localized states

are not exactly stationary. This makes possible for the boundary currents to appear in the transition regions between the domains with different values of the parity-breaking parameter.

Our treatment would be valid only as long as the adiabatic approximation holds. However, it may be expected to break during the evolution of the parity-breaking medium. E.g. the breakdown will happen if two levels become close to each other (either with avoided crossing or with actual crossing of the levels thanks to the non-unitary character of the evolution). Another interesting non-adiabatic process may be expected when the level disappears, leading to the annihilation of the boundary currents into the outgoing particles with potentially rich phenomenology. The adiabatic approximation also depends on the axial chemical potential within the fireball to be sufficiently small compared to the spatial variations at its boundary. We plan to explore these questions further within the generalized chiral perturbation theory [34, 38–40] in our future work.

Acknowledgements

We would like to thank Alexander Andrianov for the discussions and suggestions.

References

- [1] Matthew D Schwartz. *Quantum field theory and the standard model*. Cambridge University Press, 2014.
- [2] Shanzhen Chen, Yiming Li, Wenbin Qian, Yuehong Xie, Zhenwei Yang, Liming Zhang, and Yanxi Zhang. Heavy flavour physics and cp violation at lhcb: a ten-year review. *arXiv preprint arXiv:2111.14360*, 2021.
- [3] Iosif B Khriplovich and Steve K Lamoreaux. *CP violation without strangeness: electric dipole moments of particles, atoms, and molecules*. Springer Science & Business Media, 2012.
- [4] Vitaly Andreev, DG Ang, D DeMille, JM Doyle, G Gabrielse, J Haefner, NR Hutzler, Z Lasner, C Meisenhelder, BR O’Leary, et al. Improved limit on the electric dipole moment of the electron. *Nature*, 562(7727):355–360, 2018.
- [5] Anna Zakharova. Rotating and vibrating symmetric-top molecule raoch 3 in fundamental p, t-violation searches. *Physical Review A*, 105(3):032811, 2022. [Phys. Rev. A 105, 069902(E) (2022)].
- [6] Anna Zakharova and Alexander Petrov. \mathcal{P}, \mathcal{T} -odd effects for the RaOH molecule in the excited vibrational state. *Phys. Rev. A*, 103(3):032819, 2021.
- [7] Anna Zakharova, Igor Kurchavov, and Alexander Petrov. Rovibrational structure of the ytterbium monohydroxide molecule and the p, t-violation searches. *The Journal of Chemical Physics*, 155(16):164301, 2021.

- [8] Alexander Petrov and Anna Zakharova. Sensitivity of the yboh molecule to p t-odd effects in an external electric field. *Physical Review A*, 105(5):L050801, 2022.
- [9] P G Harris, C A Baker, K Green, P Iaydjiev, S Ivanov, D J R May, J M Pendlebury, D Shiers, K F Smith, M Van der Grinten, et al. New experimental limit on the electric dipole moment of the neutron. *Physical review letters*, 82(5):904, 1999.
- [10] Hai-Yang Cheng. The strong CP problem revisited. *Physics Reports*, 158(1):1–89, 1988.
- [11] Jihn E. Kim and Gianpaolo Carosi. Axions and the Strong CP Problem. *Rev. Mod. Phys.*, 82:557–602, 2010. [Rev.Mod.Phys. 91, 049902(E) (2019)].
- [12] Dmitri Kharzeev, R. D. Pisarski, and Michel H. G. Tytgat. Possibility of spontaneous parity violation in hot QCD. *Phys. Rev. Lett.*, 81:512–515, 1998.
- [13] K Buckley, T Fugleberg, and A Zhitnitsky. Can induced θ vacua be created in heavy-ion collisions? *Physical Review Letters*, 84(21):4814, 2000.
- [14] Dmitri Kharzeev. Parity violation in hot qcd: Why it can happen, and how to look for it. *Physics Letters B*, 633(2-3):260–264, 2006.
- [15] A. A. Andrianov and D. Espriu. On the possibility of P-violation at finite baryon-number densities. *Phys. Lett. B*, 663:450–455, 2008.
- [16] A. A. Andrianov, V. A. Andrianov, and D. Espriu. Spontaneous P-violation in QCD in extreme conditions. *Phys. Lett. B*, 678:416–421, 2009.
- [17] Dmitri E Kharzeev, Larry D McLerran, and Harmen J Warringa. The effects of topological charge change in heavy ion collisions: “event by event p and cp violation”. *Nuclear Physics A*, 803(3-4):227–253, 2008.
- [18] Kenji Fukushima, Dmitri E Kharzeev, and Harmen J Warringa. Chiral magnetic effect. *Physical Review D*, 78(7):074033, 2008.
- [19] P V Buividovich, Maxim N Chernodub, E V Luschevskaya, and M I Polikarpov. Numerical evidence of chiral magnetic effect in lattice gauge theory. *Physical Review D*, 80(5):054503, 2009.
- [20] BI Abelev, MM Aggarwal, Z Ahammed, AV Alakhverdyants, BD Anderson, D Arkhipkin, GS Averichev, J Balewski, O Barannikova, LS Barnby, et al. Azimuthal charged-particle correlations and possible local strong parity violation. *Physical review letters*, 103(25):251601, 2009.
- [21] Kenji Fukushima, Dmitri E Kharzeev, and Harmen J Warringa. Electric-current susceptibility and the chiral magnetic effect. *Nuclear Physics A*, 836(3-4):311–336, 2010.
- [22] Arata Yamamoto. Lattice study of the chiral magnetic effect in a chirally imbalanced matter. *Physical Review D*, 84(11):114504, 2011.

- [23] Dmitri E. Kharzeev. The Chiral Magnetic Effect and Anomaly-Induced Transport. *Prog. Part. Nucl. Phys.*, 75:133–151, 2014.
- [24] Md Rihan Haque, Alice Collaboration, et al. Measurements of the chiral magnetic effect in pb–pb collisions with alice. *Nuclear Physics A*, 982:543–546, 2019.
- [25] G Agakishiev, C Agodi, H Alvarez-Pol, A Balanda, R Bassini, G Bellia, D Belver, A Belyaev, A Blanco, M Böhmer, et al. Study of dielectron production in c+ c collisions at lagev. *Physics Letters B*, 663(1-2):43–48, 2008.
- [26] KO Lapidus and VM Emel’yanov. Low mass dilepton production in relativistic heavy ion collisions. *Physics of Particles and Nuclei*, 40(1):29–48, 2009.
- [27] Itzhak Tserruya. Electromagnetic probes. In R. Stock, editor, *Datasheet from Landolt-Börnstein - Group I Elementary Particles, Nuclei and Atoms · Volume 23: “Relativistic Heavy Ion Physics” in SpringerMaterials*. Springer-Verlag Berlin Heidelberg, Berlin Heidelberg, 2010.
- [28] AA Andrianov, VA Andrianov, D Espriu, and X Planells. Dilepton excess from local parity breaking in baryon matter. *Physics Letters B*, 710(1):230–235, 2012.
- [29] Alexander A Andrianov, Vladimir A Andrianov, Domènec Espriu, and Xumeu Planells. Analysis of dilepton angular distributions in a parity breaking medium. *Physical Review D*, 90(3):034024, 2014.
- [30] A. A. Andrianov, V. A. Andrianov, D. Espriu, A. E. Putilova, and A. V. Iakubovich. Decays of light mesons triggered by chiral chemical potential. *Acta Phys. Polon. Supp.*, 10:977, 2017.
- [31] A. A. Andrianov, V. A. Andrianov, D. Espriu, A. V. Iakubovich, and A. E. Putilova. Exotic meson decays in the environment with chiral imbalance. *EPJ Web Conf.*, 158:03012, 2017.
- [32] A. A. Andrianov, S. S. Kolevatov, and R. Soldati. Parity Breaking Medium and Squeeze Operators. *Phys. Rev. D*, 95(7):076020, 2017.
- [33] A. A. Andrianov, V. A. Andrianov, D. Espriu, and S. S. Kolevatov. Stellar matter with pseudoscalar condensates. *Eur. Phys. J. C*, 76(3):169, 2016.
- [34] Vladimir Kovalenko, Alexander Andrianov, and Vladimir Andrianov. Vector mesons spectrum in a medium with a chiral imbalance induced by the vacuum of fermions. *J. Phys. Conf. Ser.*, 1690(1):012097, 2020.
- [35] E. C. G. Stueckelberg. Theory of the radiation of photons of small arbitrary mass. *Helv. Phys. Acta*, 30:209–215, 1957.
- [36] I. A. Batalin and E. S. Fradkin. Operatorial Quantization of Dynamical Systems Subject to Second Class Constraints. *Nucl. Phys. B*, 279:514–528, 1987.
- [37] Siegfried Flügge. *Practical quantum mechanics*. Springer Science & Business Media, 2012.

- [38] A. A. Andrianov, V. A. Andrianov, and D. Espriu. Chiral perturbation theory vs. Linear Sigma Model in a chiral imbalance medium. *Particles*, 3(1):15–22, 2020.
- [39] V. A. Andrianov, A. A. Andrianov, and D. Espriu. The Chiral Medium in a Generalized Sigma Model and the Chiral Perturbation Theory. *Phys. Part. Nucl.*, 53(2):111–116, 2022.
- [40] Andrea Vioque-Rodríguez, Angel Gómez Nicola, and Domènec Espriu. Studying chiral imbalance using Chiral Perturbation Theory. *PoS*, PANIC2021:369, 2022.