

High-order approximation to generalized Caputo derivatives and generalized fractional advection-diffusion equations

Sarita Kumari^a, Rajesh K. Pandey^{a,*}, R. P. Agarwal^b

^a*Department of Mathematical Sciences, Indian Institute of Technology (BHU) Varanasi, Varanasi-221005, Uttar Pradesh, India.*

^b*Department of Mathematics, Texas A & M University-Kingsville, Kingsville, Texas, USA.*

Abstract

In this article, a high-order time-stepping scheme based on the cubic interpolation formula is considered to approximate the generalized Caputo fractional derivative (GCFD). Convergence order for this scheme is $(4 - \alpha)$, where α ($0 < \alpha < 1$) is the order of the GCFD. The local truncation error is also provided. Then, we adopt the developed scheme to establish a difference scheme for the solution of generalized fractional advection-diffusion equation with Dirichlet boundary conditions. Furthermore, we discuss about the stability and convergence of the difference scheme. Numerical examples are presented to examine the theoretical claims. The convergence order of the difference scheme is analyzed numerically, which is $(4 - \alpha)$ in time and second-order in space.

Keywords: Generalized Caputo fractional derivative, Generalized fractional advection-diffusion equation, difference scheme, stability, numerical solutions.

1. Introduction

This work include, the numerical solution of following generalized fractional advection-diffusion equation, which is studied in [1]

$$\left\{ \begin{array}{l} {}_0^C \mathcal{D}_{t, [\zeta(t), \omega(t)]}^\alpha U(x, t) = D \frac{\partial^2 U(x, t)}{\partial x^2} - A \frac{\partial U(x, t)}{\partial x} + g(x, t), \quad x \in \Omega, \quad t \in (0, T], \\ U(x, 0) = U_0(x), \quad x \in \bar{\Omega} = \Omega \cup \partial\Omega, \\ U(0, t) = \rho_1(t), \quad U(a, t) = \rho_2(t), \quad t \in (0, T], \end{array} \right. \quad (1.1)$$

where $\Omega = (0, a)$ is a bounded domain with boundary $\partial\Omega$, and the notation ${}_0^C \mathcal{D}_{t, [\zeta(t), \omega(t)]}^\alpha$ denotes the GCFD (defined in [2], and related references therein) with respect to t of order

*Corresponding author

Email addresses: saritakumari.rs.mat19@itbhu.ac.in (Sarita Kumari), rkpandey.mat@itbhu.ac.in (Rajesh K. Pandey), ravi.agarwal@tamuk.edu (R. P. Agarwal)

α is

$${}_0^C \mathcal{D}_{t;[\zeta(t),\omega(t)]}^\alpha U(t) = \frac{[\omega(t)]^{-1}}{\Gamma(1-\alpha)} \int_0^t \frac{[\omega(s)U(s)]'}{[\zeta(t)-\zeta(s)]^\alpha} ds, \quad 0 < \alpha < 1, \quad (1.2)$$

where $\partial_s = \frac{\partial}{\partial s}$, parameters $D > 0$ is the diffusivity, $A > 0$ is the advection constant and U is the solute concentration, g , U_0 , ρ_1 and ρ_2 are continuous functions on their respective domains with $U_0(0) = \rho_1(0)$ and $U_0(a) = \rho_2(0)$. Here scale, weight are sufficiently regular functions and our model (1.1) reduces to the diffusion problem when $A = 0$. Advection-diffusion equation is basically a transport problem that transport a passive scalar quantity in a fluid flow. Due to diffusion and advection, this model represents physical phenomenon of species concentration for mass transfer and temperature in heat transfer; for more details, we refer to [3, 4, 5], and [6, 7, 8, 9, 10] for further history and significance of advection-diffusion equation in physics, chemistry and biology.

Mostly used fractional derivatives in the problem formulation are the Riemann-Liouville and the Caputo derivatives [11]. In the year 2012, the generalizations of fractional integrals and derivatives were discussed by Agrawal [2]. Two functions, scale $\zeta(t)$ and weight $\omega(t)$ in one parameter, appear in the definition of the generalized fractional derivative of a function $U(t)$. If $\omega(t) = 1$ and $\zeta(t) = t$ then generalized fractional derivative reduces to the Riemann-Liouville (R-L) and the Caputo derivative, whereas if $\omega(t) = 1$, $\zeta(t) = \ln(t)$, and $\omega(t) = t^{\sigma\eta}$, $\zeta(t) = t^\sigma$ then it will convert to Hadamard [12], and modified Erdélyi-Kober fractional derivatives, respectively. In [13] studied the generalized form of R-L and the Hadamard fractional integrals, which is a special case of the Erdélyi-Kober generalized fractional derivative, and some properties of this operator. Atangana and Baleanu [14] discussed a new fractional derivative with non-singular kernel and used this derivative in the formation of fractional heat transfer model. So, we obtain different types of fractional derivatives for different choices of weight and scale functions. In generalized derivative, scale function $\zeta(t)$ manages the considered time domain, it can stretch or contract accordingly to capture the phenomena accurately over desired time range. The weight function $\omega(t)$ allows the events to be estimated differently at different time.

Over the last decade, many numerical methods were investigated to approximate the Caputo fractional derivative. For example, Mustapha [15] presented $L1$ approximation formula to solve fractional reaction-diffusion equation and second order error bound discussed on non-uniform time meshes. Alikhanov [16] constructed $L2 - 1_\sigma$ formula to approximate Caputo fractional time derivative and then used this derived scheme in solving time fractional diffusion equation with variable coefficients. Abu Arqub [17] considered reproducing kernel algorithm for approximate solution of the nonlinear time-fractional PDE's with initial and Robin boundary conditions. Li and Yan [18] discussed the idea of [19] (i.e. L_2 approximation formula for time discretization), also derived a new time discretization method with accuracy order $\mathcal{O}(\tau^{3-\alpha})$ and finite element method for spatial discretization. Cao et al. [20] presented a high-order approximation formula based on the cubic interpolation to approximate the Caputo derivative for the time fractional advection-diffusion equation. Xu and Agrawal [21] used the finite difference method (FDM) to approximate the GCFD for solving

the generalized fractional Burgers equation. Kumar et al. [22] presented $L1$ and $L2$ methods to approximate the generalized time fractional derivative which are defined as follows, respectively

$${}_0^C \mathcal{D}_{t;[\zeta(t),\omega(t)]}^\alpha U(t)|_{t=t_n} = \frac{[\omega(t)]^{-1}}{\Gamma(1-\alpha)} \sum_{l=1}^n \left(\frac{\omega_l U_l - \omega_{l-1} U_{l-1}}{\zeta_l - \zeta_{l-1}} \right) \int_{t_{l-1}}^{t_l} [\zeta(t) - \zeta(s)]^{-\alpha} (\partial_s \zeta(s)) ds + r_1^n, \quad (1.3)$$

$${}_0^C \mathcal{D}_{t;[\zeta(t),\omega(t)]}^\alpha U(t)|_{t=t_n} = \frac{[\omega(t)]^{-1}}{\Gamma(1-\alpha)} \sum_{l=1}^n \int_{t_{l-1}}^{t_l} [\zeta(t) - \zeta(s)]^{-\alpha} \partial_s (\Pi_l \omega(s) U(s)) ds + r_2^n, \quad (1.4)$$

where,

$$\Pi_l'(\omega(t)U(t)) = \zeta'(t) \left\{ \left[\frac{2\zeta(t) - \zeta_l - \zeta_{l-1}}{(\zeta_{l-2} - \zeta_{l-1})(\zeta_{l-2} - \zeta_{l-1})} \right] \omega_{l-2} U_{l-2} + \left[\frac{2\zeta(t) - \zeta_l - \zeta_{l-2}}{(\zeta_{l-1} - \zeta_{l-2})(\zeta_{l-1} - \zeta_l)} \right] \omega_{l-1} U_{l-1} + \left[\frac{2\zeta(t) - \zeta_{l-1} - \zeta_{l-2}}{(\zeta_l - \zeta_{l-2})(\zeta_l - \zeta_{l-1})} \right] \omega_l U_l \right\},$$

where, the domain $[0, T]$ was discretized into n equal subintervals i.e. $0 = t_0 < t_1 < \dots < t_N = T$ with step-size $\tau = \frac{T}{N}$, and errors $r_1^n = \mathcal{O}(\tau^{2-\alpha})$, $r_2^n = \mathcal{O}(\tau^{3-\alpha})$ were shown in [22]. In this work, we discuss the numerical scheme for GCFD with convergence rate $(4 - \alpha)$; for this accuracy we have to assume that $U'(t_0) = 0$, $U''(t_0) = 0$, $U'''(t_0) = 0$. This idea is discussed in [20] for Caputo derivative approximation, but the error bound was discussed directly.

Due to the non-local property of the fractional derivatives, the numerical solution of the fractional partial differential equations is a very difficult task [23]. Several authors have presented some precise and efficient numerical methods for fractional advection-diffusion equation. For examples, Zheng et al. [24] used finite element method (FEM) for space fractional advection-diffusion equation. Mardani et al. [25] discussed meshless moving least square method for solving the time-fractional advection-diffusion equation with variable coefficients. Cao et al. [20] proposed the higher-order approximation of the Caputo derivatives and further applied it in solving the fractional advection-diffusion equation. They used the Lagrange interpolation method to discretize time derivative and second-order central difference for the spatial derivatives. Li and Cai [26] considered a three steps process for the Caputo fractional derivative approximation, first two steps include the shifted Lubich formula derivation for infinite interval then for finite interval, and after that it is generalized to the Caputo derivative. Yadav et al. [1] discussed Taylor expansion for the approximation of the generalized time-fractional derivative to solve generalized fractional advection-diffusion equation. Tian et al. [27] presented polynomial spectral collocation method for space fractional advection-diffusion equation. In [28], authors developed explicit and implicit Euler approximations to solve variable-order fractional advection-diffusion equation on finite domain. Singh et al. [29] investigated the numerical approximation of Caputo-Prabhakar derivative and then used this approximation to solve fractional advection-diffusion equation.

Up to now, there Xu and Agrawal [21] considered the FDM for approximation of the GCFD for the generalized fractional Burgers equation. In [22], authors presented numerical scheme for the generalized fractional telegraph equation in time. Cao et al. [30] worked on the generalized time-fractional Kdv equation. Xu et al. [31] considered the solution of generalized fractional diffusion equation. Owolabi and Pindza [32] used generalized Caputo-type fractional derivative for the numerical simulation of nonlinear chaotic maps. Odibat and Baleanu [33] presented an adaptive predictor corrector algorithm to solve initial value problems with generalized Caputo fractional derivative. The main focus of this paper is to present much higher order numerical scheme to approximate GCFD as compared to [34] and [1], and also to establish the error analysis in both time and space-discretization. To the best of our knowledge, no work has been done yet for a third-order error bound of cubic interpolation formula to approximate the GCFD.

The main contributions of this work are as follows:

- (1) we extend the approximation method of Cao et al. [20] for approximating the GCFD and obtain the convergence order $(4 - \alpha)$. Further, we show that the obtained scheme reduces to the approximation scheme discussed by Cao et al. [20] for choice of the scale and the weight functions as $\zeta(t) = t$ and $\omega(t) = 1$.
- (2) we establish the full error analysis of the presented higher-order numerical scheme for the generalized time fractional derivative by using Lagrange interpolation formula.
- (3) we introduce some numerical results for different choices of scale and weight functions for the high-order time discretization scheme with convergence order $\mathcal{O}(\tau^{4-\alpha})$ for all $\alpha \in (0, 1)$ which achieves higher accuracy than the numerical methods developed in Gao et al. [35] for $\zeta(t) = t$ and $\omega(t) = 1$.

The remaining sections of the paper is arranged as follows: In Section 2, we discuss the $(4 - \alpha)$ -th order scheme to approximate the GCFD of order α of the function U . In Section 3, a higher order difference scheme to solve generalized fractional advection-diffusion equation is presented. Stability and convergence analysis are also discussed in this section. We present three numerical examples which illustrate the error and convergence order of our established numerical scheme in Section 4. Finally, Section 5, concludes some remarks.

2. Numerical scheme for the generalized Caputo fractional derivative

Motivated by the research carried out in [20, 22], this section is devoted to presenting a high-order approximation formula for the generalized Caputo-type fractional derivative using cubic interpolation polynomials.

Suppose that $U(t) \in C^4[0, T]$, and grid points $0 = t_0 < t_1 < \dots < t_N = T$ with step length $\tau = t_n - t_{n-1}$ for $1 \leq n \leq N$. For simplicity, we use $g(s) = \omega(s)U(s)$, $U(t_i) = U_i$, $\omega(t_i) = \omega_i$, and $\zeta(t_i) = \zeta_i$. The generalized Caputo fractional derivative of order α of the function $U(t)$ at grid point t_n is given by,

$${}_0^C \mathcal{D}_{t;[\zeta(t), \omega(t)]}^\alpha U_n = \frac{[\omega_n]^{-1}}{\Gamma(1 - \alpha)} \sum_{l=1}^n \int_{t_{l-1}}^{t_l} \frac{g'(s)}{[\zeta(t_n) - \zeta(s)]^\alpha} ds. \quad (2.1)$$

On the first interval $[0, t_1]$ of domain, we use continuous linear polynomial $\Pi_1 g(t)$ to approximate the function $g(t) (= \omega(t)U(t))$. Let $g(t_l) = g_l$ and the difference operator $\nabla_\tau g_l = g_l - g_{l-1}$ for $l \geq 1$. Then, we have

$$(\Pi_1 g(t))' = \frac{(g_1 - g_0)}{\tau} = \frac{(\nabla_\tau g_1)}{\tau}.$$

Thus, Eq. (2.1) yields,

$$\begin{aligned} \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \int_0^{t_1} [\zeta(t_n) - \zeta(s)]^{-\alpha} g'(s) ds &= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \frac{(\nabla_\tau g_1)}{\tau} \int_0^{t_1} [\zeta(t_n) - \zeta(s)]^{-\alpha} ds + E_\tau^1 \\ &= a_{n-1}(\nabla_\tau g_1) + E_\tau^1, \end{aligned} \quad (2.2)$$

where E_τ^1 is the truncation error on first interval and coefficients for this approximation are

$$a_{n-l} = \frac{[\omega_n]^{-1}}{\Gamma(2-\alpha)} \left\{ \frac{[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} - [\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right\}.$$

Here, we denote notation $\alpha_0^{(n)} = \frac{[\omega_n]^{-1}}{\Gamma(2-\alpha)}$, $1 \leq n \leq N$. Then

$$a_{n-l} = \alpha_0^{(n)} \left\{ \frac{[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} - [\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right\}, \quad 1 \leq l \leq n. \quad (2.3)$$

Remark 1. If the scale function ζ is a positive strictly increasing functions on the domain $[0, T]$, then following inequality holds

$$0 < [\zeta(t_n) - \zeta(t_l)]^{1-\alpha} - [\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha}, \quad 1 \leq l \leq n. \quad (2.4)$$

Since, $t_l > t_{l-1}$, then it implies $\zeta(t_l) > \zeta(t_{l-1})$ for $1 \leq l \leq n$, also $(1-\alpha) > 0$.

Remark 2. To estimate a_{n-1} , suppose that

$$\Theta = [\zeta(t_n) - \zeta(s)] \Rightarrow d\Theta = -\zeta'(s) ds = -\left(\frac{\zeta(t_l) - \zeta(t_{l-1})}{\tau} \right) ds, \quad s \in (t_{l-1}, t_l),$$

therefore,

$$\int [\zeta(t_n) - \zeta(s)]^{-\alpha} ds = \frac{-\tau}{\zeta(t_l) - \zeta(t_{l-1})} \frac{[\zeta(t_n) - \zeta(s)]^{1-\alpha}}{(1-\alpha)}.$$

On the second interval $[t_1, t_2]$, we use continuous quadratic polynomial $\Pi_2 g(t)$ to approximate the function $g(t)$, then we get

$$(\Pi_2 g(t))' = \frac{(2t - t_0 - t_1)}{2\tau^2} g_2 - \frac{(2t - t_0 - t_2)}{\tau^2} g_1 + \frac{(2t - t_1 - t_2)}{2\tau^2} g_0.$$

Thus from Eq. (2.1), we get,

$$\begin{aligned}
\frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \int_{t_1}^{t_2} [\zeta(t_n) - \zeta(s)]^{-\alpha} g'(s) ds &= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \int_{t_1}^{t_2} [\zeta(t_n) - \zeta(s)]^{-\alpha} (\Pi_2 g(s))' ds + E_\tau^2 \\
&= a_{n-2}(g_2 - g_1) + b_{n-2}(g_2 - 2g_1 + g_0) + E_\tau^2 \\
&= a_{n-2}(\nabla_\tau g_2) + b_{n-2}(\nabla_\tau^2 g_2) + E_\tau^2. \tag{2.5}
\end{aligned}$$

Here, truncation error on second interval is E_τ^2 , and

$$\begin{aligned}
b_{n-l} = \alpha_0^{(n)} \left\{ \frac{1}{(2-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{2-\alpha} - [\zeta(t_n) - \zeta(t_l)]^{2-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^2} \right] \right. \\
\left. - \frac{1}{2} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} + [\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right] \right\}, \quad 2 \leq l \leq n. \tag{2.6}
\end{aligned}$$

On the other subdomains ($l \geq 3$), we use cubic interpolation polynomial $\Pi_l g(t)$ to approximate the function $g(t)$ using four points $(t_{l-3}, g_{l-3}), (t_{l-2}, g_{l-2}), (t_{l-1}, g_{l-1}), (t_l, g_l)$.

As we know cubic interpolation polynomial is defined as,

$$\Pi_l g(t) = \sum_{r=0}^3 g_{l-r} \prod_{s=0, s \neq r}^3 \left(\frac{t - t_{l-s}}{t_{l-r} - t_{l-s}} \right),$$

then we get,

$$\begin{aligned}
(\Pi_l g(t))' &= g_{l-3} \frac{(t - t_{l-2})(t_l + t_{l-1} - 2t) + (t - t_{l-1})(t_l - t)}{6\tau^3} \\
&+ g_{l-2} \frac{(t - t_{l-3})(2t - t_{l-1} - t_l) + (t - t_l)(t - t_{l-1})}{2\tau^3} \\
&+ g_{l-1} \frac{(t - t_{l-2})(t_{l-3} + t_l - 2t) + (t - t_{l-3})(t_l - t)}{2\tau^3} \\
&+ g_l \frac{(t - t_{l-2})(2t - t_{l-3} - t_{l-1}) + (t - t_{l-1})(t - t_{l-3})}{6\tau^3}.
\end{aligned}$$

Therefore, from Eq. (2.1), we have,

$$\begin{aligned}
& \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \sum_{l=3}^n \int_{t_{l-1}}^{t_l} [\zeta(t_n) - \zeta(s)]^{-\alpha} g'(s) ds \\
&= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \sum_{l=3}^n \int_{t_{l-1}}^{t_l} [\zeta(t_n) - \zeta(s)]^{-\alpha} (\Pi_l g(s))' ds + E_\tau^n \\
&= \sum_{l=3}^n \left[A_{1,n-l} g_l + A_{2,n-l} g_{l-1} + A_{3,n-l} g_{l-2} + A_{4,n-l} g_{l-3} \right] + E_\tau^n, \tag{2.7}
\end{aligned}$$

where E_τ^n ($3 \leq n \leq N$) is the truncation error, and

$$\begin{aligned}
A_{1,n-l} = & \alpha_0^{(n)} \left\{ \frac{1}{6} \left[\frac{2[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} - 11[\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right] \right. \\
& + \frac{1}{(2-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{2-\alpha} - 2[\zeta(t_n) - \zeta(t_l)]^{2-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^2} \right] \\
& \left. + \frac{1}{(2-\alpha)(3-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{3-\alpha} - [\zeta(t_n) - \zeta(t_l)]^{3-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^3} \right] \right\}, \tag{2.8}
\end{aligned}$$

$$\begin{aligned}
A_{2,n-l} = & \alpha_0^{(n)} \left\{ \frac{1}{2} \left[\frac{6[\zeta(t_n) - \zeta(t_l)]^{1-\alpha} + [\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right] \right. \\
& + \frac{1}{(2-\alpha)} \left[\frac{5[\zeta(t_n) - \zeta(t_l)]^{2-\alpha} - 2[\zeta(t_n) - \zeta(t_{l-1})]^{2-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^2} \right] \\
& \left. + \frac{3}{(2-\alpha)(3-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_l)]^{3-\alpha} - [\zeta(t_n) - \zeta(t_{l-1})]^{3-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^3} \right] \right\}, \tag{2.9}
\end{aligned}$$

$$\begin{aligned}
A_{3,n-l} = & \alpha_0^{(n)} \left\{ -\frac{1}{2} \left[\frac{2[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} + 3[\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right] \right. \\
& + \frac{1}{(2-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{2-\alpha} - 4[\zeta(t_n) - \zeta(t_l)]^{2-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^2} \right] \\
& \left. + \frac{3}{(2-\alpha)(3-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{3-\alpha} - [\zeta(t_n) - \zeta(t_l)]^{3-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^3} \right] \right\}, \tag{2.10}
\end{aligned}$$

$$\begin{aligned}
A_{4,n-l} = & \alpha_0^{(n)} \left\{ \frac{1}{6} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} + 2[\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right] \right. \\
& + \frac{1}{(2-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_l)]^{2-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^2} \right] \\
& \left. + \frac{1}{(2-\alpha)(3-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_l)]^{3-\alpha} - [\zeta(t_n) - \zeta(t_{l-1})]^{3-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^3} \right] \right\}, \quad (2.11)
\end{aligned}$$

where, $3 \leq l \leq n$. After simplifying the Eq. (2.7), we obtain the following form

$$\begin{aligned}
& \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \sum_{l=3}^n \int_{t_{l-1}}^{t_l} [\zeta(t_n) - \zeta(s)]^{-\alpha} g'(s) ds \\
& = \sum_{l=3}^n [a_{n-l}(g_l - g_{l-1}) + b_{n-l}(g_l - 2g_{l-1} + g_{l-2}) \\
& \quad + c_{n-l}(g_l - 3g_{l-1} + 3g_{l-2} - g_{l-3})] + E_\tau^n \\
& = \sum_{l=3}^n [a_{n-l}(\nabla_\tau g_l) + b_{n-l}(\nabla_\tau^2 g_l) + c_{n-l}(\nabla_\tau^3 g_l)] + E_\tau^n. \quad (2.12)
\end{aligned}$$

Here, we introduce another coefficient c_{n-l} , which is defined as

$$\begin{aligned}
c_{n-l} = & \alpha_0^{(n)} \left\{ \frac{1}{(2-\alpha)(3-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{3-\alpha} - [\zeta(t_n) - \zeta(t_l)]^{3-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^3} \right] \right. \\
& - \frac{1}{(2-\alpha)} \left[\frac{[\zeta(t_n) - \zeta(t_l)]^{2-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]^2} \right] \\
& \left. - \frac{1}{6} \left[\frac{[\zeta(t_n) - \zeta(t_{l-1})]^{1-\alpha} + 2[\zeta(t_n) - \zeta(t_l)]^{1-\alpha}}{[\zeta(t_l) - \zeta(t_{l-1})]} \right] \right\}, \quad 3 \leq l \leq n, \quad (2.13)
\end{aligned}$$

and coefficients a_{n-l} , b_{n-l} are defined in Eqs. (2.3), (2.6), respectively and Eq. (2.12) gives more compact form of Eq. (2.7). Such forms of coefficients were missing in [20]. From this we can easily discuss properties of coefficients.

Motivated by [20] (developed for Caputo derivative), a new numerical scheme for the generalized Caputo-type fractional derivative of order α of the function $U(t)$ at grid point t_n

with the help of the equations (2.2), (2.5), (2.7), is defined by

$$\begin{aligned}
{}^{\mathcal{H}}\mathcal{D}_{t;[\zeta(t),\omega(t)]}^{\alpha}U(t)|_{t=t_n} &= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \int_0^{t_n} \frac{g'(s)}{[\zeta(t_n) - \zeta(s)]^{\alpha}} ds \\
&= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \left[\int_0^{t_1} \frac{g'(s)}{[\zeta(t_n) - \zeta(s)]^{\alpha}} ds + \int_{t_1}^{t_2} \frac{g'(s)}{[\zeta(t_n) - \zeta(s)]^{\alpha}} ds \right. \\
&\quad \left. + \sum_{l=3}^n \int_{l-1}^{t_l} \frac{g'(s)}{[\zeta(t_n) - \zeta(s)]^{\alpha}} ds \right] \\
&= \sum_{l=0}^n \lambda_l \omega_{n-l} U_{n-l}, \tag{2.14}
\end{aligned}$$

with $g(s) = \omega(s)U(s)$.

Lemma 2.1. For any $\alpha \in (0, 1)$ and $U(t) \in \mathcal{C}^4[0, T]$, then

$${}^{\mathcal{C}}\mathcal{D}_{t;[\zeta(t),\omega(t)]}^{\alpha}U_n = {}^{\mathcal{H}}\mathcal{D}_{t;[\zeta(t),\omega(t)]}^{\alpha}U_n + E_{\tau}^n, \quad n = 1, 2, \dots, N,$$

where, ${}^{\mathcal{H}}\mathcal{D}_{t;[\zeta(t),\omega(t)]}^{\alpha}$ is the approximation of GCFD and $|E_{\tau}^n| = \mathcal{O}(\tau^{4-\alpha})$.

For distinct value of n , the coefficients in (2.14) can be expressed as below,
For $n = 1$,

$$\begin{cases} \lambda_0 = a_0, \\ \lambda_1 = -a_0. \end{cases}$$

For $n = 2$,

$$\begin{cases} \lambda_0 = a_0 + b_0, \\ \lambda_1 = a_1 - a_0 - 2b_0, \\ \lambda_2 = -a_1 + b_0. \end{cases}$$

For $n = 3$,

$$\begin{cases} \lambda_0 = A_{1,0}, \\ \lambda_1 = A_{2,0} + a_1 + b_1, \\ \lambda_2 = A_{3,0} + a_2 - a_1 - 2b_1, \\ \lambda_3 = A_{4,0} - a_2 + b_1. \end{cases}$$

For $n = 4$,

$$\begin{cases} \lambda_0 = A_{1,0}, \\ \lambda_1 = A_{1,1} + A_{2,0}, \\ \lambda_2 = A_{2,1} + A_{3,0} + a_2 + b_2, \\ \lambda_3 = A_{3,1} + A_{4,0} + a_3 - a_2 - 2b_2, \\ \lambda_4 = A_{4,1} - a_3 + b_2. \end{cases}$$

For $n = 5$,

$$\begin{cases} \lambda_0 = A_{1,0}, \\ \lambda_1 = A_{1,1} + A_{2,0}, \\ \lambda_2 = A_{1,2} + A_{2,1} + A_{3,0}, \\ \lambda_3 = A_{2,2} + A_{3,1} + A_{4,0} + a_3 + b_3, \\ \lambda_4 = A_{3,2} + A_{4,1} + a_4 - a_3 - 2b_3, \\ \lambda_5 = A_{4,2} - a_4 + b_3. \end{cases}$$

For $n \geq 6$,

$$\begin{cases} \lambda_0 = A_{1,0}, \\ \lambda_1 = A_{1,1} + A_{2,0}, \\ \lambda_2 = A_{1,2} + A_{2,1} + A_{3,0}, \\ \lambda_l = A_{1,l} + A_{2,l-1} + A_{3,l-2} + A_{4,l-3} \quad (3 \leq l \leq n-3), \\ \lambda_{n-2} = a_{n-2} + b_{n-2} + A_{2,n-3} + A_{3,n-4} + A_{4,n-5}, \\ \lambda_{n-1} = a_{n-1} - a_{n-2} - 2b_{n-2} + A_{3,n-3} + A_{4,n-4}, \\ \lambda_n = -a_{n-1} + b_{n-2} + A_{4,n-3}. \end{cases} \quad (2.15)$$

Lemma 2.2. *If the scale function ζ fulfills (2.4), and the weight function ω is non-negative and non-decreasing on uniform time grids, then*

(i)[30] *Linear approximation coefficient satisfies, $1 \leq l \leq n$,*

$$0 < a_{n-1} < \dots < a_{n-l} < a_{n-l-1} < \dots < a_1 < a_0.$$

(ii) *Quadratic approximation coefficient satisfies, $2 \leq l \leq n$,*

$$0 < b_{n-2} < \dots < b_{n-l} < b_{n-l-1} < \dots < b_1 < b_0.$$

Proof. (i) If scale function ζ is continuous on respective domain then by using mean-value theorem, there exist $\hat{x}_l \in (t_{l-1}, t_l)$ such that

$$\frac{1}{\tau} \int_{t_{l-1}}^{t_l} [\zeta(t_n) - \zeta(s)]^{-\alpha} ds = [\zeta(t_n) - \zeta(\hat{x}_l)]^{-\alpha}, \quad 1 \leq l \leq n. \quad (2.16)$$

Since, $[\zeta(t_n) - \zeta(s)]^{-\alpha}$ is a monotone increasing function, we easily get our required result.

(ii) Let $\eta(s) = [\zeta(t_n) - \zeta(s)]^{1-\alpha}$, suppose scale function ζ is sufficiently smooth on domain $[0, T]$ then mean-value theorem yields

$$\begin{aligned}
2 \int_{t_{l-1}}^{t_l} \eta(s) ds - (\eta(t_{l-1}) + \eta(t_l)) &= 2\eta(\tilde{x}_l) - (\eta(t_{l-1}) + \eta(t_l)), \quad \tilde{x}_l \in (t_{l-1}, t_l), \\
&= -\theta(\eta'(t_l) - \eta'(t_{l-1})), \quad 0 < \theta < 1 \\
&= -\theta\tau\eta''(\nu_l), \quad \nu_l \in (t_{l-1}, t_l), \\
&= \frac{\theta\alpha(1-\alpha)}{\tau} [\zeta(t_l) - \zeta(t_{l-1})]^2 [\zeta(t_n) - \zeta(\nu_l)]^{-\alpha-1} > 0.
\end{aligned} \tag{2.17}$$

Using (2.17), we can easily get that $b_{n-l} > 0$ for positive strictly increasing weight function $\omega(t)$. Since, $[\zeta(t_n) - \zeta(s)]^{-\alpha-1}$ is a monotone increasing function on temporal domain $[0, T]$, so we get the desired result. \square

Lemma 2.3. *Suppose that the scale function ζ is positive and strictly increasing, and the weight function ω is non-negative and non-decreasing, then for each $\alpha \in (0, 1)$, the following conditions hold for coefficients in (2.15)*

- (1) $\lambda_0 > 0, \quad \forall n \geq 1,$
- (2) $\sum_{l=0}^n \lambda_l = 0.$

Proof. (1) If $n = 1$, then

$$\lambda_0 = a_0 = \alpha_0^{(n)} [\zeta_1 - \zeta_0]^{-\alpha},$$

as scale function is strictly increasing, therefore $\zeta_{n-1} < \zeta_n$, for $n \geq 1$. Which implies that $\lambda_0 = a_0 > 0$.

If $n = 2$, then $\lambda_0 = a_0 + b_0$,
since,

$$b_0 = \alpha_0^{(n)} \left\{ \left(\frac{1}{2-\alpha} - \frac{1}{2} \right) [\zeta_2 - \zeta_1]^{-\alpha} \right\} > 0,$$

and we have already shown that $a_0 > 0$, therefore $\lambda_0 = a_0 + b_0 > 0$.

If $n \geq 3$, then for $0 < \alpha < 1$,

$$\lambda_0 = A_{1,0} = \alpha_0^{(n)} \left\{ \left(\frac{1}{3} + \frac{1}{(2-\alpha)} + \frac{1}{(2-\alpha)(3-\alpha)} \right) [\zeta_n - \zeta_{n-1}]^{-\alpha} \right\} > 0.$$

(2) If $n = 1$, then $\lambda_0 = -\lambda_1 = a_0 > 0$, for $0 < \alpha < 1$. It implies that $\lambda_0 + \lambda_1 = 0$.

If $n = 2$, then there exist a $\alpha \in (0, 1)$, by numerical analysis

$$\lambda_0 + \lambda_1 + \lambda_2 = (a_0 + b_0) + (a_1 - a_0 + 2b_0) + (-a_1 + b_0) = 0.$$

If $n \geq 3$, then

$$\begin{aligned}
\sum_{l=0}^n \lambda_l &= A_{1,0} + A_{1,1} + A_{2,0} + A_{1,2} + A_{2,1} + A_{3,0} \\
&+ \sum_{l=3}^{n-3} (A_{1,l} + A_{2,l-1} + A_{3,l-2} + A_{4,l-3}) + a_{n-2} + b_{n-2} \\
&+ A_{2,n-3} + A_{3,n-4} + A_{4,n-5} + a_{n-1} - a_{n-2} - 2b_{n-2} + A_{3,n-3} \\
&+ A_{4,n-4} - a_{n-1} + b_{n-2} + A_{4,n-3} \\
&= \sum_{j=0}^{n-3} (A_{1,j} + A_{2,j} + A_{3,j} + A_{4,j}) = 0.
\end{aligned}$$

□

Lemma 2.4. *If scale function ζ is a Lipschitz function on interval $[t_{l-1}, t_l]$ with Lipschitz constant L , then*

$$|\zeta_l - \zeta_{l-1}| \leq L\tau, \quad 1 \leq l \leq n.$$

2.1. Truncation error for generalized Caputo derivative term

For truncation error of approximation of the generalized Caputo derivative defined in (2.14), for simplicity, suppose that function $g(t) = \omega(t)U(t)$ such that $g(t) \in C^4((0, T])$, and $\zeta(t) = v$ this implies $t = \zeta^{-1}(v)$. Therefore, $g(v) = \omega(\zeta^{-1}(v))U(\zeta^{-1}(v))$.

Theorem 1. *A triangle inequality gives the bound*

$$|{}_0^C \mathcal{D}_{t;[\zeta(t), \omega(t)]}^\alpha g_n - {}^H \mathcal{D}_{t;[\zeta(t), \omega(t)]}^\alpha g_n| \leq \sum_{n=1}^N |E_\tau^n|,$$

with

$$E_\tau^n = \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \int_{\zeta_{l-1}}^{\zeta_l} [\zeta_n - v]^{-\alpha} [g(v) - \Pi_l g(v)]' dv, \quad 1 \leq l \leq n.$$

$$\begin{aligned}
(1) \quad |E_\tau^1| &\leq \frac{\alpha[\omega_n]^{-1}}{\Gamma(1-\alpha)} \left[\frac{1}{8\alpha} + \frac{1}{2(1-\alpha)(2-\alpha)} \right] \max_{t_0 \leq v \leq t_1} |g^{(2)}(v)| L^{2-\alpha}(\tau)^{2-\alpha}, \quad n=1, \\
(2) \quad |E_\tau^2| &\leq \frac{\alpha[\omega_n]^{-1}}{\Gamma(1-\alpha)} \left\{ \frac{1}{12} \max_{t_0 \leq v \leq t_1} |g^{(2)}(v)| (t_2 - t_1)^{-\alpha-1} L^{2-\alpha}(\tau)^3 + \left[\frac{1}{12} \right. \right. \\
&\quad \left. \left. + \frac{1}{3(1-\alpha)(2-\alpha)} \left(\frac{1}{2} + \frac{1}{(3-\alpha)} \right) \right] \max_{t_0 \leq v \leq t_2} |g^{(3)}(v)| L^{3-\alpha}(\tau)^{3-\alpha} \right\}, \quad n=2, \\
(3) \quad |E_\tau^n| &\leq \frac{\alpha[\omega_n]^{-1}}{\Gamma(1-\alpha)} \left\{ \frac{1}{72} \max_{t_0 \leq x \leq t_2} |g^{(3)}(x)| (t_n - t_2)^{-\alpha-1} L^{3-\alpha}(\tau)^4 + \left[\frac{3}{128\alpha} \right. \right. \\
&\quad \left. \left. + \frac{1}{12(1-\alpha)(2-\alpha)} \left(1 + \frac{3}{(3-\alpha)} + \frac{3}{(3-\alpha)(4-\alpha)} \right) \right] \max_{t_0 \leq x_1 \leq t_n} |g^{(4)}(x_1)| L^{4-\alpha}(\tau)^{4-\alpha} \right\}, \\
&\hspace{25em} n \geq 3.
\end{aligned}$$

Proof. (1)[22] For $n=1$, the scheme (2.14) is linear approximation of GCFD, and convergence order for this approximation formula is $\mathcal{O}(\tau^{2-\alpha})$.

(2)[22] For $n=2$, the scheme (2.14) is quadratic approximation of GCFD, and here convergence order of approximation formula is $\mathcal{O}(\tau^{3-\alpha})$.

(3) For $n \geq 3$, By Lagrange interpolation remainder theorem, we use quadratic interpolation function $\Pi_2 g(v)$ to interpolate $g(v)$ using node points (ζ_0, g_0) , (ζ_1, g_1) , (ζ_2, g_2) on interval $[\zeta_0, \zeta_2]$ and cubic interpolation function $\Pi_l g(v)$ depends on (ζ_{l-3}, g_{l-3}) , (ζ_{l-2}, g_{l-2}) , (ζ_{l-1}, g_{l-1}) , (ζ_l, g_l) to interpolate $g(v)$ on $[\zeta_{l-3}, \zeta_l]$ as follows

$$g(v) - \Pi_2 g(v) = \frac{g^{(3)}(\eta_1)}{3!} (v - \zeta_0)(v - \zeta_1)(v - \zeta_2), \quad v \in [\zeta_0, \zeta_2], \quad \eta_1 \in (\zeta_0, \zeta_2), \quad (2.18)$$

$$g(v) - \Pi_l g(v) = \frac{g^{(4)}(\eta_l)}{4!} (v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l), \quad v \in [\zeta_{l-3}, \zeta_l], \quad \eta_l \in (\zeta_{l-3}, \zeta_l), \quad (2.19)$$

where $3 \leq l \leq n$.

Now,

$$\begin{aligned}
E_\tau^n &= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \left[\int_{\zeta_0}^{\zeta_2} [g(v) - \Pi_2 g(v)]' [\zeta_n - v]^{-\alpha} dv \right. \\
&\quad \left. + \sum_{l=3}^n \int_{\zeta_{l-1}}^{\zeta_l} [g(v) - \Pi_l g(v)]' [\zeta_n - v]^{-\alpha} dv \right] \\
&= \frac{[\omega_n]^{-1}}{\Gamma(1-\alpha)} \left\{ [g(v) - \Pi_2 g(v)] [\zeta_n - v]^{-\alpha} \Big|_{\zeta_0}^{\zeta_2} \right. \\
&\quad - \alpha \int_{\zeta_0}^{\zeta_2} [g(v) - \Pi_2 g(v)] [\zeta_n - v]^{-\alpha-1} dv \\
&\quad + \sum_{l=3}^n \left[[g(v) - \Pi_l g(v)] [\zeta_n - v]^{-\alpha} \Big|_{\zeta_{l-1}}^{\zeta_l} \right. \\
&\quad \left. - \alpha \int_{\zeta_{l-1}}^{\zeta_l} [g(v) - \Pi_l g(v)] [\zeta_n - v]^{-\alpha-1} dv \right] \Big\} \\
&= -\frac{\alpha [\omega_n]^{-1}}{\Gamma(1-\alpha)} \left\{ \int_{\zeta_0}^{\zeta_2} [g(v) - \Pi_2 g(v)] [\zeta_n - v]^{-\alpha-1} dv \right. \\
&\quad \left. + \sum_{l=3}^n \int_{\zeta_{l-1}}^{\zeta_l} [g(v) - \Pi_l g(v)] [\zeta_n - v]^{-\alpha-1} dv \right\} \tag{2.20}
\end{aligned}$$

Since, from (2.18) and (2.19)

$$\begin{aligned}
[g(v) - \Pi_2 g(v)] [\zeta_n - v]^{-\alpha} \Big|_{\zeta_0}^{\zeta_2} &= \frac{g^{(3)}(\eta_1)}{3!} (v - \zeta_0)(v - \zeta_1)(v - \zeta_2) [\zeta_n - v]^{-\alpha} \Big|_{\zeta_0}^{\zeta_2} = 0, \\
[g(v) - \Pi_l g(v)] [\zeta_n - v]^{-\alpha} \Big|_{\zeta_{l-1}}^{\zeta_l} &= \frac{g^{(4)}(\eta_l)}{4!} (v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l) \Big|_{\zeta_{l-1}}^{\zeta_l} = 0.
\end{aligned}$$

Consider the first integration of Eq. (2.20)

$$\begin{aligned}
&\left| \int_{\zeta_0}^{\zeta_2} [g(v) - \Pi_2 g(v)] [\zeta_n - v]^{-\alpha-1} dv \right| \\
&= \left| \int_{\zeta_0}^{\zeta_2} \frac{g^{(3)}(\eta_1)}{3!} (v - \zeta_0)(v - \zeta_1)(v - \zeta_2) [\zeta_n - v]^{-\alpha-1} dv \right| \\
&\leq \frac{1}{6} \max_{\zeta_0 \leq \eta_1 \leq \zeta_2} |g^{(3)}(\eta_1)| (\zeta_n - \zeta_2)^{-\alpha-1} \frac{(\zeta_0 - \zeta_2)^3 (\zeta_0 - 2\zeta_1 + \zeta_2)}{12} \\
&\leq \frac{1}{6} \max_{\zeta_0 \leq \eta_1 \leq \zeta_2} |g^{(3)}(\eta_1)| (\zeta_n - \zeta_2)^{-\alpha-1} \frac{(\zeta_0 - \zeta_2)^3 (\zeta_0 - \zeta_1)}{12}.
\end{aligned}$$

Using Lemma (2.4), we have following inequality

$$\left| \int_{\zeta_0}^{\zeta_2} [g(v) - \Pi_2 g(v)] [\zeta_n - v]^{-\alpha-1} dv \right| \leq \frac{1}{72} \max_{t_0 \leq x \leq t_2} |g^{(3)}(x)| (t_n - t_2)^{-\alpha-1} L^{3-\alpha} (\tau)^4. \quad (2.21)$$

Consider the second integration of Eq. (2.20)

$$\begin{aligned} & \left| \sum_{l=3}^n \int_{\zeta_{l-1}}^{\zeta_l} [g(v) - \Pi_l g(v)] [\zeta_n - v]^{-\alpha-1} dv \right| \\ & \leq \left| \sum_{l=3}^{n-1} \int_{\zeta_{l-1}}^{\zeta_l} \frac{g^{(4)}(\eta_l)}{4!} (v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l) [\zeta_n - v]^{-\alpha-1} dv \right| \\ & \quad + \left| \int_{\zeta_{n-1}}^{\zeta_n} \frac{g^{(4)}(\eta_n)}{4!} (v - \zeta_{n-3})(v - \zeta_{n-2})(v - \zeta_{n-1})(v - \zeta_n) [\zeta_n - v]^{-\alpha-1} dv \right|. \end{aligned} \quad (2.22)$$

Consider first part of RHS of Eq. (2.22)

$$\begin{aligned} & \left| \sum_{l=3}^{n-1} \int_{\zeta_{l-1}}^{\zeta_l} \frac{g^{(4)}(\eta_l)}{4!} (v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l) [\zeta_n - v]^{-\alpha-1} dv \right| \\ & \leq \frac{1}{24} \max_{\zeta_2 \leq \eta_2 \leq \zeta_{n-1}} |g^{(4)}(\eta_2)| \tilde{f}(\zeta_{l-3}, \zeta_{l-2}, \zeta_{l-1}, \zeta_l) \int_{\zeta_2}^{\zeta_{n-1}} [\zeta_n - v]^{-\alpha-1} dv \\ & \leq \frac{1}{24\alpha} \max_{\zeta_2 \leq \eta_2 \leq \zeta_{n-1}} |g^{(4)}(\eta_2)| \tilde{f}(\zeta_{l-3}, \zeta_{l-2}, \zeta_{l-1}, \zeta_l) (\zeta_n - \zeta_{n-1})^{-\alpha}. \end{aligned}$$

Here, $\tilde{f}(\zeta_{l-3}, \zeta_{l-2}, \zeta_{l-1}, \zeta_l)$ is the $\max_{\zeta_2 \leq v \leq \zeta_{n-1}} [(v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l)]$. Using Lemma (2.4), we get the following inequality

$$\begin{aligned} & \left| \sum_{l=3}^{n-1} \int_{\zeta_{l-1}}^{\zeta_l} \frac{g^{(4)}(\eta_l)}{4!} (v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l) [\zeta_n - v]^{-\alpha-1} dv \right| \\ & \leq \frac{3}{128\alpha} \max_{t_2 \leq x_1 \leq t_{n-1}} |g^{(4)}(x_1)| L^{4-\alpha} (\tau)^{4-\alpha}. \end{aligned} \quad (2.23)$$

Remark 3. If scale function ζ fulfills the condition of Lipschitz function such that $|\zeta_l - \zeta_{l-1}| \leq L\tau$ then, $\max_{\zeta_2 \leq v \leq \zeta_{n-1}} |(v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l)|$ is obtained at $v = \zeta_{l-2} + \frac{L\tau}{2}$, therefore

$$\tilde{f}(\zeta_{l-3}, \zeta_{l-2}, \zeta_{l-1}, \zeta_l) = \max_{\zeta_2 \leq v \leq \zeta_{n-1}} [(v - \zeta_{l-3})(v - \zeta_{l-2})(v - \zeta_{l-1})(v - \zeta_l)] \leq \frac{9}{16} L^4 (\tau)^4.$$

Consider the second part of RHS of Eq. (2.22)

$$\begin{aligned}
& \left| \int_{\zeta_{n-1}}^{\zeta_n} \frac{g^{(4)}(\eta_n)}{4!} (v - \zeta_{n-3})(v - \zeta_{n-2})(v - \zeta_{n-1})(v - \zeta_n)[\zeta_n - v]^{-\alpha-1} dv \right| \\
& \leq \frac{1}{24} \max_{\zeta_{n-1} \leq \eta_3 \leq \zeta_n} |g^{(4)}(\eta_3)| \left| \int_{\zeta_{n-1}}^{\zeta_n} (v - \zeta_{n-3})(v - \zeta_{n-2})(v - \zeta_{n-1})[\zeta_n - v]^{-\alpha} dv \right| \\
& \leq \frac{1}{12(1-\alpha)(2-\alpha)} \left(1 + \frac{3}{(3-\alpha)} + \frac{3}{(3-\alpha)(4-\alpha)} \right) \max_{\zeta_{n-1} \leq \eta_3 \leq \zeta_n} |g^{(4)}(\eta_3)| (\zeta_n - \zeta_{n-1})^{4-\alpha} \\
& \leq \frac{1}{12(1-\alpha)(2-\alpha)} \left(1 + \frac{3}{(3-\alpha)} + \frac{3}{(3-\alpha)(4-\alpha)} \right) \max_{t_{n-1} \leq x_2 \leq t_n} |g^{(4)}(x_2)| L^{4-\alpha}(\tau)^{4-\alpha}.
\end{aligned} \tag{2.24}$$

Now combining (2.21), (2.23) and (2.24), then error bound is

$$\begin{aligned}
|E_\tau^n| \leq \frac{\alpha [\omega_n]^{-1}}{\Gamma(1-\alpha)} & \left\{ \frac{1}{72} \max_{t_0 \leq x \leq t_2} |g^{(3)}(x)| (t_n - t_2)^{-\alpha-1} L^{3-\alpha}(\tau)^4 \right. \\
& + \left[\frac{3}{128\alpha} + \frac{1}{12(1-\alpha)(2-\alpha)} \left(1 + \frac{3}{(3-\alpha)} \right. \right. \\
& \left. \left. + \frac{3}{(3-\alpha)(4-\alpha)} \right) \right] \max_{t_0 \leq x_1 \leq t_n} |g^{(4)}(x_1)| L^{4-\alpha}(\tau)^{4-\alpha} \left. \right\}.
\end{aligned} \tag{2.25}$$

□

3. Numerical scheme for the generalized fractional advection-diffusion equation

In this section, we study the numerical scheme for solving the generalized fractional advection-diffusion defined by (1.1).

Let $u(x, t) = \omega(t)U(x, t) - \omega(0)U_0(x)$. We rewrite Eq. (1.1) to a similar equation using unity weight function and same scale function $\zeta(t)$. Then, Eq. (1.1) converts into the following form:

$$\begin{cases} {}_0^C \mathcal{D}_{t;[\zeta(t),1]}^\alpha u(x, t) = D \frac{\partial^2 u(x,t)}{\partial x^2} - A \frac{\partial u(x,t)}{\partial x} + f(x, t), & x \in \Omega, \quad t \in (0, T], \\ u(x, 0) = 0, & x \in \bar{\Omega} = \Omega \cup \partial\Omega, \\ u(0, t) = \phi_1(t), \quad u(a, t) = \phi_2(t), & t \in (0, T], \end{cases} \tag{3.1}$$

where $\phi_1(t) = \omega(t)\rho_1(t) - \omega(0)\rho_1(0)$, $\phi_2(t) = \omega(t)\rho_2(t) - \omega(0)\rho_2(0)$, and $f(x, t) = D\omega(0)U_0''(x) - A\omega(0)U_0'(x) + \omega(t)g(x, t)$.

For the uniform spatial mesh, let $0 = x_0 < x_1 < \dots < x_M = a$ of the interval $[0, a]$ with step size $h = \frac{a}{M}$ where M denotes number of subintervals, the grid points $x_0 + ih$ ($0 \leq i \leq M$), and $\tau = \frac{T}{N}$ be the step-size in temporal direction with grids $t_n = n\tau$ ($0 = t_0 < t_1 < \dots < t_n = T$).

Now, we discretize our problem (3.1) at (x_i, t_n) , then we get

$${}_0^C \mathcal{D}_{t;[\zeta(t),1]}^\alpha u(x_i, t_n) = Du_{xx}(x_i, t_n) - Au_x(x_i, t_n) + f(x_i, t_n). \quad (3.2)$$

In Eq. (3.2), for fixed t_n and $1 \leq i \leq M - 1$, the first and second order spatial derivatives discretized by using the following central difference approximations:

$$\frac{\partial u(x_i, t_n)}{\partial x} = \frac{u_n^{i+1} - u_n^{i-1}}{2h} + \mathcal{O}(h^2), \quad (3.3)$$

$$\frac{\partial^2 u(x_i, t_n)}{\partial x^2} = \frac{u_n^{i+1} - 2u_n^i + u_n^{i-1}}{h^2} + \mathcal{O}(h^2). \quad (3.4)$$

With the help of Equation (2.14), we get an approximation of the generalized Caputo-type fractional derivative term in (3.2) as follows:

$$\begin{aligned} {}^H \mathcal{D}_{t;[\zeta(t),1]}^\alpha u(x_i, t_n) &= \lambda_0 u(x_i, t_n) + \lambda_1 u(x_i, t_{n-1}) + \lambda_2 u(x_i, t_{n-2}) + \sum_{l=3}^{n-3} \lambda_{n-l} u(x_i, t_l) \\ &\quad + \lambda_{n-2} u(x_i, t_2) + \lambda_{n-1} u(x_i, t_1) + \lambda_n u(x_i, t_0) + \mathcal{O}(\tau^{4-\alpha}). \end{aligned} \quad (3.5)$$

where λ_l defined in Eq. (2.15). Next, we use Eq. (3.3), Eq. (3.4), and Eq. (3.5) in discretized equation (3.2), yields

$$\sum_{l=0}^n \lambda_{n-l} u(x_i, t_l) = Du_{xx}(x_i, t_n) - Au_x(x_i, t_n) + f_n^i + e_n^i. \quad (3.6)$$

where, $|e_n^i| \leq \tilde{c}(\tau^{4-\alpha} + h^2)$ for some constant \tilde{c} .

Now, we use numerical approximation u_n^i of $u(x_i, t_n)$ to neglect the truncation error term in Equation (3.6), then we determine the following finite difference scheme:

$$\begin{aligned} \lambda_0 u_n^i + \lambda_1 u_{n-1}^i + \sum_{l=3}^{n-2} \lambda_{n-l} u_l^i + \lambda_{n-2} u_2^i + \lambda_{n-1} u_1^i + \lambda_n u_0^i &= D \frac{u_n^{i+1} - 2u_n^i + u_n^{i-1}}{h^2} \\ &\quad - A \frac{u_n^{i+1} - u_n^{i-1}}{2h} + f_n^i, \end{aligned} \quad (3.7)$$

where $1 \leq n \leq N$, $1 \leq i \leq M - 1$. That is,

$$\begin{cases} \left(\frac{D}{h^2} + \frac{A}{2h}\right)u_1^{i-1} - \left(\lambda_0 + \frac{2D}{h^2}\right)u_1^i + \left(\frac{D}{h^2} - \frac{A}{2h}\right)u_1^{i+1} = \lambda_1 u_0^i - f_1^i, & n = 1, \\ \left(\frac{D}{h^2} + \frac{A}{2h}\right)u_2^{i-1} - \left(\lambda_0 + \frac{2D}{h^2}\right)u_2^i + \left(\frac{D}{h^2} - \frac{A}{2h}\right)u_2^{i+1} = \lambda_1 u_1^i + \lambda_2 u_0^i - f_2^i, & n = 2, \\ \left(\frac{D}{h^2} + \frac{A}{2h}\right)u_n^{i-1} - \left(\lambda_0 + \frac{2D}{h^2}\right)u_n^i + \left(\frac{D}{h^2} - \frac{A}{2h}\right)u_n^{i+1} = \lambda_1 u_{n-1}^i + \sum_{l=0}^{n-2} \lambda_{n-l} u_l^i - f_n^i, & n \geq 3. \\ u_0^i = 0, & 0 \leq i \leq M, \\ u_n^0 = \phi_1(t_n), \quad u_n^M = \phi_2(t_n), & 0 \leq n \leq N, \end{cases} \quad (3.8)$$

We rewrite the matrix form of above equation as follows:

$$\begin{cases} KU_1 = \lambda_1 U_0 - F_1 + H_1, & n = 1, \\ KU_2 = \lambda_1 U_1 + \lambda_2 U_0 - F_2 + H_2, & n = 2, \\ KU_n = \lambda_1 U_{n-1} + \sum_{l=0}^{n-2} \lambda_{n-l} U_l - F_n + H_n, & n \geq 3, \end{cases} \quad (3.9)$$

where the matrix as well as vectors of Equation (3.9) are defined as follows:

$$K = \text{tri} \left[\frac{D}{h^2} + \frac{A}{2h}, -\lambda_0 - \frac{2D}{h^2}, \frac{D}{h^2} - \frac{A}{2h} \right]_{(M-1) \times (M-1)},$$

$$U_n = (u_n^1, u_n^2, \dots, u_n^{M-1})^T,$$

$$F_n = (f_n^1, f_n^2, \dots, f_n^{M-1})^T,$$

$$H_n = \left(\left(-\frac{D}{h^2} - \frac{A}{2h}\right)u_n^0, 0, \dots, 0, \left(-\frac{D}{h^2} + \frac{A}{2h}\right)u_n^M \right)^T, \quad 1 \leq n \leq N.$$

Remark 4. Since the coefficient matrix A is a tridiagonal and strictly diagonally dominant then $\det(A) \neq 0$ (Levy-Desplanques theorem). Therefore at each time level t_n , the proposed scheme (3.8) has an unique solution for $1 \leq n \leq N$.

Theorem 2. The local truncation error of difference scheme (3.7) at (x_i, t_n) , $1 \leq i \leq M - 1$, $1 \leq n \leq N$ is

$$|e_n^i| \leq C(h^2 + \tau^{4-\alpha}), \quad (3.10)$$

where C is the positive constant independent of the time and space step sizes.

Proof. From Eqs. (4.7) the LTE of difference scheme (3.7) is

$$\begin{aligned}
e_n^i &= \lambda_0 u(x_i, t_n) + \lambda_1 u(x_i, t_{n-1}) + \lambda_2 u(x_i, t_{n-2}) + \sum_{l=3}^{n-3} \lambda_{n-l} u(x_i, t_l) + \lambda_{n-2} u(x_i, t_2) \\
&+ \lambda_{n-1} u(x_i, t_1) + \lambda_n u(x_i, t_0) - D \frac{u_n^{i+1} - 2u_n^i + u_n^{i-1}}{h^2} + A \frac{u_n^{i+1} - u_n^{i-1}}{2h} - f_n^i \\
&= \left[\lambda_0 u(x_i, t_n) + \lambda_1 u(x_i, t_{n-1}) + \lambda_2 u(x_i, t_{n-2}) + \sum_{l=2}^{n-3} \lambda_{n-l} u(x_i, t_l) + \lambda_{n-1} u(x_i, t_1) \right. \\
&+ \left. \lambda_n u(x_i, t_0) - {}^{\mathcal{H}} \mathcal{D}_{t;[\zeta(t),1]}^\alpha u(x_i, t_n) \right] - D \left[\frac{u_n^{i+1} - 2u_n^i + u_n^{i-1}}{h^2} - \frac{\partial^2 u(x_i, t_n)}{\partial x^2} \right] \\
&+ A \left[\frac{u_n^{i+1} - u_n^{i-1}}{2h} - \frac{\partial u(x_i, t_n)}{\partial x} \right] \\
&= \mathcal{O}(\tau^{4-\alpha}) - D \mathcal{O}(h^2) + A \mathcal{O}(h^2) = \mathcal{O}(\tau^{4-\alpha} + h^2).
\end{aligned}$$

□

In next theorem we use $L^2(\Omega)$ -space with the norm $\|\cdot\|_2$ and the inner product $\langle \cdot, \cdot \rangle$.

Theorem 3. (see [21]) *If tridiagonal matrix elements satisfy the inequality*

$$\lambda_1 \leq (\lambda_0)^{M-3} \left(\lambda_0 + \frac{D}{h^2} - \frac{A}{2h} \right) \left(\lambda_0 + \frac{D}{h^2} + \frac{A}{2h} \right), \quad (3.11)$$

then finite difference scheme is stable.

Proof. The Eq. (3.9) can be rewritten as following, for $1 \leq n \leq N$

$$KU_n = \lambda_1 U_{n-1} + \sum_{l=0}^{n-2} \lambda_{n-l} U_l - F_n + H_n,$$

let,

$$V_n = \sum_{l=0}^{n-2} \lambda_{n-l} U_l - F_n + H_n.$$

Since matrix K is invertible then above equation can be rewritten as

$$U_n = \lambda_1 K^{-1} U_{n-1} + K^{-1} V_n.$$

Using recurrence relation, we get the following equation

$$\begin{aligned}
U_n &= (\lambda_1 K^{-1})^2 U_{n-2} + (\lambda_1 K^{-1}) K^{-1} V_{n-1} + K^{-1} V_n \\
&= (\lambda_1 K^{-1})^n U_0 + (\lambda_1 K^{-1})^{n-1} K^{-1} V_1 + (\lambda_1 K^{-1})^{n-2} K^{-1} V_2 + \dots + K^{-1} V_n.
\end{aligned} \quad (3.12)$$

Let \tilde{U}_n is the approximate solution of Eq. (3.9), then we define error at grid point $t_n = n\tau$

$$e_n = U_n - \tilde{U}_n, \quad 0 \leq n \leq N.$$

Then, we obtain

$$e_n = (\lambda_1 K^{-1})^n e_0, \quad 1 \leq n \leq N.$$

From definition of compatible matrix norm

$$\|e_n\| \leq \|(\lambda_1 K^{-1})^n\| \|e_0\|.$$

Since,

$$\|(\lambda_1 K^{-1})^n\| \leq \|(\lambda_1 K^{-1})\| \|(\lambda_1 K^{-1})^{n-1}\| \leq \dots \leq \|(\lambda_1 K^{-1})\|^n$$

According to Ostrowski theorem ([36], Theorem 3.1), let $K = [a_{i,j}]_{M-1 \times M-1}$, then determinant of K satisfies

$$\begin{aligned} |\det(K)| &\geq \prod_{i=1}^{M-1} \left(|a_{i,i}| - \sum_{j=1, j \neq i}^{M-1} |a_{i,j}| \right) \\ &= (\lambda_0)^{M-3} \left(\lambda_0 + \frac{D}{h^2} - \frac{A}{2h} \right) \left(\lambda_0 + \frac{D}{h^2} + \frac{A}{2h} \right). \end{aligned} \quad (3.13)$$

Using assumed inequality (3.11) into (3.13), we get $|\det(K^{-1})| \leq \frac{1}{\lambda_1}$,

it implies that,

$$|\lambda_1| \|K^{-1}\| \leq 1.$$

Therefore,

$$|e_n| \leq |e_0|.$$

Thus, the numerical scheme is stable. \square

Theorem 4. *The solution U_n^i of the difference scheme (3.7) satisfies*

$$\max_{i,n} |U(x_i, t_n) - \tilde{U}(x_i, t_n)| \leq C(h^2 + \tau^{4-\alpha}) \quad (3.14)$$

for some constant C .

Proof. The truncation error for difference scheme at $(x_i, t_n) \in [0, a] \times [0, T]$ is

$$|e_n^i| \leq C(h^2 + \tau^{4-\alpha})$$

by Theorem 2. Now using theorem of stability, it implies

$$\max_{i,n} |U(x_i, t_n) - \tilde{U}(x_i, t_n)| \leq C|U(x_i, t_0) - \tilde{U}(x_i, t_0)| \quad (3.15)$$

We obtain the desired result easily after using the truncation error as discussed in Theorem 2. \square

4. Numerical results

In this section, we will check the numerical accuracy of the proposed schemes (2.14) and difference scheme (3.8), also verify the theoretical convergence order discussed in Theorem (4). Here, we provide three examples to numerically support our theory; in the first example we check convergence order and absolute error of approximation for the GCFD, while last two problems get the form (3.1) to describe accuracy and maximum absolute error of the difference scheme. We provide two Tables (7) and (10) to compare our scheme (3.8) with Gao et al. ([35], Example 4.1) and Cao et al. ([20], Example 5.1), respectively for particular choice of scale $\zeta(t) = t$ and weight $\omega(t) = 1$ functions. All numerical results are implemented in MATLAB R2018b.

To calculate the maximum absolute error E_∞ and error E_2 corresponding to the L_2 -norm, we use following formulas, respectively.

$$E_\infty(M, N) = \max_{1 \leq i \leq M-1} |U_N^i - u_N^i|,$$

$$E_2(M, N) = \left(h \sum_{i=1}^{M-1} |U_N^i - u_N^i|^2 \right)^{1/2},$$

where $\{U_n^i\}$ is the exact solution of advection diffusion equation and $\{u_n^i\}$ is the approximate solution at the point (x_i, t_n) .

Moreover, the convergence order in space and time direction for the described difference scheme corresponding to L_∞ -norm can be evaluated using the following formulas. R_x is the order of convergence in space side and R_t for temporal side.

$$R_x = \frac{\log(E(2M, N)) - \log(E(M, N))}{\log(2)},$$

and

$$R_t = \frac{\log(E(M, 2N)) - \log(E(M, N))}{\log(2)}.$$

Example 4.1. [35] Take function $u(t) = t^{4+\alpha}$, $t \in [0, 1]$, for $0 < \alpha < 1$. Determine the α -th order GCFD for $u(t)$ at $T = 1$ numerically.

The maximum absolute error and rate of convergence for the scheme (2.14) to approximate the GCFD of function $u(t)$ for $\alpha = 0.2, 0.5, 0.8$ with uniform time steps $1/10, 1/20, 1/40, 1/80, 1/160$ are calculated and shown in following tables.

Table 1 shows that the errors with respect to L_∞ -norm and convergence rates in time, and these data are found after calculating the classical Caputo derivative (i.e. taking $\zeta(t) = t$, $\omega(t) = 1$) of function $u(t)$ with the help of scheme (2.14). From this Table, we can see that the errors of our scheme (2.14) obtaining from GCFD approximation are lesser in compare to scheme developed in Gao et al. [35] for approximation of Caputo derivative and convergence rate of our scheme is $(4 - \alpha)$, while accuracy for time derivative in [35] is $(3 - \alpha)$. In

Table 2, to compute the maximum errors E_∞ and order of convergence in time direction, we take $\omega(t) = e^t$ while scale function is fixed with t . From Table 3, we validate the convergence for $\zeta(t) = t$, $\omega(t) = t + 1$ and the CPU time in seconds for $\alpha = 0.8$ are discussed. Table 4 shows that maximum errors and order of convergence for different choice of weight $\omega(t) = t^{0.5}$, t , t^4 , e^{2t} , scale is $\zeta(t) = t$ and $\alpha = 1/3$. In all cases, accuracy in time is obtained as $(4 - \alpha)$ for the scheme (2.14), which is higher than [34, 1].

Table 1: E_∞ errors and convergence rates R_t for Example 4.1, when $\zeta(t) = t$, $\omega(t) = 1$ and different α 's.

N	$\alpha = 0.2$		$\alpha = 0.5$		$\alpha = 0.5$ [35]	
	E_∞ error	R_t	E_∞ error	R_t	E_∞ error	R_t
10	1.6978e-04		1.5401e-03		1.3507e-02	
20	1.3130e-05	3.6928	1.4383e-04	3.4206	2.6121e-03	2.3704
40	9.9792e-07	3.7178	1.3116e-05	3.4550	4.8618e-04	2.4256
80	7.4966e-08	3.7346	1.1811e-06	3.4731	8.8645e-05	2.4554
160	5.5944e-09	3.7442	1.0560e-07	3.4835	1.5975e-05	2.4722

Table 2: E_∞ errors and convergence rates R_t for Example 4.1, when $\zeta(t) = t$, $\omega(t) = e^t$ and different α 's.

N	$\alpha = 0.2$		$\alpha = 0.5$		$\alpha = 0.8$		$\alpha = 0.8$
	E_∞ error	R_t	E_∞ error	R_t	E_∞ error	R_t	CPU time (s)
10	7.5552e-04		6.3075e-03		3.2184e-02		11.354861
20	7.1075e-05	3.4101	6.7873e-04	3.2162	4.2148e-03	2.9328	33.306962
40	5.9370e-06	3.5815	6.6677e-05	3.3476	5.0375e-04	3.0647	82.772072
80	4.6934e-07	3.6610	6.2491e-06	3.4155	5.7497e-05	3.1312	180.064565
160	3.5650e-08	3.7186	5.7027e-07	3.4539	6.4090e-06	3.1653	419.978865

Table 3: E_∞ errors and convergence rates R_t for Example 4.1, when $\zeta(t) = t$, $\omega(t) = t + 1$ and different α 's.

N	$\alpha = 0.2$		$\alpha = 0.5$		$\alpha = 0.8$		$\alpha = 0.8$
	E_∞ error	R_t	E_∞ error	R_t	E_∞ error	R_t	CPU time (s)
10	3.5019e-04		3.1752e-03		1.7251e-02		10.194270
20	3.0621e-05	3.5155	3.1426e-04	3.3368	2.0904e-03	3.0448	25.931157
40	2.4466e-06	3.6457	2.9499e-05	3.4132	2.3982e-04	3.1238	65.689186
80	1.8844e-07	3.6986	2.6976e-06	3.4509	2.6799e-05	3.1617	155.623905
160	1.4248e-08	3.7252	2.4326e-07	3.4711	2.9560e-06	3.1805	384.953845

Table 4: E_∞ errors and convergence rates R_t for Example 4.1, when $\zeta(t) = t$, $\alpha = \frac{1}{3}$ and different weight functions.

N	$\omega(t) = t^{0.5}$		$\omega(t) = t$		$\omega(t) = t^4$		$\omega(t) = e^{2t}$	
	E_∞ error	R_t	E_∞ error	R_t	E_∞ error	R_t	R_t	R_t
10	9.5450e-04		1.7465e-03		3.2557e-02			
20	8.4374e-05	3.4999	1.5027e-04	3.5388	2.0348e-03	4.0000	3.2332	
40	7.0877e-06	3.5734	1.2845e-05	3.5483	1.2749e-04	3.9964	3.4225	
80	5.8166e-07	3.6071	1.0650e-06	3.5923	1.1169e-05	3.5129	3.5220	
160	4.7123e-08	3.6257	8.6813e-08	3.6167	9.3941e-07	3.5716	3.5764	




Figure 1: Errors plot of the numerical results for the Example (4.1) at final time $T = 1$ for different values of α (Left side (1a) for $\alpha = 0.5$; Right side (1b) for $\alpha = 0.8$).


Example 4.2. [20] We take the following generalized fractional advection-diffusion equation:

$$\begin{cases} {}_0^C \mathcal{D}_{t;[\zeta(t),\omega(t)]}^\alpha u(x,t) = \frac{\partial^2 u(x,t)}{\partial x^2} - \frac{\partial u(x,t)}{\partial x} + f(x,t), & (x,t) \in (0,1) \times (0,1), \\ u(x,0) = 0, & x \in (0,1), \\ u(0,t) = t^{6+\alpha}, u(1,t) = et^{6+\alpha}, & t \in (0,1], \end{cases}$$


where $f(x,t) = e^{xt} \frac{\Gamma(7+\alpha)}{720}$. When $\zeta(t) = t$ and $\omega(t) = 1$, then $u(x,t) = e^{xt} t^{6+\alpha}$ is the exact solution.

To solve this example, we use the numerical scheme defined in (3.8). The maximum errors at time $t = 1$ for different values of α with different step sizes, and rate of convergence in time direction and space direction are displayed in Tables 5 and 6. In Table 5, we set $h = \frac{1}{2000}$ and describe the numerical errors and convergence rates in time R_t for different values of N . In Table 6, we fix $\tau = \frac{1}{500}$ and present the numerical errors and spatial


convergence rates R_x for different values of M . It is shown that our scheme (3.8) gives $(4 - \alpha)$ -order convergence in temporal direction and second-order convergence in spatial direction.




(a) Exact solution for $\alpha = 0.95$



(b) Numerical solution for $\alpha = 0.95$



(c) Numerical solution for $\alpha = 0.6$




(d) Numerical solution for $\alpha = 0.45$


Figure 2: Exact and approximate solutions of Example 4.2 for different α 's with $M = N = 200$ and $\omega(t) = 1$, $\zeta(t) = t$.

Table 5: Errors E_∞ and E_2 with convergence rates in time for Example 4.2, when $h = 1/2000$ and different α 's.

α	N	E_∞ error	R_t	E_2 error	R_t
0.8	8	1.6385e-02		7.4195e-03	
	16	2.2920e-03	2.8377	1.3514e-03	2.4569
	32	2.8191e-04	3.0233	1.8556e-04	2.8645
	64	3.2609e-05	3.1119	2.2573e-05	3.0392
	128	3.6574e-06	3.1564	2.5937e-06	3.1215
0.5	8	3.7940e-03		1.8168e-03	
	16	4.2889e-04	3.1451	2.5900e-04	2.8104
	32	4.3064e-05	3.3160	2.8647e-05	3.1765
	64	4.0768e-06	3.4010	2.8348e-06	3.3371
	128	3.7173e-07	3.4551	2.6407e-07	3.4243
0.2	8	5.4498e-04		2.7293e-04	
	16	5.1093e-05	3.4150	3.1455e-05	3.1172
	32	4.2949e-06	3.5724	2.8822e-06	3.4480
	64	3.3828e-07	3.6663	2.3627e-07	3.6087
	128	2.2628e-08	3.9020	1.6173e-08	3.8687



(a) $\zeta(t) = e^t$, $\omega(t) = 1$



(b) $\zeta(t) = t^{1/2}$, $\omega(t) = 1$

Figure 3: Numerical solutions of Example 4.2 with different choices of weight functions $\omega(t)$ and scale functions $\zeta(t)$, and $M = N = 200$ with $\alpha = 0.9$.

Clearly visible from above figures that scale function $\zeta(t)$ can stretch or contract the domain.

Table 6: The maximum errors and convergence rates R_x for Example 4.2, when $\tau = 1/500$ and different α 's.

M	$\alpha = 0.2$		$\alpha = 0.5$		$\alpha = 0.8$	
	E_∞ error	R_x	E_∞ error	R_x	E_∞ error	R_x
8	2.3437e-04		2.1275e-04		1.8078e-04	
16	5.8814e-05	1.9945	5.3334e-05	1.9960	4.5242e-05	1.9985
32	1.4733e-05	1.9971	1.3373e-05	1.9958	1.1319e-05	1.9990
64	3.6832e-06	2.0001	3.3408e-06	2.0010	2.7940e-06	2.0183
128	9.2088e-07	1.9999	8.3276e-07	2.0042	6.6258e-07	2.0762

Table 7: The maximum errors and convergence rates R_t of Example 4.1 in [35] (page no 43) for different values of α with $\tau = 1/2000$.

α	N	Current scheme		$L1 - 2$ [35]	
		E_∞	R_t	E_∞	R_t
0.9	10	2.2536e-03		1.8600e-02	
	20	3.2727e-04	2.7837	4.7228e-03	1.9776
	40	4.1906e-05	2.9653	1.1488e-03	2.0395
	80	5.1072e-06	3.0366	2.7367e-04	2.0697
	160	6.1092e-07	3.0635	6.4520e-05	2.0846
0.5	10	2.5470e-04		2.4936e-03	
	20	2.6005e-05	3.2919	4.8366e-04	2.3662
	40	2.4615e-06	3.4012	9.0163e-05	2.4234
	80	2.2827e-07	3.4307	1.6457e-05	2.4539
	160	2.3662e-08	3.2701	2.9700e-06	2.4701

Example 4.3. Take the following fractional advection-diffusion equation:

$$\begin{cases} {}_0^C \mathcal{D}_{t, [\zeta(t), \omega(t)]}^\alpha u(x, t) = \frac{\partial^2 u(x, t)}{\partial x^2} - \frac{\partial u(x, t)}{\partial x} + f(x, t), & (x, t) \in (0, 1) \times (0, 1), \\ u(x, 0) = 0, & x \in (0, 1), \\ u(0, t) = 0, \\ u(1, t) = t^7(\sin 1), & t \in (0, 1], \end{cases}$$

where $f(x, t) = \frac{\Gamma 8}{\Gamma(8-\alpha)} t^{7-\alpha} \sin(x) + t^7(\sin(x) + \cos(x))$. When $\zeta(t) = t$ and $\omega(t) = 1$, the exact solution is $u(x, t) = t^7 \sin(x)$.

To solve this PDE, we use our difference scheme (3.8). Here, two Tables 8, 9 are given in support of numerical results. In Table 8, we take fixed space step size $h = 1/2000$ and display maximum absolute errors and errors E_2 with respect to norm L_2 , also rate of convergence for temporal direction with different time steps $N = 8, 16, 32, 64, 128$. In Table 9, we expressed maximum errors and convergence order in space direction to set time steps fix with $\tau = 1/500$ and taking different $M = 8, 16, 32, 64, 128$.

Table 8: Errors E_∞ and E_2 with convergence rates in time for Example 4.3, when $h = 1/2000$ and different α 's.

α	N	E_∞ error	R_t	E_2 error	R_t
0.8	8	5.2117e-03		2.2239e-03	
	16	7.4012e-04	2.8159	4.1857e-04	2.4095
	32	9.1743e-05	3.0121	5.8367e-05	2.8422
	64	1.0658e-05	3.1057	7.1562e-06	3.0279
	128	1.2018e-06	3.1486	8.2814e-07	3.1112
0.5	8	1.4554e-03		6.2805e-04	
	16	1.7116e-04	3.0880	9.7274e-05	2.6907
	32	1.7546e-05	3.2862	1.1191e-05	3.1198
	64	1.6840e-06	3.3812	1.1323e-06	3.3049
	128	1.5961e-07	3.3993	1.1013e-07	3.3620
0.2	8	2.5522e-04		1.1012e-04	
	16	2.5499e-05	3.3233	1.4486e-05	2.9264
	32	2.2221e-06	3.5204	1.4170e-06	3.3538
	64	1.8445e-07	3.5906	1.2401e-07	3.5144
	128	1.8801e-08	3.2944	1.3014e-08	3.2523

Table 9: The maximum errors and convergence rates R_x for Example 4.3, when $\tau = 1/500$ and different α 's.

M	$\alpha = 0.2$		$\alpha = 0.5$		$\alpha = 0.8$	
	E_∞ error	R_x	E_∞ error	R_x	E_∞ error	R_x
8	3.0308e-04		2.7311e-04		2.3102e-04	
16	7.6029e-05	1.9951	6.8533e-05	1.9946	5.8003e-05	1.9938
32	1.9050e-05	1.9968	1.7180e-05	1.9961	1.4560e-05	1.9941
64	4.7625e-06	2.0000	4.2962e-06	1.9996	3.6518e-06	1.9953
128	1.1909e-06	1.9997	1.0752e-06	1.9985	9.2444e-07	1.9820

Table 10: The maximum errors and convergence rates R_t for Example 5.1 in [20], when $h = 1/6000$ and spatial convergence rates R_x , when $\tau = 1/200$.

$\alpha = 0.368$			Cao et al. [20]	$\alpha = 0.368$			Cao et al. [20]
N	E_∞ error	R_t	R_t	M	E_∞ error	R_x	R_x
8	8.1539e-04			4	8.8034e-04		
16	7.7720e-05	3.3911	3.4031	8	2.2559e-04	1.9643	1.9643
32	6.8845e-06	3.4969	3.4972	16	5.6583e-05	1.9953	1.9953
64	5.8739e-07	3.5510	3.5528	32	1.4173e-05	1.9972	1.9972
128	4.9904e-08	3.5571	3.5836	64	3.5359e-06	2.0030	2.0030

In Table 10, we validate the proposed scheme with [20] (Example 5.1). Firstly, we present the max error, the convergence order with fixed $h = \frac{1}{6000}$ and varying N with 8, 16,

32, 64, 128 for $\alpha = 0.368$. It is noted that the convergence order of our scheme (3.8) for temporal direction is almost the same as [20]. After that, we expressed maximum error and convergence rate for spatial dimension to set $\tau = 1/200$ and changing $M = 4, 8, 16, 32, 64$ for same value of α , we observe that the spatial convergence order of our numerical scheme is same as [20]. Thus, the scheme presented in [20] becomes a particular case of the proposed scheme (3.8) for $\zeta(t) = t$ and $\omega(t) = 1$. Also, we can compute numerical results to for different suitable choices of scale and weight functions.

5. Conclusion

A high-order numerical scheme based on cubic interpolation formula is discussed for approximation of GCFD of α -th order. Properties of discretized coefficients are analyzed and local truncation error in approximation of GCFD is also discussed. Further, we establish a difference scheme for generalized fractional advection-diffusion equation using the developed approximation formula for GCFD. The stability and convergence of this established scheme to approximate the time fractional generalized advection-diffusion equation are studied. Order of accuracy for the difference scheme is $\mathcal{O}(\tau^{4-\alpha} + h^2)$, where step-sizes τ in time and h in space direction. The convergence order of the difference scheme is described by some numerical experiments.

Numerical calculations reveal that the proposed difference scheme has $(4 - \alpha)$ -th order convergence in time; and in space direction, it has second order convergence. The temporal rate of convergence of the scheme for generalized fractional advection-diffusion equation is highest accuracy to date. In addition, the developed scheme can be directly used to get the other Caputo-type of time fractional advection-diffusion equations by selecting the suitable scale $\zeta(t)$ and weight $\omega(t)$ functions in the generalized Caputo-type fractional derivative. The developed scheme is tested for the cases having smooth solutions of the considered fractional advection-diffusion equation. However, the case of the nonsmooth solutions will be presented in our future works.

Declarations

Funding: The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

Conflict of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Swati Yadav, Rajesh K Pandey, Anil K Shukla, and Kamlesh Kumar. High-order approximation for generalized fractional derivative and its application. *International Journal of Numerical Methods for Heat & Fluid Flow*, 29(9):3515–3534, 2019.
- [2] Om P Agrawal. Some generalized fractional calculus operators and their applications in integral equations. *Fractional Calculus and Applied Analysis*, 15(4):700–711, 2012.
- [3] David A Benson, Stephen W Wheatcraft, and Mark M Meerschaert. Application of a fractional advection-dispersion equation. *Water resources research*, 36(6):1403–1412, 2000.
- [4] Liuzong Zhou and HM Selim. Application of the fractional advection-dispersion equation in porous media. *Soil Science Society of America Journal*, 67(4):1079–1084, 2003.
- [5] Dumitru Baleanu, Guo-Cheng Wu, and Sheng-Da Zeng. Chaos analysis and asymptotic stability of generalized caputo fractional differential equations. *Chaos, Solitons & Fractals*, 102:99–105, 2017.
- [6] Debasis Dan, Chris Mueller, Kun Chen, and James A Glazier. Solving the advection-diffusion equations in biological contexts using the cellular potts model. *Physical Review E*, 72(4):041909, 2005.
- [7] JG Verwer, JG Blom, and W Hundsdorfer. An implicit-explicit approach for atmospheric transport-chemistry problems. *Applied Numerical Mathematics*, 20(1-2):191–209, 1996.
- [8] Mehdi Dehghan. Weighted finite difference techniques for the one-dimensional advection–diffusion equation. *Appl. Math. Comput.*, 147(2):307–319, 2004.
- [9] Akbar Mohebbi and Mehdi Dehghan. High-order compact solution of the one-dimensional heat and advection–diffusion equations. *Applied mathematical modelling*, 34(10):3071–3084, 2010.
- [10] Kolade M Owolabi. High-dimensional spatial patterns in fractional reaction-diffusion system arising in biology. *Chaos, Solitons & Fractals*, 134:109723, 2020.
- [11] Igor Podlubny. *Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*. Elsevier, 1998.
- [12] A Kilbas Anatoly. Hadamard-type fractional calculus. *Journal of the Korean Mathematical Society*, 38(6):1191–1204, 2001.
- [13] Sebastien Gaboury, Richard Tremblay, and Benoît-Jean Fugère. Some relations involving a generalized fractional derivative operator. *Journal of Inequalities and Applications*, 2013(1):1–9, 2013.
- [14] Abdon Atangana and Dumitru Baleanu. New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model. *arXiv preprint arXiv:1602.03408*, 2016.
- [15] Kassem Mustapha. An l1 approximation for a fractional reaction-diffusion equation, a second-order error analysis over time-graded meshes. *SIAM Journal on Numerical Analysis*, 58(2):1319–1338, 2020.
- [16] Anatoly A Alikhanov. A new difference scheme for the time fractional diffusion equation. *Journal of Computational Physics*, 280:424–438, 2015.
- [17] Omar Abu Arqub. Numerical solutions for the robin time-fractional partial differential equations of heat and fluid flows based on the reproducing kernel algorithm. *International Journal of Numerical Methods for Heat & Fluid Flow*, 28(4):828–856, 2018.
- [18] Zhiqiang Li and Yubin Yan. Error estimates of high-order numerical methods for solving time fractional partial differential equations. *Fractional Calculus and Applied Analysis*, 21(3):746–774, 2018.
- [19] Chunwan Lv and Chuanju Xu. Error analysis of a high order method for time-fractional diffusion equations. *SIAM Journal on Scientific Computing*, 38(5):A2699–A2724, 2016.
- [20] Jianxiong Cao, Changpin Li, and YangQuan Chen. High-order approximation to caputo derivatives and caputo-type advection-diffusion equations (ii). *Fractional calculus and Applied analysis*, 18(3):735–761, 2015.
- [21] Yufeng Xu and Om P Agrawal. Numerical solutions and analysis of diffusion for new generalized fractional burgers equation. *Fractional Calculus and Applied Analysis*, 16(3):709–736, 2013.
- [22] Kamlesh Kumar, Rajesh K Pandey, and Farheen Sultana. Numerical schemes with convergence for generalized fractional integro-differential equations. *Journal of Computational and Applied Mathematics*, 388:113318, 2021.
- [23] Kai Diethelm. An efficient parallel algorithm for the numerical solution of fractional differential equations. *Fractional Calculus and Applied Analysis*, 14(3):475–490, 2011.

- [24] Yunying Zheng, Changpin Li, and Zhengang Zhao. A note on the finite element method for the space-fractional advection diffusion equation. *Computers & Mathematics with Applications*, 59(5):1718–1726, 2010.
- [25] A Mardani, Mohammad Reza Hooshmandasl, Mohammad Hossein Heydari, and Carlo Cattani. A meshless method for solving the time fractional advection–diffusion equation with variable coefficients. *Computers & Mathematics with Applications*, 75(1):122–133, 2018.
- [26] Changpin Li and Min Cai. High-order approximation to caputo derivatives and caputo-type advection–diffusion equations: Revisited. *Numerical functional analysis and optimization*, 38(7):861–890, 2017.
- [27] WenYi Tian, Weihua Deng, and Yujiang Wu. Polynomial spectral collocation method for space fractional advection–diffusion equation. *Numerical Methods for Partial Differential Equations*, 30(2):514–535, 2014.
- [28] Pinghui Zhuang, Fawang Liu, Vo Anh, and Ian Turner. Numerical methods for the variable-order fractional advection-diffusion equation with a nonlinear source term. *SIAM Journal on Numerical Analysis*, 47(3):1760–1781, 2009.
- [29] Deeksha Singh, Farheen Sultana, and Rajesh K Pandey. Approximation of caputo-prabhakar derivative with application in solving time fractional advection-diffusion equation. *International Journal for Numerical Methods in Fluids*, 2022.
- [30] Wen Cao, Yufeng Xu, and Zhoushun Zheng. Finite difference/collocation method for a generalized time-fractional kdv equation. *Applied Sciences*, 8(1):42, 2018.
- [31] Yufeng Xu, Zhimin He, and Om P Agrawal. Numerical and analytical solutions of new generalized fractional diffusion equation. *Computers & Mathematics with Applications*, 66(10):2019–2029, 2013.
- [32] Kolade M Owolabi and Edson Pindza. Numerical simulation of chaotic maps with the new generalized caputo-type fractional-order operator. *Results in Physics*, 38:105563, 2022.
- [33] Zaid Odibat and Dumitru Baleanu. Numerical simulation of initial value problems with generalized caputo-type fractional derivatives. *Applied Numerical Mathematics*, 156:94–105, 2020.
- [34] Yufeng Xu, Zhimin He, and Qinwu Xu. Numerical solutions of fractional advection–diffusion equations with a kind of new generalized fractional derivative. *International Journal of Computer Mathematics*, 91(3):588–600, 2014.
- [35] Guang-hua Gao, Zhi-zhong Sun, and Hong-wei Zhang. A new fractional numerical differentiation formula to approximate the caputo fractional derivative and its applications. *Journal of Computational Physics*, 259:33–50, 2014.
- [36] Yufeng Xu, Zhimin He, and Qinwu Xu. Numerical solutions of fractional advection–diffusion equations with a kind of new generalized fractional derivative. *International Journal of Computer Mathematics*, 91(3):588–600, 2014.