

GENERALISED POWER SERIES DETERMINED BY LINEAR RECURRENCE RELATIONS

LOTHAR SEBASTIAN KRAPP, SALMA KUHLMANN, AND MICHELE SERRA

10 June 2022

ABSTRACT. Within the field of Laurent series with rational coefficients, those series representing rational functions can be characterised via linear recurrence relations of their coefficients. Inspired by this result, we establish a notion of generalised linear recurrence relations for fields of generalised power series. We study distinguished Hahn fields that are determined by generalised linear recurrence relations and apply our results to the study of automorphism groups of such fields.

1. INTRODUCTION

The global aim of this work is to describe distinguished Hahn fields. In [2] we focus on a description via k -hulls of \mathcal{F} (where \mathcal{F} is a set of supports in the value group) and by characterising the Hahn fields via properties of \mathcal{F} .

We still have the same global aim, but now our description uses another focus: what distinguished Hahn fields can be characterised as the solution sets of generalised linear recurrence relations? The motivation comes from the rational Hahn field. In the next paragraphs we gather together known results about the rational Hahn field, for the reader's convenience¹.

Let $\mathbb{Q}((t))$ denote the power series field over the rational numbers in one variable t and let $\mathbb{Q}(t)$ denote the field of rational functions over \mathbb{Q} in the variable t , i.e. the field of fractions of $\mathbb{Q}[t]$. For $a \in \mathbb{Q}((t))$ and $i \in \mathbb{Z}$, we denote by a_i the i -th coefficient of the power series expansion of a . The following result exhibits a way for determining whether a power series in $\mathbb{Q}((t))$ is already contained in $\mathbb{Q}(t)$ by means of linear recurrence relations (in fact, one can also choose any other coefficient field k rather than \mathbb{Q} as coefficient field. See [8, page 91 ff.] for further details).

Part of this work was carried out while all three authors were generously hosted by the Fields Institute, in Toronto, during the month of June 2022.

2010 *Mathematics Subject Classification*: 13J05 (12J20 16W60 06F20)

¹Readers familiar with this material can skip to Section 2

Fact 1.1. *Let*

$$a = \sum_{i=k}^{\infty} a_i t^i \in \mathbb{Q}((t))$$

for some $k \in \mathbb{Z}$. Then $a \in \mathbb{Q}(t)$ if and only if the following holds: there exist $m \in \mathbb{N}$ and $c_1, \dots, c_m \in \mathbb{Q}$ such that for any $n > m$ the linear recurrence relation with constant coefficients

$$(1.1) \quad a_n = \sum_{j=1}^m c_j a_{n-j}$$

holds.

Remark 1.2. In Fact 1.1, the element $a \in \mathbb{Q}(t)$ is not uniquely determined by m and c_1, \dots, c_m . For instance, we can consider $m = c_1 = 1$. Then any $a = \sum_{i=k}^d a_i t^i \in \mathbb{Q}(t)$ whose degree d is less than 1, such as $a = 1$ or $a = t^{-1}$, satisfies (1.1).

Conversely, also m and c_1, \dots, c_m are not uniquely determined by a . For instance, if we consider $a = t^{-1}$, then we could also set $m = 2$, $c_1 = 0$ and $c_2 = 1$. \square

By setting $c_0 = -1$, the linear recurrence relation (1.1) can be reformulated as

$$(1.2) \quad 0 = \sum_{j=0}^m c_j a_{n-j}.$$

We thus obtain the following consequence of Fact 1.1, which is a special case of the generalisation we will prove in Proposition 3.6.

Corollary 1.3. *Let*

$$a = \sum_{i \in \mathbb{Z}} a_i t^i \in \mathbb{Q}((t)).$$

Then $a \in \mathbb{Q}(t)$ if and only if the following holds: there exist $\ell \in \mathbb{N}$, $k_0, \dots, k_\ell \in \mathbb{Z}$ with $k_0 < \dots < k_\ell$ and $c_0, \dots, c_\ell \in \mathbb{Q}$, not all equal to 0, such that for any $n \in \mathbb{Z} \setminus \{k_0, \dots, k_\ell\}$ the following linear recurrence relation with constant coefficients holds

$$(1.3) \quad \sum_{j=0}^{\ell} c_j a_{n-j} = 0.$$

Note that the linear recurrence relation (1.3) in Corollary 1.3 needs to hold for all but finitely many coefficients, whereas the linear recurrence relation (1.1) in Fact 1.1 only has to be satisfied for all coefficients whose index is greater than m . The aforementioned results can be reformulated by means of Hankel matrices.

Definition 1.4. (i) A **Hankel matrix** is a (possibly infinite) matrix with constant anti-diagonals. For instance, a Hankel matrix $(H_{ij})_{i,j \in \mathbb{N}}$ indexed by \mathbb{N} satisfies for all $i, j, n \in \mathbb{N}$ with $i \geq j \geq 0$ and $n \leq i - j$ that $H_{ij} = H_{i-n, j+n}$. The entry H_{ij} thus only depends on the sum $i + j$.

(ii) Let $a \in \mathbb{Q}((t))$. Then the **Hankel matrix associated** to a is the matrix $\mathcal{H}^a = (\mathcal{H}_{ij}^a)_{i,j \in \mathbb{N}} \in \mathbb{Q}^{\mathbb{N} \times \mathbb{N}}$ with $\mathcal{H}_{ij}^a = a_{i+j-v(a)}$ for any $i, j \in \mathbb{N}$.

For instance, the Hankel matrix of the power series $e^t := \sum_{n=0}^{\infty} \frac{t^n}{n!} \in \mathbb{Q}((t))$ is

$$\mathcal{H}^{e^t} = \begin{pmatrix} \frac{1}{0!} & \frac{1}{1!} & \frac{1}{2!} & \cdots \\ \frac{1}{1!} & \frac{1}{2!} & \frac{1}{3!} & \cdots \\ \frac{1}{2!} & \frac{1}{3!} & \frac{1}{4!} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \in \mathbb{Q}^{\mathbb{N} \times \mathbb{N}}$$

Remark 1.5. Notice that distinct power series may have the same associated Hankel matrix. In fact, for any $a, b \in \mathbb{Q}((t))$ we have $\mathcal{H}^a = \mathcal{H}^b$ if and only if there exists $n \in \mathbb{Z}$ such that $a = t^n b$. \square

The linear recurrence relation (1.3) express that the Hankel matrix \mathcal{H}^a has only finitely many linearly independent columns, as the following result shows.

Corollary 1.6. *Let $a \in \mathbb{Q}((t))$. Then $a \in \mathbb{Q}(t)$ if and only if the rank of \mathcal{H}^a is finite.*

Proof. We may assume that $a \in \mathbb{Q}[t]$ with $a_0 \neq 0$, as otherwise we can replace a it by $t^k a$ for some $k \in \mathbb{Z}$ (see Remark 1.5). Hence, a is of the form $a = \sum_{i=0}^{\infty} a_i t^i$ with $a_0 \neq 0$. For any $j \in \mathbb{N}$, denote by $\mathcal{H}_{.j}^a$ the j -th column of \mathcal{H}^a . By Corollary 1.3, we have to verify that the rank of \mathcal{H}^a is finite if and only if there exist $\ell \in \mathbb{N}$, $k_0, \dots, k_\ell \in \mathbb{Z}$ with $k_0 < \dots < k_\ell$ and $c_0, \dots, c_\ell \in \mathbb{Q}$, not all equal to 0, such that for any $n \in \mathbb{Z} \setminus \{k_0, \dots, k_\ell\}$ equation (1.3) holds.

First suppose that such ℓ , k_i and c_i exist, and set $d = \max\{k_\ell + 1, \ell\}$. Then for any $n \in \mathbb{N}$ with $n \geq d$ we have

$$\sum_{j=0}^{\ell} c_j \mathcal{H}_{.n-j}^a = 0.$$

As not all scalars c_j are equal to zero, this shows that $\{\mathcal{H}_{.n-\ell}^a, \dots, \mathcal{H}_{.n}^a\}$ is linearly dependent over k and $\mathcal{H}_{.n}^a$ lies in the span of $\{\mathcal{H}_{.n-\ell}^a, \dots, \mathcal{H}_{.n-1}^a\}$. Hence, any column of \mathcal{H}^a lies in the span of $\{\mathcal{H}_{.0}^a, \dots, \mathcal{H}_{.d-1}^a\}$, showing that the rank of \mathcal{H}^a is at most d .

For the converse, suppose that \mathcal{H}^a has finite rank. Let $\ell \in \mathbb{N}$ such that any column of \mathcal{H}^a lies in the span of $\{\mathcal{H}_{.0}^a, \dots, \mathcal{H}_{.\ell-1}^a\}$. Moreover, set $k_i = i$ for $i \in \{0, \dots, \ell\}$. Then there are $c_0, \dots, c_\ell \in \mathbb{Q}$, not all equal to 0, such

that

$$\sum_{j=0}^{\ell} c_j \mathcal{H}_{\ell-j}^a = 0.$$

Now let $n \in \mathbb{Z} \setminus \{0, \dots, \ell\}$. If $n \leq -1$, then trivially $\sum_{j=0}^{\ell} c_j a_{n-j} = 0$, as $v(a) = 0$. Otherwise, $n \geq \ell + 1$ and we obtain

$$\sum_{j=0}^{\ell} c_j a_{n-j} = \sum_{j=0}^{\ell} c_j \mathcal{H}_{n-\ell, \ell-j}^a = 0,$$

as required. \square

2. GENERALISED LINEAR RECURRENCE RELATIONS

Our aim is to generalise the approach described for the field $\mathbb{Q}((t))$ to a general field $k((G))$ of generalised formal power series. We will define a notion of linear recurrence and characterise subfields of $k((G))$ that are determined by linear recurrence sequences, in a sense that we explain below.

Throughout this work, let k be a field, let G be an ordered abelian group and let \mathcal{F} be a family of well-ordered subsets of G . We denote by \mathbb{N} the set of natural numbers with 0. In most other regards, we follow the notation and terminology of [2].

For a partial function $s: G \rightarrow k$, the **support** of s is given by $\text{supp}(s) = \{g \in \text{dom}(s) \mid s(g) \neq 0\}$, where $\text{dom}(s)$ denotes the domain of s . We denote by $k((G))$ the set of all (total) functions $a: G \rightarrow k$ whose support $\text{supp}(a)$ is a well-ordered subset of G . For any $g \in G$, we denote by t^g the characteristic function mapping g to 1 and everything else to 0, and we call t^g a (monic) **monomial** of $k((G))$. This way we can express an element $a \in k((G))$ as a power series $a = \sum_{g \in G} a_g t^g$, where $a_g = a(g) \in k$. For any power series $a, b \in k((G))$, their sum is given by $a + b = \sum_{g \in G} (a_g + b_g) t^g$ and their product by $ab = \sum_{g \in G} c_g t^g$ with $c_g = \sum_{h \in G} a_h b_{g-h}$. Note that $\text{supp}(a + b) \subseteq \text{supp}(a) \cup \text{supp}(b)$ and $\text{supp}(ab) \subseteq \text{supp}(a) \oplus \text{supp}(b)$. These operations make $k((G))$ a field (cf. [1, 7]). The **group ring** $k[G] = \{a \in k((G)) \mid \text{supp}(a) \text{ is finite}\}$ is a subring of $k((G))$. Its field of fractions $k(G) = \text{ff}(k[G])$ will be important in the sequel.

Definition 2.1. We call $k((G))$ the **maximal Hahn field** with coefficient field k and value group G . We call $k(G)$ the **minimal Hahn field** and, generally, we call a **Hahn field** any field K such that $k(G) \subseteq K \subseteq k((G))$.

Now let K be a Hahn field. We define the **canonical valuation** by $v := v_{\min}: K^\times \rightarrow G$, $a \mapsto \min \text{supp}(a)$. To this we associate the valuation invariants: the valuation ring $R_K = \{a \in K \mid v(a) \geq 0\}$, the valuation ideal $I_K = \{a \in K \mid v(a) > 0\}$ and the residue field $\overline{K} = R_K/I_K \simeq k$.

In this section, we introduce the main subject of study of this work: subsets of $k((G))$ that are determined by generalised linear recurrence relations.

We first set up the required concepts as well as notations and then establish several algebraic properties of sets determined by generalised linear recurrence relations depending on the underlying set of generalised linear recurrence sequences.

Definition 2.2. A **(generalised) linear recurrence sequence** in $k((G))$ is a partial function s from G to k whose domain $\text{dom}(s)$ is a well-ordered subset of G . The order type of its domain corresponds to an ordinal number, which we denote by $\text{ot}(s)$. We say that s is **trivial** if $s(g) = 0$ for any $g \in \text{dom}(s)$. The set of all linear recurrence sequences in $k((G))$ is denoted by $\text{LRS}(k, G)$.

Definition 2.3. A linear recurrence sequence $s \in \text{LRS}(k, G)$ is called **0-free** if $s(g) \neq 0$ for any $g \in \text{dom}(s)$ (i.e. $\text{supp}(s) = \text{dom}(s)$).

Notation 2.4. Let $s \in \text{LRS}(k, G)$ and let $a \in k((G))$.

- (i) Let $\alpha = \text{ot}(s)$. Then s can be expressed as a sequence $(g_i, s_i)_{i < \alpha}$ in $G \times k$, where $(g_i)_{i < \alpha}$ is strictly increasing such that $\{g_i \mid i < \alpha\} = \text{dom}(s)$ and $s_i = s(g_i)$ for any $i < \alpha$.

- (ii) We say that

$$s^* = \sum_{g \in \text{dom}(s)} s(g)t^g \in k((G))$$

is the **power series associated** to s . Regarded as functions, s^* is an extension of s whose domain is $\text{dom}(s^*) = G$. It has the property

$$s^*(g) = \begin{cases} s(g) & \text{for } g \in \text{dom}(s), \\ 0 & \text{for } g \in G \setminus \text{dom}(s). \end{cases}$$

Thus, $\text{supp}(s^*) = \text{dom}(s) \setminus s^{-1}(0)$. Moreover, $\text{supp}(s^*) = \text{supp}(s)$.

- (iii) We set a^* to be the **linear recurrence sequence associated** to a with domain $\text{supp}(a)$ mapping each $g \in \text{supp}(a)$ to $a(g)$. Note that a^* is 0-free and $\text{supp}(a^*) = \text{supp}(a)$.

- (iv) We write s^{**} for $(s^*)^*$ and a^{**} for $(a^*)^*$. Note that $a^{**} = a$. Note further that s^{**} is the restriction of s to the subset $\text{supp}(s)$ of $\text{dom}(s)$. Thus, s^{**} is the largest 0-free linear recurrence subsequence of s .

Definition 2.5. Let $s = (g_i, s_i)_{i < \alpha} \in \text{LRS}(k, G)$. We define $\langle s \rangle$ to be the set of all power series $a \in k((G))$ such that for any

$$h \in G \setminus \text{dom}(s)$$

the following (generalised) linear recurrence relation holds:

$$(2.1) \quad \sum_{i < \alpha} s(g_i)a_{h-g_i} = 0.$$

(Note that $h - g_i$ is strictly decreasing, whence the sum in (2.1) is finite.)

We say that s **determines** $\langle s \rangle$. For a set $S \subseteq \text{LRS}(k, G)$ we set

$$\langle S \rangle = \bigcup_{s \in S} \langle s \rangle$$

and say that S determines $\langle S \rangle$.

For a set $A \subseteq k((G))$ we say that A is **determined** by linear recurrence relations if there exists a set S of linear recurrence sequences such that $\langle S \rangle = A$.

Example 2.6. For any trivial linear recurrence sequence s in $k((G))$ we have $\langle s \rangle = k((G))$. Hence, any $S \subseteq \text{LRS}(k, G)$ containing a trivial linear recurrence sequence determines $k((G))$. \square

For any non-trivial linear recurrence sequence s , its associated power-series s^* is non-zero. In this case, the following lemma gives a description of the set determined by s .

Lemma 2.7 (Main Lemma). *Let $s \in \text{LRS}(k, G)$ be non-trivial. Then*

$$(2.2) \quad \langle s \rangle = \left\{ \frac{b}{s^*} \mid b \in k((G)), \text{supp}(b) \subseteq \text{dom}(s) \right\}.$$

Proof. Set $s = (g_i, s_i)_{i < \alpha}$. Let $b \in k((G))$ with $\text{supp}(b) \subseteq \text{dom}(s)$ and set $a = \frac{b}{s^*}$. Further, let $h \in G \setminus \text{dom}(s)$. Then

$$\sum_{i < \alpha} s(g_i) a_{h-g_i} = (s^* a)_h = b_h = 0,$$

as $h \notin \text{dom}(s) \supseteq \text{supp}(b)$. Hence, $a \in \langle s \rangle$, showing that

$$\langle s \rangle \supseteq \left\{ \frac{b}{s^*} \mid b \in k((G)), \text{supp}(b) \subseteq \text{dom}(s) \right\}.$$

For the converse, let $a \in k((G))$ with $a \in \langle s \rangle$. Set

$$b_h := (s^* a)_h = \sum_{i < \alpha} s(g_i) a_{h-g_i}$$

for any $h \in G$. Then for any $h \in G \setminus \text{dom}(s)$ we have $b_h = 0$, as $a \in \langle s \rangle$. Hence, $\text{supp}(b) \subseteq \text{dom}(s)$. Since for any $h \in G$ we have $b_h = (s^* a)_h$, we obtain $b = s^* a$. Hence, $a = \frac{b}{s^*}$, as required. \square

Since (2.2) is a useful and important characterisations of sets determined by a single non-trivial linear recurrence sequence s , we will use it throughout this work without referring to Lemma 2.7 each time.

Proposition 2.8. *Let $s \in \text{LRS}(k, G)$. Then $\langle s \rangle \subseteq k((G))$ is a vector space over k containing k .*

Proof. If s is trivial, then $\langle s \rangle = k((G))$, which is even a field extension of k . Otherwise, for any $a, b \in k((G))$ with $\text{supp}(a), \text{supp}(b) \subseteq \text{dom}(s)$ and any $\lambda \in k$ we have

$$\frac{a}{s^*} + \lambda \cdot \frac{b}{s^*} = \frac{a + \lambda b}{s^*}.$$

Now $\text{supp}(a + \lambda b) \subseteq \text{supp}(a) \cup \text{supp}(b) \subseteq \text{dom}(s)$. Hence, also $\frac{a + \lambda b}{s^*}$ lies in $\langle s \rangle$.

Finally, note that $\text{supp}(s^*) \subseteq \text{dom}(s)$, as $\text{supp}(s^*) = \text{dom}(s) \setminus s^{-1}(0)$. Hence, $1 = \frac{s^*}{s^*} \in \langle s \rangle$. Since $\langle s \rangle$ is closed under multiplication with elements from k , we obtain $k \subseteq \langle s \rangle$. \square

While Proposition 2.8 shows that a single linear recurrence sequence determines a k -vector subspace of $k((G))$, a non-empty set S of linear recurrence sequences does not even necessarily determine a subset of $k((G))$ that is closed under addition, as the following example shows.

Example 2.9. Consider the power series $a = 1 + t$ and $b = 1 - t$ in $\mathbb{Q}((\mathbb{Z}))$ and let $S = \{a^*, b^*\}$ be the set of associated linear recurrence sequences. Then

$$\begin{aligned} \langle S \rangle &= \left\{ \frac{c}{a} \mid c \in \mathbb{Q}((\mathbb{Z})), \text{supp}(c) \subseteq \{0, 1\} \right\} \cup \left\{ \frac{d}{b} \mid d \in \mathbb{Q}((\mathbb{Z})), \text{supp}(d) \subseteq \{0, 1\} \right\} \\ &= \left\{ \frac{c_0 + c_1 t}{1+t} \mid c_0, c_1 \in \mathbb{Q} \right\} \cup \left\{ \frac{d_0 + d_1 t}{1-t} \mid d_0, d_1 \in \mathbb{Q} \right\}. \end{aligned}$$

Thus, we have $\frac{1}{1+t}, \frac{1}{1-t} \in \langle S \rangle$. However,

$$\frac{1}{1+t} + \frac{1}{1-t} = \frac{2}{1-t^2} \notin \langle S \rangle,$$

as neither the equation $\frac{2}{1-t^2} = \frac{c_0 + c_1 t}{1+t}$ nor the equation $\frac{2}{1-t^2} = \frac{d_0 + d_1 t}{1-t}$ has a solution over \mathbb{Q} . Hence, $\langle S \rangle$ is not closed under addition.

Note that for a similar reason we have $\frac{1}{1+t} \cdot \frac{1}{1-t} = \frac{1}{1-t^2} \notin \langle S \rangle$, whence $\langle S \rangle$ is also not closed under multiplication. \square

We now establish sufficient conditions for a set determined by linear recurrence relations to be closed under addition and multiplication as well as taking multiplicative inverses of non-zero elements.

Lemma 2.10. *Let $S \subseteq \text{LRS}(k, G)$. Suppose that for any $s_1, s_2 \in S$ the following $r \in \text{LRS}(k, G)$ with domain² $\text{dom}(r) = \text{dom}(s_1) + \text{dom}(s_2)$ is also contained in S : for any $g \in \text{dom}(s_1) + \text{dom}(s_2)$ set*

$$r(g) := \begin{cases} (s_1^* \cdot s_2^*)(g) & \text{if } g \in \text{supp}(s_1^* s_2^*), \\ 0 & \text{if } g \in (\text{dom}(s_1) + \text{dom}(s_2)) \setminus \text{supp}(s_1^* s_2^*). \end{cases}$$

Then $\langle S \rangle$ is closed under addition and multiplication.

Proof. Let $s_1, s_2 \in S$, let $a_1 \in \langle s_1 \rangle$ and let $a_2 \in \langle s_2 \rangle$. We may assume that s_1 and s_2 are non-trivial, as otherwise we already have $\langle S \rangle = k((G))$. Fix $b_1, b_2 \in k((G))$ with $a_1 = \frac{b_1}{s_1^*}$ and $a_2 = \frac{b_2}{s_2^*}$, where $\text{supp}(b_1) \subseteq \text{dom}(s_1)$ and $\text{supp}(b_2) \subseteq \text{dom}(s_2)$. Then

$$a_1 + a_2 = \frac{b_1 s_2^* + b_2 s_1^*}{s_1^* s_2^*} \quad \text{and} \quad a_1 a_2 = \frac{b_1 b_2}{s_1^* s_2^*}.$$

Note that by definition of r , we have $r^* = s_1^* s_2^*$, as $\text{supp}(s_1^* s_2^*) \subseteq \text{supp}(s_1^*) + \text{supp}(s_2^*) \subseteq \text{dom}(s_1) + \text{dom}(s_2)$. Thus,

$$a_1 + a_2 = \frac{b_1 s_2^* + b_2 s_1^*}{r^*} \quad \text{and} \quad a_1 a_2 = \frac{b_1 b_2}{r^*}.$$

²Note that $\text{dom}(r)$ is well ordered, as the sum of two well-ordered subsets of G is also well-ordered.

Moreover,

$$\begin{aligned}
\text{supp}(b_1 s_2^* + b_2 s_1^*) &\subseteq \text{supp}(b_1 s_2^*) \cup \text{supp}(b_2 s_1^*) \\
&\subseteq [\text{supp}(b_1) + \text{supp}(s_2^*)] \cup [\text{supp}(b_2) + \text{supp}(s_1^*)] \\
&\subseteq [\text{dom}(s_1) + \text{dom}(s_2)] \cup [\text{dom}(s_2) + \text{dom}(s_1)] \\
&= \text{dom}(r)
\end{aligned}$$

and

$$\text{supp}(b_1 b_2) \subseteq \text{supp}(b_1) + \text{supp}(b_2) \subseteq \text{dom}(s_1) + \text{dom}(s_2) = \text{dom}(r).$$

Hence,

$$a_1 + a_2, a_1 a_2 \in \langle r \rangle \subseteq \langle S \rangle.$$

□

Lemma 2.11. *Let $S \subseteq \text{LRS}(k, G)$. Suppose that for any $s \in S$ any non-trivial $r \in \text{LRS}(k, G)$ with domain $\text{dom}(r) = \text{dom}(s)$ is also contained in S . Then $\langle S \rangle$ is closed under taking multiplicative inverses of non-zero elements.*

Proof. Let $s \in S$. If s is trivial, then $\langle S \rangle = k((G))$ and we are done. Otherwise, let $a \in \langle s \rangle \setminus \{0\}$ and let $b \in k((G))$ with $\text{supp}(b) \subseteq \text{dom}(s)$ and

$$a = \frac{b}{s^*}.$$

Let $r \in S$ be the linear recurrence sequence with domain $\text{dom}(s)$ given by

$$r(g) := \begin{cases} b(g) & \text{for } g \in \text{supp}(b), \\ 0 & \text{for } g \in \text{dom}(s) \setminus \text{supp}(b). \end{cases}$$

Then $r^* = b$. Since $\text{supp}(s^*) \subseteq \text{dom}(s) = \text{dom}(r)$, we obtain

$$a^{-1} = \frac{s^*}{b} = \frac{s^*}{r^*} \in \langle r \rangle \subseteq S,$$

as required. □

Combining Lemma 2.10 and Lemma 2.11, we obtain the following sufficient conditions on a set $S \subseteq \text{LRS}(k, G)$ in order that $\langle S \rangle$ is a subfield of $k((G))$.

Proposition 2.12. *Let $S \subseteq \text{LRS}(k, G)$ be non-empty. Suppose that for any $s_1, s_2 \in S$ every non-trivial linear recurrence sequence in $k((G))$ with domain $\text{dom}(s_1)$ and every non-trivial linear recurrence sequence in $k((G))$ with domain $\text{dom}(s_1) + \text{dom}(s_2)$ are also contained in S . Then $\langle S \rangle$ is a subfield of $k((G))$ containing k .*

Proof. Since S is non-empty, Proposition 2.8 shows that $k \subseteq \langle S \rangle$. Lemma 2.11 immediately implies that $\langle S \rangle$ is closed under taking multiplicative inverses of non-zero elements. If S contains a trivial sequence, then $\langle S \rangle = k((G))$. Otherwise, the sequence r with $\text{dom}(r) = \text{dom}(s_1) + \text{dom}(s_2)$ in Lemma 2.10 is non-trivial for any $s_1, s_2 \in S$, whence $\langle S \rangle$ is closed under addition and multiplication. □

3. LINEAR RECURRENCE HAHN FIELDS

Proposition 2.12 enables us to consider specific subfields of $k((G))$ determined by linear recurrence relations. More specifically, if S consists of *all* non-trivial linear recurrence sequences with a particular support, then $\langle S \rangle$ forms a subfield of $k((G))$. We make this precise in the following. Recall that \mathcal{F} always denotes a family of well-ordered subsets of G .

Definition 3.1. We denote by $S(\mathcal{F})$ the set of all non-trivial linear recurrence sequences $s \in \text{LRS}(k, G)$ with $\text{dom}(s) \in \mathcal{F}$. A linear recurrence sequence $s \in S(\mathcal{F})$ is called an \mathcal{F} -**sequence**.

Proposition 2.12 immediately implies the following.

Proposition 3.2. *Suppose that \mathcal{F} is non-empty and closed under sums (i.e. $A, B \in \mathcal{F}$ implies $A+B \in \mathcal{F}$). Then $\langle S(\mathcal{F}) \rangle$ is a subfield of $k((G))$ containing k .*

Proposition 3.3. *Let \mathcal{F} be the family of all finite subsets of G . Then $\langle S(\mathcal{F}) \rangle = k(G)$.*

Proof. By Proposition 3.2, $\langle S(\mathcal{F}) \rangle$ is a field containing k . Let $g \in G$. Then the sequence s with domain $\{0, g\}$ given by $s(0) = 1$ and $s(g) = 0$ is contained in $S(\mathcal{F})$. Hence,

$$t^g = \frac{t^g}{1} = \frac{t^g}{s^*} \in \langle s \rangle \subseteq \langle S(\mathcal{F}) \rangle.$$

This shows that $k(G) \subseteq \langle S(\mathcal{F}) \rangle$.

Conversely, let $a \in \langle S(\mathcal{F}) \rangle$. Then there exist $s \in S(\mathcal{F})$ and $b \in k((G))$ with $\text{supp}(b) \subseteq \text{dom}(s)$ and $a = \frac{b}{s^*}$. Since s has a finite domain, both s^* and b have finite support. Hence, $a \in k(G)$, as required. \square

Proposition 3.3 implies the following.

Corollary 3.4. *Suppose that \mathcal{F} contains all finite subsets of G . Then $k(G) \subseteq \langle S(\mathcal{F}) \rangle$. In particular, if $\langle S(\mathcal{F}) \rangle$ is a field, then it is a Hahn field.*

Corollary 3.4 can be used to construct an example of a Hahn field that is not determined by linear recurrence relations.

Lemma 3.5. *Let $G \neq \{0\}$ and let S be a set of linear recurrence sequences in $k((G))$ such that $k(G) \subsetneq \langle S \rangle$. Then $\langle S \rangle$ is uncountable.*

Proof. Corollary 3.4 implies that S contains a linear recurrence sequence s with infinite domain. If s is trivial, then $\langle S \rangle = k((G))$, which is always uncountable. Otherwise,

$$\left\{ \frac{a}{s^*} \mid a \in k((G)), \text{supp}(a) \subseteq \text{dom}(s) \right\} \subseteq \langle s \rangle \subseteq \langle S \rangle.$$

Hence, for every $B \subseteq \text{dom}(s)$, the set $\langle s \rangle$ contains the power series

$$\frac{\sum_{g \in B} t^g}{s^*}.$$

Thus, $\langle s \rangle$ contains at least $|\mathcal{P}(\text{dom}(s))| = 2^{|\text{dom}(s)|} \geq 2^{\aleph_0}$ elements, whence $\langle S \rangle$ is uncountable. \square

While Proposition 3.3 shows that any minimal Hahn field is determined by linear recurrence relations, and thus also every countable minimal Hahn field such as $\mathbb{Q}(\mathbb{Z})$, any countable Hahn field that is *not* the minimal Hahn field is *not* determined by linear recurrence relations, as Lemma 3.5 shows. For instance, let C be the relative algebraic closure of $\mathbb{Q}(\mathbb{Z})$ inside $\mathbb{Q}(\!(\mathbb{Z})\!)$. Since $\sqrt{1+t} \in C \setminus \mathbb{Q}(\mathbb{Z})$, we have that C is a countable non-minimal Hahn field and thus not determined by linear recurrence relations.

We now come to the generalisation of Corollary 1.3 to fields of generalised power series.

Proposition 3.6. *Let $c = \sum_{g \in G} c_g t^g \in k(\!(G)\!)$. Then $c \in k(G)$ if and only if the following holds: there exist $n \in \mathbb{N}$, finite sets $A = \{g_0, \dots, g_n\} \subseteq G$ with $g_0 < \dots < g_n$ and $B = \{b_0, \dots, b_n\} \subseteq k$, not all b_i equal to 0, such that for any $h \in G \setminus A$ we have*

$$\sum_{j=0}^n b_j c_{h-g_j} = 0.$$

Proof. This is basically a consequence of Proposition 3.3: Let \mathcal{F} be the set of all finite subsets of G . Suppose that $c \in k(G) = \langle S(\mathcal{F}) \rangle$. Then there exist $n \in \mathbb{N}$, $A = \{g_0, \dots, g_n\} \in \mathcal{F}$ and a non-trivial linear recurrence sequence $s = (g_i, b_i)_{i < n+1}$ in $S(\mathcal{F})$ with $c \in \langle s \rangle$. Hence, for any $h \in G \setminus \text{dom}(s) = G \setminus A$ we have

$$\sum_{j=0}^n b_j c_{h-g_j} = 0.$$

Conversely, suppose that there exist $n \in \mathbb{N}$, finite sets $A = \{g_0, \dots, g_n\} \subseteq G$ with $g_0 < \dots < g_n$ and $B = \{b_0, \dots, b_n\} \subseteq k$, not all b_i equal to 0, such that for any $h \in G \setminus A$ we have

$$\sum_{j=0}^n b_j c_{h-g_j} = 0.$$

Then $c \in \langle s \rangle \subseteq \langle S(\mathcal{F}) \rangle = k(G)$ for $s = (g_i, b_i)_{i < n+1} \in S$. □

4. RELATION TO RAYNER FIELDS

Following the terminology of [2, Definition 2.4], we call

$$k(\!(\mathcal{F})\!) = \{a \in k(\!(G)\!) \mid \text{supp}(a) \in \mathcal{F}\}$$

the *k-hull* of \mathcal{F} in $k(\!(G)\!)$.

Not all *k-hulls* are determined by linear recurrence relations: Choose \mathcal{F} such that it does not contain $\{0\}$. Then $k \not\subseteq k(\!(\mathcal{F})\!)$, whence $k(\!(\mathcal{F})\!)$ is not determined by linear recurrence relations by Proposition 2.8. However, under certain conditions on \mathcal{F} , we can always find a set of linear recurrence sequences $S_{\mathcal{F}}$ such that $\langle S_{\mathcal{F}} \rangle = k(\!(\mathcal{F})\!)$.

Notation 4.1. Let $A \in \mathcal{F}$. We denote by \mathcal{F}_A the family consisting of all subsets of $A \cup \{0\}$. Moreover, we let $s_A \in \text{LRS}(k, G)$ with domain $\text{dom}(s) = A \cup \{0\}$ mapping 0 to 1 and everything else to 0, and set $S_{\mathcal{F}} = \{s_A \mid A \in \mathcal{F}\}$.

Note that for any $A \in \mathcal{F}$, we have $s_A^* = 1 \in k((G))$.

Lemma 4.2. *Let $A \subseteq G$ be well-ordered. Then $\langle s_A \rangle = k((\mathcal{F}_A))$.*

Proof. Let $a \in \langle s_A \rangle$. Then there exists $b \in k((G))$ with $\text{supp}(b) \subseteq \text{dom}(s_A) = A \cup \{0\}$ such that $a = \frac{b}{s_A^*} = b$. Hence, $a = b \in k((\mathcal{F}_A))$.

Conversely, let $a \in k((\mathcal{F}_A))$. Then $\text{supp}(a) \in \mathcal{F}_A$ and thus $\text{supp}(a) \subseteq A \cup \{0\} = \text{dom}(s_A)$. Hence, $a = \frac{a}{s_A^*} \in \langle s_A \rangle$. \square

Proposition 4.3. *Suppose that \mathcal{F} is closed under subsets (i.e. $B \subseteq A \in \mathcal{F}$ implies $B \in \mathcal{F}$) and unions with $\{0\}$ (i.e. $A \in \mathcal{F}$ implies $A \cup \{0\} \in \mathcal{F}$). Then*

$$\langle S_{\mathcal{F}} \rangle = k((\mathcal{F})).$$

Proof. We may suppose that \mathcal{F} is non-empty, as otherwise $\langle S_{\mathcal{F}} \rangle = \emptyset = k((\mathcal{F}))$. Let $A \in \mathcal{F}$. Then $\mathcal{F}_A \subseteq \mathcal{F}$. Lemma 4.2 implies $\langle s_A \rangle = k((\mathcal{F}_A)) \subseteq k((\mathcal{F}))$, establishing $\langle S_{\mathcal{F}} \rangle \subseteq k((\mathcal{F}))$. Conversely, let $A \in \mathcal{F}$. Again by Lemma 4.2, we have

$$k((\{A\})) \subseteq k((\mathcal{F}_A)) = \langle s_A \rangle.$$

Hence,

$$k((\mathcal{F})) = \bigcup_{A \in \mathcal{F}} k((\{A\})) \subseteq \bigcup_{A \in \mathcal{F}} \langle s_A \rangle = \langle S_{\mathcal{F}} \rangle.$$

\square

Remark 4.4. In [2, Conditions 2.1], several properties of \mathcal{F} are established such that the k -hull $k((\mathcal{F}))$ forms a distinguished substructure of $k((G))$. If \mathcal{F} satisfies the conditions (S2), (S3) and (S4), then the assumption of Proposition 4.3 on \mathcal{F} holds. Hence, any k -hull $k((\mathcal{F}))$ whose underlying family \mathcal{F} satisfies (S2), (S3) and (S4) is determined by $S_{\mathcal{F}}$. In particular, every Rayner field (see [2, Definition 3.2]) is determined by linear recurrence relations. \square

Not every Hahn field that is determined by linear recurrence relations is also a Rayner field. For instance, Proposition 3.3 shows that $k(G)$ is always determined by linear recurrence relations. However, $k(G)$ is not necessarily a Rayner field, as [9, Corollary 3.3.51] shows for the case $k = \mathbb{Q}$ and $G = \mathbb{Z}$.

Following the notation in [2], we use the convention that for any well-ordered subset $A \subseteq G$ we denote by $k((A))$ all elements from $k((G))$ whose support lies in A .

Example 4.5. Remark 4.4 gives us several examples of distinguished subfields of $k((G))$ that are determined by linear recurrence relations:

- (i) If \mathcal{F} consists of all finite subsets of G , then $\langle S_{\mathcal{F}} \rangle = k[G]$.

- (ii) If \mathcal{F} consists of all well-ordered subsets of $G^{\geq 0}$, then we obtain $\langle S_{\mathcal{F}} \rangle = k((G^{\geq 0}))$.
- (iii) Let κ be an uncountable cardinal. If \mathcal{F} consists of all well-ordered subsets of G whose cardinality is strictly less than κ , then $\langle S_{\mathcal{F}} \rangle = k((G))_{\kappa}$. Likewise, if \mathcal{F} consists of all well-ordered subsets of $G^{\geq 0}$ whose maximal cardinality is strictly less than κ , then $\langle S_{\mathcal{F}} \rangle = k((G^{\geq 0}))_{\kappa}$. See [4] for further details on κ -bounded power series.
- (iv) Note that $k((G^{>0}))$ is not determined by linear recurrence relations, as any set that is determined by linear recurrence relations contains k by Proposition 2.8. \square

Proposition 4.6. *Suppose that $k \neq \mathbb{F}_2$ and that $\{0\} \in \mathcal{F}$. Suppose further that $k((\mathcal{F}))$ is a subgroup of $(k((G)), +)$. Then $k((\mathcal{F})) = \langle S_{\mathcal{F}} \rangle$.*

Proof. This follows immediately from [2, Proposition 3.4] and Remark 4.4. \square

In the last part of this section, we relate $\langle S_{\mathcal{F}} \rangle$ to $\langle S(\mathcal{F}) \rangle$, which was introduced in Definition 3.1. If \mathcal{F} consist of all finite subsets of G , then by Proposition 3.3 we have $\langle S(\mathcal{F}) \rangle = k(G)$, and thus by Example 4.5 (i)

$$\langle S(\mathcal{F}) \rangle = \text{ff}(k[G]) = \text{ff}(k((\mathcal{F}))) = \text{ff}(\langle S_{\mathcal{F}} \rangle).$$

The following result shows that $\langle S(\mathcal{F}) \rangle = \text{ff}(\langle S_{\mathcal{F}} \rangle)$ also holds for a wider class of families \mathcal{F} .

Proposition 4.7. *Let \mathcal{F} contain $\{0\}$ and be closed under subsets, unions and sums, i.e. let \mathcal{F} satisfy (S2), (S3), (S4) and (A2) of [2, Conditions 2.1]. Then $\langle S(\mathcal{F}) \rangle = \text{ff}(\langle S_{\mathcal{F}} \rangle)$.*

Proof. By Proposition 3.2, $\langle S(\mathcal{F}) \rangle$ is a field; by Proposition 4.3 $\langle S_{\mathcal{F}} \rangle = k((\mathcal{F}))$. For the inclusion $\langle S(\mathcal{F}) \rangle \supseteq \text{ff}(\langle S_{\mathcal{F}} \rangle)$, it thus suffices to show that $k((\mathcal{F})) \subseteq \langle S(\mathcal{F}) \rangle$. To this end, let $a \in k((G))$ with $\text{supp}(a) \in \mathcal{F}$. Then $s_{\text{supp}(a)}$ has domain $\text{supp}(a) \cup \{0\}$ and thus this sequence lies in $S(\mathcal{F})$. Hence,

$$\frac{a}{s_{\text{supp}(a)}^*} = \frac{a}{1} \in \langle S(\mathcal{F}) \rangle.$$

For the inclusion $\langle S(\mathcal{F}) \rangle \subseteq \text{ff}(\langle S_{\mathcal{F}} \rangle)$, let $s \in S(\mathcal{F})$ and let $a \in k((G))$ with $\text{supp}(a) \subseteq \text{dom}(s) \in \mathcal{F}$. Since \mathcal{F} is closed under subsets, also $\text{supp}(a), \text{supp}(s) \in \mathcal{F}$. Hence,

$$\frac{a}{s^*} \in \text{ff}(\langle S_{\mathcal{F}} \rangle),$$

as required. \square

Corollary 4.8. *Let \mathcal{F} be a Rayner field family. Then $\langle S(\mathcal{F}) \rangle = \langle S_{\mathcal{F}} \rangle = k((\mathcal{F}))$.*

5. LIFTING PROPERTY

Recall from Section 2 that we call a field K Hahn field if $k(G) \subseteq K \subseteq k((G))$. Recall further that K is endowed with the canonical valuation v . Let $\text{Aut } K$ be the group of field automorphisms of K . We say that σ is **valuation preserving** if, for all $a, b \in K$, we have that $v(a) = v(b)$ implies $v(\sigma(a)) = v(\sigma(b))$. For a characterisation of order preserving automorphisms, in terms of the valuation invariants of K , see [5, Theorem 4.2]. The set of valuation preserving automorphisms of K forms a subgroup of $\text{Aut } K$ that we denote by $v\text{-Aut } K$. Let $\sigma \in v\text{-Aut } K$. We canonically associate to σ the order preserving automorphism $\sigma_G \in o\text{-Aut } G$ given by $\sigma_G(v(a)) = v(\sigma(a))$, for any $a \in K$, as well as the automorphism $\sigma_k \in \text{Aut } k$ given by $\sigma_k(x) = (\sigma(x))_0$, for any $x \in k$. This yields a homomorphism

$$\Phi_K: v\text{-Aut } K \rightarrow \text{Aut } k \times o\text{-Aut } G, \quad \sigma \mapsto (\sigma_k, \sigma_G).$$

Notice that, for the maximal Hahn field $k((G))$, the map $\Phi_{k((G))}$ admits the canonical section:

$$\Psi_c: \text{Aut } k \times o\text{-Aut } G \rightarrow \begin{array}{ccc} v\text{-Aut } k((G)) & & \\ (\sigma_k \times \sigma_G) & \mapsto & \left(\begin{array}{ccc} \Psi_c(\sigma_k, \sigma_G): & k((G)) & \rightarrow & k((G)) \\ & \sum a_g t^g & \mapsto & \sum \sigma_k(a) t^{\sigma_G(g)} \end{array} \right) \end{array}$$

Definition 5.1. We say that a Hahn field K has the **canonical lifting property** if, for all $(\sigma_k, \sigma_G) \in \text{Aut } k \times o\text{-Aut } G$, we have $\Psi_c(\sigma_k, \sigma_G) \in v\text{-Aut } K$, i.e. if $\Psi_{c,K}: (\sigma_k, \sigma_G) \mapsto \Psi_c(\sigma_k, \sigma_G)|_K$ is a section of Φ_K .

For a general and detailed study of the canonical lifting property (and others) we refer the reader to [6, 9]. For easier reference, we report here a result that characterises Rayner fields with the canonical lifting property.

Proposition 5.2 ([6, Proposition 3.6.6]). *Let \mathcal{F} be a Rayner field family and let $K = k((\mathcal{F}))$ be the corresponding Rayner field. Then K has the canonical lifting property if and only if \mathcal{F} is stable under $o\text{-Aut } G$, i.e. if $A \in \mathcal{F}$ and $\tau \in o\text{-Aut } G$, then $\tau(A) \in \mathcal{F}$.*

Let s be a linear recurrence sequence in $k((G))$. We set \tilde{s} to be the linear recurrence sequence with domain $\text{dom}(s)$ with

$$\tilde{s} := \begin{cases} \tilde{s}(g) = s(g) & \text{for } g \in \text{supp}(s), \\ \tilde{s}(g) = 1 & \text{for } g \in \text{dom}(s) \setminus \text{supp}(s). \end{cases}$$

Then $\text{dom}(\tilde{s}) = \text{supp}(\tilde{s}) = \text{dom}(s)$ and $\tilde{s}^{**} = \tilde{s}$.

Definition 5.3. Let $\varphi: k((G)) \rightarrow k((G))$. We define a map $\varphi^*: \text{LRS}(k, G) \mapsto \text{LRS}(k, G)$ as follows: For any $s \in \text{LRS}(k, G)$ we let

$$\text{dom}(\varphi^*(s)) := \text{supp}(\varphi(\tilde{s}^*)).$$

For any $g \in \text{dom}(\varphi^*(s))$, we set

$$\varphi^*(s)(g) := \begin{cases} \varphi(\tilde{s}^*)(g) & \text{if } g \in \text{supp}(\varphi(\tilde{s}^*)), \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 5.4. *Let φ be an automorphism on $k((G))$ that preserves supports, i.e. for any $a, b \in k((G))$ we have*

$$\text{supp}(a) \subseteq \text{supp}(b) \text{ if and only if } \text{supp}(\varphi(a)) \subseteq \text{supp}(\varphi(b)).$$

Then for any linear recurrence sequence s in $k((G))$ we have

$$\varphi(\langle s \rangle) = \langle \varphi^*(s) \rangle.$$

Proof. First let $a \in k((G))$ with $\text{supp}(a) \subseteq \text{dom}(s)$. In order to establish $\varphi(\langle s \rangle) \subseteq \langle \varphi^*(s) \rangle$, we have to show that there exists $b \in k((G))$ with $\text{supp}(b) \subseteq \text{dom}(\varphi^*(s))$ such that

$$\varphi\left(\frac{a}{s^*}\right) = \frac{b}{\varphi^*(s)^*}.$$

Set

$$b = \frac{\varphi(a) \cdot \varphi^*(s)^*}{\varphi(s^*)}.$$

We have to verify that $\text{supp}(b) \subseteq \text{dom}(\varphi^*(s))$.

Claim: $\varphi^*(s)^* = \varphi(s^*)$.

By definition of φ^* we obtain

$$\begin{aligned} \text{supp}(\varphi^*(s)) &= \text{dom}(\varphi^*(s)) \cap \text{supp}(\varphi(s^*)) \\ &= \text{supp}(\varphi(\tilde{s}^*)) \cap \text{supp}(\varphi(s^*)). \end{aligned}$$

Since $\text{supp}(\tilde{s}^*) \supseteq \text{supp}(s^*)$, we obtain

$$\text{supp}(\varphi^*(s)) = \text{supp}(\varphi(s^*)),$$

as φ is support-preserving. Now for any $g \in \text{supp}(\varphi(s^*))$, we have

$$\varphi^*(s)(g) = \varphi(s^*)(g),$$

establishing the claim.

Now since $\text{supp}(a) \subseteq \text{dom}(s) = \text{supp}(\tilde{s}^*)$, we obtain

$$\text{supp}(b) = \text{supp}(\varphi(a)) \subseteq \text{supp}(\varphi(\tilde{s}^*)) = \text{dom}(\varphi^*(s)),$$

as required.

The converse follows by similar arguments: If $b \in k((G))$ with $\text{supp}(b) \subseteq \text{dom}(\varphi^*(s)) = \text{supp}(\varphi(\tilde{s}^*))$, then

$$\text{supp}(\varphi^{-1}(b)) \subseteq \text{supp}(\tilde{s}^*) = \text{dom}(s).$$

Hence, for $a = \varphi^{-1}(b)$ we obtain

$$\frac{b}{\varphi^*(s)^*} = \frac{\varphi(a)}{\varphi(s^*)} = \varphi\left(\frac{a}{s^*}\right) \in \varphi(\langle s \rangle).$$

□

For any automorphism ψ on the ordered abelian group G we denote by $\tilde{\psi} := \Psi_c(\text{id}_k, \psi)$ the canonical lift of ψ to $k((G))$.

Lemma 5.5. *Let S be a set of linear recurrence sequences and let ψ be an automorphism on G . Suppose that for any $s = (g_i, s_i)_{i < \alpha} \in S$ also $s^\psi := (\psi(g_i), s_i)_{i < \alpha} \in S$. Then for any $a \in \langle S \rangle$ we have $\tilde{\psi}(a) \in \langle S \rangle$.*

Proof. Let $s = (g_i, s_i)_{i < \alpha} \in S$ and let $a \in k((G))$ with $\text{supp}(a) \subseteq \text{dom}(s) = \{g_i \mid i < \alpha\}$. Then

$$\text{supp}(\tilde{\psi}(a)) = \psi(\text{supp}(a)) \subseteq \psi(\text{dom}(s)) = \{\psi(g_i) \mid i < \alpha\} = \text{dom}(s^\psi).$$

Moreover,

$$\tilde{\psi}(s^*) = \tilde{\psi}\left(\sum_{i < \alpha} s_i t^{g_i}\right) = \sum_{i < \alpha} s_i t^{\psi(g_i)} = (s^\psi)^*.$$

Hence,

$$\tilde{\psi}\left(\frac{a}{s}\right) = \frac{\tilde{\psi}(a)}{\tilde{\psi}(s^*)} = \frac{\tilde{\psi}(a)}{(s^\psi)^*} \in \langle s^\psi \rangle \subseteq \langle S \rangle.$$

□

Theorem 5.6. *Suppose that \mathcal{F} has the following properties:*

- (i) *Every finite subset of G is contained in \mathcal{F} .*
- (ii) *\mathcal{F} is closed under sums.*
- (iii) *For any $\psi \in o\text{-Aut}(G)$ and any $A \in \mathcal{F}$ we have $\psi(A) \in \mathcal{F}$.*

Then $\langle S(\mathcal{F}) \rangle$ is a Hahn field with the canonical lifting property.

Proof. Let $K = \langle S(\mathcal{F}) \rangle$. The first and second condition ensure that K is a Hahn field (see Proposition 3.2 and Corollary 3.4). By Lemma 5.5, the third condition ensures that K has the canonical lifting property □

While Theorem 5.6 presents a class of Hahn fields with the canonical lifting property that are determined by linear recurrence relations, not every Hahn field determined by linear recurrence relations has the canonical lifting property. In the following, we present a Rayner field – which is determined by linear recurrence relations by Remark 4.4 – that does not have the canonical lifting property.

Lemma 5.7. *Let A_1, \dots, A_m be finitely many subsets of G and let \mathcal{F} consist of all well-ordered subsets of $\langle g_1, \dots, g_n, A_1, \dots, A_m \rangle$, where $g_1, \dots, g_n \in G$. Then \mathcal{F} is a Rayner field family.*

Proof. \mathcal{F} is non-empty as $\{0\} \in \mathcal{F}$, and by its definition it is closed under subsets, finite unions and addition with a singleton $\{g\}$ for any $g \in G$. Since $\{\{g\} \mid g \in G\} \subseteq \mathcal{F}$, we also have $\langle \bigcup_{A \in \mathcal{F}} A \rangle = G$. Finally, let $A \in \mathcal{F}$ with $A \subseteq G^{\geq 0}$. Then $\bigoplus_{n \in \mathbb{N}} A \subseteq \langle A \rangle \subseteq \langle g_1, \dots, g_n, A_1, \dots, A_m \rangle$. □

Example 5.8. Let $G = \prod_{\mathbb{Z}} \mathbb{Q}$ (cf. [4, Section 0.1]) and let $A_1 = \{-\frac{1}{p_i} \mathbb{1}_1\}$, where $p_i \in \mathbb{N}$ denotes the i -th prime number. Let \mathcal{F} consist of all well-ordered subsets of subgroups of G of the form $\langle g_1, \dots, g_n, A_1 \rangle$, where $g_1, \dots, g_n \in G$. By Lemma 5.7, \mathcal{F} is a Rayner field family. Since A_1 is well-ordered, we have $A_1 \in \mathcal{F}$. Now consider the automorphism on G induced by $\mathbb{1}_n \mapsto \mathbb{1}_{n-1}$. Then $\tau(A_1) = \{-\frac{1}{p_i} \mathbb{1}_0\}$. We show that $\tau(A_1)$ cannot be a subset of a subgroup of G of the form $\langle g_1, \dots, g_n, A_1 \rangle$, whence $\tau(A_1) \notin \mathcal{F}$.

To this end, let $g_1, \dots, g_n \in G$ be arbitrary and let $h \in \langle g_1, \dots, g_n, A_1 \rangle$. Then there exist $a_1, \dots, a_\ell \in A_1$ and $r_1, \dots, r_n, s_1, \dots, s_\ell \in \mathbb{Z}$ such that

$$h = r_1 g_1 + \dots + r_n g_n + s_1 a_1 + \dots + s_\ell a_\ell.$$

Evaluating at 0, we obtain

$$h(0) = r_1 g_1(0) + \dots + r_n g_n(0),$$

as $a_i(0) = 0$ for any $i \in \{1, \dots, \ell\}$. Let q_1, \dots, q_m be all primes dividing at least one of the denominators of $g_1(0), \dots, g_n(0) \in \mathbb{Q}$ expressed in their reduced fraction form and let $p \in \mathbb{N}$ be a prime number such that $p > q_i$ for all $i \in \{1, \dots, m\}$. Then $g = -\frac{1}{p} \mathbf{1}_0 \in \tau(A_1)$. Hence, $g(0) = -\frac{1}{p}$. However, the denominator of $r_1 g_1(0) + \dots + r_n g_n(0)$ in reduced fraction form is not divisible by p . Hence, $g(0)$ cannot be expressed as a \mathbb{Z} -linear combination of $g_1(0), \dots, g_n(0)$. This shows that $g \notin \langle g_1, \dots, g_n, A_1 \rangle$, as required. \square

REFERENCES

- [1] H. HAHN, ‘Über die nichtarchimedischen Grössensysteme’, *Kais. Akad. Wiss. Wien, S.-B. IIa* 116 (1907) 601–655.
- [2] L. S. KRAPP, S. KUHLMANN and M. SERRA, ‘On Rayner structures’, *Comm. Algebra* 50 (2022) 940–948, doi:10.1080/00927872.2021.1976789.
- [3] S. KUHLMANN, *Ordered Exponential Fields*, Fields Inst. Monogr. 12 (Amer. Math. Soc., Providence, RI, 2000), doi:10.1090/fim/012.
- [4] S. KUHLMANN and S. SHELAH, ‘ κ -bounded exponential-logarithmic power series fields’, *Ann. Pure Appl. Logic* 136 (2005) 284–296, doi:10.1016/j.apal.2005.04.001.
- [5] S. KUHLMANN, M. MATUSINSKI and F. POINT, ‘The Valuation Difference Rank of a Quasi-Ordered Difference Field’, *Groups, Modules, and Model Theory – Surveys and Recent Developments* (eds M. Droste, L. Fuchs, B. Goldsmith and L. Strüngmann; Springer, Cham, 2017) 399–414, doi:10.1007/978-3-319-51718-6_23.
- [6] S. KUHLMANN and M. SERRA, ‘The automorphism group of a valued field of generalised power series’, *J. Algebra* 605 (2022) 339–376, doi:10.1016/j.jalgebra.2022.04.023.
- [7] B. H. NEUMANN, ‘On ordered division rings’, *Trans. Amer. Math. Soc.* 66 (1949), 202–252, doi:10.1090/S0002-9947-1949-0032593-5.
- [8] P. RIBENBOIM, *The Theory of Classical Valuations*, Springer Monogr. Math. (Springer, New York, 1999).
- [9] M. SERRA, ‘Automorphism groups of Hahn groups and Hahn fields’, Doctoral Thesis, Universität Konstanz, 2021.

FACHBEREICH MATHEMATIK UND STATISTIK, UNIVERSITÄT KONSTANZ, 78457 KONSTANZ, GERMANY

Email address: `sebastian.krapp@uni-konstanz.de`

FACHBEREICH MATHEMATIK UND STATISTIK, UNIVERSITÄT KONSTANZ, 78457 KONSTANZ, GERMANY

Email address: `salma.kuhlmann@uni-konstanz.de`

FACHBEREICH MATHEMATIK UND STATISTIK, UNIVERSITÄT KONSTANZ, 78457 KONSTANZ, GERMANY

Email address: `michele.serra@uni-konstanz.de`