

UNKNOTTING NUMBER AND CABLING

JENNIFER HOM, TYE LIDMAN, AND JUNGHWAN PARK

ABSTRACT. The unknotting number of knots is a difficult quantity to compute, and even its behavior under basic satelliting operations is not understood. We establish a lower bound on the unknotting number of cable knots and iterated cable knots purely in terms of the winding number of the pattern. The proof uses Alishahi-Eftekhary's bounds on unknotting number from knot Floer homology together with Hanselman-Watson's computation of the knot Floer homology of cables in terms of immersed curves in the punctured torus.

1. INTRODUCTION

The unknotting number $u(K)$ of a knot K is one of the simplest knot invariants, yet not much is known about its properties. For instance, we do not know how the unknotting number changes with respect to the connected sum. Scharlemann [Sch85] proved that composite knots have unknotting number at least 2, and there are some other partial known results e.g. [Yan08, AE20], but the problem remains very much open.

The connected sum operation is one of the simplest satellite constructions, and we know just as little about most other satellite operations, apart from special cases like Whitehead doubling. In this article, we study the behavior of the unknotting number under another simple satellite operation, *cabling*. Let $K_{p,q}$ denote the (p, q) -cable of K , where p denotes the longitudinal winding, and let $K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}$ denote the iterated cable of K . We may assume that $p > 1$ and $p_i > 1$ since $K_{p,q}$ is isotopic to $K_{-p, -q}$ and $K_{1,q}$ is isotopic to K .

Theorem 1.1. *If K is a nontrivial knot, then $u(K_{p,q}) \geq p$ and $u(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq p_1 p_2 \dots p_m$.*

We make a remark that it was previously shown by Scharlemann-Thompson [ST88, Corollary 3.3] that $u(K_{p,q}) \geq 2$ for any nontrivial knot K . In fact, they showed that a satellite knot of nonzero winding number in the pattern has unknotting number at least 2 (in particular, it recovers the result of Scharlemann [Sch85] mentioned above).

When K is a torus knot T_{p_0, q_0} , we have the following stronger statement. Since $u(K) = u(-K)$ where $-K$ is the reverse of the mirror image of a knot K , we assume that T_{p_0, q_0} is a positive torus knot.

Theorem 1.2. *If $1 < p_0 < q_0$ and $q_0 \neq 3$, then $u(T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}) \geq p_0 p_1 \dots p_m - 1$.*

The main tool we use to establish these lower bounds is the torsion order of a knot (see Definition 2.3) coming from knot Floer homology [OS04b, Ras03]. The torsion order of a knot K is denoted by $\text{Ord}(K)$ and has many topological applications. In particular, Alishahi-Eftekhary [AE20, Theorem 1.1] (see also [Zem19]) showed that if K is a knot, then

$$u(K) \geq \text{Ord}(K).$$

We quickly sketch the idea of the proof. The knot Floer homology of a knot can be recast in the immersed curves interpretation [HRW16, HRW18] of bordered Floer homology [LOT18]. The bordered Floer homology of the exterior of a knot K in S^3 can be thought of roughly as an immersed multicurve in the boundary of the knot exterior (with a single point removed). The torsion order can be morally thought of as the maximal number of times this multicurve wraps consecutively in the direction of the meridian of the knot. Hanselman-Watson [HW19] gives a direct topological formula for transforming the immersed curve for K into the immersed curve for $K_{p,q}$,

The first author was partially supported by NSF grants DMS-1552285 and DMS-2104144.

The second author was partially supported by NSF grants DMS-1709702 and DMS-2105469 and a Sloan Fellowship.

The third author was partially supported by Samsung Science and Technology Foundation (SSTF-BA2102-02) and the POSCO TJ Park Science Fellowship.

and we will show the winding number loosely translates into how much additional wrapping is introduced to the curves.

Using the above strategy, we will obtain the following more general bound on $\text{Ord}(K)$. The inequality itself may be of independent interest, and a similar computation for a certain family of knots can be found in [HKP20, Proposition 1.2].

Proposition 1.3. *If K is a nontrivial knot, then*

$$\text{Ord}(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq \max \{p_1 p_2 \dots p_m (\text{Ord}(K) - 1) + 1, p_1 p_2 \dots p_m\}.$$

As above, when K is a torus knot, or more generally an L -space knot, we have the following stronger statement. Note that $\text{Ord}(K) = \text{Ord}(-K)$, since the knot Floer homology of $-K$ is the dual of the knot Floer homology of K [OS06, Section 5.1].

Proposition 1.4. *If K is a nontrivial L -space knot and $K \neq T_{2,3}$, then*

$$\text{Ord}(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq p_1 \dots p_m (\text{Ord}(K) + 1) - 1.$$

Moreover, if the bridge index of K is $\text{Ord}(K) + 1$, then the above inequality is an equality.

1.1. More applications. As mentioned earlier, the torsion order of a knot has many topological applications. We take advantage of this fact and further explore the consequences of Proposition 1.3 and Proposition 1.4. Throughout the article, we will always work in the smooth category.

A knot is *slice* if it bounds a smoothly embedded disk in B^4 . Note that the $(p, 1)$ -cable of a slice knot is also slice since there is a disk for $K_{p,1}$ constructed by taking p parallel copies of a disk bounded by K and attaching $p - 1$ half-twisted bands to them. In this construction, each parallel copy of a disk for K gives at least one local minima for the resulting disk for $K_{p,1}$. Surprisingly, we show that *any* disk bounded by $K_{p,1}$ has at least $p + 1$ local minima given that K is a nontrivial slice knot. This is done by combining Proposition 1.3 with a bound of Juhász-Miller-Zemke [JMZ20, Corollary 1.7].

Corollary 1.5. *If K is a nontrivial slice knot, then any slice disk bounded by $K_{p_1, 1; p_2, 1; \dots; p_m, 1}$ has at least $p_1 p_2 \dots p_m + 1$ local minima with respect to the radial function on B^4 restricted to the disk.*

We make a remark that it was previously shown in [HKP20, Theorem 1.1] that if a nontrivial knot K bounds a disk with 2 local minima, then any disk bounded by $K_{p,1}$ has at least $p + 1$ local minima (in fact, they also show that there is a disk for $K_{p,1}$ with $p + 1$ local minima).

We also provide applications to concordance. Two knots K and J are *concordant* if $K \# -J$ is slice, and the set of concordance classes of knots form a group called the *knot concordance group* \mathcal{C} under the operation of connected sum. Every element in \mathcal{C} is represented by infinitely many knots, and it is natural to ask which knot is the simplest representative of the class. There are many natural ways to measure the complexity of a knot, and it turns out that the torus knots, and more generally algebraic knots, those that are one component links of isolated singularities of complex algebraic plane curves, have the minimal complexity among all the knots that are concordant to them with respect to various geometric knot invariants.

For instance, the resolution of the local Thom conjecture by Kronheimer-Mrowka [KM93, Corollary 1.3] implies that each algebraic knot minimizes both the 3-genus and the unknotting number in this sense. Moreover, Juhász-Miller-Zemke [JMZ20, Corollary 1.10] proved that this is also the case for torus knots with respect to the braid index and the bridge index (see also [FK17, Theorem 1.3]). This was done by finding a lower bound for the braid index and the bridge index in terms of the torsion order [JMZ20, Corollary 1.8]. Using Proposition 1.4, we extend their result to a large family of iterated torus knots including all algebraic knots. For a knot K , its bridge index is denoted by $\text{br}(K)$ and its braid index is denoted by $\text{b}(K)$. Again, since $\text{b}(K) = \text{b}(-K)$ and $\text{br}(K) = \text{br}(-K)$, we assume that T_{p_0, q_0} is a positive torus knot.

Corollary 1.6. *Let K be either an algebraic knot or an iterated torus knot $T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}$ where $T_{p_0, q_0} \neq T_{2,3}$. If J is concordant to K , then $\text{b}(J) \geq \text{b}(K)$ and $\text{br}(J) \geq \text{br}(K)$.*

We believe that the statement should also be true even when $T_{p_0, q_0} = T_{2,3}$, though one would need to use a different method than ours to prove this since there are examples where the torsion order does not give an optimal bound for the bridge number (e.g., $\text{Ord}(T_{2,3; 2,3}) = 2$ whereas $\text{br}(T_{2,3; 2,3}) - 1 = 3$).

We provide one more 4-dimensional application. Recall that the 4-genus of a knot K , denoted by $g_4(K)$, is defined to be the minimal genus of an oriented smooth surface properly embedded in B^4 bounded by K . It is well known that the 4-genus gives a lower bound for the unknotting number, and the two invariants coincide for the closures of positive braids [Sto03]. Hence for a large family of iterated torus knots, including algebraic knots, the 4-genus is equal to the unknotting number. This is not true for all iterated torus knots. In fact, we show the following.

Corollary 1.7. *There exists a family of iterated torus knots $\{K_i\}$, such that the gap between $u(K_i)$ and $g_4(K_i)$ is arbitrarily large.*

Lastly, we give an application to the untwisting number, a generalization of the unknotting number. A *null-homologous twist* is an operation on a knot, where we perform 1 or -1 surgery on an unknotted curve that is null-homologous in the knot complement. The *untwisting number* of a knot K , denoted by $u_t(K)$, is defined to be the minimum number of null-homologous twists required to convert K to the unknot (for related work see e.g. [MD88, Inc16, Inc17, Liv21, McC21b, McC21a]). By definition, it is clear that for any nontrivial knot K , we have

$$u(K) \geq u_t(K) \geq 1.$$

In [Inc16, Theorem 1.3], it was shown that if K is an unknotting number one knot with $\tau(K) > 0$, where τ is the concordance invariant of [OS03], then $u(K_{p,1}) \geq p$ and $u_t(K_{p,1}) = 1$. The fact that τ gives a lower bound for the unknotting number [OS03] and a cabling formula for τ of [Hom14a, Theorem 1] imply that $u(K_{p,1}) \geq p$. The equality $u_t(K_{p,1}) = 1$ follows easily and is left to the reader. Theorem 1.1 directly implies that the above result generalizes to all unknotting number one knots:

Corollary 1.8. *If K is an unknotting number one knot, then $u(K_{p,1}) \geq p$ and $u_t(K_{p,1}) = 1$. □*

1.2. Some open problems. We end this section with a few open problems on the unknotting number. Given a nontrivial knot K , one easily obtains a crude upper bound on $u(K_{p,q})$ as follows. First, choose an unknotting sequence for K . For each crossing change for K , perform the corresponding p^2 crossing changes on $K_{p,q}$. Then the resulting knot is a torus knot $T_{p,q'}$ where q' depends on the unknotting number of K and also the sign of the crossing changes that were performed to unknot K . By further performing the minimal number of crossing changes needed to turn $T_{p,q'}$ into the unknot, we obtain a bound.

For instance, if K is a knot that can be turned into the unknot by changing a positive crossing to a negative crossing (e.g., $T_{2,3}$), then this crude bound tells us that $u(K_{2,1}) \leq 5$. Similarly, if K turns into the unknot by changing a negative crossing to a positive crossing, then we have $u(K_{2,1}) \leq 6$. Unfortunately, in either case these bounds are far from sharp. For instance, it is a straightforward exercise to find a sequence of crossing changes for $T_{2,3;2,1}$ and show that $u(T_{2,3;2,1}) \leq 3$. Combined with Theorem 1.1 (or [ST88]), we see that $u(T_{2,3;2,1})$ is either 2 or 3. To our surprise, it seems that the unknotting number of $T_{2,3;2,1}$ is not known. We ask the following questions.

Question 1.9. *Is there a nontrivial knot K where we know the value of $u(K_{2,1})$? More generally, is there a formula for the unknotting number of cabled knots?*

Finally, we remark that it is likely that the ideas in this paper can be generalized to other satellite operations. In particular, work of Chen [Che19] describes the knot Floer homology of satellite knots with (1,1)-patterns in the solid torus in terms of immersed curves similar to the work of Hanselman-Watson [HW19] for cables used here. It would be interesting to apply similar techniques to obtain bounds on the unknotting number of such satellite knots. Furthermore, it is natural to ask how the winding number of a pattern governs the unknotting number of a satellite knot. In line with Scharlemann-Thompson's result that a nontrivial satellite of nonzero winding number has unknotting number at least 2 [ST88], we pose

Conjecture 1.10. *Suppose that K is a satellite of a nontrivial knot, where the pattern has winding number p . Then, $u(K) \geq p + 1$.*

Using Theorem 1.1, we are able to give further evidence for Conjecture 1.10 being true using Theorem 1.1. An analogous statement for the Levine-Tristram signature [Lev69, Tri69] can be proved using [Lit84, Theorem 1].

Corollary 1.11. *Let P be a nontrivial pattern with winding number $p \geq 1$. There exists a constant C_P such that for any companion K with $\text{Ord}(K) \geq C_P$, we have $u(P(K)) \geq p + 1$.*

Proof. Let $C_{p,1}$ be the $(p, 1)$ -cabling pattern in the solid torus. Choose a sequence of c crossing changes in the solid torus taking $C_{p,1}$ to P . Therefore, $u(K_{p,1}) \leq u(P(K)) + c$. Moreover, if $u(K_{p,1}) \geq c + p + 1$, then $u(P(K)) \geq p + 1$. The result now follows from Proposition 1.3. \square

ORGANIZATION

In Section 2, we review and develop the relevant connections between knot Floer complexes and the immersed curve machinery of Hanselman-J. Rasmussen-Watson. In Section 3, we use the immersed curve description of cabling due to Hanselman-Watson to prove Propositions 1.3 and 1.4. Lastly, in Section 4, we prove the applications of the propositions including Theorems 1.1 and 1.2.

ACKNOWLEDGEMENTS

We would like to thank Peter Feller for helpful conversations.

2. KNOT-LIKE COMPLEXES

We will assume that the reader is familiar with knot Floer homology [OS04b, Ras03]. We will view our knot Floer complexes as chain complexes over the ring $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ or $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$. See [Hom20, Section 3] for an expository overview. Chain complexes over the latter ring have an interpretation as immersed curves in the punctured torus. The goal of this section is to describe this relationship, which is described in [HRW18, Section 4] (see also [KWZ20]), and to prove a technical result obstructing certain types of curves from being realized by such chain complexes.

2.1. Modules over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$ and immersed curves. We will consider the bigraded ring $\mathbb{F}[\mathcal{U}, \mathcal{V}]$, where $\text{gr}(\mathcal{U}) = (-2, 0)$ and $\text{gr}(\mathcal{V}) = (0, -2)$. We refer to the first component of gr as $\text{gr}_{\mathcal{U}}$ and the second component as $\text{gr}_{\mathcal{V}}$.

Every chain complex C over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ is homotopy equivalent to one that is *reduced*; that is, that $\partial x = 0 \pmod{(\mathcal{U}, \mathcal{V})}$ for all $x \in C$. Without loss of generality, we will assume throughout that our chain complexes are reduced.

Definition 2.1. A *knot-like complex* C over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ is a free, finitely generated, bigraded chain complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ such that

- (1) $H_*(C/\mathcal{U})/\mathcal{V}$ -torsion $\cong \mathbb{F}[\mathcal{V}]$ where $\text{gr}_{\mathcal{U}}(1) = 0$,
- (2) $H_*(C/\mathcal{V})/\mathcal{U}$ -torsion $\cong \mathbb{F}[\mathcal{U}]$ where $\text{gr}_{\mathcal{V}}(1) = 0$.

Remark 2.2. Similarly, we may define a *knot-like complex* C over the ring $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$ (see [DHST21, Definition 3.1]). In the present paper, we work over both rings, as Proposition 2.8 below relies on the existence of a nonzero term in the image of $\mathcal{U}\mathcal{V}$.

Knot-like complexes arise as the knot Floer complexes of knots in S^3 . That is, if K is a knot in S^3 , then the knot Floer complex $CFK(K)$ is a knot-like complex. See, for example, [Hom20, Section 3.2].

We let $H^-(C)$ denote $H_*(C/\mathcal{V})$. Note that when $C = CFK(K)$, we have that $H^-(C) = HFK^-(K)$. The Alexander grading is given by $\frac{1}{2}(\text{gr}_{\mathcal{U}} - \text{gr}_{\mathcal{V}})$.

Definition 2.3. The *torsion order* of a knot-like complex C is

$$\text{Ord}(C) = \min\{n \mid \mathcal{U}^n \cdot \text{Tor}_{\mathcal{U}} H^-(C) = 0\}.$$

The *torsion order* of a knot K is $\text{Ord}(K) = \text{Ord}(CFK(K))$, where $CFK(K)$ is the knot Floer complex of K .

Theorem 2.4 (Theorem 1.1 of [AE20]). *The torsion order of a knot K provides a lower bound on the unknotting number of K :*

$$u(K) \geq \text{Ord}(K). \tag{2.1}$$

Let $\{z_i\}$ be a basis over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ for a knot-like complex C . Let $\langle \partial z_i, z_j \rangle$ denote the coefficient of z_j in ∂z_i . If $\langle \partial z_i, z_j \rangle = \mathcal{U}^n$, then we say that there is a *horizontal arrow of length n* from z_i to z_j . If $\langle \partial z_i, z_j \rangle = \mathcal{V}^n$, then we say that there is a *vertical arrow of length n* from z_i to z_j . If $\langle \partial z_i, z_j \rangle \in \text{Im}(\mathcal{U}\mathcal{V})$, then we say there is a *diagonal arrow* from z_i to z_j . (Observe that $\langle \partial z_i, z_j \rangle$ must be a monomial in \mathcal{U} and \mathcal{V} for grading reasons.)

We now consider certain particularly nice bases for knot-like complexes. A basis $\{x_i\}_{i=0}^{2N}$ for a knot-like complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ is called *horizontally simplified* if $\partial x_i = 0 \pmod{\mathcal{V}}$ for i even and $\partial x_i = \mathcal{U}^{n_i} x_{i+1} \pmod{\mathcal{V}}$ for i odd. See Figure 1 for a graphical depiction.




FIGURE 1. A graphical depiction of a horizontally simplified basis modulo (\mathcal{V}) .

Similarly, we say that a basis $\{y_i\}_{i=0}^{2N}$ for a knot-like complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ is *vertically simplified* if $\partial y_i = 0 \pmod{\mathcal{U}}$ for i even and $\partial y_i = \mathcal{V}^{m_i} y_{i+1} \pmod{\mathcal{U}}$ for i odd. A horizontally simplified basis and a vertically simplified basis for a knot-like complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$ is defined in a similar way.

Remark 2.5. Note that if $\{y_i\}$ is a vertically simplified basis and j is odd, then $\langle \partial y_i, y_j \rangle = 0 \pmod{\mathcal{U}}$ for all i .

We can always choose a vertically simplified basis for C , or, if we prefer, a horizontally simplified basis for C ; see, for example, [LOT18, Proposition 11.57].

In many situations, it suffices to work with chain complexes over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$ rather than $\mathbb{F}[\mathcal{U}, \mathcal{V}]$. We now describe the relationship between chain complexes over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$ and immersed curves. Our immersed curves will sit in $(\mathbb{R}/\mathbb{Z}) \times \mathbb{R}$ with punctures at each lattice point $(0, n + \frac{1}{2})$. The punctures may be thought of as pegs around which the immersed curves wind. For convenience, we identify $(\mathbb{R}/\mathbb{Z}) \times \mathbb{R}$ with $([-\frac{1}{2}, \frac{1}{2}] / -\frac{1}{2} \sim \frac{1}{2}) \times \mathbb{R}$ and work with the latter.

Let $\{x_i\}$ be a horizontally simplified basis for a knot-like complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$. Each basis element x_i corresponds to a point on the vertical line $\{0\} \times \mathbb{R}$; we abuse notation and also denote this point by x_i . The vertical coordinate of x_i is given by the Alexander grading of x_i . A horizontal arrow of length n_i from x_i to x_{i+1} , i odd, gives rise to an arc of length n_i from x_i to x_{i+1} to the right of the vertical line $\{0\} \times \mathbb{R}$, where x_i is n_i units below x_{i+1} . We call such an arc a *right arc of length n_i* . We also include an arc from x_0 to $(\frac{1}{2}, 0)$, which we call the *essential right arc*.

Similarly, if $\{y_i\}$ is a vertically simplified basis for a knot-like complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$, each basis element y_i corresponds to a point on the vertical line $\{0\} \times \mathbb{R}$; we again abuse notation and denote the point by y_i . The vertical coordinate of y_i is again given by the Alexander grading of y_i . A vertical arrow of length m_i from y_i to y_{i+1} , i odd, gives rise to an arc of length m_i from y_i to y_{i+1} to the left of the vertical line $\{0\} \times \mathbb{R}$, where now y_i is m_i units above y_{i+1} . We call such an arc a *left arc of length m_i* . We also include an arc from y_0 to $(-\frac{1}{2}, 0)$, which we call the *essential left arc*.

Thus, given a knot-like complex with a simultaneously horizontally and vertically simplified basis, it is straightforward to find an associated immersed curve, which will be a concatenation of right and left arcs of various lengths, together with the essential arcs. See Figure 2 for some examples.

In the absence of a simultaneously horizontally and vertically simplified basis, one obtains an immersed curve together with a nontrivial local system. (Note that there exist knot-like complexes that do not admit a simultaneously horizontally and vertically simplified basis; see, for example, [Hom15, Figure 3].) As noted at the end of Section 1.1 of [HRW18], a local system can be expanded by replacing the associated curve with parallel copies together with crossover arrows, which can be confined to a prescribed part of the curve. Note that if the crossover arrows are confined to a right arc, then the corresponding basis is vertically simplified.

We denote the component of the immersed curve containing the essential right arc (and hence also the essential left arc) γ_0 . As noted on page 2 of [HW19], for an immersed curve associated to the complement of a knot in S^3 , the component γ_0 always carries the trivial 1-dimensional local system (otherwise the rank of \widehat{HF} of the meridional filling of K would be greater than one; compare with items (1) and (2) of Definition 2.1). We call the




FIGURE 2. Examples of chain complexes over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ and their associated immersed curves.

right arc of γ_0 containing y_0 the *initial right arc*. Note that the initial right arc abuts the essential left arc. See Figure 2(d).

Conversely, an immersed curve (possibly with a nontrivial local system) gives a chain complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}\mathcal{V})$, essentially by reversing the above procedure; see [KWZ20, Theorem 2]. If the immersed curve is the bordered invariant associated to the complement of a knot in S^3 , then the associated chain complex is the mod $(\mathcal{U}\mathcal{V})$ reduction of a knot-like complex. Each intersection point of the immersed curve with the vertical axis yields a basis element z_i , and we have a horizontal (resp. vertical) arrow of length n from z_i to z_j exactly when there is an upwards right arc (resp. downwards left arc) of length n from z_i to z_j . Here, we do not give the immersed curve an orientation; we only look locally at an arc and consider right (resp. left) arcs as upwards (resp. downwards).

Remark 2.6. Note that under this correspondence between immersed curves and chain complexes, if we restrict to the component γ_0 (which always carries the trivial 1-dimensional local system), we may assume that our basis is simultaneously vertically and horizontally simplified.

Suppose one is given a right arc which is either noninitial or initial such that y_0 is not contained in $\{0\} \times [-\frac{1}{2}, \frac{1}{2}]$. Observe that given a right arc of length n , there are two options for the behavior at each end of the arc: it may turn up or down. These four possibilities are illustrated in Figure 3, and denoted $\eta_n^{\pm\pm}$, where the first superscript describes the behavior of the top end, the second superscript describes the behavior of the bottom end, and the subscript denotes the length of the arc. (We follow the notation of [HW19, proof of Proposition 5].) An analogous statement holds for the ends of a left arc.




FIGURE 3. The four different types of end behavior of a right arc when the right arc is either noninitial or initial such that y_0 is not contained in $\{0\} \times [-\frac{1}{2}, \frac{1}{2}]$. Here, $n = 3$.

For the initial right arc such that y_0 is contained in $\{0\} \times [-\frac{1}{2}, \frac{1}{2}]$, we have five more cases, denoted by η^0 , $\eta_n^{\pm 0}$, and $\eta_n^{0\pm}$ as in Figure 4. (Note that η^0 is a degenerate initial right arc.) As before the first superscript describes the behavior of the top end, the second superscript describes the behavior of the bottom end, and the subscript denotes the length of the arc.




FIGURE 4. The five different types of behavior of the initial right arc such that y_0 is contained in $\{0\} \times [-\frac{1}{2}, \frac{1}{2}]$. Here, $n = 3$.

The following lemma is a consequence of the relationship between a knot-like complex and its associated immersed curve. From now on, we assume that the immersed curves are pulled tight so that for each arc there is at least one peg between two intersection points between the arc and $\{0\} \times \mathbb{R}$. (Note that this is the immersed curve analogue of assuming that a chain complex is reduced.)

Lemma 2.7. *Let C be a knot-like complex and suppose that $H^-(C) = \mathbb{F}[U] \oplus \bigoplus_{i=1}^N \mathbb{F}[U]/(\mathcal{U}^{n_i})$. Then for each i , the immersed curve associated to C has a right arc of length n_i . Conversely, an immersed curve with a right arc of length n comes from a knot-like complex C with an $\mathbb{F}[U]/(\mathcal{U}^n)$ summand in $H^-(C)$.*

Proof. Consider a horizontally simplified basis $\{x_i\}$ for C . Then it follows from the above discussion that each pair $\{x_i, x_{i+1}\}$, i odd, where $\partial x_i = \mathcal{U}^{n_i} x_{i+1} \pmod{\mathcal{V}}$ gives rise to a right arc of length n_i . The converse statement follows similarly. \square

2.2. Concordance invariants and immersed curves. The concordance invariants τ [OS03] and ε [Hom14b] have simple interpretations in the immersed curve setting. Consider the component γ_0 associated to the complement of a knot K . Begin at $(-\frac{1}{2}, 0)$ and follow γ_0 to the right to y_0 . Then $\tau(K)$ is the floor of the sum of $\frac{1}{2}$ and the vertical coordinate of y_0 (i.e., the sum of $\frac{1}{2}$ and the vertical coordinate of the peg right below y_0 .) If γ_0 turns downward at y_0 , then $\varepsilon(K) = 1$. If γ_0 turns upwards at y_0 , then $\varepsilon(K) = -1$. If γ_0 consists of a horizontal line, then $\varepsilon(K) = 0$. For example, in Figure 2(d), we have that $\tau(K) = 4$ and $\varepsilon(K) = 1$. In fact, using [Hom17, Theorem 1] Hanselman-Watson proved that γ_0 is a concordance invariant [HW19, Proposition 2].

2.3. An obstruction to certain immersed curves. The following proposition states that if $\{y_i\}$ is a vertically simplified basis for a knot-like complex and $i > 0$ is even and j is odd, then there cannot be a horizontal arrow of length one from y_i to y_j . The proof relies on the fact that $\partial^2 = 0$. See Figure 5.

Proposition 2.8. *Let $\{y_i\}$ be a vertically simplified basis for a knot-like complex. Then for all even $i > 0$ and odd j , we have $\langle \partial y_i, y_j \rangle = 0 \pmod{\mathcal{U}^2, \mathcal{V}}$.*

Proof. Suppose for the sake of contradiction that $\langle \partial y_i, y_j \rangle \neq 0 \pmod{\mathcal{U}^2, \mathcal{V}}$ for some even $i > 0$ and odd j . Since our basis is assumed to be reduced and since our chain complexes are graded, it follows that $\langle \partial y_i, y_j \rangle = \mathcal{U}$.

We claim that $\langle \partial^2 y_{i-1}, y_j \rangle \neq 0$, contradicting that $\partial^2 = 0$. Indeed,

$$\partial y_{i-1} = \mathcal{V}^{m_{i-1}} y_i + \mathcal{U} \sum_{k \in K} \mathcal{U}^{a_k} \mathcal{V}^{b_k} y_{\ell_k}$$

for some nonnegative integers a_k, b_k and index set K . It follows that

$$\begin{aligned} \langle \partial^2 y_{i-1}, y_j \rangle &= \langle \mathcal{V}^{m_{i-1}} \partial y_i + \mathcal{U} \sum_{k \in K} \mathcal{U}^{a_k} \mathcal{V}^{b_k} \partial y_{\ell_k}, y_j \rangle \\ &= \langle \mathcal{V}^{m_{i-1}} \partial y_i, y_j \rangle + \langle \mathcal{U} \sum_{k \in K} \mathcal{U}^{a_k} \mathcal{V}^{b_k} \partial y_{\ell_k}, y_j \rangle. \end{aligned}$$

The first term above on the right hand side is $\mathcal{V}^{m_{i-1}} \mathcal{U}$, but by Remark 2.5, the second term is $0 \pmod{\mathcal{U}^2}$, completing the proof. \square




FIGURE 5. A graphical depiction of the proof of Proposition 2.8. Since the basis is vertically simplified, there is no other term in $\partial^2 y_{i-1}$ that can cancel with $\mathcal{U} \mathcal{V}^{m_{i-1}} y_j$. It is key that the horizontal arrow between y_i and y_j is length one; if it were longer, there could be a sequence of two arrows (each with a nonzero horizontal component) from y_{i-1} to y_j making $\partial^2 y_{i-1} = 0$.

Corollary 2.9. *With notation as in Figure 3, a noninitial right arc of the form η_1^-+ cannot occur.*

Proof. The result follows immediately from Proposition 2.8 and the discussion in Section 2.1, translating between a chain complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{UV})$ and an immersed curve. Namely, a noninitial right arc η_1^{-+} corresponds to a vertically simplified basis $\{y_i\}$ with a horizontal arrow of length one from y_i to y_j , $i > 0$ even and j odd, which by Proposition 2.8 cannot occur. \square

We have an analogous result when $\varepsilon = 1$ and the initial right arc of the form $\eta_1^{\pm+}$ and η_1^{0+} .

Lemma 2.10. *With notation as in Figure 3 and Figure 4, if $\varepsilon = 1$, then an initial right arc of the form $\eta_1^{\pm+}$ or η_1^{0+} cannot occur.*

Proof. The result follows from Lemma 3.7 of [Hom15], which, in the language of immersed curves, states that if $\varepsilon = 1$ and $\eta_n^{\pm+}$ or η_n^{0+} is the initial right arc, then $n \geq 2$. (We remind the reader that the sign conventions of the a_i -invariants in [Hom15] differ from the conventions in [DHST21].) Since the initial right arc is a part of γ_0 , by Remark 2.6 we may assume we are working with a horizontally and vertically simplified basis.

For the convenience of the reader, we sketch the proof. If $\varepsilon = 1$, then an initial right arc of the form $\eta_n^{\pm+}$ or η_n^{0+} translates to a vertically simplified basis $\{y_i\}$ with a horizontal arrow of length n from y_{i+1} to y_0 and a vertical arrow of length m_i from y_i to y_{i+1} , i odd. If $n = 1$, then since $\{y_i\}$ is vertically simplified, we have $\langle \partial^2 y_i, y_0 \rangle \neq 0 \pmod{(\mathcal{UV})}$, contradicting that $\partial^2 = 0$. Hence, the result follows. \square

3. CABLES

In this section, we use [HW19, Theorem 1] to study the effect of cabling on $HF\bar{K}^-(K)$ by understanding the effect of cabling on right arcs of the immersed curve associated to K , from which Theorems 1.1 and 1.2 follow readily. Note that the following proposition only describes a subset of the right arcs produced by (p, q) -cabling.

Proposition 3.1. *Consider the immersed curve associated to a knot-like complex.*

(1) *Consider a noninitial right arc of the form $\eta_n^{\pm\pm}$. Under the operation of (p, q) -cabling:*

- (a) *A right arc η_n^{-} yields a right arc of length pn .*
- (b) *A right arc η_n^{-+} yields a right arc of length $pn - p + 1$.*
- (c) *A right arc η_n^{+-} yields a right arc of length $pn + p - 1$.*
- (d) *A right arc η_n^{++} yields a right arc of length pn .*

Moreover, in this case the resulting arc is noninitial and has the same form as the original right arc.

(2) *Suppose $\varepsilon = 1$ and consider an initial right arc of the form $\eta_n^{\pm\pm}$ or $\eta_n^{0\pm}$. Under the operation of (p, q) -cabling:*

- (a) *Right arcs η_n^{-} , η_n^{+-} , and η_n^{0-} yield right arcs of length*
 - $\cdot pn$ *for* $q < p(2\tau - 1)$,
 - $\cdot pn + p - 2p\tau + q - 1$ *for* $p(2\tau - 1) < q < 2p\tau$,
 - $\cdot pn + p - 1$ *for* $2p\tau < q$.
- (b) *Right arcs η_n^{-+} , η_n^{++} , and η_n^{0+} yield right arcs of length*
 - $\cdot pn - p + 1$ *for* $q < p(2\tau - 1)$,
 - $\cdot pn - 2p\tau + q$ *for* $p(2\tau - 1) < q < 2p\tau$,
 - $\cdot pn$ *for* $2p\tau < q$.

Furthermore, the right arc obtained in each case from (2a) and (2b) is noninitial and the second superscript does not change.

Before we prove Proposition 3.1, we first show how it implies Propositions 1.3 and 1.4, whose statements we recall.

Proposition 3.2. *If K is a nontrivial knot, then*

$$\text{Ord}(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq \max\{p_1 p_2 \dots p_m (\text{Ord}(K) - 1) + 1, p_1 p_2 \dots p_m\}.$$

Proof. Let K be a nontrivial knot. Since $\text{Ord}(K) = \text{Ord}(-K)$ [OS06, Section 5.1] and $\varepsilon(K) = -\varepsilon(-K)$ [Hom14a, Proposition 3.6], by mirroring if necessary, we may assume that $\varepsilon(K) \geq 0$. As observed in [AE20], since knot Floer homology detects the unknot [OS04a, Theorem 1.2], it follows that $\text{Ord}(K) \geq 1$. Writing $\text{Ord}(K) = n$,

Lemma 2.7 implies that the immersed curve associated to K has a right arc of the form $\eta_n^{\pm\pm}$ or $\eta_n^{0\pm}$ with $n \geq 1$. We begin with the case of $m = 1$.

We first consider the case where we have a noninitial right arc of the form $\eta_n^{\pm\pm}$ with $n \geq 1$. We consider the four cases in part (1) of Proposition 3.1 under the operation of (p_1, q_1) -cabling:

- (a) A right arc of the form η_n^{--} yields a right arc of length $p_1 n$. Since $n \geq 1$, we have that $p_1 n \geq \max\{p_1(n-1) + 1, p_1\}$.
- (b) A right arc of the form η_n^{-+} yields a right arc of length $p_1(n-1) + 1$. Corollary 2.9 states that η_1^{-+} cannot occur, hence $n \geq 2$, which implies that $p_1(n-1) + 1 \geq \max\{p_1(n-1) + 1, p_1\}$.
- (c) A right arc of the form η_n^{+-} yields a right arc of length $p_1(n+1) - 1$. Since $n \geq 1$, we have that $p_1(n+1) - 1 \geq \max\{p_1(n-1) + 1, p_1\}$.
- (d) A right arc of the form η_n^{++} yields a right arc of length $p_1 n$. Since $n \geq 1$, we have that $p_1 n \geq \max\{p_1(n-1) + 1, p_1\}$.

We see that in all cases, we have a right arc of length at least $\max\{p_1(n-1) + 1, p_1\}$.

Now suppose that $\eta_n^{\pm\pm}$ is the initial right arc, which implies that $\varepsilon(K) \neq 0$; hence $\varepsilon(K) = 1$. The desired result now follows from part (2) of Proposition 3.1. That is, we obtain a right arc of length at least $\max\{p_1(n-1) + 1, p_1\}$ in all cases. This is clear for all cases except (2b) when $q < p(2\tau - 1)$. Lemma 2.10 states that η_1^{-+} cannot occur; hence in (2b), $n \geq 2$. Thus, in all cases, we have a right arc of length at least $\max\{p_1(n-1) + 1, p_1\}$. This completes the proof for $m = 1$.

For $m > 1$, we consider two cases separately. If $n > 1$, then the resulting new arc for K_{p_1, q_1} is noninitial by Proposition 3.1 and has length at least $p_1(n-1) + 1 = \max\{p_1(n-1) + 1, p_1\}$. Repeating the analysis from (a), (b), (c), and (d), we see that

$$\text{Ord}(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq p_1 p_2 \dots p_m (n-1) + 1.$$

If $n = 1$, as we have seen from above that the arc that we have started with cannot be a noninitial arc with the form η_1^{-+} , an initial arc of the form $\eta_1^{\pm+}$, or an initial arc of the form η_1^{0+} . Hence the resulting new arc for K_{p_1, q_1} is noninitial with the form η_n^{--} , η_n^{+-} , or η_n^{++} by Proposition 3.1 and has length at least $p_1 = \max\{p_1(n-1) + 1, p_1\}$. Then repeating the analysis from (a), (c), and (d), we see that

$$\text{Ord}(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq p_1 p_2 \dots p_m.$$

This concludes the proof. \square

Proposition 3.3. *Let K be an L-space knot and $K \neq T_{2,3}$. If $J = K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}$, then γ_0 associated to the complement of J contains a right arc with length at least $p_1 \dots p_m (\text{Ord}(K) + 1) - 1$. In particular,*

$$\text{Ord}(J) \geq p_1 \dots p_m (\text{Ord}(K) + 1) - 1.$$

Moreover, if the bridge index of K is $\text{Ord}(K) + 1$, then the above inequality is an equality.

Proof. Suppose K is an L-space knot and $K \neq T_{2,3}$. Recall from [OS05, Theorem 1.2] (see also, for example, [HM17, Section 7]) that for an L-space knot, the knot Floer complex is determined by the Alexander polynomial. Moreover, they showed that the Alexander polynomial of K has the following form:

$$\Delta_K(t) = \sum_{k=0}^{2\ell} (-1)^k t^{\alpha_k}$$

where $\alpha_0, \dots, \alpha_{2\ell}$ is a decreasing sequence of integers and $\alpha_0 - \alpha_{2\ell}$ is twice the genus of K [OS05, Corollary 1.3]. Moreover, since $T_{2,3}$ is the unique genus one L-space knot [Ni07, BZ67, GAn70] and $K \neq T_{2,3}$, we have that $\alpha_0 - \alpha_{2\ell} > 2$. Lastly, Hedden-Watson [HW18, Corollary 9] showed that $\alpha_0 = \alpha_1 + 1$ and $\alpha_{2\ell-1} = \alpha_{2\ell} + 1$. In particular, there is an α_i such that $\alpha_1 > \alpha_i > \alpha_{2\ell-1}$.

The torsion order of an L-space knot may be computed easily from its Alexander polynomial [JMZ20, Lemma 5.1]. More precisely, we have

$$\text{Ord}(K) = \max\{\alpha_{i-1} - \alpha_i \mid 1 \leq i \leq 2\ell\}. \quad (3.1)$$

Moreover, it is well known that the knot Floer complex of an L-space knot has a special form. More precisely, the differentials in the complex form a staircase (see e.g. [HM17, Section 7]). Hence, following the discussion in

Section 2.1, it can be easily verified that in fact there is a noninitial right arc of the form η_n^{+-} in γ_0 associated to K where $n = \text{Ord}(K)$. Part (1c) of Proposition 3.1 implies that the immersed curve γ_0 associated to K_{p_1, q_1} has a right arc of the form η^{+-} of length $p_1 n + p_1 - 1 = p_1(n + 1) - 1$. The desired inequality now follows from repeatedly applying part (1c) of Proposition 3.1.

For the equality, it is a classical theorem of Schubert [Sch54] that the bridge index of J is equal to $p_1 p_2 \cdots p_m$ times the bridge index of K . Moreover, in [JMZ20, Corollary 1.8] they prove that the torsion order of a knot is less than the bridge index of the knot. Hence, we conclude that if the bridge index of K is $\text{Ord}(K) + 1$, then the torsion order of J is $p_1 \cdots p_m (\text{Ord}(K) + 1) - 1$. \square

We now prove Proposition 3.1.

Proof of Proposition 3.1. Recall that [HW19, Theorem 1] states that the immersed curve $\gamma_{p,q}$ corresponding to the (p, q) -cable of K can be obtained from the immersed curve γ for K by the following three step procedure:

- (i) Draw p copies of γ from left to right, each scaled vertically by a factor of p , staggered in height such that each copy is q units below the preceding copy to the left.
- (ii) Connect the loose ends of successive copies of the curve.
- (iii) Translate the pegs horizontally so that they lie in the same vertical line, carrying the curves along with them. Pull the curves tight so that there is at least one peg between two intersection points of each arc and $\{0\} \times \mathbb{R}$. Observe that the left end is $(p-1)q$ units higher than the right end. We shift the left end $\frac{(p-1)q}{2}$ units down and the right end $\frac{(p-1)q}{2}$ units up, so that the two ends points have the same vertical coordinate. We normalize the vertical coordinates so that these two end points are at $(-\frac{1}{2}, 0)$ and $(\frac{1}{2}, 0)$, respectively.

For a picture of Step (i) see Figure 6 and for Step (ii), see Figure 8. We will analyze the contribution of the rightmost copy, namely the p th copy, of γ from Step (i) to the curve $\gamma_{p,q}$. We begin by considering the four different types $\eta_n^{\pm\pm}$ of right arc, and assume that $\eta_n^{\pm\pm}$ is a noninitial right arc. While there may be a nontrivial local system present, this will not be relevant to the proof since this part of the argument only makes use of right arcs and we may confine the nontrivial local system to left arcs.

We first consider (1a), that is, a right arc η_n^{-} , scaled vertically by a factor of p . Since η_n^{-} is noninitial, we see that after translating the pegs horizontally and pulling tight, the image of η_n^{-} spans pn pegs, as each of the n pegs originally spanned by η_n^{-} have now become p pegs (consisting of the original peg and $p-1$ pegs below it), for a total of pn pegs.

We next consider (1b), that is, a right arc η_n^{-+} , scaled vertically by a factor of p . When we translate the pegs horizontally, we see that the image of η_n^{-+} spans $p(n-1) + 1$ pegs, as each of the first $n-1$ pegs originally spanned by η_n^{-+} have now become p pegs (consisting of the original peg and $p-1$ pegs below it), plus we still have the last of the original n pegs, for a total of $p(n-1) + 1 = pn - p + 1$ pegs.

We next consider (1c), that is, a right arc η_n^{+-} , scaled vertically by a factor of p . When we translate the pegs horizontally, we see that the image of η_n^{+-} spans $pn + p - 1$ pegs, as each of the original n pegs has now become p pegs (consisting of the original peg and the $p-1$ pegs below it), plus an additional $p-1$ pegs above the first of the original pegs.

Lastly, we consider (1d), a right arc of the form η_n^{++} ; the result follows as in the case of η_n^{-} .

Furthermore, it is clear that if $\eta_n^{\pm\pm}$ is a noninitial right arc, then the right arc obtained in each of the above four cases is a noninitial right arc and the form does not change. This completes the proof for noninitial right arcs, i.e. Part (1) of the proposition.

We next consider the initial right arcs $\eta_n^{\pm\pm}$ and $\eta_n^{0\pm}$ with $\varepsilon = 1$. We only give a detailed proof for the case η_n^{-} with $\varepsilon = 1$. The rest of the cases can be proved similarly. Note that given an initial right arc of the form η_n^{-} and $\varepsilon = 1$, we have $\tau > 0$. Further, we will not need to consider local systems since we will now restrict to working with γ_0 .

Let u_0 be the peg at $(0, \tau + \frac{1}{2})$, u'_0 be the peg at $(0, \tau - \frac{1}{2})$, v_0 be the peg at $(0, \tau - \frac{1}{2} - n)$, and w_0 be the peg at $(0, -\tau + \frac{1}{2})$ for γ . Note that u_0 and u'_0 are the pegs that are located right above and below the intersection point between the essential left arc and the vertical line $\{0\} \times \mathbb{R}$, respectively, and w_0 is the peg right above the intersection point between the essential right arc and the vertical line $\{0\} \times \mathbb{R}$ (see Section 2.2 and Figure 7).




FIGURE 6. The rightmost copy of a right arc of γ from Step (i), with the four different types of end behavior. Here, $n = 3$ and $p = 2$.





FIGURE 7. Pegs u_0 , u'_0 , v_0 , and w_0 of an initial right arc of the form η_n^{--} with $\varepsilon = 1$ and $\tau > 0$.

Here, we are using the fact that there is a symmetry of the immersed curves under 180 degree rotation [HRW18, Theorem 7]. Lastly, note that there are the two intersection points between the initial right arc and the vertical line $\{0\} \times \mathbb{R}$, and v_0 is the peg right below the bottom intersection point (see Figure 7).

Let u and u' be the pegs for the immersed curve $\gamma_{p,q}$ in the p th copy of γ from Step (i) of Hanselman-Watson's procedure that correspond to u_0 and u'_0 , respectively. The peg that corresponds to v_0 in the p th copy of γ is denoted by v and the peg that corresponds to w_0 in the $(p-1)$ th copy of γ is denoted by w (see Figure 8). By abusing notation, the vertical coordinates of the pegs u , u' , v , and w will also be denoted by u , u' , v , and w , respectively. With this notation, note that we have $u' < u$.

FIGURE 8. Pegs u , u' , v , and w for the immersed curve $\gamma_{p,q}$ for three cases.

As depicted in Figure 8, we consider the following three cases:

$$w < u', \quad u' < w < u, \quad \text{and} \quad u < w.$$

We leave it as an easy exercise for the readers to verify that these are the exact cases that appear on (2a), i.e. $w < u'$ if and only if $q < p(2\tau - 1)$, $u' < w < u$ if and only if $p(2\tau - 1) < q < 2p\tau$, and $u < w$ if and only if $2p\tau < q$.

Suppose that $w < u'$ which corresponds to Figure 8 (1). By translating the pegs horizontally, we see that the image of the p th copy of η_n^- from Step (i) takes the form η^- and, as in the noninitial right arc case from (1a), it can be checked that it spans pn pegs. To be more precise, the image spans all the pegs between u' and v including u' and excluding v . Following steps (i), (ii), and (iii), it can be easily computed that the vertical coordinate of u' minus the vertical coordinate of v is pn . Hence, we conclude that the resulting arc is noninitial and takes the form η_{pn}^- .

Suppose that $u' < w < u$ which corresponds to Figure 8 (2). In this case, the image of the p th copy of γ takes the form η^{+-} . Moreover, the image spans all the pegs between w and v excluding both w and v . As before, it can be easily computed that the vertical coordinate of w minus the vertical coordinate of v is $pn + p - 2p\tau + q$. We conclude that the resulting arc is noninitial and takes the form $\eta_{pn+p-2p\tau+q-1}^{+-}$.

Lastly, suppose that $u < w$ which corresponds to Figure 8 (3). Again, the image of the p th copy of γ takes the form η^{+-} . In this case, the image spans all the pegs between u and v excluding both u and v . Since the vertical coordinate of u minus the vertical coordinate of w is $pn + p$, the resulting arc is noninitial and takes the form η_{pn+p-1}^{+-} . This completes the proof of the proposition for initial right arcs of the form η_n^- . As mentioned above, the remaining cases can be addressed similarly. \square

4. APPLICATIONS

In this section, we prove the main theorems and applications stated in the introduction. Most of these follow directly from Proposition 3.2 and 3.3. We first prove Theorems 1.1 and 1.2. We prove them together, since the proofs are similar. We recall the statements.

Theorem 4.1. *If K is a nontrivial knot, then $u(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq p_1 p_2 \dots p_m$. Moreover, if $1 < p_0 < q_0$ and $q_0 \neq 3$, then $u(T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}) \geq p_0 p_1 \dots p_m - 1$.*

Proof. From Proposition 3.2, we have $\text{Ord}(K_{p_1, q_1; p_2, q_2; \dots; p_m, q_m}) \geq p_1 p_2 \dots p_m$. Moreover, since a positive torus knot T_{p_0, q_0} is an L -space knot with $\text{Ord}(T_{p_0, q_0}) = p_0 - 1$ (see e.g. [JMZ20, Corollary 5.3]), by Proposition 3.3 we

have $\text{Ord}(T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}) \geq p_0 p_1 \dots p_m - 1$. The proof is then completed by using the unknotting bound of Alishahi-Eftekhary in (2.1). \square

Next we prove Corollary 1.5, whose statement we recall.

Corollary 4.2. *If K is a nontrivial slice knot, then any slice disk bounded by $K_{p_1, 1; p_2, 1; \dots; p_m, 1}$ has at least $p_1 p_2 \dots p_m + 1$ local minima with respect to the radial function on B^4 restricted to the disk.*

Proof. Let K be a nontrivial slice knot. By Proposition 3.2, we have $\text{Ord}(K_{p_1, 1; p_2, 1; \dots; p_m, 1}) \geq p_1 p_2 \dots p_m$. Juhász-Miller-Zemke [JMZ20, Corollary 1.7] bound the number of local minima of a slice disk from below by the torsion order, which completes the proof. \square

Recall that for a knot K , its bridge index is denoted by $\text{br}(K)$ and its braid index is denoted by $\text{b}(K)$. We prove Corollary 1.6, whose statement we recall.

Corollary 4.3. *Let K be either an algebraic knot or an iterated torus knot $T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}$ where $T_{p_0, q_0} \neq T_{2, 3}$. If J is concordant to K , then $\text{b}(J) \geq \text{b}(K)$ and $\text{br}(J) \geq \text{br}(K)$.*

Proof. First, assume that K is an iterated torus knot $T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}$ where $T_{p_0, q_0} \neq T_{2, 3}$. As before, since $\text{Ord}(T_{p_0, q_0}) = p_0 - 1$, by Proposition 3.3 we conclude that γ_0 associated to K contains a right arc with length at least $p_0 p_1 \dots p_m - 1$. By Schubert [Sch54], we see that $\text{br}(K) = \text{b}(K) = p_0 p_1 \dots p_m$ (see also [Wil92]). Since γ_0 is a concordance invariant, if J is concordant to K , then $\text{Ord}(J) \geq p_0 p_1 \dots p_m - 1$. Now, the proof is complete by using the bound $\text{br}(J) \geq \text{Ord}(J) - 1$ [JMZ20, Corollary 1.8] and the fact that $\text{br}(J) \leq \text{b}(J)$.

Now, assume that K is an algebraic knot. For this we do not need to use Proposition 3.3. Recall that if K is an algebraic knot then it has the following form $T_{p_0, q_0; p_1, q_1; \dots; p_m, q_m}$ where $q_{i+1} > p_i q_i p_{i+1}$ (see e.g. [EN85]). In particular, it is an L -space knot [Hed09]. Using this fact, it is an easy exercise to verify that $\alpha_1 - \alpha_2 = p_0 p_1 \dots p_m - 1$ where

$$\Delta_K(t) = \sum_{k=0}^{2\ell} (-1)^k t^{\alpha_k}$$

and $\alpha_0, \dots, \alpha_{2\ell}$ is a decreasing sequence of integers. Then by (3.1) and the bound of [JMZ20, Corollary 1.8], we conclude that γ_0 associated to K contains a right arc with length $p_0 p_1 \dots p_m - 1$. As before, since γ_0 is a concordance invariant, the proof is complete. \square

Remark 4.4. The above proof shows that if K is an algebraic knot or an iterated torus knot, then $\text{br}(K) = \text{b}(K) = \text{Ord}(K) + 1$.

Lastly, we restate and prove Corollary 1.7.

Corollary 4.5. *There exists a family of iterated torus knots $\{K_i\}$, such that the gap between $u(K_i)$ and $g_4(K_i)$ is arbitrarily large.*

Proof. Let K be $T_{2, -5; 2, 5}$. Then there is a genus 6 Seifert surface Σ for K obtained by taking two parallel copies of the genus 2 Seifert surface for $T_{2, -5}$ and connecting them with 5 half-twisted bands. Note that $T_{2, 5} \# T_{2, -5}$ sits on Σ and it bounds a genus 4 surface embedded in Σ . Since $T_{2, 5} \# T_{2, -5}$ is a slice knot, if we surger along $T_{2, 5} \# T_{2, -5}$ on Σ we obtain a genus 2 surface bounded by K properly embedded in B^4 . Hence we conclude $g_4(K) \leq 2$.

In fact, we show that $g_4(K) = 2$. Note that $\nu^+(T_{2, -5}) = 0$ (see e.g. [BCG17]), where ν^+ is the concordance invariant introduced by Hom-Wu [HW16]. Also, they show that ν^+ gives a lower bound on the 4-genus. Then by a cabling formula for ν^+ of Wu [Wu16, Theorem 1.1], we conclude that $\nu^+(K) = 2$ and $g_4(K) = 2$.

Let i be a positive integer and K_i be $K_{i, 1}$. Then we have that $g_4(K_i) = 2i$, where the upper bound is obtained by taking n parallel copies of the genus 2 surface bounded by K , and the lower bound is given by the inequality by Sato [Sat18, Corollary 1.4] for ν^+ . Lastly, by Theorem 1.2 and the fact that $u(K) = u(-K)$, we have that $u(K_i) \geq 4i - 1$. This completes the proof. \square

REFERENCES

- [AE20] Akram Alishahi and Eaman Eftekhary. Knot Floer homology and the unknotting number. *Geom. Topol.*, 24(5):2435–2469, 2020.
- [BCG17] József Bodnár, Daniele Celoria, and Marco Golla. A note on cobordisms of algebraic knots. *Algebr. Geom. Topol.*, 17(4):2543–2564, 2017.
- [BZ67] Gerhard Burde and Heiner Zieschang. Neuwirthsche Knoten und Flächenabbildungen. *Abh. Math. Sem. Univ. Hamburg*, 31:239–246, 1967.
- [Che19] Wenzhao Chen. Knot Floer homology of satellite knots with (1,1)-patterns. arXiv:1912.07914, 2019.
- [DHST21] Irving Dai, Jennifer Hom, Matthew Stoffregen, and Linh Truong. More concordance homomorphisms from knot Floer homology. *Geom. Topol.*, 25(1):275–338, 2021.
- [EN85] David Eisenbud and Walter Neumann. *Three-dimensional link theory and invariants of plane curve singularities*, volume 110 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1985.
- [FK17] Peter Feller and David Kratovich. On cobordisms between knots, braid index, and the epsilon-invariant. *Math. Ann.*, 369(1-2):301–329, 2017.
- [GAn70] Francisco González-Acuña. Dehn’s construction on knots. *Bol. Soc. Mat. Mexicana (2)*, 15:58–79, 1970.
- [Hed09] Matthew Hedden. On knot Floer homology and cabling. II. *Int. Math. Res. Not. IMRN*, (12):2248–2274, 2009.
- [HKP20] Jennifer Hom, Sungkyung Kang, and JungHwan Park. Ribbon knots, cabling, and handle decompositions. arXiv:2003.02832, to appear: *Math. Res. Lett.*, 2020.
- [HM17] Kristen Hendricks and Ciprian Manolescu. Involutive Heegaard Floer homology. *Duke Math. J.*, 166(7):1211–1299, 2017.
- [Hom14a] Jennifer Hom. Bordered Heegaard Floer homology and the tau-invariant of cable knots. *J. Topol.*, 7(2):287–326, 2014.
- [Hom14b] Jennifer Hom. The knot Floer complex and the smooth concordance group. *Comment. Math. Helv.*, 89(3):537–570, 2014.
- [Hom15] Jennifer Hom. An infinite-rank summand of topologically slice knots. *Geom. Topol.*, 19(2):1063–1110, 2015.
- [Hom17] Jennifer Hom. A survey on Heegaard Floer homology and concordance. *J. Knot Theory Ramifications*, 26(2):1740015, 24, 2017.
- [Hom20] Jennifer Hom. Lectures notes on Heegaard Floer homology, 2020. arXiv:2008.01836.
- [HRW16] Jonathan Hanselman, Jacob Rasmussen, and Liam Watson. Bordered Floer homology for manifolds with torus boundary via immersed curves. arXiv:1604.03466, 2016.
- [HRW18] Jonathan Hanselman, Jacob Rasmussen, and Liam Watson. Heegaard Floer homology for manifolds with torus boundary: properties and examples. arXiv:1810.10355, 2018.
- [HW16] Jennifer Hom and Zhongtao Wu. Four-ball genus bounds and a refinement of the Ozsváth-Szabó tau invariant. *J. Symplectic Geom.*, 14(1):305–323, 2016.
- [HW18] Matthew Hedden and Liam Watson. On the geography and botany of knot Floer homology. *Selecta Math. (N.S.)*, 24(2):997–1037, 2018.
- [HW19] Jonathan Hanselman and Liam Watson. Cabling in terms of immersed curves. arXiv:1908.04397, to appear: *Geom. Topol.*, 2019.
- [Inc16] Kenan Ince. The untwisting number of a knot. *Pacific J. Math.*, 283(1):139–156, 2016.
- [Inc17] Kenan Ince. Untwisting information from Heegaard Floer homology. *Algebr. Geom. Topol.*, 17(4):2283–2306, 2017.
- [JMZ20] András Juhász, Maggie Miller, and Ian Zemke. Knot cobordisms, bridge index, and torsion in Floer homology. *J. Topol.*, 13(4):1701–1724, 2020.
- [KM93] Peter B. Kronheimer and Tomasz S. Mrowka. Gauge theory for embedded surfaces. I. *Topology*, 32(4):773–826, 1993.
- [KWZ20] Artem Kotelskiy, Liam Watson, and Claudius Zibrowius. A mnemonic for the Lipshitz-Ozsváth-Thurston correspondence. arXiv:2005.02792, to appear: *Algebr. Geom. Topol.*, 2020.
- [Lev69] Jerome Levine. Knot cobordism groups in codimension two. *Comment. Math. Helv.*, 44:229–244, 1969.
- [Lit84] Richard A. Litherland. Cobordism of satellite knots. In *Four-manifold theory (Durham, N.H., 1982)*, volume 35 of *Contemp. Math.*, pages 327–362. Amer. Math. Soc., Providence, RI, 1984.
- [Liv21] Charles Livingston. Null-homologous unknottings. In *Topology and geometry—a collection of essays dedicated to Vladimir G. Turaev*, volume 33 of *IRMA Lect. Math. Theor. Phys.*, pages 59–68. Eur. Math. Soc., Zürich, [2021] ©2021.
- [LOT18] Robert Lipshitz, Peter S. Ozsváth, and Dylan P. Thurston. Bordered Heegaard Floer homology. *Mem. Amer. Math. Soc.*, 254(1216):viii+279, 2018.
- [McC21a] Duncan McCoy. Gaps between consecutive untwisting numbers. *Glasg. Math. J.*, 63(1):59–65, 2021.
- [McC21b] Duncan McCoy. Null-homologous twisting and the algebraic genus. In *2019–20 MATRIX annals*, volume 4 of *MATRIX Book Ser.*, pages 147–165. Springer, Cham, [2021] ©2021.
- [MD88] Yves Mathieu and Michel Domergue. Chirurgies de Dehn de pente ± 1 sur certains noeuds dans les 3-variétés. *Math. Ann.*, 280(3):501–508, 1988.
- [Ni07] Yi Ni. Knot Floer homology detects fibred knots. *Invent. Math.*, 170(3):577–608, 2007.
- [OS03] Peter S. Ozsváth and Zoltán Szabó. Knot Floer homology and the four-ball genus. *Geom. Topol.*, 7:615–639, 2003.
- [OS04a] Peter S. Ozsváth and Zoltán Szabó. Holomorphic disks and genus bounds. *Geom. Topol.*, 8:311–334, 2004.
- [OS04b] Peter S. Ozsváth and Zoltán Szabó. Holomorphic disks and knot invariants. *Adv. Math.*, 186(1):58–116, 2004.
- [OS05] Peter S. Ozsváth and Zoltán Szabó. On knot Floer homology and lens space surgeries. *Topology*, 44(6):1281–1300, 2005.

- [OS06] Peter S. Ozsváth and Zoltán Szabó. Holomorphic triangles and invariants for smooth four-manifolds. *Adv. Math.*, 202(2):326–400, 2006.
- [Ras03] Jacob Rasmussen. *Floer homology and knot complements*. ProQuest LLC, Ann Arbor, MI, 2003. Thesis (Ph.D.)–Harvard University.
- [Sat18] Kouki Sato. A full-twist inequality for the ν^+ -invariant. *Topology Appl.*, 245:113–130, 2018.
- [Sch54] Horst Schubert. Über eine numerische Knoteninvariante. *Math. Z.*, 61:245–288, 1954.
- [Sch85] Martin G. Scharlemann. Unknotting number one knots are prime. *Invent. Math.*, 82(1):37–55, 1985.
- [ST88] Martin G. Scharlemann and Abigail Thompson. Unknotting number, genus, and companion tori. *Math. Ann.*, 280(2):191–205, 1988.
- [Sto03] Alexander Stoimenow. Positive knots, closed braids and the Jones polynomial. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 2(2):237–285, 2003.
- [Tri69] A. G. Tristram. Some cobordism invariants for links. *Proc. Cambridge Philos. Soc.*, 66:251–264, 1969.
- [Wil92] Robert F. Williams. The braid index of generalized cables. *Pacific J. Math.*, 155(2):369–375, 1992.
- [Wu16] Zhongtao Wu. A cabling formula for the ν^+ invariant. *Proc. Amer. Math. Soc.*, 144(9):4089–4098, 2016.
- [Yan08] Zhiqing Yang. Unknotting number of the connected sum of n identical knots. *J. Knot Theory Ramifications*, 17(3):253–255, 2008.
- [Zem19] Ian Zemke. Link cobordisms and functoriality in link Floer homology. *J. Topol.*, 12(1):94–220, 2019.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332
Email address: `hom@math.gatech.edu`

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC, 27607
Email address: `tlid@math.ncsu.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, DAEJEON 34141, REPUBLIC OF KOREA
Email address: `jungpark0817@kaist.ac.kr`