

Feynman Integrals of Grassmannians

Tai-Fu Feng^{*a,b,c,d}, Hai-Bin Zhang^{†a,b,c}, Chao-Hsi Chang^{‡e,f,g}

^a*Department of Physics, Hebei University, Baoding, 071002, China*

^b*Hebei Key Laboratory of High-precision Computation and
Application of Quantum Field Theory, Baoding, 071002, China*

^c*Department of Physics, Guangxi University, Nanning, 530004, China*

^d*Department of Physics, Chongqing University, Chongqing, 401331, China*

^e*Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Science, Beijing, 100190, China*

^f*CCAST (World Laboratory), P.O.Box 8730, Beijing, 100190, China and*

^g*School of Physical Sciences, University of Chinese
Academy of Sciences, Beijing 100049, China*

Abstract

We embed Feynman integrals in the subvarieties of Grassmannians through homogenization of the integrands in projective space, then obtain GKZ-systems satisfied by those scalar integrals. The Feynman integral can be written as linear combinations of the hypergeometric functions of a fundamental solution system in neighborhoods of regular singularities of the GKZ-system, whose linear combination coefficients are determined by the integral on an ordinary point or some regular singularities. Taking some Feynman diagrams as examples, we elucidate in detail how to obtain the fundamental solution systems of Feynman integrals in neighborhoods of regular singularities. Furthermore we also present the parametric representations of Feynman integrals of the 2-loop self-energy diagrams which are convenient to embed in the subvarieties of Grassmannians.

PACS numbers: 02.30.Jr, 11.10.-z, 12.38.Bx

Keywords: Feynman integral, Grassmannian, GKZ-hypergeometric system

* email:fengt@hbu.edu.cn

† email:zhanghb@hbu.edu.cn

‡ email:zhangzx@itp.ac.cn

I. INTRODUCTION

Although the discovery of Higgs with mass around 125 GeV [1, 2] indicates a great success of the Standard Model (SM), the updated experimental data on the anomalous magnetic moment (MDM) of muon [3] and mass of the charged gauge boson [4] seem to herald the signals of new physics beyond. Before the collision energy of the center of mass of the running collider reaches the threshold energy of new physics, the SM theoretical predictions of various physical observations should be accurately calculated [5]. Through comparison between the experimental data and corresponding SM evaluations, the energy scale of new physics and strength of the coupling of new physics particles to fields of the SM can be roughly estimated.

To obtain the SM evaluations of physical observables accurately, one has to compute the scalar integrals of corresponding Feynman diagrams involved at first, which is a challenging work in radiative corrections in quantum field theory. In dimensional regularization, any scalar integral can be expanded as a Laurent series around $\varepsilon = 0$ where the time-space dimension $D = 4 - 2\varepsilon$. For example, the Feynman integral of the 1-loop triangle diagram C_0 is expanded as

$$C_0 = C_0^{(0)} + \sum_{i=1}^{\infty} C_0^{(i)} \varepsilon^i, \quad (1)$$

around $D = 4$, and the well-known result of $C_0^{(0)}$ is presented in the literature[6]. However, evaluating $C_0^{(i)}$ precisely is also necessary to obtain high order radiative corrections to the amplitude self-consistently when the virtual interactions originate from the counter terms[7]. Considering Feynman integrals as the generalized hypergeometric functions [8], one finds that the D -module of a Feynman diagram [9] is isomorphic to Gel'fand-Kapranov-Zelevinsky (GKZ) D -module [10–14]. Correspondingly Feynman integrals satisfy indeed the systems of holonomic linear partial differential equations (PDEs) [15] whose singularities are determined by the Landau singularities.

During the past decades, some progresses have been made on the subject. Under the assumption of zero virtual masses, the Feynman integral of the 1-loop triangle diagram is given as a linear combination of the fourth kind of Appell functions [16] whose arguments are the dimensionless ratios among the external momentum squared, and is simplified further as a linear combination of the Gauss functions ${}_2F_1$ through the quadratic transforma-

tion [17] in the literature [18]. Taking some special assumptions on parameter space, the authors of Ref. [19] obtain the multiple hypergeometric expressions of the scalar integral C_0 through Mellin-Barnes representations. An algorithm to evaluate the scalar integrals of 1-loop vertex-type Feynman diagrams is presented in Ref. [20], and a geometrical interpretation of the analytic expressions of the scalar integrals of 1-loop N -point Feynman diagrams is also given in Ref. [21]. Certainly the expressions of C_0 function can also be derived from expressions of the 1-loop massive N -point Feynman diagrams [22, 23]. In addition Feynman integrals of ladder diagrams with 3 or 4 external lines can be evaluated through the parametric representation and Mellin-Barnes contour integrals [24]. Actually the 1-loop 2-point function B_0 can be expressed as a linear combination of the Gauss functions ${}_2F_1$ through the recurrence relations respecting time-space dimension, the 1-loop 3-point function C_0 similarly is formulated as a linear combination of the Appell functions F_1 , and the 1-loop 4-point function D_0 is formulated as a linear combination of the Lauricella functions F_s [25, 26], respectively. The expression of C_0 is convenient for analytic continuation and numerical evaluation because continuation of F_1 has been analyzed thoroughly. However, how to perform continuation of the Lauricella function F_s outside its convergent domain is still a challenge. Expressing Feynman integrals as generalized hypergeometric functions in dimension regularization, the authors of Ref. [27] analyze the Laurent expansion of these hypergeometric functions around $D = 4$. The differential-reduction algorithm to evaluate those hypergeometric functions can be found in Refs. [28–32]. A hypergeometric system of linear PDEs is given through the corresponding Mellin-Barnes representation [33]. Some irreducible master integrals of sunset and bubble Feynman diagrams are explicitly evaluated through Mellin-Barnes contour in Ref. [34].

GKZ-hypergeometric systems of Feynman integrals with codimension= 0, 1 are presented in Refs. [35, 36] through Lee-Pomeransky parametric representations [37]. To construct canonical series solutions with suitable independent variables, one should compute the restricted D -module of GKZ-hypergeometric system originating from Lee-Pomeransky representations on corresponding hyperplane in the parameter space [38–40]. Certainly GKZ-hypergeometric systems of some Feynman diagrams are obtained from Mellin-Barnes representations [41–43] through Miller’s transformation [44, 45]. Ref.[46] explores the idea of bootstrap Feynman integrals using integrability, Ref.[47] applies the GKZ description

of periods to solve the l-loop banana amplitudes with their general mass dependence, and Ref. [48] completely clarifies the analytic structure of all banana amplitudes with arbitrary masses, respectively. The updated summary on some classical and modern aspects of hypergeometric differential equations is given in literature [49]. A mathematical package for integrating families of Feynman integrals order-by order in the dimensional regulator from their GKZ-systems is given in literature [50], and a specialized integration algorithm for parametric Feynman integrals with tame kinematics is also presented in literature [51]. Actually Cohen-Macaulay Property of Feynman Integral indicates that the process of finding a series representation of these integrals is fully algorithmic [52], and regular singularities of Feynman Integral are determined by its Landau Discriminant [53]. Ref. [54] introduces a class of polytopes to analyze the structure of UV and IR divergences of general Feynman integrals in Schwinger parameter space. Furthermore, Ref. [55] elaborates on the connection among GKZ-systems, de Rham theory for twisted cohomology groups, and Pfaffian equations for Feynman Integrals. Ref. [56] presents a Mathematica package to find linear transformations for some classes of multivariable hypergeometric functions. In literature [57] the class of N loop massive scalar self-energy diagrams with $N + 1$ propagators is studied, and the new convergent series representations for the 2-loop sunset diagram with 3 different propagator masses and external momentum are obtained in Ref.[58]. The relationship between Feynman diagrams and hypergeometric functions is reviewed in the Ref. [59].

Since the fundamental solution systems of neighborhoods of different regular singularities are different, and many equivalent fundamental solution systems can be constructed corresponding to neighborhoods of same regular singularity. When a Feynman integral is expressed as a linear combination of hypergeometric functions of the fundamental solution system, there are obvious differences among the equivalent representations accordingly. Finding a framework for understanding these rambling results is undoubtedly important on both theoretical and phenomenological aspects. The fundamental solution systems of neighborhoods of different regular singularities can be regarded as analytic continuations of each other. Thus the analytic expression of Feynman integral can be obtained in the whole parameter space.

Nevertheless, one only obtains the fundamental solution systems in neighborhoods of the regular singularities $0, \infty$ using the GKZ-hypergeometric system of a general compact

manifold. Taking the Feynman integral as a linear combinations of hypergeometric functions from the fundamental solution systems around different regular singularities and embedding it in subvarieties of Grassmannians, one finds that the dual space of the GKZ-system satisfied by the Feynman integral in the splitting coordinates determines the integer lattice matrices, then writes the exponent matrices compatible with the integer lattice matrices. A pair of integer lattice and exponent matrices uniquely determines a generalized hypergeometric function where the geometric representation [60] of its exponent matrix can clearly tell us whether or not the hypergeometric function of a different pair of integer lattice and exponent matrices is proportional to it in the nonempty intersection of their convergent regions.

Some hypergeometric functions are defined on the Grassmannian $G_{k,n}$ [61] in a natural way, where the Grassmannian $G_{k,n}$ is the manifold of all $(k-1)$ -dimensional projective subspaces of $(n-1)$ -dimensional projective space ($k < n$) [62]. In order to describe Grassmannian exactly, one can understand the manifold $Z_{k,n}$ of $k \times n$ matrices of rank k being a bundle space over the base space $G_{k,n}$ whose projection $Z_{k,n} \rightarrow G_{k,n}$ assigns to each matrix $Z_{k,n}$ the k -dimensional projective subspace spanned by the row vectors of this matrix. We consider an arbitrary smooth function on the manifold $Z_{k,n}$ satisfying the homogeneous condition

$$f(\chi_1 z_1, \dots, \chi_n z_n) = \prod_{i=1}^n \chi_i^{\alpha_i - 1} f(z_1, \dots, z_n), \chi_i \neq 0, \quad (2)$$

where z_i , ($i = 1, \dots, n$) denotes the i -th column vector of the $k \times n$ matrix $Z_{k,n}$, and

$$\sum_{i=1}^n \alpha_i = n - k. \quad (3)$$

Let $t = (t_1, \dots, t_k)$ denote the local coordinates of the k -dimensional projective subspace, and the volume element

$$\omega(t) = \sum_{i=1}^k (-)^{i-1} dt_1 \wedge \dots \wedge dt_{i-1} \wedge dt_{i+1} \wedge \dots \wedge dt_k \quad (4)$$

thus the integral

$$F(z) = \int_S f(t \cdot z) \omega(t) \quad (5)$$

satisfies the homogeneous conditions

$$\begin{aligned} F(g \cdot z) &= |\det g|^{-1} F(z), \quad g \in GL(k, C), \\ F(z \cdot \chi) &= \prod_{i=1}^n \chi_i^{\alpha_i - 1} F(z), \quad \chi = \text{diag}(\chi_1, \dots, \chi_n). \end{aligned} \quad (6)$$

Here S is an arbitrary $(k-1)$ -dimensional hypersurface enclosing the origin. In this approach elements of the matrix $Z_{k \times n}$ are named as the splitting local coordinates. Obviously the first condition is equivalent to the system of partial differential equations (PDEs) [63–66]

$$\sum_{j=1}^n z_{ij} \frac{\partial F}{\partial z_{i'j}} = -\delta_{i,i'} F, \quad i \in [1, k], \quad i' \in [1, k], \quad (7)$$

with $\delta_{i,i'}$ denoting the Kronecker symbol, and the second condition of Eq.(6) is equivalent to the system of PDEs

$$\sum_{i=1}^k z_{ij} \frac{\partial F}{\partial z_{ij}} = (\alpha_j - 1)F, \quad j \in [1, n], \quad (8)$$

respectively. Using the GKZ-system presented above, we can construct the fundamental solution system which is composed of the hypergeometric functions of local splitting coordinates z_{ij} [65, 66].

After embedding parametric representations of Feynman integrals in projective space, we formulate the Feynman integrals as hypergeometric functions over Grassmannians. Taking Feynman integrals of the 1-loop self-energy, the massless 1-loop triangle, and the 2-loop vacuum diagrams as examples, we illuminate how to construct canonical series solutions from those relevant GKZ-systems [65, 66], whose convergent regions of those series can be derived by Horn's study [67]. Finally parametric representations of the 2-loop self energy diagrams with 3, 4 and 5 propagators are presented respectively. Those Feynman integrals are embedded in the subvarieties of Grassmannians by the parametric representations directly. The adopted algorithm here is applied to obtain the fundamental solution systems of the GKZ-systems associating with the subvarieties of the Grassmannian $G_{3,6}$ in Ref. [65], and all techniques in detail of this approach can be found in the long review article [66]. In principle, we can embed the parametric representation of a given Feynman integral into infinite subvarieties of Grassmannians. However, the dimension of dual space of the GKZ-systems of most subvarieties is much larger than the number of independent dimensionless ratios between the square of external momentum and the square of virtual masses. Correspondingly convergent regions of the obtained hypergeometric series solutions are empty. In addition, we don't state those mathematical concepts and theorems that have been used in our analyses here, because they can be found in some well-known mathematical textbooks [68–76].

The generally strategy for analysing Feynman integral includes following steps in this approach. First we embed parametric representation of Feynman integral in the subvarieties

of proper Grassmannians, and write out the GKZ-system satisfied by the Feynman integral. Second the fundamental solution systems are constructed in neighborhoods of regular singularities of the GKZ-system. The integration constants, i.e. the combination coefficients, are determined from the Feynman integral with some special kinematic parameters [42, 77, 78].

Our presentation is organized as following. After embedding the Feynman integrals in the subvarieties of Grassmannians appropriately, we derive the fundamental solution systems of Feynman integrals of the 1-loop vacuum and 1-loop self energy in section II, respectively. Then we present in detail how to embed parametric representation of the 1-loop massless triangle diagrams in section III, and several equivalent fundamental solution systems are presented in different neighborhoods of regular singularities. In section IV, we embed Feynman integral of the 2-loop vacuum in the subvarieties of Grassmannian $G_{4,8}$, then present the hypergeometric series in neighborhoods of regular singularities. Finally parametric representations of the 2-loop self energy diagrams with 3, 4 and 5 propagators are presented respectively in section V, and the conclusions are summarized in section VI.

II. THE 1-LOOP SELF-ENERGY

We first use the well-known 1-loop bubble diagram to illustrate how to embed the corresponding Feynman integral into the subvarieties of Grassmannians, then construct its fundamental solution system. The Feynman integral of the 1-loop bubble diagram is written as

$$\begin{aligned} A_{1VA}(m^2) &= \left(\Lambda_{\text{RE}}^2\right)^{2-D/2} \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2 - m^2} \\ &= -\frac{i\Gamma(1 - D/2) \left(\Lambda_{\text{RE}}^2\right)^{2-D/2}}{(4\pi)^{D/2} \left(m^2\right)^{1-D/2}}, \end{aligned} \quad (9)$$

with Λ_{RE} denoting the renormalization scale. After performing the Wick rotation and integrating over angles,

$$A_{1VA}(m^2) = -\frac{i\left(\Lambda_{\text{RE}}^2\right)^{2-D/2}}{(4\pi)^{D/2}\Gamma(\frac{D}{2})} \int_0^\infty \frac{dt_1 t_1^{D/2-1}}{t_1 + m^2}. \quad (10)$$

Introducing the homogeneous coordinate $t_2 = 1$, one reformulates the scalar integral as

$$A_{1VA}(m^2) = -\frac{i\left(\Lambda_{\text{RE}}^2\right)^{2-D/2}}{(4\pi)^{D/2}\Gamma(\frac{D}{2})} \int \frac{\omega(t) t_1^{D/2-1} t_2^{-D/2}}{t_1 + t_2 m^2}, \quad (11)$$

with the volume element of projective line $\omega(t) = t_2 dt_1 - t_1 dt_2$. Eq.(11) can be taken as an integral on the Grassmannian $G_{2,3}$ with splitting local coordinates as

$$A^{1V} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & m^2 \end{pmatrix}, \quad (12)$$

whose geometric representation is 3 non-overlapping points on a projective line. In the matrix A^{1V} , the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , the first column represents the power function $t_1^{D/2-1}$, the second column represents the power function $t_2^{-D/2}$, and the third column represents the power function $(z_{1,3}t_1 + z_{2,3}t_2)^{-1} = (t_1 + t_2 m^2)^{-1}$, respectively. In the above splitting coordinates, the Feynman integral of the 1-loop bubble satisfies the following GKZ-system

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,3}\}A_{1VA} &= -A_{1VA}, \\ \{\vartheta_{2,2} + \vartheta_{2,3}\}A_{1VA} &= -A_{1VA}, \\ \vartheta_{1,1}A_{1VA} &= \left(\frac{D}{2} - 1\right)A_{1VA}, \\ \vartheta_{2,2}A_{1VA} &= -\frac{D}{2}A_{1VA}, \\ \{\vartheta_{1,3} + \vartheta_{2,3}\}A_{1VA} &= -A_{1VA}, \end{aligned} \quad (13)$$

where the Euler operator $\vartheta_{i,j} = z_{i,j} \partial / \partial z_{i,j}$. The dual space of the above GKZ-system is the null space $\{0\}$. Thus the exponent matrix is written uniquely as

$$\begin{pmatrix} \frac{D}{2} - 1 & 0 & -\frac{D}{2} \\ 0 & -\frac{D}{2} & \frac{D}{2} - 1 \end{pmatrix}, \quad (14)$$

and the matrix of integer lattice is the zero matrix $0_{2 \times 3}$. Using the above matrices of exponent and integer lattice, one obtains the unique solution of the GKZ-system satisfied by $A_{1VA}(m^2)$ as

$$A_{1VA}(m^2) = \frac{C_{1VA}^{(12)}}{\Gamma(1 - \frac{D}{2})\Gamma(\frac{D}{2})} (m^2)^{D/2-1} \propto \frac{1^{-D/2}}{\Gamma(1 - \frac{D}{2})\Gamma(\frac{D}{2})} (m^2)^{D/2-1}, \quad (15)$$

with the integration constant

$$C_{1VA}^{(12)} = -\frac{i\Gamma^2(1 - \frac{D}{2})\Gamma(\frac{D}{2})}{(4\pi)^{D/2}} (\Lambda_{\text{RE}}^2)^{2-D/2}. \quad (16)$$

The above solution is obtained where we take the 1st and 2nd column vectors as the basis of the projective line. Because

$$\begin{aligned}\det A_{13}^{1V} &= \det \begin{pmatrix} 1 & 1 \\ 0 & m^2 \end{pmatrix} = m^2 \neq 0, \\ \det A_{23}^{1V} &= \det \begin{pmatrix} 0 & 1 \\ 1 & m^2 \end{pmatrix} = -1 \neq 0,\end{aligned}\tag{17}$$

we can also choose $\{(1, 0)^T, (1, m^2)^T\}$ or $\{(0, 1)^T, (1, m^2)^T\}$ as the basis of the projective line respectively. As we choose the vectors $\{(1, 0)^T, (1, m^2)^T\}$ as the basis of the projective line, the splitting local coordinate matrix is written as

$$\begin{pmatrix} 1 & -\frac{1}{m^2} & 0 \\ 0 & \frac{1}{m^2} & 1 \end{pmatrix}.\tag{18}$$

Using the unique matrix of exponents

$$\begin{pmatrix} \frac{D}{2} - 1 & -\frac{D}{2} & 0 \\ 0 & 0 & -1 \end{pmatrix},\tag{19}$$

and the integer lattice matrix $0_{2 \times 3}$, one obtains the single solution as

$$\begin{aligned}A_{1VA}(m^2) &= \frac{C_{1VA}^{(13)}}{\det(A_{13}^{1V})\Gamma(1 - \frac{D}{2})\Gamma(1)} \left(-\frac{1}{m^2}\right)^0 \left(\frac{1}{m^2}\right)^{-D/2} \\ &= \frac{C_{1VA}^{(13)}}{\Gamma(1 - \frac{D}{2})} (m^2)^{D/2-1}.\end{aligned}\tag{20}$$

Matching with Eq.(9), we get the integration constant as

$$C_{1VA}^{(13)} = -\frac{i\Gamma^2(1 - \frac{D}{2})}{(4\pi)^{D/2}} (\Lambda_{\text{RE}}^2)^{2-D/2}.\tag{21}$$

Similarly taking the vectors $\{(0, 1)^T, (1, m^2)^T\}$ as the basis of the projective line, we write the splitting local coordinate matrix as

$$\begin{pmatrix} -m^2 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}.\tag{22}$$

Using the matrix of exponents in Eq.(19) and the integer lattice $0_{2 \times 3}$, we write the solution as

$$\begin{aligned}A_{1VA}(m^2) &= \frac{C_{1VA}^{(23)}}{\det(A_{23}^{1V})\Gamma(\frac{D}{2})\Gamma(1)} (-m^2)^{D/2-1} \\ &= -\frac{(-)^{D/2-1} C_{1VA}^{(23)}}{\Gamma(\frac{D}{2})} (m^2)^{D/2-1}.\end{aligned}\tag{23}$$

From Eq.(9), the integration constant is given as

$$C_{1VA}^{(23)} = \frac{i(-1)^{1-D/2}\Gamma(1-\frac{D}{2})\Gamma(\frac{D}{2})}{(4\pi)^{D/2}}(\Lambda_{\text{RE}}^2)^{2-D/2}. \quad (24)$$

Note that on the Grassmannian $G_{2,3}$, there is only one choice of the exponent matrix compatible with the integer lattice $0_{2 \times 3}$. This is the reason why three different choices on the basis of the projective line induce the same solution of the GKZ-system which is proportional to $(m^2)^{D/2-1}$.

It should be emphasized that the matrix of splitting local coordinates in Eq.(12) is not uniquely determined. In fact, the columns of the matrix represent 3 distinct points on the projective line. Multiplying each column by a non-zero constant does not change the corresponding geometric representation of the matrix Eq.(12). Correspondingly those fundamental solutions are completely consistent with each other up to a constant scalar multiple.

In the following we apply this method to present the subvarieties of Grassmannians embedded by the Feynman integral of the 1-loop self-energy diagram, then construct the fundamental solution systems which are composed of linear independent hypergeometric functions in neighborhoods of the regular singularities. As mentioned above, the analytic expression of the Feynman integral is expressed as the linear combinations of these hypergeometric functions of the fundamental solution systems in different coordinate neighborhoods of regular singularities, whose combination coefficients are determined by the Feynman integral on an ordinary point or some regular singularities. In the dimension regularization, the Feynman integral of the 1-loop self-energy diagram is written as

$$A_{1SE}(p^2, m_1^2, m_2^2) = (\Lambda_{\text{RE}}^2)^{2-D/2} \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - m_1^2)((q+p)^2 - m_2^2)}. \quad (25)$$

Adopting Feynman parametric representation, we get the integral of zero virtual masses as

$$\begin{aligned} A_{1SE}(p^2, 0, 0) &= \frac{i(-1)^{D/2-2}\Gamma(2-\frac{D}{2})(\Lambda_{\text{RE}}^2)^{2-D/2}}{(4\pi)^{D/2}} \\ &\quad \times \int_0^1 dt_1 t_1^{D/2-2} (p^2 t_1 - p^2)^{D/2-2} \\ &= \frac{i(-1)^{D/2-2}\Gamma(2-\frac{D}{2})(\Lambda_{\text{RE}}^2)^{2-D/2}}{(4\pi)^{D/2}} \\ &\quad \times \int_0^1 \omega(t) t_1^{D/2-2} t_2^{2-D} (p^2 t_1 + p^2 t_2)^{D/2-2}. \end{aligned} \quad (26)$$

In the last equation $t_2 = -1$, and $\omega(t)$ is the volume element of the projective line. Eq.(26) can be embedded in the subvariety of the Grassmannian $G_{2,3}$ with splitting local coordinates as

$$A^{1SE} = \begin{pmatrix} 1 & 0 & p^2 \\ 0 & 1 & p^2 \end{pmatrix}. \quad (27)$$

In the matrix above, the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , the first column represents the power function $t_1^{D/2-2}$, the second column represents the power function t_2^{2-D} , and the third column represents the power function $(z_{1,3}t_1 + z_{2,3}t_2)^{D/2-2} = (t_1p^2 + t_2p^2)^{D/2-2}$, respectively. The Feynman integral satisfies the following GKZ-system

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,3}\}A_{1SE} &= -A_{1SE}, \\ \{\vartheta_{2,2} + \vartheta_{2,3}\}A_{1SE} &= -A_{1SE}, \\ \vartheta_{1,1}A_{1SE} &= \left(\frac{D}{2} - 2\right)A_{1SE}, \\ \vartheta_{2,2}A_{1SE} &= (2 - D)A_{1SE}, \\ \{\vartheta_{1,3} + \vartheta_{2,3}\}A_{1SE} &= \left(\frac{D}{2} - 2\right)A_{1SE}. \end{aligned} \quad (28)$$

Obviously the dual space of the GKZ-system of Eq.(28) is the null space $\{0\}$ which is consistent with the exponent matrix

$$\begin{pmatrix} \frac{D}{2} - 2 & 0 & 1 - \frac{D}{2} \\ 0 & 2 - D & D - 3 \end{pmatrix}. \quad (29)$$

Using the matrix of exponents, one obtains the solution of the GKZ-system of Eq.(28)

$$\begin{aligned} A_{1SE}(p^2, 0, 0) &= \frac{C_{1SE}^{(0)}}{\Gamma(2 - \frac{D}{2})\Gamma(D - 2)} (p^2)^{1-D/2} (p^2)^{D-3} \\ &= \frac{C_{1SE}^{(0)}}{\Gamma(2 - \frac{D}{2})\Gamma(D - 2)} (p^2)^{D/2-2}, \end{aligned} \quad (30)$$

with the integration constant

$$C_{1SE}^{(0)} = \frac{i\Gamma^2(2 - \frac{D}{2})\Gamma^2(1 - \frac{D}{2})(\Lambda_{RE}^2)^{2-D/2}}{(4\pi)^{D/2}}. \quad (31)$$

The solution Eq.(30) is obtained where we take the first and second column vectors of Eq.(27) as the basis of the projective line. Because

$$\begin{aligned}\det A_{13}^{1SE} &= \det \begin{pmatrix} 1 & p^2 \\ 0 & p^2 \end{pmatrix} = p^2 \neq 0, \\ \det A_{23}^{1SE} &= \det \begin{pmatrix} 0 & p^2 \\ 1 & p^2 \end{pmatrix} = -p^2 \neq 0,\end{aligned}\quad (32)$$

we can choose $\{(1, 0)^T, (p^2, p^2)^T\}$ or $\{(0, 1)^T, (p^2, p^2)^T\}$ as the basis of the projective line respectively. Using two different choices of the basis of the projective line, the solutions of the GKZ-system are all proportional to $(p^2)^{D/2-2}$.

Along with increasing of the number of independent arguments of the Feynman integral, the dimension of dual space of the GKZ-system increases. The bases of the dual space determine the PDEs satisfied by the Feynman integral in the combined local coordinates.

A. The fundamental solution systems as $m_1^2 = 0$

As $m_1^2 = 0$, $p^2 \neq 0$, and $m_2^2 \neq 0$, Feynman parametric representation of the 1-loop self energy is given as

$$\begin{aligned}A_{1SE}(p^2, 0, m_2^2) &= \frac{i\Gamma(2 - \frac{D}{2})(\Lambda_{\text{RE}}^2)^{2-D/2}}{(-)^{2-D/2}(4\pi)^{D/2}} \int_0^1 dt_1 dt_3 \frac{t_1^{D/2-2} \delta(t_1 + t_3 - 1)}{(t_3 p^2 - m_2^2)^{2-D/2}} \\ &= \frac{i\Gamma(2 - \frac{D}{2})(\Lambda_{\text{RE}}^2)^{2-D/2}}{(-)^{1-D/2}(4\pi)^{D/2}} \int \omega_3(t) \frac{t_1^{D/2-2} t_2^{2-D} \delta(t_1 + t_2 + t_3)}{(t_3 p^2 + t_2 m_2^2)^{2-D/2}},\end{aligned}\quad (33)$$

where

$$\omega_k(t) = \sum_{i=1}^k (-)^{i-1} t_i dt_1 \wedge \cdots \widehat{dt_i} \wedge \cdots dt_k \quad (34)$$

denotes the volume element of $(k-1)$ -dimension projective space, and the hat symbol implies that the wedge product does not involve dt_i . Eq.(33) can be taken as an integral on the subvariety of the Grassmannian $G_{3,5}$ with splitting local coordinates as

$$A^{(1)} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & m_2^2 \\ 0 & 0 & 1 & 1 & p^2 \end{pmatrix}. \quad (35)$$

Here the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , and the third row corresponds to the integration variable t_3 , respectively. Meanwhile the first column represents the power function $t_1^{D/2-2}$, the second column represents the power function t_2^{2-D} , the third column represents the power function $t_3^0 = 1$, the fourth column represents the function $\delta(t_1 + t_2 + t_3)$, and the fifth column represents $(z_{1,5}t_1 + z_{2,5}t_2 + z_{3,5}t_3)^{D/2-2} = (t_2m_2^2 + t_3p^2)^{D/2-2}$, respectively. The geometric description of the matrix of Eq.(35) can be drawn in 2-dimension projective plane [60] as Fig.1. Accordingly the Feynman integral satisfies the following GKZ-system in splitting




FIG. 1: The geometric description of the matrix in Eq.(35) on the projective plane CP^2 , where the points $1, \dots, 5$ denote the indices of columns of the 3×5 matrix.

local coordinates

$$\begin{aligned}
\{\vartheta_{1,1} + \vartheta_{1,4}\}A_{1SE}(p^2, 0, m_2^2) &= -A_{1SE}(p^2, 0, m_2^2), \\
\{\vartheta_{2,2} + \vartheta_{2,4} + \vartheta_{2,5}\}A_{1SE}(p^2, 0, m_2^2) &= -A_{1SE}(p^2, 0, m_2^2), \\
\{\vartheta_{3,3} + \vartheta_{3,4} + \vartheta_{3,5}\}A_{1SE}(p^2, 0, m_2^2) &= -A_{1SE}(p^2, 0, m_2^2), \\
\vartheta_{1,1}A_{1SE}(p^2, 0, m_2^2) &= \left(\frac{D}{2} - 2\right)A_{1SE}(p^2, 0, m_2^2), \\
\vartheta_{2,2}A_{1SE}(p^2, 0, m_2^2) &= (2 - D)A_{1SE}(p^2, 0, m_2^2), \\
\vartheta_{3,3}A_{1SE}(p^2, 0, m_2^2) &= 0,
\end{aligned}$$

$$\begin{aligned}
\{\vartheta_{1,4} + \vartheta_{2,4} + \vartheta_{3,4}\} A_{1SE}(p^2, 0, m_2^2) &= -A_{1SE}(p^2, 0, m_2^2), \\
\{\vartheta_{2,5} + \vartheta_{3,5}\} A_{1SE}(p^2, 0, m_2^2) &= \left(\frac{D}{2} - 2\right) A_{1SE}(p^2, 0, m_2^2),
\end{aligned} \tag{36}$$

which induces the exponent matrix as

$$\begin{pmatrix} \frac{D}{2} - 2 & 0 & 0 & 1 - \frac{D}{2} & 0 \\ 0 & 2 - D & 0 & \alpha_{2,4} & \alpha_{2,5} \\ 0 & 0 & 0 & \alpha_{3,4} & \alpha_{3,5} \end{pmatrix}. \tag{37}$$

Where the exponents $\alpha_{i,j}$ satisfy the following relations

$$\begin{aligned}
\alpha_{2,4} + \alpha_{2,5} &= D - 3, \quad \alpha_{3,4} + \alpha_{3,5} = -1, \\
\alpha_{2,4} + \alpha_{3,4} &= \frac{D}{2} - 2, \quad \alpha_{2,5} + \alpha_{3,5} = \frac{D}{2} - 2.
\end{aligned} \tag{38}$$

Furthermore, the dual space of the GKZ-system of Eq.(36) is spanned by a 3×5 matrix $(0_{3 \times 3} | E_3^{(1)})$ with

$$E_3^{(1)} = \begin{pmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{pmatrix}. \tag{39}$$

The base of the dual space implies the Feynman integral also satisfying the PDE as

$$\frac{\partial^2 A_{1SE}}{\partial z_{2,4} \partial z_{3,5}}(p^2, 0, m_2^2) = \frac{\partial^2 A_{1SE}}{\partial z_{2,5} \partial z_{3,4}}(p^2, 0, m_2^2). \tag{40}$$

When the exponent matrix is given, the Feynman integral can be formally expressed as

$$A_{1SE}(p^2, 0, m_2^2) = \prod_{i,j} z_{i,j}^{\alpha_{i,j}} \varphi\left(\frac{m_2^2}{p^2}\right), \tag{41}$$

where $\varphi(x)$ satisfies the PDE in the combined coordinate x as

$$\begin{aligned}
&x \left[\alpha_{2,4} \alpha_{3,5} + (1 - \alpha_{2,4} - \alpha_{3,5}) x \frac{d}{dx} - x^2 \frac{d^2}{dx^2} \right] \varphi(x) \\
&= \left[\alpha_{2,5} \alpha_{3,4} + (1 + \alpha_{2,5} + \alpha_{3,4}) x \frac{d}{dx} - x^2 \frac{d^2}{dx^2} \right] \varphi(x).
\end{aligned} \tag{42}$$

As $\alpha_{2,5} \alpha_{3,4} = 0$ this equation recovers the well-known Gauss equation. Note that the combined local coordinate $x = m_2^2/p^2 = z_{2,5} z_{3,4} / z_{2,4} z_{3,5}$ is determined by the base of the dual

space. The integer lattice $(0_{3 \times 3} | nE_3^{(1)})$ ($n \geq 0$) is compatible with two choices of the exponents. The first choice is written as

$$\alpha_{2,4} = 0, \alpha_{2,5} = D - 3, \alpha_{3,4} = \frac{D}{2} - 2, \alpha_{3,5} = 1 - \frac{D}{2}, \quad (43)$$

whose geometric description is presented as Fig.2(1) with $a = 3$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 5)\}$. The second choice is

$$\alpha_{2,4} = \frac{D}{2} - 1, \alpha_{2,5} = \frac{D}{2} - 2, \alpha_{3,4} = -1, \alpha_{3,5} = 0, \quad (44)$$

whose geometric description is presented as Fig.2(2) with $(a, b) = (2, 5)$, $(c, d, e) = (1, 3, 4)$.




FIG. 2: The geometric description of the single orbits in Eq.(43),Eq.(44),Eq.(47), and Eq.(48) on the projective plane CP^2 , where the points $1, \dots, 5$ denote the indices of columns of the 3×5 matrix.

Adopting the matrix of integer lattice $(0_{3 \times 3} | nE_3^{(1)})$ and the corresponding matrices of exponents, we obtain two linear independent hypergeometric functions as

$$\begin{aligned} \psi_{\{1,2,3\}}^{(1)}(p^2, 0, m_2^2) &\sim (p^2)^{1-D/2} (m_2^2)^{D-3} \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)}{n!} \left(\frac{p^2}{m_2^2}\right)^n \\ &\sim (p^2)^{1-D/2} (m_2^2 - p^2)^{D-3}, \\ \psi_{\{1,2,3\}}^{(2)}(p^2, 0, m_2^2) &\sim (m_2^2)^{D/2-2} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(\frac{D}{2} + n)} \left(\frac{p^2}{m_2^2}\right)^n, \end{aligned} \quad (45)$$

where the intersection of their convergent regions is $|p^2/m_2^2| \leq 1$. Two linearly independent Gaussian functions constitute the fundamental solution system of the GKZ-system of Eq.(36) in the region $|p^2/m_2^2| \leq 1$ which is consistent with the results in the literature. As $y_i = 0$ in

Eq.(46) of Ref. [41], the fourth type Appell functions of the last two terms are transformed into the Gauss functions above. In the first equation of Eq.(45), we apply the well-known reduction [70]

$${}_2F_1 \left(\begin{matrix} a, b \\ b \end{matrix} \middle| x \right) = (1-x)^{-a}. \quad (46)$$

Actually the last expression can be taken as the analytical continuation of the hypergeometric series in the whole complex plane except the regular singularities $p^2 = 0$ and $p^2 - m_2^2 = 0$.

Similarly the matrix of integer lattice $(0_{3 \times 3} | -nE_3^{(1)})$ ($n \geq 0$) permits to two possibilities of the exponents. The first possibility is

$$\alpha_{2,4} = D - 3, \alpha_{2,5} = 0, \alpha_{3,4} = 1 - \frac{D}{2}, \alpha_{3,5} = \frac{D}{2} - 2, \quad (47)$$

whose geometric description is presented as Fig.2(2) with $(a, b) = (3, 5)$, $(c, d, e) = (1, 2, 4)$.

The second possibility of the exponents is

$$\alpha_{2,4} = \frac{D}{2} - 2, \alpha_{2,5} = \frac{D}{2} - 1, \alpha_{3,4} = 0, \alpha_{3,5} = -1, \quad (48)$$

whose geometric description is drawn as Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, respectively. Similarly using the matrix of integer lattice and the matrices of exponents, one obtains two linear independent hypergeometric functions as

$$\begin{aligned} \psi_{\{1,2,3\}}^{(3)}(p^2, 0, m_2^2) &\sim (p^2)^{D/2-2} \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)}{n!} \left(\frac{m_2^2}{p^2}\right)^n \\ &\sim (p^2)^{1-D/2} (p^2 - m_2^2)^{D-3}, \\ \psi_{\{1,2,3\}}^{(4)}(p^2, 0, m_2^2) &\sim \frac{(m_2^2)^{D/2-1}}{p^2} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(\frac{D}{2} + n)} \left(\frac{m_2^2}{p^2}\right)^n, \end{aligned} \quad (49)$$

where the intersection of their convergent region is $|m_2^2/p^2| \leq 1$. These two linearly independent Gaussian functions constitute the fundamental solution system of the GKZ-system of Eq.(36) in the region $|m_2^2/p^2| \leq 1$ which is consistent with the results in the literature. Choosing $y_i = 0$ in Eq.(45) of Ref. [41], the fourth type Appell functions of the second and the fourth terms are reduced as the Gauss functions above.

If the GKZ-system of Eq.(36) is embedded in a compact n -dimensional manifold, we can only construct the hypergeometric series in neighborhoods of the regular singularities $m_2^2/p^2 = 0, \infty$. In order to construct the hypergeometric series in neighborhoods of the

regular singularity $m_2^2/p^2 = 1$, we embed the GKZ-system of Eq.(36) into the subvariety of the Grassmannian $G_{3,5}$ where the splitting local coordinates are presented in Eq.(35).

Because $\det(A_{\{1,2,5\}}^{(1)}) = p^2$ where the matrix $A_{\{1,2,5\}}^{(1)}$ denotes the submatrix of $A^{(1)}$ composed of the first-, second-, and fifth columns, one can choose the first, the second, and the fifth column vectors spanning the projective plane. Obviously we have

$$\left(A_{\{1,2,5\}}^{(1)}\right)^{-1} \cdot A^{(1)} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & -\frac{m_2^2}{p^2} & 1 - \frac{m_2^2}{p^2} & 0 \\ 0 & 0 & \frac{1}{p^2} & \frac{1}{p^2} & 1 \end{pmatrix}, \quad (50)$$

and the matrix of exponents

$$\begin{pmatrix} \frac{D}{2} - 2 & 0 & 0 & 1 - \frac{D}{2} & 0 \\ 0 & 2 - D & \alpha_{2,3} & \alpha_{2,4} & 0 \\ 0 & 0 & \alpha_{3,3} & \alpha_{3,4} & \frac{D}{2} - 2 \end{pmatrix}. \quad (51)$$

Where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{2,3} + \alpha_{2,4} &= D - 3, \quad \alpha_{3,3} + \alpha_{3,4} = 1 - \frac{D}{2}, \\ \alpha_{2,3} + \alpha_{3,3} &= 0, \quad \alpha_{2,4} + \alpha_{3,4} = \frac{D}{2} - 2. \end{aligned} \quad (52)$$

Correspondingly there are two choices on the matrix of integer lattice whose submatrix composed of the third and fourth columns is formulated as $\pm nE_3^{(1)}$ with integer $n \geq 0$, and other elements are all zero. The matrix of integer lattice $nE_3^{(1)}$ is compatible with two choices of the exponents. The first choice is

$$\alpha_{2,3} = 0, \quad \alpha_{2,4} = D - 3, \quad \alpha_{3,3} = 0, \quad \alpha_{3,4} = 1 - \frac{D}{2}, \quad (53)$$

whose geometric representation is presented by Fig.2(2) with $(a, b) = (3, 5)$, $(c, d, e) = (1, 2, 4)$. The second choice is

$$\alpha_{2,3} = \frac{D}{2} - 1, \quad \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{3,3} = 1 - \frac{D}{2}, \quad \alpha_{3,4} = 0, \quad (54)$$

whose geometric representation is drawn as Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, respectively. Adopting the matrix of integer lattice and the matrices of exponents, we obtain a fundamental solution system composed of two linear independent

hypergeometric functions as

$$\begin{aligned}
\psi_{\{1,2,5\}}^{(1)}(p^2, 0, m_2^2) &\sim (p^2)^{1-D/2}(p^2 - m_2^2)^{D-3} \\
&\times \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)}{n!\Gamma(1-n)\Gamma(2-\frac{D}{2}+n)} \left(\frac{m_2^2}{m_2^2-p^2}\right)^n \\
&\sim (p^2)^{1-D/2}(m_2^2-p^2)^{D-3}, \\
\psi_{\{1,2,5\}}^{(2)}(p^2, 0, m_2^2) &\sim \left(\frac{m_2^2}{p^2}\right)^{D/2-1}(m_2^2-p^2)^{D/2-2} \\
&\times \sum_{n=0}^{\infty} \frac{\Gamma(2-\frac{D}{2}+n)\Gamma(\frac{D}{2}-1+n)}{n!\Gamma(\frac{D}{2}+n)} \left(\frac{m_2^2}{m_2^2-p^2}\right)^n, \quad (55)
\end{aligned}$$

where the intersection of convergent regions of two hypergeometric functions is $|m_2^2/(m_2^2 - p^2)| \leq 1$. In fact the basis $\{1, 2, 5\}$ of the project plane composed of the first, second and fifth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 3\}$ through the permutation $(\widehat{35})$, accordingly the geometric description of the exponents in Eq.(53) is derived from that of the exponents in Eq.(47) through the permutation $(\widehat{35})$. Because the geometric description of Eq.(53) is same as that of Eq.(47), the corresponding function $\psi_{\{1,2,5\}}^{(1)}$ is proportionate to $\psi_{\{1,2,3\}}^{(3)}$ in the nonempty intersection of their convergent regions. Similarly the geometric description of the exponents in Eq.(54) is obtained through the permutation $(\widehat{35})$ from that of Eq.(48). Because the geometric description of Eq.(54) is same as that of Eq.(48), the function $\psi_{\{1,2,5\}}^{(2)}$ is proportionate to $\psi_{\{1,2,3\}}^{(4)}$ in the nonempty intersection of their convergent regions. Actually $\psi_{\{1,2,5\}}^{(2)} \sim \psi_{\{1,2,3\}}^{(4)}$ is consistent with the well-known relation [70]

$${}_2F_1\left(\begin{matrix} a, b \\ c \end{matrix} \middle| x\right) = (1-x)^{-a} {}_2F_1\left(\begin{matrix} a, c-b \\ c \end{matrix} \middle| \frac{x}{x-1}\right). \quad (56)$$

Similarly the matrix of integer lattice $-nE_3^{(1)}$ ($n \geq 0$) permits two possibilities of the exponents. The first choice is

$$\alpha_{2,3} = D-3, \alpha_{2,4} = 0, \alpha_{3,3} = 3-D, \alpha_{3,4} = \frac{D}{2} - 2, \quad (57)$$

whose geometric representation is presented in Fig.2(1) with $a = 5$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 3)\}$. And the second possibility is

$$\alpha_{2,3} = 0, \alpha_{2,4} = D-3, \alpha_{3,3} = 0, \alpha_{3,4} = 1 - \frac{D}{2}, \quad (58)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (2, 3)$, $(c, d, e) = (1, 4, 5)$, respectively. Adopting the matrix of integer lattice and the matrices of exponents,

we obtain a fundamental solution system composed of two linear independent hypergeometric functions as

$$\begin{aligned}\psi_{\{1,2,5\}}^{(3)}(p^2, 0, m_2^2) &\sim (m_2^2)^{D-3}(p^2)^{1-D/2} \\ &\quad \times \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)\Gamma(2-\frac{D}{2}+n)}{n!\Gamma(4-D+n)} \left(\frac{m_2^2-p^2}{m_2^2}\right)^n, \\ \psi_{\{1,2,5\}}^{(4)}(p^2, 0, m_2^2) &\sim (p^2)^{1-D/2}(m_2^2-p^2)^{D-3},\end{aligned}\tag{59}$$

whose convergent region is $|(m_2^2-p^2)/m_2^2| \leq 1$.

We take the first, third, and fifth columns as a basis spanning the projective plane because $\det(A_{\{1,3,5\}}^{(1)}) = -m_2^2$ where the matrix $A_{\{1,3,5\}}^{(1)}$ denotes the submatrix of $A^{(1)}$ composed of the first, third, and fifth columns. It is easily derived

$$\left(A_{\{1,3,5\}}^{(1)}\right)^{-1} \cdot A^{(1)} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & -\frac{p^2}{m_2^2} & 1 & 1-\frac{p^2}{m_2^2} & 0 \\ 0 & \frac{1}{m_2^2} & 0 & \frac{1}{m_2^2} & 1 \end{pmatrix}.\tag{60}$$

Accordingly the matrix of exponents is written as

$$\begin{pmatrix} \frac{D}{2}-2 & 0 & 0 & 1-\frac{D}{2} & 0 \\ 0 & \alpha_{2,2} & 0 & \alpha_{2,4} & 0 \\ 0 & \alpha_{3,2} & 0 & \alpha_{3,4} & \frac{D}{2}-2 \end{pmatrix},\tag{61}$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{2,2} + \alpha_{2,4} &= -1, \quad \alpha_{3,2} + \alpha_{3,4} = 1 - \frac{D}{2}, \\ \alpha_{2,2} + \alpha_{3,2} &= 2 - D, \quad \alpha_{2,4} + \alpha_{3,4} = \frac{D}{2} - 2.\end{aligned}\tag{62}$$

There are two choices on the matrix of integer lattice whose submatrix composed of the second and fourth columns is formulated as $\pm nE_3^{(1)}$ ($n \geq 0$), and other elements are all zero. The matrix of integer lattice $nE_3^{(1)}$ permits two possible choices of the exponents. The first choice is

$$\alpha_{2,2} = 0, \quad \alpha_{2,4} = -1, \quad \alpha_{3,2} = 2 - D, \quad \alpha_{3,4} = \frac{D}{2} - 1,\tag{63}$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (2, 5)$, $(c, d, e) = (1, 3, 4)$. The second choice of the exponents is

$$\alpha_{2,2} = 1 - \frac{D}{2}, \quad \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{3,2} = 1 - \frac{D}{2}, \quad \alpha_{3,4} = 0,\tag{64}$$

whose geometric representation is presented in Fig.2(1) with $a = 3$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 5)\}$, respectively. Adopting the matrix of integer lattice $nE_3^{(1)}$ and the matrices of exponents, we obtain a fundamental solution system composed of two hypergeometric solutions as

$$\begin{aligned} \psi_{\{1,3,5\}}^{(1)}(p^2, 0, m_2^2) &\sim \frac{(m_2^2)^{D/2-1}}{p^2 - m_2^2} \sum_{n=0}^{\infty} \frac{\Gamma(D-2+n)}{\Gamma(\frac{D}{2}+n)} \left(\frac{p^2}{p^2 - m_2^2}\right)^n, \\ \psi_{\{1,3,5\}}^{(2)}(p^2, 0, m_2^2) &\sim \left(\frac{m_2^2}{p^2}\right)^{D/2-1} (p^2 - m_2^2)^{D/2-2} \\ &\quad \times \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)\Gamma(\frac{D}{2} - 1 + n)}{n!\Gamma(2 - \frac{D}{2} + n)} \left(\frac{p^2}{p^2 - m_2^2}\right)^n \\ &\sim (p^2)^{1-D/2} (p^2 - m_2^2)^{D-3}, \end{aligned} \quad (65)$$

which are convergent in the region $|p^2/(p^2 - m_2^2)| \leq 1$. In fact the basis $\{1, 3, 5\}$ of the projective plane is obtained from the basis $\{1, 2, 3\}$ through the permutation $(\overline{25})$, accordingly the geometric description of the exponents Eq.(63) is derived from that of the exponents Eq.(44) through the permutation $(\overline{25})$. Because the geometric description of Eq.(63) is same as that of Eq.(44), the corresponding function $\psi_{\{1,2,5\}}^{(1)}$ is proportionate to $\psi_{\{1,2,3\}}^{(2)}$ in the nonempty intersection of their convergent regions. Similarly the geometric description of the exponents Eq.(64) is obtained through the permutation $(\overline{25})$ from that of Eq.(43). Because the geometric description of Eq.(64) is same as that of Eq.(43), the function $\psi_{\{1,2,5\}}^{(2)}$ is proportionate to $\psi_{\{1,2,3\}}^{(1)}$ in the nonempty intersection of their convergent regions. Actually $\psi_{\{1,3,5\}}^{(2)} \sim \psi_{\{1,2,3\}}^{(1)}$ is consistent with the well-known relation Eq.(56).

Similarly the matrix of integer lattice $-nE_3^{(1)}$ permits two possibilities of the exponents. The first one is

$$\alpha_{2,2} = -1, \alpha_{2,4} = 0, \alpha_{3,2} = 3 - D, \alpha_{3,4} = \frac{D}{2} - 2, \quad (66)$$

where the geometric representation is presented in Fig.2(1) with $a = 5$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 3)\}$. The second possibility is

$$\alpha_{2,2} = 2 - D, \alpha_{2,4} = D - 3, \alpha_{3,2} = 0, \alpha_{3,4} = 1 - \frac{D}{2}, \quad (67)$$

where the geometric representation is presented in Fig.2(2) with $(a, b) = (2, 3)$, $(c, d, e) = (1, 4, 5)$, respectively. Adopting the matrix of integer lattice and the matrices of exponents,

we obtain a fundamental solution system composed of two linear independent hypergeometric solutions as

$$\begin{aligned}\psi_{\{1,3,5\}}^{(3)}(p^2, 0, m_2^2) &\sim \frac{(m_2^2)^{D/2-1}}{p^2} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(4 - D + n)} \left(\frac{p^2 - m_2^2}{p^2}\right)^n, \\ \psi_{\{1,3,5\}}^{(4)}(p^2, 0, m_2^2) &\sim \frac{(p^2 - m_2^2)^{D-3} (m_2^2)^{D/2-1}}{(p^2)^{D-2}} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n)}{n!} \left(\frac{p^2 - m_2^2}{p^2}\right)^n \\ &\sim (p^2 - m_2^2)^{D-3} (p^2)^{1-D/2},\end{aligned}\quad (68)$$

whose convergent region is $|1 - m_2^2/p^2| \leq 1$. Certainly the basis $\{1, 3, 5\}$ of the projective plane composed of the first, third and fifth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 5\}$ through the permutation $(\widehat{23})$, accordingly the geometric description of the exponents Eq.(66) is derived from that of the exponents Eq.(57) through the permutation $(\widehat{23})$. Because the geometric description of Eq.(66) is same as that of Eq.(57), the function $\psi_{\{1,3,5\}}^{(3)}$ is proportionate to $\psi_{\{1,2,5\}}^{(3)}$ in the nonempty intersection of their convergent regions. Similarly the geometric description of the exponents Eq.(67) is obtained through the permutation $(\widehat{23})$ from that of Eq.(58). Because the geometric description of Eq.(67) is same as that of Eq.(58), the function $\psi_{\{1,3,5\}}^{(4)}$ is proportionate to $\psi_{\{1,2,5\}}^{(4)}$ in the nonempty intersection of their convergent regions. Actually $\psi_{\{1,3,5\}}^{(3)} \sim \psi_{\{1,2,5\}}^{(3)}$ is consistent with the well-known relation Eq.(56).

Because $\det(A_{\{2,3,4\}}^{(1)}) = 1$ where the matrix $A_{\{2,3,4\}}^{(1)}$ denotes the submatrix of $A^{(1)}$ composed of the second, third, and fourth columns, one derives

$$\left(A_{\{2,3,4\}}^{(1)}\right)^{-1} \cdot A^{(1)} = \begin{pmatrix} -1 & 1 & 0 & 0 & m_2^2 \\ -1 & 0 & 1 & 0 & p^2 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}. \quad (69)$$

Correspondingly the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 2 - D & 0 & 0 & \alpha_{1,5} \\ \alpha_{2,1} & 0 & 0 & 0 & \alpha_{2,5} \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}, \quad (70)$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,1} + \alpha_{1,5} &= D - 3, \quad \alpha_{2,1} + \alpha_{2,5} = -1, \\ \alpha_{1,1} + \alpha_{2,1} &= \frac{D}{2} - 2, \quad \alpha_{1,5} + \alpha_{2,5} = \frac{D}{2} - 2.\end{aligned}\quad (71)$$

There are two choices on the matrix of integer lattice whose submatrix composed of the first and fifth columns is formulated as $\pm nE_3^{(3)}$ ($n \geq 0$) with

$$E_3^{(3)} = \begin{pmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{pmatrix}, \quad (72)$$

and other elements are all zero. The matrix of integer lattice $nE_3^{(3)}$ is compatible with two choices of the exponents. The first choice is

$$\alpha_{1,1} = 0, \alpha_{1,5} = D - 3, \alpha_{2,1} = \frac{D}{2} - 2, \alpha_{2,5} = 1 - \frac{D}{2}, \quad (73)$$

whose geometric representation is presented in Fig.2(1) with $a = 3$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 5)\}$. The second choice is

$$\alpha_{1,1} = \frac{D}{2} - 1, \alpha_{1,5} = \frac{D}{2} - 2, \alpha_{2,1} = -1, \alpha_{2,5} = 0. \quad (74)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (2, 5)$, $(c, d, e) = (1, 3, 4)$, respectively. Adopting the matrix of integer lattice $nE_3^{(3)}$ and the matrices of exponents, we obtain a fundamental solution system composed by two linear independent hypergeometric functions as

$$\begin{aligned} \psi_{\{2,3,4\}}^{(1)}(p^2, 0, m_2^2) &\sim (p^2)^{1-D/2} (m_2^2)^{D-3} \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)}{n!} \left(\frac{p^2}{m_2^2}\right)^n \\ &\sim (p^2)^{1-D/2} (m_2^2 - p^2)^{D-3}, \\ \psi_{\{2,3,4\}}^{(2)}(p^2, 0, m_2^2) &\sim (m_2^2)^{D/2-2} \sum_{n=0}^{\infty} \frac{\Gamma(2-\frac{D}{2}+n)}{\Gamma(\frac{D}{2}+n)} \left(\frac{p^2}{m_2^2}\right)^n, \end{aligned} \quad (75)$$

which are convergent in the region is $|p^2/m_2^2| \leq 1$. Similarly the basis $\{2, 3, 4\}$ of the projective plane composed of the second, third and fourth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 3\}$ through the permutation $(\widehat{14})$, or equivalently from the basis $\{1, 3, 5\}$ through the permutation $(\widehat{14})(25)$. Accordingly the geometric description of Eq.(73) is taken from that of Eq.(43) through the permutation $(\widehat{14})$, or from that of Eq.(64) through the permutation $(\widehat{14})(25)$. Because the geometric description of Eq.(73) is same as that of Eq.(43) and Eq.(64), the functions $\psi_{\{1,2,3\}}^{(1)} \sim \psi_{\{1,2,5\}}^{(2)} \sim \psi_{\{2,3,4\}}^{(1)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of Eq.(74) is taken from that of Eq.(44) through the permutation $(\widehat{14})$, or from that of Eq.(63) through

the permutation $(14)\widehat{(25)}$. Because the geometric description of Eq.(74) is same as that of Eq.(44) and Eq.(63), the functions $\psi_{\{1,2,3\}}^{(2)} \sim \psi_{\{1,2,5\}}^{(1)} \sim \psi_{\{2,3,4\}}^{(2)}$ in nonempty intersection of their convergent regions.

The matrix of integer lattice $-nE_3^{(3)}$ permits two possibilities of the exponents. The first possibility is given as

$$\alpha_{1,1} = D - 3, \alpha_{1,5} = 0, \alpha_{2,1} = 1 - \frac{D}{2}, \alpha_{2,5} = \frac{D}{2} - 2, \quad (76)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (3, 5)$, $(c, d, e) = (1, 2, 4)$. Meanwhile the second possibility is

$$\alpha_{1,1} = \frac{D}{2} - 2, \alpha_{1,5} = \frac{D}{2} - 1, \alpha_{2,1} = 0, \alpha_{2,5} = -1, \quad (77)$$

whose geometric representation is drawn in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, respectively. Adopting the matrix of integer lattice $-nE_3^{(3)}$ and the matrices of exponents, we construct a fundamental solution system composed of two linear independent hypergeometric functions as

$$\begin{aligned} \psi_{\{2,3,4\}}^{(3)}(p^2, 0, m_2^2) &\sim (p^2)^{D/2-2} \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)}{n!} \left(\frac{m_2^2}{p^2}\right)^n, \\ &\sim (p^2)^{1-D/2} (p^2 - m_2^2)^{D-3}, \\ \psi_{\{2,3,4\}}^{(4)}(p^2, 0, m_2^2) &\sim \frac{(m_2^2)^{D/2-1}}{p^2} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(\frac{D}{2} + n)} \left(\frac{m_2^2}{p^2}\right)^n, \end{aligned} \quad (78)$$

which are convergent in the region is $|m_2^2/p^2| \leq 1$. Similarly the basis $\{2, 3, 4\}$ of the projective plane composed of the second, third and fourth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 3\}$ through the permutation $(\widehat{14})$, or equivalently from the basis $\{1, 2, 5\}$ through the permutation $(14)\widehat{(35)}$. Similarly the geometric description of Eq.(76) is taken from that of Eq.(47) through the permutation $(\widehat{14})$, or from that of Eq.(53) through the permutation $(14)\widehat{(25)}$. Because the geometric description of Eq.(76) is same as that of Eq.(47) and Eq.(53), the functions $\psi_{\{1,2,3\}}^{(3)} \sim \psi_{\{1,2,5\}}^{(1)} \sim \psi_{\{2,3,4\}}^{(3)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of Eq.(77) is derived from that of Eq.(48) through the permutation $(\widehat{14})$, or from that of Eq.(54) through the permutation $(14)\widehat{(25)}$. Because the geometric description of Eq.(77) is same as that of Eq.(48) and Eq.(54), the functions $\psi_{\{1,2,3\}}^{(4)} \sim \psi_{\{1,2,5\}}^{(2)} \sim \psi_{\{2,3,4\}}^{(4)}$ in nonempty intersection of their convergent regions.

Because $\det(A_{\{2,4,5\}}^{(1)}) = -p^2$ where the matrix $A_{\{2,4,5\}}^{(1)}$ denotes the submatrix of $A^{(1)}$ composed of the second, fourth, and fifth columns, one derives

$$\left(A_{\{2,4,5\}}^{(1)}\right)^{-1} \cdot A^{(1)} = \begin{pmatrix} -1 + \frac{m_2^2}{p^2} & 1 & -\frac{m_2^2}{p^2} & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ -\frac{1}{p^2} & 0 & \frac{1}{p^2} & 0 & 1 \end{pmatrix}. \quad (79)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 2 - D & \alpha_{1,3} & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ \alpha_{3,1} & 0 & \alpha_{3,3} & 0 & \frac{D}{2} - 2 \end{pmatrix}, \quad (80)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,3} &= D - 3, \quad \alpha_{3,1} + \alpha_{3,3} = 1 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{3,1} &= \frac{D}{2} - 2, \quad \alpha_{1,3} + \alpha_{3,3} = 0. \end{aligned} \quad (81)$$

Correspondingly there are two choices on the matrix of integer lattice whose submatrix composed of the first and third columns is formulated as $\pm nE_3^{(2)}$ ($n \geq 0$), with

$$E_3^{(2)} = \begin{pmatrix} 1 & -1 \\ 0 & 0 \\ -1 & 1 \end{pmatrix}, \quad (82)$$

and other elements are all zero. The matrix of integer lattice $nE_3^{(2)}$ permits two possibilities of the exponents. The first possibility is

$$\alpha_{1,1} = 0, \quad \alpha_{1,3} = D - 3, \quad \alpha_{3,1} = \frac{D}{2} - 2, \quad \alpha_{3,3} = 3 - D, \quad (83)$$

whose geometric representation is presented in Fig.2(1) with $a = 5$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 3)\}$. The second possibility is

$$\alpha_{1,1} = D - 3, \quad \alpha_{1,3} = 0, \quad \alpha_{3,1} = 1 - \frac{D}{2}, \quad \alpha_{3,3} = 0, \quad (84)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (2, 3)$, $(c, d, e) = (1, 4, 5)$, respectively. Adopting the matrix of integer lattice and the matrices of exponents,

we obtain a fundamental solution system composed of two linear independent hypergeometric solutions as

$$\begin{aligned} \psi_{\{2,4,5\}}^{(1)}(p^2, 0, m_2^2) &\sim (m_2^2)^{D-3}(p^2)^{1-D/2} \\ &\times \sum_{n=0}^{\infty} \frac{\Gamma(3-D+n)\Gamma(2-\frac{D}{2}+n)}{n!\Gamma(4-D+n)} \left(\frac{m_2^2-p^2}{m_2^2}\right)^n, \\ \psi_{\{2,4,5\}}^{(2)}(p^2, 0, m_2^2) &\sim (p^2)^{1-D/2}(m_2^2-p^2)^{D-3}, \end{aligned} \quad (85)$$

whose convergent region is $|(m_2^2-p^2)/m_2^2| \leq 1$. Similarly the basis $\{2, 4, 5\}$ of the projective plane composed of the second, fourth and fifth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 5\}$ through the permutation $\widehat{(14)}$, or equivalently obtained from the basis $\{1, 3, 5\}$ through the permutation $(14)\widehat{(23)}$. Accordingly the geometric description of Eq.(83) is obtained from that of Eq.(57) through the permutation $\widehat{(14)}$, or obtained from that of Eq.(66) through the permutation $(14)\widehat{(23)}$. Because the geometric description of Eq.(83) is same as that of Eq.(57) and Eq.(66), the functions $\psi_{\{1,2,5\}}^{(3)} \sim \psi_{\{1,3,5\}}^{(3)} \sim \psi_{\{2,4,5\}}^{(1)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of Eq.(84) is obtained from that of Eq.(58) through the permutation $\widehat{(14)}$, or obtained from that of Eq.(67) through the permutation $(14)\widehat{(23)}$. Because the geometric description of Eq.(84) is same as that of Eq.(58) and Eq.(67), the functions $\psi_{\{1,2,5\}}^{(4)} \sim \psi_{\{1,3,5\}}^{(4)} \sim \psi_{\{2,4,5\}}^{(2)}$ in nonempty intersection of their convergent regions. Including appropriate scalar factors, each hypergeometric function is analytic continuation of other two functions in the union of their convergent regions.

The matrix of integer lattice $-nE_3^{(2)}$ permits two choices of the exponents. The first choice is

$$\alpha_{1,1} = D - 3, \quad \alpha_{1,3} = 0, \quad \alpha_{3,1} = 1 - \frac{D}{2}, \quad \alpha_{3,3} = 0, \quad (86)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (3, 5)$, $(c, d, e) = (1, 2, 4)$. Meanwhile the second choice is

$$\alpha_{1,1} = \frac{D}{2} - 2, \quad \alpha_{1,3} = \frac{D}{2} - 1, \quad \alpha_{3,1} = 0, \quad \alpha_{3,3} = 1 - \frac{D}{2}, \quad (87)$$

whose geometric representation is presented in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, respectively. Adopting the matrix of integer lattice and the matrices of

exponents, we obtain a fundamental solution system composed of two linear independent hypergeometric functions as

$$\begin{aligned} \psi_{\{2,4,5\}}^{(3)}(p^2, 0, m_2^2) &\sim (p^2)^{1-D/2}(m_2^2 - p^2)^{D-3}, \\ \psi_{\{2,4,5\}}^{(4)}(p^2, 0, m_2^2) &\sim \left(\frac{m_2^2}{p^2}\right)^{D/2-1}(m_2^2 - p^2)^{D/2-2} \\ &\quad \times \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)\Gamma(\frac{D}{2} - 1 + n)}{n!\Gamma(\frac{D}{2} + n)} \left(\frac{m_2^2}{m_2^2 - p^2}\right)^n, \end{aligned} \quad (88)$$

which are convergent in the region $|m_2^2/(m_2^2 - p^2)| \leq 1$. Because the basis $\{2, 4, 5\}$ of the projective plane composed of the second, fourth and fifth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 3\}$ through the permutation $(14)\widehat{(35)}$, or equivalently obtained from the basis $\{1, 2, 5\}$ through the permutation $\widehat{(14)}$. Accordingly the geometric description of Eq.(86) is taken from that of Eq.(47) through the permutation $(14)\widehat{(35)}$, or taken from that of Eq.(53) through the permutation $\widehat{(14)}$. Since the geometric description of Eq.(86) is same as that of Eq.(47) and Eq.(53), the functions $\psi_{\{1,2,3\}}^{(1)} \sim \psi_{\{1,2,5\}}^{(1)} \sim \psi_{\{2,4,5\}}^{(3)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of Eq.(87) is taken from that of Eq.(48) through the permutation $(14)\widehat{(35)}$, or taken from that of Eq.(54) through the permutation $\widehat{(14)}$. Similarly the geometric description of Eq.(87) is same as that of Eq.(48) and Eq.(54), the functions $\psi_{\{1,2,3\}}^{(2)} \sim \psi_{\{1,2,5\}}^{(2)} \sim \psi_{\{2,4,5\}}^{(4)}$ in nonempty intersection of their convergent regions.

Because $\det(A_{\{3,4,5\}}^{(1)}) = m_2^2$ where the matrix $A_{\{3,4,5\}}^{(1)}$ denotes the submatrix of $A^{(1)}$ composed of the third, fourth, and fifth columns, we get

$$\left(A_{\{3,4,5\}}^{(1)}\right)^{-1} \cdot A^{(1)} = \begin{pmatrix} -1 + \frac{p^2}{m_2^2} & -\frac{p^2}{m_2^2} & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ -\frac{1}{m_2^2} & \frac{1}{m_2^2} & 0 & 0 & 1 \end{pmatrix}. \quad (89)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ \alpha_{3,1} & \alpha_{3,2} & 0 & 0 & \frac{D}{2} - 2 \end{pmatrix}, \quad (90)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,2} &= -1, \quad \alpha_{3,1} + \alpha_{3,2} = 1 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{3,1} &= \frac{D}{2} - 2, \quad \alpha_{1,2} + \alpha_{3,2} = 2 - D. \end{aligned} \quad (91)$$

Correspondingly there are two choices on the matrix of integer lattice whose submatrix composed of the first and second columns is formulated as $\pm nE_3^{(2)}$ ($n \geq 0$), and other elements are all zero. The matrix of integer lattice $nE_3^{(2)}$ permits two possibilities of the exponents. The first possibility is

$$\alpha_{1,1} = 0, \alpha_{1,2} = -1, \alpha_{3,1} = \frac{D}{2} - 2, \alpha_{3,2} = 3 - D, \quad (92)$$

whose geometric representation is presented in Fig.2(1) with $a = 5$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 3)\}$. The second possibility is

$$\alpha_{1,1} = D - 3, \alpha_{1,2} = 2 - D, \alpha_{3,1} = 1 - \frac{D}{2}, \alpha_{3,2} = 0, \quad (93)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (2, 3)$, $(c, d, e) = (1, 4, 5)$, respectively. Adopting the matrix of integer lattice $nE_3^{(2)}$ and the matrices of exponents, we obtain a fundamental solution system composed of two linear independent hypergeometric solutions as

$$\begin{aligned} \psi_{\{3,4,5\}}^{(1)}(p^2, 0, m_2^2) &\sim \frac{(m_2^2)^{D/2-1}}{p^2} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(4 - D + n)} \left(\frac{p^2 - m_2^2}{p^2}\right)^n, \\ \psi_{\{3,4,5\}}^{(2)}(p^2, 0, m_2^2) &\sim \frac{(p^2 - m_2^2)^{D-3} (m_2^2)^{D/2-1}}{(p^2)^{D-2}} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n)}{n!} \left(\frac{p^2 - m_2^2}{p^2}\right)^n \\ &\sim (p^2 - m_2^2)^{D-3} (p^2)^{1-D/2}, \end{aligned} \quad (94)$$

which are convergent in the region $|1 - m_2^2/p^2| \leq 1$. Obviously the basis $\{3, 4, 5\}$ of the projective plane composed of the third, fourth and fifth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 5\}$ through the permutation $(14)\widehat{(23)}$, or equivalently obtained from the basis $\{1, 3, 5\}$ through the permutation $\widehat{(14)}$, or equivalently obtained from the basis $\{2, 3, 5\}$ through the permutation $\widehat{(23)}$, respectively. Accordingly the geometric description of Eq.(92) is taken from that of Eq.(57) through the permutation $(14)\widehat{(23)}$, or taken from that of Eq.(66) through the permutation $\widehat{(14)}$, or taken from that of Eq.(83) through the permutation $\widehat{(23)}$, respectively. Because the geometric description of Eq.(92) is same as that of Eq.(57), Eq.(66), and Eq.(83), the functions $\psi_{\{1,2,5\}}^{(3)} \sim \psi_{\{1,3,5\}}^{(3)} \sim \psi_{\{2,4,5\}}^{(1)} \sim \psi_{\{3,4,5\}}^{(1)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of Eq.(93) is obtained from that of Eq.(58) through the permutation $(14)\widehat{(23)}$, or obtained from that of Eq.(67) through the permutation $\widehat{(14)}$, or obtained from that of Eq.(84) through

the permutation $(\widehat{23})$, respectively. Because the geometric description of Eq.(93) is same as that of Eq.(58), Eq.(67), and Eq.(84), the functions $\psi_{\{1,2,5\}}^{(4)} \sim \psi_{\{1,3,5\}}^{(4)} \sim \psi_{\{2,4,5\}}^{(2)} \sim \psi_{\{3,4,5\}}^{(2)}$ in nonempty intersection of their convergent regions. Including appropriate scalar factors, each hypergeometric function is analytic continuation of other functions in the union of their convergent regions.

The matrix of integer lattice $-nE_3^{(2)}$ permits two choices of the exponents. The first choice is

$$\alpha_{1,1} = -1, \alpha_{1,2} = 0, \alpha_{3,1} = \frac{D}{2} - 1, \alpha_{3,2} = 2 - D, \quad (95)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (2, 5)$, $(c, d, e) = (1, 3, 4)$. And the second choice is

$$\alpha_{1,1} = \frac{D}{2} - 2, \alpha_{1,2} = 1 - \frac{D}{2}, \alpha_{3,1} = 0, \alpha_{3,2} = 1 - \frac{D}{2}. \quad (96)$$

whose geometric representation is presented in Fig.2(1) with $a = 3$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 5)\}$, respectively. Adopting the matrix of integer lattice $-nE_3^{(2)}$ and the matrices of exponents, we obtain a fundamental solution system composed of two hypergeometric solutions as

$$\begin{aligned} \psi_{\{3,4,5\}}^{(3)}(p^2, 0, m_2^2) &\sim \frac{(m_2^2)^{D/2-1}}{p^2 - m_2^2} \sum_{n=0}^{\infty} \frac{\Gamma(D-2+n)}{\Gamma(\frac{D}{2}+n)} \left(\frac{p^2}{p^2 - m_2^2}\right)^n, \\ \psi_{\{3,4,5\}}^{(4)}(p^2, 0, m_2^2) &\sim \left(\frac{m_2^2}{p^2}\right)^{D/2-1} (p^2 - m_2^2)^{D/2-2} \\ &\quad \times \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2}-1+n)}{n!} \left(\frac{p^2}{p^2 - m_2^2}\right)^n \\ &= (p^2)^{1-D/2} (p^2 - m_2^2)^{D-3}, \end{aligned} \quad (97)$$

which are convergent in the region $|p^2/(p^2 - m_2^2)| \leq 1$. Since the basis $\{3, 4, 5\}$ of the projective plane composed of the third, fourth and fifth columns of the matrix Eq.(35) is obtained from the basis $\{1, 2, 3\}$ through the permutation $(14)(25)$, or equivalently obtained from the basis $\{1, 3, 5\}$ through the permutation $(\widehat{14})$, or equivalently obtained from the basis $\{2, 3, 4\}$ through the permutation $(\widehat{25})$, respectively. Accordingly the geometric description of Eq.(95) is taken from that of Eq.(44) through the permutation $(14)(25)$, or taken from that of Eq.(63) through the permutation $(\widehat{14})$, or taken from that of Eq.(74) through the permutation $(\widehat{25})$, respectively. Because the geometric description of Eq.(95) is same as

that of Eq.(44), Eq.(63), and Eq.(74), the functions $\psi_{\{1,2,3\}}^{(2)} \sim \psi_{\{1,3,5\}}^{(1)} \sim \psi_{\{2,3,4\}}^{(2)} \sim \psi_{\{3,4,5\}}^{(3)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of Eq.(96) is taken from that of Eq.(43) through the permutation $(14)\widehat{(25)}$, or taken from that of Eq.(64) through the permutation $(\widehat{14})$, or taken from that of Eq.(73) through the permutation $(\widehat{25})$, respectively. Because the geometric description of Eq.(96) is same as that of Eq.(43), Eq.(64), and Eq.(73), the functions $\psi_{\{1,2,3\}}^{(1)} \sim \psi_{\{1,3,5\}}^{(2)} \sim \psi_{\{2,3,4\}}^{(1)} \sim \psi_{\{3,4,5\}}^{(4)}$ in nonempty intersection of their convergent regions.

B. The fundamental solution system in general case

As the 1-loop self energy diagram involves 2 nonzero virtual masses, the integer lattice is spanned by two linear independent vectors, and the number of exponential matrices compatible with them also increases accordingly.

Using Feynman parametric representation, we write the 1-loop self energy in the general case as

$$\begin{aligned}
A^{1S}(p^2, m_1^2, m_2^2) &= \left(\Lambda_{\text{RE}}^2\right)^{2-D/2} \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - m_1^2)((q+p)^2 - m_2^2)} \\
&= \frac{i\left(\Lambda_{\text{RE}}^2\right)^{2-D/2} \Gamma(2 - \frac{D}{2})}{(-)^{2-D/2} (4\pi)^{D/2}} \int_0^1 dt_1 dt_2 \delta(t_1 + t_2 - 1) \\
&\quad \times (t_1 t_2 p^2 - t_1 m_1^2 - t_2 m_2^2)^{D/2-2} \\
&= -\frac{i\left(\Lambda_{\text{RE}}^2\right)^{2-D/2} \Gamma(2 - \frac{D}{2})}{(-)^{2-D/2} (4\pi)^{D/2}} \int \omega_3(t) \delta(t_1 + t_2 + t_3) \\
&\quad \times t_3^{2-D} (t_1 t_2 p^2 + t_1 t_3 m_1^2 + t_2 t_3 m_2^2)^{D/2-2}. \tag{98}
\end{aligned}$$

The integral can be embedded in the subvariety of the Grassmannian $G_{3,6}$ where the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , and the third row corresponds to the integration variable t_3 , respectively. Meanwhile the first column represents the power function $t_1^0 = 1$, the second column represents the power function $t_2^0 = 1$, the third column represents the power function t_3^{2-D} , the fourth column represents the function $\delta(t_1 + t_2 + t_3)$. In order to embed the homogeneous polynomial $(t_1 t_2 p^2 + t_1 t_3 m_1^2 + t_2 t_3 m_2^2)$ as the fifth and sixth columns of the Grassmannian $G_{3,6}$, we rewrite

$$t_1 t_2 p^2 + t_1 t_3 m_1^2 + t_2 t_3 m_2^2$$

$$= z_{1, \widehat{\sigma}_1(5)} z_{2, \widehat{\sigma}_1(6)} t_1 t_2 + z_{1, \widehat{\sigma}_2(5)} z_{3, \widehat{\sigma}_2(6)} t_1 t_3 + z_{2, \widehat{\sigma}_3(5)} z_{3, \widehat{\sigma}_3(6)} t_2 t_3 , \quad (99)$$

where $\widehat{\sigma}_i$, ($i = 1, 2, 3$) are elements of the permutation group $S_2 = \{\widehat{e}, \widehat{(56)}\}$ on the column indices 5, 6. Taking $\widehat{\sigma}_1 = \widehat{\sigma}_3 = \widehat{(56)}$, $\widehat{\sigma}_2 = \widehat{e}$, we have

$$z_{1,6} z_{2,5} = p^2 , \quad z_{1,5} z_{3,6} = m_1^2 , \quad z_{2,6} z_{3,5} = m_2^2 . \quad (100)$$

The solution of Eq.(100)

$$z_{1,5} = z_{2,5} = z_{3,5} = 1 , \quad z_{1,6} = p^2 , \quad z_{3,6} = m_1^2 , \quad z_{2,6} = m_2^2 , \quad (101)$$

indicates that the splitting local coordinates are written as

$$A^{(1S)} = \begin{pmatrix} 1 & 0 & 0 & 1 & 1 & p^2 \\ 0 & 1 & 0 & 1 & 1 & m_2^2 \\ 0 & 0 & 1 & 1 & 1 & m_1^2 \end{pmatrix} , \quad (102)$$

whose geometric description is drawn in Fig.3. Taking Feynman parametric representation




FIG. 3: The geometric description of the matrix in Eq.(102) on the projective plane CP^2 , where the points $1, \dots, 6$ denote the indices of columns of the 3×6 matrix, and 4, 5 denote a double point.

of the 1-loop self energy as a function of the splitting local coordinates,

$$A^{1S}(p^2, m_1^2, m_2^2) = \Psi^{1S}(z_1, \dots, z_6) , \quad (103)$$

we obviously have

$$\Psi^{1S}(\chi_1 z_1, \dots, \chi_6 z_6) = \chi_3^{2-D} \chi_4^{-1} \chi_5^{D/2-2} \chi_6^{D/2-2} \Psi^{1S}(z_1, \dots, z_6). \quad (104)$$

Where z_i ($i = 1, \dots, 6$) denotes the i -th column vector in the matrix Eq.(102), and χ_i is a nonzero constant, respectively. If we choose the first, second and third column vectors of $A^{(1S)}$ as the basis of the projective plane, constructed hypergeometric functions contain redundant dependent variables. However $\det(A_{\{1,2,4\}}^{(1S)}) = 1$ where the matrix $A_{\{1,2,4\}}^{(1S)}$ denotes the submatrix of $A^{(1S)}$ composed of the first, second, and fourth columns, one derives

$$Z_{3 \times 6} = \left(A_{\{1,2,4\}}^{(1S)} \right)^{-1} \cdot A^{(1S)} = \begin{pmatrix} 1 & 0 & -1 & 0 & 0 & p^2 - m_1^2 \\ 0 & 1 & -1 & 0 & 0 & m_2^2 - m_1^2 \\ 0 & 0 & 1 & 1 & 1 & m_1^2 \end{pmatrix}. \quad (105)$$

Accordingly, the Feynman integral satisfies the following GKZ-system in splitting local coordinates

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,3} + \vartheta_{1,6}\} A_{1SE}(p^2, m_1^2, m_2^2) &= -A_{1SE}(p^2, m_1^2, m_2^2), \\ \{\vartheta_{2,2} + \vartheta_{2,3} + \vartheta_{2,6}\} A_{1SE}(p^2, m_1^2, m_2^2) &= -A_{1SE}(p^2, m_1^2, m_2^2), \\ \{\vartheta_{3,3} + \vartheta_{3,4} + \vartheta_{3,5} + \vartheta_{3,6}\} A_{1SE}(p^2, m_1^2, m_2^2) &= -A_{1SE}(p^2, m_1^2, m_2^2), \\ \vartheta_{1,1} A_{1SE}(p^2, m_1^2, m_2^2) &= 0, \\ \vartheta_{2,2} A_{1SE}(p^2, m_1^2, m_2^2) &= 0, \\ \{\vartheta_{1,3} + \vartheta_{2,3} + \vartheta_{3,3}\} A_{1SE}(p^2, m_1^2, m_2^2) &= (2 - D) A_{1SE}(p^2, m_1^2, m_2^2), \\ \vartheta_{3,4} A_{1SE}(p^2, m_1^2, m_2^2) &= -A_{1SE}(p^2, m_1^2, m_2^2), \\ \vartheta_{3,5} A_{1SE}(p^2, m_1^2, m_2^2) &= \left(\frac{D}{2} - 2\right) A_{1SE}(p^2, m_1^2, m_2^2), \\ \{\vartheta_{1,6} + \vartheta_{2,6} + \vartheta_{3,6}\} A_{1SE}(p^2, m_1^2, m_2^2) &= \left(\frac{D}{2} - 2\right) A_{1SE}(p^2, m_1^2, m_2^2). \end{aligned} \quad (106)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & 0 & \alpha_{1,3} & 0 & 0 & \alpha_{1,6} \\ 0 & 0 & \alpha_{2,3} & 0 & 0 & \alpha_{2,6} \\ 0 & 0 & \alpha_{3,3} & -1 & \frac{D}{2} - 2 & \alpha_{3,6} \end{pmatrix}, \quad (107)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,3} + \alpha_{1,6} &= -1, \quad \alpha_{2,3} + \alpha_{2,6} = -1, \quad \alpha_{3,3} + \alpha_{3,6} = 2 - \frac{D}{2}, \\ \alpha_{1,3} + \alpha_{2,3} + \alpha_{3,3} &= 2 - D, \quad \alpha_{1,6} + \alpha_{2,6} + \alpha_{3,6} = \frac{D}{2} - 2. \end{aligned} \quad (108)$$

Correspondingly there are 12 choices on the matrix of integer lattice whose submatrix composed of the third and sixth columns is formulated as $\pm n_1 E_3^{(i)} \pm n_2 E_3^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 3), (2, 3)\}$, and other elements are all zero. Furthermore, the matrix

$$E_3^{(1)} = \begin{pmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad (109)$$

and the matrices $E_3^{(2)}$, $E_3^{(3)}$ are presented in Eq.(82) and Eq.(72), respectively. Note that $E_3^{(3)} = E_3^{(2)} - E_3^{(1)}$, two linear independent bases of integer lattice matrices $E_3^{(1)}$, $E_3^{(2)}$ induce two independent PDEs as:

$$\begin{aligned} \frac{\partial^2 A^{(1S)}}{\partial z_{1,3} \partial z_{3,6}}(p^2, m_1^2, m_2^2) &= \frac{\partial^2 A^{(1S)}}{\partial z_{1,6} \partial z_{3,3}}(p^2, m_1^2, m_2^2), \\ \frac{\partial^2 A^{(1S)}}{\partial z_{2,3} \partial z_{3,6}}(p^2, m_1^2, m_2^2) &= \frac{\partial^2 A^{(1S)}}{\partial z_{2,6} \partial z_{3,3}}(p^2, m_1^2, m_2^2). \end{aligned} \quad (110)$$

When the exponential matrix is given, the Feynman integral can be formally expressed as

$$A_{1SE}(p^2, m_1^2, m_2^2) = \prod_{i,j} z_{i,j}^{\alpha_{i,j}} \varphi\left(1 - \frac{p^2}{m_1^2}, 1 - \frac{m_2^2}{m_1^2}\right), \quad (111)$$

where $(i, j) \in \{(1, 3), (1, 6), (2, 3), (2, 6), (3, 3), (3, 6)\}$, and $\varphi(x_1, x_2)$ satisfies the PDEs in the combined local coordinates $x_1 = z_{1,6} z_{3,3} / (z_{1,3} z_{3,6}) = 1 - p^2/m_1^2$, $x_2 = z_{2,6} z_{3,3} / (z_{2,3} z_{3,6}) = 1 - m_2^2/m_1^2$ as

$$\begin{aligned} &x_1 \left[\alpha_{1,3} \alpha_{3,6} + (1 - \alpha_{1,3} - \alpha_{3,6}) x_1 \frac{\partial}{\partial x_1} - \alpha_{1,3} x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_1^2 \frac{\partial^2}{\partial x_1^2} \right] \varphi(x_1, x_2) \\ &= \left[\alpha_{1,6} \alpha_{3,3} + (1 + \alpha_{1,6} + \alpha_{3,3}) x_1 \frac{\partial}{\partial x_1} + \alpha_{1,6} x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_1^2 \frac{\partial^2}{\partial x_1^2} \right] \varphi(x_1, x_2), \\ &x_2 \left[\alpha_{2,3} \alpha_{3,6} - \alpha_{2,3} x_1 \frac{\partial}{\partial x_1} + (1 - \alpha_{2,3} - \alpha_{3,6}) x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right] \varphi(x_1, x_2) \\ &= \left[\alpha_{2,6} \alpha_{3,3} + \alpha_{2,6} x_1 \frac{\partial}{\partial x_1} + (1 + \alpha_{2,6} + \alpha_{3,3}) x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right] \varphi(x_1, x_2). \end{aligned} \quad (112)$$

As $\alpha_{1,6} = \alpha_{2,6} = 0$, the PDEs are changed into the first type Appell PDEs. The holonomic rank of the ideal composed of the PDEs is 3, which indicates the fundamental solution system constituted by three linear independent hypergeometric functions in neighborhoods of regular singularities. It is known that the fourth type Appell function cannot be embedded into Grassmannians [14]. Nevertheless, including the appropriate power factor and the

transformation of the variables, one proves that the Feynman integral presented by the fourth Appell functions [41, 42] satisfies the above PDEs. Those matrices of integer lattice permit 12 possibilities of the exponents. Corresponding to the integer lattice $n_1 E_3^{(3)} + n_2 E_3^{(2)}$ the choice of exponents is written as

$$\alpha_{1,3} = 1 - \frac{D}{2}, \alpha_{1,6} = \frac{D}{2} - 2, \alpha_{2,3} = -1, \alpha_{2,6} = 0, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,6} = 0, \quad (113)$$

whose geometric representation is presented in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 6), (4, 5)\}$,




FIG. 4: The geometric descriptions of the single orbits in Eq.(105) on the projective plane CP^2 , where the points $1, \dots, 6$ denote the indices of columns of the 3×6 matrix.

$\{e, f\} = \{2, 3\}$. As the integer lattice is chosen as $n_1 E_3^{(3)} + n_2 E_3^{(1)}$, the compatible matrix of exponents is

$$\alpha_{1,3} = 0, \alpha_{1,6} = -1, \alpha_{2,3} = -\frac{D}{2}, \alpha_{2,6} = \frac{D}{2} - 1, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,6} = 0, \quad (114)$$

whose geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 2$, $d = 3$, $\{e, f\} = \{1, 6\}$. The exponents of the integer lattice $n_1 E_3^{(2)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,6} = -1, \alpha_{2,3} = 0, \alpha_{2,6} = -1, \alpha_{3,3} = 2 - D, \alpha_{3,6} = \frac{D}{2}, \quad (115)$$

whose geometric representation is presented in Fig.4(4) with $(a, b, c) = (3, 4, 5)$, $\{d, e, f\} =$

$\{1, 2, 6\}$. Accordingly three linear independent hypergeometric functions are obtained

$$\begin{aligned}
\psi_{\{1,2,4\}}^{(1)}(p^2, m_1^2, m_2^2) &\sim (m_1^2 - p^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - m_2^2}{m_1^2 - p^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - p^2}\right)^{n_2} \\
&\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(2)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D/2-1}}{p^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - m_2^2}{m_1^2 - p^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - m_2^2}\right)^{n_2} \\
&\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(3)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_1^2)^{D/2}}{(p^2 - m_1^2)(m_2^2 - m_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1 + n_2)}{\Gamma(\frac{D}{2} + 1 + n_1 + n_2)} \\
&\quad \times \left(\frac{m_1^2}{m_1^2 - p^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - m_2^2}\right)^{n_2}, \tag{116}
\end{aligned}$$

whose intersection of their convergent regions is nonempty proper subset of the whole parameter space. Three hypergeometric functions constitute a fundamental solution system of the GKZ-system in the nonempty intersection.

Similarly the exponents of the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(2)}$ are

$$\alpha_{1,3} = 2 - D, \alpha_{1,6} = D - 3, \alpha_{2,3} = 0, \alpha_{2,6} = -1, \alpha_{3,3} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \tag{117}$$

whose geometric representation is drawn in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 3), (4, 5)\}$, $\{e, f\} = \{2, 6\}$. Corresponding to the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(1)}$ the matrix of exponents is

$$\alpha_{1,3} = -1, \alpha_{1,6} = 0, \alpha_{2,3} = 3 - D, \alpha_{2,6} = D - 4, \alpha_{3,3} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \tag{118}$$

whose geometric representation is given in Fig.4(2) with $(a, b) = (4, 5)$, $c = 2$, $d = 6$, $\{e, f\} = \{1, 3\}$. Meanwhile the exponents of the integer lattice $-n_1 E_3^{(2)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,6} = 0, \alpha_{2,3} = -1, \alpha_{2,6} = 0, \alpha_{3,3} = 4 - D, \alpha_{3,6} = \frac{D}{2} - 2, \tag{119}$$

whose geometric representation is presented in Fig.4(4) with $(a, b, c) = (4, 5, 6)$, $\{d, e, f\} =$

$\{1, 2, 3\}$. Correspondingly the linear independent hypergeometric functions are

$$\begin{aligned}
\psi_{\{1,2,4\}}^{(4)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_1^2)^{2-D/2}(m_1^2 - p^2)^{D-3}}{m_2^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - p^2}{m_1^2 - m_2^2}\right)^{n_1} \left(\frac{m_1^2 - p^2}{m_1^2}\right)^{n_2} \\
&\sim \frac{(m_1^2 - p^2)^{D-3}(p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(5)}(p^2, m_1^2, m_2^2) &\sim (m_2^2 - m_1^2)^{D-4}(m_1^2)^{2-D/2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - p^2}{m_1^2 - m_2^2}\right)^{n_1} \left(\frac{m_1^2 - m_2^2}{m_1^2}\right)^{n_2} \\
&\sim \frac{(m_2^2 - m_1^2)^{D-3}(m_2^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(6)}(p^2, m_1^2, m_2^2) &\sim (m_1^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{\Gamma(5 - D + n_1 + n_2)} \\
&\quad \times \left(\frac{m_1^2 - p^2}{p^2}\right)^{n_1} \left(\frac{m_1^2 - m_2^2}{p^2}\right)^{n_2},
\end{aligned} \tag{120}$$

whose intersection of their convergent regions is a nonempty proper subset of the parameter space. The hypergeometric functions constitute a fundamental solution system of the GKZ-system in a nonempty proper subset of the whole parameter space.

The exponents of the integer lattice $n_1 E_3^{(3)} - n_2 E_3^{(2)}$ are

$$\alpha_{1,3} = 3 - D, \alpha_{1,6} = D - 4, \alpha_{2,3} = -1, \alpha_{2,6} = 0, \alpha_{3,3} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \tag{121}$$

whose geometric representation is plotted in Fig.4(2) with $(a, b) = (4, 5)$, $c = 1$, $d = 6$, $\{e, f\} = \{2, 3\}$. In a similar way the exponent matrix of the integer lattice $n_1 E_3^{(3)} - n_2 E_3^{(1)}$ is

$$\alpha_{1,3} = 0, \alpha_{1,6} = -1, \alpha_{2,3} = 2 - D, \alpha_{2,6} = D - 3, \alpha_{3,3} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \tag{122}$$

whose geometric representation is shown in Fig.4(1) with $\{(a, b), (c, d)\} = \{(2, 3), (4, 5)\}$, $\{e, f\} = \{1, 6\}$. Meanwhile the exponent matrix of $n_1 E_3^{(2)} - n_2 E_3^{(1)}$ is given by

$$\alpha_{1,3} = 0, \alpha_{1,6} = -1, \alpha_{2,3} = -1, \alpha_{2,6} = 0, \alpha_{3,3} = 3 - D, \alpha_{3,6} = \frac{D}{2} - 1, \tag{123}$$

whose geometric representation is presented in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 6), (2, 3)\}$. Accordingly the hypergeometric functions are

$$\begin{aligned}
\psi_{\{1,2,4\}}^{(7)}(p^2, m_1^2, m_2^2) &\sim (m_1^2)^{2-D/2} (p^2 - m_1^2)^{D-4} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - m_2^2}{m_1^2 - p^2}\right)^{n_1} \left(\frac{m_1^2 - p^2}{m_1^2}\right)^{n_2} \\
&\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(8)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{2-D/2}}{p^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - m_2^2}{m_1^2 - p^2}\right)^{n_1} \left(\frac{m_1^2 - m_2^2}{m_1^2}\right)^{n_2} \\
&\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(9)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_1^2)^{D/2-1}}{p^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 3 + n_1 - n_2)}{\Gamma(\frac{D}{2} + n_1 - n_2)} \\
&\quad \times \left(\frac{m_1^2}{m_1^2 - p^2}\right)^{n_1} \left(\frac{m_1^2 - m_2^2}{m_1^2}\right)^{n_2}. \tag{124}
\end{aligned}$$

Although the intersection of the convergent regions of these series is the empty set, the sets of the hypergeometric functions $\{\psi_{\{1,2,4\}}^{(1)}, \psi_{\{1,2,4\}}^{(8)}, \psi_{\{1,2,4\}}^{(9)}\}$ and $\{\psi_{\{1,2,4\}}^{(6)}, \psi_{\{1,2,4\}}^{(7)}, \psi_{\{1,2,4\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Corresponding to the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(2)}$, the matrix of exponents is

$$\alpha_{1,3} = -\frac{D}{2}, \alpha_{1,6} = \frac{D}{2} - 1, \alpha_{2,3} = 0, \alpha_{2,6} = -1, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,6} = 0, \tag{125}$$

whose geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 1$, $d = 3$, $\{e, f\} = \{2, 6\}$. Similarly the exponents of the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,6} = 0, \alpha_{2,3} = 1 - \frac{D}{2}, \alpha_{2,6} = \frac{D}{2} - 2, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,6} = 0, \tag{126}$$

whose geometric representation is plotted in Fig.4(1) with $\{(a, b), (c, d)\} = \{(2, 6), (4, 5)\}$, $\{e, f\} = \{1, 3\}$. In addition the exponents of the integer lattice $-n_1 E_3^{(2)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,6} = 0, \alpha_{2,3} = 0, \alpha_{2,6} = -1, \alpha_{3,3} = 3 - D, \alpha_{3,6} = \frac{D}{2} - 1, \tag{127}$$

whose geometric representation is shown in Fig.4(3) with $(a, b) = (4, 5)$, $c = 2$, $d = 3$, $\{(c, d), (e, f)\} = \{(1, 3), (2, 6)\}$. We obtain the hypergeometric functions according to the exponents and integer lattices as

$$\begin{aligned}
\psi_{\{1,2,4\}}^{(10)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D/2-1}}{m_2^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - p^2}{m_1^2 - m_2^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - p^2}\right)^{n_2} \\
&\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(11)}(p^2, m_1^2, m_2^2) &\sim (m_2^2 - m_1^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\
&\quad \times \left(\frac{m_1^2 - p^2}{m_1^2 - m_2^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - m_2^2}\right)^{n_2} \\
&\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(12)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_1^2)^{D/2-1}}{m_2^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1 - n_2)}{\Gamma(4 - D + n_1 - n_2)} \\
&\quad \times \left(\frac{m_1^2 - p^2}{m_1^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - m_2^2}\right)^{n_2}. \tag{128}
\end{aligned}$$

Certainly the intersection of the convergent regions of these series is the empty set. However, the sets of the hypergeometric functions $\{\psi_{\{1,2,4\}}^{(3)}, \psi_{\{1,2,4\}}^{(10)}, \psi_{\{1,2,4\}}^{(11)}\}$ and $\{\psi_{\{1,2,4\}}^{(4)}, \psi_{\{1,2,4\}}^{(11)}, \psi_{\{1,2,4\}}^{(12)}\}$ constitute the fundamental solution systems of the corresponding nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{\{1,3,4\}}^{(1S)}) = -1$ where the matrix $A_{\{1,3,4\}}^{(1S)}$ denotes the submatrix of $A^{(1S)}$ composed of the first, third, and fourth columns, one derives

$$\left(A_{\{1,3,4\}}^{(1S)}\right)^{-1} \cdot A^{(1S)} = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & m_1^2 - m_2^2 \\ 0 & -1 & 1 & 0 & 0 & p^2 - m_2^2 \\ 0 & 1 & 0 & 1 & 1 & m_2^2 \end{pmatrix}. \tag{129}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & 0 & 0 & 0 & \alpha_{1,6} \\ 0 & \alpha_{2,2} & 2 - D & 0 & 0 & \alpha_{2,6} \\ 0 & \alpha_{3,2} & 0 & -1 & \frac{D}{2} - 2 & \alpha_{3,6} \end{pmatrix}, \tag{130}$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,2} + \alpha_{1,6} &= -1, \quad \alpha_{2,2} + \alpha_{2,6} = D - 3, \quad \alpha_{3,2} + \alpha_{3,6} = 2 - \frac{D}{2}, \\ \alpha_{1,2} + \alpha_{2,2} + \alpha_{3,2} &= 0, \quad \alpha_{1,6} + \alpha_{2,6} + \alpha_{3,6} = \frac{D}{2} - 2.\end{aligned}\quad (131)$$

There are 12 choices on the matrix of integer lattice whose the submatrix composed of the second and sixth columns is formulated as $\pm n_1 E_3^{(i)} \pm n_2 E_3^{(j)}$, where $n_{1,2} \geq 0$, and $(i, j) = \{(1, 2), (1, 3), (2, 3)\}$, and other elements are all zero.

Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents. Corresponding to the integer lattice $n_1 E_3^{(3)} + n_2 E_3^{(2)}$, the exponents are

$$\alpha_{1,2} = 1 - \frac{D}{2}, \quad \alpha_{1,6} = \frac{D}{2} - 2, \quad \alpha_{2,2} = D - 3, \quad \alpha_{2,6} = 0, \quad \alpha_{3,2} = 2 - \frac{D}{2}, \quad \alpha_{3,6} = 0 \quad (132)$$

whose geometric representation is drawn in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 6), (4, 5)\}$, $\{e, f\} = \{2, 3\}$. The exponents of the integer lattice $n_1 E_3^{(3)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = 0, \quad \alpha_{1,6} = -1, \quad \alpha_{2,2} = \frac{D}{2} - 2, \quad \alpha_{2,6} = \frac{D}{2} - 1, \quad \alpha_{3,2} = 2 - \frac{D}{2}, \quad \alpha_{3,6} = 0, \quad (133)$$

whose geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 3$, $d = 2$, $\{e, f\} = \{1, 6\}$. In a similar way the exponents of the integer lattice $n_1 E_3^{(2)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = 0, \quad \alpha_{1,6} = -1, \quad \alpha_{2,2} = 0, \quad \alpha_{2,6} = D - 3, \quad \alpha_{3,2} = 0, \quad \alpha_{3,6} = 2 - \frac{D}{2}, \quad (134)$$

whose geometric representation is drawn in Fig.4(4) with $(a, b, c) = (2, 4, 5)$, $\{d, e, f\} = \{1, 3, 6\}$. Accordingly the hypergeometric functions are

$$\begin{aligned}\psi_{\{1,3,4\}}^{(1)}(p^2, m_1^2, m_2^2) &\sim (p^2 - m_2^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - D + n_1) \Gamma(\frac{D}{2} - 2 + n_2)}{n_1! n_2!} \\ &\quad \times \left(\frac{m_2^2 - m_1^2}{m_2^2 - p^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - p^2}\right)^{n_2} \\ &\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\ \psi_{\{1,3,4\}}^{(2)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D/2-1}}{p^2 - m_2^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 - n_2)}{n_2! \Gamma(\frac{D}{2} + n_1 - n_2)} \\ &\quad \times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{m_2^2 - m_1^2}{m_2^2 - p^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_1^2}\right)^{n_2}, \\ \psi_{\{1,3,4\}}^{(3)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2)^{2-D/2} (m_2^2 - m_1^2)^{D-3}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(4 - D + n_2)}{n_2! \Gamma(1 - n_1 - n_2)}\end{aligned}$$

$$\begin{aligned} & \times \frac{1}{\Gamma(3 - \frac{D}{2} + n_1 + n_2)} \left(\frac{m_2^2}{m_2^2 - p^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_1^2}\right)^{n_2} \\ & \sim \frac{(m_2^2)^{2-D/2} (m_2^2 - m_1^2)^{D-3}}{m_2^2 - p^2}, \end{aligned} \quad (135)$$

whose intersection is a nonempty proper subset of the whole parameter space. In other words, three hypergeometric functions constitute a fundamental solution system of the GKZ-system in the nonempty intersection.

The matrix of exponents of the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(2)}$ is

$$\alpha_{1,2} = 0, \alpha_{1,6} = -1, \alpha_{2,2} = 0, \alpha_{2,6} = D - 3, \alpha_{3,2} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \quad (136)$$

whose geometric representation is shown in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 2), (4, 5)\}$, $\{e, f\} = \{3, 6\}$. Similarly the exponents of the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,6} = 0, \alpha_{2,2} = 1, \alpha_{2,6} = D - 4, \alpha_{3,2} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \quad (137)$$

whose geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 3$, $d = 6$, $\{e, f\} = \{1, 2\}$. Furthermore the exponents of the integer lattice $-n_1 E_3^{(2)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,6} = 0, \alpha_{2,2} = D - 3, \alpha_{2,6} = 0, \alpha_{3,2} = 4 - D, \alpha_{3,6} = \frac{D}{2} - 2, \quad (138)$$

where the geometric representation is plotted in Fig.4(4) with $(a, b, c) = (4, 5, 6)$, $\{d, e, f\} = \{1, 2, 3\}$. The corresponding hypergeometric functions are presented as

$$\begin{aligned} \psi_{\{1,3,4\}}^{(4)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_1^2 - m_2^2)^{D-3} (m_2^2)^{2-D/2}}{p^2 - m_2^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - D + n_1)}{n_1! n_2!} \\ & \quad \times \Gamma\left(\frac{D}{2} - 2 + n_2\right) \left(\frac{m_2^2 - p^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2 - p^2}{m_2^2}\right)^{n_2} \\ & \sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\ \psi_{\{1,3,4\}}^{(5)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^D}{(m_2^2)^{D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + n_1 + n_2) \Gamma(1 + n_2)}{\Gamma(3 + n_1 + n_2) \Gamma(D - 1 + n_2)} \\ & \quad \times \left(\frac{m_2^2 - p^2}{m_2^2}\right)^{n_1} \left(\frac{m_2^2 - m_1^2}{m_2^2}\right)^{n_2}, \\ \psi_{\{1,3,4\}}^{(6)}(p^2, m_1^2, m_2^2) & \sim (m_2^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{n_2! \Gamma(5 - D + n_1 + n_2)} \\ & \quad \times \Gamma(3 - D + n_2) \left(\frac{m_2^2 - p^2}{m_2^2}\right)^{n_1} \left(\frac{m_2^2 - m_1^2}{m_2^2}\right)^{n_2}, \end{aligned} \quad (139)$$

whose intersection is a nonempty proper subset of the whole parameter space. Since the differences of the exponents between Eq.(136) and Eq.(137) are all integers, the hypergeometric solution of the exponents of Eq.(137) is written as

$$\begin{aligned}
\psi_{\{1,3,4\}}^{(5)'}(p^2, m_1^2, m_2^2) &\sim (m_2^2 - m_1^2)^{D-4} (m_2^2)^{2-D/2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(4 - D + n_1 - n_2)}{n_2! \Gamma(2 + n_1 - n_2)} \\
&\quad \times \Gamma\left(\frac{D}{2} - 2 + n_2\right) \left(\frac{m_2^2 - p^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2 - m_1^2}{m_2^2}\right)^{n_2} \\
&\sim -\psi_{\{1,3,4\}}^{(4)}(p^2, m_1^2, m_2^2) \\
&\quad + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(2 - D)} \psi_{\{1,3,4\}}^{(5)}(p^2, m_1^2, m_2^2). \tag{140}
\end{aligned}$$

For the integer lattice $n_1 E_3^{(3)} - n_2 E_3^{(2)}$, those exponents are written as

$$\alpha_{1,2} = 3 - D, \alpha_{1,6} = D - 4, \alpha_{2,2} = D - 3, \alpha_{2,6} = 0, \alpha_{3,2} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \tag{141}$$

whose geometric representation is drawn in Fig.4(2) with $(a, b) = (4, 5)$, $c = 1$, $d = 6$, $\{e, f\} = \{2, 3\}$. Corresponding to the integer lattice $n_1 E_3^{(3)} - n_2 E_3^{(1)}$, the exponents are

$$\alpha_{1,2} = 0, \alpha_{1,6} = -1, \alpha_{2,2} = 0, \alpha_{2,6} = D - 3, \alpha_{3,2} = 0, \alpha_{3,6} = 2 - \frac{D}{2}, \tag{142}$$

where the geometric representation of the exponent matrix is presented in Fig.4(1) with $\{(a, b), (c, d)\} = \{(2, 3), (4, 5)\}$, $\{e, f\} = \{1, 6\}$. For the integer lattice $n_1 E_3^{(2)} - n_2 E_3^{(1)}$, the matrix of exponents is given by

$$\alpha_{1,2} = 0, \alpha_{1,6} = -1, \alpha_{2,2} = D - 3, \alpha_{2,6} = 0, \alpha_{3,2} = 3 - D, \alpha_{3,6} = \frac{D}{2} - 1, \tag{143}$$

whose geometric representation is shown in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 6), (2, 3)\}$.

Correspondingly the hypergeometric functions are

$$\begin{aligned}
\psi_{\{1,3,4\}}^{(7)}(p^2, m_1^2, m_2^2) &\sim (p^2 - m_2^2)^{D-4} (m_2^2)^{2-D/2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - D + n_1) \Gamma(\frac{D}{2} - 2 + n_2)}{n_1! n_2!} \\
&\quad \times \left(\frac{m_2^2 - m_1^2}{m_2^2 - p^2}\right)^{n_1} \left(\frac{m_2^2 - p^2}{m_2^2}\right)^{n_2} \\
&\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,2,4\}}^{(8)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(\frac{D}{2} - 2 + n_2)}{n_2! \Gamma(1 - n_1 - n_2)}
\end{aligned}$$

$$\begin{aligned}
& \times \frac{1}{\Gamma(D-2+n_1+n_2)} \left(\frac{m_2^2-m_1^2}{m_2^2-p^2}\right)^{n_1} \left(\frac{m_2^2-m_1^2}{m_2^2}\right)^{n_2} \\
& \sim \frac{(m_2^2-m_1^2)^{D-3}(m_2^2)^{2-D/2}}{m_2^2-p^2}, \\
\psi_{\{1,3,4\}}^{(9)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2)^{D/2-1}}{m_2^2-p^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D-3+n_1-n_2)\Gamma(3-D+n_2)}{n_2! \Gamma(\frac{D}{2}+n_1-n_2)} \\
& \times \left(\frac{m_2^2}{m_2^2-p^2}\right)^{n_1} \left(\frac{m_2^2-m_1^2}{m_2^2}\right)^{n_2} \\
& \sim \frac{(m_2^2)^{D/2-1}(m_1^2-p^2)^{D-3}}{(m_2^2-p^2)^{D-2}} \\
& \times \sum_{n_1=0}^{\infty} \frac{\Gamma(D-3+n_1)}{\Gamma(\frac{D}{2}+n_1)} \left(\frac{m_2^2}{m_2^2-p^2}\right)^{n_1}, \tag{144}
\end{aligned}$$

Although the intersection of the convergent regions of these series is the empty set, the sets of the hypergeometric functions $\{\psi_{\{1,3,4\}}^{(1)}, \psi_{\{1,3,4\}}^{(8)}, \psi_{\{1,3,4\}}^{(9)}\}$ and $\{\psi_{\{1,3,4\}}^{(6)}, \psi_{\{1,3,4\}}^{(7)}, \psi_{\{1,3,4\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

In a similar way, the matrix of exponents of the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(2)}$ is

$$\alpha_{1,2} = \frac{D}{2} - 2, \quad \alpha_{1,6} = 1 - \frac{D}{2}, \quad \alpha_{2,2} = 0, \quad \alpha_{2,6} = D - 3, \quad \alpha_{3,2} = 2 - \frac{D}{2}, \quad \alpha_{3,6} = 0 \tag{145}$$

whose geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 1$, $d = 2$, $\{e, f\} = \{3, 6\}$. The exponents of the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = -1, \quad \alpha_{1,6} = 0, \quad \alpha_{2,2} = \frac{D}{2} - 1, \quad \alpha_{2,6} = \frac{D}{2} - 2, \quad \alpha_{3,2} = 2 - \frac{D}{2}, \quad \alpha_{3,6} = 0, \tag{146}$$

whose geometric representation is drawn in Fig.4(1) with $\{(a, b), (c, d)\} = \{(3, 6), (4, 5)\}$, $\{e, f\} = \{1, 2\}$. Finally the choice on exponents of the integer lattice $-n_1 E_3^{(2)} + n_2 E_3^{(1)}$ is

$$\alpha_{1,2} = -1, \quad \alpha_{1,6} = 0, \quad \alpha_{2,2} = 0, \quad \alpha_{2,6} = D - 3, \quad \alpha_{3,2} = 1, \quad \alpha_{3,6} = 1 - \frac{D}{2}, \tag{147}$$

whose geometric representation is presented in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 2), (3, 6)\}$. Accordingly the hypergeometric functions are written as

$$\begin{aligned}
\psi_{\{1,3,4\}}^{(10)}(p^2, m_1^2, m_2^2) & \sim (p^2 - m_2^2)^{1-D/2} (m_1^2 - m_2^2)^{D-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3-D+n_1)}{n_1! n_2!} \\
& \times \Gamma\left(\frac{D}{2} - 2 + n_2\right) \left(\frac{m_2^2 - p^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - p^2}\right)^{n_2}
\end{aligned}$$

$$\begin{aligned}
& \sim \frac{(p^2 - m_1^2)^{D-3}(p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,3,4\}}^{(11)}(p^2, m_1^2, m_2^2) & \sim (m_2^2 - m_1^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{n_2! \Gamma(\frac{D}{2} + n_1 + n_2)} \\
& \quad \times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{m_2^2 - p^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_1^2}\right)^{n_2}, \\
\psi_{\{1,3,4\}}^{(12)}(p^2, m_1^2, m_2^2) & \sim (m_2^2)^{3-D/2} (m_2^2 - m_1^2)^{D-5} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(5 - D + n_1 + n_2)}{\Gamma(3 + n_1 + n_2)} \\
& \quad \times \frac{\Gamma(1 + n_2)}{\Gamma(4 - \frac{D}{2} + n_2)} \left(\frac{m_2^2 - p^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_1^2}\right)^{n_2}, \tag{148}
\end{aligned}$$

Certainly the intersection of the convergent regions of these series is a nonempty proper subset of the whole parameter space. Furthermore the sets of the hypergeometric functions $\{\psi_{\{1,3,4\}}^{(3)}, \psi_{\{1,3,4\}}^{(10)}, \psi_{\{1,3,4\}}^{(11)}\}$ and $\{\psi_{\{1,3,4\}}^{(4)}, \psi_{\{1,3,4\}}^{(11)}, \psi_{\{1,3,4\}}^{(12)}\}$ also constitute the fundamental solution systems of the corresponding nonempty proper subsets of the whole parameter space, respectively. Similarly the hypergeometric solution of the exponents of Eq.(147) is

$$\begin{aligned}
\psi_{\{1,3,4\}}^{(12)'}(p^2, m_1^2, m_2^2) & \sim (m_2^2)^{1-D/2} (m_2^2 - p^2)^{D-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n_1 - n_2)}{n_2! \Gamma(2 + n_1 - n_2)} \\
& \quad \times \Gamma(3 - D + n_2) \left(\frac{m_2^2 - m_1^2}{m_2^2}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - p^2}\right)^{n_2} \\
& \sim -\psi_{\{1,3,4\}}^{(10)}(p^2, m_1^2, m_2^2) \\
& \quad + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(\frac{D}{2} - 3)} \psi_{\{1,3,4\}}^{(12)}(p^2, m_1^2, m_2^2). \tag{149}
\end{aligned}$$

Because $\det(A_{\{2,3,4\}}^{(1S)}) = 1$, we choose the second, third, fourth column vectors composing of the basis of the projective plane, and find

$$\left(A_{\{2,3,4\}}^{(1S)}\right)^{-1} \cdot A^{(1S)} = \begin{pmatrix} -1 & 1 & 0 & 0 & 0 & m_2^2 - p^2 \\ -1 & 0 & 1 & 0 & 0 & m_1^2 - p^2 \\ 1 & 0 & 0 & 1 & 1 & p^2 \end{pmatrix}. \tag{150}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & 0 & 0 & 0 & \alpha_{1,6} \\ \alpha_{2,1} & 0 & 2 - D & 0 & 0 & \alpha_{2,6} \\ \alpha_{3,1} & 0 & 0 & -1 & \frac{D}{2} - 2 & \alpha_{3,6} \end{pmatrix}, \tag{151}$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,1} + \alpha_{1,6} &= -1, \quad \alpha_{2,1} + \alpha_{2,6} = D - 3, \quad \alpha_{3,1} + \alpha_{3,6} = 2 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{2,1} + \alpha_{3,1} &= 0, \quad \alpha_{1,6} + \alpha_{2,6} + \alpha_{3,6} = \frac{D}{2} - 2.\end{aligned}\quad (152)$$

Correspondingly there are 12 choices on the matrix of integer lattice whose submatrix composed of the first and sixth columns is formulated as $\pm n_1 E_3^{(i)} \pm n_2 E_3^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 3), (2, 3)\}$, and other elements are all zero. The exponents are obtained from those corresponding to $\psi_{\{1,3,4\}}^{(i)}$ through the exchanging $\alpha_{i,2} \rightarrow \alpha_{i,1}$, $i = 1, 2, 3$. Certainly the geometric representations of those exponent matrices are obtained from that of the hypergeometric functions by the permutation $(\widehat{12})$. The hypergeometric functions are accordingly written as

$$\psi_{\{2,3,4\}}^{(i)}(p^2, m_1^2, m_2^2) = \psi_{\{1,3,4\}}^{(i)}(p^2, m_2^2, m_1^2), \quad (i = 1, \dots, 12). \quad (153)$$

Since $\det(A_{\{1,4,6\}}^{(1S)}) = m_1^2 - m_2^2$, we choose the first, fourth, and sixth column vectors composing of the basis of the projective plane, and find

$$\left(A_{\{1,4,6\}}^{(1S)}\right)^{-1} \cdot A^{(1S)} = \begin{pmatrix} 1 & -\frac{m_1^2 - p^2}{m_1^2 - m_2^2} & -\frac{p^2 - m_2^2}{m_1^2 - m_2^2} & 0 & 0 & 0 \\ 0 & \frac{m_1^2}{m_1^2 - m_2^2} & -\frac{m_2^2}{m_1^2 - m_2^2} & 1 & 1 & 0 \\ 0 & -\frac{1}{m_1^2 - m_2^2} & \frac{1}{m_1^2 - m_2^2} & 0 & 0 & 1 \end{pmatrix}. \quad (154)$$

Accordingly the matrix of exponents is

$$\begin{pmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & 0 & 0 & 0 \\ 0 & \alpha_{2,2} & \alpha_{2,3} & -1 & \frac{D}{2} - 2 & 0 \\ 0 & \alpha_{3,2} & \alpha_{3,3} & 0 & 0 & \frac{D}{2} - 2 \end{pmatrix}, \quad (155)$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,2} + \alpha_{1,3} &= -1, \quad \alpha_{2,2} + \alpha_{2,3} = 2 - \frac{D}{2}, \quad \alpha_{3,2} + \alpha_{3,3} = 1 - \frac{D}{2}, \\ \alpha_{1,2} + \alpha_{2,2} + \alpha_{3,2} &= 0, \quad \alpha_{1,3} + \alpha_{2,3} + \alpha_{3,3} = 2 - D.\end{aligned}\quad (156)$$

There are 12 choices on the matrix of integer lattice whose submatrix composed of the second and third columns is formulated as $\pm n_1 E_3^{(i)} \pm n_2 E_3^{(j)}$, where $n_{1,2} \geq 0$, and $(i, j) =$

$\{(1, 2), (1, 3), (2, 3)\}$, and other elements are all zero. Corresponding to the integer lattice $n_1 E_3^{(3)} + n_2 E_3^{(2)}$, the choice of exponents is written as

$$\alpha_{1,2} = D - 3, \alpha_{1,3} = 2 - D, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = 1 - \frac{D}{2}, \alpha_{3,3} = 0, \quad (157)$$

whose geometric representation is presented in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 3), (4, 5)\}$, $\{e, f\} = \{2, 6\}$. In a similar way the exponents of the integer lattice $n_1 E_3^{(3)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = \frac{D}{2} - 1, \alpha_{2,3} = 3 - D, \alpha_{3,2} = 1 - \frac{D}{2}, \alpha_{3,3} = 0, \quad (158)$$

whose geometric representation is shown in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 3), (2, 6)\}$. The exponents of the integer lattice $n_1 E_3^{(2)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = 1 - \frac{D}{2}, \quad (159)$$

whose geometric representation is plotted in Fig.4(1) with $\{(a, b), (c, d)\} = \{(2, 6), (4, 5)\}$, $\{e, f\} = \{1, 3\}$. Three linear independent hypergeometric functions are

$$\begin{aligned} \psi_{\{1,4,6\}}^{(1)}(p^2, m_1^2, m_2^2) &\sim \frac{[(m_1^2 - p^2)(m_1^2 - m_2^2)]^{D-3}}{(p^2 - m_2^2)^{D-2}(m_1^2)^{D/2-2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2!} \\ &\quad \times \Gamma\left(\frac{D}{2} - 1 + n_2\right) \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_1} \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_2} \\ &\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\ \psi_{\{1,4,6\}}^{(2)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{D/2-1}}{(m_2^2)^{D-3} (p^2 - m_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1 - n_2)}{\Gamma(4 - D + n_1 - n_2) n_2!} \\ &\quad \times \Gamma\left(\frac{D}{2} - 1 + n_2\right) \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_1} \left(\frac{m_1^2}{m_2^2}\right)^{n_2} \\ &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{D/2-1}}{(m_2^2)^{D-3} (p^2 - m_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1)}{\Gamma(4 - D + n_1) n_2!} \\ &\quad \times \Gamma\left(\frac{D}{2} - 1 + n_2\right) \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_1} \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_2} \\ &\sim \frac{[(m_2^2 - m_1^2)(p^2 - m_2^2)]^{D/2-2} (m_1^2)^{D/2-1}}{(m_2^2)^{D-3}} \\ &\quad \times \sum_{n_1=0}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1)}{\Gamma(4 - D + n_1)} \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_1}, \\ \psi_{\{1,4,6\}}^{(3)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(\frac{D}{2} - 2 + n_2)}{n_2! \Gamma(1 - n_1 - n_2)} \end{aligned}$$

$$\begin{aligned}
& \times \frac{1}{\Gamma(2 - \frac{D}{2} + n_1 + n_2)} \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_1} \left(\frac{m_1^2}{m_2^2}\right)^{n_2} \\
& \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2}, \tag{160}
\end{aligned}$$

whose intersection is a nonempty proper subset of the whole parameter space. In other words, those hypergeometric functions constitute a fundamental solution system of the GKZ-system of PDEs in a proper subset of the whole parameter space.

Corresponding to the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(2)}$, the matrix of exponents is

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = 1 - \frac{D}{2}, \tag{161}$$

whose geometric representation is presented in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 2), (4, 5)\}$, $\{e, f\} = \{3, 6\}$. Similarly the exponents of the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 1, \alpha_{2,3} = 1 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = 1 - \frac{D}{2}, \tag{162}$$

whose geometric representation is drawn in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 2), (3, 6)\}$. Meanwhile the exponents of the integer lattice $-n_1 E_3^{(2)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = \frac{D}{2} - 1, \alpha_{3,3} = 2 - D, \tag{163}$$

whose geometric representation is presented in Fig.4(1) with $\{(a, b), (c, d)\} = \{(3, 6), (4, 5)\}$, $\{e, f\} = \{1, 2\}$. The corresponding hypergeometric functions are

$$\begin{aligned}
\psi_{\{1,4,6\}}^{(4)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2!} \\
& \times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{m_1^2 (p^2 - m_2^2)}{m_2^2 (p^2 - m_1^2)}\right)^{n_1} \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_2} \\
& \sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,4,6\}}^{(5)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{3-D/2}}{m_1^2 (p^2 - m_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + 1 + n_1 + n_2)}{\Gamma(3 + n_1 + n_2)} \\
& \times \frac{\Gamma(1 + n_2)}{\Gamma(4 - \frac{D}{2} + n_2)} \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_1^2}\right)^{n_2}, \\
\psi_{\{1,4,6\}}^{(6)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D/2-2} (m_2^2)^{D/2-1}}{(p^2 - m_1^2) (m_1^2)^{D/2-2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1 + n_2)}{\Gamma(\frac{D}{2} + n_1 + n_2) n_2!} \\
& \times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_1^2}\right)^{n_2}
\end{aligned}$$

$$\begin{aligned}
& \sim \frac{(m_2^2 - m_1^2)^{D/2-2} (m_2^2)^{D/2-1}}{(p^2 - m_1^2) (m_1^2)^{D/2-2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1)}{\Gamma(\frac{D}{2} + n_1) n_2!} \\
& \quad \times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_2} \\
& \sim \frac{(m_2^2 - p^2)^{D/2-2} (m_2^2)^{D/2-1}}{(p^2 - m_1^2) (p^2)^{D/2-2}} \\
& \quad \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1)}{\Gamma(\frac{D}{2} + n_1)} \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_1}, \tag{164}
\end{aligned}$$

whose intersection of convergent regions is a nonempty proper subset of the whole parameter space. Correspondingly the hypergeometric functions with the exponents of Eq.(162) is written as

$$\begin{aligned}
\psi_{\{1,4,6\}}^{(5)'}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - p^2)^{D-3} p^2}{(m_1^2 - p^2) (m_2^2)^{D/2-1}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n_1 - n_2)}{\Gamma(2 + n_1 - n_2) n_2!} \\
& \quad \times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{p^2(m_1^2 - m_2^2)}{m_2^2(m_1^2 - p^2)}\right)^{n_1} \left(\frac{m_2^2}{p^2}\right)^{n_2} \\
& \sim \psi_{\{1,4,6\}}^{(4)}(p^2, m_1^2, m_2^2) \\
& \quad + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(\frac{D}{2} - 3)} \psi_{\{1,4,6\}}^{(5)}(p^2, m_1^2, m_2^2). \tag{165}
\end{aligned}$$

As the integer lattice is chosen as $n_1 E_3^{(3)} - n_2 E_3^{(2)}$, the exponents are written as

$$\alpha_{1,2} = \frac{D}{2} - 2, \quad \alpha_{1,3} = 1 - \frac{D}{2}, \quad \alpha_{2,2} = 2 - \frac{D}{2}, \quad \alpha_{2,3} = 0, \quad \alpha_{3,2} = 0, \quad \alpha_{3,3} = 1 - \frac{D}{2} \tag{166}$$

whose geometric representation is plotted in Fig.4(2) with $(a, b) = (4, 5)$, $c = 1$, $d = 2$, $\{e, f\} = \{3, 6\}$. The exponents of the integer lattice $n_1 E_3^{(3)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = 0, \quad \alpha_{1,3} = -1, \quad \alpha_{2,2} = 0, \quad \alpha_{2,3} = 2 - \frac{D}{2}, \quad \alpha_{3,2} = 0, \quad \alpha_{3,3} = 1 - \frac{D}{2}, \tag{167}$$

whose geometric representation is shown in Fig.4(4) with $(a, b, c) = (2, 4, 5)$, $(d, e, f) = (1, 3, 6)$. The exponents of the integer lattice $n_1 E_3^{(2)} - n_2 E_3^{(1)}$ are given by

$$\alpha_{1,2} = 0, \quad \alpha_{1,3} = -1, \quad \alpha_{2,2} = 2 - \frac{D}{2}, \quad \alpha_{2,3} = 0, \quad \alpha_{3,2} = \frac{D}{2} - 2, \quad \alpha_{3,3} = 3 - D, \tag{168}$$

where the geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 6$, $d = 2$, $(e, f) = (1, 3)$. Correspondingly, the hypergeometric functions are

$$\psi_{\{1,4,6\}}^{(7)}(p^2, m_1^2, m_2^2) \sim \frac{(p^2 - m_1^2)^{D/2-2} (m_2^2 - m_1^2)^{D-3}}{(p^2 - m_2^2)^{D/2-1} (m_1^2)^{D/2-2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2!}$$

$$\begin{aligned}
& \times \Gamma\left(\frac{D}{2} - 1 + n_2\right) \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_1} \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_2} \\
& \sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{1,4,6\}}^{(8)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma\left(\frac{D}{2} - 1 + n_2\right)}{n_2! \Gamma(1 - n_1 - n_2)} \\
& \times \frac{1}{\Gamma\left(3 - \frac{D}{2} + n_1 + n_2\right)} \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_1} \left(\frac{m_2^2}{m_1^2}\right)^{n_2} \\
& \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2}, \\
\psi_{\{1,4,6\}}^{(9)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma\left(2 - \frac{D}{2} + n_1 - n_2\right)}{\Gamma(4 - D + n_1 - n_2) n_2!} \\
& \times \Gamma\left(\frac{D}{2} - 2 + n_2\right) \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_1} \left(\frac{m_2^2}{m_1^2}\right)^{n_2} \\
& \sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma\left(2 - \frac{D}{2} + n_1\right)}{\Gamma(4 - D + n_1) n_2!} \\
& \times \Gamma\left(\frac{D}{2} - 2 + n_2\right) \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_1} \left(\frac{m_2^2(p^2 - m_1^2)}{m_1^2(p^2 - m_2^2)}\right)^{n_2} \\
& \sim \frac{(m_2^2 - m_1^2)^{D/2-1} (m_2^2 - p^2)^{D/2-3}}{(p^2)^{D/2-2}} \\
& \times \sum_{n_1=0}^{\infty} \frac{\Gamma\left(2 - \frac{D}{2} + n_1\right)}{\Gamma(4 - D + n_1)} \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_1}. \tag{169}
\end{aligned}$$

Although the intersection of the convergent regions of these series is the empty subset, the sets of the hypergeometric functions $\{\psi_{\{1,4,6\}}^{(1)}, \psi_{\{1,4,6\}}^{(8)}, \psi_{\{1,4,6\}}^{(9)}\}$ and $\{\psi_{\{1,4,6\}}^{(6)}, \psi_{\{1,4,6\}}^{(7)}, \psi_{\{1,4,6\}}^{(8)}\}$ constitute the fundamental solution systems of the corresponding nonempty proper subsets of the whole parameter space, respectively.

The matrix of exponents of the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(2)}$ is

$$\alpha_{1,2} = \frac{D}{2} - 1, \quad \alpha_{1,3} = -\frac{D}{2}, \quad \alpha_{2,2} = 0, \quad \alpha_{2,3} = 2 - \frac{D}{2}, \quad \alpha_{3,2} = 1 - \frac{D}{2}, \quad \alpha_{3,3} = 0, \tag{170}$$

whose geometric representation is drawn in Fig.4(2) with $(a, b) = (4, 5)$, $c = 1$, $d = 3$, $\{e, f\} = \{2, 6\}$. The exponents of the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,2} = -1, \quad \alpha_{1,3} = 0, \quad \alpha_{2,2} = \frac{D}{2}, \quad \alpha_{2,3} = 2 - D, \quad \alpha_{3,2} = 1 - \frac{D}{2}, \quad \alpha_{3,3} = 0, \tag{171}$$

whose geometric representation is presented in Fig.4(4) with $(a, b, c) = (3, 4, 5)$, $(d, e, f) =$

(1, 2, 6). Finally the choice on exponents of integer lattice $-n_1 E_3^{(2)} + n_2 E_3^{(1)}$ is

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 1, \alpha_{3,3} = -\frac{D}{2}, \quad (172)$$

whose geometric representation is plotted in Fig.4(2) with $(a, b) = (4, 5)$, $c = 6$, $d = 3$, $(e, f) = (1, 2)$. Three corresponding power series are presented as

$$\begin{aligned} \psi_{\{1,4,6\}}^{(10)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D/2-1} (m_2^2 - m_1^2)^{D-3}}{(m_2^2)^{D/2-2} (p^2 - m_2^2)^{D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2!} \\ &\quad \times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{m_1^2 (p^2 - m_2^2)}{m_2^2 (p^2 - m_1^2)}\right)^{n_1} \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_2} \\ &\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\ \psi_{\{1,4,6\}}^{(11)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{D/2}}{(p^2 - m_1^2) (m_2^2)^{D-2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1 + n_2)}{\Gamma(1 + \frac{D}{2} + n_1 + n_2)} \\ &\quad \times \frac{\Gamma(\frac{D}{2} - 1 + n_2)}{n_2!} \left(\frac{m_1^2 (p^2 - m_2^2)}{m_2^2 (p^2 - m_1^2)}\right)^{n_1} \left(\frac{m_1^2}{m_2^2}\right)^{n_2} \\ &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_1^2)^{D/2}}{(p^2 - m_1^2) (m_2^2)^{D-2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1)}{\Gamma(1 + \frac{D}{2} + n_1)} \\ &\quad \times \frac{\Gamma(\frac{D}{2} - 1 + n_2)}{n_2!} \left(\frac{m_1^2 (p^2 - m_2^2)}{m_2^2 (p^2 - m_1^2)}\right)^{n_1} \left(\frac{p^2 - m_1^2}{p^2 - m_2^2}\right)^{n_2} \\ &\sim \frac{(m_2^2 - m_1^2)^{D/2-2} (p^2 - m_2^2)^{D/2-1} (m_1^2)^{D/2}}{(p^2 - m_1^2) (m_2^2)^{D-2}} \\ &\quad \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1)}{\Gamma(1 + \frac{D}{2} + n_1)} \left(\frac{m_1^2 (p^2 - m_2^2)}{m_2^2 (p^2 - m_1^2)}\right)^{n_1}, \\ \psi_{\{1,4,6\}}^{(12)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_1^2)^2 (m_2^2 - m_1^2)^{D-3}}{(m_2^2)^{2-D/2} (p^2 - m_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + n_1 + n_2) \Gamma(1 + n_2)}{\Gamma(3 + n_1 + n_2) \Gamma(-1 - \frac{D}{2} + n_2)} \\ &\quad \times \left(\frac{m_1^2 (p^2 - m_2^2)}{m_2^2 (p^2 - m_1^2)}\right)^{n_1} \left(\frac{m_1^2}{m_2^2}\right)^{n_2}. \end{aligned} \quad (173)$$

Certainly intersection of convergent regions of these series is the empty set, and the hypergeometric function with the exponents of Eq.(172) is written as

$$\begin{aligned} \psi_{\{1,4,6\}}^{(12)'}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{p^2 - m_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + n_1 - n_2)}{\Gamma(2 + n_1 - n_2) n_2!} \\ &\quad \times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{p^2 - m_2^2}{p^2 - m_1^2}\right)^{n_1} \left(\frac{m_1^2}{m_2^2}\right)^{n_2} \\ &\sim \psi_{\{1,4,6\}}^{(10)}(p^2, m_1^2, m_2^2) \\ &\quad + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(\frac{D}{2} + 2)} \psi_{\{1,4,6\}}^{(12)}(p^2, m_1^2, m_2^2). \end{aligned} \quad (174)$$

The sets of the hypergeometric functions $\{\psi_{\{1,4,6\}}^{(3)}, \psi_{\{1,4,6\}}^{(10)}, \psi_{\{1,4,6\}}^{(11)}\}$ and $\{\psi_{\{1,4,6\}}^{(4)}, \psi_{\{1,4,6\}}^{(11)}, \psi_{\{1,4,6\}}^{(12)}\}$ evidently constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{\{2,4,6\}}^{(1S)}) = -p^2 + m_1^2$, and

$$\left(A_{\{2,4,6\}}^{(1S)}\right)^{-1} \cdot A^{(1S)} = \begin{pmatrix} -\frac{m_1^2 - m_2^2}{m_1^2 - p^2} & 1 & -\frac{m_2^2 - p^2}{m_1^2 - p^2} & 0 & 0 & 0 \\ \frac{m_1^2}{m_1^2 - p^2} & 0 & -\frac{p^2}{m_1^2 - p^2} & 1 & 1 & 0 \\ -\frac{1}{m_1^2 - p^2} & 0 & \frac{1}{m_1^2 - p^2} & 0 & 0 & 1 \end{pmatrix}, \quad (175)$$

the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} & 0 & 0 & 0 \\ \alpha_{2,1} & 0 & \alpha_{2,3} & -1 & \frac{D}{2} - 2 & 0 \\ \alpha_{3,1} & 0 & \alpha_{3,3} & 0 & 0 & \frac{D}{2} - 2 \end{pmatrix}, \quad (176)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,3} &= -1, \quad \alpha_{2,1} + \alpha_{2,3} = 2 - \frac{D}{2}, \quad \alpha_{3,1} + \alpha_{3,3} = 1 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{2,1} + \alpha_{3,1} &= 0, \quad \alpha_{1,3} + \alpha_{2,3} + \alpha_{3,3} = 2 - D. \end{aligned} \quad (177)$$

There are 12 choices on the matrix of integer lattice whose submatrix composed of the first and third columns is formulated as $\pm n_1 E_3^{(i)} \pm n_2 E_3^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 3), (2, 3)\}$, and other elements are all zero. The exponents are obtained from those corresponding to $\psi_{\{1,4,6\}}^{(i)}$ through the exchanging $\alpha_{i,2} \rightarrow \alpha_{i,1}$, $i = 1, 2, 3$. Certainly the geometric representations of those exponent matrices are obtained from that of the hypergeometric functions by the permutation $(\widehat{12})$. The hypergeometric functions are

$$\psi_{\{2,4,6\}}^{(i)}(p^2, m_1^2, m_2^2) = \psi_{\{1,4,6\}}^{(i)}(p^2, m_2^2, m_1^2), \quad (i = 1, \dots, 12). \quad (178)$$

Because $\det(A_{\{3,4,6\}}^{(1S)}) = m_2^2 - m_1^2$, we choose the third, fourth, and sixth columns as the basis of the projective plane, and get

$$\left(A_{\{3,4,6\}}^{(1S)}\right)^{-1} \cdot A^{(1S)} = \begin{pmatrix} \frac{m_1^2 - m_2^2}{m_2^2 - p^2} & -\frac{m_1^2 - p^2}{m_2^2 - p^2} & 1 & 0 & 0 & 0 \\ \frac{m_2^2}{m_2^2 - p^2} & -\frac{p^2}{m_2^2 - p^2} & 0 & 1 & 1 & 0 \\ -\frac{1}{m_2^2 - p^2} & \frac{1}{m_2^2 - p^2} & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (179)$$

Certainly the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & 2-D & 0 & 0 & 0 \\ \alpha_{2,1} & \alpha_{2,2} & 0 & -1 & \frac{D}{2}-2 & 0 \\ \alpha_{3,1} & \alpha_{3,2} & 0 & 0 & 0 & \frac{D}{2}-2 \end{pmatrix}, \quad (180)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,2} &= D - 3, \quad \alpha_{2,1} + \alpha_{2,2} = 2 - \frac{D}{2}, \quad \alpha_{3,1} + \alpha_{3,2} = 1 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{2,1} + \alpha_{3,1} &= 0, \quad \alpha_{1,2} + \alpha_{2,2} + \alpha_{3,2} = 0. \end{aligned} \quad (181)$$

There are 12 choices on the matrix of integer lattice whose submatrix composed of the first and second columns is formulated as $\pm n_1 E_3^{(i)} \pm n_2 E_3^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 3), (2, 3)\}$, and other elements are all zero. As the integer lattice is chosen as $n_1 E_3^{(3)} + n_2 E_3^{(2)}$, the matrix of exponents is written as

$$\alpha_{1,1} = D - 3, \quad \alpha_{1,2} = 0, \quad \alpha_{2,1} = 2 - \frac{D}{2}, \quad \alpha_{2,2} = 0, \quad \alpha_{3,1} = 1 - \frac{D}{2}, \quad \alpha_{3,2} = 0, \quad (182)$$

whose geometric representation is plotted in Fig.4(1) with $\{(a, b), (c, d)\} = \{(2, 3), (4, 5)\}$, $\{e, f\} = \{1, 6\}$. In a similar way the exponents of the integer lattice $n_1 E_3^{(3)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,1} = 0, \quad \alpha_{1,2} = D - 3, \quad \alpha_{2,1} = \frac{D}{2} - 1, \quad \alpha_{2,2} = 3 - D, \quad \alpha_{3,1} = 1 - \frac{D}{2}, \quad \alpha_{3,2} = 0, \quad (183)$$

where the geometric representation is presented in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 6), (2, 3)\}$. The exponents of the integer lattice $n_1 E_3^{(2)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,1} = 0, \quad \alpha_{1,2} = D - 3, \quad \alpha_{2,1} = 0, \quad \alpha_{2,2} = 2 - \frac{D}{2}, \quad \alpha_{3,1} = 0, \quad \alpha_{3,2} = 1 - \frac{D}{2}, \quad (184)$$

whose geometric representation is drawn in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 6), (4, 5)\}$, $\{e, f\} = \{2, 3\}$. Using the integer lattices and corresponding exponents, the hypergeometric functions are

$$\begin{aligned} \psi_{\{3,4,6\}}^{(1)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \sum_{n_1, n_2} \frac{(-)^{n_1+n_2} \Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2! \Gamma(1 - n_1 - n_2)} \\ &\quad \times \frac{\Gamma(\frac{D}{2} - 1 + n_2)}{\Gamma(D - 2 + n_1 + n_2)} \left(\frac{p^2(m_2^2 - m_1^2)}{m_2^2(p^2 - m_1^2)}\right)^{n_1} \left(\frac{m_2^2 - m_1^2}{p^2 - m_1^2}\right)^{n_2} \\ &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2}, \end{aligned}$$

$$\begin{aligned}
\psi_{\{3,4,6\}}^{(2)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D-3} (m_2^2)^{D/2-1}}{(m_2^2 - p^2)(p^2)^{D-3}} \\
&\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1 - n_2) \Gamma(3 - D + n_1)}{n_1! n_2! \Gamma(4 - D + n_1 - n_2)} \\
&\times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{p^2(m_2^2 - m_1^2)}{m_2^2(p^2 - m_1^2)}\right)^{n_1} \left(\frac{m_2^2}{p^2}\right)^{n_2}, \\
\psi_{\{3,4,6\}}^{(3)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2} \\
&\times \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(3 - D + n_1) \Gamma(\frac{D}{2} - 2 + n_2)}{n_1! n_2! \Gamma(1 - n_1 - n_2)} \\
&\times \frac{1}{\Gamma(2 - \frac{D}{2} + n_1 + n_2)} \left(\frac{m_2^2 - m_1^2}{p^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{p^2}\right)^{n_2} \\
&\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \tag{185}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. In other words, those hypergeometric functions constitute a fundamental solution system of the GKZ-system in a proper subset of the whole parameter space.

Corresponding to the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(2)}$, the matrix of exponents is

$$\alpha_{1,1} = 0, \alpha_{1,2} = D - 3, \alpha_{2,1} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{3,1} = 0, \alpha_{3,2} = 1 - \frac{D}{2}, \tag{186}$$

whose geometric representation is shown in Fig.4(1) with $\{(a, b), (c, d)\} = \{(1, 3), (4, 5)\}$, $\{e, f\} = \{2, 6\}$. Similarly the exponents of the integer lattice $-n_1 E_3^{(3)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,1} = D - 3, \alpha_{1,2} = 0, \alpha_{2,1} = 3 - D, \alpha_{2,2} = \frac{D}{2} - 1, \alpha_{3,1} = 0, \alpha_{3,2} = 1 - \frac{D}{2}, \tag{187}$$

whose geometric representation is presented in Fig.4(3) with $(a, b) = (4, 5)$, $\{(c, d), (e, f)\} = \{(1, 3), (2, 6)\}$. The exponents of the integer lattice $-n_1 E_3^{(2)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,1} = D - 3, \alpha_{1,2} = 0, \alpha_{2,1} = 2 - \frac{D}{2}, \alpha_{2,2} = 0, \alpha_{3,1} = 1 - \frac{D}{2}, \alpha_{3,2} = 0, \tag{188}$$

whose geometric representation is drawn in Fig.4(1) with $\{(a, b), (c, d)\} = \{(2, 6), (4, 5)\}$, $\{e, f\} = \{1, 3\}$. Accordingly the hypergeometric functions are written as

$$\begin{aligned}
\psi_{\{3,4,6\}}^{(4)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2} \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2! \Gamma(1 - n_1 - n_2)} \\
&\times \frac{\Gamma(\frac{D}{2} - 1 + n_2)}{\Gamma(D - 2 + n_1 + n_2)} \left(\frac{m_2^2(p^2 - m_1^2)}{p^2(m_2^2 - m_1^2)}\right)^{n_1} \left(\frac{p^2 - m_1^2}{m_2^2 - m_1^2}\right)^{n_2}
\end{aligned}$$

$$\begin{aligned}
& \sim \frac{(p^2 - m_1^2)^{D-3}(p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{3,4,6\}}^{(5)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D-3}(p^2)^{D/2-1}}{(m_2^2)^{D-3}(m_2^2 - p^2)} \\
& \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1 - n_2)\Gamma(3 - D + n_1)}{n_1!n_2!\Gamma(4 - D + n_1 - n_2)} \\
& \times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{m_2^2(p^2 - m_1^2)}{p^2(m_2^2 - m_1^2)}\right)^{n_1} \left(\frac{m_2^2}{p^2}\right)^{n_2}, \\
\psi_{\{3,4,6\}}^{(6)}(p^2, m_1^2, m_2^2) & \sim \frac{(m_2^2 - m_1^2)^{D-3}(m_2^2)^{2-D/2}}{m_2^2 - p^2} \\
& \times \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2}\Gamma(3 - D + n_1)\Gamma(\frac{D}{2} - 2 + n_2)}{n_1!n_2!\Gamma(1 - n_1 - n_2)} \\
& \times \frac{1}{\Gamma(2 - \frac{D}{2} + n_1 + n_2)} \left(\frac{p^2 - m_1^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{p^2}{m_2^2}\right)^{n_2} \\
& \sim \frac{(m_2^2 - m_1^2)^{D-3}(m_2^2)^{2-D/2}}{m_2^2 - p^2},
\end{aligned} \tag{189}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The fact indicates that those hypergeometric functions constitute a fundamental solution system of the GKZ-system in a proper subset of the whole parameter space.

As the integer lattice is chosen as $n_1 E_3^{(3)} - n_2 E_3^{(2)}$ the exponents are

$$\alpha_{1,1} = \frac{D}{2} - 2, \alpha_{1,2} = \frac{D}{2} - 1, \alpha_{2,1} = 2 - \frac{D}{2}, \alpha_{2,2} = 0, \alpha_{3,1} = 0, \alpha_{3,2} = 1 - \frac{D}{2} \tag{190}$$

whose geometric representation is plotted in Fig.4(2) with $(a, b) = (4, 5)$, $c = 3$, $d = 1$, $\{e, f\} = \{2, 6\}$. Corresponding to the integer lattice $n_1 E_3^{(3)} - n_2 E_3^{(1)}$, the exponents are

$$\alpha_{1,1} = 0, \alpha_{1,2} = D - 3, \alpha_{2,1} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{3,1} = 0, \alpha_{3,2} = 1 - \frac{D}{2}, \tag{191}$$

whose geometric representation is presented in Fig.4(4) with $(a, b, c) = (1, 4, 5)$, $(d, e, f) = (2, 3, 6)$. Similarly the exponents of the integer lattice $n_1 E_3^{(2)} - n_2 E_3^{(1)}$ are

$$\alpha_{1,1} = 0, \alpha_{1,2} = D - 3, \alpha_{2,1} = 2 - \frac{D}{2}, \alpha_{2,2} = 0, \alpha_{3,1} = \frac{D}{2} - 2, \alpha_{3,2} = 3 - D, \tag{192}$$

whose geometric representation is presented in Fig.4(2) with $(a, b) = (4, 5)$, $c = 6$, $d = 1$, $(e, f) = (2, 3)$. Applying those matrices of exponents and integer lattices, we obtain three

linear independent hypergeometric functions

$$\begin{aligned}
\psi_{\{3,4,6\}}^{(7)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D/2-1} (m_2^2 - m_1^2)^{D/2-2}}{(m_2^2 - p^2)(m_2^2)^{D/2-2}} \\
&\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1 - n_2) \Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_1 - n_2)} \\
&\times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{p^2(m_2^2 - m_1^2)}{m_2^2(p^2 - m_1^2)}\right)^{n_1} \left(\frac{p^2 - m_1^2}{m_2^2 - m_1^2}\right)^{n_2}, \\
\psi_{\{3,4,6\}}^{(8)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2} \\
&\times \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(3 - D + n_1) \Gamma(\frac{D}{2} - 1 + n_2)}{n_1! n_2! \Gamma(1 - n_1 - n_2) \Gamma(3 - \frac{D}{2} + n_1 + n_2)} \\
&\times \left(\frac{p^2(m_2^2 - m_1^2)}{m_2^2(p^2 - m_1^2)}\right)^{n_1} \left(\frac{p^2}{m_2^2}\right)^{n_2} \\
&\sim \frac{(p^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{p^2 - m_2^2}, \\
\psi_{\{3,4,6\}}^{(9)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \\
&\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 - n_2) \Gamma(3 - D + n_1)}{n_1! n_2! \Gamma(4 - D + n_1 - n_2)} \\
&\times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{m_2^2 - m_1^2}{p^2 - m_1^2}\right)^{n_1} \left(\frac{p^2}{m_2^2}\right)^{n_2}. \tag{193}
\end{aligned}$$

Although the intersection of the convergent regions of these series is the empty set, the sets of the hypergeometric functions $\{\psi_{\{3,4,6\}}^{(1)}, \psi_{\{3,4,6\}}^{(8)}, \psi_{\{3,4,6\}}^{(9)}\}$ and $\{\psi_{\{3,4,6\}}^{(6)}, \psi_{\{3,4,6\}}^{(7)}, \psi_{\{3,4,6\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

As the integer lattice is chosen as $-n_1 E_3^{(3)} + n_2 E_3^{(2)}$, the matrix of exponents is

$$\alpha_{1,1} = \frac{D}{2} - 1, \quad \alpha_{1,2} = \frac{D}{2} - 2, \quad \alpha_{2,1} = 0, \quad \alpha_{2,2} = 2 - \frac{D}{2}, \quad \alpha_{3,1} = 1 - \frac{D}{2}, \quad \alpha_{3,2} = 0 \tag{194}$$

whose geometric representation is plotted in Fig.4(2) with $(a, b) = (4, 5)$, $c = 3$, $d = 2$, $\{e, f\} = \{1, 6\}$. Corresponding to the integer lattice $-n_1 E_3^{(3)} + n_2 E_3^{(1)}$, the matrix of exponents is

$$\alpha_{1,1} = D - 3, \quad \alpha_{1,2} = 0, \quad \alpha_{2,1} = 2 - \frac{D}{2}, \quad \alpha_{2,2} = 0, \quad \alpha_{3,1} = 1 - \frac{D}{2}, \quad \alpha_{3,2} = 0, \tag{195}$$

whose geometric representation is presented in Fig.4(4) with $(a, b, c) = (2, 4, 5)$, $(d, e, f) =$

(1, 3, 6). The exponents of the integer lattice $-n_1 E_3^{(2)} + n_2 E_3^{(1)}$ are

$$\alpha_{1,1} = D - 3, \alpha_{1,2} = 0, \alpha_{2,1} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{3,1} = 3 - D, \alpha_{3,2} = \frac{D}{2} - 2, \quad (196)$$

whose geometric representation is shown in Fig.4(2) with $(a, b) = (4, 5)$, $c = 6$, $d = 2$, $(e, f) = (1, 3)$. Accordingly the hypergeometric series are

$$\begin{aligned} \psi_{\{3,4,6\}}^{(10)}(p^2, m_1^2, m_2^2) &\sim \frac{(p^2 - m_1^2)^{D/2-2} (m_2^2 - m_1^2)^{D/2-1}}{(m_2^2 - p^2)(p^2)^{D/2-2}} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1 - n_2) \Gamma(\frac{D}{2} - 2 + n_1)}{n_1! n_2! \Gamma(\frac{D}{2} - 1 + n_1 - n_2)} \\ &\times \Gamma(\frac{D}{2} - 1 + n_2) \left(\frac{m_2^2(p^2 - m_1^2)}{p^2(m_2^2 - m_1^2)}\right)^{n_1} \left(\frac{m_2^2 - m_1^2}{p^2 - m_1^2}\right)^{n_2}, \\ \psi_{\{3,4,6\}}^{(11)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{(-)^{n_1+n_2} \Gamma(3 - D + n_1) \Gamma(\frac{D}{2} - 1 + n_2)}{n_1! n_2! \Gamma(1 - n_1 - n_2) \Gamma(3 - \frac{D}{2} + n_1 + n_2)} \\ &\times \left(\frac{m_2^2(p^2 - m_1^2)}{p^2(m_2^2 - m_1^2)}\right)^{n_1} \left(\frac{m_2^2}{p^2}\right)^{n_2} \\ &\sim \frac{(m_2^2 - m_1^2)^{D-3} (m_2^2)^{2-D/2}}{m_2^2 - p^2}, \\ \psi_{\{3,4,6\}}^{(12)}(p^2, m_1^2, m_2^2) &\sim \frac{(m_2^2 - m_1^2)^{D-3} (p^2)^{2-D/2}}{m_2^2 - p^2} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 - n_2) \Gamma(3 - D + n_1)}{n_1! n_2! \Gamma(4 - D + n_1 - n_2)} \\ &\times \Gamma(\frac{D}{2} - 2 + n_2) \left(\frac{p^2 - m_1^2}{m_2^2 - m_1^2}\right)^{n_1} \left(\frac{m_2^2}{p^2}\right)^{n_2}. \end{aligned} \quad (197)$$

Although the intersection of the convergent regions of these series is the empty set, the sets of the hypergeometric functions $\{\psi_{\{3,4,6\}}^{(3)}, \psi_{\{3,4,6\}}^{(10)}, \psi_{\{3,4,6\}}^{(11)}\}$ and $\{\psi_{\{3,4,6\}}^{(4)}, \psi_{\{3,4,6\}}^{(11)}, \psi_{\{3,4,6\}}^{(12)}\}$ also constitute the fundamental solution systems of the corresponding nonempty proper subsets of the whole parameter space, respectively.

In principle, we can embed the Feynman parametric representation of the 1-loop self energy into infinite subvarieties of the Grassmannian $G_{3,6}$. However, the dimension of dual space of the GKZ-systems of most subvarieties is much larger than the number of independent dimensionless ratios between the square of outgoing momentum and the square of virtual masses. Correspondingly convergent regions of the obtained hypergeometric series solutions are empty.

III. THE 1-LOOP TRIANGLE WITH ZERO VIRTUAL MASSES

The Feynman integral of the 1-loop triangle diagram with zero virtual masses is represented as the linear combination of the generalized hypergeometric functions F_4 in literature [16] through α -parametric representation for the denominators. Here, we construct the equivalent fundamental solution system through embedding the Feynman integral on Grassmannians.

Denoting three incoming momenta as p_i ($i = 1, 2, 3$), we obtain the Feynman parametric representation of the diagram as

$$\begin{aligned}
C(p_1^2, p_2^2, p_3^2) &= \left(\Lambda_{\text{RE}}^2\right)^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2(q-p_1)^2(q+p_2)^2} \\
&= -i \left(\Lambda_{\text{RE}}^2\right)^{4-D} \frac{(-)^{D/2-3} \Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \int_0^1 dt_1 dt_2 dt_3 \\
&\quad \times \frac{\delta(t_1 + t_2 + t_3 - 1)}{(t_1 t_2 p_3^2 + t_2 t_3 p_1^2 + t_1 t_3 p_2^2)^{3-D/2}}, \tag{198}
\end{aligned}$$

where $p_1 + p_2 + p_3 = 0$. As $p_2 \neq 0$, $p_3 \neq 0$, but $p_2^2 = p_3^2 = 0$, the Feynman parametric representation is simplified as

$$\begin{aligned}
C(p_1^2, 0, 0) &= -i \left(\Lambda_{\text{RE}}^2\right)^{4-D} \frac{(-)^{D/2-3} \Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \\
&\quad \times \int_0^1 dt_1 dt_2 t_2^{D/2-3} (p_1^2 - p_1^2 t_1 - p_1^2 t_2)^{D/2-3} \\
&= i \left(\Lambda_{\text{RE}}^2\right)^{4-D} \frac{\Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \\
&\quad \times \int \omega_3(t) t_2^{D/2-3} t_3^{3-D} (p_1^2 t_1 + p_1^2 t_2 + p_1^2 t_3)^{D/2-3}. \tag{199}
\end{aligned}$$

Eq.(199) can be embedded in the subvariety of the Grassmannian $G_{3,4}$ with splitting local coordinates as

$$A^{(1)} = \begin{pmatrix} 1 & 0 & 0 & p_1^2 \\ 0 & 1 & 0 & p_1^2 \\ 0 & 0 & 1 & p_1^2 \end{pmatrix}. \tag{200}$$

Here the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , and the third row corresponds to the integration variable t_3 , respectively. Meanwhile the first column represents the power function $t_1^0 = 1$, the second column represents the power function $t_2^{D/2-3}$, and the third column represents the

power function t_3^{3-D} , the fourth column represents the function $p_1^2(t_1 + t_2 + t_3)$, respectively. Taking Feynman parametric representation of the 1-loop triangle diagram as a function of the splitting local coordinates in Eq.(200),

$$C(p_1^2, 0, 0) = \Psi^{1T}(z_1, \dots, z_4), \quad (201)$$

we obviously have

$$\Psi^{1T}(\chi_1 z_1, \dots, \chi_4 z_4) = \chi_2^{D/2-3} \chi_3^{3-D} \chi_4^{D/2-3} \Psi^{1T}(z_1, \dots, z_4). \quad (202)$$

Where z_i ($i = 1, \dots, 4$) denotes the i -th column vector in the matrix Eq.(200), and χ_i is a nonzero constant, respectively. Eq.(202) implies that the special Feynman integral satisfies following GKZ-system

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,4}\} C(p_1^2, 0, 0) &= -C(p_1^2, 0, 0), \\ \{\vartheta_{2,2} + \vartheta_{2,4}\} C(p_1^2, 0, 0) &= -C(p_1^2, 0, 0), \\ \{\vartheta_{3,3} + \vartheta_{3,4}\} C(p_1^2, 0, 0) &= -C(p_1^2, 0, 0), \\ \vartheta_{1,1} C(p_1^2, 0, 0) &= 0, \\ \vartheta_{2,2} C(p_1^2, 0, 0) &= \left(\frac{D}{2} - 3\right) C(p_1^2, 0, 0), \\ \vartheta_{3,3} C(p_1^2, 0, 0) &= (3 - D) C(p_1^2, 0, 0), \\ \{\vartheta_{1,4} + \vartheta_{2,4} + \vartheta_{3,4}\} C(p_1^2, 0, 0) &= \left(\frac{D}{2} - 3\right) C(p_1^2, 0, 0). \end{aligned} \quad (203)$$

Obviously the dual space of the GKZ-system of Eq.(203) is the null space $\{0\}$, and compatible with the exponent matrix

$$\begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & \frac{D}{2} - 3 & 0 & 2 - \frac{D}{2} \\ 0 & 0 & 3 - D & D - 4 \end{pmatrix}. \quad (204)$$

Using the above matrices of exponent and integer lattice, one obtains the unique solution of the GKZ-system as

$$C(p_1^2, 0, 0) = \frac{C_{1T}}{\Gamma(3 - \frac{D}{2})\Gamma(D - 3)} (p_1^2)^{D/2-3}, \quad (205)$$

where C_{1T} is the integration constant. In the case $p_1 = -p_2 \neq 0$, $p_3 = 0$, the Feynman parametric representation is modified as

$$C(p_1^2, p_1^2, 0) = -i(\Lambda_{\text{RE}}^2)^{4-D} \frac{(-)^{D/2-3} \Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \int_0^1 dt_1 dt_2 (1 - t_1 - t_2)^{D/2-3}$$

$$\begin{aligned}
& \times (t_1 p_1^2 + t_1 p_1^2)^{D/2-3} \\
& = i \left(\Lambda_{\text{RE}}^2 \right)^{4-D} \frac{\Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \int \omega(t) t_3^{3-D} (t_1 + t_2 + t_3)^{D/2-3} \\
& \quad \times (t_1 p_1^2 + t_2 p_1^2)^{D/2-3}, \tag{206}
\end{aligned}$$

which can be embedded in the subvariety of the Grassmannian $G_{3,5}$ with splitting local coordinates as

$$A^{(1)'} = \begin{pmatrix} 1 & 0 & 0 & 1 & p_1^2 \\ 0 & 1 & 0 & 1 & p_1^2 \\ 0 & 0 & 1 & 1 & 0 \end{pmatrix}. \tag{207}$$

In the above splitting coordinates, the Feynman integral of 1-loop massless triangle satisfies the following GKZ-system

$$\begin{aligned}
& \{\vartheta_{1,1} + \vartheta_{1,4} + \vartheta_{1,5}\} C(p_1^2, p_1^2, 0) = -C(p_1^2, p_1^2, 0), \\
& \{\vartheta_{2,2} + \vartheta_{2,4} + \vartheta_{2,5}\} C(p_1^2, p_1^2, 0) = -C(p_1^2, p_1^2, 0), \\
& \{\vartheta_{3,3} + \vartheta_{3,4}\} C(p_1^2, p_1^2, 0) = -C(p_1^2, p_1^2, 0), \\
& \vartheta_{1,1} C(p_1^2, p_1^2, 0) = 0, \\
& \vartheta_{2,2} C(p_1^2, p_1^2, 0) = 0, \\
& \vartheta_{3,3} C(p_1^2, p_1^2, 0) = (3 - D) C(p_1^2, p_1^2, 0), \\
& \{\vartheta_{1,4} + \vartheta_{2,4} + \vartheta_{3,4}\} C(p_1^2, p_1^2, 0) = \left(\frac{D}{2} - 3\right) C(p_1^2, p_1^2, 0), \\
& \{\vartheta_{1,5} + \vartheta_{2,5}\} C(p_1^2, p_1^2, 0) = \left(\frac{D}{2} - 3\right) C(p_1^2, p_1^2, 0). \tag{208}
\end{aligned}$$

Because the submatrix of Eq.(207) composed of the first, fourth, fifth columns and that composed of the second, fourth, fifth columns satisfy

$$\begin{aligned}
\det A_{145}^{(1)'} &= \det \begin{pmatrix} 1 & 1 & p_1^2 \\ 0 & 1 & p_1^2 \\ 0 & 1 & 0 \end{pmatrix} = -p_1^2 \neq 0, \\
\det A_{245}^{(1)'} &= \det \begin{pmatrix} 0 & 1 & p_1^2 \\ 1 & 1 & p_1^2 \\ 0 & 1 & 0 \end{pmatrix} = p_1^2 \neq 0, \tag{209}
\end{aligned}$$

we can choose $\{(1, 0, 0)^T, (1, 1, 1)^T, (p_1^2, p_1^2, 0)^T\}$ or $\{(0, 1, 0)^T, (1, 1, 1)^T, (p_1^2, p_1^2, 0)^T\}$ spanning the projective plane respectively. As we choose the vectors $\{(1, 0, 0)^T, (1, 1, 1)^T, (p_1^2, p_1^2, 0)^T\}$

as the basis of the projective plane, splitting local coordinates are written as

$$\begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & \frac{1}{p_1^2} & -\frac{1}{p_1^2} & 0 & 1 \end{pmatrix}, \quad (210)$$

with the exponent matrix

$$\begin{pmatrix} 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 2 - \frac{D}{2} & \frac{D}{2} - 3 & 0 \\ 0 & 1 & 1 - \frac{D}{2} & 0 & \frac{D}{2} - 3 \end{pmatrix}. \quad (211)$$

Using the above matrix of exponents and the integer lattice $0_{3 \times 5}$, one obtains the unique solution as

$$\begin{aligned} C(p_1^2, p_1^2, 0) &= \frac{C_{1T}^{(145)}}{\det(A_{145}^{(1)'})\Gamma(2)\Gamma(2 - \frac{D}{2})} (p_1^2)^{D/2-2} \\ &= -\frac{C_{1T}^{(145)}}{\Gamma(2 - \frac{D}{2})} (p_1^2)^{D/2-3}. \end{aligned} \quad (212)$$

Similarly taking the vectors $\{(0, 1, 0)^T, (1, 1, 1)^T, (p_1^2, p_1^2, 0)^T\}$ as the basis of the projective plane, we write the splitting local coordinates as

$$\begin{pmatrix} -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ \frac{1}{p_1^2} & 0 & -\frac{1}{p_1^2} & 0 & 1 \end{pmatrix}, \quad (213)$$

with the exponent matrix

$$\begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 - \frac{D}{2} & \frac{D}{2} - 3 & 0 \\ 1 & 0 & 1 - \frac{D}{2} & 0 & \frac{D}{2} - 3 \end{pmatrix}. \quad (214)$$

Using the above matrix of exponents and the integer lattice $0_{3 \times 5}$, we write the unique solution as

$$\begin{aligned} C(p_1^2, p_1^2, 0) &= \frac{C_{1T}^{(245)}}{\det(A_{245}^{(1)'})\Gamma(2)\Gamma(2 - \frac{D}{2})} (p_1^2)^{D/2-2} \\ &= \frac{C_{1T}^{(245)}}{\Gamma(2 - \frac{D}{2})} (p_1^2)^{D/2-3}. \end{aligned} \quad (215)$$

A. The fundamental solution systems as $p_3 \neq 0, p_3^2 = 0$

As $p_3 \neq 0, p_3^2 = 0$, the Feynman parametric representation is

$$C(p_1^2, p_2^2, 0) = i(\Lambda_{\text{RE}}^2)^{4-D} \frac{\Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \int \omega(t)_3 \frac{t_3^{3-D} (t_1 + t_2 + t_3)^{D/2-3}}{(t_2 p_1^2 + t_1 p_2^2)^{3-D/2}}. \quad (216)$$

The integral can be embedded in the subvariety of the Grassmannian $G_{3,5}$ which local coordinates are written as

$$A^{(2)} = \begin{pmatrix} 1 & 0 & 0 & p_2^2 & 1 \\ 0 & 1 & 0 & p_1^2 & 1 \\ 0 & 0 & 1 & 0 & 1 \end{pmatrix}. \quad (217)$$

Here three rows correspond to the integration variables t_i ($i = 1, 2, 3$), the first column represents the power function $t_1^0 = 1$, the second column represents the power function $t_2^0 = 1$, the third column represents the power function t_3^{3-D} , the fourth column represents the function $(t_2 p_1^2 + t_1 p_2^2)^{D/2-3}$, and the fifth column represents $(t_1 + t_2 + t_3)^{D/2-3}$, respectively. The geometric description of the matrix is similarly presented in Fig.1 where $(2, 3, 5)$ is changed into $(1, 2, 4)$, $\{1, 4\}$ is changed into $\{3, 5\}$, respectively. Accordingly, the Feynman integral satisfies the following GKZ-system in splitting local coordinates

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,4} + \vartheta_{1,5}\} C(p_1^2, p_2^2, 0) &= -C(p_1^2, p_2^2, 0), \\ \{\vartheta_{2,2} + \vartheta_{2,4} + \vartheta_{2,5}\} C(p_1^2, p_2^2, 0) &= -C(p_1^2, p_2^2, 0), \\ \{\vartheta_{3,3} + \vartheta_{3,5}\} C(p_1^2, p_2^2, 0) &= -C(p_1^2, p_2^2, 0), \\ \vartheta_{1,1} C(p_1^2, p_2^2, 0) &= 0, \\ \vartheta_{2,2} C(p_1^2, p_2^2, 0) &= 0, \\ \vartheta_{3,3} C(p_1^2, p_2^2, 0) &= (3 - D) C(p_1^2, p_2^2, 0), \\ \{\vartheta_{1,4} + \vartheta_{2,4}\} C(p_1^2, p_2^2, 0) &= (\frac{D}{2} - 3) C(p_1^2, p_2^2, 0), \\ \{\vartheta_{1,5} + \vartheta_{2,5} + \vartheta_{3,5}\} C(p_1^2, p_2^2, 0) &= (\frac{D}{2} - 3) C(p_1^2, p_2^2, 0). \end{aligned} \quad (218)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & 0 & 0 & \alpha_{1,4} & \alpha_{1,5} \\ 0 & 0 & 0 & \alpha_{2,4} & \alpha_{2,5} \\ 0 & 0 & 3 - D & 0 & D - 4 \end{pmatrix}, \quad (219)$$

with

$$\begin{aligned}\alpha_{1,4} + \alpha_{1,5} &= -1, \quad \alpha_{2,4} + \alpha_{2,5} = -1, \\ \alpha_{1,4} + \alpha_{2,4} &= \frac{D}{2} - 3, \quad \alpha_{1,5} + \alpha_{2,5} = 1 - \frac{D}{2}.\end{aligned}\quad (220)$$

The dual space of the GKZ-system is spanned by the integer matrix whose submatrix composed of the fourth and fifth columns is formulated as $E_3^{(3)}$, and other elements are all zero.

The basis of the dual space implies the Feynman integral satisfying the PDE as

$$\frac{\partial^2 C}{\partial z_{1,4} \partial z_{2,5}}(p_1^2, p_2^2, 0) = \frac{\partial^2 C}{\partial z_{1,5} \partial z_{2,4}}(p_1^2, p_2^2, 0). \quad (221)$$

When the exponent matrix is given, the Feynman integral can be formally expressed as

$$C(p_1^2, p_2^2, 0) = \prod_{i,j} z_{i,j}^{\alpha_{i,j}} \varphi\left(\frac{p_1^2}{p_2^2}\right), \quad (222)$$

where $\varphi(x)$ satisfies the PDE

$$\begin{aligned}& x \left[\alpha_{1,4} \alpha_{2,5} + (1 - \alpha_{1,4} - \alpha_{2,5}) x \frac{d}{dx} - x^2 \frac{d^2}{dx^2} \right] \varphi(x) \\ &= \left[\alpha_{1,5} \alpha_{2,4} + (1 + \alpha_{1,5} + \alpha_{2,4}) x \frac{d}{dx} - x^2 \frac{d^2}{dx^2} \right] \varphi(x).\end{aligned}\quad (223)$$

As $\alpha_{1,5} \alpha_{2,4} = 0$ this equation recovers the well-known Gauss equation.

There are two choices on the matrix of integer lattice whose submatrix composed of the fourth and fifth columns is formulated as $\pm n E_3^{(3)}$ with $n \geq 0$. The matrix of integer lattice $n E_3^{(3)}$ permits two choices on the exponents. The first one is

$$\alpha_{1,4} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{2,4} = \frac{D}{2} - 3, \quad \alpha_{2,5} = 2 - \frac{D}{2}, \quad (224)$$

whose geometric representation is plotted in Fig.2(2) with $(a, b) = (2, 4)$, $\{c, d, e\} = \{1, 3, 5\}$.

The second choice of exponents is

$$\alpha_{1,4} = \frac{D}{2} - 2, \quad \alpha_{1,5} = 1 - \frac{D}{2}, \quad \alpha_{2,4} = -1, \quad \alpha_{2,5} = 0, \quad (225)$$

whose geometric description is presented in Fig.2(1) with $a = 1$, $\{(b, c), (d, e)\} = \{(2, 4), (3, 5)\}$, respectively. The corresponding hypergeometric functions are evidently written as

$$\begin{aligned}\psi_{\{1,2,3\}}^{(1)}(p_1^2, p_2^2, 0) &\sim (p_1^2)^{D/2-3} \sum_{n=0}^{\infty} \left(\frac{p_2^2}{p_1^2}\right)^n \sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2}, \\ \psi_{\{1,2,3\}}^{(2)}(p_1^2, p_2^2, 0) &\sim \frac{(p_2^2)^{D/2-2}}{p_1^2} \sum_{n=0}^{\infty} \left(\frac{p_2^2}{p_1^2}\right)^n \sim \frac{(p_2^2)^{D/2-2}}{p_1^2 - p_2^2},\end{aligned}\quad (226)$$

where the intersection of their convergent regions is $|p_2^2/p_1^2| \leq 1$. The linearly independent Gauss functions constitute the fundamental solution system of the GKZ-system of Eq.(218) in the region $|p_2^2/p_1^2| \leq 1$ which is consistent with the results in the literature. As $\mu = \nu = \rho = 1$ and $k^2 = 0$ in Eq.(7) of Ref. [16], the fourth type Appell functions of the second and third terms are modified into the Gauss functions in Eq.(226).

The integer lattice $-nE_3^{(3)}$ permits two choices of the exponents. The first choice is

$$\alpha_{1,4} = -1, \alpha_{1,5} = 0, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{2,5} = 1 - \frac{D}{2}, \quad (227)$$

whose geometric representation is drawn in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, and the second one is

$$\alpha_{1,4} = \frac{D}{2} - 3, \alpha_{1,5} = 2 - \frac{D}{2}, \alpha_{2,4} = 0, \alpha_{2,5} = -1. \quad (228)$$

whose geometric representation is presented in Fig.2(2) with $(a, b) = (1, 4)$, $\{c, d, e\} = \{2, 3, 5\}$. The hypergeometric functions are

$$\begin{aligned} \psi_{\{1,2,3\}}^{(3)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_2^2} \sum_{n=0}^{\infty} \left(\frac{p_1^2}{p_2^2}\right)^n \sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2}, \\ \psi_{\{1,2,3\}}^{(4)}(p_1^2, p_2^2, 0) &\sim (p_2^2)^{D/2-3} \sum_{n=0}^{\infty} \left(\frac{p_1^2}{p_2^2}\right)^n \sim \frac{(p_2^2)^{D/2-2}}{p_1^2 - p_2^2}, \end{aligned} \quad (229)$$

whose intersection of their convergent regions is $|p_1^2/p_2^2| \leq 1$. The projective plane can also spanned by the first, second, and fifth columns because $\det(A_{\{1,2,5\}}^{(2)}) = 1$, the hypergeometric solutions based on the matrix $(A_{\{1,2,5\}}^{(2)})^{-1} \cdot A^{(2)}$ are same as those presented above.

Because $\det(A_{\{1,3,4\}}^{(2)}) = -p_1^2$, where the matrix $A_{\{1,3,4\}}^{(2)}$ denotes the submatrix of $A^{(2)}$ composed of the first, third, and fourth columns, one derives

$$(A_{\{1,3,4\}}^{(2)})^{-1} \cdot A^{(2)} = \begin{pmatrix} 1 & -\frac{p_2^2}{p_1^2} & 0 & 0 & 1 - \frac{p_2^2}{p_1^2} \\ 0 & 0 & 1 & 0 & 1 \\ 0 & \frac{1}{p_1^2} & 0 & 1 & \frac{1}{p_1^2} \end{pmatrix}. \quad (230)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & 0 & 0 & \alpha_{1,5} \\ 0 & 0 & 3 - D & 0 & D - 4 \\ 0 & \alpha_{3,2} & 0 & \frac{D}{2} - 3 & \alpha_{3,5} \end{pmatrix}, \quad (231)$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,2} + \alpha_{1,5} &= -1, \quad \alpha_{3,2} + \alpha_{3,5} = 2 - \frac{D}{2}, \\ \alpha_{1,2} + \alpha_{3,2} &= 0, \quad \alpha_{1,5} + \alpha_{3,5} = 1 - \frac{D}{2}.\end{aligned}\tag{232}$$

There are two choices on the matrix of integer lattice whose submatrix composed of the second and fifth columns is formulated as $\pm nE_3^{(2)}$ with $n \geq 0$, and other elements are all zero. The matrix of integer lattice $nE_3^{(2)}$ permits two choices on the exponents. The first choice is

$$\alpha_{1,2} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{3,2} = 0, \quad \alpha_{3,5} = 2 - \frac{D}{2},\tag{233}$$

whose geometric description is presented in Fig.2(2) with $(a, b) = (2, 4)$, $\{c, d, e\} = \{1, 3, 5\}$. The second choice is

$$\alpha_{1,2} = \frac{D}{2} - 2, \quad \alpha_{1,5} = 1 - \frac{D}{2}, \quad \alpha_{3,2} = 2 - \frac{D}{2}, \quad \alpha_{3,5} = 0,\tag{234}$$

whose geometric description is presented in Fig.2(1) with $a = 1$, $\{(b, c), (d, e)\} = \{(2, 4), (3, 5)\}$, respectively. Two hypergeometric functions are evidently written as

$$\begin{aligned}\psi_{\{1,3,4\}}^{(1)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2} \sum_{n=0}^{\infty} \frac{1}{\Gamma(3 - \frac{D}{2} + n)\Gamma(1 - n)} \left(\frac{p_2^2}{p_1^2 - p_2^2}\right)^n \\ &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2}, \\ \psi_{\{1,3,4\}}^{(2)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2 p_2^2)^{D/2-2}}{(p_1^2 - p_2^2)^{D/2-1}} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n)}{n!} \left(\frac{p_2^2}{p_2^2 - p_1^2}\right)^n \\ &\sim \frac{(p_2^2)^{D/2-2}}{p_1^2 - p_2^2},\end{aligned}\tag{235}$$

whose intersection of their convergent regions is $|p_2^2/(p_2^2 - p_1^2)| \leq 1$.

Corresponding to the integer lattice $-nE_3^{(2)}$, the first choice of exponents is

$$\alpha_{1,2} = -1, \quad \alpha_{1,5} = 0, \quad \alpha_{3,2} = 1, \quad \alpha_{3,5} = 1 - \frac{D}{2},\tag{236}$$

whose geometric representation is drawn in Fig.2(1) with $a = 4$, $\{(b, c), (d, e)\} = \{(1, 2), (3, 5)\}$, and the second choice is

$$\alpha_{1,2} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{3,2} = 0, \quad \alpha_{3,5} = 2 - \frac{D}{2}.\tag{237}$$

where the geometric representation is drawn in Fig.2(2) with $(a, b) = (1, 2)$, $\{c, d, e\} = \{3, 4, 5\}$. The corresponding hypergeometric functions are written as

$$\begin{aligned}\psi_{\{1,3,4\}}^{(3)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_2^2} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n)}{\Gamma(2 + n)} \left(\frac{p_2^2 - p_1^2}{p_2^2}\right)^n \\ &\sim \frac{(p_2^2)^{D/2-2} - (p_1^2)^{D/2-2}}{p_1^2 - p_2^2}, \\ \psi_{\{1,3,4\}}^{(4)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2} \sum_{n=1}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n)}{\Gamma(n)} \left(\frac{p_2^2 - p_1^2}{p_2^2}\right)^n \\ &\sim \frac{(p_2^2)^{D/2-2}}{p_1^2},\end{aligned}\tag{238}$$

whose intersection of their convergent regions is $|(p_2^2 - p_1^2)/p_2^2| \leq 1$.

Since $\det(A_{\{2,3,4\}}^{(2)}) = p_2^2$, where the matrix $A_{\{2,3,4\}}^{(2)}$ denotes the submatrix of $A^{(2)}$ composed of the second, third, and fourth columns, we have

$$\left(A_{\{2,3,4\}}^{(2)}\right)^{-1} \cdot A^{(2)} = \begin{pmatrix} -\frac{p_1^2}{p_2^2} & 1 & 0 & 0 & 1 - \frac{p_1^2}{p_2^2} \\ 0 & 0 & 1 & 0 & 1 \\ \frac{1}{p_2^2} & 0 & 0 & 1 & \frac{1}{p_2^2} \end{pmatrix}.\tag{239}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & 0 & 0 & \alpha_{1,5} \\ 0 & 0 & 3 - D & 0 & D - 4 \\ \alpha_{3,1} & 0 & 0 & \frac{D}{2} - 3 & \alpha_{3,5} \end{pmatrix},\tag{240}$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,1} + \alpha_{1,5} &= -1, \quad \alpha_{3,1} + \alpha_{3,5} = 2 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{3,1} &= 0, \quad \alpha_{1,5} + \alpha_{3,5} = 1 - \frac{D}{2}.\end{aligned}\tag{241}$$

There are two choices on the matrix of integer lattice whose submatrix composed of the first and fifth columns is formulated as $\pm nE_3^{(2)}$ with $n \geq 0$, and other elements are all zero. The matrix of integer lattice $nE_3^{(2)}$ permits two choices on the exponents. The first choice is

$$\alpha_{1,1} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{3,1} = 0, \quad \alpha_{3,5} = 2 - \frac{D}{2},\tag{242}$$

whose geometric description is presented in Fig.2(2) with $(a, b) = (1, 4)$, $\{c, d, e\} = \{2, 3, 5\}$.

The second choice is

$$\alpha_{1,1} = \frac{D}{2} - 2, \quad \alpha_{1,5} = 1 - \frac{D}{2}, \quad \alpha_{3,1} = 2 - \frac{D}{2}, \quad \alpha_{3,5} = 0,\tag{243}$$

whose geometric description is presented in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, respectively.

The first choice of exponents of the integer lattice $-nE_3^{(2)}$ is

$$\alpha_{1,1} = -1, \alpha_{1,5} = 0, \alpha_{3,1} = 1, \alpha_{3,5} = 1 - \frac{D}{2}, \quad (244)$$

whose geometric representation is drawn in Fig.2(1) with $a = 4$, $\{(b, c), (d, e)\} = \{(1, 2), (3, 5)\}$. The second choice is

$$\alpha_{1,1} = 0, \alpha_{1,5} = -1, \alpha_{3,1} = 0, \alpha_{3,5} = 2 - \frac{D}{2}. \quad (245)$$

whose geometric representation is drawn in Fig.2(2) with $(a, b) = (1, 2)$, $\{c, d, e\} = \{3, 4, 5\}$. The corresponding hypergeometric functions are evidently written as

$$\psi_{\{2,3,4\}}^{(i)}(p_1^2, p_2^2, 0) \sim \psi_{\{1,3,4\}}^{(i)}(p_2^2, p_1^2, 0), \quad (i = 1, 2, 3, 4). \quad (246)$$

One obviously has $\det(A_{\{1,4,5\}}^{(2)}) = p_1^2$, thus derives

$$\left(A_{\{1,4,5\}}^{(2)}\right)^{-1} \cdot A^{(2)} = \begin{pmatrix} 1 & -\frac{p_2^2}{p_1^2} & -1 + \frac{p_2^2}{p_1^2} & 0 & 0 \\ 0 & \frac{1}{p_1^2} & -\frac{1}{p_1^2} & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{pmatrix}, \quad (247)$$

where the matrix $A_{\{1,4,5\}}^{(2)}$ denotes the submatrix of $A^{(2)}$ composed of the first, fourth, and fifth columns. Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & 0 & 0 \\ 0 & \alpha_{2,2} & \alpha_{2,3} & \frac{D}{2} - 3 & 0 \\ 0 & 0 & 2 - \frac{D}{2} & 0 & \frac{D}{2} - 3 \end{pmatrix}, \quad (248)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,2} + \alpha_{1,3} &= -1, \quad \alpha_{2,2} + \alpha_{2,3} = 2 - \frac{D}{2}, \\ \alpha_{1,2} + \alpha_{2,2} &= 0, \quad \alpha_{1,3} + \alpha_{2,3} = 1 - \frac{D}{2}. \end{aligned} \quad (249)$$

There are two choices on the matrix of integer lattice whose submatrix composed of the second and third columns is formulated as $\pm nE_3^{(1)}$ with $n \geq 0$, and other elements are all

zero. The matrix of integer lattice $nE_3^{(1)}$ permits two choices on the exponents. The first choice is

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \quad (250)$$

whose geometric description is presented in Fig.2(2) with $(a, b) = (2, 4)$, $\{c, d, e\} = \{1, 3, 5\}$.

The second one is

$$\alpha_{1,2} = \frac{D}{2} - 2, \alpha_{1,3} = 1 - \frac{D}{2}, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \quad (251)$$

whose geometric description is presented in Fig.2(1) with $a = 1$, $\{(b, c), (d, e)\} = \{(2, 4), (3, 5)\}$, respectively. The hypergeometric functions are

$$\begin{aligned} \psi_{\{1,4,5\}}^{(1)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2} \sum_{n=0}^{\infty} \frac{1}{\Gamma(3 - \frac{D}{2} + n)\Gamma(1 - n)} \left(\frac{p_2^2}{p_1^2 - p_2^2}\right)^n \\ &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2}, \\ \psi_{\{1,4,5\}}^{(2)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2 p_2^2)^{D/2-2}}{(p_1^2 - p_2^2)^{D/2-1}} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n)}{n!} \left(\frac{p_2^2}{p_2^2 - p_1^2}\right)^n \\ &\sim \frac{(p_2^2)^{D/2-2}}{p_1^2 - p_2^2}, \end{aligned} \quad (252)$$

whose intersection of their convergent regions is $|p_2^2/(p_2^2 - p_1^2)| \leq 1$.

Corresponding to the integer lattice $-nE_3^{(2)}$, the first choice of exponents is

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 1, \alpha_{2,3} = 1 - \frac{D}{2}, \quad (253)$$

whose geometric representation is drawn in Fig.2(1) with $a = 4$, $\{(b, c), (d, e)\} = \{(1, 2), (3, 5)\}$. The second choice is

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \quad (254)$$

whose geometric representation is drawn in Fig.2(2) with $(a, b) = (1, 2)$, $\{c, d, e\} = \{3, 4, 5\}$.

The hypergeometric functions are presented as

$$\begin{aligned} \psi_{\{1,4,5\}}^{(3)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_2^2} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n)}{\Gamma(2 + n)} \left(\frac{p_2^2 - p_1^2}{p_2^2}\right)^n \\ &\sim \frac{(p_2^2)^{D/2-2} - (p_1^2)^{D/2-2}}{p_2^2 - p_1^2}, \\ \psi_{\{1,4,5\}}^{(4)}(p_1^2, p_2^2, 0) &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2}, \end{aligned} \quad (255)$$

whose intersection of their convergent regions is $|(p_2^2 - p_1^2)/p_2^2| \leq 1$.

Since $\det(A_{\{2,4,5\}}^{(2)}) = -p_2^2$, one derives

$$\left(A_{\{2,4,5\}}^{(2)}\right)^{-1} \cdot A^{(2)} = \begin{pmatrix} -\frac{p_1^2}{p_2^2} & 1 & -1 + \frac{p_1^2}{p_2^2} & 0 & 0 \\ \frac{1}{p_2^2} & 0 & -\frac{1}{p_2^2} & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{pmatrix}, \quad (256)$$

where the matrix $A_{\{2,4,5\}}^{(2)}$ denotes the submatrix of $A^{(2)}$ consisted in the second, fourth, and fifth columns. Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} & 0 & 0 \\ \alpha_{2,1} & 0 & \alpha_{2,3} & \frac{D}{2} - 3 & 0 \\ 0 & 0 & 2 - \frac{D}{2} & 0 & \frac{D}{2} - 3 \end{pmatrix}, \quad (257)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,3} &= -1, \quad \alpha_{2,1} + \alpha_{2,3} = 2 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{2,1} &= 0, \quad \alpha_{1,3} + \alpha_{2,3} = 1 - \frac{D}{2}. \end{aligned} \quad (258)$$

There are two choices on the matrix of integer lattice whose submatrix composed of the first and third columns is formulated as $\pm nE_3^{(3)}$ with $n \geq 0$, and other elements are all zero. The matrix of integer lattice $nE_3^{(3)}$ permits two choices on exponents. The first one is

$$\alpha_{1,1} = 0, \quad \alpha_{1,3} = -1, \quad \alpha_{2,1} = 0, \quad \alpha_{2,3} = 2 - \frac{D}{2}, \quad (259)$$

whose geometric description is presented in Fig.2(2) with $(a, b) = (1, 4)$, $\{c, d, e\} = \{2, 3, 5\}$.

The second choice is

$$\alpha_{1,1} = \frac{D}{2} - 2, \quad \alpha_{1,3} = 1 - \frac{D}{2}, \quad \alpha_{2,1} = 2 - \frac{D}{2}, \quad \alpha_{2,3} = 0, \quad (260)$$

whose geometric description is presented in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$, respectively.

The first choice of exponents of the integer lattice $-nE_3^{(3)}$ is

$$\alpha_{1,1} = -1, \quad \alpha_{1,3} = 0, \quad \alpha_{2,1} = 1, \quad \alpha_{2,3} = 1 - \frac{D}{2}, \quad (261)$$

whose geometric representation is drawn in Fig.2(1) with $a = 4$, $\{(b, c), (d, e)\} = \{(1, 2), (3, 5)\}$. The second choice is

$$\alpha_{1,1} = 0, \quad \alpha_{1,3} = -1, \quad \alpha_{2,1} = 0, \quad \alpha_{2,3} = 2 - \frac{D}{2}. \quad (262)$$

whose geometric representation is drawn in Fig.2(2) with $(a, b) = (1, 2)$, $\{c, d, e\} = \{3, 4, 5\}$.

The hypergeometric functions are

$$\psi_{\{2,4,5\}}^{(i)}(p_1^2, p_2^2, 0) \sim \psi_{\{1,4,5\}}^{(i)}(p_2^2, p_1^2, 0), \quad (i = 1, 2, 3, 4). \quad (263)$$

B. The fundamental solution systems in general case

In the general case, the Feynman parametric representation is

$$\begin{aligned} C(p_1^2, p_2^2, p_3^2) &= i \left(\Lambda_{\text{RE}}^2 \right)^{4-D} \frac{(-)^{D/2-3} \Gamma(3 - \frac{D}{2})}{(4\pi)^{D/2}} \int_0^1 \omega_4(t) t_4^{3-D} \\ &\times \frac{\delta(t_1 + t_2 + t_3 + t_4)}{(t_1 t_2 p_3^2 + t_2 t_3 p_1^2 + t_1 t_3 p_2^2)^{3-D/2}}. \end{aligned} \quad (264)$$

The integral can be embedded in the subvariety of the Grassmannian $G_{4,7}$ where the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , the third row corresponds to the integration variable t_3 , and the fourth row corresponds to the integration variable t_4 , respectively. Meanwhile the first column represents the power function $t_1^0 = 1$, the second column represents the power function $t_2^0 = 1$, the third column represents the power function $t_3^0 = 1$, the fourth column represents the power function t_4^{3-D} , the seventh column represents the function $\delta(t_1 + t_2 + t_3)$. In order to embed the homogeneous polynomial $t_1 t_2 p_3^2 + t_2 t_3 p_1^2 + t_1 t_3 p_2^2$ as the fifth and sixth columns of the Grassmannian $G_{4,7}$, we rewrite

$$\begin{aligned} &t_1 t_2 p_3^2 + t_2 t_3 p_1^2 + t_1 t_3 p_2^2 \\ &= z_{1, \widehat{\sigma_1(5)}} z_{2, \widehat{\sigma_1(6)}} t_1 t_2 + z_{1, \widehat{\sigma_2(5)}} z_{3, \widehat{\sigma_2(6)}} t_1 t_3 + z_{2, \widehat{\sigma_3(5)}} z_{3, \widehat{\sigma_3(6)}} t_2 t_3 \\ &\quad + z_{1, \widehat{\sigma_4(5)}} z_{4, \widehat{\sigma_4(6)}} t_1 t_4 + z_{2, \widehat{\sigma_5(5)}} z_{4, \widehat{\sigma_5(6)}} t_2 t_4 + z_{3, \widehat{\sigma_6(5)}} z_{4, \widehat{\sigma_6(6)}} t_3 t_4, \end{aligned} \quad (265)$$

where $\widehat{\sigma}_i$, $(i = 1, \dots, 6)$ are elements of the permutation group $S_2 = \{\widehat{e}, \widehat{(56)}\}$ on the column indices 5, 6. Taking $\widehat{\sigma}_1 = \widehat{\sigma}_3 = \widehat{\sigma}_4 = \widehat{\sigma}_5 = \widehat{\sigma}_6 = \widehat{(56)}$, $\widehat{\sigma}_2 = \widehat{e}$, we have

$$\begin{aligned} z_{1,6} z_{2,5} &= p_3^2, \quad z_{1,5} z_{3,6} = p_2^2, \quad z_{2,6} z_{3,5} = p_1^2, \\ z_{1,6} z_{4,5} &= 0, \quad z_{2,6} z_{4,5} = 0, \quad z_{3,6} z_{4,5} = 0. \end{aligned} \quad (266)$$

The solution of Eq.(266)

$$\begin{aligned} z_{1,5} &= p_2^2, \quad z_{2,5} = p_3^2, \quad z_{3,5} = p_1^2, \quad z_{4,5} = 0, \\ z_{1,6} &= z_{2,6} = z_{3,6} = 1, \end{aligned} \quad (267)$$

does not constrain the element $z_{4,6}$. For convenience we choose $z_{4,6} = 0$ or $z_{4,6} = 1$, respectively. The integral can be embedded in the subvariety of the Grassmannian $G_{4,7}$ where splitting local coordinates are written as

$$A^{(1T)} = \begin{pmatrix} 1 & 0 & 0 & 0 & p_2^2 & 1 & 1 \\ 0 & 1 & 0 & 0 & p_3^2 & 1 & 1 \\ 0 & 0 & 1 & 0 & p_1^2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}, \quad (268)$$

as $z_{4,6} = 1$, whose geometric description is presented in Fig.5. Taking Feynman parametric representation of the 1-loop massless triangle diagram as a function of the splitting local coordinates,

$$C(p_1^2, p_2^2, p_3^2) = \Psi^{1T}(z_1, \dots, z_7), \quad (269)$$

we obviously have

$$\Psi^{1T}(\chi_1 z_1, \dots, \chi_7 z_7) = \chi_4^{3-D} \chi_5^{D/2-3} \chi_6^{D/2-3} \chi_7^{-1} \Psi^{1T}(z_1, \dots, z_7). \quad (270)$$

Where z_i ($i = 1, \dots, 7$) denotes the i -th column vector in the matrix Eq.(268), and χ_i is a nonzero constant, respectively.

If we choose the first, second, third, and fourth column vectors of $A^{(1T)}$ in Eq.(268) as the basis of the projective plane, the convergent region of the fundamental solution system is the empty subset of the whole parameter space because those constructed hypergeometric functions contain redundant dependent variables. However $\det(A_{\{1,2,3,6\}}^{(1T)}) = 1$ where the matrix $A_{\{1,2,3,6\}}^{(1T)}$ denotes the submatrix of $A^{(1T)}$ composed of the first, second, third and sixth columns, one derives the splitting local coordinates as

$$Z_{4 \times 7} = \left(A_{\{1,2,3,6\}}^{(1T)} \right)^{-1} A^{(1T)} = \begin{pmatrix} 1 & 0 & 0 & -1 & p_2^2 & 0 & 0 \\ 0 & 1 & 0 & -1 & p_3^2 & 0 & 0 \\ 0 & 0 & 1 & -1 & p_1^2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}. \quad (271)$$

Accordingly, the Feynman integral satisfies the following GKZ-system

$$\{\vartheta_{1,1} + \vartheta_{1,4} + \vartheta_{1,5}\} C(p_1^2, p_2^2, p_3^2) = -C(p_1^2, p_2^2, p_3^2),$$




FIG. 5: The geometric description of the matroid $A^{(1T)}$ in Eq.(268) on the projective plane CP^4 , where the points $1, \dots, 7$ denote the indices of columns of the 3×7 matrix. In the representation, point 4 and double point 6,7 lie out the hyperplane $\{1, 2, 3, 5\}$.

$$\begin{aligned}
& \{\vartheta_{2,2} + \vartheta_{2,4} + \vartheta_{2,5}\}C(p_1^2, p_2^2, p_3^2) = -C(p_1^2, p_2^2, p_3^2), \\
& \{\vartheta_{3,3} + \vartheta_{3,4} + \vartheta_{3,5}\}C(p_1^2, p_2^2, p_3^2) = -C(p_1^2, p_2^2, p_3^2), \\
& \{\vartheta_{4,4} + \vartheta_{4,6} + \vartheta_{4,7}\}C(p_1^2, p_2^2, p_3^2) = -C(p_1^2, p_2^2, p_3^2), \\
& \vartheta_{1,1}C(p_1^2, p_2^2, p_3^2) = 0, \\
& \vartheta_{2,2}C(p_1^2, p_2^2, p_3^2) = 0, \\
& \vartheta_{3,3}C(p_1^2, p_2^2, p_3^2) = 0, \\
& \{\vartheta_{1,4} + \vartheta_{2,4} + \vartheta_{3,4} + \vartheta_{4,4}\}C(p_1^2, p_2^2, p_3^2) = (3 - D)C(p_1^2, p_2^2, p_3^2), \\
& \{\vartheta_{1,5} + \vartheta_{2,5} + \vartheta_{3,5}\}C(p_1^2, p_2^2, p_3^2) = \left(\frac{D}{2} - 3\right)C(p_1^2, p_2^2, p_3^2), \\
& \vartheta_{4,6}C(p_1^2, p_2^2, p_3^2) = \left(\frac{D}{2} - 3\right)C(p_1^2, p_2^2, p_3^2), \\
& \vartheta_{4,7}C(p_1^2, p_2^2, p_3^2) = -C(p_1^2, p_2^2, p_3^2), \tag{272}
\end{aligned}$$

which induces the matrix of exponents as

$$\begin{pmatrix} 0 & 0 & 0 & \alpha_{1,4} & \alpha_{1,5} & 0 & 0 \\ 0 & 0 & 0 & \alpha_{2,4} & \alpha_{2,5} & 0 & 0 \\ 0 & 0 & 0 & \alpha_{3,4} & \alpha_{3,5} & 0 & 0 \\ 0 & 0 & 0 & 3 - \frac{D}{2} & 0 & \frac{D}{2} - 3 & -1 \end{pmatrix}, \quad (273)$$

with

$$\begin{aligned} \alpha_{1,4} + \alpha_{1,5} &= -1, \quad \alpha_{2,4} + \alpha_{2,5} = -1, \quad \alpha_{3,4} + \alpha_{3,5} = -1, \\ \alpha_{1,4} + \alpha_{2,4} + \alpha_{3,4} &= -\frac{D}{2}, \quad \alpha_{1,5} + \alpha_{2,5} + \alpha_{3,5} = \frac{D}{2} - 3. \end{aligned} \quad (274)$$

Corresponding to the matrix of local coordinates in Eq.(271), we get 12 choices on the matrix




FIG. 6: The geometric descriptions of single orbits in Eq.(268) on the projective plane CP^4 , where point 4 and double point (6, 7) lie out the hyperplane $\{1, 2, 3, 5\}$.

of integer lattice whose submatrix composed of the fourth and fifth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other elements are all zero. Here the bases of integer lattice are written as

$$E_4^{(1)} = \begin{pmatrix} 1 & -1 \\ -1 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad E_4^{(2)} = \begin{pmatrix} 1 & -1 \\ 0 & 0 \\ -1 & 1 \\ 0 & 0 \end{pmatrix}, \quad E_4^{(3)} = \begin{pmatrix} 1 & -1 \\ 0 & 0 \\ 0 & 0 \\ -1 & 1 \end{pmatrix},$$

$$E_4^{(4)} = \begin{pmatrix} 0 & 0 \\ 1 & -1 \\ -1 & 1 \\ 0 & 0 \end{pmatrix}, \quad E_4^{(5)} = \begin{pmatrix} 0 & 0 \\ 1 & -1 \\ 0 & 0 \\ -1 & 1 \end{pmatrix}, \quad E_4^{(6)} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & -1 \\ -1 & 1 \end{pmatrix}. \quad (275)$$

Note that $E_4^{(4)} = E_4^{(2)} - E_4^{(1)}$, two linear independent integer lattice matrices $E_4^{(1)}$ and $E_4^{(4)}$ induce two independent PDEs in splitting local coordinates as:

$$\begin{aligned} \frac{\partial^2 C}{\partial z_{1,4} \partial z_{2,5}}(p_1^2, p_2^2, p_3^2) &= \frac{\partial^2 C}{\partial z_{1,5} \partial z_{2,4}}(p_1^2, p_2^2, p_3^2), \\ \frac{\partial^2 C}{\partial z_{2,4} \partial z_{3,5}}(p_1^2, p_2^2, p_3^2) &= \frac{\partial^2 C}{\partial z_{2,5} \partial z_{3,4}}(p_1^2, p_2^2, p_3^2). \end{aligned} \quad (276)$$

When the exponents are given, the Feynman integral can be formally expressed as

$$C(p_1^2, p_2^2, p_3^2) = \prod_{i,j} z_{i,j}^{\alpha_{i,j}} \varphi\left(\frac{p_1^2}{p_3^2}, \frac{p_2^2}{p_3^2}\right), \quad (277)$$

where $(i, j) \in \{(1, 4), (1, 5), (2, 4), (2, 5), (3, 4), (3, 5)\}$, and $\varphi(x_1, x_2)$ satisfies the PDEs in the combined coordinate $x_1 = z_{2,4} z_{3,5} / (z_{1,4} z_{2,5}) = p_1^2 / p_3^2$, $x_2 = z_{1,5} z_{2,4} / (z_{1,4} z_{2,5}) = p_2^2 / p_3^2$ as

$$\begin{aligned} &x_1 \left[\alpha_{1,4} \alpha_{2,5} + (1 - \alpha_{1,4} - \alpha_{2,5}) x_1 \frac{\partial}{\partial x_1} - \alpha_{1,4} x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_1^2 \frac{\partial^2}{\partial x_1^2} \right] \varphi(x_1, x_2) \\ &= \left[\alpha_{2,4} \alpha_{3,5} + (1 + \alpha_{2,4} + \alpha_{3,5}) x_1 \frac{\partial}{\partial x_1} + \alpha_{2,4} x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_1^2 \frac{\partial^2}{\partial x_1^2} \right] \varphi(x_1, x_2), \\ &x_2 \left[\alpha_{1,4} \alpha_{2,5} - \alpha_{1,4} x_1 \frac{\partial}{\partial x_1} + (1 - \alpha_{1,4} - \alpha_{2,5}) x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right] \varphi(x_1, x_2) \\ &= \left[\alpha_{1,5} \alpha_{2,4} + \alpha_{1,5} x_1 \frac{\partial}{\partial x_1} + (1 + \alpha_{1,5} + \alpha_{2,4}) x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right] \varphi(x_1, x_2). \end{aligned} \quad (278)$$

As $\alpha_{1,5} = \alpha_{2,4} = 0$, the PDEs above are changed into the first type Appell PDEs. The holonomic rank of the ideal composed of the partial differential operators of above PDEs is 3, which indicates that three linear independent hypergeometric functions constitute the fundamental solution system in neighborhoods of regular singular points. It is known that the fourth type Appell function cannot be embedded into the subvarieties of Grassmannians [14]. Nevertheless, including the appropriate power factor, the Feynman integral presented through the fourth Appell functions [16, 41, 42] satisfies the PDEs above. Those matrices of integer lattice permit 12 possibilities of the exponents.

The exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are

$$\alpha_{1,4} = 2 - \frac{D}{2}, \quad \alpha_{1,5} = \frac{D}{2} - 3, \quad \alpha_{2,4} = -1, \quad \alpha_{2,5} = 0, \quad \alpha_{3,4} = -1, \quad \alpha_{3,5} = 0, \quad (279)$$

whose geometric representation is presented in Fig.6(1) with $(a, b) = (1, 5)$, $\{c, d\} = \{2, 3\}$.

As the integer lattice is chosen as $n_1 E_4^{(1)} + n_2 E_4^{(4)}$, the matrix of exponents is

$$\alpha_{1,4} = 0, \alpha_{1,5} = -1, \alpha_{2,4} = 1 - \frac{D}{2}, \alpha_{2,5} = \frac{D}{2} - 2, \alpha_{3,4} = -1, \alpha_{3,5} = 0, \quad (280)$$

whose geometric representation is shown in Fig.6(2) with $\{a, b, c\} = \{1, 2, 5\}$, $d = 3$. When the integer lattice is chosen as $n_1 E_4^{(2)} + n_2 E_4^{(4)}$, the exponents are

$$\alpha_{1,4} = 0, \alpha_{1,5} = -1, \alpha_{2,4} = 0, \alpha_{2,5} = -1, \alpha_{3,4} = -\frac{D}{2}, \alpha_{3,5} = \frac{D}{2} - 1, \quad (281)$$

whose geometric representation is drawn in Fig.6(3) with $a = 3$, $(b, c, d) = (1, 2, 5)$. Accordingly the constructed hypergeometric functions are

$$\begin{aligned} \psi_{\{1,2,3,6\}}^{(1)}(p_1^2, p_2^2, p_3^2) &\sim (p_2^2)^{D/2-3} \sum_{n_1, n_2}^{\infty} \left(\frac{p_3^2}{p_2^2}\right)^{n_1} \left(\frac{p_1^2}{p_2^2}\right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\ \psi_{\{1,2,3,6\}}^{(2)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_3^2)^{D/2-2}}{p_2^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_3^2}{p_2^2}\right)^{n_1} \left(\frac{p_1^2}{p_3^2}\right)^{n_2} \\ &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\ \psi_{\{1,2,3,6\}}^{(3)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{p_2^2 p_3^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_1^2}{p_2^2}\right)^{n_1} \left(\frac{p_1^2}{p_3^2}\right)^{n_2} \\ &\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)}, \end{aligned} \quad (282)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space.

The exponents corresponding to the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(2)}$ are

$$\alpha_{1,4} = -\frac{D}{2}, \alpha_{1,5} = \frac{D}{2} - 1, \alpha_{2,4} = 0, \alpha_{2,5} = -1, \alpha_{3,4} = 0, \alpha_{3,5} = -1, \quad (283)$$

whose geometric representation is plotted in Fig.6(3) with $a = 1$, $\{b, c, d\} = \{2, 3, 5\}$. The exponents of the inter lattice $-n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = -1, \alpha_{1,5} = 0, \alpha_{2,4} = 1 - \frac{D}{2}, \alpha_{2,5} = \frac{D}{2} - 2, \alpha_{3,4} = 0, \alpha_{3,5} = -1, \quad (284)$$

whose geometric representation is plotted in Fig.6(2) with $\{a, b, c\} = \{2, 3, 5\}$, $d = 1$. Similarly the exponents of the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = -1, \alpha_{1,5} = 0, \alpha_{2,4} = -1, \alpha_{2,5} = 0, \alpha_{3,4} = 2 - \frac{D}{2}, \alpha_{3,5} = \frac{D}{2} - 3, \quad (285)$$

whose geometric representation is presented in Fig.6(1) with $(a, b) = (3, 5)$, $\{c, d\} = \{1, 2\}$.

The corresponding hypergeometric functions are

$$\begin{aligned}
\psi_{\{1,2,3,6\}}^{(4)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_2^2)^{D/2-1}}{p_1^2 p_3^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2}{p_3^2}\right)^{n_1} \left(\frac{p_2^2}{p_1^2}\right)^{n_2} \\
&\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\
\psi_{\{1,2,3,6\}}^{(5)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_3^2)^{D/2-2}}{p_1^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2}{p_3^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2} \\
&\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,2,3,6\}}^{(6)}(p_1^2, p_2^2, p_3^2) &\sim (p_1^2)^{D/2-3} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2}{p_1^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2} \\
&\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)}, \tag{286}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The fact indicates that those hypergeometric functions constitute a fundamental solution system of the GKZ-system in the proper subset of the whole parameter space.

Corresponding to the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(2)}$, the exponents are

$$\alpha_{1,4} = 1 - \frac{D}{2}, \alpha_{1,5} = \frac{D}{2} - 2, \alpha_{2,4} = -1, \alpha_{2,5} = 0, \alpha_{3,4} = 0, \alpha_{3,5} = -1, \tag{287}$$

whose geometric representation is drawn in Fig.6(2) with $\{a, b, c\} = \{1, 3, 5\}$, $d = 2$. The exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = 0, \alpha_{1,5} = -1, \alpha_{2,4} = -\frac{D}{2}, \alpha_{2,5} = \frac{D}{2} - 1, \alpha_{3,4} = 0, \alpha_{3,5} = -1, \tag{288}$$

whose geometric representation is presented in Fig.6(3) with $a = 2$, $\{b, c, d\} = \{1, 3, 5\}$. As the integer lattice is chosen as $n_1 E_4^{(2)} - n_2 E_4^{(4)}$, the exponents are

$$\alpha_{14} = 0, \alpha_{15} = -1, \alpha_{24} = -1, \alpha_{25} = 0, \alpha_{34} = 1 - \frac{D}{2}, \alpha_{35} = \frac{D}{2} - 2, \tag{289}$$

whose geometric representation is presented in Fig.6(2) with $\{a, b, c\} = \{1, 3, 5\}$, $d = 2$.

Those hypergeometric functions are

$$\psi_{\{1,2,3,6\}}^{(7)}(p_1^2, p_2^2, p_3^2) \sim \frac{(p_2^2)^{D/2-2}}{p_1^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_3^2}{p_2^2}\right)^{n_1} \left(\frac{p_2^2}{p_1^2}\right)^{n_2}$$

$$\begin{aligned}
& \sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)} , \\
\psi_{\{1,2,3,6\}}^{(8)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_3^2)^{D/2-1}}{p_1^2 p_2^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_3^2}{p_2^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2} \\
& \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} , \\
\psi_{\{1,2,3,6\}}^{(9)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-2}}{p_2^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_1^2}{p_2^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2} \\
& \sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)} . \tag{290}
\end{aligned}$$

Although the intersection of the convergent regions of these series is the empty subset of the whole parameter space, the sets of the hypergeometric functions $\{\psi_{\{1,2,3,6\}}^{(1)}, \psi_{\{1,2,3,6\}}^{(8)}, \psi_{\{1,2,3,6\}}^{(9)}\}$ and $\{\psi_{\{1,2,3,6\}}^{(6)}, \psi_{\{1,2,3,6\}}^{(7)}, \psi_{\{1,2,3,6\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

The exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are

$$\alpha_{1,4} = 1 - \frac{D}{2} , \alpha_{1,5} = \frac{D}{2} - 2 , \alpha_{2,4} = 0 , \alpha_{2,5} = -1 , \alpha_{3,4} = -1 , \alpha_{3,5} = 0 , \tag{291}$$

whose geometric representation is presented in Fig.6(2) with $\{a, b, c\} = \{1, 2, 5\}$, $d = 3$. Similarly the exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = -1 , \alpha_{1,5} = 0 , \alpha_{2,4} = 2 - \frac{D}{2} , \alpha_{2,5} = \frac{D}{2} - 3 , \alpha_{3,4} = -1 , \alpha_{3,5} = 0 , \tag{292}$$

whose geometric representation is plotted in Fig.6(1) with $(a, b) = (2, 5)$, $\{c, d\} = \{1, 3\}$. Corresponding to integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(4)}$, the exponents are

$$\alpha_{1,4} = -1 , \alpha_{1,5} = 0 , \alpha_{2,4} = 0 , \alpha_{2,5} = -1 , \alpha_{3,4} = 1 - \frac{D}{2} , \alpha_{3,5} = \frac{D}{2} - 2 , \tag{293}$$

whose geometric representation is drawn in Fig.6(2) with $\{a, b, c\} = \{2, 3, 5\}$, $d = 1$. The corresponding hypergeometric functions are

$$\begin{aligned}
\psi_{\{1,2,3,6\}}^{(10)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_2^2)^{D/2-2}}{p_3^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2}{p_3^2}\right)^{n_1} \left(\frac{p_1^2}{p_2^2}\right)^{n_2} \\
& \sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)} , \\
\psi_{\{1,2,3,6\}}^{(11)}(p_1^2, p_2^2, p_3^2) & \sim (p_3^2)^{D/2-3} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2}{p_3^2}\right)^{n_1} \left(\frac{p_1^2}{p_3^2}\right)^{n_2}
\end{aligned}$$

$$\begin{aligned}
& \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,2,3,6\}}^{(12)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-2}}{p_3^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2}{p_1^2}\right)^{n_1} \left(\frac{p_1^2}{p_3^2}\right)^{n_2} \\
& \sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)}. \tag{294}
\end{aligned}$$

Certainly the intersection of the convergent regions of these series is the empty subset of the whole parameter space, the sets of the hypergeometric functions $\{\psi_{\{1,2,3,6\}}^{(3)}, \psi_{\{1,2,3,6\}}^{(10)}, \psi_{\{1,2,3,6\}}^{(11)}\}$ and $\{\psi_{\{1,2,3,6\}}^{(4)}, \psi_{\{1,2,3,6\}}^{(11)}, \psi_{\{1,2,3,6\}}^{(12)}\}$ constitute the fundamental solution systems of the corresponding nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{\{1,2,5,6\}}^{(1T)}) = p_1^2$ where the matrix $A_{\{1,2,5,6\}}^{(1T)}$ denotes the submatrix of $A^{(1T)}$ composed of the first, second, fifth and sixth columns, we have

$$\left(A_{\{1,2,5,6\}}^{(1T)}\right)^{-1} A^{(1T)} = \begin{pmatrix} 1 & 0 & -\frac{p_2^2}{p_1^2} & -1 + \frac{p_2^2}{p_1^2} & 0 & 0 & 0 \\ 0 & 1 & -\frac{p_3^2}{p_1^2} & -1 + \frac{p_3^2}{p_1^2} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{p_1^2} & -\frac{1}{p_1^2} & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}. \tag{295}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & 0 & \alpha_{1,3} & \alpha_{1,4} & 0 & 0 & 0 \\ 0 & 0 & \alpha_{2,3} & \alpha_{2,4} & 0 & 0 & 0 \\ 0 & 0 & \alpha_{3,3} & \alpha_{3,4} & \frac{D}{2} - 3 & 0 & 0 \\ 0 & 0 & 0 & 3 - \frac{D}{2} & 0 & \frac{D}{2} - 3 & -1 \end{pmatrix}, \tag{296}$$

where the matrix elements satisfy the relations

$$\begin{aligned}
\alpha_{1,3} + \alpha_{1,4} &= -1, \quad \alpha_{2,3} + \alpha_{2,4} = -1, \quad \alpha_{3,3} + \alpha_{3,4} = 2 - \frac{D}{2}, \\
\alpha_{1,3} + \alpha_{2,3} + \alpha_{3,3} &= 0, \quad \alpha_{1,4} + \alpha_{2,4} + \alpha_{3,4} = -\frac{D}{2}. \tag{297}
\end{aligned}$$

Corresponding to the matrix of splitting local coordinates in Eq.(295), we find 12 choices on the matrix of integer lattice whose submatrix composed of the third and fourth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, with $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other elements are all zero. The exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are

$$\alpha_{1,3} = \frac{D}{2} - 1, \quad \alpha_{1,4} = -\frac{D}{2}, \quad \alpha_{2,3} = -1, \quad \alpha_{2,4} = 0, \quad \alpha_{3,3} = 2 - \frac{D}{2}, \quad \alpha_{3,4} = 0, \tag{298}$$

where the geometric representation is presented in Fig.6(3) with $a = 1$, $\{b, c, d\} = \{2, 3, 5\}$. The exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,4} = -1, \alpha_{2,3} = \frac{D}{2} - 2, \alpha_{2,4} = 1 - \frac{D}{2}, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,4} = 0, \quad (299)$$

whose geometric representation is shown in Fig.6(2) with $\{a, b, c\} = \{2, 3, 5\}$, $d = 1$. In a similar way, the exponents of the integer lattice $n_1 E_4^{(2)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,4} = -1, \alpha_{2,3} = 0, \alpha_{2,4} = -1, \alpha_{3,3} = 0, \alpha_{3,4} = 2 - \frac{D}{2}, \quad (300)$$

whose geometric representation is plotted in Fig.6(1) with $(a, b) = (3, 5)$, $\{c, d\} = \{1, 2\}$. The corresponding hypergeometric functions are evidently written as

$$\begin{aligned} \psi_{\{1,2,5,6\}}^{(1)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2 p_2^2)^{D/2-1}}{p_3^2 (p_2^2 - p_1^2)^{D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_2^2 (p_1^2 - p_3^2)}{p_3^2 (p_2^2 - p_1^2)} \right)^{n_1} \left(\frac{p_2^2}{p_2^2 - p_1^2} \right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\ \psi_{\{1,2,5,6\}}^{(2)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1} (p_3^2)^{D/2-2}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)^{D/2-1}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_2^2 (p_3^2 - p_1^2)}{p_3^2 (p_2^2 - p_1^2)} \right)^{n_1} \left(\frac{p_3^2}{p_3^2 - p_1^2} \right)^{n_2} \\ &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\ \psi_{\{1,2,5,6\}}^{(3)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)} \sum_{n_1, n_2}^{\infty} \frac{1}{\Gamma(3 - \frac{D}{2} + n_1 + n_2)} \\ &\quad \times \frac{1}{\Gamma(1 - n_1 - n_2)} \left(\frac{p_2^2}{p_2^2 - p_1^2} \right)^{n_1} \left(\frac{p_3^2}{p_3^2 - p_1^2} \right)^{n_2} \\ &\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)}, \end{aligned} \quad (301)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. In other words, those hypergeometric functions constitute a fundamental solution system of the GKZ-system in the proper subset of the whole parameter space.

The matrix of exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(2)}$ is

$$\alpha_{1,3} = 0, \alpha_{1,4} = -1, \alpha_{2,3} = 0, \alpha_{2,4} = -1, \alpha_{3,3} = 0, \alpha_{3,4} = 2 - \frac{D}{2}, \quad (302)$$

whose geometric representation is presented in Fig.6(1) with $(a, b) = (1, 3)$, $\{c, d\} = \{2, 5\}$. Meanwhile the exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,4} = 0, \alpha_{2,3} = 1, \alpha_{2,4} = -2, \alpha_{3,3} = 0, \alpha_{3,4} = 2 - \frac{D}{2}, \quad (303)$$

whose geometric representation is plotted in Fig.6(2) with $\{a, b, c\} = \{1, 2, 3\}$, $d = 5$. Similarly the exponents of the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,4} = 0, \alpha_{2,3} = -1, \alpha_{2,4} = 0, \alpha_{3,3} = 2, \alpha_{3,4} = -\frac{D}{2}, \quad (304)$$

whose geometric representation is presented in Fig.6(3) with $a = 5$, $\{b, c, d\} = \{1, 2, 3\}$. Three linear independent hypergeometric functions are

$$\begin{aligned} \psi_{\{1,2,5,6\}}^{(4)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_3^2(p_2^2 - p_1^2)}{p_2^2(p_3^2 - p_1^2)}\right)^{n_1} \left(\frac{p_2^2 - p_1^2}{p_2^2}\right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\ \psi_{\{1,2,5,6\}}^{(5)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1} p_3^2}{p_2^2 (p_1^2 - p_3^2)^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_3^2(p_2^2 - p_1^2)}{p_2^2(p_3^2 - p_1^2)}\right)^{n_1} \left(\frac{p_3^2 - p_1^2}{p_3^2}\right)^{n_2} \\ &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\ \psi_{\{1,2,5,6\}}^{(6)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{p_2^2 p_3^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + n_1 + n_2)}{\Gamma(2 + n_1 + n_2)} \\ &\quad \times \left(\frac{p_2^2 - p_1^2}{p_2^2}\right)^{n_1} \left(\frac{p_3^2 - p_1^2}{p_3^2}\right)^{n_2}, \end{aligned} \quad (305)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space.

Corresponding to the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(2)}$, the exponents are

$$\alpha_{1,3} = 1, \alpha_{1,4} = -2, \alpha_{2,3} = -1, \alpha_{2,4} = 0, \alpha_{3,3} = 0, \alpha_{3,4} = 2 - \frac{D}{2}, \quad (306)$$

whose geometric representation is shown in Fig.6(2) with $\{a, b, c\} = \{1, 2, 3\}$, $d = 5$. Accordingly, the exponents of the integer lattice of $n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,4} = -1, \alpha_{2,3} = 0, \alpha_{2,4} = -1, \alpha_{3,3} = 0, \alpha_{3,4} = 2 - \frac{D}{2}, \quad (307)$$

whose geometric representation is plotted in Fig.6(1) with $(a, b) = (2, 3)$, $\{c, d\} = \{1, 5\}$. The matrix of exponents of the integer lattice $n_1 E_4^{(2)} - n_2 E_4^{(4)}$ is given by

$$\alpha_{1,3} = 0, \alpha_{1,4} = -1, \alpha_{2,3} = -1, \alpha_{2,4} = 0, \alpha_{3,3} = 1, \alpha_{3,4} = 1 - \frac{D}{2}, \quad (308)$$

whose geometric representation is plotted in Fig.6(2) with $\{a, b, c\} = \{2, 3, 5\}$, $d = 1$. The corresponding hypergeometric functions are presented as

$$\begin{aligned} \psi_{\{1,2,5,6\}}^{(7)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1} p_2^2}{p_3^2 (p_1^2 - p_2^2)^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_2^2 (p_1^2 - p_3^2)}{p_3^2 (p_1^2 - p_2^2)} \right)^{n_1} \left(\frac{p_2^2 - p_1^2}{p_2^2} \right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\ \psi_{\{1,2,5,6\}}^{(8)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_2^2 (p_3^2 - p_1^2)}{p_3^2 (p_2^2 - p_1^2)} \right)^{n_1} \left(\frac{p_3^2 - p_1^2}{p_3^2} \right)^{n_2} \\ &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\ \psi_{\{1,2,5,6\}}^{(9)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{p_3^2 (p_1^2 - p_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 - n_1 + n_2)}{\Gamma(2 - n_1 + n_2)} \\ &\quad \times \left(\frac{p_2^2}{p_2^2 - p_1^2} \right)^{n_1} \left(\frac{p_3^2 - p_1^2}{p_3^2} \right)^{n_2} \\ &\sim \frac{(p_1^2)^{D/2-1}}{p_3^2 (p_1^2 - p_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_2^2 (p_3^2 - p_1^2)}{p_3^2 (p_2^2 - p_1^2)} \right)^{n_1} \left(\frac{p_3^2 - p_1^2}{p_3^2} \right)^{n_2} \\ &\sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)}, \end{aligned} \quad (309)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. Furthermore, the sets of the hypergeometric functions $\{\psi_{\{1,2,5,6\}}^{(1)}, \psi_{\{1,2,5,6\}}^{(8)}, \psi_{\{1,2,5,6\}}^{(9)}\}$ and $\{\psi_{\{1,2,5,6\}}^{(6)}, \psi_{\{1,2,5,6\}}^{(7)}, \psi_{\{1,2,5,6\}}^{(8)}\}$ constitute also the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

The matrix of exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(2)}$ is

$$\alpha_{1,3} = \frac{D}{2} - 2, \alpha_{1,4} = 1 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{2,4} = -1, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,4} = 0, \quad (310)$$

whose geometric representation is presented in Fig.6(2) with $\{a, b, c\} = \{1, 3, 5\}$, $d = 2$. Correspondingly the exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,4} = 0, \alpha_{2,3} = \frac{D}{2} - 1, \alpha_{2,4} = -\frac{D}{2}, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{3,4} = 0, \quad (311)$$

whose geometric representation is plotted in Fig.6(3) with $a = 2$, $\{b, c, d\} = \{1, 3, 5\}$. Finally the exponents of the integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,4} = 0, \alpha_{2,3} = 0, \alpha_{2,4} = -1, \alpha_{3,3} = 1, \alpha_{3,4} = 1 - \frac{D}{2}, \quad (312)$$

whose geometric representation is presented in Fig.6(2) with $\{a, b, c\} = \{1, 3, 5\}$, $d = 2$. Those hypergeometric functions are formulated as

$$\begin{aligned} \psi_{\{1,2,5,6\}}^{(10)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1} (p_2^2)^{D/2-2}}{(p_3^2 - p_1^2) (p_2^2 - p_1^2)^{D/2-1}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_3^2 (p_2^2 - p_1^2)}{p_2^2 (p_3^2 - p_1^2)} \right)^{n_1} \left(\frac{p_2^2}{p_2^2 - p_1^2} \right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2) (p_2^2 - p_3^2)}, \\ \psi_{\{1,2,5,6\}}^{(11)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2 p_3^2)^{D/2-1}}{p_2^2 (p_3^2 - p_1^2)^{D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2!} \\ &\quad \times \left(\frac{p_3^2 (p_2^2 - p_1^2)}{p_2^2 (p_3^2 - p_1^2)} \right)^{n_1} \left(\frac{p_3^2}{p_3^2 - p_1^2} \right)^{n_2} \\ &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2) (p_3^2 - p_2^2)}, \\ \psi_{\{1,2,5,6\}}^{(12)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-2} (p_3^2)^2}{(p_1^2 - p_3^2)^2 (p_2^2 - p_3^2)} \sum_{n=0}^{\infty} \frac{\Gamma(1+n)}{\Gamma(4 - \frac{D}{2} + n)} \left(\frac{p_3^2}{p_3^2 - p_1^2} \right)^n, \end{aligned} \quad (313)$$

where the intersection of their convergent regions is a nonempty subset of the whole parameter space. The hypergeometric function of the exponents of Eq.(312) is

$$\begin{aligned} \psi_{\{1,2,5,6\}}^{(12)'}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{p_2^2 (p_1^2 - p_3^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n_1 - n_2)}{\Gamma(2 + n_1 - n_2)} \\ &\quad \times \left(\frac{p_2^2 - p_1^2}{p_2^2} \right)^{n_1} \left(\frac{p_3^2}{p_3^2 - p_1^2} \right)^{n_2} \\ &\sim \psi_{\{1,2,5,6\}}^{(10)}(p_1^2, p_2^2, p_3^2) \\ &\quad + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi (\frac{D}{2} - 3)} \psi_{\{1,2,5,6\}}^{(12)}(p_1^2, p_2^2, p_3^2). \end{aligned} \quad (314)$$

The sets of the hypergeometric functions $\{\psi_{\{1,2,5,6\}}^{(3)}, \psi_{\{1,2,5,6\}}^{(10)}, \psi_{\{1,2,5,6\}}^{(11)}\}$ and $\{\psi_{\{1,2,5,6\}}^{(4)}, \psi_{\{1,2,5,6\}}^{(11)}, \psi_{\{1,2,5,6\}}^{(12)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{1,3,5,6})^{(1T)} = -p_3^2$, where the matrix $A_{\{1,3,5,6\}}^{(1T)}$ denotes the submatrix of $A^{(1T)}$ composed of the first, third, fifth and sixth columns, one finds

$$\left(A_{\{1,3,5,6\}}^{(1T)}\right)^{-1} \cdot A^{(1T)} = \begin{pmatrix} 1 & -\frac{p_2^2}{p_3^2} & 0 & -1 + \frac{p_2^2}{p_3^2} & 0 & 0 & 0 \\ 0 & -\frac{p_1^2}{p_3^2} & 1 & -1 + \frac{p_1^2}{p_3^2} & 0 & 0 & 0 \\ 0 & \frac{1}{p_3^2} & 0 & -\frac{1}{p_3^2} & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}. \quad (315)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & 0 & \alpha_{1,4} & 0 & 0 & 0 \\ 0 & \alpha_{2,2} & 0 & \alpha_{2,4} & 0 & 0 & 0 \\ 0 & \alpha_{3,2} & 0 & \alpha_{3,4} & \frac{D}{2} - 3 & 0 & 0 \\ 0 & 0 & 0 & 3 - \frac{D}{2} & 0 & \frac{D}{2} - 3 & -1 \end{pmatrix}, \quad (316)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,2} + \alpha_{1,4} &= -1, \quad \alpha_{2,2} + \alpha_{2,4} = -1, \quad \alpha_{3,2} + \alpha_{3,4} = 2 - \frac{D}{2}, \\ \alpha_{1,2} + \alpha_{2,2} + \alpha_{3,2} &= 0, \quad \alpha_{1,4} + \alpha_{2,4} + \alpha_{3,4} = -\frac{D}{2}. \end{aligned} \quad (317)$$

Correspondingly there are 12 choices on the matrix of integer lattice whose submatrix composed of the second and fourth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents whose geometric descriptions are obtained by the permutation $(\widehat{23})$ of that of $\psi_{\{1,2,5,6\}}^{(i)}$. Correspondingly the hypergeometric functions are

$$\psi_{\{1,3,5,6\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,2,5,6\}}^{(i)}(p_3^2, p_2^2, p_1^2), \quad i = 1, \dots, 12. \quad (318)$$

Because $\det(A_{2,3,5,6})^{(1T)} = p_2^2$, where the matrix $A_{\{2,3,5,6\}}^{(1T)}$ denotes the submatrix of $A^{(1T)}$

composed of the second, third, fifth and sixth columns, one finds

$$\left(A_{\{2,3,5,6\}}^{(1T)}\right)^{-1} \cdot A^{(1T)} = \begin{pmatrix} -\frac{p_3^2}{p_2^2} & 1 & 0 & -1 + \frac{p_3^2}{p_2^2} & 0 & 0 & 0 \\ -\frac{p_1^2}{p_2^2} & 0 & 1 & -1 + \frac{p_1^2}{p_2^2} & 0 & 0 & 0 \\ \frac{1}{p_2^2} & 0 & 0 & -\frac{1}{p_2^2} & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}. \quad (319)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & 0 & \alpha_{1,4} & 0 & 0 & 0 \\ \alpha_{2,1} & 0 & 0 & \alpha_{2,4} & 0 & 0 & 0 \\ \alpha_{3,1} & 0 & 0 & \alpha_{3,4} & \frac{D}{2} - 3 & 0 & 0 \\ 0 & 0 & 0 & 3 - \frac{D}{2} & 0 & \frac{D}{2} - 3 & -1 \end{pmatrix}, \quad (320)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,4} &= -1, \quad \alpha_{2,1} + \alpha_{2,4} = -1, \quad \alpha_{3,1} + \alpha_{3,4} = 2 - \frac{D}{2}, \\ \alpha_{1,1} + \alpha_{2,1} + \alpha_{3,1} &= 0, \quad \alpha_{1,4} + \alpha_{2,4} + \alpha_{3,4} = -\frac{D}{2}. \end{aligned} \quad (321)$$

Correspondingly there are 12 choices on the matrix of integer lattice whose submatrix composed of the second and fourth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents whose geometric descriptions are obtained by the permutation $(\widehat{123})$ of that of $\psi_{\{1,2,5,6\}}^{(i)}$. Correspondingly the hypergeometric functions are

$$\psi_{\{2,3,5,6\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,2,5,6\}}^{(i)}(p_2^2, p_3^2, p_1^2), \quad i = 1, \dots, 12. \quad (322)$$

Since the Equation Eq.(266) does not depend on the splitting local coordinate $z_{4,6}$, we can embed the integral into the subvariety of the Grassmannian $G_{4,7}$ where the local coordinates are

$$A^{(1T)'} = \begin{pmatrix} 1 & 0 & 0 & 0 & p_2^2 & 1 & 1 \\ 0 & 1 & 0 & 0 & p_3^2 & 1 & 1 \\ 0 & 0 & 1 & 0 & p_1^2 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}. \quad (323)$$




FIG. 7: The geometric description of the matroid $A^{(1T)^\vee}$ in Eq.(323) on the projective plane CP^4 , where the points $1, \dots, 7$ denote the indices of columns of the 3×7 matrix. In the representation, the points $5, 6$ belong to the hyperplane $\{1, 2, 3\}$, the line $\{4, 6, 7\}$ intersects with the hyperplane $\{1, 2, 3\}$ at the point 6 , respectively.

Taking $\hat{\sigma}_1 = \hat{\sigma}_3 = \widehat{(56)}$, $\hat{\sigma}_2 = \hat{\sigma}_4 = \hat{\sigma}_5 = \hat{\sigma}_6 = \hat{e}$ in Eq.(265), we have

$$\begin{aligned} z_{1,6} z_{2,5} &= p_3^2, & z_{1,5} z_{3,6} &= p_2^2, & z_{2,6} z_{3,5} &= p_1^2, \\ z_{1,5} z_{4,6} &= 0, & z_{2,5} z_{4,6} &= 0, & z_{3,5} z_{4,6} &= 0. \end{aligned} \quad (324)$$

The solution of Eq.(324)

$$\begin{aligned} z_{1,5} &= p_2^2, & z_{2,5} &= p_3^2, & z_{3,5} &= p_1^2, & z_{4,5} &= z_{4,6} = 0, \\ z_{1,6} &= z_{2,6} &= z_{3,6} &= 1, \end{aligned} \quad (325)$$

corresponds to the subvariety of the Grassmannian $G_{4,7}$ in Eq.(323). Accordingly we obtain the hypergeometric functions which satisfy the PDEs presented in Eq.(278) in combined local coordinates. To shorten the length of text, we collect the hypergeometric functions in appendix A.

The matrices of Eq.(268) and Eq.(323) belong to the embedding of the same parametric representation Eq.(264) of the 1-loop triangle diagram with zero virtual masses on different subvarieties of the Grassmannian $G_{4,7}$. Correspondingly, constructed fundamental solution

systems either belong to the equivalent fundamental solution systems of the neighborhoods of the same regular singularity, or belong to the fundamental solution systems of neighborhoods of different regular singularities.

IV. THE 2-LOOP VACUUM

A. The fundamental system with one zero virtual mass

As one virtual mass $m_1 = 0$, the scalar integral is simplified as

$$A_{2Vac}(0, m_2^2, m_3^2) = -\frac{\Gamma(\frac{D}{2} - 1)\Gamma(3 - D)\mu^{4\epsilon}}{(4\pi)^D\Gamma(\frac{D}{2})} \int \omega_3(t)t_1^{-1}(t_2t_3)^{1-D/2} \times \frac{\delta(t_1 + t_2 + t_3)}{(t_2m_2^2 + t_3m_3^2)^{3-D}}. \quad (326)$$

The integral can be embedded in the subvariety of the Grassmannian $G_{3,5}$ whose splitting local coordinates are written as

$$A^{2V0} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & m_2^2 \\ 0 & 0 & 1 & 1 & m_3^2 \end{pmatrix}, \quad (327)$$

whose geometric description is presented in Fig.1. In the matrix of local coordinates, the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , the third row corresponds to the integration variable t_3 , the first column represents the power function t_1^{-1} , the second column represents the power function $t_2^{1-D/2}$, the third column represents the power function $t_3^{1-D/2}$, the fourth column represents the function $\delta(t_1 + t_2 + t_3)$, and the fifth column represents the power function $(t_2m_2^2 + t_3m_3^2)^{D-3}$, respectively.

Accordingly, the Feynman integral satisfies the following GKZ-system

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,4}\}A_{2Vac}(0, m_2^2, m_3^2) &= -A_{2Vac}(0, m_2^2, m_3^2), \\ \{\vartheta_{2,2} + \vartheta_{2,4} + \vartheta_{2,5}\}A_{2Vac}(0, m_2^2, m_3^2) &= -A_{2Vac}(0, m_2^2, m_3^2), \\ \{\vartheta_{3,3} + \vartheta_{3,4} + \vartheta_{3,5}\}A_{2Vac}(0, m_2^2, m_3^2) &= -A_{2Vac}(0, m_2^2, m_3^2), \\ \vartheta_{1,1}A_{2Vac}(0, m_2^2, m_3^2) &= -A_{2Vac}(0, m_2^2, m_3^2), \\ \vartheta_{2,2}A_{2Vac}(0, m_2^2, m_3^2) &= (1 - \frac{D}{2})A_{2Vac}(0, m_2^2, m_3^2), \end{aligned}$$

$$\begin{aligned}
\vartheta_{3,3}A_{2Vac}(0, m_2^2, m_3^2) &= (1 - \frac{D}{2})A_{2Vac}(0, m_2^2, m_3^2), \\
\{\vartheta_{1,4} + \vartheta_{2,4} + \vartheta_{3,4}\}A_{2Vac}(0, m_2^2, m_3^2) &= -A_{2Vac}(0, m_2^2, m_3^2), \\
\{\vartheta_{2,5} + \vartheta_{3,5}\}A_{2Vac}(0, m_2^2, m_3^2) &= (D - 3)A_{2Vac}(0, m_2^2, m_3^2).
\end{aligned} \tag{328}$$

Certainly the exponent matrix is written as

$$\begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 1 - \frac{D}{2} & 0 & \alpha_{2,4} & \alpha_{2,5} \\ 0 & 0 & 1 - \frac{D}{2} & \alpha_{3,4} & \alpha_{3,5} \end{pmatrix}, \tag{329}$$

with

$$\begin{aligned}
\alpha_{2,4} + \alpha_{2,5} &= \frac{D}{2} - 2, \quad \alpha_{3,4} + \alpha_{3,5} = \frac{D}{2} - 2, \\
\alpha_{2,4} + \alpha_{3,4} &= -1, \quad \alpha_{2,5} + \alpha_{3,5} = D - 3.
\end{aligned} \tag{330}$$

Furthermore, the dual space of the GKZ-system of Eq.(328) is spanned by a 3×5 matrix $(0_{3 \times 3} | E_3^{(1)})$. The basis of the dual space implies the Feynman integral satisfying the PDE

$$\frac{\partial^2 A_{2Vac}}{\partial z_{2,4} \partial z_{3,5}}(0, m_2^2, m_3^2) = \frac{\partial^2 A_{2Vac}}{\partial z_{2,5} \partial z_{3,4}}(0, m_2^2, m_3^2). \tag{331}$$

When the exponents are given, the Feynman integral can be formally expressed as

$$A_{2Vac}(0, m_2^2, m_3^2) = \prod_{i,j} z_{i,j}^{\alpha_{i,j}} \varphi\left(\frac{m_3^2}{m_2^2}\right), \tag{332}$$

where $\varphi(x)$ satisfies the PDE in the combined coordinate $x = z_{2,4}z_{3,5}/z_{2,5}z_{3,4}$ as

$$\begin{aligned}
&x \left[\alpha_{2,4} \alpha_{3,5} + (1 - \alpha_{2,4} - \alpha_{3,5})x \frac{d}{dx} - x^2 \frac{d^2}{dx^2} \right] \varphi(x) \\
&= \left[\alpha_{2,5} \alpha_{3,4} + (1 + \alpha_{2,5} + \alpha_{3,4})x \frac{d}{dx} - x^2 \frac{d^2}{dx^2} \right] \varphi(x).
\end{aligned} \tag{333}$$

At $\alpha_{2,5} \alpha_{3,4} = 0$ this equation recovers the well-known Gauss equation. The integer lattice $(0_{3 \times 3} | nE_3^{(1)})$ ($n \geq 0$) is compatible with two choices of the exponents. The first choice is

$$\alpha_{2,4} = 0, \quad \alpha_{2,5} = \frac{D}{2} - 2, \quad \alpha_{3,4} = -1, \quad \alpha_{3,5} = \frac{D}{2} - 1, \tag{334}$$

whose geometric representation is plotted in Fig.2(1) with $a = 3$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 5)\}$. The second choice is

$$\alpha_{2,4} = 1 - \frac{D}{2}, \quad \alpha_{2,5} = D - 3, \quad \alpha_{3,4} = \frac{D}{2} - 2, \quad \alpha_{3,5} = 0, \tag{335}$$

whose geometric representation is depicted in Fig.2(2) with $(a, b) = (2, 5)$, $\{c, d, e\} = \{1, 3, 4\}$. The corresponding hypergeometric functions are

$$\begin{aligned}\psi_{\{1,2,3\}}^{(1)}(m_2^2, m_3^2) &\sim (m_2^2)^{D/2-2}(m_3^2)^{D/2-1} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(\frac{D}{2} + n)} \left(\frac{m_3^2}{m_2^2}\right)^n, \\ \psi_{\{1,2,3\}}^{(2)}(m_2^2, m_3^2) &\sim (m_2^2)^{D-3} \sum_{n=0}^{\infty} \frac{\Gamma(3 - D + n)}{n!} \left(\frac{m_3^2}{m_2^2}\right)^n \\ &\sim (m_2^2 - m_3^2)^{D-3},\end{aligned}\quad (336)$$

whose the intersection of their convergent regions is $|m_3^2/m_2^2| \leq 1$. Two linearly independent Gauss functions constitute the fundamental solution system of the GKZ-system of Eq.(36) in the region $|m_3^2/m_2^2| \leq 1$ which is consistent with the results in the literature. When $y_i = 0$ in Eq.(46) of Ref. [41], the fourth type Appell functions of the last two terms are simplified as the Gauss functions above.

The third choice of the exponents is

$$\alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{2,5} = 0, \quad \alpha_{3,4} = 1 - \frac{D}{2}, \quad \alpha_{3,5} = D - 3, \quad (337)$$

whose geometric representation is drawn in Fig.2(2) with $(a, b) = (3, 5)$, $\{c, d, e\} = \{1, 2, 4\}$. Similarly the fourth choice of the exponents is

$$\alpha_{2,4} = -1, \quad \alpha_{2,5} = \frac{D}{2} - 1, \quad \alpha_{3,4} = 0, \quad \alpha_{3,5} = \frac{D}{2} - 2, \quad (338)$$

whose geometric representation is shown in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$. The hypergeometric functions are constructed accordingly

$$\begin{aligned}\psi_{\{1,2,3\}}^{(3)}(m_2^2, m_3^2) &\sim (m_3^2)^{D-3} \sum_{n=0}^{\infty} \frac{\Gamma(3 - D + n)}{n!} \left(\frac{m_2^2}{m_3^2}\right)^n \\ &\sim (m_2^2 - m_3^2)^{D-3}, \\ \psi_{\{1,2,3\}}^{(4)}(m_2^2, m_3^2) &\sim (m_2^2)^{D/2-1}(m_3^2)^{D/2-2} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(\frac{D}{2} + n)} \left(\frac{m_2^2}{m_3^2}\right)^n,\end{aligned}\quad (339)$$

where the intersection of their convergent regions is $|m_2^2/m_3^2| \leq 1$.

Because $\det(A_{\{1,2,5\}}^{2V0}) = m_3^2$, thus

$$(A_{\{1,2,5\}}^{2V0})^{-1} \cdot A^{2V0} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & -\frac{m_2^2}{m_3^2} & 1 - \frac{m_2^2}{m_3^2} & 0 \\ 0 & 0 & \frac{1}{m_3^2} & \frac{1}{m_3^2} & 1 \end{pmatrix}. \quad (340)$$

The matrix of exponents is written as

$$\begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 1 - \frac{D}{2} & \alpha_{2,3} & \alpha_{2,4} & 0 \\ 0 & 0 & \alpha_{3,3} & \alpha_{3,4} & D - 3 \end{pmatrix}, \quad (341)$$

with

$$\begin{aligned} \alpha_{2,3} + \alpha_{2,4} &= \frac{D}{2} - 2, \quad \alpha_{3,3} + \alpha_{3,4} = 2 - D, \\ \alpha_{2,3} + \alpha_{3,3} &= 1 - \frac{D}{2}, \quad \alpha_{2,4} + \alpha_{3,4} = -1. \end{aligned} \quad (342)$$

Corresponding to the matrix of local coordinates in Eq.(340), one has two choices on the matrix of integer lattice whose submatrix composed of the third and fourth columns is formulated as $\pm nE_3^{(1)}$, with $n \geq 0$, and other elements are all zero. We have two choices on the exponents of the integer lattice $nE_3^{(1)}$. The first choice is

$$\alpha_{2,3} = 0, \quad \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{3,3} = 1 - \frac{D}{2}, \quad \alpha_{3,4} = 1 - \frac{D}{2}, \quad (343)$$

where the geometric representation is presented in Fig.2(2) with $(a, b) = (3, 5)$, $\{c, d, e\} = \{1, 2, 4\}$. The second choice of the exponents is

$$\alpha_{2,3} = \frac{D}{2} - 1, \quad \alpha_{2,4} = -1, \quad \alpha_{3,3} = 2 - D, \quad \alpha_{3,4} = 0, \quad (344)$$

whose geometric representation is plotted in Fig.2(1) with $a = 2$, $\{(b, c), (d, e)\} = \{(1, 4), (3, 5)\}$. The corresponding hypergeometric functions are

$$\begin{aligned} \psi_{\{1,2,5\}}^{(1)}(m_2^2, m_3^2) &\sim (m_3^2)^{D/2-1} (m_3^2 - m_2^2)^{D/2-2} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n)}{n!} \left(\frac{m_2^2}{m_2^2 - m_3^2}\right)^n \\ &\sim (m_3^2 - m_2^2)^{D-3}, \\ \psi_{\{1,2,5\}}^{(2)}(m_2^2, m_3^2) &\sim \frac{(m_2^2 m_3^2)^{D/2-1}}{m_3^2 - m_2^2} \sum_{n=0}^{\infty} \frac{\Gamma(D - 2 + n)}{\Gamma(\frac{D}{2} + n)} \left(\frac{m_2^2}{m_2^2 - m_3^2}\right)^n, \end{aligned} \quad (345)$$

where the intersection of their convergent regions is $|m_2^2/m_2^2 - m_3^2| \leq 1$. In fact the basis $\{1, 2, 5\}$ of the projective plane composed of the first, second and fifth columns of the matrix Eq.(326) is obtained from the basis $\{1, 2, 3\}$ through the permutation $(\widehat{35})$, accordingly the geometric description of the exponents Eq.(343) is derived from that of the exponents Eq.(337) through the permutation $(\widehat{35})$. Because the geometric description of Eq.(343)

is same as that of Eq.(337), $\psi_{\{1,2,5\}}^{(1)}$ is proportional to $\psi_{\{1,2,3\}}^{(3)}$ in nonempty intersection of their convergent regions. Similarly the geometric description of the exponents Eq.(344) is obtained through the permutation $\widehat{(35)}$ from that of Eq.(338). Because the geometric description of Eq.(344) is same as that of Eq.(338), $\psi_{\{1,2,5\}}^{(2)}$ is proportional to $\psi_{\{1,2,3\}}^{(4)}$ in nonempty intersection of their convergent regions. Actually $\psi_{\{1,2,5\}}^{(2)} \sim \psi_{\{1,2,3\}}^{(4)}$ is consistent with the well-known relation Eq.(56).

There are two choices on the exponents of the integer lattice $-nE_3^{(1)}$. The first choice is

$$\alpha_{2,3} = \frac{D}{2} - 2, \alpha_{2,4} = 0, \alpha_{3,3} = 3 - D, \alpha_{3,4} = -1, \quad (346)$$

where the geometric representation is drawn in Fig.2(1) with $a = 5$, $\{(b, c), (d, e)\} = \{(1, 4), (2, 3)\}$. The second choice is

$$\alpha_{2,3} = 1 - \frac{D}{2}, \alpha_{2,4} = D - 3, \alpha_{3,3} = 0, \alpha_{3,4} = 2 - D, \quad (347)$$

whose geometric representation is plotted in Fig.2(2) with $(a, b) = (2, 3)$, $\{c, d, e\} = \{1, 4, 5\}$. Accordingly the hypergeometric functions are

$$\begin{aligned} \psi_{\{1,2,5\}}^{(3)}(m_2^2, m_3^2) &\sim (m_2^2)^{D/2-2} (m_3^2)^{D/2-1} \sum_{n=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n)}{\Gamma(4 - D + n)} \left(\frac{m_2^2 - m_3^2}{m_2^2}\right)^n, \\ \psi_{\{1,2,5\}}^{(4)}(m_2^2, m_3^2) &\sim \frac{(m_3^2)^{D/2-1} (m_3^2 - m_2^2)^{D-3}}{(m_2^2)^{D/2-1}} \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n)}{n!} \left(\frac{m_2^2 - m_3^2}{m_2^2}\right)^n \\ &\sim (m_3^2 - m_2^2)^{D-3}. \end{aligned} \quad (348)$$

Similarly $\det(A_{\{1,3,5\}}^{2V0}) = -m_2^2$, thus

$$(A_{\{1,3,5\}}^{2V0})^{-1} \cdot A^{2V0} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & -\frac{m_3^2}{m_2^2} & 1 & 1 - \frac{m_3^2}{m_2^2} & 0 \\ 0 & \frac{1}{m_2^2} & 0 & \frac{1}{m_2^2} & 1 \end{pmatrix}. \quad (349)$$

The matrix of exponents is written as

$$\begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & \alpha_{2,2} & 1 - \frac{D}{2} & \alpha_{2,4} & 0 \\ 0 & \alpha_{3,2} & 0 & \alpha_{3,4} & D - 3 \end{pmatrix}, \quad (350)$$

where the entries satisfy the constraints of Eq.(342) with the replacement $\alpha_{i,3} \rightarrow \alpha_{i,2}$, $i = 2, 3$. The geometric descriptions are obtained from that of $\psi_{\{1,2,5\}}^{(i)}$ by the permutation $\widehat{(23)}$. The

corresponding hypergeometric functions are

$$\psi_{\{1,3,5\}}^{(i)}(m_2^2, m_3^2) \sim \psi_{\{1,2,5\}}^{(i)}(m_3^2, m_2^2), \quad i = 1, 2, 3, 4. \quad (351)$$

Because $\det(A_{\{2,3,4\}}^{2V0}) = 1$, thus

$$(A_{\{2,3,4\}}^{2V0})^{-1} \cdot A^{2V0} = \begin{pmatrix} -1 & 1 & 0 & 0 & m_2^2 \\ -1 & 0 & 1 & 0 & m_3^2 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}. \quad (352)$$

The matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 1 - \frac{D}{2} & 0 & 0 & \alpha_{1,5} \\ \alpha_{2,3} & 0 & 1 - \frac{D}{2} & 0 & \alpha_{2,5} \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}, \quad (353)$$

with

$$\begin{aligned} \alpha_{1,1} + \alpha_{1,5} &= \frac{D}{2} - 2, \quad \alpha_{2,1} + \alpha_{2,5} = \frac{D}{2} - 2, \\ \alpha_{1,1} + \alpha_{2,1} &= -1, \quad \alpha_{1,5} + \alpha_{2,5} = D - 3. \end{aligned} \quad (354)$$

Corresponding to the local coordinates in Eq.(352), one finds two choices on the matrix of integer lattice whose submatrix composed of the first and fifth columns is formulated as $\pm nE_3^{(1)}$, where $n \geq 0$, and other elements are all zero. Basing on the matrices of integer lattice, we have 4 choices on the exponents which are obtained that of $\psi_{\{1,2,3\}}^{(i)}$ through the replacements $\alpha_{2,4} \rightarrow \alpha_{1,1}$, $\alpha_{2,5} \rightarrow \alpha_{1,5}$, $\alpha_{3,4} \rightarrow \alpha_{2,1}$, and $\alpha_{3,5} \rightarrow \alpha_{2,5}$, respectively. Certainly the hypergeometric functions are

$$\psi_{\{2,3,4\}}^{(i)}(m_2^2, m_3^2) \sim \psi_{\{1,2,3\}}^{(i)}(m_2^2, m_3^2), \quad i = 1, 2, 3, 4. \quad (355)$$

B. The fundamental solution systems in the general case

In the general case, the Feynman parametric representation is written as

$$\begin{aligned} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= \frac{i^{4-D} \exp\{i\pi(2-D)/2\} \Gamma(3-D) \mu^{4\epsilon}}{(4\pi)^D} \int_0^\infty \prod_{i=1}^3 dt_i \\ &\times (t_1 t_2 t_3)^{1-D/2} \frac{\delta(t_1 t_2 t_3 - t_1 t_2 - t_1 t_3 - t_2 t_3)}{(t_1 m_1^2 + t_2 m_2^2 + t_3 m_3^2)^{3-D}} \end{aligned}$$

$$\begin{aligned}
&= -\frac{i^{4-D} \exp\{i\pi(2-D)/2\} \Gamma(3-D) \mu^{4\epsilon}}{(4\pi)^D} \\
&\quad \times \int \omega(t) \frac{(t_1 t_2 t_3)^{1-D/2} t_4^{D/2-1}}{(t_1 m_1^2 + t_2 m_2^2 + t_3 m_3^2)^{3-D}} \\
&\quad \times \delta(t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4) \tag{356}
\end{aligned}$$

The integral can be embedded in the subvariety of the Grassmannian $G_{4,8}$ where the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , the third row corresponds to the integration variable t_3 , and the fourth row corresponds to the integration variable t_4 , respectively. Meanwhile the first column represents the power function $t_1^{1-D/2}$, the second column represents the power function $t_2^{1-D/2}$, the third column represents the power function $t_3^{1-D/2}$, the fourth column represents the power function $t_4^{D/2-1}$, and the eighth column represents the function $(t_1 m_1^2 + t_2 m_2^2 + t_3 m_3^2)^{D-3}$, respectively. In order to embed the homogeneous function $\delta(t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4)$ as the fifth, sixth, and seventh columns of the Grassmannian $G_{4,8}$, we rewrite

$$\begin{aligned}
&t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4 \\
&= z_{1, \widehat{\tau_1(5)}} z_{2, \widehat{\tau_1(6)}} z_{3, \widehat{\tau_1(7)}} t_1 t_2 t_3 + z_{1, \widehat{\tau_2(5)}} z_{2, \widehat{\tau_2(6)}} z_{4, \widehat{\tau_2(7)}} t_1 t_2 t_4 \\
&\quad + z_{1, \widehat{\tau_3(5)}} z_{3, \widehat{\tau_3(6)}} z_{4, \widehat{\tau_3(7)}} t_1 t_3 t_4 + z_{2, \widehat{\tau_4(5)}} z_{3, \widehat{\tau_4(6)}} z_{4, \widehat{\tau_4(7)}} t_2 t_3 t_4, \tag{357}
\end{aligned}$$

where $\widehat{\tau_i}$, ($i = 1, 2, 3, 4$) are elements of the permutation group S_3 on the column indices 5, 6, 7. Taking $\widehat{\tau_1} = \widehat{\tau_4} = \widehat{e}$, $\widehat{\tau_2} = (\widehat{567})$, and $\widehat{\tau_3} = (\widehat{576})$, we have

$$z_{1,5} z_{2,6} z_{3,7} = z_{2,5} z_{3,6} z_{4,7} = z_{1,6} z_{2,7} z_{4,5} = z_{1,7} z_{3,5} z_{4,6} = 1. \tag{358}$$

The solution of Eq.(358)

$$z_{i,j} = 1, \quad i = 1, \dots, 4, j = 5, 6, 7, \tag{359}$$

indicates that the Feynman parametric representation can be embedded on the subvariety of the Grassmannian $G_{4,8}$

$$A^{(2V)} = \begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 1 & 1 & m_1^2 \\ 0 & 1 & 0 & 0 & 1 & 1 & 1 & m_2^2 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 & m_3^2 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \end{pmatrix}, \tag{360}$$

whose geometric description is presented in Fig.8. Taking parametric representation of the 2-loop vacuum as a function of the splitting local coordinates,

$$A_{2Vac}(m_1^2, m_2^2, m_3^2) = \Psi^{2V}(z_1, \dots, z_8), \quad (361)$$

we obviously have

$$\Psi^{2V}(\chi_1 z_1, \dots, \chi_8 z_8) = (\chi_1 \chi_2 \chi_3)^{1-D/2} \chi_4^{D/2-1} (\chi_5 \chi_6 \chi_7)^{-1} \chi_8^{D-3} \Psi^{2V}(z_1, \dots, z_8). \quad (362)$$

Where z_i ($i = 1, \dots, 8$) denotes the i -th column vector in the matrix Eq.(360), and χ_i is a nonzero constant, respectively. If we choose the first, second, third, and fourth column




FIG. 8: The geometric description of the matroid $A^{(2V)}$ in Eq.(360) on the projective plane CP^4 , where the points $1, \dots, 8$ denote the indices of columns of the 4×8 matrix. In the representation, the point 8 belong to the hyperplane $\{1, 2, 3\}$, points $\{5, 6, 7\}$ are combined into a triple point, respectively.

vectors of $A^{(2V)}$ as the basis of the projective plane, the hypergeometric functions contains redundant interdependent arguments. However $\det(A_{\{1,2,3,5\}}^{(2V)}) = 1$ where the matrix $A_{\{1,2,3,5\}}^{(2V)}$ denotes the submatrix of $A^{(2V)}$ composed of the first, second, third and fifth columns, one

derives

$$\left(A_{\{1,2,3,5\}}^{(2V)} \right)^{-1} \cdot A^{(2V)} = \begin{pmatrix} 1 & 0 & 0 & -1 & 0 & 0 & 0 & m_1^2 \\ 0 & 1 & 0 & -1 & 0 & 0 & 0 & m_2^2 \\ 0 & 0 & 1 & -1 & 0 & 0 & 0 & m_3^2 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \end{pmatrix}. \quad (363)$$

Accordingly, the Feynman integral satisfies the following GKZ-system in the splitting coordinates above

$$\begin{aligned} \{\vartheta_{1,1} + \vartheta_{1,4} + \vartheta_{1,8}\} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \{\vartheta_{2,2} + \vartheta_{2,4} + \vartheta_{2,8}\} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \{\vartheta_{3,3} + \vartheta_{3,4} + \vartheta_{3,8}\} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \{\vartheta_{4,4} + \vartheta_{4,5} + \vartheta_{4,6} + \vartheta_{4,7}\} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \vartheta_{1,1} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= \left(1 - \frac{D}{2}\right) A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \vartheta_{2,2} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= \left(1 - \frac{D}{2}\right) A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \vartheta_{3,3} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= \left(1 - \frac{D}{2}\right) A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \{\vartheta_{1,4} + \vartheta_{2,4} + \vartheta_{3,4} + \vartheta_{4,4}\} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= \left(\frac{D}{2} - 1\right) A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \vartheta_{4,5} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \vartheta_{4,6} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \vartheta_{4,7} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= -A_{2Vac}(m_1^2, m_2^2, m_3^2), \\ \{\vartheta_{1,8} + \vartheta_{2,8} + \vartheta_{3,8}\} A_{2Vac}(m_1^2, m_2^2, m_3^2) &= (D - 3) A_{2Vac}(m_1^2, m_2^2, m_3^2). \end{aligned} \quad (364)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 1 - \frac{D}{2} & 0 & 0 & \alpha_{1,4} & 0 & 0 & 0 & \alpha_{1,8} \\ 0 & 1 - \frac{D}{2} & 0 & \alpha_{2,4} & 0 & 0 & 0 & \alpha_{2,8} \\ 0 & 0 & 1 - \frac{D}{2} & \alpha_{3,4} & 0 & 0 & 0 & \alpha_{3,8} \\ 0 & 0 & 0 & 2 & -1 & -1 & -1 & 0 \end{pmatrix}, \quad (365)$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,4} + \alpha_{1,8} &= \frac{D}{2} - 2, \quad \alpha_{2,4} + \alpha_{2,8} = \frac{D}{2} - 2, \quad \alpha_{3,4} + \alpha_{3,8} = \frac{D}{2} - 2, \\ \alpha_{1,4} + \alpha_{2,4} + \alpha_{3,4} &= \frac{D}{2} - 3, \quad \alpha_{1,8} + \alpha_{2,8} + \alpha_{3,8} = D - 3. \end{aligned} \quad (366)$$

Corresponding to the matrix of local coordinates in Eq.(363), we have 12 choices on the




FIG. 9: The geometric descriptions of single orbits in Eq.(366) on the projective plane CP^4 , where single point 4 and triple point $\{5,6,7\}$ lie outside the hyperplane $\{1,2,3,8\}$.

matrix of integer lattice whose submatrix composed of the fourth and eighth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other elements are all zero. Note that $E_4^{(1)} = E_4^{(2)} - E_4^{(4)}$, two linear independent integer lattice matrices $E_4^{(2)}, E_4^{(4)}$ induce two independent PDEs as:

$$\begin{aligned} \frac{\partial^2 A_{2Vac}}{\partial z_{1,4} \partial z_{3,8}}(m_1^2, m_2^2, m_3^2) &= \frac{\partial^2 A_{2Vac}}{\partial z_{1,8} \partial z_{3,4}}(m_1^2, m_2^2, m_3^2), \\ \frac{\partial^2 A_{2Vac}}{\partial z_{2,4} \partial z_{3,8}}(m_1^2, m_2^2, m_3^2) &= \frac{\partial^2 A_{2Vac}}{\partial z_{2,8} \partial z_{3,4}}(m_1^2, m_2^2, m_3^2). \end{aligned} \quad (367)$$

When the exponents are given, the Feynman integral can be formally expressed as

$$A_{2Vac}(m_1^2, m_2^2, m_3^2) = \prod_{i,j} z_{i,j}^{\alpha_{i,j}} \varphi\left(\frac{m_1^2}{m_3^2}, \frac{m_2^2}{m_3^2}\right), \quad (368)$$

where $(i, j) \in \{(1, 4), (1, 8), (2, 4), (2, 8), (3, 4), (3, 8)\}$, and $\varphi(x_1, x_2)$ satisfies the PDEs in the combined coordinate $x_1 = z_{1,8} z_{3,4} / (z_{1,4} z_{3,8})$, $x_2 = z_{2,8} z_{3,4} / (z_{2,4} z_{3,8})$ as

$$\begin{aligned} &x_1 \left[\alpha_{1,4} \alpha_{3,8} + (1 - \alpha_{1,4} - \alpha_{3,8}) x_1 \frac{\partial}{\partial x_1} - \alpha_{1,4} x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_1^2 \frac{\partial^2}{\partial x_1^2} \right] \varphi(x_1, x_2) \\ &= \left[\alpha_{1,8} \alpha_{3,4} + (1 + \alpha_{1,8} + \alpha_{3,4}) x_1 \frac{\partial}{\partial x_1} + \alpha_{1,8} x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_1^2 \frac{\partial^2}{\partial x_1^2} \right] \varphi(x_1, x_2), \end{aligned}$$

$$\begin{aligned}
& x_2 \left[\alpha_{2,4} \alpha_{3,8} - \alpha_{2,4} x_1 \frac{\partial}{\partial x_1} + (1 - \alpha_{2,4} - \alpha_{3,8}) x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right] \varphi(x_1, x_2) \\
& = \left[\alpha_{2,8} \alpha_{3,4} + \alpha_{2,8} x_1 \frac{\partial}{\partial x_1} + (1 + \alpha_{2,8} + \alpha_{3,4}) x_2 \frac{\partial}{\partial x_2} + x_1 x_2 \frac{\partial^2}{\partial x_1 \partial x_2} + x_2^2 \frac{\partial^2}{\partial x_2^2} \right] \varphi(x_1, x_2). \quad (369)
\end{aligned}$$

As $\alpha_{1,8} = \alpha_{2,8} = 0$, the PDEs above are changed into the first type Appell PDEs. The holonomic rank of the ideal composed by the partial differential operators above is 3, which indicates three linear independent hypergeometric functions constituting the fundamental solution systems in neighborhoods of regular singularities. It is known that the fourth type Appell function cannot be embedded into Grassmannians [14]. Nevertheless, including the appropriate power factor, the Feynman integral presented through the fourth Appell functions [41, 42] satisfies the above PDEs. Those matrices of integer lattice permit 12 choices of the exponents. Corresponding to the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(2)}$, the choice of exponents is written as

$$\alpha_{1,4} = 1 - \frac{D}{2}, \quad \alpha_{1,8} = D - 3, \quad \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{2,8} = 0, \quad \alpha_{3,4} = \frac{D}{2} - 2, \quad \alpha_{3,8} = 0 \quad (370)$$

whose geometric representation is presented in Fig.9(1) with $(a, b) = (1, 8)$, $\{c, d\} = \{2, 3\}$. In a similar way the exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = 0, \quad \alpha_{1,8} = \frac{D}{2} - 2, \quad \alpha_{2,4} = -1, \quad \alpha_{2,8} = \frac{D}{2} - 1, \quad \alpha_{3,4} = \frac{D}{2} - 2, \quad \alpha_{3,8} = 0, \quad (371)$$

whose geometric representation is plotted in Fig.9(2) with $\{a, b, c\} = \{1, 2, 8\}$, $d = 3$. Corresponding to the integer lattice $n_1 E_4^{(2)} + n_2 E_4^{(4)}$, the exponents are

$$\alpha_{1,4} = 0, \quad \alpha_{1,8} = \frac{D}{2} - 2, \quad \alpha_{2,4} = 0, \quad \alpha_{2,8} = \frac{D}{2} - 2, \quad \alpha_{3,4} = \frac{D}{2} - 3, \quad \alpha_{3,8} = 1, \quad (372)$$

whose geometric representation is presented in Fig.9(3) with $a = 3$, $(b, c, d) = (1, 2, 8)$. The hypergeometric functions are evidently written as

$$\begin{aligned}
\psi_{\{1,2,3,5\}}^{(1)}(m_1^2, m_2^2, m_3^2) & \sim (m_1^2)^{D-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2!} \\
& \quad \times \frac{\Gamma(3 - D + n_1 + n_2)}{\Gamma(2 - \frac{D}{2} + n_1 + n_2)} \left(\frac{m_2^2}{m_1^2}\right)^{n_1} \left(\frac{m_3^2}{m_1^2}\right)^{n_2}, \\
\psi_{\{1,2,3,5\}}^{(2)}(m_1^2, m_2^2, m_3^2) & \sim (m_1^2)^{D/2-2} (m_2^2)^{D/2-1} \\
& \quad \times \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2) \Gamma(1 + n_1)}{n_2! \Gamma(1 + n_1 + n_2) \Gamma(\frac{D}{2} + n_1)}
\end{aligned}$$

$$\begin{aligned}
& \times \Gamma\left(2 - \frac{D}{2} + n_2\right) \left(\frac{m_2^2}{m_1^2}\right)^{n_1} \left(\frac{m_3^2}{m_1^2}\right)^{n_2}, \\
\psi_{\{1,2,3,5\}}^{(3)}(m_1^2, m_2^2, m_3^2) & \sim (m_1^2 m_2^2)^{D/2-2} m_3^2 \sum_{n_1, n_2}^{\infty} \frac{\Gamma\left(2 - \frac{D}{2} + n_1\right) \Gamma\left(2 - \frac{D}{2} + n_2\right)}{n_1! n_2!} \\
& \times \frac{\Gamma\left(3 - \frac{D}{2} + n_1 + n_2\right)}{\Gamma(2 + n_1 + n_2)} \left(\frac{m_3^2}{m_1^2}\right)^{n_1} \left(\frac{m_3^2}{m_2^2}\right)^{n_2}, \tag{373}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. In other words, three linear independent hypergeometric functions constitute a fundamental solution system of the GKZ-system in the nonempty proper subset of the whole parameter space. Because non-positive integers are the first order zeros of $\Gamma^{-1}(z)$, the exponents of Eq.(371) induce the following hypergeometric function

$$\begin{aligned}
\psi_{\{1,2,3,5\}}^{(2)'}(m_1^2, m_2^2, m_3^2) & \sim \frac{\sin^3 \frac{\pi D}{2}}{\pi^3} (m_1^2)^{D/2-2} (m_2^2)^{D/2-1} \\
& \times \sum_{n_1=0}^{\infty} \sum_{n_2=1+n_1}^{\infty} \frac{\Gamma\left(1 - \frac{D}{2} - n_1 + n_2\right)}{n_1! n_2! \Gamma(-n_1 + n_2)} \\
& \times \Gamma\left(2 - \frac{D}{2} + n_1\right) \Gamma\left(2 - \frac{D}{2} + n_2\right) \left(\frac{m_2^2}{m_1^2}\right)^{n_1} \left(\frac{m_3^2}{m_2^2}\right)^{n_2} \\
& + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon \sin^2 \frac{\pi D}{2}}{\pi^3} (m_1^2)^{D/2-2} (m_2^2)^{D/2-1} \\
& \times \sum_{n_2=0}^{\infty} \sum_{n_1=n_2}^{\infty} \frac{\Gamma(1 + n_1 - n_2)}{n_1! n_2! \Gamma\left(\frac{D}{2} + n_1 - n_2\right)} \\
& \times \Gamma\left(2 - \frac{D}{2} + n_1\right) \Gamma\left(2 - \frac{D}{2} + n_2\right) \left(\frac{m_2^2}{m_1^2}\right)^{n_1} \left(\frac{m_3^2}{m_2^2}\right)^{n_2} \\
& \sim \frac{\sin^3 \frac{\pi D}{2}}{\pi^3} \psi_{\{1,2,3,5\}}^{(3)}(m_1^2, m_2^2, m_3^2) \\
& + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon \sin^2 \frac{\pi D}{2}}{\pi^3} \psi_{\{1,2,3,5\}}^{(2)}(m_1^2, m_2^2, m_3^2). \tag{374}
\end{aligned}$$

The exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(2)}$ are

$$\alpha_{1,4} = \frac{D}{2} - 3, \quad \alpha_{1,8} = 1, \quad \alpha_{2,4} = 0, \quad \alpha_{2,8} = \frac{D}{2} - 2, \quad \alpha_{3,4} = 0, \quad \alpha_{3,8} = \frac{D}{2} - 2, \tag{375}$$

whose geometric representation is plotted in Fig.9(3) with $a = 1$, $\{b, c, d\} = \{2, 3, 8\}$. Similarly the exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are given by

$$\alpha_{1,4} = \frac{D}{2} - 2, \quad \alpha_{1,8} = 0, \quad \alpha_{2,4} = -1, \quad \alpha_{2,8} = \frac{D}{2} - 1, \quad \alpha_{3,4} = 0, \quad \alpha_{3,8} = \frac{D}{2} - 2, \tag{376}$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{2, 3, 8\}$, $d = 1$.

Corresponding to the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(4)}$, the exponents are

$$\alpha_{1,4} = \frac{D}{2} - 2, \alpha_{1,8} = 0, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{2,8} = 0, \alpha_{3,4} = 1 - \frac{D}{2}, \alpha_{3,8} = D - 3 \quad (377)$$

whose geometric representation is depicted in Fig.9(1) with $(a, b) = (3, 8)$, $\{c, d\} = \{1, 2\}$.

Accordingly, the hypergeometric functions are

$$\begin{aligned} \psi_{\{1,2,3,5\}}^{(4)}(m_1^2, m_2^2, m_3^2) &\sim m_1^2 (m_2^2 m_3^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2!} \\ &\quad \times \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{\Gamma(2 + n_1 + n_2)} \left(\frac{m_1^2}{m_2^2}\right)^{n_1} \left(\frac{m_1^2}{m_3^2}\right)^{n_2}, \\ \psi_{\{1,2,3,5\}}^{(5)}(m_1^2, m_2^2, m_3^2) &\sim (m_2^2)^{D/2-1} (m_3^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{n_1! \Gamma(1 + n_1 + n_2)} \\ &\quad \times \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(1 + n_2)}{\Gamma(\frac{D}{2} + n_2)} \left(\frac{m_1^2}{m_3^2}\right)^{n_1} \left(\frac{m_2^2}{m_3^2}\right)^{n_2}, \\ \psi_{\{1,2,3,5\}}^{(6)}(m_1^2, m_2^2, m_3^2) &\sim (m_3^2)^{D-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2!} \\ &\quad \times \frac{\Gamma(3 - D + n_1 + n_2)}{\Gamma(2 - \frac{D}{2} + n_1 + n_2)} \left(\frac{m_1^2}{m_3^2}\right)^{n_1} \left(\frac{m_2^2}{m_3^2}\right)^{n_2}, \end{aligned} \quad (378)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space.

Similarly the exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(2)}$ are

$$\alpha_{1,4} = -1, \alpha_{1,8} = \frac{D}{2} - 1, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{2,8} = 0, \alpha_{3,4} = 0, \alpha_{3,8} = \frac{D}{2} - 2, \quad (379)$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{1, 3, 8\}$, $d = 2$. The

exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are given as

$$\alpha_{1,4} = 0, \alpha_{1,8} = \frac{D}{2} - 2, \alpha_{2,4} = \frac{D}{2} - 3, \alpha_{2,8} = 1, \alpha_{3,4} = 0, \alpha_{3,8} = \frac{D}{2} - 2, \quad (380)$$

whose geometric representation is drawn in Fig.9(3) with $a = 2$, $\{b, c, d\} = \{1, 3, 8\}$. Mean-

while the exponents of integer lattice $n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = 0, \alpha_{1,8} = \frac{D}{2} - 2, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{2,8} = 0, \alpha_{3,4} = -1, \alpha_{3,8} = \frac{D}{2} - 1, \quad (381)$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{1, 3, 8\}$, $d = 2$. The

corresponding hypergeometric functions are

$$\psi_{\{1,2,3,5\}}^{(7)}(m_1^2, m_2^2, m_3^2) \sim (m_1^2)^{D/2-1} (m_3^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{n_1! \Gamma(1 + n_1 + n_2)}$$

$$\begin{aligned}
& \times \frac{\Gamma(2 - \frac{D}{2} + n_1)\Gamma(1 + n_2)}{\Gamma(\frac{D}{2} + n_2)} \left(\frac{m_2^2}{m_3^2}\right)^{n_1} \left(\frac{m_1^2}{m_3^2}\right)^{n_2}, \\
\psi_{\{1,2,3,5\}}^{(8)}(m_1^2, m_2^2, m_3^2) & \sim (m_1^2 m_3^2)^{D/2-2} m_2^2 \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1)\Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2!} \\
& \times \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{\Gamma(2 + n_1 + n_2)} \left(\frac{m_2^2}{m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_3^2}\right)^{n_2}, \\
\psi_{\{1,2,3,5\}}^{(9)}(m_1^2, m_2^2, m_3^2) & \sim (m_1^2)^{D/2-2} (m_3^2)^{D/2-1} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{n_2! \Gamma(1 + n_1 + n_2)} \\
& \times \frac{\Gamma(1 + n_1)\Gamma(2 - \frac{D}{2} + n_2)}{\Gamma(\frac{D}{2} + n_1)} \left(\frac{m_3^2}{m_1^2}\right)^{n_1} \left(\frac{m_2^2}{m_1^2}\right)^{n_2}, \tag{382}
\end{aligned}$$

where the intersection of their convergent regions is empty. However, the sets of the hypergeometric functions $\{\psi_{\{1,2,3,5\}}^{(1)}, \psi_{\{1,2,3,5\}}^{(8)}, \psi_{\{1,2,3,5\}}^{(9)}\}$ and $\{\psi_{\{1,2,3,5\}}^{(6)}, \psi_{\{1,2,3,5\}}^{(7)}, \psi_{\{1,2,3,5\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

The exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are

$$\alpha_{1,4} = -1, \alpha_{1,8} = \frac{D}{2} - 1, \alpha_{2,4} = 0, \alpha_{2,8} = \frac{D}{2} - 2, \alpha_{3,4} = \frac{D}{2} - 2, \alpha_{3,8} = 0, \tag{383}$$

whose geometric representation is plotted in Fig.9(2) with $\{a, b, c\} = \{1, 2, 8\}$, $d = 3$. The exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,4} = \frac{D}{2} - 2, \alpha_{1,8} = 0, \alpha_{2,4} = 1 - \frac{D}{2}, \alpha_{2,8} = D - 3, \alpha_{3,4} = \frac{D}{2} - 2, \alpha_{3,8} = 0 \tag{384}$$

where the geometric representation is presented in Fig.9(1) with $(a, b) = (2, 8)$, $\{c, d\} = \{1, 3\}$. Corresponding to the integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(4)}$, the exponents are

$$\alpha_{1,4} = \frac{D}{2} - 2, \alpha_{1,8} = 0, \alpha_{2,4} = 0, \alpha_{2,8} = \frac{D}{2} - 2, \alpha_{3,4} = -1, \alpha_{3,8} = \frac{D}{2} - 1, \tag{385}$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{2, 3, 8\}$, $d = 1$. Accordingly the hypergeometric functions are evidently written as

$$\begin{aligned}
\psi_{\{1,2,3,5\}}^{(10)}(m_1^2, m_2^2, m_3^2) & \sim (m_1^2)^{D/2-1} (m_2^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 + n_2)}{n_2! \Gamma(1 + n_1 + n_2)} \\
& \times \frac{\Gamma(1 + n_1)\Gamma(2 - \frac{D}{2} + n_2)}{\Gamma(\frac{D}{2} + n_1)} \left(\frac{m_1^2}{m_2^2}\right)^{n_1} \left(\frac{m_3^2}{m_2^2}\right)^{n_2}, \\
\psi_{\{1,2,3,5\}}^{(11)}(m_1^2, m_2^2, m_3^2) & \sim (m_2^2)^{D-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1)\Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2!}
\end{aligned}$$

$$\begin{aligned}
& \times \frac{\Gamma(3-D+n_1+n_2)}{\Gamma(2-\frac{D}{2}+n_1+n_2)} \left(\frac{m_1^2}{m_2^2}\right)^{n_1} \left(\frac{m_3^2}{m_2^2}\right)^{n_2}, \\
\psi_{\{1,2,3,5\}}^{(12)}(m_1^2, m_2^2, m_3^2) & \sim (m_2^2)^{D/2-2} (m_3^2)^{D/2-1} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2-\frac{D}{2}+n_1+n_2)}{n_1! \Gamma(1+n_1+n_2)} \\
& \times \frac{\Gamma(2-\frac{D}{2}+n_1) \Gamma(1+n_2)}{\Gamma(\frac{D}{2}+n_2)} \left(\frac{m_1^2}{m_2^2}\right)^{n_1} \left(\frac{m_3^2}{m_2^2}\right)^{n_2}, \quad (386)
\end{aligned}$$

where the intersection of their convergent regions is empty. Nevertheless, the sets of the hypergeometric functions $\{\psi_{\{1,2,3,5\}}^{(3)}, \psi_{\{1,2,3,5\}}^{(10)}, \psi_{\{1,2,3,5\}}^{(11)}\}$ and $\{\psi_{\{1,2,3,5\}}^{(4)}, \psi_{\{1,2,3,5\}}^{(11)}, \psi_{\{1,2,3,5\}}^{(12)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{\{1,2,5,8\}})^{(2V)} = -m_3^2$, where the matrix $A_{\{1,2,5,8\}}^{(2V)}$ denotes the submatrix of $A^{(2V)}$ composed of the first, second, fifth and eighth columns, one finds

$$\left(A_{\{1,2,5,8\}}^{(2V)}\right)^{-1} \cdot A^{(2V)} = \begin{pmatrix} 1 & 0 & -\frac{m_1^2}{m_2^2} & -1 + \frac{m_1^2}{m_2^2} & 0 & 0 & 0 & 0 \\ 0 & 1 & -\frac{m_2^2}{m_3^2} & -1 + \frac{m_2^2}{m_3^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & 0 & \frac{1}{m_3^2} & -\frac{1}{m_3^2} & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (387)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 1 - \frac{D}{2} & 0 & \alpha_{1,3} & \alpha_{1,4} & 0 & 0 & 0 & 0 \\ 0 & 1 - \frac{D}{2} & \alpha_{2,3} & \alpha_{2,4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & -1 & -1 & -1 & 0 \\ 0 & 0 & \alpha_{4,3} & \alpha_{4,4} & 0 & 0 & 0 & D-3 \end{pmatrix}, \quad (388)$$

where the matrix elements satisfy the relations

$$\begin{aligned}
\alpha_{1,3} + \alpha_{1,4} &= \frac{D}{2} - 2, \quad \alpha_{2,3} + \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{4,3} + \alpha_{4,4} = 2 - D, \\
\alpha_{1,3} + \alpha_{2,3} + \alpha_{4,3} &= 1 - \frac{D}{2}, \quad \alpha_{1,4} + \alpha_{2,4} + \alpha_{4,4} = \frac{D}{2} - 3. \quad (389)
\end{aligned}$$

Corresponding to the matrix of splitting local coordinates in Eq.(387), one finds 12 choices on the matrix of integer lattice whose submatrix composed of the third and fourth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 3), (1, 5), (3, 5)\}$, and other elements are all zero.

Corresponding to the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(3)}$, the exponents are written as

$$\alpha_{1,3} = 1, \quad \alpha_{1,4} = \frac{D}{2} - 3, \quad \alpha_{2,3} = \frac{D}{2} - 2, \quad \alpha_{2,4} = 0, \quad \alpha_{4,3} = 2 - D, \quad \alpha_{4,4} = 0, \quad (390)$$

whose geometric representation is drawn in Fig.9(3) with $a = 1$, $\{b, c, d\} = \{2, 3, 8\}$. In a similar way the exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,4} = \frac{D}{2} - 2, \alpha_{2,3} = \frac{D}{2} - 1, \alpha_{2,4} = -1, \alpha_{4,3} = 2 - D, \alpha_{4,4} = 0, \quad (391)$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{2, 3, 8\}$, $d = 1$. The exponents of the integer lattice $n_1 E_4^{(3)} + n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,4} = \frac{D}{2} - 2, \alpha_{2,3} = 0, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{4,3} = 1 - \frac{D}{2}, \alpha_{4,4} = 1 - \frac{D}{2} \quad (392)$$

whose geometric representation is plotted in Fig.9(1) with $(a, b) = (3, 8)$, $\{c, d\} = \{1, 2\}$. The corresponding hypergeometric functions are evidently written as

$$\begin{aligned} \psi_{\{1,2,5,8\}}^{(1)}(m_1^2, m_2^2, m_3^2) &\sim \frac{m_1^2 m_3^2 (m_2^2)^{D/2-2}}{(m_1^2 - m_3^2)^{3-D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{n_1! n_2! \Gamma(2 + n_1 + n_2)} \\ &\quad \times \Gamma(2 - \frac{D}{2} + n_1) \Gamma(D - 2 + n_2) \\ &\quad \times \left(\frac{m_1^2 (m_2^2 - m_3^2)}{m_2^2 (m_1^2 - m_3^2)} \right)^{n_1} \left(\frac{m_1^2}{m_1^2 - m_3^2} \right)^{n_2}, \\ \psi_{\{1,2,5,8\}}^{(2)}(m_1^2, m_2^2, m_3^2) &\sim \frac{(m_2^2)^{D/2-1} m_3^2 (m_1^2 - m_3^2)^{D/2-2}}{m_2^2 - m_3^2} \\ &\quad \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D - 2 + n_1 + n_2) \Gamma(2 - \frac{D}{2} + n_1)}{n_1! \Gamma(1 + n_1 + n_2)} \\ &\quad \times \frac{\Gamma(1 + n_2)}{\Gamma(\frac{D}{2} + n_2)} \left(\frac{m_1^2}{m_1^2 - m_3^2} \right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_3^2} \right)^{n_2}, \\ \psi_{\{1,2,5,8\}}^{(3)}(m_1^2, m_2^2, m_3^2) &\sim \left((m_1^2 - m_3^2) (m_2^2 - m_3^2) \right)^{D/2-2} m_3^2 \\ &\quad \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{n_1! n_2! \Gamma(2 + n_1 + n_2)} \\ &\quad \times \Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2) \\ &\quad \times \left(\frac{m_1^2}{m_1^2 - m_3^2} \right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_3^2} \right)^{n_2}, \end{aligned} \quad (393)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space, and three linear independent hypergeometric functions constitute a fundamental solution system of the GKZ-system in the nonempty intersection.

The exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(3)}$ are

$$\alpha_{1,3} = 1 - \frac{D}{2}, \alpha_{1,4} = D - 3, \alpha_{2,3} = 0, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{4,3} = 0, \alpha_{4,4} = 2 - D, \quad (394)$$

whose geometric representation is presented in Fig.9(1) with $(a, b) = (1, 3)$, $\{c, d\} = \{2, 8\}$.

Similarly the exponents corresponding to the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(5)}$ are given by

$$\alpha_{1,3} = \frac{D}{2} - 2, \alpha_{1,4} = 0, \alpha_{2,3} = 3 - D, \alpha_{2,4} = \frac{3D}{2} - 5, \alpha_{4,3} = 0, \alpha_{4,4} = 2 - D \quad (395)$$

whose geometric representation is shown in Fig.9(2) with $\{a, b, c\} = \{1, 2, 3\}$, $d = 8$. Finally the exponents of the integer lattice $-n_1 E_4^{(3)} - n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = \frac{D}{2} - 2, \alpha_{1,4} = 0, \alpha_{2,3} = \frac{D}{2} - 2, \alpha_{2,4} = 0, \alpha_{4,3} = 5 - \frac{3D}{2}, \alpha_{4,4} = \frac{D}{2} - 3 \quad (396)$$

whose geometric representation is presented in Fig.9(3) with $a = 8$, $\{b, c, d\} = \{1, 2, 3\}$.

Accordingly the hypergeometric functions are summarized as

$$\begin{aligned} \psi_{\{1,2,5,8\}}^{(4)}(m_1^2, m_2^2, m_3^2) &\sim \frac{m_3^2(m_2^2 m_3^2)^{D/2-2}}{(m_1^2)^{D/2-1}(m_1^2 - m_3^2)^{3-D}} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2) \Gamma(2 - \frac{D}{2} + n_1) \Gamma(D - 2 + n_2)}{n_1! n_2! \Gamma(D - 2 + n_1 + n_2)} \\ &\times \left(\frac{m_2^2(m_1^2 - m_3^2)}{m_1^2(m_2^2 - m_3^2)} \right)^{n_1} \left(\frac{m_1^2 - m_3^2}{m_1^2} \right)^{n_2}, \\ \psi_{\{1,2,5,8\}}^{(5)}(m_1^2, m_2^2, m_3^2) &\sim \frac{(m_1^2)^{D/2-2} m_3^2}{(m_2^2)^{D-3} (m_2^2 - m_3^2)^{5-3D/2}} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(5 - \frac{3D}{2} - 1 + n_1 - n_2) \Gamma(2 - \frac{D}{2} + n_1) \Gamma(D - 2 + n_2)}{n_1! n_2! \Gamma(4 - D + n_1 - n_2)} \\ &\times \left(\frac{m_2^2(m_1^2 - m_3^2)}{m_1^2(m_2^2 - m_3^2)} \right)^{n_1} \left(\frac{m_2^2 - m_3^2}{m_2^2} \right)^{n_2}, \\ \psi_{\{1,2,5,8\}}^{(6)}(m_1^2, m_2^2, m_3^2) &\sim m_3^2 (m_1^2 m_2^2)^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(2 - \frac{D}{2} + n_2)}{n_1! n_2!} \\ &\times \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{\Gamma(6 - \frac{3D}{2} + n_1 + n_2)} \left(\frac{m_1^2 - m_3^2}{m_1^2} \right)^{n_1} \left(\frac{m_2^2 - m_3^2}{m_2^2} \right)^{n_2}, \end{aligned} \quad (397)$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The hypergeometric functions constitute a fundamental solution system of the GKZ-system in a nonempty proper subset of the whole parameter space.

Similarly the exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(3)}$ are written as

$$\alpha_{1,3} = 3 - D, \alpha_{1,4} = \frac{3D}{2} - 5, \alpha_{2,3} = \frac{D}{2} - 2, \alpha_{2,4} = 0, \alpha_{4,3} = 0, \alpha_{4,4} = 2 - D \quad (398)$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{1, 2, 3\}$, $d = 8$. The exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,4} = \frac{D}{2} - 2, \alpha_{2,3} = 1 - \frac{D}{2}, \alpha_{2,4} = D - 3, \alpha_{4,3} = 0, \alpha_{4,4} = 2 - D \quad (399)$$

whose geometric representation is drawn in Fig.9(1) with $(a, b) = (2, 3)$, $\{c, d\} = \{1, 8\}$. Accordingly the exponents of the integer lattice $n_1 E_4^{(3)} - n_2 E_4^{(5)}$ are given by

$$\alpha_{1,3} = 0, \alpha_{1,4} = \frac{D}{2} - 2, \alpha_{2,3} = \frac{D}{2} - 2, \alpha_{2,4} = 0, \alpha_{4,3} = 3 - D, \alpha_{4,4} = -1, \quad (400)$$

where the geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{2, 3, 8\}$, $d = 1$.

The corresponding hypergeometric functions are presented as

$$\begin{aligned} \psi_{\{1,2,5,8\}}^{(7)}(m_1^2, m_2^2, m_3^2) &\sim \frac{(m_1^2)^{3-D} (m_2^2)^{D/2-2} m_3^2}{(m_1^2 - m_3^2)^{5-3D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(D - 2 + n_2)}{n_1! n_2!} \\ &\times \frac{\Gamma(5 - \frac{3D}{2} + n_1 - n_2)}{\Gamma(4 - D + n_1 - n_2)} \left(\frac{m_1^2 (m_2^2 - m_3^2)}{m_2^2 (m_1^2 - m_3^2)} \right)^{n_1} \left(\frac{m_1^2 - m_3^2}{m_1^2} \right)^{n_2}, \\ \psi_{\{1,2,5,8\}}^{(8)}(m_1^2, m_2^2, m_3^2) &\sim \frac{m_3^2 (m_1^2 - m_3^2)^{D/2-2}}{(m_2^2)^{D/2-1} (m_2^2 - m_3^2)^{3-D}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1) \Gamma(D - 2 + n_2)}{n_1! n_2!} \\ &\times \frac{\Gamma(\frac{D}{2} - 1 + n_1 + n_2)}{\Gamma(D - 2 + n_1 + n_2)} \left(\frac{m_1^2 (m_2^2 - m_3^2)}{m_2^2 (m_1^2 - m_3^2)} \right)^{n_1} \left(\frac{m_2^2 - m_3^2}{m_2^2} \right)^{n_2}, \\ \psi_{\{1,2,5,8\}}^{(9)}(m_1^2, m_2^2, m_3^2) &\sim \frac{m_1^2 m_3^2 (m_2^2)^{D/2-2}}{(m_1^2 - m_3^2)^{3-D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{n_1! n_2! \Gamma(1 + n_1 + n_2)} \\ &\times \Gamma(D - 2 + n_1) \Gamma(2 - \frac{D}{2} + n_2) \\ &\times \left(\frac{m_1^2}{m_1^2 - m_3^2} \right)^{n_1} \left(\frac{m_1^2 (m_2^2 - m_3^2)}{m_2^2 (m_1^2 - m_2^2)} \right)^{n_2}, \end{aligned} \quad (401)$$

where the intersection of their convergent regions is empty. However, the sets of the hypergeometric functions $\{\psi_{\{1,2,5,8\}}^{(1)}, \psi_{\{1,2,5,8\}}^{(8)}, \psi_{\{1,2,5,8\}}^{(9)}\}$ and $\{\psi_{\{1,2,5,8\}}^{(6)}, \psi_{\{1,2,5,8\}}^{(7)}, \psi_{\{1,2,5,8\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

The exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(3)}$ are

$$\alpha_{1,3} = \frac{D}{2} - 1, \alpha_{1,4} = -1, \alpha_{2,3} = 0, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{4,3} = 2 - D, \alpha_{4,4} = 0, \quad (402)$$

whose geometric representation is depicted in Fig.9(2) with $\{a, b, c\} = \{1, 2, 8\}$, $d = 3$. The corresponding exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(5)}$ are given as

$$\alpha_{1,3} = \frac{D}{2} - 2, \alpha_{1,4} = 0, \alpha_{2,3} = 1, \alpha_{2,4} = \frac{D}{2} - 3, \alpha_{4,3} = 2 - D, \alpha_{4,4} = 0, \quad (403)$$

whose geometric representation is shown in Fig.9(3) with $a = 2$, $\{b, c, d\} = \{1, 3, 8\}$. Finally, the exponents of the integer lattice $-n_1 E_4^{(3)} + n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = \frac{D}{2} - 2, \alpha_{1,4} = 0, \alpha_{2,3} = 0, \alpha_{2,4} = \frac{D}{2} - 2, \alpha_{4,3} = 3 - D, \alpha_{4,4} = -1, \quad (404)$$

whose geometric representation is presented in Fig.9(2) with $\{a, b, c\} = \{1, 3, 8\}$, $d = 2$.

Three hypergeometric functions are evidently written as

$$\begin{aligned}
\psi_{\{1,2,5,8\}}^{(10)}(m_1^2, m_2^2, m_3^2) &\sim \frac{(m_1^2)^{D/2-1}(m_2^2)^{D/2-2}m_3^2}{m_1^2 - m_3^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(D-2+n_1+n_2)}{\Gamma(1+n_1+n_2)} \\
&\quad \times \frac{\Gamma(2-\frac{D}{2}+n_1)\Gamma(1+n_2)}{n_1!\Gamma(\frac{D}{2}+n_2)} \left(\frac{m_2^2}{m_2^2 - m_3^2}\right)^{n_1} \left(\frac{m_1^2}{m_1^2 - m_3^2}\right)^{n_2}, \\
\psi_{\{1,2,5,8\}}^{(11)}(m_1^2, m_2^2, m_3^2) &\sim \frac{m_2^2 m_3^2 (m_1^2)^{D/2-2}}{(m_2^2 - m_3^2)^{3-D/2}} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3-\frac{D}{2}+n_1+n_2)}{\Gamma(2+n_1+n_2)} \\
&\quad \times \frac{\Gamma(2-\frac{D}{2}+n_1)\Gamma(D-2+n_2)}{n_1!n_2!} \\
&\quad \times \left(\frac{m_2^2(m_1^2 - m_3^2)}{m_1^2(m_2^2 - m_3^2)}\right)^{n_1} \left(\frac{m_2^2}{m_2^2 - m_3^2}\right)^{n_2}, \\
\psi_{\{1,2,5,8\}}^{(12)}(m_1^2, m_2^2, m_3^2) &\sim m_3^2 (m_1^2(m_2^2 - m_3^2))^{D/2-2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2-\frac{D}{2}+n_1+n_2)}{\Gamma(1+n_1+n_2)} \\
&\quad \times \frac{\Gamma(1+n_1)\Gamma(2-\frac{D}{2}+n_2)}{n_2!\Gamma(4-D+n_1)} \\
&\quad \times \left(\frac{m_1^2 - m_3^2}{m_1^2}\right)^{n_1} \left(\frac{m_2^2(m_1^2 - m_3^2)}{m_1^2(m_2^2 - m_3^2)}\right)^{n_2}, \tag{405}
\end{aligned}$$

where the intersection of their convergent regions is empty. Nevertheless, the sets of the hypergeometric functions $\{\psi_{\{1,2,5,8\}}^{(3)}, \psi_{\{1,2,5,8\}}^{(10)}, \psi_{\{1,2,5,8\}}^{(11)}\}$ and $\{\psi_{\{1,2,5,8\}}^{(4)}, \psi_{\{1,2,5,8\}}^{(11)}, \psi_{\{1,2,5,8\}}^{(12)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{\{1,3,5,8\}})^{(2V)} = m_2^2$, where the matrix $A_{\{1,3,5,8\}}^{(2V)}$ denotes the submatrix of $A^{(2V)}$ composed of the first, third, fifth and eighth columns, one finds

$$\left(A_{\{1,3,5,8\}}^{(2V)}\right)^{-1} \cdot A^{(2V)} = \begin{pmatrix} 1 & -\frac{m_1^2}{m_2^2} & 0 & -1 + \frac{m_1^2}{m_2^2} & 0 & 0 & 0 & 0 \\ 0 & -\frac{m_3^2}{m_2^2} & 1 & -1 + \frac{m_3^2}{m_2^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 0 & \frac{1}{m_2^2} & 0 & -\frac{1}{m_2^2} & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{406}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 1 - \frac{D}{2} & \alpha_{1,2} & 0 & \alpha_{1,4} & 0 & 0 & 0 & 0 \\ 0 & \alpha_{2,2} & 1 - \frac{D}{2} & \alpha_{2,4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & -1 & -1 & -1 & 0 \\ 0 & \alpha_{4,2} & 0 & \alpha_{4,4} & 0 & 0 & 0 & D-3 \end{pmatrix}, \tag{407}$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,2} + \alpha_{1,4} &= \frac{D}{2} - 2, \quad \alpha_{2,2} + \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{4,2} + \alpha_{4,4} = 2 - D, \\ \alpha_{1,2} + \alpha_{2,2} + \alpha_{4,2} &= 1 - \frac{D}{2}, \quad \alpha_{1,4} + \alpha_{2,4} + \alpha_{4,4} = \frac{D}{2} - 3.\end{aligned}\quad (408)$$

Corresponding to the matrix of splitting local coordinates in Eq.(406), we have 12 choices on the matrix of integer lattice whose submatrix composed of the second and fourth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 3), (1, 5), (3, 5)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents whose geometric descriptions are obtained from that of $\psi_{\{1,2,5,8\}}^{(i)}$ by the permutation $(\overline{23})$. Correspondingly the hypergeometric functions are

$$\psi_{\{1,3,5,8\}}^{(i)}(m_1^2, m_2^2, m_3^2) = \psi_{\{1,2,5,8\}}^{(i)}(m_1^2, m_3^2, m_2^2) \quad (409)$$

Similarly $\det(A_{\{2,3,5,8\}})^{(2V)} = -m_1^2$, where the matrix $A_{\{2,3,5,8\}}^{(2V)}$ denotes the submatrix of $A^{(2V)}$ composed of the second, third, fifth and eighth columns, one finds

$$\left(A_{\{2,3,5,8\}}^{(2V)}\right)^{-1} \cdot A^{(2V)} = \begin{pmatrix} -\frac{m_2^2}{m_1^2} & 1 & 0 & -1 + \frac{m_2^2}{m_1^2} & 0 & 0 & 0 & 0 \\ -\frac{m_3^2}{m_1^2} & 0 & 1 & -1 + \frac{m_3^2}{m_1^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ \frac{1}{m_1^2} & 0 & 0 & -\frac{1}{m_1^2} & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (410)$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 1 - \frac{D}{2} & 0 & \alpha_{1,4} & 0 & 0 & 0 & 0 \\ \alpha_{2,1} & 0 & 1 - \frac{D}{2} & \alpha_{2,4} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & -1 & -1 & -1 & 0 \\ \alpha_{4,1} & 0 & 0 & \alpha_{4,4} & 0 & 0 & 0 & D - 3 \end{pmatrix}, \quad (411)$$

where the matrix elements satisfy the relations

$$\begin{aligned}\alpha_{1,1} + \alpha_{1,4} &= \frac{D}{2} - 2, \quad \alpha_{2,1} + \alpha_{2,4} = \frac{D}{2} - 2, \quad \alpha_{4,1} + \alpha_{4,4} = 2 - D, \\ \alpha_{1,1} + \alpha_{2,1} + \alpha_{4,1} &= 1 - \frac{D}{2}, \quad \alpha_{1,4} + \alpha_{2,4} + \alpha_{4,4} = \frac{D}{2} - 3.\end{aligned}\quad (412)$$

Correspondingly there are 12 choices on the matrix of integer lattice whose submatrix composed of the first and fourth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, (i, j)

$\in \{(1, 3), (1, 5), (3, 5)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents whose geometric descriptions are obtained from that of $\psi_{\{1,2,5,8\}}^{(i)}$ by the permutation $(\widehat{13})$. Correspondingly the hypergeometric functions are

$$\psi_{\{2,3,5,8\}}^{(i)}(m_1^2, m_2^2, m_3^2) = \psi_{\{1,2,5,8\}}^{(i)}(m_3^2, m_2^2, m_1^2). \quad (413)$$

In view of combinatorial geometry [60], the points with the same splitting local coordinates in the projective space can be taken as one point whose exponent is the sum of the exponents of the coincident points. Nevertheless, in the matrix of splitting local coordinates specifying a column for each homogeneous factor in the integrand allows us to apply more combinatorial information to efficiently obtain fundamental solution systems of neighborhoods of different regular singularities.

V. PARAMETRIC REPRESENTATIONS OF THE 2-LOOP SELF-ENERGIES

To embed the Feynman integrals into the subvarieties of Grassmannians through the Feynman parametric representation is not practical because sizes of corresponding Grassmannians are too large to obtain the hypergeometric functions efficiently. In order to construct hypergeometric functions efficiently, we adopt the α -parametric representation [79] of the Feynman integrals for multiloop diagrams. The diagrams of the 2-loop self-energies are drawn in Fig.10, and the Feynman integral of the 2-loop self energy diagram with 3 propagators can be expressed as

$$\begin{aligned} A_{2L3}(p^2, m_1^2, m_2^2, m_3^2) &= -\frac{i^{6-2D}\Gamma(3-D)(\Lambda_{\text{RE}}^2)^{3-D}}{(4\pi)^D} \int \omega_4(t) (t_1 t_2 t_3)^{1-D/2} \\ &\quad \times t_4^{D/2-1} \delta(t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4) \\ &\quad \times [t_1 m_1^2 + t_2 m_2^2 + t_3 m_3^2 + t_4 p^2]^{D-3}. \end{aligned} \quad (414)$$

The integral can be embedded in the subvariety of the Grassmannian $G_{4,8}$ where the first row corresponds to the integration variable t_1 , the second row corresponds to the integration variable t_2 , the third row corresponds to the integration variable t_3 , and the fourth row corresponds to the integration variable t_4 , respectively. Meanwhile the first column represents the power function $t_1^{1-D/2}$, the second column represents the power function $t_2^{1-D/2}$, the




FIG. 10: The skeleton diagrams of two-loop self-energies.

third column represents the power function $t_3^{1-D/2}$, the fourth column represents the power function $t_4^{D/2-1}$, the eighth column represents the function $(t_1 m_1^2 + t_2 m_2^2 + t_3 m_3^2 + t_4 p^2)^{D-3}$. In order to embed the homogeneous function $\delta(t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4)$ as the fifth, sixth, and seventh columns of the Grassmannian $G_{4,8}$, we rewrite

$$\begin{aligned}
& t_1 t_2 t_3 + t_1 t_2 t_4 + t_1 t_3 t_4 + t_2 t_3 t_4 \\
&= z_{1, \hat{\tau}_1(5)} z_{2, \hat{\tau}_1(6)} z_{3, \hat{\tau}_1(7)} t_1 t_2 t_3 + z_{1, \hat{\tau}_2(5)} z_{2, \hat{\tau}_2(6)} z_{4, \hat{\tau}_2(7)} t_1 t_2 t_4 \\
&\quad + z_{1, \hat{\tau}_3(5)} z_{3, \hat{\tau}_3(6)} z_{4, \hat{\tau}_3(7)} t_1 t_3 t_4 + z_{2, \hat{\tau}_4(5)} z_{3, \hat{\tau}_4(6)} z_{4, \hat{\tau}_4(7)} t_2 t_3 t_4, \tag{415}
\end{aligned}$$

where $\hat{\tau}_i$, ($i = 1, 2, 3, 4$) are elements of the permutation group S_3 on the column indices 5, 6, 7. Taking $\hat{\tau}_1 = \hat{\tau}_4 = \hat{e}$, $\hat{\tau}_2 = (\widehat{567})$, and $\hat{\tau}_3 = (\widehat{576})$, we have

$$z_{1,5} z_{2,6} z_{3,7} = z_{2,5} z_{3,6} z_{4,7} = z_{1,6} z_{2,7} z_{4,5} = z_{1,7} z_{3,5} z_{4,6} = 1. \tag{416}$$

The solution of Eq.(358)

$$z_{i,j} = 1, \quad i = 1, \dots, 4, j = 5, 6, 7, \tag{417}$$

The integral is embedded in the subvariety of the Grassmannian $G_{4,8}$ whose splitting local coordinates are written as

$$\left(\begin{array}{cccccccc}
1 & 0 & 0 & 0 & 1 & 1 & 1 & m_1^2 \\
0 & 1 & 0 & 0 & 1 & 1 & 1 & m_2^2 \\
0 & 0 & 1 & 0 & 1 & 1 & 1 & m_3^2 \\
0 & 0 & 0 & 1 & 1 & 1 & 1 & p^2
\end{array} \right). \tag{418}$$

Similarly the Feynman integral of the 2-loop diagram with 4 propagators can be written as

$$\begin{aligned}
A_{2L4}(p^2) = & -\frac{i^{4-D}\Gamma(4-D)\exp\left\{\frac{i\pi(2-D)}{2}\right\}\left(\Lambda_{\text{RE}}^2\right)^{4-D}}{(4\pi)^D}\int\prod_{i=1}^4\omega_5(t)\delta(t_1t_2t_3 \\
& +t_1t_2t_4+t_1t_3t_4+t_1t_2t_5+t_1t_3t_5+t_2t_3t_5+t_2t_4t_5+t_3t_4t_5) \\
& \times\left[t_1t_2+t_1t_3+t_2t_3+t_2t_4+t_3t_4\right]^{1-D/2}\left[\sum_{i=1}^4t_im_i^2+t_5p^2\right]^{D-4}, \quad (419)
\end{aligned}$$

and the Feynman integral of the 2-loop self energy with 5 propagators can be accordingly expressed as

$$\begin{aligned}
A_{2L5}(p^2) = & \frac{\exp\left\{\frac{i\pi(8-2D)}{2}\right\}\Gamma(5-D)\left(\Lambda_{\text{RE}}^2\right)^{5-D}}{(4\pi)^D}\int\omega_6(t) \\
& \times\left[t_1(t_2+t_4+t_5)+(t_2+t_4)(t_3+t_5)+t_3t_5\right]^{1-D/2} \\
& \times\delta\left(t_1t_2(t_3+t_4)+(t_1+t_2)(t_3t_4+t_3t_5+t_4t_5)\right. \\
& \left.+t_1t_6(t_2+t_4+t_5)+t_6(t_2+t_4)(t_3+t_5)+t_3t_5t_6\right) \\
& \times\left[\sum_{i=1}^5t_im_i^2+t_6p^2\right]^{D-5}, \quad (420)
\end{aligned}$$

respectively. Using the integrals in projective space above, we obtain the fundamental solution systems in neighborhoods of different regular singularities. Concrete results in detail are presented elsewhere.

VI. SUMMARY

The Feynman integrals can be taken as functions on the subvarieties of Grassmannians through homogenizing the parametric representation. Thus the GKZ-systems satisfied by the Feynman integrals are obtained in splitting local coordinates accordingly. The dual PDEs of GKZ-system determine the matrices of integer lattice and corresponding exponent matrices in splitting local coordinates. Actually the dual PDEs can be written as the PDEs satisfied by these scalar integrals in combined local coordinates. Using those matrices of integer lattice and the corresponding exponent matrices, we obtain the fundamental solution systems in neighborhoods of the regular singularities.

Taking the diagrams of the 1-loop self-energy, the 1-loop triangle with massless virtual particles, and the 2-loop vacuum as examples, we elucidate how to construct the funda-

mental solution systems of the corresponding Feynman integral in neighborhoods of the regular singularities. For most multi-loop diagrams, the sizes of Grassmannians embedded by Feynman integrals are too large to construct the fundamental systems in neighborhoods of the regular singularities efficiently through the Feynman parametric representations. To efficiently derive the fundamental solution system, we can embed the corresponding scalar integrals into the subvarieties of Grassmannians using the α -parametric representation. In the last section, we present the parametric representations of diagrams of the 2-loop self energy with 3 propagators, 4 propagators, and 5 propagators, respectively. Generally the Feynman integrals can be written as linear combinations of those hypergeometric functions of the fundamental solution systems in the neighborhoods of regular singularities. The integration constants, i.e. the combination coefficients, are determined from the Feynman integral on an ordinary point or some regular singularities. The concrete expressions of the Feynman integrals together with their partial derivatives of some regular singularities can be evaluated by the Gegenbauer polynomials [77] or Bessel functions [42, 78].

Acknowledgments

The work has been supported partly by the National Natural Science Foundation of China (NNSFC) with Grant No. 12075074, No. 11535002, No. 11745006, No. 11675239, No. 11821505, No. 11705045, No. 11947302 and No. 12075301, the Natural Science Foundation of Guangxi Autonomous Region with Grant No. 2022GXNSFDA035068, and the Natural Science Foundation of Hebei Province with Grant No. A2022201017, the youth top-notch talent support program of the Hebei Province. Furthermore, the author (C.-H. Chang) is also supported by Key Research program of Frontier Sciences, CAS, Grant No. QYZDY-SSW-SYS006.

Appendix A: The fundamental solution systems of $A^{(1T)'$

When we choose the first, second, third, fourth column vectors of $A^{(1T)'}$ as the basis of the projective plane, the corresponding hypergeometric functions contain redundant dependent indeterminates. Because $\det(A_{\{1,2,4,6\}}^{(1T)'}) = -1$ where the matrix $A_{\{1,2,4,6\}}^{(1T)'}$ denotes the submatrix

of $A^{(1T)'}$ composed of the first, second, fourth and sixth columns, one derives

$$\left(A_{\{1,2,4,6\}}^{(1T)'} \right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} 1 & 0 & -1 & 0 & p_2^2 - p_1^2 & 0 & 0 \\ 0 & 1 & -1 & 0 & p_3^2 - p_1^2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & p_1^2 & 1 & 1 \end{pmatrix}. \quad (\text{A1})$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & 0 & \alpha_{1,3} & 0 & \alpha_{1,5} & 0 & 0 \\ 0 & 0 & \alpha_{2,3} & 0 & \alpha_{2,5} & 0 & 0 \\ 0 & 0 & 0 & 3 - D & 0 & 0 & D - 4 \\ 0 & 0 & \alpha_{4,3} & 0 & \alpha_{4,5} & \frac{D}{2} - 3 & 3 - D \end{pmatrix}, \quad (\text{A2})$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,3} + \alpha_{1,5} &= -1, \quad \alpha_{2,3} + \alpha_{2,5} = -1, \quad \alpha_{4,3} + \alpha_{4,5} = \frac{D}{2} - 1, \\ \alpha_{1,3} + \alpha_{2,3} + \alpha_{4,3} &= 0, \quad \alpha_{1,5} + \alpha_{2,5} + \alpha_{4,5} = \frac{D}{2} - 3. \end{aligned} \quad (\text{A3})$$

Corresponding to the matrix of local coordinates in Eq.(A1), we find 12 choices on the matrix of integer lattice whose submatrix composed of the third and fifth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 3), (1, 5), (3, 5)\}$, and other elements are all zero. Basing on the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(3)}$, we obtain the exponents as

$$\alpha_{1,3} = 2 - \frac{D}{2}, \quad \alpha_{1,5} = \frac{D}{2} - 3, \quad \alpha_{2,3} = -1, \quad \alpha_{2,5} = 0, \quad \alpha_{4,3} = \frac{D}{2} - 1, \quad \alpha_{4,5} = 0, \quad (\text{A4})$$

where the geometric representation is presented in Fig.11(1) with $(a, b) = (1, 5)$, $\{c, d\} = \{2, 3\}$. In a similar way the exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{2,3} = 1 - \frac{D}{2}, \quad \alpha_{2,5} = \frac{D}{2} - 2, \quad \alpha_{4,3} = \frac{D}{2} - 1, \quad \alpha_{4,5} = 0, \quad (\text{A5})$$

whose geometric representation is plotted in Fig.11(2) with $a = 2$, $b = 3$, $(c, d) = (1, 5)$. The exponents corresponding to the integer lattice $n_1 E_4^{(3)} + n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{2,3} = 0, \quad \alpha_{2,5} = -1, \quad \alpha_{4,3} = 0, \quad \alpha_{4,5} = \frac{D}{2} - 1, \quad (\text{A6})$$




FIG. 11: The geometric descriptions of single orbits in Eq.(323) on the projective plane CP^4 , where the line $\{4,6,7\}$ intersects with the hyperplane $\{1,2,3\}$ at the point 6.

whose geometric representation is drawn in Fig.11(3) with $a = 3$, $(b, c, d) = (1, 2, 5)$. Three linear independent hypergeometric functions are evidently written as

$$\begin{aligned}
\psi_{\{1,2,4,6\}}^{(1)}(p_1^2, p_2^2, p_3^2) &\sim (p_2^2 - p_1^2)^{D/2-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\
&\quad \times \left(\frac{p_3^2 - p_1^2}{p_2^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2}{p_1^2 - p_2^2}\right)^{n_2} \\
&\sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\
\psi_{\{1,2,4,6\}}^{(2)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_3^2 - p_1^2)^{D/2-2}}{p_2^2 - p_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\
&\quad \times \left(\frac{p_3^2 - p_1^2}{p_2^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2}{p_1^2 - p_3^2}\right)^{n_2} \\
&\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,2,4,6\}}^{(3)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \sum_{n_1, n_2}^{\infty} \\
&\quad \times \left(\frac{p_1^2}{p_1^2 - p_2^2}\right)^{n_1} \left(\frac{p_1^2}{p_1^2 - p_3^2}\right)^{n_2}
\end{aligned}$$

$$\begin{aligned} & \times \frac{1}{\Gamma(1 - n_1 - n_2)\Gamma(\frac{D}{2} + n_1 + n_2)} \\ & \sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)}, \end{aligned} \quad (\text{A7})$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. In other words, those hypergeometric functions constitute a fundamental solution system of the GKZ-system in a proper subset of the whole parameter space. Although the embedded subvariety of Grassmannian Eq.(323) is different from the subvariety of Eq.(268), the constructed hypergeometric function satisfies the same PDEs of combined coordinates in Eq.(278).

The exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(3)}$ are presented as

$$\alpha_{1,3} = 0, \alpha_{1,5} = -1, \alpha_{2,3} = 0, \alpha_{2,5} = -1, \alpha_{4,3} = 0, \alpha_{4,5} = \frac{D}{2} - 1, \quad (\text{A8})$$

whose geometric representation is presented in Fig.11(1) with $(a, b) = (1, 3)$, $\{c, d\} = \{2, 5\}$. Similarly the exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,5} = 0, \alpha_{2,3} = 1, \alpha_{2,5} = -2, \alpha_{4,3} = 0, \alpha_{4,5} = \frac{D}{2} - 1, \quad (\text{A9})$$

whose geometric representation is drawn in Fig.11(2) with $a = 2$, $b = 5$, $(c, d) = (1, 3)$. Corresponding to the integer lattice $-n_1 E_4^{(3)} - n_2 E_4^{(5)}$, the exponents are

$$\alpha_{1,3} = -1, \alpha_{1,5} = 0, \alpha_{2,3} = -1, \alpha_{2,5} = 0, \alpha_{4,3} = 2, \alpha_{4,5} = \frac{D}{2} - 3, \quad (\text{A10})$$

whose geometric representation is plotted in Fig.11(3) with $a = 5$, $(b, c, d) = (1, 2, 3)$. The hypergeometric functions are evidently presented as

$$\begin{aligned} \psi_{\{1,2,4,6\}}^{(4)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\ & \times \left(\frac{p_2^2 - p_1^2}{p_3^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2 - p_2^2}{p_1^2}\right)^{n_2} \\ & \sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\ \psi_{\{1,2,4,6\}}^{(5)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-1}}{(p_3^2 - p_1^2)^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\ & \times \left(\frac{p_2^2 - p_1^2}{p_3^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2 - p_3^2}{p_1^2}\right)^{n_2} \end{aligned}$$

$$\begin{aligned}
& \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,2,4,6\}}^{(6)}(p_1^2, p_2^2, p_3^2) & \sim (p_1^2)^{D/2-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{\Gamma(3 + n_1 + n_2)} \\
& \quad \times \left(\frac{p_1^2 - p_2^2}{p_1^2}\right)^{n_1} \left(\frac{p_1^2 - p_3^2}{p_1^2}\right)^{n_2}, \tag{A11}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The hypergeometric functions constitute a fundamental solution system of the GKZ-system of PDEs in a proper subset of the whole parameter space.

As the integer lattice is chosen as $n_1 E_4^{(1)} - n_2 E_4^{(3)}$, the exponents are

$$\alpha_{1,3} = 1, \alpha_{1,5} = -2, \alpha_{2,3} = -1, \alpha_{2,5} = 0, \alpha_{4,3} = 0, \alpha_{4,5} = \frac{D}{2} - 1, \tag{A12}$$

whose geometric representation is presented in Fig.11(2) with $a = 1, b = 5, \{c, d\} = \{2, 3\}$.

The exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(5)}$ are given by

$$\alpha_{1,3} = 0, \alpha_{1,5} = -1, \alpha_{2,3} = 0, \alpha_{2,5} = -1, \alpha_{4,3} = 0, \alpha_{4,5} = \frac{D}{2} - 1, \tag{A13}$$

whose geometric representation is plotted in Fig.11(1) with $(a, b) = (2, 3), \{c, d\} = \{1, 5\}$.

The exponents of the integer lattice $n_1 E_4^{(3)} - n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,5} = -1, \alpha_{2,3} = -1, \alpha_{2,5} = 0, \alpha_{4,3} = 1, \alpha_{4,5} = \frac{D}{2} - 2, \tag{A14}$$

whose geometric representation is presented in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 5), (2, 3)\}$. Three hypergeometric functions are obtained as

$$\begin{aligned}
\psi_{\{1,2,4,6\}}^{(7)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\
& \quad \times \left(\frac{p_3^2 - p_1^2}{p_2^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2 - p_2^2}{p_1^2}\right)^{n_2} \\
& \sim \frac{(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_2^2 - p_3^2)}, \\
\psi_{\{1,2,4,6\}}^{(8)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_1^2 - p_3^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\
& \quad \times \left(\frac{p_3^2 - p_1^2}{p_2^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2 - p_3^2}{p_1^2}\right)^{n_2} \\
& \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,2,4,6\}}^{(9)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2}}{(p_1^2 - p_2^2)^2(p_3^2 - p_2^2)} \sum_{n=0}^{\infty} \frac{\Gamma(1+n)}{\Gamma(\frac{D}{2} + 1 + n)} \left(\frac{p_1^2}{p_1^2 - p_2^2}\right)^n, \tag{A15}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The hypergeometric function of the exponents of Eq.(A14) is

$$\begin{aligned}
\psi_{\{1,2,4,6\}}^{(9)'}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_1^2}{p_2^2 - p_1^2}\right)^{n_1} \left(\frac{p_3^2 - p_1^2}{p_1^2}\right)^{n_2} \\
&\quad \times \frac{1}{\Gamma(2 - n_1 + n_2)\Gamma(\frac{D}{2} - 1 + n_1 - n_2)} \\
&\sim -\psi_{\{1,2,4,6\}}^{(8)}(p_1^2, p_2^2, p_3^2) \\
&\quad + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(\frac{D}{2} + 1)} \psi_{\{1,2,4,6\}}^{(9)}(p_1^2, p_2^2, p_3^2). \tag{A16}
\end{aligned}$$

Furthermore, the sets of the hypergeometric functions $\{\psi_{\{1,2,4,6\}}^{(1)}, \psi_{\{1,2,4,6\}}^{(8)}, \psi_{\{1,2,4,6\}}^{(9)}\}$ and $\{\psi_{\{1,2,4,6\}}^{(6)}, \psi_{\{1,2,4,6\}}^{(7)}, \psi_{\{1,2,4,6\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

The exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(3)}$ are

$$\alpha_{1,3} = 1 - \frac{D}{2}, \quad \alpha_{1,5} = \frac{D}{2} - 2, \quad \alpha_{2,3} = 0, \quad \alpha_{2,5} = -1, \quad \alpha_{4,3} = \frac{D}{2} - 1, \quad \alpha_{4,5} = 0, \tag{A17}$$

whose geometric representation is presented in Fig.11(2) with $a = 1, b = 3, \{c, d\} = \{2, 5\}$.

Corresponding to the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(5)}$, the exponents are

$$\alpha_{1,3} = -1, \quad \alpha_{1,5} = 0, \quad \alpha_{2,3} = 2 - \frac{D}{2}, \quad \alpha_{2,5} = \frac{D}{2} - 3, \quad \alpha_{4,3} = \frac{D}{2} - 1, \quad \alpha_{4,5} = 0, \tag{A18}$$

whose geometric representation is depicted in Fig.11(1) with $(a, b) = (2, 5), \{c, d\} = \{1, 3\}$.

Finally, the exponents of the integer lattice $-n_1 E_4^{(3)} + n_2 E_4^{(5)}$ are

$$\alpha_{1,3} = -1, \quad \alpha_{1,5} = 0, \quad \alpha_{2,3} = 0, \quad \alpha_{2,5} = -1, \quad \alpha_{4,3} = 1, \quad \alpha_{4,5} = \frac{D}{2} - 2, \tag{A19}$$

whose geometric representation is depicted in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 3), (2, 5)\}$.

Three hypergeometric functions are evidently written as

$$\begin{aligned}
\psi_{\{1,2,4,6\}}^{(10)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_2^2 - p_1^2)^{D/2-2}}{p_3^2 - p_1^2} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \\
&\quad \times \left(\frac{p_2^2 - p_1^2}{p_3^2 - p_1^2}\right)^{n_1} \left(\frac{p_1^2 - p_2^2}{p_1^2}\right)^{n_2} \\
&\sim \frac{(p_2^2)^{D/2-1} (p_2^2 - p_1^2)^{D/2-2}}{(p_1^2)^{D/2-1} (p_2^2 - p_3^2)}, \\
\psi_{\{1,2,4,6\}}^{(11)}(p_1^2, p_2^2, p_3^2) &\sim (p_3^2 - p_1^2)^{D/2-3} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!}
\end{aligned}$$

$$\begin{aligned}
& \times \left(\frac{p_2^2 - p_1^2}{p_3^2 - p_1^2} \right)^{n_1} \left(\frac{p_1^2}{p_1^2 - p_3^2} \right)^{n_2} \\
& \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,2,4,6\}}^{(12)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-2}}{p_1^2 - p_2^2} \sum_{n_1, n_2}^{\infty} \left(\frac{p_2^2 - p_1^2}{p_1^2} \right)^{n_1} \left(\frac{p_1^2}{p_3^2 - p_1^2} \right)^{n_2} \\
& \times \frac{1}{\Gamma(2 + n_1 - n_2) \Gamma(\frac{D}{2} - 1 - n_1 + n_2)} \\
& \sim \frac{(p_1^2)^{D/2-1}}{(p_1^2 - p_2^2)^2} \sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_1)}{n_1!} \\
& \times \left(\frac{p_1^2 - p_2^2}{p_1^2} \right)^{n_1} \left(\frac{p_1^2 - p_2^2}{p_1^2 - p_3^2} \right)^{n_2} \\
& \sim \frac{(p_1^2 - p_3^2)(p_2^2)^{D/2-1}}{(p_2^2 - p_1^2)^2 (p_2^2 - p_3^2)}, \tag{A20}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. In addition, the sets of the hypergeometric functions $\{\psi_{\{1,2,4,6\}}^{(3)}, \psi_{\{1,2,4,6\}}^{(10)}, \psi_{\{1,2,4,6\}}^{(11)}\}$ and $\{\psi_{\{1,2,4,6\}}^{(4)}, \psi_{\{1,2,4,6\}}^{(11)}, \psi_{\{1,2,4,6\}}^{(12)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Because $\det(A_{\{1,3,4,6\}}^{(1T)'}) = 1$ where the matrix $A_{\{1,3,4,6\}}^{(1T)'}$ denotes the submatrix of $A^{(1T)'}$ composed of the first, third, fourth and sixth column vectors, one derives

$$\left(A_{\{1,3,4,6\}}^{(1T)'} \right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} 1 & -1 & 0 & 0 & p_2^2 - p_3^2 & 0 & 0 \\ 0 & -1 & 0 & 0 & p_1^2 - p_3^2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & p_3^2 & 1 & 1 \end{pmatrix}. \tag{A21}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & 0 & 0 & \alpha_{1,5} & 0 & 0 \\ 0 & \alpha_{2,2} & 0 & 0 & \alpha_{2,5} & 0 & 0 \\ 0 & 0 & 0 & 3 - D & 0 & 0 & D - 4 \\ 0 & \alpha_{4,2} & 0 & 0 & \alpha_{4,5} & \frac{D}{2} - 3 & 3 - D \end{pmatrix}, \tag{A22}$$

where the matrix elements satisfy similar relations presented in Eq.(275) through the replacement $\alpha_{i3} \rightarrow \alpha_{i2}$, $i = 1, 2, 4$. Correspondingly there are 12 choices on the matrix of integer lattice whose submatrix composed of the second and fifth columns is formulated as $\pm n_1 E_4^{(i)}$

$\pm n_2 E_4^{(j)}$, with $n_{1,2} \geq 0$, $(i, j) \in \{(1, 3), (1, 5), (3, 5)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents which are obtained through replacements $\alpha_{i_3} \rightarrow \alpha_{i_2}$ of the functions $\psi_{\{1,2,4,6\}}^{(i)}$. The geometric representations are also acquired through the permutation $(\widehat{23})$. The corresponding hypergeometric functions are finally written as

$$\psi_{\{1,3,4,6\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,2,4,6\}}^{(i)}(p_3^2, p_2^2, p_1^2), \quad (i = 1, \dots, 12). \quad (\text{A23})$$

Because $\det(A_{\{2,3,4,6\}}^{(1T)'}) = -1$, we have

$$\left(A_{\{2,3,4,6\}}^{(1T)'}\right)^{-1} \cdot A_{\{2,3,4,6\}}^{(1T)'} = \begin{pmatrix} -1 & 1 & 0 & 0 & p_3^2 - p_2^2 & 0 & 0 \\ -1 & 0 & 0 & 0 & p_1^2 - p_2^2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & p_2^2 & 1 & 1 \end{pmatrix}. \quad (\text{A24})$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & 0 & 0 & \alpha_{1,5} & 0 & 0 \\ \alpha_{2,1} & 0 & 0 & 0 & \alpha_{2,5} & 0 & 0 \\ 0 & 0 & 0 & 3 - D & 0 & 0 & D - 4 \\ \alpha_{4,1} & 0 & 0 & 0 & \alpha_{4,5} & \frac{D}{2} - 3 & 3 - D \end{pmatrix}, \quad (\text{A25})$$

where the matrix elements satisfy similar relations presented in Eq.(275) through the replacement $\alpha_{i_3} \rightarrow \alpha_{i_1}$, $i = 1, 2, 4$. Corresponding to the matrix of splitting local coordinates in Eq.(A1), we have 12 choices on the matrix of integer lattice whose submatrix composed of the first and fifth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 3), (1, 5), (3, 5)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents which are obtained through replacements $\alpha_{i_3} \rightarrow \alpha_{i_1}$ of that of the function $\psi_{\{1,2,4,6\}}^{(i)}$. The geometric representations are also acquired through the permutation $(\widehat{13})$ accordingly. The hypergeometric functions are

$$\psi_{\{2,3,4,6\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,2,4,6\}}^{(i)}(p_2^2, p_3^2, p_1^2), \quad (i = 1, \dots, 12). \quad (\text{A26})$$

Obviously $\det(A_{\{1,2,6,7\}}^{(1T)'}) = 1$ where the matrix $A_{\{1,2,6,7\}}^{(1T)'}$ denotes the submatrix of $A_{\{1,2,6,7\}}^{(1T)'}$

composed of the first, second, sixth and seventh columns, one has

$$\left(A_{\{1,2,6,7\}}^{(1T)'}\right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} 1 & 0 & -1 & 0 & p_2^2 - p_1^2 & 0 & 0 \\ 0 & 1 & -1 & 0 & p_3^2 - p_1^2 & 0 & 0 \\ 0 & 0 & 1 & -1 & p_1^2 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}. \quad (\text{A27})$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & 0 & \alpha_{1,3} & 0 & \alpha_{1,5} & 0 & 0 \\ 0 & 0 & \alpha_{2,3} & 0 & \alpha_{2,5} & 0 & 0 \\ 0 & 0 & \alpha_{3,3} & 3 - D & \alpha_{3,5} & \frac{D}{2} - 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad (\text{A28})$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,3} + \alpha_{1,5} &= -1, \quad \alpha_{2,3} + \alpha_{2,5} = -1, \quad \alpha_{3,3} + \alpha_{3,5} = \frac{D}{2} - 1, \\ \alpha_{1,3} + \alpha_{2,3} + \alpha_{3,3} &= 0, \quad \alpha_{1,5} + \alpha_{2,5} + \alpha_{3,5} = \frac{D}{2} - 3. \end{aligned} \quad (\text{A29})$$

Certainly there are 12 choices on the matrix of integer lattice whose submatrix composed of the third and fifth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents. The exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are written as

$$\alpha_{1,3} = 2 - \frac{D}{2}, \quad \alpha_{1,5} = \frac{D}{2} - 3, \quad \alpha_{2,3} = -1, \quad \alpha_{2,5} = 0, \quad \alpha_{3,3} = \frac{D}{2} - 1, \quad \alpha_{3,5} = 0, \quad (\text{A30})$$

whose geometric representation is plotted in Fig.11(1) with $(a, b) = (1, 5)$, $\{c, d\} = \{2, 3\}$. The exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{2,3} = 1 - \frac{D}{2}, \quad \alpha_{2,5} = \frac{D}{2} - 2, \quad \alpha_{3,3} = \frac{D}{2} - 1, \quad \alpha_{3,5} = 0, \quad (\text{A31})$$

whose geometric representation is presented in Fig.11(2) with $a = 2$, $b = 3$, $(c, d) = (1, 5)$. In a similar way, the exponents of the integer lattice $n_1 E_4^{(2)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = 0, \quad \alpha_{1,5} = -1, \quad \alpha_{2,3} = 0, \quad \alpha_{2,5} = -1, \quad \alpha_{3,3} = 0, \quad \alpha_{3,5} = \frac{D}{2} - 1, \quad (\text{A32})$$

whose geometric representation is depicted in Fig.11(3) with $a = 3$, $(b, c, d) = (1, 2, 5)$.

The exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(2)}$ are presented as

$$\alpha_{1,3} = 0, \alpha_{1,5} = -1, \alpha_{2,3} = 0, \alpha_{2,5} = -1, \alpha_{3,3} = 0, \alpha_{3,5} = \frac{D}{2} - 1, \quad (\text{A33})$$

whose geometric representation is depicted in Fig.11(1) with $(a, b) = (1, 3)$, $\{c, d\} = \{2, 5\}$. Corresponding to the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(4)}$, the exponents are given by

$$\alpha_{1,3} = -1, \alpha_{1,5} = 0, \alpha_{2,3} = 1, \alpha_{2,5} = -2, \alpha_{3,3} = 0, \alpha_{3,5} = \frac{D}{2} - 1, \quad (\text{A34})$$

whose geometric representation is presented in Fig.11(2) with $a = 2, b = 5, (c, d) = (1, 3)$. Similarly the exponents of the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,5} = 0, \alpha_{2,3} = -1, \alpha_{2,5} = 0, \alpha_{3,3} = 2, \alpha_{3,5} = \frac{D}{2} - 3, \quad (\text{A35})$$

whose geometric representation is plotted in Fig.11(3) with $a = 5, (b, c, d) = (1, 2, 3)$.

Corresponding to the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(2)}$, the exponents are

$$\alpha_{1,3} = 1, \alpha_{1,5} = -2, \alpha_{2,3} = -1, \alpha_{2,5} = 0, \alpha_{3,3} = 0, \alpha_{3,5} = \frac{D}{2} - 1, \quad (\text{A36})$$

whose geometric representation is depicted in Fig.11(2) with $a = 1, b = 5, \{c, d\} = \{2, 3\}$. The exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = 0, \alpha_{1,5} = -1, \alpha_{2,3} = 0, \alpha_{2,5} = -1, \alpha_{3,3} = 0, \alpha_{3,5} = \frac{D}{2} - 1, \quad (\text{A37})$$

whose geometric representation is presented in Fig.11(1) with $(a, b) = (2, 3)$, $\{c, d\} = \{1, 5\}$. Similarly the exponents of the integer lattice $n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are given by

$$\alpha_{1,3} = 0, \alpha_{1,5} = -1, \alpha_{2,3} = -1, \alpha_{2,5} = 0, \alpha_{3,3} = 1, \alpha_{3,5} = \frac{D}{2} - 2, \quad (\text{A38})$$

whose geometric representation is presented in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 5), (2, 3)\}$. Meanwhile the exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are

$$\alpha_{1,3} = 1 - \frac{D}{2}, \alpha_{1,5} = \frac{D}{2} - 2, \alpha_{2,3} = 0, \alpha_{2,5} = -1, \alpha_{3,3} = \frac{D}{2} - 1, \alpha_{3,5} = 0, \quad (\text{A39})$$

whose geometric representation is presented in Fig.11(2) with $a = 1, b = 3, \{c, d\} = \{2, 5\}$. The matrix of exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(4)}$ is formulated as

$$\alpha_{1,3} = -1, \alpha_{1,5} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{2,5} = \frac{D}{2} - 3, \alpha_{3,3} = \frac{D}{2} - 1, \alpha_{3,5} = 0, \quad (\text{A40})$$

whose geometric representation is drawn in Fig.11(1) with $(a, b) = (2, 5)$, $\{c, d\} = \{1, 3\}$. Finally, the exponents of the integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,3} = -1, \alpha_{1,5} = 0, \alpha_{2,3} = 0, \alpha_{2,5} = -1, \alpha_{3,3} = 1, \alpha_{3,5} = \frac{D}{2} - 2, \quad (\text{A41})$$

whose geometric representation is depicted in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 3), (2, 5)\}$.

The corresponding hypergeometric functions are evidently written as

$$\psi_{\{1,2,6,7\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,2,4,6\}}^{(i)}(p_1^2, p_2^2, p_3^2), \quad (i = 1, \dots, 12). \quad (\text{A42})$$

Furthermore $\det(A_{\{1,3,6,7\}}^{(1T)'}) = -1$, $\det(A_{\{2,3,6,7\}}^{(1T)'}) = 1$, the calculation in detail implies

$$\begin{aligned} \psi_{\{1,3,6,7\}}^{(i)}(p_1^2, p_2^2, p_3^2) &= \psi_{\{1,2,4,6\}}^{(i)}(p_3^2, p_2^2, p_1^2), \\ \psi_{\{2,3,6,7\}}^{(i)}(p_1^2, p_2^2, p_3^2) &= \psi_{\{1,2,4,6\}}^{(i)}(p_2^2, p_3^2, p_1^2). \end{aligned} \quad (\text{A43})$$

Because $\det(A_{\{1,4,5,6\}}^{(1T)'}) = p_3^2 - p_1^2$, where the matrix $A_{\{1,4,5,6\}}^{(1T)'}$ denotes the submatrix of $A^{(1T)'}$ composed of the first, fourth, fifth and sixth columns, we get

$$\left(A_{\{1,4,5,6\}}^{(1T)'} \right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} 1 & -\frac{p_2^2 - p_1^2}{p_3^2 - p_1^2} & -\frac{p_3^2 - p_2^2}{p_3^2 - p_1^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & \frac{1}{p_3^2 - p_1^2} & -\frac{1}{p_3^2 - p_1^2} & 0 & 1 & 0 & 0 \\ 0 & -\frac{p_1^2}{p_3^2 - p_1^2} & \frac{p_3^2}{p_3^2 - p_1^2} & 0 & 0 & 1 & 1 \end{pmatrix}. \quad (\text{A44})$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 - D & 0 & 0 & D - 4 \\ 0 & \alpha_{3,2} & \alpha_{3,3} & 0 & \frac{D}{2} - 3 & 0 & 0 \\ 0 & \alpha_{4,2} & \alpha_{4,3} & 0 & 0 & \frac{D}{2} - 3 & 3 - D \end{pmatrix}, \quad (\text{A45})$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,2} + \alpha_{1,3} &= -1, \alpha_{3,2} + \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{4,2} + \alpha_{4,3} = \frac{D}{2} - 1, \\ \alpha_{1,2} + \alpha_{3,2} + \alpha_{4,2} &= 0, \alpha_{1,3} + \alpha_{3,3} + \alpha_{4,3} = 0. \end{aligned} \quad (\text{A46})$$

Corresponding to the matrix of splitting local coordinates in Eq.(A44), one obtains 12 choices on the matrix of integer lattice whose submatrix composed of the third and fifth columns

is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(2, 3), (2, 6), (3, 6)\}$, and other elements are all zero.

The exponents of the integer lattice $n_1 E_4^{(2)} + n_2 E_4^{(3)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{3,2} = 2 - \frac{D}{2}, \alpha_{3,3} = 0, \alpha_{4,2} = \frac{D}{2} - 1, \alpha_{4,3} = 0, \quad (\text{A47})$$

whose geometric representation is presented in Fig.11(1) with $(a, b) = (1, 3)$, $\{c, d\} = \{2, 5\}$.

The exponents of the integer lattice $n_1 E_4^{(2)} + n_2 E_4^{(6)}$ are

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{3,2} = 1 - \frac{D}{2}, \alpha_{3,3} = 1, \alpha_{4,2} = \frac{D}{2} - 1, \alpha_{4,3} = 0, \quad (\text{A48})$$

whose geometric representation is depicted in Fig.11(2) with $a = 5$, $b = 2$, $(c, d) = (1, 3)$.

The exponents of the integer lattice $n_1 E_4^{(3)} + n_2 E_4^{(6)}$ are presented by

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{3,2} = 0, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{4,2} = 0, \alpha_{4,3} = \frac{D}{2} - 1, \quad (\text{A49})$$

whose geometric representation is presented in Fig.11(3) with $a = 2$, $(b, c, d) = (1, 3, 5)$. The corresponding hypergeometric functions are obviously written as

$$\begin{aligned} \psi_{\{1,4,5,6\}}^{(1)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1) \Gamma(1 - \frac{D}{2} + n_2)}{n_1! n_2!} \\ &\quad \times \left(\frac{p_1^2 - p_2^2}{p_3^2 - p_2^2} \right)^{n_1} \left(\frac{p_3^2(p_1^2 - p_2^2)}{p_1^2(p_3^2 - p_2^2)} \right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_3^2 - p_2^2)}, \\ \psi_{\{1,4,5,6\}}^{(2)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-3} (p_3^2)^2}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{\Gamma(3 + n_1 + n_2)} \\ &\quad \times \frac{\Gamma(1 + n_2)}{\Gamma(4 - \frac{D}{2} + n_2)} \left(\frac{p_3^2(p_1^2 - p_2^2)}{p_1^2(p_3^2 - p_2^2)} \right)^{n_1} \left(\frac{p_3^2}{p_1^2} \right)^{n_2}, \\ \psi_{\{1,4,5,6\}}^{(3)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2! \Gamma(1 - n_1 - n_2) \Gamma(\frac{D}{2} + n_1 + n_2)} \\ &\quad \times (-)^{n_1 + n_2} \left(\frac{p_3^2(p_1^2 - p_2^2)}{p_1^2(p_3^2 - p_2^2)} \right)^{n_1} \left(\frac{p_3^2}{p_1^2} \right)^{n_2} \\ &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \end{aligned} \quad (\text{A50})$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. In addition the hypergeometric function of the exponents of Eq.(A48) is

expressed as

$$\begin{aligned}
\psi_{\{1,4,5,6\}}^{(2)'}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n_1 - n_2)}{\Gamma(2 + n_1 - n_2)} \\
&\times \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2!} \left(\frac{p_1^2 - p_2^2}{p_3^2 - p_2^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2} \\
&\sim \psi_{\{1,4,5,6\}}^{(1)}(p_1^2, p_2^2, p_3^2) \\
&+ \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(4 - \frac{D}{2})} \psi_{\{1,4,5,6\}}^{(2)}(p_1^2, p_2^2, p_3^2). \tag{A51}
\end{aligned}$$

The exponents of the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(3)}$ are presented as

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{3,2} = 0, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{4,2} = 0, \alpha_{4,3} = \frac{D}{2} - 1, \tag{A52}$$

whose geometric representation is plotted in Fig.11(1) with $(a, b) = (1, 2)$, $\{c, d\} = \{3, 5\}$. Similarly the exponents of the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(6)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{3,2} = 1, \alpha_{3,3} = 1 - \frac{D}{2}, \alpha_{4,2} = 0, \alpha_{4,3} = \frac{D}{2} - 1, \tag{A53}$$

whose geometric representation is presented in Fig.11(2) with $a = 5$, $b = 3$, $(c, d) = (1, 2)$. Corresponding to the integer lattice $-n_1 E_4^{(3)} - n_2 E_4^{(6)}$, the exponents are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{3,2} = 2 - \frac{D}{2}, \alpha_{3,3} = 0, \alpha_{4,2} = \frac{D}{2} - 1, \alpha_{4,3} = 0, \tag{A54}$$

whose geometric representation is plotted in Fig.11(3) with $a = 3$, $(b, c, d) = (1, 2, 5)$. The corresponding hypergeometric functions are obviously written as

$$\begin{aligned}
\psi_{\{1,4,5,6\}}^{(4)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1) \Gamma(1 - \frac{D}{2} + n_2)}{n_1! n_2!} \\
&\times \left(\frac{p_3^2 - p_2^2}{p_1^2 - p_2^2}\right)^{n_1} \left(\frac{p_1^2(p_3^2 - p_2^2)}{p_3^2(p_1^2 - p_2^2)}\right)^{n_2} \\
&\sim \frac{(p_2^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,4,5,6\}}^{(5)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^2 (p_3^2)^{D/2-3}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(3 - \frac{D}{2} + n_1 + n_2)}{\Gamma(3 + n_1 + n_2)} \\
&\times \frac{\Gamma(1 + n_2)}{\Gamma(4 - \frac{D}{2} + n_2)} \left(\frac{p_1^2(p_3^2 - p_2^2)}{p_3^2(p_1^2 - p_2^2)}\right)^{n_1} \left(\frac{p_1^2}{p_3^2}\right)^{n_2}, \\
\psi_{\{1,4,5,6\}}^{(6)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_2)}{n_2! \Gamma(1 - n_1 - n_2) \Gamma(\frac{D}{2} + n_1 + n_2)}
\end{aligned}$$

$$\begin{aligned}
& \times (-)^{n_1+n_2} \left(\frac{p_1^2(p_3^2 - p_2^2)}{p_3^2(p_1^2 - p_2^2)} \right)^{n_1} \left(\frac{p_1^2}{p_3^2} \right)^{n_2} \\
& \sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)}, \tag{A55}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The fact indicates that those linear independent hypergeometric functions constitute a fundamental solution system of the GKZ-system in the proper subset of the whole parameter space. Actually the hypergeometric function of the exponents of Eq.(A53) is

$$\begin{aligned}
\psi_{\{1,4,5,6\}}^{(5)'}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_3^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 1 + n_1 - n_2) \Gamma(1 - \frac{D}{2} + n_2)}{n_2! \Gamma(2 + n_1 - n_2)} \\
& \times \left(\frac{p_3^2 - p_2^2}{p_1^2 - p_2^2} \right)^{n_1} \left(\frac{p_1^2}{p_3^2} \right)^{n_2} \\
& \sim \psi_{\{1,4,5,6\}}^{(4)}(p_1^2, p_2^2, p_3^2) \\
& + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi (\frac{D}{2} - 3)} \psi_{\{1,4,5,6\}}^{(5)}(p_1^2, p_2^2, p_3^2). \tag{A56}
\end{aligned}$$

Similarly the exponents of the integer lattice $n_1 E_4^{(2)} - n_2 E_4^{(3)}$ are

$$\alpha_{1,2} = \frac{D}{2} - 2, \alpha_{1,3} = 1 - \frac{D}{2}, \alpha_{3,2} = 2 - \frac{D}{2}, \alpha_{3,3} = 0, \alpha_{4,2} = 0, \alpha_{4,3} = \frac{D}{2} - 1 \tag{A57}$$

whose geometric representation is drawn in Fig.11(2) with $a = 1, b = 3, \{c, d\} = \{2, 5\}$. The matrix of exponents of the integer lattice $n_1 E_4^{(2)} - n_2 E_4^{(6)}$ is formulated as

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{3,2} = 0, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{4,2} = 0, \alpha_{4,3} = \frac{D}{2} - 1, \tag{A58}$$

whose geometric representation is plotted in Fig.11(1) with $(a, b) = (2, 5), \{c, d\} = \{1, 3\}$. The exponents of the integer lattice $n_1 E_4^{(3)} - n_2 E_4^{(6)}$ are given by

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{3,2} = 2 - \frac{D}{2}, \alpha_{3,3} = 0, \alpha_{4,2} = \frac{D}{2} - 2, \alpha_{4,3} = 1, \tag{A59}$$

where the geometric representation is presented in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 3), (2, 5)\}$. The corresponding hypergeometric functions are obviously written as

$$\begin{aligned}
\psi_{\{1,4,5,6\}}^{(7)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_3^2)^{D/2-1} (p_2^2 - p_1^2)^{D/2-2}}{(p_3^2 - p_1^2) (p_3^2 - p_2^2)^{D/2-1}} \\
& \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1) \Gamma(1 - \frac{D}{2} + n_2)}{n_1! n_2!}
\end{aligned}$$

$$\begin{aligned}
& \times \left(\frac{p_1^2 - p_2^2}{p_3^2 - p_2^2} \right)^{n_1} \left(\frac{p_1^2(p_3^2 - p_2^2)}{p_3^2(p_1^2 - p_2^2)} \right)^{n_2} \\
& \sim \frac{(p_2^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,4,5,6\}}^{(8)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \\
& \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2! \Gamma(1 - n_1 - n_2) \Gamma(3 - \frac{D}{2} + n_1 + n_2)} \\
& \times (-)^{n_1+n_2} \left(\frac{p_1^2 - p_2^2}{p_3^2 - p_2^2} \right)^{n_1} \left(\frac{p_1^2}{p_3^2} \right)^{n_2} \\
& \sim \frac{(p_3^2)^{D/2-1}}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)}, \\
\psi_{\{1,4,5,6\}}^{(9)}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2} p_3^2}{p_3^2(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \\
& \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + n_1 + n_2) \Gamma(1 + n_2)}{\Gamma(3 + n_1 + n_2) \Gamma(\frac{D}{2} + 1 + n_2)} \\
& \times \left(\frac{p_1^2 - p_2^2}{p_3^2 - p_2^2} \right)^{n_1} \left(\frac{p_1^2}{p_3^2} \right)^{n_2}, \tag{A60}
\end{aligned}$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The hypergeometric function of the exponents in Eq.(A59) is

$$\begin{aligned}
\psi_{\{1,4,5,6\}}^{(9)'}(p_1^2, p_2^2, p_3^2) & \sim \frac{(p_1^2)^{D/2-2} p_3^2}{(p_3^2 - p_1^2)(p_3^2 - p_2^2)} \\
& \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 - n_2) \Gamma(\frac{D}{2} - 2 + n_2)}{n_2! \Gamma(2 + n_1 - n_2)} \\
& \times \left(\frac{p_3^2(p_1^2 - p_2^2)}{p_1^2(p_3^2 - p_2^2)} \right)^{n_1} \left(\frac{p_1^2}{p_3^2} \right)^{n_2} \\
& \sim \frac{(p_2^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_3^2 - p_2^2)} \\
& + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi(\frac{D}{2} + 1)} \psi_{\{1,4,5,6\}}^{(9)}(p_1^2, p_2^2, p_3^2), \tag{A61}
\end{aligned}$$

Furthermore, the sets of the hypergeometric functions $\{\psi_{\{1,4,5,6\}}^{(1)}, \psi_{\{1,4,5,6\}}^{(8)}, \psi_{\{1,4,5,6\}}^{(9)}\}$ and $\{\psi_{\{1,4,5,6\}}^{(6)}, \psi_{\{1,4,5,6\}}^{(7)}, \psi_{\{1,4,5,6\}}^{(8)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Meanwhile the exponents of the integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(3)}$ are

$$\alpha_{1,2} = 1 - \frac{D}{2}, \quad \alpha_{1,3} = \frac{D}{2} - 2, \quad \alpha_{3,2} = 0, \quad \alpha_{3,3} = 2 - \frac{D}{2}, \quad \alpha_{4,2} = \frac{D}{2} - 1, \quad \alpha_{4,3} = 0 \tag{A62}$$

whose geometric representation is depicted in Fig.11(2) with $a = 1, b = 2, \{c, d\} = \{3, 5\}$.

The exponents of the integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(6)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{3,2} = 2 - \frac{D}{2}, \alpha_{3,3} = 0, \alpha_{4,2} = \frac{D}{2} - 1, \alpha_{4,3} = 0, \quad (\text{A63})$$

whose geometric representation is plotted in Fig.11(1) with $(a, b) = (3, 5), \{c, d\} = \{1, 2\}$.

Finally, the exponents of the integer lattice $-n_1 E_4^{(3)} + n_2 E_4^{(6)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{3,2} = 0, \alpha_{3,3} = 2 - \frac{D}{2}, \alpha_{4,2} = 1, \alpha_{4,3} = \frac{D}{2} - 2, \quad (\text{A64})$$

whose geometric representation is shown in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 2), (3, 5)\}$.

The corresponding hypergeometric functions are obviously written as

$$\begin{aligned} \psi_{\{1,4,5,6\}}^{(10)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1} (p_3^2 - p_2^2)^{D/2-2}}{(p_2^2 - p_1^2)^{D/2-1} (p_3^2 - p_1^2)} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} - 2 + n_1) \Gamma(1 - \frac{D}{2} + n_2)}{n_1! n_2!} \\ &\times \left(\frac{p_3^2 - p_2^2}{p_1^2 - p_2^2}\right)^{n_1} \left(\frac{p_3^2 (p_1^2 - p_2^2)}{p_1^2 (p_3^2 - p_2^2)}\right)^{n_2} \\ &\sim \frac{(p_2^2)^{D/2-1}}{(p_1^2 - p_2^2)(p_3^2 - p_2^2)}, \\ \psi_{\{1,4,5,6\}}^{(11)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(1 - \frac{D}{2} + n_2)}{n_2! \Gamma(1 - n_1 - n_2) \Gamma(3 - \frac{D}{2} + n_1 + n_2)} \\ &\times (-)^{n_1+n_2} \left(\frac{p_3^2 - p_2^2}{p_1^2 - p_2^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2} \\ &\sim \frac{(p_1^2)^{D/2-1}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)}, \\ \psi_{\{1,4,5,6\}}^{(12)}(p_1^2, p_2^2, p_3^2) &\sim \frac{(p_3^2)^{D/2}}{p_1^2 (p_2^2 - p_1^2) (p_3^2 - p_1^2)} \\ &\times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(\frac{D}{2} + n_1 + n_2) \Gamma(1 + n_2)}{\Gamma(3 + n_1 + n_2) \Gamma(\frac{D}{2} + 1 + n_2)} \\ &\times \left(\frac{p_3^2 - p_2^2}{p_1^2 - p_2^2}\right)^{n_1} \left(\frac{p_3^2}{p_1^2}\right)^{n_2}, \end{aligned} \quad (\text{A65})$$

where the intersection of their convergent regions is a nonempty proper subset of the whole parameter space. The hypergeometric function of the exponents of Eq.(A64) is

$$\psi_{\{1,4,5,6\}}^{(12)'}(p_1^2, p_2^2, p_3^2) \sim \frac{p_1^2 (p_3^2)^{D/2-2}}{(p_2^2 - p_1^2)(p_3^2 - p_1^2)}$$

$$\begin{aligned}
& \times \sum_{n_1, n_2}^{\infty} \frac{\Gamma(2 - \frac{D}{2} + n_1 - n_2) \Gamma(\frac{D}{2} - 2 + n_2)}{n_2! \Gamma(2 + n_1 - n_2)} \\
& \times \left(\frac{p_1^2 (p_3^2 - p_2^2)}{p_3^2 (p_1^2 - p_2^2)} \right)^{n_1} \left(\frac{p_3^2}{p_1^2} \right)^{n_2} \\
& \sim \frac{(p_2^2)^{D/2-1}}{(p_3^2 - p_2^2)(p_1^2 - p_2^2)} \\
& + \lim_{\epsilon \rightarrow 0} \frac{\sin \pi \epsilon}{\sin \pi (\frac{D}{2} + 1)} \psi_{\{1,4,5,6\}}^{(12)}(p_1^2, p_2^2, p_3^2), \tag{A66}
\end{aligned}$$

Additionally the sets of the hypergeometric functions $\{\psi_{\{1,4,5,6\}}^{(3)}, \psi_{\{1,4,5,6\}}^{(10)}, \psi_{\{1,4,5,6\}}^{(11)}\}$ and $\{\psi_{\{1,4,5,6\}}^{(4)}, \psi_{\{1,4,5,6\}}^{(11)}, \psi_{\{1,4,5,6\}}^{(12)}\}$ constitute the fundamental solution systems of the nonempty proper subsets of the whole parameter space, respectively.

Similarly $\det(A_{2,4,5,6}^{(1T)'}) = p_1^2 - p_2^2$, and

$$\left(A_{\{2,4,5,6\}}^{(1T)'} \right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} -\frac{p_3^2 - p_1^2}{p_2^2 - p_1^2} & 1 & -\frac{p_2^2 - p_3^2}{p_2^2 - p_1^2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ \frac{1}{p_2^2 - p_1^2} & 0 & -\frac{1}{p_2^2 - p_1^2} & 0 & 1 & 0 & 0 \\ -\frac{p_1^2}{p_2^2 - p_1^2} & 0 & \frac{p_2^2}{p_2^2 - p_1^2} & 0 & 0 & 1 & 1 \end{pmatrix}. \tag{A67}$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 - D & 0 & 0 & D - 4 \\ \alpha_{3,1} & 0 & \alpha_{3,3} & 0 & \frac{D}{2} - 3 & 0 & 0 \\ \alpha_{4,1} & 0 & \alpha_{4,3} & 0 & 0 & \frac{D}{2} - 3 & 3 - D \end{pmatrix}, \tag{A68}$$

where the matrix elements satisfy the relations which are obtained by the replacement $\alpha_{i,2} \rightarrow \alpha_{i,1}$, with $(i = 1, 3, 4)$ in Eq.(A46). Correspondingly geometric representations of those exponents are obtained from that of $\psi_{\{1,4,5,6\}}^{(i)}$ through the permutation $(\widehat{12})$. The constructed hypergeometric functions are

$$\psi_{\{2,4,5,6\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,4,5,6\}}^{(i)}(p_1^2, p_3^2, p_2^2), \quad (i = 1, \dots, 12). \tag{A69}$$

Because $\det(A_{3,4,5,6}^{(1T)'}) = p_2^2 - p_3^2$, and

$$\left(A_{\{3,4,5,6\}}^{(1T)'} \right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} -\frac{p_1^2 - p_3^2}{p_2^2 - p_3^2} & -\frac{p_2^2 - p_1^2}{p_2^2 - p_3^2} & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ \frac{1}{p_2^2 - p_3^2} & -\frac{1}{p_2^2 - p_3^2} & 0 & 0 & 1 & 0 & 0 \\ -\frac{p_3^2}{p_2^2 - p_3^2} & \frac{p_2^2}{p_2^2 - p_3^2} & 0 & 0 & 0 & 1 & 1 \end{pmatrix}, \tag{A70}$$

the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3-D & 0 & 0 & D-4 \\ \alpha_{3,1} & \alpha_{3,2} & 0 & 0 & \frac{D}{2}-3 & 0 & 0 \\ \alpha_{4,1} & \alpha_{4,2} & 0 & 0 & 0 & \frac{D}{2}-3 & 3-D \end{pmatrix}, \quad (\text{A71})$$

where the matrix elements satisfy the relations which are obtained by the replacement $\alpha_{i,3} \rightarrow \alpha_{i,1}$, with $(i = 1, 3, 4)$ in Eq.(A46). Correspondingly the geometric representations of those matrices of exponents are obtained from that of $\psi_{\{1,4,5,6\}}^{(i)}$ through the permutation $(\widehat{132})$. The constructed hypergeometric functions are

$$\psi_{\{3,4,5,6\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,4,5,6\}}^{(i)}(p_3^2, p_1^2, p_2^2), \quad (i = 1, \dots, 12). \quad (\text{A72})$$

Because $\det(A_{\{1,5,6,7\}}^{(1T)'}) = p_3^2 - p_1^2$, where the matrix $A_{\{1,5,6,7\}}^{(1T)'}$ denotes the submatrix of $A^{(1T)'}$ composed of the first, fifth, sixth and seventh columns, one finds

$$\left(A_{\{1,5,6,7\}}^{(1T)'}\right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} 1 & -\frac{p_2^2 - p_1^2}{p_3^2 - p_1^2} & -\frac{p_3^2 - p_2^2}{p_3^2 - p_1^2} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{p_3^2 - p_1^2} & -\frac{1}{p_3^2 - p_1^2} & 0 & 1 & 0 & 0 \\ 0 & -\frac{p_1^2}{p_3^2 - p_1^2} & \frac{p_3^2}{p_3^2 - p_1^2} & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}. \quad (\text{A73})$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & 0 & 0 & 0 & 0 \\ 0 & \alpha_{2,2} & \alpha_{2,3} & 0 & \frac{D}{2}-3 & 0 & 0 \\ 0 & \alpha_{3,2} & \alpha_{3,3} & 3-D & 0 & \frac{D}{2}-3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad (\text{A74})$$

where the matrix elements satisfy the relations

$$\begin{aligned} \alpha_{1,2} + \alpha_{1,3} &= -1, \quad \alpha_{2,2} + \alpha_{2,3} = 2 - \frac{D}{2}, \quad \alpha_{3,2} + \alpha_{3,3} = \frac{D}{2} - 1, \\ \alpha_{1,2} + \alpha_{2,2} + \alpha_{3,2} &= 0, \quad \alpha_{1,3} + \alpha_{2,3} + \alpha_{3,3} = 0. \end{aligned} \quad (\text{A75})$$

Corresponding to the matrix of local coordinates in Eq.(A44), one obtains 12 choices on the matrix of integer lattice whose submatrix composed of the third and fifth columns is formulated as $\pm n_1 E_4^{(i)} \pm n_2 E_4^{(j)}$, where $n_{1,2} \geq 0$, $(i, j) \in \{(1, 2), (1, 4), (2, 4)\}$, and other

elements are all zero. Basing on the matrices of integer lattice, we have 12 choices on the matrix of exponents. The exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are written as

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = \frac{D}{2} - 1, \alpha_{3,3} = 0, \quad (\text{A76})$$

whose geometric representation is presented in Fig.11(1) with $(a, b) = (1, 3)$, $\{c, d\} = \{2, 5\}$. In a similar way the exponents of the integer lattice $n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 1 - \frac{D}{2}, \alpha_{2,3} = 1, \alpha_{3,2} = \frac{D}{2} - 1, \alpha_{3,3} = 0, \quad (\text{A77})$$

whose geometric representation is plotted in Fig.11(2) with $a = 5$, $b = 2$, $(c, d) = (1, 3)$. The corresponding exponents of the integer lattice $n_1 E_4^{(2)} + n_2 E_4^{(4)}$ are presented by

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = \frac{D}{2} - 1, \quad (\text{A78})$$

whose geometric representation is shown in Fig.11(3) with $a = 2$, $(b, c, d) = (1, 3, 5)$.

The exponents of the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(2)}$ are presented as

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = \frac{D}{2} - 1, \quad (\text{A79})$$

whose geometric representation is depicted in Fig.11(1) with $(a, b) = (1, 2)$, $\{c, d\} = \{3, 5\}$.

Corresponding to the integer lattice $-n_1 E_4^{(1)} - n_2 E_4^{(4)}$, the exponents are given by

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 1, \alpha_{2,3} = 1 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = \frac{D}{2} - 1, \quad (\text{A80})$$

whose geometric representation is drawn in Fig.11(2) with $a = 5$, $b = 3$, $(c, d) = (1, 2)$.

Similarly the exponents of the integer lattice $-n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = \frac{D}{2} - 1, \alpha_{3,3} = 0, \quad (\text{A81})$$

whose geometric representation is presented in Fig.11(3) with $a = 3$, $(b, c, d) = (1, 2, 5)$.

Meanwhile the exponents of the integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(2)}$ are written as

$$\alpha_{1,2} = \frac{D}{2} - 2, \alpha_{1,3} = 1 - \frac{D}{2}, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = 0, \alpha_{3,3} = \frac{D}{2} - 1 \quad (\text{A82})$$

whose geometric representation is presented in Fig.11(2) with $a = 1$, $b = 3$, $\{c, d\} = \{2, 5\}$.

The corresponding exponents of integer lattice $n_1 E_4^{(1)} - n_2 E_4^{(4)}$ are

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 0, \alpha_{3,3} = \frac{D}{2} - 1, \quad (\text{A83})$$

whose geometric representation is drawn in Fig.11(1) with $(a, b) = (2, 5)$, $\{c, d\} = \{1, 3\}$. The exponents corresponding to the integer lattice $n_1 E_4^{(2)} - n_2 E_4^{(4)}$ are given by

$$\alpha_{1,2} = 0, \alpha_{1,3} = -1, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = \frac{D}{2} - 2, \alpha_{3,3} = 1, \quad (\text{A84})$$

whose geometric representation is depicted in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 3), (2, 5)\}$.

The exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(2)}$ are

$$\alpha_{1,2} = 1 - \frac{D}{2}, \alpha_{1,3} = \frac{D}{2} - 2, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = \frac{D}{2} - 1, \alpha_{3,3} = 0 \quad (\text{A85})$$

whose geometric representation is presented in Fig.11(2) with $a = 1, b = 2, \{c, d\} = \{3, 5\}$.

The corresponding exponents of the integer lattice $-n_1 E_4^{(1)} + n_2 E_4^{(4)}$ are presented by

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 2 - \frac{D}{2}, \alpha_{2,3} = 0, \alpha_{3,2} = \frac{D}{2} - 1, \alpha_{3,3} = 0, \quad (\text{A86})$$

whose geometric representation is drawn in Fig.11(1) with $(a, b) = (3, 5)$, $\{c, d\} = \{1, 2\}$.

Finally the exponents of the integer lattice $-n_1 E_4^{(2)} + n_2 E_4^{(4)}$ are

$$\alpha_{1,2} = -1, \alpha_{1,3} = 0, \alpha_{2,2} = 0, \alpha_{2,3} = 2 - \frac{D}{2}, \alpha_{3,2} = 1, \alpha_{3,3} = \frac{D}{2} - 2, \quad (\text{A87})$$

whose geometric representation is plotted in Fig.11(4) with $\{(a, b), (c, d)\} = \{(1, 2), (3, 5)\}$.

The corresponding hypergeometric functions are evidently written as

$$\psi_{\{1,5,6,7\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,4,5,6\}}^{(i)}(p_1^2, p_2^2, p_3^2), \quad (i = 1, \dots, 12). \quad (\text{A88})$$

Similarly $\det(A_{2,5,6,7}^{(1T)'}) = p_1^2 - p_2^2$, and

$$\left(A_{\{2,5,6,7\}}^{(1T)'} \right)^{-1} \cdot A^{(1T)'} = \begin{pmatrix} -\frac{p_3^2 - p_1^2}{p_2^2 - p_1^2} & 1 & -\frac{p_2^2 - p_3^2}{p_2^2 - p_1^2} & 0 & 0 & 0 & 0 \\ \frac{1}{p_2^2 - p_1^2} & 0 & -\frac{1}{p_2^2 - p_1^2} & 0 & 1 & 0 & 0 \\ -\frac{p_1^2}{p_2^2 - p_1^2} & 0 & \frac{p_2^2}{p_2^2 - p_1^2} & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}. \quad (\text{A89})$$

Obviously the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & 0 & \alpha_{1,3} & 0 & 0 & 0 & 0 \\ \alpha_{2,1} & 0 & \alpha_{2,3} & 0 & \frac{D}{2} - 3 & 0 & 0 \\ \alpha_{3,1} & 0 & \alpha_{3,3} & 3 - D & 0 & \frac{D}{2} - 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad (\text{A90})$$

where the matrix elements satisfy the relations which are obtained by the replacement $\alpha_{i,2} \rightarrow \alpha_{i,1}$, with $(i = 1, 3, 4)$ in Eq.(A75). Correspondingly the geometric representations of those exponent matrices are obtained from that of exponents of $\psi_{\{1,4,5,6\}}^{(i)}$ through the permutation $(\widehat{12})$. The constructed hypergeometric functions are

$$\psi_{\{2,5,6,7\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,4,5,6\}}^{(i)}(p_1^2, p_3^2, p_2^2), \quad (i = 1, \dots, 12). \quad (\text{A91})$$

Because $\det(A_{3,5,6,7}^{(1T)'}) = p_2^2 - p_3^2$, and

$$\left(A_{\{3,5,6,7\}}^{(1T)'}\right)^{-1} \cdot A_{\{3,5,6,7\}}^{(1T)'} = \begin{pmatrix} -\frac{p_1^2 - p_3^2}{p_2^2 - p_3^2} & -\frac{p_2^2 - p_1^2}{p_2^2 - p_3^2} & 1 & 0 & 0 & 0 & 0 \\ \frac{1}{p_2^2 - p_3^2} & -\frac{1}{p_2^2 - p_3^2} & 0 & 0 & 1 & 0 & 0 \\ -\frac{p_3^2}{p_2^2 - p_3^2} & \frac{p_2^2}{p_2^2 - p_3^2} & 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \end{pmatrix}, \quad (\text{A92})$$

the matrix of exponents is written as

$$\begin{pmatrix} \alpha_{1,1} & \alpha_{1,2} & 0 & 0 & 0 & 0 & 0 \\ \alpha_{3,1} & \alpha_{3,2} & 0 & 0 & \frac{D}{2} - 3 & 0 & 0 \\ \alpha_{4,1} & \alpha_{4,2} & 0 & 3 - D & 0 & \frac{D}{2} - 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad (\text{A93})$$

where the matrix elements satisfy the relations which are obtained by the replacement $\alpha_{i,3} \rightarrow \alpha_{i,1}$, with $(i = 1, 3, 4)$ in Eq.(A75). Correspondingly the geometric representations of those exponent matrices are obtained from that of exponents of $\psi_{\{1,4,5,6\}}^{(i)}$ through the permutation $(\widehat{132})$. The constructed hypergeometric functions are

$$\psi_{\{3,5,6,7\}}^{(i)}(p_1^2, p_2^2, p_3^2) = \psi_{\{1,4,5,6\}}^{(i)}(p_3^2, p_1^2, p_2^2), \quad (i = 1, \dots, 12). \quad (\text{A94})$$

-
- [1] CMS Collaboration, S. Chatrchyan et al., *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, published in Phys. Lett. **B716**(2012)30.
- [2] ATLAS Collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC* published in Phys. Lett. **B716**(2012)1.

- [3] Muon g-2 Collaboration, B. Abi et al., *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, published in Phys. Rev. Lett. **126**(2021)141801.
- [4] CDF Collaboration, T. Aaltonen et al., *High-precision measurement of the W boson mass with the CDF II detector* published in Science **376**(2022)170.
- [5] G. Heinrich, *Collider Physics at the Precision Frontier*, published in Phys. Rept. **922**(2021)1-69.
- [6] G. t'Hooft, M. Veltman, *Scalar One Loop Integrals*, published in Nucl. Phys. B **153**(1979)365.
- [7] J. Collins, *Renormalization*, Cambridge Univ. Press, 1984.
- [8] T. Regge, *Algebraic Topology Methods in the Theory of Feynman Relativistic Amplitudes*, In: *Battelle Rencontres - 1967 Lectures in Math. and Phys.*, edited by C. M. DeWitt and J. A. Wheeler, 1967, pp. 433-458.
- [9] E. Nasrollahpoursamami, *Periods of Feynman Diagrams and GKZ D-modules*, arXiv:1605.04970[math-ph].
- [10] I. M. Gel'fand, *General theory of hypergeometric functions*, published in Soviet Math. Dokl. **33**(1986)573.
- [11] I. M. Gel'fand, M. I. Graev, and A. V. Zelevinsky, *Holonomic systems of equations and series of hypergeometric type*, published in Soviet Math. Dokl. **36**(1988)5.
- [12] I. M. Gel'fand, A. V. Zelevinsky, and M. M. Kapranov, *Hypergeometric functions and toral manifold*, published in Soviet Math. Dokl. **37**(1988)678.
- [13] I. M. Gel'fand, M. M. Kapranov, and A. V. Zelevinsky, *Generalized Euler integrals and A-hypergeometric systems*, published in Adv. in Math. **84**(1990)255.
- [14] I. M. Gelfand, A. V. Zelevinskii, and M. M. kapranov, *Hypergeometric functions and toric varieties*, published in Functional. Anal. Appl. **23**(1989)94-106.
- [15] M. Kashiwara, T. Kawai, *Holonomic Systems of Linear Differential Equations and Feynman Integrals*, Publications of the Research Institute for Math. Sci. **12**(1976)131.
- [16] E. E. Boos, and A. I. Davydychev, *A method for calculating vertex-type Feynman integrals*, published in Vestn. Mosk. Univ. **28**(1987)8.
- [17] A. I. Davydychev, and J. B. Tausk, *Two loop selfenergy diagrams with different masses and the momentum expansion*, published in Nucl. Phys. B **397**(1993)123.
- [18] A. I. Davydychev, *Explicit results for all orders of the epsilon expansion of certain massive*

- and massless diagrams*, published in Phys. Rev. D**61**(2000)087701.
- [19] E. E. Boos, and A. I. Davydychev, *A Method of evaluating massive Feynman integrals*, published in Theor. Math. Phys. **89**(1991)1052.
- [20] A. I. Davydychev, *Recursive algorithm of evaluating vertex type Feynman integrals*, published in J. Phys. A**25**(1992)5587.
- [21] A. I. Davydychev, *Geometrical methods in loop calculations and the three-point function*, published in Nucl. Instrum. Meth. A**559**(2006)293.
- [22] A. I. Davydychev, *General results for massive N point Feynman diagrams with different masses*, published in J. Math. Phys.**33**(1992)358.
- [23] A. I. Davydychev, *Some exact results for N point massive Feynman integrals*, published in J. Math. Phys.**32**(1991)1052.
- [24] N. I. Ussyukina, and A. I. Davydychev, *An Approach to the evaluation of three and four point ladder diagrams*, published in Phys. Lett. B**298**(1993)363.
- [25] O. V. Tarasov, *Application and explicit solution of recurrence relations with respect to space-time dimension*, published in Nucl. Phys. B (Proc. Suppl)**89**(2000)237.
- [26] J. Fleischer, F. Jegerlehner, and O. V. Tarasov, *A New hypergeometric representation of one loop scalar integrals in d dimensions*, published in Nucl. Phys. B**672**(2003)303.
- [27] M. Yu. Kalmykov, B. A. Kniehl, *Towards all-order Laurent expansion of generalized hypergeometric functions around rational values of parameters*, published in Nucl. Phys. B**809**(2009)365.
- [28] V. V. Bytev, M. Yu. Kalmykov, B. A. Kniehl, *Differential reduction of generalized hypergeometric functions from Feynman diagrams: One-variable case*, published in Nucl. Phys. B**836**(2010)129.
- [29] M. Yu. Kalmykov, B. A. Kniehl, *Counting master integrals: Integration by parts versus differential reduction*, published in Phys. Lett. B**702**(2011)268.
- [30] V. V. Bytev, M. Yu. Kalmykov, B. A. Kniehl, *HYPERDIRE, HYPERgeometric functions Differential REduction: MATHEMATICA-based packages for differential reduction of generalized hypergeometric functions ${}_pF_{p-1}$, F_1 , F_2 , F_3 , F_4* , published in Comput. Phys. Commun**184**(2013)2332.
- [31] V. Bytev, M. Kalmykov, *HYPERDIRE hypergeometric functions differential reduction:*

- Mathematica-based packages for the differential reduction of generalized hypergeometric function: Horn-type hypergeometric functions of two variables*, published in *Comput. Phys. Commun***189**(2015)128.
- [32] V. V. Bytev, M. Yu. Kalmykov, *HYPERDIRE-HYPERgeometric functions Differential REDuction: Mathematica-based packages for the differential reduction of generalized hypergeometric functions: Lauricella function F_c of three variables*, published in *Comput. Phys. Commun***206**(2016)78.
- [33] M. Y. Kalmykov, and B. A. Kniehl, *Mellin-Barnes representations of Feynman diagrams, linear systems of differential equations, and polynomial solutions*, published in *Phys. Lett. B***714**(2012)103.
- [34] M. Y. Kalmykov, and B. A. Kniehl, *Counting the number of master integrals for sunrise diagrams via the Mellin-Barnes representation*, published in *JHEP***1707**(2017)031.
- [35] L. Cruz, *Feynman integrals as A-hypergeometric functions*, published in *JHEP***1912**(2019)123 (arXiv:1907.00507 [math-ph]).
- [36] R. Klausen, *Hypergeometric Series Representations of Feynman integrals by GKZ Hypergeometric Systems*, published in *JHEP***2004**(2020)121 (arXiv:1910.08651 [hep-th]).
- [37] R. N. Lee, A. A. Pomeransky, *Critical points and number of master integrals*, published in *JHEP***1311**(2013)165.
- [38] T. Oaku, *Adv. in Appl. Math.* **19**(1997)61.
- [39] U. Walther, *J. of Pure and Applied Algebra.* **139**(1999)303.
- [40] T. Oaku, N. Takayama, *J. of Pure and Applied Algebra.* **156**(2001)267.
- [41] T.-F. Feng, C.-H. Chang, J.-B. Chen, and H.-B. Zhang, *GKZ-hypergeometric systems for Feynman integrals*, published in *Nucl. Phys. B***953**(2020)114952.
- [42] T.-F. Feng, C.-H. Chang, J.-B. Chen, Z.-H. Gu, and H.-B. Zhang, *Evaluating Feynman integrals by the hypergeometry*, published in *Nucl. Phys. B***927**(2018)516.
- [43] T.-F. Feng, C.-H. Chang, J.-B. Chen, and H.-B. Zhang, *The system of partial differential equations for the C_0 function*, published in *Nucl. Phys. B***940**(2019)130.
- [44] W. Miller Jr., *J. Math. Mech.* **17**(1968)1143.
- [45] W. Miller Jr., *SIAM. J. Math. Anal.* **3**(1972)31.
- [46] F. Loebbert, D. Müller, and H. Münkler, *Yangian Bootstrap for Conformal Feynman Integrals*,

- published in Phys. Rev. D **101**(2020)066006.
- [47] A. Klemm, C. Nega, and R. Safari, *The l -loop Banana Amplitude from GKZ Systems and relative Calabi-Yau Periods*, published in JHEP **2004**(2020)088.
- [48] K. Bönisch, F. Fischbach, A. Klemm, C. Nega, and R. Safari, *Analytic Structure of all Loop Banana Amplitudes*, published in JHEP **2105**(2021)066.
- [49] T. Reichelt, M. Schulze, C. Sevenheck, and U. Walther, *Algebraic aspects of hypergeometric differential equations*, e-Print: 2004.07262[math.AG].
- [50] M. Hidding, *DiffExp, a Mathematica package for computing Feynman integrals in terms of one-dimensional series expansions*, published in Comput. Phys. Commun. **269**(2021)108125.
- [51] M. Borinsky, *Tropical Monte Carlo quadrature for Feynman integrals*, arXiv: 2008.12310[math-ph].
- [52] F. Tellander, M. Helmer, *Cohen-Macaulay Property of Feynman Integrals*, arXiv: 2108.01410[hep-th].
- [53] S. Mizera, and S. Telen, *Landau Discriminants*, arXiv: 2109.08036[math-ph].
- [54] N. Arkani-Hamed, A. Hillman, and S. Mizera, *Feynman Polytopes and the tropical geometry of UV and IR divergence*, published in Phys. Rev. D **105**(2022)125013.
- [55] V. Chestnov, F. Gasparotto, M. K. Mandal, P. Mastrolia, S. J. Matsubara-Heo, H. J. Munch, and N. Takayamac, *Macaulay Matrix for Feynman Integrals: Linear Relations and Intersection Numbers*, arXiv: 2204.12983[hep-th].
- [56] B. Ananthanarayan, S. Bera, S. Friot, T. Pathak, *Olsson.wl: a Mathematica Mathematica package for the computation of linear transformations of multivariable hypergeometric functions*, arXiv: 2201.01189[cs.MS].
- [57] F. A. Berends, M. Böhm, M. Buza, and R. Scharf, Z. Phys. C **63**(1994)227.
- [58] B. Ananthanarayan, S. Friot, and S. Ghosh, *New series representations for the two-loop massive sunset diagram*, published in Eur. Phys. J. C **80**(2020)606.
- [59] M. Kalmykov, V. Bytev, B. Kniehl, S.-O. Moch, B. Ward, and S. Yost, *Hypergeometric Functions and Feynman Diagrams*, arXiv:2012.14492 [hep-th].
- [60] J. Oxley, *Matroid Theory*, (Oxford Univ. Press 2011).
- [61] I. M. Gelfand, *General theory of hypergeometric functions*, published in Sov. Math. Dokl. **33**(1986)573-577.

- [62] J. T. Schwartz, *Differential Geometry and Topology*, Gordon and Breach Science Publishers, 1968.
- [63] I. M. Gelfand, S. G. Gindikin, and M. I. Graev, *Integral geometry in affine and projective spaces*, published in J. Sov. Math. **18**(1980)39-167.
- [64] I. M. Gelfand, and S. I. Gelfand, *Generalized hypergeometric equations*, published in Sov. Math. Dokl. **33**(1986)643-646.
- [65] I. M. Gelfand, and M. I. Graev, *Hypergeometric functions associated with the Grassmanian $G_{3,6}$* , published in Math.USSR. Sb. **66**(1990)1-40.
- [66] I. M. Gelfand, M. I. Graev, and V. S. Retakh, *General hypergeometric systems of equations and series of hypergeometric type*, published in Russ. Math. Surveys **47:4**(1992)1-88.
- [67] J. Horn, Math. Ann. **34**(1889)544.
- [68] M. Saito, B. Sturmfels, N. Takayama, *Gröbner Deformations of Hypergeometric Differential Equations*, Springer 2000.
- [69] M. E. Taylor, *Partial differential equations* (Springer, Heidelberg 2012).
- [70] L. J. Slater, *Generalised Hypergeometric Functions* (Cambridge University Press 1966).
- [71] J. Leray, *Le Calcul Différentiel et Intégral sur Une Variété Analytique Complexe*, Bull. Soc. Math. **87**(1959)81.
- [72] D. Cox, J. Little, D. O'Shea, *Ideals, Varieties and Algorithms* (Springer, New York 1991).
- [73] D. Cox, J. Little, D. O'Shea, *Using Algebraic Geometry* (Springer, New York 1998).
- [74] B. Sturmfels, *Gröbner Bases and Convex Polytopes*, University Lecture Notes, Vol. 8. American Mathematical Society, Providence 1995.
- [75] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry* (Springer, New York 1995).
- [76] S. C. Coutinho, *A Primer of Algebraic D-modules*, London Mathematical Society Student Text 33, Cambridge Univ. Press, Cambridge 1995.
- [77] K. G. Chetyrkin, A. L. Kataev, and F. V. Tkachov, *New Approach to Evaluation of Multiloop Feynman Integrals: The Gegenbauer Polynomial x Space Technique*, published in Nucl. Phys. **B174**(1980)345.
- [78] Z. H. Gu, H. B. Zhang, and T. F. Feng, *Hypergeometric expression for a three-loop vacuum integral*, published in Int. J. Mod. Phys. **A35**(2020)2050089.

- [79] V. A. Smirnov, *Analytic Tools for Feynman Integrals*, (Springer, Heidelberg 2012), and references therein.