

ON THE ERDŐS-TUZA-VALTR CONJECTURE

JINEON BAEK

ABSTRACT. The Erdős-Szekeres conjecture states that any set of more than 2^{n-2} points in the plane with no three on a line contains the vertices of a convex n -gon. Erdős, Tuza, and Valtr strengthened the conjecture by stating that any set of more than $\sum_{i=n-b}^{a-2} \binom{n-2}{i}$ points in a plane either contains the vertices of a convex n -gon, a points lying on a concave downward curve, or b points lying on a concave upward curve. They also showed that the generalization is actually equivalent to the Erdős-Szekeres conjecture.

We prove the first new case of the Erdős-Tuza-Valtr conjecture since the original 1935 paper of Erdős and Szekeres. Namely, we show that any set of $\binom{n-1}{2} + 2$ points in the plane with no three points on a line and no two points sharing the same x -coordinate either contains 4 points lying on a concave downward curve or the vertices of a convex n -gon.

1. INTRODUCTION

A set of n points on a plane is in a *convex position* if they form the vertices of a convex polygon. For simplicity, denote any set of n points in convex position as an *n -gon*. The well-known Erdős-Szekeres conjecture is the following. Here, a finite set of points on a plane is in *general position* if no three points are on a line and no two points share the same x -coordinate¹.

Definition 1.1. For any $n \geq 3$, let $N(n)$ be the maximum number of points on a plane in a general position with no subset forming an n -gon.

Conjecture 1 (Erdős-Szekeres [4]). For any $n \geq 3$, $N(n) = 2^{n-2}$.

In 1935, Erdős and Szekeres showed $N(n) \leq \binom{2n-4}{n-2} = O(4^n/\sqrt{n})$ [3]. In 1960, they provided a construction of exactly 2^{n-2} points with no n points in convex position, showing the lower bound $N(n) \geq 2^{n-2}$ [4]. The upper bound of $N(n)$ stayed in the magnitude of $O(4^n/\sqrt{n})$ despite many improvements [2, 8, 14, 15, 9, 11] until Suk [12] proved an upper bound of $N(n) \leq 2^{n+o(n)}$ in 2016. The best upper bound so far is $N(n) \leq 2^{n+O(\sqrt{n \log n})}$ [7], and the precise equality $N(n) = 2^{n-2}$ remains neither proven or disproven to this date.

Now we state the generalization of Conjecture 1 by Erdős, Tuza, and Valtr [5].

Definition 1.2. An *a -cap* (resp. *a -cup*) is a set of a points lying on the graph of a downwardly (resp. upwardly) convex function (see Figure 1).

Definition 1.3. Call any tuple (n, a, b) of integers satisfying $2 \leq a, b \leq n \leq a+b-2$ a *triplet*. For any triplet (n, a, b) , define $N(n, a, b)$ as the maximum number of points on a plane in general position with no subset forming an n -gon, a -cap, or b -cup.

¹It is not usually required for the x -coordinates to be different when stating the Erdős-Szekeres conjecture. However, the extra assumption does not hurt generality as we can rotate the point set slightly if there is any overlap in the x -coordinates.




FIGURE 1. A 4-cap, 5-cup and 6-gon (from left to right)

Conjecture 2 (Erdős-Tuza-Valtr [5]). For any triplet (n, a, b) , we have

$$N(n, a, b) = \sum_{i=n-b}^{a-2} \binom{n-2}{i}.$$

Definition 1.4. For any triplet (n, a, b) , let $P(n, a, b)$ be the statement that $N(n, a, b) = \sum_{i=n-b}^{a-2} \binom{n-2}{i}$. That is, $P(n, a, b)$ is the special case of Conjecture 1 with the triplet (n, a, b) .

Then the special case $P(n, n, n)$ of Conjecture 2 is Conjecture 1, as n -caps/cups are n -gons and the conjectured sum equals 2^{n-2} . Interestingly, they showed that the strengthened Conjecture 2 is actually *implied by* the original Conjecture 1 as well.

Theorem 1.5 ([5]). Conjecture 1 and Conjecture 2 are equivalent. Specifically, for any $n \geq 3$ and any triplets (n, a, b) , (n, a', b') with $a \geq a'$ and $b \geq b'$, the statement $P(n, a, b)$ implies $P(n, a', b')$.

So any counterexample of Conjecture 2 for one triplet (n, a, b) would disprove Conjecture 1 immediately. Indeed, such a line of attack [1] succeeded in disproving a set-theoretic generalization [13] of Conjecture 1 by disproving an analogous set-theoretic generalization of Conjecture 2 for the triplet $(n, 4, n)$.

It is then natural to ask whether the original Conjecture 2 will hold for the same triplet $(n, 4, n)$ or not. We show that it does.

Theorem 1.6 (Main Theorem). For any $n \geq 3$, the statement $P(n, 4, n)$ holds. That is, $\binom{n-1}{2} + 2$ points on a plane in general position determine either a 4-cap or a n -gon.

Related, Erdős and Szekeres proved the following *cups-caps theorem* in 1935 [3].

Theorem 1.7 (Erdős-Szekeres). For any $a, b \geq 2$, any set of more than $\binom{a+b-4}{a-2}$ points in general position contains either an a -cap or b -cup.

For any $a, b \geq 2$, the cases $P(a+b-3, a, b)$ and $P(a+b-2, a, b)$ of Conjecture 2 are immediate consequences of this theorem. To see this, observe that any $(a+b-3)$ -gon either contains an a -cap on top or a b -cup on the bottom, and that the conjectured values $N(a+b-3, a, b) = \binom{a+b-5}{a-3} + \binom{a+b-5}{a-2} = \binom{a+b-4}{a-2}$ and $N(a+b-2, a, b) = \binom{a+b-4}{a-2}$ match with the cups-caps theorem. Our Theorem 1.6 is not a consequence of the cups-caps theorem as $n < a+b-3$ in the triplet $(n, 4, n)$.

By Theorem 1.5, for triplets (n, a, b) with a fixed n , the cases $P(n, a, b)$ of Conjecture 2 form a pyramid of implications from top to bottom (Figure 2). On the top, we have the Erdős-Szekeres conjecture (Conjecture 1), and at the bottom, we have the cups-caps theorem (Theorem 1.7). Our theorem lies strictly in the middle of them (Theorem 1.6). To the best of our knowledge, this is the first new instance of Conjecture 2 proven so far since the original 1935 paper [3] of Erdős and Szekeres.




FIGURE 2. A schematic diagram of the implications between Conjecture 1, Theorem 1.7 and our Theorem 1.6.

The proof of main Theorem 1.6 generalizes to a purely combinatorial model of convexity (Theorem 2.7). Section 2 describes the combinatorial setup we mainly work with. Section 3 describes the α -statistic, a function for understanding a -cap, b -cup free configurations especially with nearly maximal number of points. Section 4 introduces (α, β) -plane, an useful notion for understanding 4-cap free configurations, and motivates the definition of *interweaved laced cups* (Definition 4.4 and 4.5). Finally, Section 5 proves main Theorem 1.6 by finding interweaved laced cups with induction (Theorem 5.10).

2. A COMBINATORIAL MODEL OF CONVEXITY

We introduce a purely combinatorial model of convexity that we mainly work with. It is a slight modification of the one first suggested in [13] and explored further in [6], [10] and [1].

Definition 2.1. In this paper, a *configuration* is a finite set S of elements called *points* or *vertices* equipped with the following structures.

- A linear ordering $<$ of the vertices.
- For any subset of S with size 3, an arbitrary assignment of whether they form a cap or a cup.

In standard terms, a configuration is a bi-coloring of a finite, complete 3-uniform hypergraph with a prescribed linear ordering of vertices.

Given a finite set of points in general position, we can make a configuration by ordering the points in their increasing x -coordinates and assigning each size 3 subset to be either a cap or cup according to its position in the plane. Let's say that

a configuration is *realizable* if it can be constructed from actual points in this way. The notion of caps and cups in plane are generalized to arbitrary configurations which may not be realizable.

Definition 2.2. Denote any set of vertices $x_1 < x_2 < \dots < x_a$ of a configuration as simply $x_1x_2 \dots x_a$ in the increasing order.

In a configuration, a set $C = x_1x_2 \dots x_a$ of a vertices forms an a -cup (resp. a -cap) if any three consecutive points $x_{i-1}x_ix_{i+1}$ form a cup (resp. cap) for any $1 < i < a$. In particular, we allow 1-cups and 1-caps. The *size* of C , also denoted $|C|$, is the number of vertices a in C .

We say that x_1 is the *starting point* of C and x_a is the *ending point* of C . The points x_1 and x_a are the *endpoints* of C . Equivalently, we say that C starts with x_1 and ends with x_a , or that C is a cap (resp. cup) from x_1 to x_a . Call any pair $x < y$ of vertices in a configuration an *edge*. If the size a of C is at least 2, we say that C *starts* with the edge x_1x_2 and *ends* with the edge $x_{a-1}x_a$.

For a cap (resp. cup) C from vertex s to vertex t , say that C *extends to left* with the vertex x (or edge xs) if xC is also a cap (resp. cup). Likewise, say that C *extends to right* with the vertex x (or edge tx) if Cx forms a cap (resp. cup).

We also introduce the *mirror reflection* of a configuration. For realizable configurations, this corresponds to reflecting the points along the y -axis.

Definition 2.3. The *mirror reflection* S^{op} of a configuration S is the configuration with the same vertex set and assignments of 3-caps and 3-cups, but the prescribed linear ordering reversed. In this way, a cap (resp. cup) C in S naturally corresponds to the reflected cap (resp. cup) C^{op} in S^{op} with the same set of vertices.

We introduce two possible generalizations of an n -gon in this combinatorial model of convexity.

Definition 2.4. In a configuration, a *weak* (a, b) -gon is a pair of a -cap C_1 and b -cup C_2 sharing the same endpoints x and y . An *weak* n -gon is any weak (a, b) -gon with $a + b = n + 2$.

On the other hand, a *strong* (a, b) -gon is a weak (a, b) -gon (C_1, C_2) with the additional constraint that $C_1 \cap C_2 = \{x, y\}$. An *strong* n -gon is any strong (a, b) -gon with $a + b = n + 2$.

An (a, b) -gon or n -gon denotes the weak (a, b) -gon or n -gon by default.

The only difference is that a weak n -gon allows the cap and cup to have overlapping vertices in the middle, while a strong n -gon does not. Note also that there is no difference in any realizable configurations. We use the following notion.

Definition 2.5. A configuration is a -cap (resp. b -cup or weak/strong n -gon) free if it has no a -cap (resp. b -cup or weak/strong n -gon) in the configuration.

Peters and Szekeres [13] proposed to generalize Conjecture 1 by stating that for any $n \geq 2$, the maximum size of a *strong* n -gon free configuration is 2^{n-2} , and supplied a computer proof for the case $n = 6$. Later, Balko and Valtr [1] found counterexamples for $n = 7$ and 8 with SAT solvers by finding counterexamples for analogues of Conjecture 2. We instead propose to use the definition of a *weak* n -gon instead to generalize Conjecture 1.

Conjecture 3 (Set-theoretic Erdős-Szekeres). For any $n \geq 2$, the maximum size $\widehat{N}(n)$ of a *weak* n -gon free configuration is equal to 2^{n-2} .

From now on, we omit the word *weak* when we mention any (a, b) -gon or n -gon. We can also state the analogous generalization of Conjecture 2 to arbitrary configurations.

Conjecture 4 (Set-theoretic Erdős-Tuza-Valtr). For any triplet (n, a, b) , the maximum size $\widehat{N}(n, a, b)$ of a weak n -gon, a -cap and b -cup free configuration is equal to $\sum_{i=n+2-b}^a \binom{n-2}{i-2}$.

The generalization of cups-caps theorem (Theorem 1.7) to arbitrary configurations is well-known (e.g. [10]).

Theorem 2.6 (Set-theoretic cups-caps). For any $a, b \geq 2$, the maximum size of an a -cap and b -cup free configuration is $\binom{a+b-4}{a-2}$.

Our main result is the proof of Conjecture 4 for the case $(n, a, b) = (n, 4, n)$. It is the generalization of Theorem 1.6 to arbitrary configurations.

Theorem 2.7. For any $n \geq 3$, any configuration of size $\binom{n-1}{2} + 2$ contains either a 4-cap, n -cup or a $(3, n-1)$ -gon.

3. STRUCTURE OF a -CAP, b -CUP FREE CONFIGURATIONS

In this section, we discuss some properties of a configuration S that avoids a -caps and b -cups. They will be useful especially when the size of S is close to the maximum possible value $\binom{a+b-4}{a-2}$ shown in Theorem 2.6. We do not claim originality of the results in this section - it is mostly a recasting of the definitions introduced in [10].

First, we define the *slope labeling* of an a -cap free configuration S .

Definition 3.1. A *slope labeling* s of an a -cap free configuration S is the assignment of an integer $s(xy) \in \{1, 2, \dots, a-2\}$ to all the edges xy of S so that it satisfies the following.

- For any points $x < y < z$ in S , $s(xy) \leq s(yz)$ implies that xyz is a 3-cup.

The value $s(xy)$ assigned to the edge xy is the *label* of xy .

For the actual slope $s_{\mathbb{R}}(xy) \in \mathbb{R}$ of an edge in a realizable configuration S , both the following properties would hold.

- For any points $x < y < z$ in S , $s_{\mathbb{R}}(xy) > s_{\mathbb{R}}(yz)$ implies that xyz is a 3-cap.
- For any points $x < y < z$ in S , $s_{\mathbb{R}}(xy) < s_{\mathbb{R}}(yz)$ implies that xyz is a 3-cup.

A slope labeling restricts the possible values of a ‘slope’ (label) to a much smaller set $\{1, 2, \dots, a-2\}$ at the cost of giving up the first property. That is, edges of strictly decreasing labels may not form a cap (see the cup ABF in Figure 3). It is also impossible in general to assign a slope labeling that satisfies both the properties. However, any a -cap free configuration has a slope labeling that satisfies the second property.

Theorem 3.2. For any a -cap free configuration S , a slope labeling always exists. In particular, for any edge e , let $c(e)$ be the maximum length of a cap starting with e ; the function $c(e) - 1$ is a slope labeling.

Proof. For any edge e , its assigned label i is in between 1 and $a-2$ inclusive by definition. Assume by contrary that there is a 3-cap xyz in S with the label s of xy less than or equal to the label t of yz . As the label of edge yz is t , there is a cap

C of size $t + 1$ starting with yz . Since xyz forms a cap, the cap C extends to a cap xC of size $t + 2$ that starts with xy . By the definition of s , we have $s + 1 \geq t + 2$, which contradicts the hypothesis $s \leq t$. \square




FIGURE 3. A 4-cap and 4-cup free realizable configuration with the slope labeling of Theorem 3.2. The labels of the edges AF , BD , CD and CE are 1, 2, 2 and 2 respectively.

For the rest of the paper, we will fix a large a -cap free configuration S and a slope labeling of S so that there is no confusion in the label of an arbitrary edge of S . The following lemma is immediate.

Lemma 3.3. For any a -cap free configuration S with slope labeling s , the restriction of s to an arbitrary subset S' of S is a slope labeling of S' .

Assume any a -cap, b -cup free configuration S with a slope labeling s by integers from 1 to $a - 2$. For each point p of S , we assign its α -statistic which is a tuple $\alpha(p) = (\alpha_1(p), \alpha_2(p), \dots, \alpha_{a-2}(p))$ of natural numbers.

Definition 3.4. Fix any a -cap, b -cup free configuration S with a slope labeling. For all $p \in S$ and $1 \leq i \leq a - 2$, define the integer $\alpha_i(p)$ as maximum length of a cup that ends with the point p and an edge of slope $\leq i$. If there is no such cup, then let $\alpha_i(p) = 1$. In particular, the rightmost point p has the value $\alpha^i(p) = 1$ for any i . The α -statistic of a point p in S is the tuple $\alpha(p) = (\alpha_1(p), \alpha_2(p), \dots, \alpha_{a-2}(p))$.

Definition 3.5. For any integers $a, b \geq 2$, define the grid simplex

$$T_{a,b} := \{(x_1, x_2, \dots, x_{a-2}) \in \mathbb{N}^{a-2} : 1 \leq x_1 \leq \dots \leq x_{a-2} \leq b - 1\}.$$

Theorem 3.6. For any a -cap, b -cup free configuration S with a fixed slope labeling s , the α -statistic of S satisfies the following.

- (1) For any point p , $1 \leq \alpha_1(p) \leq \dots \leq \alpha_{a-2}(p) \leq b - 1$. Consequently, the α -statistic α is a map from S to $T_{a,b}$.

- (2) For any two points $x < y$ connected by an edge of label i , $\alpha_i(x) < \alpha_i(y)$ holds. So in particular, α is injective.

Proof. We check the two conditions the α -statistic has to satisfy. The inequality $1 \leq \alpha_1(p) \leq \dots \leq \alpha_{a-2}(p) \leq b - 1$ for any point p follows directly from the definitions. The cup of length $\alpha_i(x)$ that ends with the vertex x and an edge of label $\leq i$ can be extended to the right with the edge xy of label i , so that $\alpha_i(x) + 1 \leq \alpha_i(y)$ in any of the two given definitions of α_i . \square

Note that the size of $T_{a,b}$ is $\binom{a+b-4}{a-2}$. For a quick proof, note that any $(x_1, x_2, \dots, x_{a-2}) \in T_{a,b}$ corresponds bijectively to an arbitrary subset $\{x_i + i - 1 : 1 \leq i \leq a - 2\}$ of $\{1, 2, \dots, a + b - 4\}$ of size $a - 2$. Consequently, α being injective immediately proves Theorem 2.6 that $|S| \leq \binom{a+b-4}{a-2}$. Therefore, if the size of S is nearly equal to the maximum size $\binom{a+b-4}{a-2}$, we can expect the map α to be almost bijective with some exceptional ‘holes’ in $T_{a,b}$.

Let us end this section with a remark on the α -statistic of the mirror configuration.

Definition 3.7. For any a -cap, b -cup free configuration S and its slope labeling s , define the mirror reflection s^{op} of the slope labeling s as the following.

$$s^{\text{op}}(yx) = a - 1 - s(xy)$$

We leave it as an exercise to show that s^{op} is a proper slope labeling of S^{op} .

Lemma 3.8. The mirror reflection s^{op} of a slope labeling s of S is a slope labeling of S^{op} .

Remark 3.9. We use the mirror reflection of a labeling extensively to exploit the symmetry without loss of generality. For example, take an arbitrary 4-cap free configuration S . Its slope labeling s has values in $\{1, 2\}$. So for any edge xy , if its label $s(xy)$ is 1 (resp. 2) then the labeling $s^{\text{op}}(yx)$ of its reflection is 2 (resp. 1). Therefore, if the statement we want to show regarding S is invariant under reflection, then we can safely assume that the edge connecting x and y is labeled 1 by reflecting S if necessary.

4. THE (α, β) -PLANE

In this section, we focus our attention on an arbitrary 4-cap, n -cup free configuration S with a fixed slope labeling and α -statistic.

In our configuration S , we only have edges of label 1 and 2, and the following is an immediate consequence of Definition 3.1. We will use this extensively without further mentions.

Corollary 4.1. In an arbitrary 4-cap free configuration with slope labeling, any edge of label 1 extends a cup to the left. That is, for any edge pq of label 1 and a cup C that starts with q , the sequence pC is also a cup.

Likewise, any edge of label 2 extends a cup to the right. That is, for any edge pq of label 2 and a cup C that ends with p , the sequence Cq is also a cup.

Also, we define aliases for the α -statistic of S .




FIGURE 4. The (α, β) -plane of the 4-cap, 4-cup free configuration in Figure 3 with the slope labels of some edges.

Definition 4.2. For an arbitrary 4-cap, n -cup free configuration S with a fixed slope labeling and α -statistic $p \mapsto (\alpha_1(p), \alpha_2(p))$, define aliases $\alpha = \alpha_1$ and $\beta = \alpha_2$.

Thus, $\alpha(p) = \alpha_1(p)$ is the maximum length of a cup that ends with the vertex p and an edge of label 1 ($\alpha(p) = 1$ if there is no such cup). The value $\beta(p) = \alpha_2(p)$ is the maximum length of any cup that ends with the vertex p .

The α -statistic $\alpha(p) = (\alpha(p), \beta(p))$ maps S to the triangular grid set

$$T_{4,n} := \{(a, b) \in \mathbb{N}^2 : 1 \leq a \leq b \leq n-1\}$$

injective by Theorem 3.6. With this, if the size of S is $|T_{4,n}| - k = \binom{n-1}{2} - k$ where k is small, it helps to identify S with the grid points $T_{4,n}$ with k missing holes (see Figure 4). Call such a diagram an (α, β) -plane of the 4-cap, n -cup free configuration S . The following is a direct consequence of Theorem 3.6, and we will use it extensively without mentioning.

Corollary 4.3. Assume an arbitrary 4-cap, n -cup free configuration S .

For any points p, q with $\beta(p) = \beta(q)$,

- p and q are always connected with an edge of label 1
- and $p < q$ if and only if $\alpha(p) < \alpha(q)$.

Consequently, in an (α, β) -plane any horizontal edge is labeled 1, and each column is sorted in the increasing order of vertices from left to right.

Likewise, for any points p, q with $\alpha(p) = \alpha(q)$,

- (1) p and q are always connected with an edge of label 2
- (2) and $p < q$ if and only if $\beta(p) < \beta(q)$.

Consequently, in an (α, β) -plane any vertical edges is labeled 2, and each row is sorted in the increasing order of vertices from bottom to top.

We will now use the concept of the (α, β) -plane to introduce and motivate several key notions for our proof. Take a 4-cap, n -cup free configuration S . For simplicity, assume at first that S of size $\binom{n}{2}$ (the maximum possible size), so that every location in the (α, β) -plane is occupied. Then we can identify a point p in S with its location $(\alpha(p), \beta(p))$ in the (α, β) -plane.

Take any $1 \leq k \leq n - 1$ and consider the path of vertices

$$C_k = (1, k), (2, k), \dots, (k - 1, k), (k, k), (k, k + 1), \dots, (k, n - 1)$$

in the (α, β) -plane. By Corollary 4.3, the first $k - 1$ edges in this path are labeled 1, and the last $n - k - 1$ edges are labeled 2, so these points must form an $(n - 1)$ -cup C_k from start $p_k = (1, k)$ to end $q_k = (k, n - 1)$. For any pair of $(n - 1)$ -cups C_k and C_l with $k < l$, the ordering of their endpoints is then $p_k < p_l \leq q_k < q_l$. Motivated by this, we introduce the following definition of *interweaved cups*.

Definition 4.4. Two cups C_1 and C_2 running from p to r and q to s respectively are *interweaved* if $p < q \leq r < s$ holds.

So if $|S| = \binom{n}{2}$, we get $n - 1$ different $(n - 1)$ -cups C_1, \dots, C_{n-1} all mutually interweaved. Moreover, for each cup C_k we have the k -cup D_k of vertices

$$D_k = (1, 1), (1, 2), \dots, (1, k)$$

that ends with the left endpoint p_k of C_k and the $(n - k)$ -cup E_k of vertices

$$E_k = (k, n - 1), (k + 1, n - 1), \dots, (n - 1, n - 1)$$

that starts with the right endpoint q_k of C_k . Motivated by this, we introduce the following notion of *laced cups*.

Definition 4.5. A $(n - 1)$ -cup C from p to q is *laced* if there exists a cup C_p that ends with p , and a cup C_q that starts with q , so that $|C_p| + |C_q| = n - 1$.²

So if $|S| = \binom{n}{2} = \binom{n-1}{2} + (n-1)$, then all of our mutually interweaved $(n - 1)$ -cups C_k are laced. The importance of this concept will be shown in Lemma 5.2 which shows that if S contains just *two* interweaved laced $(n - 1)$ -cups, then S contains a $(3, n - 1)$ -gon. We expect that for $|S| = \binom{n-1}{2} + d$ with any $1 \leq d \leq n - 1$, we can find d mutually interweaved laced $(n - 1)$ -cups (Conjecture 5). Roughly, this amounts to saying that an additional hole in the (α, β) -plane only destroys one of the mutually interweaved laced $(n - 1)$ -cups.

We show this for $d = 2$ (Theorem 5.10) and this is sufficient to show the main theorem that for $|S| = \binom{n-1}{2} + 2$ we can always find an n -gon (Theorem 2.7). The proof for case $d = 2$ requires a delicate inductive argument and several lemmas about interweaved laced $(n - 1)$ -cups; we now turn to stating and proving those lemmas. Before doing so, we briefly remark that the concept of interweaved cups and of laced cups are symmetric under mirror reflection.

Lemma 4.6. Two cups C_1, C_2 are interweaved in configuration S if and only if C_2^{op} and C_1^{op} are interweaved in S^{op} . An $(n - 1)$ -cup C is laced in configuration S if and only if C^{op} is laced in S^{op} .

²One might expect the sum to be n given the above motivation, but it turns out that $n - 1$ is sufficient.




FIGURE 5. Figure for the proof of Lemma 5.1. The cup P ends at a and the cup Q is from b to c .




FIGURE 6. Figure for the proof of Lemma 5.2. The cup C_p ends with p and C_r starts with r .

5. INTERWEAVED LACED CUPS

We first show that in a 4-cap free configuration, a pair of interweaved laced $(n-1)$ -cups from p to r and q to s respectively (so that $p < q \leq r < s$) is sufficient to force an n -gon. The following covers the degenerate case $q = r$.

Lemma 5.1 (Balko and Valtr [1]). Take any $n \geq 3$ and a 4-cap, n -cup free configuration S . If the ending point x of an $(n-1)$ -cup C_1 is also the starting point of another $(n-1)$ -cup C_2 , then S contains an $(3, n-1)$ -gon.

Proof. (See Figure 5) Say $C_1 = Px$ where P is an $(n-2)$ -cup that ends with some vertex a . Likewise, say that $C_2 = xQ$ where Q is an $(n-2)$ -cup that starts with b . Note that the statement to prove and the definitions introduced so far are symmetric under reflection. So without loss of generality, we can assume that ab is labeled 1. Now aQ is a $(n-1)$ -cup because ab has label 1. Say that Q ends with the point c .

If axc is a cap, then we find the $(3, n-1)$ -gon formed by axc and aC_2' and we are done. If axc is a cup, then the $(n-1)$ -cup C_1 extends to the right with vertex c , contradicting that S is n -cup free. \square

Now we show the general case $q \leq r$.

Lemma 5.2. Take any 4-cap, n -cup free configuration S where $n \geq 3$. If S contains a pair of interweaved laced $(n-1)$ -cups, then S contains an $(3, n-1)$ -gon.

Proof. (See Figure 6) Let C_1 and C_2 be the pair of interweaved laced $(n-1)$ -cups from p to r and q to s respectively, so that $p < q \leq r < s$. If $q = r$, we are done by Lemma 5.1, so we can assume $q < r$. As the setup is symmetric along reflection




FIGURE 7. Figure for the proof of Lemma 5.3. The cups P , Q and R contains the endpoints a , b and c , and d respectively.

(e.g. Remark 3.9 and 4.6), we can assume that the edge qr is of label 1 without loss of generality.

As C_1 is laced we have a cup C_p ending with p and a cup C_r starting with r so that $|C_p| + |C_r| = n - 1$. Observe that the edge pq is of label 2, or otherwise we can extend the $(n - 1)$ -cup C_2 to left with pq and reach contradiction.

If pqr is a cap, then the cap pqr and the cup C_1 forms a $(3, n - 1)$ -gon and we are done. Now assume that pqr is a cup. Since pq is of label 2, the cup C_p extends to right with q . Since qr is of label 1, the cup C_r extends to left with q as well. Now the cups C_pq and qC_r are joined along the vertex q . Since pqr is a cup, they all connect to make a cup C_pqC_r of size $|C_p| + |C_r| + 1 = n$, leading to contradiction. \square

We prepare more lemmas and observations before starting the main proof. In particular, we will use the following special case to find a pair of interweaved laced cups. Note that for this lemma we have $n \geq 4$ instead of $n \geq 3$.

Lemma 5.3. Take any 4-cap, n -cup free configuration S with $n \geq 4$. Assume an $(n - 1)$ -cup C from x to y in S . Also, assume an $(n - 2)$ -cup C_x that ends with x , and an $(n - 2)$ -cup C_y that starts with y . Then S contains a pair of interweaved laced $(n - 1)$ -cups.

Proof. Using the symmetry of the statement along mirror reflection, we can assume that xy is labeled 1 without loss of generality.

Let $C_x = Px$, $C = xQy$ and $C_y = yR$ so that P, Q, R are $(n - 3)$ -cups (see Figure 7). Let a be the ending point of P and d be the starting point of R . Say that Q is from point b to c . As $|P|, |Q|, |R| \geq 1$, the points a, b, c and d are all well-defined and we have $a < x < b \leq c < y < d$ (there is a possibility for $b = c$ when $n = 4$).

The sequence xyR is an $(n - 1)$ -cup because xy is labeled 1 and $C_y = yR$ is $(n - 2)$ -cup. It is also laced because x is the endpoint of the $(n - 2)$ -cup $C_x = Px$. So we have two laced $(n - 1)$ -cups $xQy = C$ and xyR .

Do case analysis on the label of ab . Assume first that ab is labeled 1. Then aQy is an $(n - 1)$ -cup. It is also laced because a is a 1-cup and yR is a $(n - 2)$ -cup. aQy interweaves with the laced $(n - 1)$ -cup xyR because $a < x < y$ and the endpoint of R is strictly on the right of y .

Assume now that ab is labeled 2. We do another case analysis on the label of by . If by is labeled 1, then byR is a laced $(n - 1)$ -cup because it is a $(n - 1)$ -cup and Pb is a $(n - 2)$ -cup. Now the laced $(n - 1)$ -cups xQy and byR are interweaved because $x < b < y$ and the endpoint of R is strictly on the right of y . If by is labeled 2, then Pby is an $(n - 1)$ -cup, and it is also laced because yR is an $(n - 2)$ -cup. Now laced

$(n-1)$ -cups Pby and xyR are interweaved, because the starting point of P comes before x , $x < y$, and the endpoint of R is strictly on the right side of y . The proof is now done. \square

We introduce the following terminologies for a 4-cap, n -cup free configuration S of size at least $\binom{n-1}{2} + 1$. Note that the following definition does not depend on a particular choice of a slope labeling of S .

Definition 5.4. Assume an arbitrary 4-cap, n -cup free configuration S of size at least $\binom{n-1}{2} + 1$ with $n \geq 3$.

Define $L(S)$ as the set of left endpoints of all $(n-1)$ -cups in S , and $R(S)$ as the set of right endpoints of all $(n-1)$ -cups in S . Let p_S be the rightmost point of $L(S)$ and q_S be the leftmost point of $R(S)$. By Theorem 2.6, there is at least one $(n-1)$ -cup in S , so that $L(S), R(S)$ are nonempty and the points p_S, q_S are well-defined.

Remark 5.5. Note that under the assumption of Definition 5.4, $L(S^{\text{op}}) = R(S)$, $R(S^{\text{op}}) = L(S)$, $p_{S^{\text{op}}} = q_S$ and $q_{S^{\text{op}}} = p_S$.

Lemma 5.6. Under the assumption of Definition 5.4, we always have $p_S \leq q_S$. In other words, the sets $L(S)$ and $R(S)$ are separated in the increasing order with at most one overlap.

Proof. Assume the contrary that $p_S > q_S$. Take any slope labeling of S . If the edge $q_S p_S$ is of label 1 we can extend an $(n-1)$ -cup starting with p_S to the left by q_S , finding an n -cup in S . Likewise, if the edge $q_S p_S$ is of label 2 we can extend an $(n-1)$ -cup starting with q_S to the right by p_S , finding an n -cup in S . Both cases lead to contradiction. \square

We rule out the following case where we can find a pair of interweaved laced $(n-1)$ -cups immediately as well.

Lemma 5.7. Assume an arbitrary 4-cap, n -cup free configuration S of size at least $\binom{n-1}{2} + 1$. If a $(n-2)$ -cup in S ends with p_S and a $(n-2)$ -cup in S starts with q_S , then S contains a pair of interweaved laced $(n-1)$ -cups.

Proof. An $(n-1)$ -cup C_l from some $p' \in L(S)$ to q_S exists by the definition of q_S and $L(S)$. Likewise, an $(n-1)$ -cup C_r from p_S to some $q' \in R(S)$ exists by the definition of p_S and $R(S)$. Note that C_l (or C_r) is laced by the existence of a $(n-2)$ -cup that ends with p_S (or that starts with q_S).

We have $p' \leq p_S$ by the definition of p_S , $p_S \leq q_S$ by Lemma 5.6 and $q_S \leq q'$ by the definition of q_S . If $p' = p_S$, then we can apply lemma 5.3 to C_l to conclude the proof. If $q_S = q'$, then we can apply lemma 5.3 to C_r to conclude the proof. If none of such equalities hold, then $p' < p_S \leq q_S < q'$ holds so C_l and C_r forms a pair of interweaved laced $(n-1)$ -cups. \square

We define the rows of S in the (α, β) -plane and show that each row is nonempty.

Definition 5.8. Fix an arbitrary 4-cap, n -cup free configuration S of size at least $\binom{n-1}{2} + 1$ with $n \geq 3$ and a fixed slope labeling and α -statistic. For any $1 \leq i \leq n-1$, define

$$R_i(S) = \{p \in S : \beta(p) = i\}$$

which is the i 'th row of the (α, β) -plane of S from the bottom. Note that $R_{n-1}(S) = R(S)$ in particular by definition.

Lemma 5.9. Under the assumption of Definition 5.8, each row $R_i(S)$ is nonempty for all $1 \leq i \leq n-1$.

Proof. Define the set $R_{\geq i}(S) = \{p \in S : \beta(p) \geq i\}$ for all $1 \leq i \leq n-1$. By Theorem 2.6, there is at least one $(n-1)$ -cup C in S , so that the right endpoint of C is contained in $R_{\geq i}(S)$ for all i . Therefore, the minimum (leftmost) point x_i of $R_{\geq i}(S)$ exists for all i .

We show that $x_i < x_{i+1}$ for all $1 \leq i < n-1$. As x_{i+1} is in $R_{\geq(i+1)}(S)$, we can take a $(i+1)$ -cup Cx_{i+1} that ends with x_{i+1} . Let y be the ending point of the i -cup C . Then $y \in R_{\geq i}(S)$ by definition and also $y < x_{i+1}$ because Cx_{i+1} forms a cup. This implies that $x_i = \min R_{\geq i}(S) \leq y < x_{i+1}$.

Because S is n -cup free, the equality $R_{n-1}(S) = R_{\geq(n-1)}(S)$ holds and the set is nonempty. For all $1 \leq i < n-1$, the set $R_i(S) = R_{\geq(i+1)}(S) \setminus R_{\geq i}(S)$ contains x_i because $x_i < x_{i+1}$. This concludes the proof. \square

Now we are ready to prove the existence of a pair of interweaved laced $(n-1)$ -cups by induction.

Theorem 5.10. For any $n \geq 3$, any 4-cap, n -cup free configuration of size $\binom{n-1}{2} + 2$ contains a pair of interweaved laced $(n-1)$ -cups.

Proof. We proceed by induction. The base case $n = 3$ can be checked as the following. For any configuration of size $\binom{2}{2} + 2 = 3$, if the points are x, y, z in the increasing order, then xy and yz forms a pair of interweaved laced 2-cups by definition.

Now we show the inductive step. Fix an arbitrary $n \geq 4$ and the following inductive hypothesis (*).

- (*) Any 4-cap, $(n-1)$ -cup free configuration of size $\binom{n-2}{2} + 2$ contains a pair of interweaved laced $(n-2)$ -cups.

Fix an arbitrary 4-cap, n -cup free configuration S of size $\binom{n-1}{2} + 2$. Our goal now is to show that S contains a pair of interweaved laced $(n-1)$ -cups.

By Lemma 5.7, we are already done if p_S is an endpoint of a $(n-2)$ -cup and q_S is the starting point of an $(n-2)$ -cup at the same time. It remains for us to show the case where either p_S is not the endpoint of a $(n-2)$ -cup, or q_S is not the starting point of an $(n-2)$ -cup. Without loss of generality (e.g. Remark 5.5), we can assume that q_S is not the starting point of an $(n-2)$ -cup by reflecting S in the other case. Now fix a specified slope labeling and α -statistic of S to use (Theorem 3.2).

We define the set S' on which we will apply the inductive hypothesis (*) as the following. For each $1 \leq i \leq n-1$, define x_i as the leftmost point of the i 'th row $R_i(S)$ of the (α, β) -plane (Definition 5.8). By Lemma 5.9, each x_i is well-defined. Define the set $\Delta = \{x_1, x_2, \dots, x_{n-2}\}$ of size $n-2$. Note that the point x_{n-1} is excluded from Δ by definition, and that $x_{n-1} = q_S$ by Definition 5.4. Define the set S' of size $\binom{n-2}{2} + 2$ as $S' = S \setminus \Delta$ (see Figure 8). We will show later that S' is $(n-1)$ -cup free, so that we can apply the induction hypothesis (*) to S' .

We assumed without loss of generality that q_S is not the starting point of an $(n-2)$ -cup. Using it, we show the following consequences.




FIGURE 8. A hypothetical (α, β) -plane of a 4-cap, n -cup free set S of size $\binom{n-1}{2} + 2$ with $n = 7$. There are $n - 3 = 4$ ‘holes’ not in S drawn as white points. The set Δ of size $n - 2 = 5$ is marked with circles.

(**) Let x be the starting point of an arbitrary $(n - 2)$ -cup C in S' and let $i = \beta(x)$. Then the following hold.

- (1) $x < q_S$
- (2) $x \notin R_{n-1}(S)$ and $i \leq n - 2$
- (3) $x_i < x$ and the edge $x_i x$ is labeled 1, so $x_i C$ is an $(n - 1)$ -cup in S
- (4) $\alpha(x_i) = 1$ and $\alpha(x) = 2$

First we show $x < q_S$, the first item of (**). Assume otherwise that $x \geq q_S$. Then the case $x = q_S$ directly contradicts our assumption, so we should have $x > q_S$. If the edge $q_S x$ is labeled 1, then $q_S C$ is an $(n - 1)$ -cup so it also contradicts our assumption. So $q_S x$ should be labeled 2. But in this case, as q_S is the ending point of some $(n - 1)$ -cup D by definition, Dx is an n -cup which contradicts that S is n -cup free. This concludes the proof of $x < q_S$ by contradiction.

As q_S is the minimum value of $R(S) = R_{n-1}(S)$, we also get $x \notin R_{n-1}(S)$ and $i = \beta(x) \leq n - 2$ (the second item of (**)). Therefore, $x_i \in \Delta$ and as x_i is minimal in the row $R_i(S)$ that also contains x , we have $x_i < x$ and the edge $x_i x$ is labeled 1. Consequently, the $(n - 1)$ -cup C extends to left with x_i (the third item of (**)).

If $\alpha(x_i) > 1$, then we can extend the $(n - 1)$ -cup $x_i C$ to left in S , so it contradicts that S is n -cup free. So $\alpha(x_i) = 1$. Next, we show that $\alpha(x) = 2$. Because the edge $x_i p$ is labeled 1, we have $\alpha(p) \geq 2$. If $\alpha(p) > 2$, then we have a 3-cup D that ends with p_ϵ and an edge with label 1. Now the cup D and C_ϵ meets at p_ϵ , and they form a cup because D ends with an edge with label 1. The cup is of size $|D| + |C_\epsilon| - 1 = n$ and we reach contradiction. So the equality $\alpha(p_\epsilon) = 2$ holds (the fourth item of (**)). This concludes the proof of (**).

We now show that S' is $(n - 1)$ -cup free. Assume otherwise that S' contains an $(n - 1)$ -cup C_0 from some point x to y . Let $i = \beta(x)$. Then since x is also the starting point of some $(n - 2)$ -cup in S' , the property (**) implies that the edge $x_i x$ has label 1. Now the $(n - 1)$ -cup C_0 in S' extends to left with x_i in S , contradicting that S is n -cup free.

As S' is $(n-1)$ -cup free, we can apply the inductive hypothesis (*). Doing so, we find a pair (C_1, C_2) of interweaved laced $(n-2)$ -cups in S' , each from p_1 to r_1 and p_2 to r_2 so that $p_1 < p_2 \leq r_1 < r_2$. Define $i_1 = \beta(p_1)$ and $i_2 = \beta(p_2)$. By the property (**) applied to C_1 and C_2 respectively, we have the following.

- $p_1, p_2 < q_S$
- $i_1, i_2 < n-1$
- $x_{i_1}p_1$ and $x_{i_2}p_2$ are edges with label 1
- $x_{i_1}C_1$ and $x_{i_2}C_2$ are $(n-1)$ -cups in S
- $\alpha(x_{i_1}) = \alpha(x_{i_2}) = 1$ and $\alpha(p_1) = \alpha(p_2) = 2$

We show that the pair $(x_{i_1}C_1, x_{i_2}C_2)$ of $(n-1)$ -cups is interweaved and laced. This completes the inductive step and the whole proof.

First, we show that the pair is interweaved. That is, whether $x_{i_1} < x_{i_2} \leq r_1 < r_2$ is true. By the assumptions $x_{i_2} < p_2 \leq r_1 < r_2$, it only remains for us to show that $x_{i_1} < x_{i_2}$. Applying Corollary 4.3 to $p_1 < p_2$ and $\alpha(p_1) = \alpha(p_2) = 2$, we have $\beta(p_1) < \beta(p_2)$ which is exactly $i_1 < i_2$ by definition. Applying Corollary 4.3 again to $\beta(x_{i_1}) = i_1 < i_2 = \beta(x_{i_2})$ and $\alpha(x_{i_1}) = \alpha(x_{i_2}) = 1$, we have $x_{i_1} < x_{i_2}$. This completes the proof that the pair $(x_{i_1}C_1, x_{i_2}C_2)$ is interweaved.

Now we show that each cup in the pair $(x_{i_1}C_1, x_{i_2}C_2)$ is laced. Let $\epsilon \in \{1, 2\}$ be any of the index 1 or 2. We extend the laced $(n-2)$ -cup C_ϵ in S' to show that $x_{i_\epsilon}C_\epsilon$ in S is also laced. Because C_ϵ from p_ϵ to r_ϵ is laced in S' , there are cups C_{p_ϵ} and C_{r_ϵ} in S' that ends with p_ϵ and starts with r_ϵ respectively, so that $|C_{p_\epsilon}| + |C_{r_\epsilon}| = n-2$.

Say that C_{p_ϵ} starts with a point z in S' . Then $z \leq p_\epsilon < q_S$ so $\beta(z) < n-1$ by the definition of q_S . Consequently, we have $\alpha(z) \geq 2$ as $\beta(z) < n-1$ and z is not in Δ . So z is the endpoint of an edge yz of label 1 in S . We can extend C_{p_ϵ} to left to a cup yC_{p_ϵ} of size $|C_{p_\epsilon}| + 1$ in S , and the cup yC_{p_ϵ} ends with p_ϵ . This implies that $|C_{p_\epsilon}| + 1 \leq \beta(p_\epsilon) = i_\epsilon = \beta(x_{i_\epsilon})$. So in particular, $\beta(x_{i_\epsilon}) + |C_{r_i}| \geq |C_{p_\epsilon}| + |C_{r_\epsilon}| + 1 = n-1$ and $x_{i_\epsilon}C_\epsilon$ is a laced $(n-1)$ -cup for any of $\epsilon = 1$ or 2 . This concludes the proof. □

We obtain the main theorem (Theorem 2.7) immediately as the following.

Proof of Theorem 2.7. Combine Theorem 5.10 with Lemma 5.2. □

We end with a conjecture that generalizes Theorem 5.10.

Conjecture 5. For any $n \geq 3$ and $1 \leq k \leq n$, any 4-cap, n -cup free configuration of size $\binom{n-1}{2} + k$ contains k mutually interweaved laced $(n-1)$ -cups.

Theorem 5.10 is a special case $k = 2$ of this conjecture. We can also prove the case $k = 1$ with a similar induction argument with the same $S' = S \setminus \Delta$. The case $k = n$ is an immediate consequence of the discussion at the end of Section 4. The proof of Theorem 5.10 fails to extend because it exploits the mirror symmetry and essentially applies the inductive hypothesis twice to force the 'rightmost' laced $(n-1)$ -cup starting with p_S and the 'leftmost' laced $(n-1)$ -cup ending with q_S . This special case is then covered by Lemma 5.3.

ACKNOWLEDGEMENT

The author is indebted to David Speyer for many illuminating discussions and his help in making the presentation much more organized and clear. The author

thanks to Andreas Holmsen for directing the author to the work of Moshkovitz and Shapira [10] that inspired the notion of (α, β) -plane. The author also thanks to Martin Balko and Pavel Valtr for their work [1] that inspired the main Theorem 1.6 and their thoughtful feedback on the early version of the draft.

REFERENCES

- [1] M. Balko and P. Valtr. A sat attack on the erdős–szekeres conjecture. *European Journal of Combinatorics*, 66:13–23, 2017.
- [2] F. R. Chung and R. L. Graham. Forced convex n -gons in the plane. *Discrete & Computational Geometry*, 19(3):367–371, 1998.
- [3] P. Erdős and G. Szekeres. A combinatorial problem in geometry. *Compositio mathematica*, 2:463–470, 1935.
- [4] P. Erdős and G. Szekeres. On some extremum problems in elementary geometry. In *Annales Univ. Sci. Budapest*, pages 3–4, 1960.
- [5] P. Erdos, Z. Tuza, and P. Valtr. Ramsey-remainder. *European Journal of Combinatorics*, 17(6):519–532, 1996.
- [6] J. Fox, J. Pach, B. Sudakov, and A. Suk. Erdős–szekeres-type theorems for monotone paths and convex bodies. *Proceedings of the London Mathematical Society*, 105(5):953–982, 2012.
- [7] A. F. Holmsen, H. N. Mojarad, J. Pach, and G. Tardos. Two extensions of the erdős–szekeres problem. *Journal of the European Mathematical Society*, 22(12):3981–3995, 2020.
- [8] D. Kleitman and L. Pachter. Finding convex sets among points in the plane. *Discrete & Computational Geometry*, 19(3):405–410, 1998.
- [9] H. N. Mojarad and G. Vlachos. An improved upper bound for the erdős–szekeres conjecture. *Discrete & Computational Geometry*, 56(1):165–180, 2016.
- [10] G. Moshkovitz and A. Shapira. Ramsey theory, integer partitions and a new proof of the erdős–szekeres theorem. *Advances in Mathematics*, 262:1107–1129, 2014.
- [11] S. Norin and Y. Yuditsky. Erdős–szekeres without induction. *Discrete & Computational Geometry*, 55(4):963–971, 2016.
- [12] A. Suk. On the erdős–szekeres convex polygon problem. *Journal of the American Mathematical Society*, 30(4):1047–1053, 2017.
- [13] G. Szekeres and L. Peters. Computer solution to the 17-point erdős–szekeres problem. *The ANZIAM Journal*, 48(2):151–164, 2006.
- [14] G. Tóth and P. Valtr. Note on the erdos-szekeres theorem. *Discrete & Computational Geometry*, 19(3):457–459, 1998.
- [15] G. Tóth and P. Valtr. The erdos-szekeres theorem: upper bounds and related results. *Combinatorial and computational geometry*, 19:557–568, 2005.

JINEON BAEK, UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, 2074 EAST HALL, 530 CHURCH STREET, ANN ARBOR, MI 48109-1043
Email address: jineon@umich.edu