

STRUCTURE OF A MAXIMAL TOTAL INDEPENDENT SET

LEWIS STANTON

ABSTRACT. Let G be a simple, connected and finite graph of order n . Denote the independence number, edge independence number and total independence number by $\alpha(G)$, $\alpha'(G)$ and $\alpha''(G)$ respectively. This paper establishes a relation between $\alpha''(G)$ with $\alpha(G)$, $\alpha'(G)$ and n . It also describes the possible structures of a total independent set of a given size.

1. INTRODUCTION

The independence number (denoted $\alpha(G)$) of a graph G is an extensively studied invariant in graph theory. There are a variety of bounds and computer algorithms which can be used to find the independence number and to construct an independent set of vertices. The same is true for a variant of the independence number known as the edge independence number (denoted $\alpha'(G)$). This is a similar invariant to the independence number but is defined on the edge set of the graph as opposed to the vertex set of a graph. The edge independence number is also well understood and is completely determined for certain classes of graphs such as Hamiltonian graphs.

The total independence number (denoted $\alpha''(G)$) is another variation of the definition of the independence number; being defined on the union of the vertex and edge sets of G . While its value can be calculated for all graphs in terms of the edge domination number of a graph, there is not much known in terms of its relationship with other invariants and the structure of a total independent set of a given size.

In this paper, we explore the total independence number and its relationship with the independence number and edge independence number. Theorem 5.1 proves that for a simple, connected and finite graph of order n , $\alpha''(G) \leq n + \left\lfloor \frac{\alpha(G) - n + 2\alpha'(G)}{2} \right\rfloor - \alpha'(G)$. Corollary 7.3 gives another bound which is that $\alpha''(G) \leq \frac{n + \alpha(G)}{2}$. In Section 7, we explore the possible structures a total independent set of a given size can have, namely the possible number of edges and vertices it can contain.

2. BASIC DEFINITIONS

In this paper, we will use the definitions from Gross, Yellen and Zhang [2]. A graph G is *finite* if its vertex set $V(G)$ and edge set $E(G)$ are finite. A graph G is *simple* if any edge has two distinct end vertices and there is at most one edge between any two distinct vertices. Finally, a graph G is *connected* if for any two distinct vertices $x, y \in V(G)$ there exists a walk in G between x and y .

We will restrict our attention to simple, connected and finite graphs; however, many of the results leading up to Theorem 5.1 may hold in a broader context. Let G be a simple, connected and finite

graph of order n . There are various notions of adjacency in a graph. Two vertices $v, v' \in V(G)$ are adjacent if there is an edge $e \in E(G)$ such that $e = v \cdot v'$. Two edges $e, e' \in E(G)$ are adjacent if they share an end vertex. Finally, a vertex $v \in V(G)$ and $e \in E(G)$ are adjacent if v is an end vertex of e .

With these definitions of adjacency, we aim to find subsets of the vertex set or edge set such that none of the elements of the subset are pairwise adjacent. A subset $A \subseteq V(G)$ is an *independent set* if the vertices in A are pairwise non-adjacent. The *independence number* of G is

$$\alpha(G) = \max\{|A| \mid A \text{ is an independent set of } G\}.$$

Similarly, a subset $A' \subseteq E(G)$ is an *edge independent set* if the edges in A' are pairwise non-adjacent. The *edge independence number* is

$$\alpha'(G) = \max\{|A'| \mid A' \text{ is an edge independent set of } G\}.$$

Finally, a subset $A'' \subseteq V(G) \cup E(G)$ is a *total independent set* if the elements of A'' are pairwise non-adjacent. The *total independence number* is

$$\alpha''(G) = \max\{|A''| \mid A'' \text{ is a total independent set of } G\}.$$

There are some connections between dominating sets and independent sets which we will use. A subset $D \subseteq E(G)$ is an *edge dominating set* if every edge $e \in E(G) \setminus D$ is adjacent to an edge in D . The *edge domination number* of G is $\gamma'(G) = \min\{|D| \mid D \text{ is an edge dominating set of } G\}$. The property that we will require is that a maximal edge independent set is also an edge dominating set.

Lemma 2.1. *Let G be a simple, connected and finite graph of order n . Let $A' \subseteq E(G)$ be an edge independent set of size $\alpha'(G)$. Then A' is also an edge dominating set.*

Proof. Let $A' \subseteq E(G)$ be a maximal edge independent set of size $\alpha'(G)$. Suppose A' is not an edge dominating set. This implies that there is an edge $e \in E(G)$ such that e is not adjacent to any edge in A' . However, this implies that the set $B' = A' \cup \{e\}$ is an edge independent set of size larger than A' , contradicting the maximality of A' . \square

We will use a simple bound on the edge independence number.

Lemma 2.2. *Let G be a simple, connected and finite graph of order n . Then $\alpha'(G) \leq \lfloor \frac{n}{2} \rfloor$.*

Proof. Let X be an edge independent set of size $\alpha'(G)$. Each edge in X can not share an end vertex with any other edge otherwise they would be adjacent to one another. Therefore, there are $2\alpha'(G)$ vertices contained in X as end points of the edges. Hence, $2\alpha'(G) \leq n$ which implies that $\alpha'(G) \leq \frac{n}{2}$. Since $\alpha'(G)$ is an integer, $\alpha'(G) \leq \lfloor \frac{n}{2} \rfloor$ as desired. \square

A *Hamiltonian cycle* is a cycle that contains each vertex exactly once. A *Hamiltonian graph* is a graph which contains a Hamiltonian cycle. For Hamiltonian graphs, we obtain a stronger result.

Lemma 2.3. *Let G be a simple, connected and finite graph of order n which is Hamiltonian. Then $\alpha'(G) = \lfloor \frac{n}{2} \rfloor$.*

Proof. Let C be a Hamiltonian cycle in G consisting of vertices x_1, \dots, x_n . We can construct a total independent set by taking every other edge in the Hamiltonian cycle. If n is even, take $x_1 \cdot x_2, x_3 \cdot x_4, \dots, x_{n-1} \cdot x_n$ as an edge independent set. If n is odd, take $x_1 \cdot x_2, x_3 \cdot x_4, \dots, x_{n-2} \cdot x_{n-1}$. In both cases, these are edge independent sets of size $\lfloor \frac{n}{2} \rfloor$ therefore $\alpha'(G) \geq \lfloor \frac{n}{2} \rfloor$. However, since $\alpha'(G) \leq \lfloor \frac{n}{2} \rfloor$ by Lemma 2.2, $\alpha'(G) = \lfloor \frac{n}{2} \rfloor$. \square

In Section 3, we provide a summary of existing results. There are two further invariants of graphs in this section which we consider.

Definition 2.4. The *arboricity* of an undirected graph G is the minimum number of subsets that $V(G)$ can be partitioned into such that each subset induces an acyclic graph. We will denote this $\beta(G)$.

Definition 2.5. The *edge arboricity* of an undirected graph G is the minimum number of subsets that $E(G)$ can be partitioned into such that each subset induces an acyclic graph. We will denote this $\beta'(G)$.

These are also both well studied invariants which tell us how dense a graph is. The value of the edge arboricity of a graph is given by the Nash-Williams theorem [1].

We also have some special types of graph which we will consider. The *discrete graph* on n vertices is the graph with $V(G) = \{x_1, \dots, x_n\}$ and $E(G) = \emptyset$. The *complete graph* K_n , $n \geq 1$ is a graph with $V(G) = \{x_1, \dots, x_n\}$ and the edge set consists of an edge for every pair of distinct vertices x_i, x_j , $1 \leq i < j \leq n$. The *complete bipartite graph* $K_{m,n}$, $n, m \geq 1$ is a graph with $V(G) = A \cup B$ where A and B are disjoint sets with $|A| = m$ and $|B| = n$. The edge set consists of all edges of the form $a \cdot b$ where $a \in A$ and $b \in B$.

In this paper, we will prove a relation between $\alpha''(G)$ with $\alpha(G)$ and $\alpha'(G)$ and use the proof of this to show the possible structures of a total independent set of given size.

3. SUMMARY OF EXISTING RESULTS

We begin with some simple relations between $\alpha''(G)$, $\alpha(G)$ and $\alpha'(G)$.

Lemma 3.1. *Let G be a simple, connected and finite graph. Then $\alpha''(G) \leq \alpha(G) + \alpha'(G)$.*

Proof. Let X be a total independent set of size $\alpha''(G)$. Let $X_v = X \cap V(G)$ be the set of vertices contained in X and $X_e = X \cap E(G)$ be the set of edges contained in X . From the definition of $\alpha(G)$ and $\alpha'(G)$, we must have that $|X_v| \leq \alpha(G)$ and $|X_e| \leq \alpha'(G)$ and hence $|X| = |X_v| + |X_e| \leq \alpha(G) + \alpha'(G)$. \square

For completeness, we state and prove a basic lower bound on the total independence number based on the independence number and edge independence number.

Lemma 3.2. *Let G be a simple, connected and finite graph. Then $\alpha''(G) \geq \max\{\alpha(G), \alpha'(G)\}$.*

Proof. Let $X_{\alpha(G)}$ be an independent set of size $\alpha(G)$. This is clearly a total independent set so $\alpha''(G) \geq \alpha(G)$. Similarly, let $X_{\alpha'(G)}$ be an edge independent set of size $\alpha'(G)$. This is clearly also a total independent set so $\alpha''(G) \geq \alpha'(G)$. Combining these together, we get that $\alpha''(G) \geq \max\{\alpha(G), \alpha'(G)\}$. \square

These two bounds do not provide much insight into the total independence number. By considering the underlying structure of a total independent set, Theorem 5.1 will provide an alternative bound on the total independence number in relation to the independence number and the edge independence number. In Section 6, we consider when this bound is strictly better than the bound in Lemma 3.1. Before doing this, we consider some other bounds based on the arboricity and edge arboricity [4].

Lemma 3.3. *Let G be a simple, connected and finite graph. Then $1 + \lfloor \frac{n}{2} \rfloor \leq \beta(G) + \alpha''(G) \leq n + 1$ and $1 + \lfloor \frac{n}{2} \rfloor \leq \beta'(G) + \alpha''(G) \leq n + \lfloor \frac{(n+1)}{8} \rfloor$.* \square

We also have an equality in terms of the edge domination number and the total independence number [3].

Lemma 3.4. *Let G be a simple, connected and finite graph. We have that $\gamma'(G) + \alpha''(G) = n$.* \square

4. PRELIMINARY RESULTS

In this Section, we set out some basic results which will help us prove the main theorem. The purpose of these lemmas is to show that the procedure we use to prove the main theorem will always work for all simple, connected and finite graphs.

Lemma 4.1. *If G is a finite, simple graph of order n that is not the discrete graph, then $\alpha(G) < n$.*

Proof. Suppose we have an independent set A of size n . Since G is not the discrete graph, there exists an edge $x \cdot y$. However, this means that x and y can not be in A as they are adjacent. \square

Lemma 4.2. *For any simple, connected and finite graph G of order n , $\alpha(G) + 2\alpha'(G) \geq n$.*

Proof. First consider the case $\alpha'(G) = \lfloor \frac{n}{2} \rfloor$. Since n is an integer, $\frac{n-1}{2} \leq \lfloor \frac{n}{2} \rfloor \leq \frac{n}{2}$ which implies that $n - 1 \leq 2 \cdot \lfloor \frac{n}{2} \rfloor \leq n$. Therefore, $2\alpha'(G) \geq n - 1$. Since $\alpha(G) \geq 1$, it follows that $2\alpha'(G) + \alpha(G) \geq n$.

Now suppose $\alpha'(G) < \lfloor \frac{n}{2} \rfloor$. Note that $\alpha'(G) < \lfloor \frac{n}{2} \rfloor$ implies that $2\alpha'(G) < n$, and so $n - 2\alpha'(G) \geq 1$. Now let F be a maximal edge independent set of size $\alpha'(G)$. In particular, F contains $2\alpha'(G)$ distinct vertices as end vertices of edges in F . Let X_F be the set of vertices which are end vertices

of edges in F . Note that $|X_F| = 2\alpha'(G)$. As F is a maximal edge independent set, by Lemma 2.1 it is also an edge dominating set. Therefore, for all $e \in E(G)$, $e = x \cdot y$ for some $x \in V(G)$ and $y \in X_F$. Now consider $V(G) \setminus X_F$. By definition, $|V(G) \setminus X_F| = n - 2\alpha'(G) \geq 1$. We want to show that $V(G) \setminus X_F$ is an independent set.

If $|V(G) \setminus X_F| = 1$ then it is trivially an independent set. If $|V(G) \setminus X_F| > 1$, consider $x, x' \in V(G) \setminus X_F$ where x, x' are distinct. Suppose x, x' are adjacent vertices, then $x \cdot x' \in E(G)$. However, this implies that either $x \in X_F$ or $x' \in X_F$ which is a contradiction. Therefore, none of the vertices in $V(G) \setminus X_F$ can be adjacent to each other so they form an independent set. Hence, $\alpha(G) \geq |V(G) \setminus X_F| = n - 2\alpha'(G)$, and so $\alpha(G) + 2\alpha'(G) \geq n$. \square

5. MAIN THEOREM

With the results in the previous section, we can now prove the main theorem. The basic idea is to partition the total independent sets by the amount of edges contained in them and construct an upper bound for each set in turn. The upper bound therefore on $\alpha''(G)$ will be the biggest of these bounds. For a set $X \subset V(G) \cup E(G)$ in the following, we will use the following notation, $X_v = X \cap V(G)$ and $X_e = X \cap E(G)$.

Theorem 5.1. *Let G be a simple, connected and finite graph of order n . Then*

$$\alpha''(G) \leq n + \left\lfloor \frac{\alpha(G) - n + 2\alpha'(G)}{2} \right\rfloor - \alpha'(G).$$

Proof. Let G be a simple, connected and finite graph and let \mathcal{X}_G be the set of total independent sets of G . Partition $\mathcal{X}_G = \mathcal{X}_{G_{\alpha'(G)}} \cup \mathcal{X}_{G_{\alpha'(G)-1}} \cdots \cup \mathcal{X}_{G_0}$ where \mathcal{X}_{G_i} is the set of total independent sets containing i edges. Let $m' = \left\lfloor \frac{\alpha(G) - n + 2\alpha'(G)}{2} \right\rfloor$. By Lemma 4.2, $m' \geq 0$. Since G is simple, connected and finite, by Lemma 4.1 $\alpha(G) < n$, which implies that $\alpha(G) - n < 0$. Therefore, $\alpha(G) - n + 2\alpha'(G) < 2\alpha'(G)$ and so $\frac{\alpha(G) - n + 2\alpha'(G)}{2} < \alpha'(G)$. By definition $m' \leq \frac{\alpha(G) - n + 2\alpha'(G)}{2}$. Hence, $m' < \alpha'(G)$. Observe each vertex can only appear once in a total independent set and we can have a maximum of $\alpha'(G)$ edges in a total independent set and $\alpha(G)$ vertices in the total independent set.

Consider $X \in \mathcal{X}_{G_{\alpha'(G)-i}}$ with $0 \leq i \leq m'$. Write X as $X = X_v \cup X_e$. By definition of $\mathcal{X}_{G_{\alpha'(G)-i}}$, $|X_e| = \alpha'(G) - i$. This means that $2(\alpha'(G) - i)$ vertices appear in X_e as end vertices. Now,

$$n - 2(\alpha'(G) - i) = n - 2\alpha'(G) + 2i \leq n - 2\alpha'(G) + 2m' \leq n - 2\alpha'(G) + 2\alpha'(G) + \alpha(G) - n = \alpha(G).$$

Therefore, there are at most $n - 2(\alpha'(G) - i)$ vertices which can be added to X_v . Hence, $|X| = |X_v| + |X_e| \leq n - 2(\alpha'(G) - i) + \alpha'(G) - i = n - \alpha'(G) + i$. Observe that this is increasing with i as i goes from 0 to m' . Hence, for all $i \in \mathbb{N}_0$ with $0 \leq i \leq m'$ and for all $X \in \mathcal{X}_{G_{\alpha'(G)-i}}$, $|X| \leq n - \alpha'(G) + m'$

Now consider $X \in \mathcal{X}_{G_{\alpha'(G)-(m'+1)}}$, write X as $X = X_v \cup X_e$. By definition of $\mathcal{X}_{G_{\alpha'(G)-(m'+1)}}$, $|X_e| = \alpha'(G) - (m' + 1)$. This means that $2(\alpha'(G) - (m' + 1))$ vertices appear in the set as end

vertices. However, m' has the property that $m' \leq \frac{\alpha(G)-n+2\alpha'(G)}{2}$ and $m' + 1 > \frac{\alpha(G)-n+2\alpha'(G)}{2}$. Therefore

$$\begin{aligned} n - 2(\alpha'(G) - (m' + 1)) &= n - 2\alpha'(G) + 2(m' + 1) > n - 2\alpha'(G) + 2\left(\frac{\alpha(G) - n + 2\alpha'(G)}{2}\right) \\ &= n - 2\alpha'(G) + 2\alpha'(G) + \alpha(G) - n = \alpha(G). \end{aligned}$$

However, we can not add more than $\alpha(G)$ vertices to the X_v . Therefore for $X \in \mathcal{X}_{G_{\alpha'(G)-i}}$ for $i > m'$, we can only add at most $\alpha(G)$ more vertices to the set X_v .

Hence, for $X \in \mathcal{X}_{G_{\alpha'(G)-i}}$ with $i > m'$ and writing X as $X = X_v \cup X_e$ where $|X_e| = \alpha'(G) - i$. It follows that $|X| = |X_v| + |X_e| \leq \alpha(G) + \alpha'(G) - i$, which is decreasing as i increases. It follows for all $i \in \mathbb{N}_0$ with $m' + 1 \leq i \leq \alpha'(G)$ and for all $X \in \mathcal{X}_{G_{\alpha'(G)-i}}$, $|X| \leq \alpha(G) + \alpha'(G) - (m' + 1)$

Now, $\frac{\alpha(G)-n+2\alpha'(G)}{2} \leq m' + \frac{1}{2}$ which implies that $\frac{\alpha(G)-n+2\alpha'(G)-1}{2} \leq m'$ and so $\alpha(G) - n + 2\alpha'(G) - 1 \leq 2m'$. Hence, $\alpha'(G) - (m' + 1) + \alpha(G) \leq n - \alpha'(G) + m'$. Therefore, for all $i \in \mathbb{N}_0$ where $0 \leq i \leq \alpha'(G)$ and for all $X \in \mathcal{X}_{G_{\alpha'(G)-i}}$, $|X| \leq n - \alpha'(G) + m'$, and so $\alpha''(G) \leq n - \alpha'(G) + m'$. \square

6. HOW GOOD IS THE ESTIMATE?

The next question to consider is how good the bound is that Theorem 5.1 provides.

Lemma 6.1. *The bound in Theorem 5.1 is sharp for all n .*

Proof. Consider the complete graph K_n where n is odd. By definition of K_n , $\alpha(K_n) = 1$ and $\alpha'(K_n) = \lfloor \frac{n}{2} \rfloor = \frac{n-1}{2}$ as n is odd. Therefore by Theorem 5.1,

$$\alpha''(G) \leq n + \left\lfloor \frac{1 - n + n - 1}{2} \right\rfloor - \frac{n-1}{2} = \frac{1}{2}(n+1) = \frac{1}{2}(n-1) + 1.$$

We can construct a total independent set of this size by first taking any maximal edge independent set. This contains $n - 1$ distinct vertices as end vertices of the edges in this edge independent set and so we can then take the total independent set to be the $\frac{n-1}{2}$ edges in the edge independent set and the vertex left over. This is a total independent set of size $\frac{1}{2}(n-1) + 1$ so $\alpha''(G) = \frac{1}{2}(n-1) + 1$ in this case.

For n even, $\alpha(K_n) = 1$ and $\alpha'(K_n) = \frac{n}{2}$. Therefore by Theorem 5.1,

$$\alpha''(G) \leq n + \left\lfloor \frac{1 - n + n}{2} \right\rfloor - \frac{n}{2} = \frac{n}{2}.$$

We can construct a total independent set of this size by taking any maximal edge independent set. \square

However, it is not always a sharp bound. Consider the graph $K_{2,3}$. It follows from the definition of $K_{2,3}$ that $\alpha(K_{2,3}) = 3$ and $\alpha'(G) = 2$. Therefore by Theorem 5.1, $\alpha''(G) \leq 5 + \lfloor \frac{3-5+4}{2} \rfloor - 2 = 4$. However, we can not construct a total independent set of this size. If we have a total independent set containing 2 edges, we can only at most add one extra vertex to the total independent set as 4

vertices are end vertices of the edges. If we have a total independent set containing no edges, we can have at most 3 vertices in the total independent set as $\alpha(K_{2,3}) = 3$.

This leaves the case of a total independent set containing 1 edge. Let $V(K_{2,3}) = A \cup B$ be the partition of the vertices with $|A| = 2$ and $|B| = 3$. The edge contained in the edge independent set is of the form $x \cdot y$ where $x \in A$ and $y \in B$. Therefore, there are three vertices left you could add to the total independent set. However, it is clear you can not add all 3 as the vertex in $A \setminus \{x\}$ is adjacent to both the vertices in B . Therefore, we can only add the two vertices in $B \setminus \{y\}$ so the total independent set has size 3. This implies that $\alpha''(G) = 3 < 4$ and the bound is not an equality.

We now show that the estimate in the theorem is the same as or better than the result in Lemma 3.1.

Lemma 6.2. *The result in Theorem 5.1 is equal to or better than the result in Lemma 3.1.*

Proof. We have that $n + m' - \alpha'(G) \leq n + 2m' - \alpha'(G) \leq n + \alpha(G) - n + 2\alpha'(G) - \alpha'(G) = \alpha(G) + \alpha'(G)$. \square

We can prove easily when this is a strict inequality based on the value of $\frac{\alpha(G) - n + 2\alpha'(G)}{2}$.

Lemma 6.3. *If $\frac{\alpha(G) - n + 2\alpha'(G)}{2} > 0$ then the result in Theorem 5.1 is strictly better than the result in Lemma 3.1.*

Proof. Case 1: $\frac{\alpha(G) - n + 2\alpha'(G)}{2} \geq 1$. In this case, $\frac{\alpha(G) - n + 2\alpha'(G)}{2} \geq 1$ which implies that $m' \geq 1$. Therefore,

$$n + m' - \alpha'(G) < n + 2m' - \alpha'(G) \leq n + \alpha(G) - n + 2\alpha'(G) - \alpha'(G) = \alpha(G) + \alpha'(G).$$

Case 2: $\frac{\alpha(G) - n + 2\alpha'(G)}{2} = \frac{1}{2}$. In this case, $\frac{\alpha(G) - n + 2\alpha'(G)}{2} = \frac{1}{2}$ which implies that $m' = 0$. We have that $0 < \frac{\alpha(G) - n + 2\alpha'(G)}{2}$ which implies that $0 < \alpha(G) - n + 2\alpha'(G)$, and so $n - \alpha'(G) < \alpha(G) + \alpha'(G)$. \square

We can show that this is in general not true when $\frac{\alpha(G) - n + 2\alpha'(G)}{2} = 0$. Consider the graph K_3 . It follows from the definition of K_3 that $\alpha(K_3) = 1$ and $\alpha'(K_3) = 1$. This satisfies the condition $\frac{\alpha(K_3) - n + 2\alpha'(K_3)}{2} = \frac{1 - 3 + 2}{2} = 0$ and we can see that $\alpha(K_3) + \alpha'(K_3) = 2$ and $n - \alpha'(G) = 2$.

7. STRUCTURE OF MAXIMAL TOTAL INDEPENDENT SET

Theorem 5.1 considers every possible structure of a total independent set and produces a bound for each such set. We can adapt this proof to provide the possible structures of a maximal total independent set.

Corollary 7.1. *Let G be a simple, connected and finite graph of order n . The number of edges e in a total independent set of size k must satisfy $k - \alpha(G) \leq e \leq n - k$.*

Proof. From the proof of Theorem 5.1, we can see that for a total independent set with $e = \alpha'(G) - i$ edges in it with $i \leq m'$, the size of a total independent set can be at most $n - \alpha'(G) + i$. Therefore, we want to consider the cases where $n - \alpha'(G) + i \geq k$. This implies that $i \geq k + \alpha'(G) - n$ and so $-i \leq n - k - \alpha'(G)$. Hence $e = \alpha'(G) - i \leq n - k$

Now for $i > m'$, from the proof of Theorem 5.1, a total independent set containing $e = \alpha'(G) - i$ edges can have size at most $\alpha'(G) - i + \alpha(G)$. Similar to the previous case, we want to consider the cases where $\alpha'(G) - i + \alpha(G) \geq k$. This implies that $i \leq \alpha'(G) + \alpha(G) - k$ and so $-i \geq k - \alpha'(G) - \alpha(G)$. Hence $e = \alpha'(G) - i \geq k - \alpha(G)$.

Combining the two, we get $k - \alpha(G) \leq e \leq n - k$. □

We note by letting $k = \alpha''(G)$, we immediately get the following result.

Corollary 7.2. *Let G be a simple, connected and finite graph of order n with total independence number $\alpha''(G)$. The amount of edges e in a largest total independent set of size $\alpha''(G)$ must satisfy $\alpha''(G) - \alpha(G) \leq e \leq n - \alpha''(G)$.* □

We can use Corollary 7.2 to easily get another bound on $\alpha''(G)$ in terms of $\alpha(G)$.

Corollary 7.3. *Let G be a simple connected and finite graph of order n with total independence number $\alpha''(G)$. Then $\alpha''(G) \leq \frac{n + \alpha(G)}{2}$.*

Proof. By Corollary 7.2, $\alpha''(G) - \alpha(G) \leq e \leq n - \alpha''(G)$ which implies that $\alpha''(G) - \alpha(G) \leq n - \alpha''(G)$. Rearranging this, we obtain $\alpha''(G) \leq \frac{n + \alpha(G)}{2}$. □

REFERENCES

- [1] Bo Liang. Chen, Makoto Matsumoto, Jian Fang. Wang, Zhong Fu. Zhang, and Jian Xun. Zhang. "A short proof of Nash-Williams theorem for the arboricity of a graph". *Graphs and Combinatorics* 10.1 (1994), pp. 27–28.
- [2] Jonathan L. Gross, Jay Yellen, and Ping Zhang. *Handbook of Graph Theory, Second Edition*. CRC Press, 2013.
- [3] Zhong Fu. Zhang, Jian Xun. Zhang, and Jingwen Li. "On The Relations Between The Parameters Of Graphs". *Taiwanese Journal of Mathematics* 10.6 (2006), pp. 1549–1556.
- [4] Zhong Fu. Zhang, Jian Xun. Zhang, Jian Fang. Wang, and Bo Liang. Chen. "Relations among arboricity, total independence numbers and total covering numbers". *Qufu Shifan Daxue Xuebao Ziran Kexue Ban* 18, no. 4 (1992), pp. 9–14.