

NON-BINARY COUNTERPARTS OF THE PREFER-SAME AND PREFER-OPPOSITE DE BRUIJN SEQUENCES

ABBAS ALHAKIM
DEPARTMENT OF MATHEMATICS
AMERICAN UNIVERSITY OF BEIRUT
BEIRUT, LEBANON

ABSTRACT. The well-known prefer-one, prefer-opposite, and prefer-same binary de Bruijn sequences are all constructed using simple preference rules. We apply the technique of preference functions of span one to define q -ary sequences that generalize the prefer-opposite and prefer-same sequences and we present some of their basic properties that are shared with their binary versions. In particular, we show that the prefer-higher sequence (the nonbinary counter-part of the prefer-one sequence) is obtained from a homomorphic image of the proposed prefer-opposite, when repetitions are cleaned up. This mirrors a known relationship between the binary versions. We also perform calculations that demonstrate that the discrepancy profile of the proposed sequences is similar to that of the binary case.

1. INTRODUCTION

Given positive integers $q > 1$ and $n \geq 1$, a q -ary de Bruijn sequence of order n is a sequence of symbols such that every pattern of size n appears exactly once as a contiguous subsequence. A small class of particularly simple greedy algorithms that generate binary de Bruijn sequences of any given order are the prefer-one algorithm, see [8, 13], the prefer-same algorithm [7, 9] and the prefer-opposite algorithm [1].

The oldest and most well-known of these is the prefer-one. This algorithm proceeds by appending bits to an initial string of n zeros in such a way that every time a bit is to be added, 1 is attempted first and appended to the sequence. If the last n bits form a word that was not encountered earlier then this appended 1 is kept. Otherwise it is replaced by a 0 and the same check is made. The process is repeated in the same fashion and it is stopped as soon as neither 1 nor 0 puts a new n -tuple. The prefer-opposite algorithm works in a similar way but favors the opposite of the most recently placed bit. Likewise, the prefer same algorithm first proposes to duplicate the most recently placed bit. In its original version, it must be started with a string of n ones followed by a string of n zeros and the algorithm must also keep track of the number of occurrences of runs of consecutive ones and zeros of all possible lengths in order to know whether or not a proposed bit is to be accepted, even when it puts a new n -tuple. [4] shows that the extra counting of runs is not necessary, provided the initial word of the sequence is the

¹This research was partially supported by the University Research Board (URB) of the American University of Beirut. Project Number 104107

Key words and phrases. De Bruijn sequence, Ford sequence, preference function, prefer-one sequence, prefer-opposite sequence, prefer-same sequence.

alternating string $1010\cdots$, thus establishing that the prefer-same algorithm is indeed of the same type as its other two sequences.

Another attractive feature of the three sequences is that they are all extremes. It is well known that the Prefer-one sequence is the lexicographically least de Bruijn sequence for a given order n . On the other hand, the prefer-opposite and prefer-same sequences respectively have the smallest and lexicographically largest run length encoding, see [4] for definition and proof.

When q is higher than 2, a generalization of the prefer-one sequence, known as the prefer-higher or non-binary Ford sequence, is classical, see Fredricksen [9]. For the other two sequences, the concept of ‘opposite’ is needed but not well defined. We will employ preference functions of span 1 to obtain natural generalization of the prefer-opposite and prefer-same sequences to the case of alphabet size q . Span-1 preference functions correspond to those de Bruijn sequences that are representable by $q \times q$ matrices. Indeed, the binary de Bruijn sequences mentioned above are representable using 2×2 matrices together with an appropriate starting pattern.

The rest of the paper is organized as follows. In Section 2 we give basic definitions and background about preference functions, with preliminary lemmas that will be essential for the rest of the paper. In Section 3 we propose two classes of nonbinary sequences which we respectively call nonbinary prefer-opposite and prefer-same sequences and we establish properties that are similar to those of the binary counterparts. In Section 4 we show that mapping the proposed prefer-opposite sequence via an appropriate homomorphism we recover the prefer-higher sequence, replicating the relationship between the binary prefer-opposite and the prefer-one sequences, via the Lempel Homomorphism.

2. PRELIMINARIES

In this section we give basic definitions and provide important lemmas that are essential for establishing the results of this paper. For $q \geq 2$, \mathcal{A} is an alphabet with q symbols that we will denote as $\{0, 1, \dots, q-1\}$. We will interchangeably refer to these symbols as digits. For an integer $n \geq 1$, a sequence of n digits will be referred to as an n -word, and denoted as $a_1 \cdots a_n$ and often as (a_1, \dots, a_n) for notational clarity. a^n denotes the word obtained by repeating the symbol a n times. A *de Bruijn digraph* of order n admits all possible n -words as *vertices*, and for two vertices $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$, there is an *edge* from \mathbf{a} to \mathbf{b} if and only if $a_i = b_{i-1}$ for $i = 2, \dots, n$. A string $a_1 \dots a_L$ of length $L \geq n$ can then be identified with a *directed path* (or simply a path) in the de Bruijn digraph with vertices $(a_1, \dots, a_n), (a_2, \dots, a_{n+1}), \dots, (a_{L-n+1}, \dots, a_L)$. A *cycle* is a path whose first and last vertices coincide. A *simple cycle* is one that does not cross a vertex, other than the first and last, more than once. The first definition below follows Golomb [11].

Definition 2.1. For an integer $n \geq 1$, a preference function is a function $P : \mathcal{A}^n \rightarrow S$, where S is the set of all permutations of the elements of \mathcal{A} . We write $P(\mathbf{x}) = (P_1, \dots, P_q)$ for every n -word \mathbf{x} .

0 → 2, 1, 0	0 → 1, 2, 0	0 → 1, 2, 0
1 → 2, 1, 0	1 → 1, 0, 2	1 → 2, 1, 0
2 → 2, 1, 0	2 → 2, 1, 0	2 → 0, 1, 2
00 → 0, 2, 1	10 → 1, 2, 0	20 → 1, 2, 0
01 → 1, 2, 0	11 → 0, 1, 2	21 → 0, 2, 1
02 → 2, 1, 0	12 → 0, 2, 1	22 → 0, 1, 2

TABLE 1. Top row: three preference diagrams of span 1. Lower row: one preference diagram of span 2.

Definition 2.2. For an integer s , $0 \leq s \leq n$, a preference function P defined on \mathcal{A}^n is said to be of span s if s is the smallest integer such that $P(x_1, \dots, x_n)$ is fully determined by (x_{n-s+1}, \dots, x_n) , for all n -words (x_1, \dots, x_n) .

Examples of preference functions of spans 0, 1, 2 for $q = 3$ are displayed as diagrams in Table 1, where the permutation to the right of an arrow is the function value of the pattern on the left of the arrow. Observe that the upper left diagram is that of a constant preference function and so it is of span 0.

The following recursive construction produces a unique finite sequence $\{a_i\}$ from an alphabet \mathcal{A} , given an arbitrary initial n -word (I_1, \dots, I_n) with $n > s$ and a preference function of span s . We denote the resulting sequence by (P, I) .

1. For $i = 1, \dots, n$ let $a_i = I_i$.
2. Suppose that a_1, \dots, a_k for some integer $k \geq n$ have been defined. Let $a_{k+1} = P_i(a_{k-s+1}, \dots, a_k)$ where i , $1 \leq i \leq q$ is the smallest integer such that the n -word $(a_{k-n+2}, \dots, a_{k+1})$ has not appeared in the sequence as a substring, if such an i exists.
3. If no such i exists, halt the program (the construction is complete.)

The following lemma is a slight generalization of Lemma 2 of Chapter 3 in Golomb [11]. The proof is essentially the same.

Lemma 2.3. Consider an arbitrary preference function P of span $s \geq 0$ and initial word $I = (I_1, \dots, I_n)$; $n > s$. Then every n -word occurs at most once in (P, I) . Furthermore, (P, I) ends with the pattern (I_1, \dots, I_{n-1}) .

It follows that the sequence (P, I) can be identified with a simple cycle in the de Bruijn digraph of order n , by removing the last pattern (I_1, \dots, I_{n-1}) and wrapping the rest around a circle. In the following two definitions, we fix an integer $s \geq 1$ and a preference function P of span s , which can equivalently be considered as a function from \mathcal{A}^s to S .

Definition 2.4. For an integer i such that $1 \leq i \leq q$, the i^{th} column function induced by P is a function from \mathcal{A}^s to \mathcal{A}^s defined as

$$g_i(x_1, \dots, x_s) = (x_2, \dots, x_s, P_i(x_1, \dots, x_s)).$$

Naturally, g_i will define at least one cycle of length $k \geq 1$. That is, a sequence of k vertices v_1, \dots, v_k in \mathcal{A}^s such that $g_i(v_j) = v_{j+1}$ for $j = 1, \dots, k-1$ and $g_i(v_k) = v_1$.

Definition 2.5. For a given i , $1 \leq i \leq q$ and a cycle C defined by g_i , we say that a word (x_1, \dots, x_s) is on C if it is simply one of the vertices on C . For an arbitrary integer $n > s$, we say that (x_1, \dots, x_n) is on C if it forms a path in C . That is, if (x_1, \dots, x_s) is a vertex on C and $x_j = P_i(x_{j-s}, \dots, x_{j-1})$ for $j = s+1, \dots, n$.

We note that a path in C in the above definition need not be a simple path. That is, it may cross itself one or more times, depending on the value of n . As an example, the word 0^n of n zeros is a path on the cycle that sends 0^s to itself, for $n > s$.

Definition 2.6. Let g_k , for some k , $1 \leq k \leq q$, be the function induced by the k^{th} column of a preference function P . To a given cycle C on g_k we associate the following sets.

- $[C]$ is the set of all vertices on C .
- The closure $\overline{[C]}$ consists of every vertex in \mathcal{A}^s such that g_k defines a path that starts with this vertex and terminates in C . That is, $(a_1, \dots, a_s) \in \overline{[C]}$ if and only if there is a minimal integer $s' \geq s$ such that for $i = 1, \dots, s' - s$, $g_k(a_i, \dots, a_{i+s-1}) = (a_{i+1}, \dots, a_{i+s})$ and $(a_{s'-s+1}, \dots, a_{s'}) \in [C]$.
- Σ is the complement of $\overline{[C]}$ in \mathcal{A}^s .

The following theorem and its corollary are stated and proved in Alhakim [5].

Theorem 2.7. *Let P be a preference function of span $s \geq 0$ such that the induced g_q has [at least] a cycle C , and let $[C]$, $\overline{[C]}$ and Σ be the sets associated with C as in Definition 2.6. For an arbitrary integer $n > s$ let $I = (I_1, \dots, I_n)$ be such that (I_1, \dots, I_{n-1}) is on C as in Definition 2.5.*

Also, assume that there is an integer q' ; $1 < q' \leq q$ with the property that Σ is closed under all functions g_k with $k \geq q'$, in the sense that whenever $(x_1, \dots, x_s) \in \Sigma$, then $g_k(x_1, \dots, x_s) \in \Sigma$ for all $k \geq q'$. Then the sequence (P, I) misses the set of words M defined as

$$\{(x_1, \dots, x_n) : ((x_1 \dots x_s) \in \Sigma) \wedge (\forall i > s, x_i = P_k(x_{i-s}, \dots, x_{i-1}) \text{ for some } k \geq q')\}$$

Furthermore, if q' is the smallest integer with the given property and if the function $\tilde{g}_{q'-1}$ (defined as $g_{q'-1}$ restricted to Σ) has no cycles, then no other word is missing.

Corollary 2.8. *Let P be a preference function of span $s \geq 0$ such that the induced function g_q has exactly one cycle C , and let $I = (I_1, \dots, I_n)$ be an initial word such that (I_1, \dots, I_{n-1}) is on C . Then (P, I) is a de Bruijn sequence.*

3. GENERALIZING THE PREFER-OPPOSITE AND PREFER-SAME SEQUENCES

In this section we will focus on preference functions P of span 1, that can be represented by $q \times q$ a matrices, where for each digit $j = 0, \dots, q - 1$, $P(j)$ is given by the $(j + 1)^{\text{st}}$ row of the matrix. When $q = 2$, the only possible matrices are

$$F_2 = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}; O_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}; S_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}; \quad \text{and } Z_2 = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}.$$

Applying Theorem 2.7 to the first matrix with initial word 0^n , the set Σ is empty and we obtain the famous prefer-one sequence, first attributed to Martin [13]. With the same initial word, the second matrix gives $q' = 2$ and $\Sigma = \{1\}$, leading to the prefer-opposite sequence, see Alhakim [1]. For the third sequence, starting with one of the alternating strings $010\dots$ or $101\dots$ leads to a full sequence. [4] shows that these sequences give a cyclic rotation of the prefer-same sequence (and bitwise complement), thus presenting

a new and simplified way to generate the latter sequence, avoiding some subtleties in the original algorithm described in Eldert [7] and Fredricksen [9]. The fourth matrix, the ‘prefer-zero’ matrix, evidently produces the complement of the prefer-one sequence when the initial word is 1^n .

As mentioned in the introduction, the prefer-opposite is already generalized to the so-called prefer-higher sequence for general alphabet size q , see Fredricksen [9]. For the other two sequences, the concept of ‘opposite’ is of course not naturally meaningful. However, we will employ preference functions of span 1–representable by $q \times q$ matrices– to obtain natural generalization of the prefer-opposite and prefer-same sequences to the case of alphabet size q .

Fix a digit $d \in \mathcal{A}$. We define the two $q \times q$ matrices $O := O^{(q,d)}$ and $S := S^{(q,d)}$ by

$$O_{ij} = i + (j + 1)d \pmod q \text{ for } i, j = 0, 1, \dots, q, \text{ and}$$

$$S_{ij} = i + (j)d \pmod q \text{ for } i, j = 0, 1, \dots, q$$

Examples of preference functions of the O type are given in the top row of Table 2, while the lower row illustrates preference functions of Type S .

Lemma 3.1. *If $\gcd(d, q) = 1$, then each of the matrices O and S defines a preference function of span 1; the column functions for both preference functions are bijective functions from \mathcal{A} to \mathcal{A} . Moreover, for all digits $a \in \mathcal{A}$, $O_q(a) = a$ and $S_1(a) = a$.*

Proof. For each i , the i^{th} row of O is $(i + d, i + 2d, \dots, i + qd)$, and the i^{th} row of S is $(i, i + d, i + 2d, \dots, i + (q - 1)d)$; where addition modulo q . Since $\gcd(d, q) = 1$, it is immediate that the q entries are distinct in both scenarios, verifying that both O and S are preference functions of span 1. Also in both scenarios, every column function has the form $g_j(i) = i + jd \pmod q$ for some fixed value of $j = 1, \dots, q$, which is evidently bijective. The other two statements of the lemma are obvious. \square

Theorem 3.2. *The sequence $\mathbf{o}_n = (O, 0^n)$, with $\gcd(d, q) = 1$, is a de Bruijn sequence missing only the words $1^n, \dots, (q - 1)^n$.*

Proof. We apply Theorem 2.7 with C being the self loop $0 \rightarrow 0$. By Lemma 3.1, $O_q(a) = a$ for all $a \in \mathcal{A}$, so that $\Sigma = \mathcal{A} \setminus \{0\}$ is closed under g_q . Since $O_q(0) = 0$, $O_{q-1}(0) \neq 0$. g_{q-1} is a bijection (by Lemma 3.1) so there exists $b \in \Sigma$ such that $g_{q-1}(b) = 0$, which is not in Σ . Therefore, $q' := q$ is minimal in the sense of Theorem 2.7. We now verify that $g_{q'-1}$ has no cycles when restricted to Σ . Noting that $\gcd(q, d(q-1)) = 1$, $g_{q'-1}(i) = i + (q-1)d$ induces exactly one cycle. Thus, $g_{q'-1}^q(0) = 0$. Let $c \in \Sigma$ be such $g_{q-1}(0) = c$. Iterating g_{q-1} we get

$$(c, g_{q-1}(c), g_{q-1}^2(c), \dots, g_{q-1}^{q-1}(c))$$

which forms a ‘‘path’’ whose elements are all different, exhausting Σ and with 0 at the end. This verifies that g_{q-1} has no cycles in Σ . By Theorem 2.7, the only missing words are as claimed. \square

Definition 3.3. We say that (y_1, \dots, y_n) is an alternating string with step l , $l = 1, \dots, q - 1$ if $y_{i+1} = y_i + ld \pmod q$ for $i = 1, \dots, n - 1$.

Theorem 3.4. *Let I be an alternating string with step $d(q - 1)$ and $\gcd(d, q) = 1$. Then the sequence $\mathbf{s}_n = (S, I)$ is a full de Bruijn sequence.*

Proof. Note that $S_q(a) = a + d(q - 1)$ for any $a \in \mathcal{A}$. Since $d(q - 1)$ is relatively prime to q , the induced function g_q cycles through all symbols in \mathcal{A} , i.e. words of size $s = 1$. It

follows that the initial I is on the cycle of g_q . Thus the statement follows immediately from Corollary 2.8. \square

The following gives an interpretation of the sequence \mathbf{o}_n as a prefer-opposite sequence. If we *define* the opposite of i to be $i+d \pmod q$, then in the construction of the sequence we first propose the opposite of the last produced digit. If the resulting n -word is new we accept the proposed digit. Otherwise we propose the opposite of the last rejected digit and so. We stop when q proposals for the same digit are rejected. Noting that S can be obtained from O by moving the last column of the latter to the first column, the preference function defined by S gives the highest preference to the most recent digit (hence prefer-same) and then proceeds by skipping the value d each time a proposed digit is rejected.

Two notions of *opposite* of i which are available for any q are the next digit $i+1$, and the *farthest* digit $i+q-1$. For $q=5$, these are illustrated in the top part of Table 2, along with a third case, $d=2$, in the middle.

0 \rightarrow 1, 2, 3, 4, 0	0 \rightarrow 2, 4, 1, 3, 0	0 \rightarrow 4, 3, 2, 1, 0
1 \rightarrow 2, 3, 4, 0, 1	1 \rightarrow 3, 0, 2, 4, 1	1 \rightarrow 0, 4, 3, 2, 1
2 \rightarrow 3, 4, 0, 1, 2	2 \rightarrow 4, 1, 3, 0, 2	2 \rightarrow 1, 0, 4, 3, 2
3 \rightarrow 4, 0, 1, 2, 3	3 \rightarrow 0, 2, 4, 1, 3	3 \rightarrow 2, 1, 0, 4, 3
4 \rightarrow 0, 1, 2, 3, 4	4 \rightarrow 1, 3, 0, 2, 4	4 \rightarrow 3, 2, 1, 0, 4
0 \rightarrow 0, 1, 2, 3, 4	0 \rightarrow 0, 2, 4, 1, 3	0 \rightarrow 0, 4, 3, 2, 1
1 \rightarrow 1, 2, 3, 4, 0	1 \rightarrow 1, 3, 0, 2, 4	1 \rightarrow 1, 0, 4, 3, 2
2 \rightarrow 2, 3, 4, 0, 1	2 \rightarrow 2, 4, 1, 3, 0	2 \rightarrow 2, 1, 0, 4, 3
3 \rightarrow 3, 4, 0, 1, 2	3 \rightarrow 3, 0, 2, 4, 1	3 \rightarrow 3, 2, 1, 0, 4
4 \rightarrow 4, 0, 1, 2, 3	4 \rightarrow 4, 1, 3, 0, 2	4 \rightarrow 4, 3, 2, 1, 0

TABLE 2. Top row: generalized prefer-opposite diagrams with $d=1, 2, 4$. Lower row: generalized prefer-same diagrams with $d=1, 2, 4$.

Besides this intuitive interpretation, the proposed sequences bear more resemblance to the binary counterparts, as we now present. We first focus on \mathbf{o}_n and later present similar results for \mathbf{s}_n . We recall here that the binary prefer-opposite sequence of order n has all words except 1^n , and it terminates with the substring $1^{n-1}0^{n-1}$, see Alhakim [1]. Proposition 3.6 below shows how this phenomenon generalizes in the current setting.

Lemma 3.5. *In the sequence \mathbf{o}_n , the digit a does not appear after the last occurrence of the pattern a^{n-1} for any $a \in \mathcal{A}$.*

Proof. The statement is trivial when $a=0$ because 0^{n-1} is the terminal pattern. Suppose the statement is false for some $a \neq 0$ and suppose that $x_1 \cdots x_{n-1}a$ is the first n -word that terminates with a after the last occurrence of a^{n-1} .

It follows that $x_1 \cdots x_{n-1}a$ must be followed by something, as it is not the last word. The next n -word is either $x_2 \cdots x_{n-1}aa$, or otherwise the word $x_2 \cdots x_{n-1}aa$ will occur afterwards in the sequence, because $O_q(a) = a$.

The same argument can now be iterated to establish that a^{n-1} will appear later in the sequence, thus contradicting the assumption that a^{n-1} appeared for the last time earlier. \square

Proposition 3.6. *For any integer d co-prime with q , the generalized prefer-opposite sequence \mathbf{o}_n terminates with the pattern*

$$\sigma_{q-2}^{n-1} \cdots \sigma_1^{n-1} \sigma_0^{n-1} 0^{n-1}$$

where $\sigma_0, \dots, \sigma_{q-2}$ are an appropriate re-arrangement of the digits $1, \dots, q-1$ determined by the function g_{q-1} .

Proof. Since g_{q-1} is cyclic, the terms of the sequence

$$g_{q-1}(0), g_{q-1}^2(0), \dots, g_{q-1}^q(0) = 0$$

are all distinct, where $g_{q-1}^{i+1}(0) = g_{q-1}^i(g_{q-1}(0))$. Let $a = g_{q-1}^i(0)$ and $b = g_{q-1}^{i+1}(0) = g_{q-1}(a)$ for an arbitrary $i < q-1$, so that a and b are distinct from 0. By the preference order, the last occurrence of a^{n-1} is succeeded by the digit b . We will show that it is in fact succeeded by b^{n-1} .

Let j be the largest integer such that a^{n-1} is succeeded by b^j . By Theorem 3.2, $1 \leq j \leq n-1$ so suppose $j < n-1$. It then follows that the n -word $a^i b^j$ that is a suffix of $a^{n-1} b^j$ has $i \geq 2$. Since $a^i b^j$ is not the terminal n -word of the sequence it must be succeeded by a symbol c . Since j is maximal, $c \neq b$. This puts the word $a^{i-1} b^j c$ and it follows that $a^{i-1} b^{j+1}$ will appear later in the sequence because $O_q(b) = b$. This contradicts Lemma 3.5 as $i-1 \geq 1$, therefore $j = n-1$.

Letting $\sigma_{q-1-i} = g_{q-1}^i(0)$, the previous paragraph establishes that the last occurrence of σ_{q-1-i}^{n-1} is succeeded by $\sigma_{q-1-i-1}^{n-1}$ for $i = 1, \dots, q-2$. When $i = q-1$, observe that $g_{q-1}(\sigma_0) = 0$, so the last occurrence of σ_0^{n-1} is followed by 0. At this point no nonzero symbol is allowed to appear, by Lemma 3.5. Thus the only remaining possibility is the terminal string 0^{n-1} and this finishes the proof. \square

As an example, the order n sequences generated by the generalized prefer-opposite diagrams in Table 2 terminate, respectively from left to right, with the patterns

$$4^{n-1} 3^{n-2} 2^{n-1} 1^{n-1} 0^{n-1}, 3^{n-1} 1^{n-1} 4^{n-2} 2^{n-1} 0^{n-1}, \text{ and } 1^{n-1} 2^{n-1} 3^{n-1} 4^{n-1} 0^{n-1}.$$

Remark 3.7. The case $n = 2$ yields the following interesting palindromic sequences

$$0012340241303142043210, 0024130432101234031420, 0043210314202413012340.$$

The reader is invited to check that a palindrome is obtained (with another zero appended at the end) if and only if $n = 2$ and q is a prime number.

3.1. Mapping \mathbf{o}_n to the prefer-higher sequence. The most striking similarity with the binary prefer-opposite is the relationship between the proposed sequence and the prefer-higher sequence, which is essentially the same relationship between the binary prefer-opposite and the prefer-one sequences, discussed by [14].

To describe this relationship let us first recall the so-called D -homomorphism used in Lempel [12] to recursively construct binary de Bruijn sequences. For two binary numbers b_1 and b_2 it is defined as $D(b_1, b_2) = b_1 + b_2 \pmod{2}$. It can more generally be defined as a function from $\{0, 1\}^l$ to $\{0, 1\}^{l-1}$, for any integer $l \geq 2$ as follows, $D(b_1, \dots, b_l) = (b_1 + b_2, b_2 + b_3, \dots, b_{l-1} + b_l)$, where addition is performed modulo 2. In fact, Rubin and Weiss [14] establish that if we map the binary prefer-opposite sequence

(b_0, \dots, b_{2^n-1}) via the D -homomorphism, getting $(\hat{b}_0, \dots, \hat{b}_{2^n-2})$, and then drop all bits \hat{b}_i such that $(\hat{b}_{i-n+2}, \dots, \hat{b}_i)$ is already a substring of $(\hat{b}_0, \dots, \hat{b}_{i-1})$, the remaining bits form the prefer-one sequence of order $n - 1$.

To state the q -ary counterpart of this result, we will use the generalized q -ary D -homomorphisms defined in Alhakim and Akinwande [2] as

$$D_\beta(x_1, \dots, x_l) = (\beta(x_2 - x_1), \dots, \beta(x_l - x_{l-1})),$$

where β is co-prime with q , subtraction is performed modulo q , and x_i is a digit in the q -ary alphabet \mathcal{A} . These homomorphisms were also used to generalize the Lempel construction to nonbinary alphabets, see [2].

Theorem 3.8. *Let $D_\beta(\mathbf{o}_n) = (\hat{x}_0, \dots, \hat{x}_{q^n-2})$ be the homomorphic image of the generalized prefer-opposite sequence $\mathbf{o}_n = (x_0, \dots, x_{q^n-1})$ by the homomorphism D_β , where $\gcd(d, q) = 1$ and $\beta = d^{-1}(q - 1)$, with d^{-1} being the inverse of d modulo q .*

Then the subsequence of $D_\beta(\mathbf{o}_n)$ obtained by deleting every digit \hat{x}_i such that $(\hat{x}_{i-n+2}, \dots, \hat{x}_i)$ is a substring of $(\hat{x}_0, \dots, \hat{x}_{i-1})$, is the prefer-higher de Bruijn sequence \mathbf{h}_{n-1} of order $n-1$.

To help the reader follow the proof more closely, we illustrate this result with an example. For $q = 3$, $d = 2$, \mathbf{o}_4 is

0000210212101020211022100201012120200220222122112111011001000120122011200111222000

As $\beta = 1$, the appropriate homomorphism is $D_1(x_1, x_2) = x_2 - x_1$, giving the image sequence $D_1(\mathbf{o}_4)$:

00022221221221220220220211211211210201200210201200210201200111110110110100100100

To best present how the prefer-higher sequence of order 3 emerges from the latter sequence, we write it as a sequence of words and cross out any repetition, keeping only the first occurrence of every word.

000 \rightarrow 002 \rightarrow 022 \rightarrow 222 \rightarrow ~~222~~ \rightarrow ~~222~~ \rightarrow 221 \rightarrow 212 \rightarrow 122 \rightarrow ~~221~~ \rightarrow ~~212~~ \rightarrow ~~122~~ \rightarrow ~~221~~ \rightarrow ~~212~~ \rightarrow ~~122~~ \rightarrow 220 \rightarrow 202 \rightarrow ~~022~~ \rightarrow ~~220~~ \rightarrow ~~202~~ \rightarrow ~~022~~ \rightarrow ~~220~~ \rightarrow ~~202~~ \rightarrow 021 \rightarrow 211 \rightarrow 112 \rightarrow 121 \rightarrow ~~211~~ \rightarrow ~~112~~ \rightarrow ~~121~~ \rightarrow ~~211~~ \rightarrow ~~112~~ \rightarrow ~~121~~ \rightarrow 210 \rightarrow 102 \rightarrow 020 \rightarrow 201 \rightarrow 012 \rightarrow 120 \rightarrow 200 \rightarrow ~~002~~ \rightarrow ~~021~~ \rightarrow ~~210~~ \rightarrow ~~102~~ \rightarrow ~~020~~ \rightarrow ~~201~~ \rightarrow ~~012~~ \rightarrow ~~120~~ \rightarrow ~~200~~ \rightarrow 001 \rightarrow 011 \rightarrow 111 \rightarrow ~~111~~ \rightarrow ~~111~~ \rightarrow 110 \rightarrow 101 \rightarrow ~~011~~ \rightarrow ~~110~~ \rightarrow ~~101~~ \rightarrow ~~011~~ \rightarrow ~~110~~ \rightarrow ~~101~~ \rightarrow 010 \rightarrow 100 \rightarrow ~~001~~ \rightarrow ~~010~~ \rightarrow ~~100~~ \rightarrow ~~001~~ \rightarrow ~~010~~ \rightarrow ~~100~~

The uncrossed words can be written compactly as the sequence

$$00022212202112102012001110100,$$

which is the prefer higher sequence of order 3.

Definition 3.9. We say that (y_1, \dots, y_n) is a translate of (x_1, \dots, x_n) if there exists $c \in \mathcal{A}$ such that $y_i = x_i + c \pmod q$ for $i = 1, \dots, n$.

Lemma 3.10. *Given two n -tuples \mathbf{v} and \mathbf{w} , $D_\beta(\mathbf{v}) = D_\beta(\mathbf{w})$ if and only if \mathbf{w} is a translate of \mathbf{v} .*

Lemma 3.11. *Every $(n - 1)$ -tuple has exactly q inverse images by D_β .*

Proof. Let $\mathbf{w}_{n-1} = (\hat{x}_1, \dots, \hat{x}_{n-1})$. For each x_1 in \mathcal{A} , define $x_i = x_{i-1} + \beta^{-1}\hat{x}_{i-1}$, $i = 2, \dots, n$. One can readily check that $D_\beta(x_1, \dots, x_n) = \mathbf{w}_{n-1}$. \square

In general, there exist $q/\gcd(q, l)$ alternating strings with step l ; see Definition 3.3, which are all translates of each other. Observe that the q alternating strings with step $l = 1$ occur consecutively in the beginning of the sequence. Interestingly, alternating strings of higher steps also aggregate together.

Lemma 3.12. *When the first alternating string with step l ; $l > 1$ occurs, all of its translates follow consecutively.*

Proof. Let $l = \hat{l}d$, where \hat{l} is between 1 and $q - 1$, inclusive. Due to the form of the preference matrix $O^{(q,d)}$, an alternating string with step l corresponds to a cycle of $g_{\hat{l}}$. That is, when the first digit of an alternating string with step l is given, all other digits are entered via the \hat{l}^{th} preference (the 1st preference being the highest). By the prefer-opposite strategy, the first alternating string is achieved when all words that have combinations of preference steps less than or equal to \hat{l} , but not purely \hat{l} , have been used. Immediately after this, preference step \hat{l} will be repeated until a digit of preference \hat{l} cannot be added without putting an already present word. This is the case when a cycle of the function $g_{\hat{l}}$ is exhausted.

There are two cases. If \hat{l} is co-prime with q , then $g_{\hat{l}}$ is periodic with period q , in which case all alternating strings of step l are placed consecutively. Otherwise, preference $\hat{l} + 1$ is used once and then \hat{l} is used again, that is, another cycle of $g_{\hat{l}}$ is entered, and so on until all cycles of $g_{\hat{l}}$ are placed, thus producing all alternating strings with step l . \square

Lemma 3.13. *A word $\mathbf{w} = (x_1, \dots, x_n)$ is succeeded by the digit $x_n + jd$ for some $j = 0, 1, \dots, q-1$ and d that is co-prime with q , if and only if $D_\beta(x_1, \dots, x_n)$ is succeeded by $q - j \pmod q$.*

Proof. $D_\beta(x_1, \dots, x_n) = (\beta(x_2 - x_1), \dots, \beta(x_n - x_{n-1}))$. If $x_n + jd$ succeeds \mathbf{w} , $D_\beta(\mathbf{w})$ is succeeded by digit $\beta(x_n + jd - x_n) = \beta jd$. Using the value of β we get $d^{-1}(q-1)jd = j(q-1) = q - j$.

Conversely, suppose $D_\beta(x_1, \dots, x_n)$ is succeeded by $q - j \pmod q$ and let y be the digit that succeeds \mathbf{w} . That is, $\beta(y - x_n) = q - j$. This implies that $y - x_n = \beta^{-1}(q - j) = d(q-1)(q-j) = dj \pmod q$ and the result follows. \square

Lemma 3.13 provides the intuition behind the result of Theorem 3.8. In the prefer-opposite sequence, the q appearances of an $(n-1)$ -tuple (x_1, \dots, x_{n-1}) in \mathbf{o}_n are succeeded respectively by $x_{n-1} + d, x_{n-1} + 2d, \dots, x_{n-1} + qd = x_{n-1}$. Their images by D_β are followed respectively by $q-1, \dots, 1, 0$, i.e., according to the prefer-higher strategy. However, the problem is more complicated. Using Lemma 3.10, we see that a translate of the former n -tuples can potentially occur in \mathbf{o}_n in the wrong location so as to mess up the prefer-higher order.

Furthermore, another main issue is to verify that, upon crossing out a sequence of one or more consecutive words, the last word before this crossed sequence teams up correctly with the next uncrossed word.

In the binary case, Rubin and Weiss [14] carry out a meticulous study to inspect various cases of which translates of a word occur (in this case the only translate of a word is its bitwise complement.) Our proof will apply the principle of strong mathematical induction, which greatly simplifies the discussion.

Proof of Theorem 3.8. Observe that the inverse images of $\mathbf{0}_{n-1}$ are all words of the form c^n . By Theorem 3.2, 0^n is the only inverse image of 0^{n-1} that occurs in \mathbf{o}_n . Moreover, any word that is not of the form c^n occurs in \mathbf{o}_n along with all of its translates.

By Lemmas 3.10 and 3.11, the collection of inverse images of all $n - 1$ tuples is the collection of n tuples. It follows that $D_\beta(\mathbf{o}_n)$ includes every $n - 1$ tuple exactly q times, except for 0^{n-1} that appears once in the beginning.

Hence, there will be exactly q^{n-1} uncrossed digits. We let $I = (i_1, i_2, \dots, i_{q^{n-1}})$ be the subsequence of $1, \dots, q^n - 1$ consisting of the indices in $\hat{\mathbf{o}}_{n-1}$ that remain after crossing repetitions. That is, $i_t \in I$ if and only if $(\hat{x}_{i_t}, \dots, \hat{x}_{i_t+n-2}) \notin \{(\hat{x}_j, \dots, \hat{x}_{j+n-2}) : j = 1, \dots, i_t - n + 1\}$.

Let us also represent the prefer-higher sequence \mathbf{h}_{n-1} as $(y_i; i = 0, \dots, q^{n-1} - 1)$, and define $\mathbf{x}_i := (x_i, \dots, x_{i+n-1})$, $\hat{\mathbf{x}}_i := (\hat{x}_i, \dots, \hat{x}_{i+n-2})$ and $\mathbf{y}_i := (y_j, \dots, y_{j+n-2})$ for $i \geq 0$ to be the i^{th} word of \mathbf{o}_n , $D_\beta(\mathbf{o}_n)$ and \mathbf{h}_{n-1} respectively. Note that $D_\beta(\mathbf{x}_i) = \hat{\mathbf{x}}_i$.

We need to verify that (1) for each t the last $n - 2$ digits $\hat{\mathbf{x}}_{i_t}$ coincide with the first $n - 2$ digits of $\hat{\mathbf{x}}_{i_{t+1}}$, and (2) the prefer higher strategy is respected. A moment's reflection shows that it is sufficient to show that $\hat{\mathbf{x}}_{i_t} = \mathbf{y}_t$ for all t ; $0 \leq t \leq q^{n-1} - 1$. Observing that this statement is true for $t = 0$, we assume that it is true for all $j \leq t$, where $0 \leq t < q^{n-1} - 1$ and proceed to prove it is true for $t + 1$.

First consider the case when $i_{t+1} = i_t + 1$. Suppose that $\hat{\mathbf{x}}_{i_{t+1}}$ ends with the digit $q - j$ for some j , then by Lemma 3.13 $\mathbf{x}_{i_{t+1}}$ has the form $(x_{i_{t+1}}, \dots, x_{i_{t+1}+n-1}, x_{i_{t+1}+n-1} + jd)$. By the prefer-opposite strategy, the $j - 1$ words $(x_{i_{t+1}}, \dots, x_{i_{t+1}+n-1}, x_{i_{t+1}+n-1} + ld)$, $1 \leq l < j$ occur earlier in \mathbf{o}_n . Since they all agree with $\mathbf{x}_{i_{t+1}}$ on the first $n - 1$ digits, their images by D_β are of the form $(\hat{x}_{i_{t+1}}, \dots, \hat{x}_{i_{t+1}+n-2}, q - l)$ and they all occur earlier. To see that $\hat{\mathbf{x}}_{i_{t+1}}$ will not be crossed out, suppose that it did occur as \mathbf{y}_s , $s \leq t$. So its first occurring inverse image in \mathbf{o}_n must be \mathbf{x}_{i_s} , by the induction hypothesis, so that $\mathbf{x}_{i_{t+1}}$ is not new, contradicting the fact that $i_t + 1 = i_{t+1}$.

When $i_{t+1} > i_t + 1$, $i_{t+1} - 1 \notin I$ and $D_\beta(\mathbf{x}_{i_{t+1}-1}) = \mathbf{y}_\tau$ for some $\tau \leq t$. By the inductive hypothesis this means that $\mathbf{y}_\tau = D_\beta(\mathbf{x}_{i_\tau})$. There are q possible $(n - 1)$ -tuples that can logically succeed \mathbf{y}_τ : $\mathbf{w}_l = (y_{\tau+1}, \dots, y_{\tau+n-2}, q - k)$; $k = 1, \dots, q$.

Since $i_\tau \leq i_t$, $i_\tau + 1 \leq i_{t+1} - 1$ so that $i_{\tau+1} \notin I$. Thus at least one of the \mathbf{w}_l occurs in $\hat{\mathbf{x}}_{i_0}, \dots, \hat{\mathbf{x}}_{i_t}$. Let j , $1 \leq j \leq q$ be the maximum integer such that \mathbf{w}_j occurs. Then by the inductive hypothesis, all \mathbf{w}_l , $l \leq j$ occur before since so far $\hat{\mathbf{x}}_{i_0}, \dots, \hat{\mathbf{x}}_{i_t}$ coincide with $\mathbf{y}_0, \dots, \mathbf{y}_t$. This also implies that $j < q$, for otherwise $i_{t+1} \notin I$ (as $D_\beta(\mathbf{x}_{i_{t+1}})$ has to go to one of the \mathbf{w}_l and has to be new.) It also follows that $D_\beta(\mathbf{x}_{i_{t+1}})$ is \mathbf{w}_{j+1} , for if it equals \mathbf{w}_k for some $k > j + 1$, $\mathbf{x}_{i_{t+1}}$ would have the form $(x_{i_{t+1}}, \dots, x_{i_{t+1}+n-2}, x_{i_{t+1}+n-2} + kd)$. The prefer-opposite strategy would then mean that $(x_{i_{t+1}}, \dots, x_{i_{t+1}+n-2}, x_{i_{t+1}+n-2} + (j + 1)d)$ occurs earlier in \mathbf{o}_n and therefore \mathbf{w}_{j+1} occurs earlier in $D_\beta(\mathbf{o}_n)$, contradicting the maximality of j .

To sum up, the prefer-opposite again implies that the n -tuples

$$(3.1) \quad \mathbf{x}_{\kappa_l} = (x_{i_{t+1}}, \dots, x_{i_{t+1}+n-2}, x_{i_{t+1}+n-2} + ld), 1 \leq l \leq j$$

all occur earlier in \mathbf{o}_n , and $\{D_\beta(\mathbf{x}_{\kappa_l}) : 1 \leq l \leq j\} = \{\mathbf{w}_1, \dots, \mathbf{w}_j\}$. The \mathbf{x}_{κ_l} are evidently all distinct, and there are two possibilities.

Case 1. Suppose $\mathbf{x}_{i_\tau} \neq \mathbf{x}_{\kappa_l}$ for any $l = 1, \dots, j$. If $\kappa_1, \dots, \kappa_j, i_\tau$ are all strictly less than i_t then their images by D_β are $\mathbf{w}_1, \dots, \mathbf{w}_{j+1}$, contradicting that $i_{t+1} \in I$. Thus $i_t \in \{\kappa_1, \dots, \kappa_j, i_\tau\}$, and by Equation (3.1) and the prefer-opposite, $i_\tau = i_t$.

Case 2. If $\mathbf{x}_{i_\tau} = \mathbf{x}_{\kappa_l}$ for some l , then \mathbf{x}_{i_τ} has the form of an alternating string with step l . Since $\tau \in I$, \mathbf{x}_{i_τ} is the first such alternating string. By Lemma 3.12, all alternating strings of step l and that are translates of each other occur consecutively. This forces the last of these strings to be $\mathbf{x}_{i_{t+1}-1}$ and thus forces i_τ to be i_t , i.e. $t = \tau$. □

4. DISCREPANCY CALCULATIONS

Cooper and Heitsch [6] define the discrepancy of a binary sequence $w = w_1 \cdots w_n$ as $disc(w) = \max_M \left| \sum_{k=0}^M (-1)^{w_k} \right|$, where the maximum is taken over all initial strings of length M . In fact, they prove that the discrepancy of the prefer-one sequence of word size n is of order $\Theta(2^n \ln n/n)$, much larger than that of a typical de Bruijn sequence. This is the case because the prefer-one scheme biases the initial part of the sequence (started with 0^n) to having more ones than zeros.

Using a variant of the above discrepancy formula, Gabric and Sawada [10] numerically studied the discrepancy of a number of binary de Bruijn sequences, including the prefer-same and prefer-opposite sequences which come out to have a low discrepancy. In this section we define a measure of discrepancy for sequences of general alphabet size and we use it to compare the prefer-higher sequence and the two proposed sequences with the binary versions.

Definition 4.1. Given is a finite sequence s from an alphabet \mathcal{A} of size q . For a letter $a \in \mathcal{A}$, we write $f_a(i)$ to denote the frequency of a in the initial subsequence of size i . The discrepancy of s is defined as

$$discrep(s) = \max_i \left(\max_{a,b} |f_a(i) - f_b(i)| \right),$$

where the inner maximum is taken over all pairs of digits a and b , while the outer maximum is taken over all initial subsequences of size i of the sequence.

We observe that this discrepancy, when applied to a de Bruijn sequence (or any periodic sequence), is a function of the initial word. When $q = 2$, this definition boils down to the formula given in [6].

Gabric and Sawada [10] conjecture that the prefer-opposite and prefer-same sequences have a discrepancy in the order of $\Theta(n^2)$. Table 3 below displays the calculated discrepancy for all three sequences for $q = 2, \dots, 5$ and for several values of n , when each sequence starts with its natural, cyclic initial word. Notably, both proposed sequences

Discrepancy(pref-same, pref-opp, pref-higher)												
n	$q = 2$			$q = 3$			$q = 4$			$q = 5$		
2	1	2	2	2	2	2	3	3	3	3	2	4
3	3	3	3	3	4	5	4	5	9	6	4	16
4	3	5	4	5	7	12	6	9	31	9	9	70
5	5	7	5	8	12	30	9	16	112	11	16	310
6	6	10	6	10	19	84	13	25	391	14	25	1346
7	10	13	14	16	27	227	18	36	1407	17	36	5807
8	12	17	28	21	37	618	23	49	5152	22	49	25994
9	16	21	56	25	48	1734	29	64	18955	-	-	-
10	20	26	110	31	61	4847	-	-	-	-	-	-
11	26	31	211	38	75	13532	-	-	-	-	-	-
12	30	37	404	44	91	38456	-	-	-	-	-	-
13	36	43	771	-	-	-	-	-	-	-	-	-
14	42	50	1474	-	-	-	-	-	-	-	-	-
15	50	57	2809	-	-	-	-	-	-	-	-	-

TABLE 3. The discrepancy of the three sequences are reported for various values of q and n using a Matlab code. Dashed entries were left as they take too long.

have discrepancies that are very similar to their binary counterparts, regardless of the value of q and n . Also, the table strongly suggests that the discrepancy of both sequences are dominated by $(n - 1)^2$ while that of the prefer-higher sequence grows essentially exponentially, for various values of q . One can even guess the patterns for the first two sequences. When $q = 2$ for instance, the discrepancy of the pref-same, for n high enough, is

$$discrep(\text{pref-same}(n)) = \begin{cases} n(n - 2)/4 & n \text{ is even} \\ (n - 1)^2/4 & (n - 1) \pmod{4} = 0, \\ (n - 1)^2/4 + 1 & (n - 1) \pmod{4} = 2 \end{cases},$$

while that of the prefer-opposite appears to be

$$discrep(\text{pref-opp}(n)) = \begin{cases} (\frac{n}{2})^2 + 1 & n \text{ is even} \\ (\frac{n-1}{2})^2 + \frac{n-1}{2} + 1 & n \text{ is odd} \end{cases}.$$

Curiously also, $discrep(\text{pref-opp}(n)) = (n - 1)^2$ for n high enough when $q = 4, 5$.

5. CONCLUSION

Relying on well-established techniques of preference functions, we introduced two nonbinary sequences with properties that parallel the well-known binary prefer-opposite and prefer-same sequences. The latter two sequences were recently generated efficiently via a simple successor rule, see Sala et al [15]. Generating the new sequences efficiently is planned as future research. Also of interest is an analytical study of the discrepancy of these sequences.

REFERENCES

- [1] A. Alhakim. A Simple Combinatorial Algorithm for de Bruijn Sequences. *The American Mathematical Monthly*, **117**, Number 8, (2010) 728-732.
- [2] Alhakim, A., Akinwande, M. A recursive construction of nonbinary de Bruijn sequences. *Des. Codes Cryptogr.* 60, 155-169 (2011).
- [3] A. Alhakim. Spans of preference functions. *Discrete Applied Mathematics*, Vol. 160, 7-8, (2012) 992-998.
- [4] A. Alhakim, E. Sala, J. Sawada. Revisiting the prefer-same and prefer-opposite de Bruijn Sequence Constructions. *Theoretical Computer Science*, (2020).
- [5] A. Alhakim, Designing preference functions for de Bruijn sequences with forbidden words. Submitted manuscript.
- [6] J. Cooper, C. Heitsch. The Discrepancy of the Lex-Least De Bruijn Sequence. *Discrete Math*, 310, 6-7, (2010) 1152-1159.
- [7] C. Eldert, H. J. Gray, H. M. Gurk, M. Rubinoff. Shifting Counters. *AIEE Trans.*, 77, (1958), 70-74.
- [8] L. R. Ford, A Cyclic Arrangement of m -tuples, Report P-1071, Rand Corp., 1957.
- [9] H. Fredricksen, A Survey of Full Length Nonlinear Shift Register Cycle Algorithms, *SIAM Review*, **24** (1982) 195-221.
- [10] D. Gabric, J. Sawada, Investigating the discrepancy property of de Bruijn sequences, *Discrete Mathematics Volume 345*, Issue 4, April 2022.
- [11] S. Golomb, *Shift Register Sequences*, Holden-Day, San Francisco, 1967.
- [12] A. Lempel, On a homomorphism of the de Bruijn graph and its applications to the design of feedback shift registers. *IEEE Trans. Comput.* 100(12), 1204-1209 (1970).
- [13] M. H. Martin, A Problem in Arrangements, *Bulletin of the American Mathematical Society*, **40** (1934) 859-864.
- [14] Rubin, A., Weiss, G. Mapping prefer-opposite to prefer-one de Bruijn sequences. *Des. Codes Cryptogr.* 85, 547-555 (2017).
- [15] Evan Sala, E., Sawada, J., Alhakim, A. Efficient constructions of the Prefer-same and Prefer-opposite de Bruijn sequences. Submitted Manuscript.

Email address: aa145@aub.edu.lb