

On the harmonious chromatic number of graphs

Gabriela Araujo-Pardo* Juan José Montellano-Ballesteros* Mika Olsen†
Christian Rubio-Montiel‡

November 18, 2022

Abstract

The harmonious chromatic number of a graph G is the minimum number of colors that can be assigned to the vertices of G in a proper way such that any two distinct edges have different color pairs. This paper gives various results on harmonious chromatic numbers related to homomorphisms, the incidence graph of finite linear systems, and the known circulant graphs.

Keywords. Harmonious homomorphism, Levi graph, circulant graph, diameter two.

Mathematics Subject Classification: 05C15, 05B25.

1 Introduction

A *harmonious coloring* of a finite graph G is a proper k -coloring such that every pair of colors appears on at most one edge. The *harmonious chromatic number* $h(G)$ of G is the minimum number k such that G has a harmonious k -coloring.

This parameter was introduced by Mitchem [14], which is a slight variation of the original definition given independently by Hopcroft and Krishnamoorthy [11] and by Frank, Harary, and Plantholt [8].

From the definition, a graph G of order n has size $m \leq \binom{h(G)}{2}$, which in turn gives the easy lower bound

$$\sqrt{2m + \frac{1}{4}} + \frac{1}{2} \leq h(G). \quad (1)$$

In [14] and [15] the authors give families of graphs that show that the lower bound (1) is tight.

It is easy to see that a proper n -coloring of G is harmonious, however, the harmonious chromatic number has an additional difficulty that it makes impossible to use usual greedy colorings to complete partial colorations even if extra colors are allowed: if we have an uncolored vertex adjacent to two different vertices with the same color, it is not possible to obtain a harmonious coloring.

Therefore, if G has order n and diameter at most two, then $h(G) = n$, see [5], for instance, complete graphs, complete bipartite graphs, the line graph of the complete graph, the Erdős-Rényi orthogonal polarity graph, or a graph with a universal vertex (stars, wheels, or connected trivially perfect graphs, etc.) have harmonious number equals to n . In the survey, [5], we can find a list of interesting results about the harmonious chromatic number and the achromatic number, a related parameter, also see Chapters 12 and 13 of [3].

The purpose of this paper is to study the harmonious chromatic number. The paper is organized as follows. In Section 2 is given results arising from homomorphisms; in Section 3 we study the harmonious chromatic number of graphs with small diameter related to incidence graphs of linear systems; and finally, in Section 4 we analyze the harmonious chromatic number of the circulant graphs.

*Instituto de Matemáticas, Universidad Nacional Autónoma de México, Mexico City, Mexico. [garaujo|juancho]@math.unam.mx.

†Departamento de Matemáticas Aplicadas y Sistemas, UAM-Cuajimalpa, Mexico City, Mexico. olsen@correo.cua.uam.mx.

‡División de Matemáticas e Ingeniería, FES Acatlán, Universidad Nacional Autónoma de México, Naucalpan, Mexico. christian.rubio@acatlan.unam.mx.

2 On homomorphisms

Let G be a graph and ς a harmonious $h(G)$ -coloring of G . Take a couple of vertices x, y in the same color class, then $d(x, y)$ is at least 3. An elementary homomorphism ϵ is obtained by identifying x and y , therefore $h(G) \leq h(\epsilon(G))$. We can repeat this process in the graph $\epsilon(G)$ with the coloring inherited from ς . A graph H of diameter two can be obtained by a sequence of these elementary homomorphisms beginning with G , see [5].

On the other hand, let G be a graph and take a couple of vertices x, y such that $d(x, y) \geq 3$. An elementary homomorphism ϵ is obtained by identifying x and y . We can repeat this process in the graph $\epsilon(G)$. A graph H of diameter two can be obtained by a sequence of these elementary homomorphisms beginning with G . The composition of the elementary homomorphisms is an homomorphism f that we can obtain by an appropriate partition $P = \{V_1, V_2, \dots, V_k\}$ of G where $V_i = f^{-1}(v_i)$ for each v_i a vertex of $f(G)$. Each set V_i is an independent set and every two sets V_i and V_j only share at most an edge, otherwise, if there are vertices $x, y \in V_i$ and $a, b \in V_j$ such that $xa, yb \in E(G)$, then some elementary homomorphism identifies x and y , a contradiction because $f(a) = f(b)$. We say f is a *harmonious homomorphism*. Therefore, the harmonious k -coloring of G corresponds exactly to the harmonious homomorphisms from G to some graph of diameter two with k vertices.

Remark 1. *A function f defines a harmonious homomorphism to some graph of diameter two with k vertices if and only if f maps vertices at distance at most 2 of G to different colors via a sequence of elementary harmonious homomorphisms.*

The following result is straightforward.

Remark 2. *If H is a harmonious homomorphic image of a graph G , then $h(G) \leq h(H)$.*

Theorem 3. *Let u and v be vertices in a graph G such that $d(u, v) \geq 3$ and let ϵ be the elementary harmonious homomorphism of G that identify the vertices u and v . Then*

$$h(\epsilon(G)) = h(G)$$

if and only if there exists an $h(G)$ -coloring of G such that u and v have the same color.

Proof. First, suppose that $h(\epsilon(G)) = h(G) = k$. Let ς' be a k -coloring of $\epsilon(G)$. Then the coloring which gives the same color to a vertex x of G that ς' assigns to x in $\epsilon(G)$ is a k -coloring of G . Because $h(G) = k$, such $h(G)$ -coloring of G gives the same color to u and v .

Now, suppose first that there is an $h(G)$ -coloring ς of G in which u and v are assigned the same color. Then the coloring ς' of $\epsilon(G)$ defined by

$$\varsigma'(x) = \begin{cases} \varsigma(x) & \text{if } x \in V(G) \setminus \{u, v\} \\ \varsigma(u) = \varsigma(v) & \text{if } x \in \{u, v\} \end{cases}$$

is an $h(G)$ -coloring of $\epsilon(G)$. Thus $h(\epsilon(G)) \leq h(G)$. By Remark 2, $h(G) \leq h(\epsilon(G))$ and the result follows. \square

Theorem 4. *If ϵ is an elementary harmonious homomorphism identifying nonadjacent vertices u and v of G such that $d(u, v) \geq 3$, then*

$$h(G) \leq h(\epsilon(G)) \leq h(G) + \min\{\deg(u), \deg(v)\} + 1.$$

Therefore, $h(G) \leq h(\epsilon(G)) \leq h(G) + \Delta(G) + 1$.

Proof. By Remark 2, $h(G) \leq h(\epsilon(G))$. Suppose that $h(G) = k$ and a k -coloring of G is given with the colors $1, 2, \dots, k$ and $\deg(u) = \min\{\deg(u), \deg(v)\}$ with $N(u) = \{u_1, u_2, \dots, u_{\deg(u)}\}$. Define a coloring ς' of $\epsilon(G)$ by

$$\varsigma'(x) = \begin{cases} \varsigma(x) & \text{if } x \in V(G) \setminus N(u) \setminus \{u, v\} \\ k + 1 & \text{if } x \in \{u, v\} \\ k + i + 1 & \text{if } x = u_i \end{cases}.$$

Since ζ' is a $(k + \deg(u) + 1)$ -coloring of $\epsilon(G)$, it follows that $h(\epsilon(G)) \leq k + \deg(u) + 1 = h(G) + \min\{\deg(u), \deg(v)\} + 1$. \square

Theorem 5. *Let G be a graph with nonadjacent vertices u and v in G such that $d(u, v) \geq 3$. Then*

$$h(G) \leq h(G + uv) \leq h(G) + 1.$$

Proof. Clearly, $h(G + uv) \geq h(G)$. On the other hand, take an $h(G)$ -coloring ζ of G . If $\zeta(u) = \zeta(v)$, we can recolor the vertex u with a new color and add the edge uv . If $\zeta(u) \neq \zeta(v)$ and then there are two possibilities: v has a neighbor colored $\zeta(u)$, hence u has not a neighbor colored $\zeta(v)$ and we can recolor the vertex u with a new color and we can add the edge uv . Otherwise, v has not a neighbor colored $\zeta(u)$ and we can recolor the vertex v with a new color and add the edge uv . \square

Beginning with a graph G , we can always perform a sequence of elementary harmonious homomorphisms until arriving at some graph of diameter two H with k vertices. As we saw, the graph H obtained in this manner corresponds to a harmonious k -coloring of G . Consequently, we have the following.

Definition 6. *The largest order of a graph of diameter two that is a harmonious homomorphic image of a graph G is the harmonious achromatic number of G , denoted by $ha(G)$.*

We have seen that for every graph G of order n ,

$$h(G) \leq ha(G) \leq n.$$

A *complete coloring* of G is a proper k -coloring such that every pair of colors appears on at least one edge. The *achromatic number* $\alpha(G)$ of G is the maximum number k such that G has a complete k -coloring.

In the case of the chromatic and the achromatic number there is an interpolation result [9], namely, a graph G has a complete proper k -coloring if and only if $\chi(G) \leq k \leq \alpha(G)$. By giving a simple example, we show that this interpolation result fails in the case of the harmonious chromatic number and the harmonious achromatic number.

Theorem 7. *Let $r \geq 3$ be a positive integer. Then there exists a graph G_r which has $h(G_r) = r + 1$ and $ha(G_r) = h(G_r) + r - 2$ but no harmonious homomorphism f exists such that $f(G_r) = k$ with*

$$h(G_r) < k < ha(G_r).$$

Proof. Let G_r be the union of the star graphs $L = K_{1, r-1}$ and $R = K_{1, r-1}$. If ϵ is an elementary harmonious homomorphism of G_r that identify the vertices u and v such that $d(u, v) \geq 3$, then u and v belong to different components. Suppose that u_i is a leaf of L for $i \in \{1, 2, \dots, r-1\}$ and u_0 is the no leaf of L , while v_j is a leaf of R for $j \in \{1, 2, \dots, r-1\}$ and v_0 is the no leaf of R .

1. ϵ identifies u_0 and v_0 . In this case, $\epsilon(G_r)$ is a star graph $K_{1, r-2}$ and then $ha(G_r) = 2r - 1$.
2. ϵ identifies u_0 and v_j for some $j \in \{1, 2, \dots, r-1\}$. Suppose that $j = r - 1$. In this case, any other elementary harmonious homomorphism has to identify some leaf u_i with some leaf v_j . Without loss of generality, ϵ_i identify u_i and v_i for $i \in \{1, 2, \dots, r-2\}$ resulting a graph with a leaf, $r - 2$ vertices of degree two and the vertices u_0 of degree r and v_0 of degree $r - 1$. The composition of the elementary harmonious homomorphism induces a harmonious homomorphism f , then $h(f(G_r)) = r + 1$.
3. The case of ϵ identifying u_i and v_0 for some $i \in \{1, 2, \dots, r-1\}$ is similar to the previous case.
4. ϵ identifies u_i and v_j , say $i = j = 1$. We have the previous case if we produce a new, harmonious homomorphism that identifies u_0 of v_0 . Hence, any other elementary harmonious homomorphism has to identify some leaf u_i with some leaf v_j . Without loss of generality, ϵ_i identify u_i and v_i for $i \in \{2, \dots, r-1\}$. The composition of the elementary harmonious homomorphism induces a harmonious homomorphism f' , then $h(f'(G_r)) = r + 1$. Therefore, $h(G_r) = r + 1$.

□

The complement \overline{G} of a graph G is the graph with vertex set $V(G)$ and two of its vertices u and v are adjacent if and only if u is not adjacent to v in G .

Theorem 8. *If ϵ is an elementary harmonious homomorphism of a graph G , then*

$$h(\overline{G}) - 1 \leq h(\overline{\epsilon(G)}) \leq h(\overline{G}).$$

Proof. To begin with, we prove that $h(\overline{G}) - 1 \leq h(\overline{\epsilon(G)})$. Suppose that $h(\epsilon(G)) = k$ for some elementary harmonious homomorphism ϵ of G that identifies two vertices u and v in G such that $d(u, v) \geq 3$. Denote by u the vertex in $\overline{\epsilon(G)}$ obtained by identifying u and v . Suppose that a harmonious k -coloring of $\overline{\epsilon(G)}$ is using the colors $1, 2, \dots, k$ such that the vertex u in $\overline{\epsilon(G)}$ has the color k . Clearly, no vertex in $\overline{\epsilon(G)}$ adjacent to u is colored k . Color the vertices of \overline{G} as follows: each vertex distinct from v has the same color assigned to that vertex in $\overline{\epsilon(G)}$ and assign v the color $k + 1$. Since no vertex in \overline{G} adjacent to u is assigned the color k , this produces a harmonious $(k + 1)$ -coloring of \overline{G} and so $h(\overline{G}) \leq k + 1 = h(\overline{\epsilon(G)}) + 1$ and the inequality follows.

Now, we show that $h(\overline{\epsilon(G)}) \leq h(\overline{G})$. Suppose that $h(\overline{G}) = l$ and let $P = \{V_1, V_2, \dots, V_l\}$ be a partition of $V(\overline{G})$ into l independent sets in G and suppose that ϵ is an elementary harmonious homomorphism of G for which $\epsilon(u) = \epsilon(v)$ for vertices u and v of G at distance at least 3. Due to u and v are not adjacent in G , then u and v are adjacent in \overline{G} and without loss of generality $u \in V_1$ and $v \in V_l$. Let $V'_1 = V_1 - \{u\}$ and $V'_i = V_i$ for $i \in \{2, \dots, l\}$. Then $P' = \{V'_1, V'_2, \dots, V'_l\}$ is a partition of $V(\overline{\epsilon(G)})$ into $l - 1$ classes if $V'_1 = \emptyset$, or l subsets if $V'_1 \neq \emptyset$. Moreover, none of $V'_2, V'_3, \dots, V'_{l-1}$ contains v , hence these sets are independent in $\overline{\epsilon(G)}$. The set V'_l is independent in $\overline{\epsilon(G)}$ because, if x and y are vertices in $V'_l - \{v\}$, then x and y are not adjacent in V'_l since x and y are not adjacent in V_l ; and if v and w are two adjacent vertices in V'_l , it follows that v and w are not adjacent in $\epsilon(G)$. Therefore, u and w are not adjacent in G , and also v and w are not adjacent in G . Therefore, v and w are adjacent in \overline{G} , a contradiction. In consequence, P' is a partition of $V(\overline{\epsilon(G)})$ into $l - 1$ or l independent sets. A similar argument works to obtain that the partition P' induces a harmonious coloring. We only have to verify that there is at most one edge between V'_i and V'_j . Suppose that v and w are vertices in V'_i and there are vertices x and y in V'_j for some $i \in \{1, 2, \dots, l - 1\}$ such that v is adjacent to x and w is adjacent to y . Therefore, v and x are not adjacent in $\epsilon(G)$, and hence both v and x are not adjacent in G and u and x are not adjacent in G . Then v is adjacent to x in \overline{G} and also y is adjacent to w in \overline{G} , a contradiction because, P induces a harmonious coloring in \overline{G} . Thus, the result follows. □

Since $ha(G) \leq n$ for any graph G of order n , we deduce that

$$ha(G) + ha(\overline{G}) \leq 2n.$$

In [13] it was proved that

$$n + 1 \leq h(G) + h(\overline{G}) \leq 2n.$$

Moreover, for any $n \geq 7$ and $r \in \{1, 2, \dots, n\}$, there exists a graph G of order n such that $h(G) + h(\overline{G}) = n + r$.

3 On graphs of small diameter

Consider a graph H of order n and diameter 2 such that deleting any edge increases the diameter, such graph is called a diameter 2-critical graph. Clearly, $G = H - e$ for any edge e of H is a graph of diameter 3 such that $h(G) = n - 1$. We have the following question: Can there be an arbitrarily large difference between $h(G)$ and n for a graph G of order n and diameter 3? Theorem 9 shows a graph of order n and diameter 3 such that its harmonious chromatic number is $\theta(\sqrt{n})$.

Theorem 9. *Let $r, s \geq 3$ be positive integers. There exists a graph G of order n and diameter 3 such that $n = r(s + 1)$ and $h(G) = r + s$. In particular, if $r = s$, then $h(G) = 2\sqrt{n} + \theta(1)$.*

Proof. Let $G_{r,s}$ be the graph obtained by adding s leaves to each vertex of the complete graph K_r . A vertex in the complete subgraph K_r has degree $r + s - 1$ in $G_{r,s}$, thus $h(G_{r,s}) \geq r + s$. Two leaves sharing a vertex are at distance two, otherwise, the vertices of degree one are at distance 3, hence the diameter is 3 and a harmonious homomorphism can be done by identifying r leaves at distance 3 to obtain a graph of order $r + s$ and diameter two, then $h(G_{r,s}) \leq r + s$. \square

Now we analyze graphs arising from finite geometries. A finite projective plane of order q has $q^2 + q + 1$ points and $q^2 + q + 1$ lines satisfying the following axioms.

1. Any two distinct points determine a line.
2. Any two distinct lines determine a point.
3. There exists four points such that no line is incident to three or four of them.

In particular, if q is a prime power there exists a projective plane that arises from a Galois field of order q , we denote the field by $GF(q) = \{g_0 = 0, g_1 = 1, g_2, \dots, g_{q-1}\}$. It is called the *algebraic projective plane*, and it is denoted by $PG(2, q)$. For further information about the algebraic projective planes, see [10]. Next, we take a projective plane $PG(2, q)$ and we will label their points and lines.

Let P be a point and let L be a line incident to P . We use P and L to label the elements of the projective plane. We call P the infinity point and L the infinity line. Since each line have $q + 1$ points, let $\{P_{g_0}, P_{g_1}, \dots, P_{g_{q-1}}\}$ be the set of points incident to L such that each point is different to P . Since each point is incident to $q + 1$ lines, let $\{L_{g_0}, L_{g_1}, \dots, L_{g_{q-1}}\}$ be the set of lines incident to P such that each line is different to L . Let $\{(g_i, g_0), (g_i, g_1), \dots, (g_i, g_{q-1})\}$ be the set of label of the points different from P incident to L_{g_i} . The remaining lines are labeled as follows. The line $[a, b]$ is adjacent to all the points (x, y) that satisfy $y = ax + b$ using the arithmetic of $GF(q)$, for all $a, b, x, y \in GF(q)$.

Now, we recall the definition of the incidence graph of a projective plane of order q , that is, a regular bipartite graph of degree $q + 1$ and partition (A, B) with $2(q^2 + q + 1)$ vertices: the set A have $q^2 + q + 1$ vertices corresponding to the points and the set B have $q^2 + q + 1$ vertices corresponding to the lines. Two vertices are adjacent if the corresponding pair point-line is incident. The such graph has a diameter 3, girth 6 and it is a $(q + 1, 6)$ -cage which has order equal to the well-known Moore bound. The incidence graph is also known as the Levi graph.

Theorem 10. *If G is the incidence graph of the algebraic projective plane of odd order q , then $q^2 + q + 1 \leq h(G) \leq q^2 + q + 2$.*

Proof. In any harmonious coloring of G , different points receive different colors, because every two points share a line, then $h(G) \geq q^2 + q + 1$.

We define the coloring in the points and lines as follows. We assign $q^2 + q + 1$ colors in the $q^2 + q + 1$ points of $PG(2, q)$. Now, if $a \neq 0$, the line $[a, b]$ is colored with the color of the point (a, b) . Since $a^2 \neq 0$, then $b \neq a^2 + b$ and (a, b) is not incident to $[a, b]$. Then the partial coloring is proper. Moreover, for $\alpha \neq 0$, if the point (α, β) belongs to the line $[a, b]$, $\beta = a\alpha + b$, then the point (a, b) is not incident to the line $[\alpha, \beta]$, otherwise, $b = a\alpha + \beta = a\alpha + a\alpha + b$, i.e., $0 = 1 + 1$, a contradiction because q is odd. Since for any pair of color classes $\{(a, b), [a, b]\}$ and $\{(m, n), [m, n]\}$ there is at most one incidence between them, the partial coloring is harmonious.

For $b \neq -1$, the line $[0, b]$ is colored with the color of the point $(0, b + 1)$, the line $[0, -1]$ is colored with the color of the point P , and the line L is colored with the color of the point $(0, 0)$. Clearly, the partial coloring is proper. If a point $(0, b + 1)$ is incident to the line $[a, b + 1]$ then the point $(a, b + 1)$ is not incident to the line $[0, b]$ since $b + 1 \neq b$. Therefore, the partial coloring is harmonious.

For $a \neq 0, -1$, the line L_a is colored with the color of the point P_{a+1} , the line L_{-1} is colored with the color of the point P_1 , and the line L_0 is colored with a new color. Clearly, the coloring is proper. Now, the point P_1 is incident to the line $[1, b]$ and $(1, b)$ is not incident to the line L_{-1} . If a point P_{a+1} is incident to the line $[a + 1, b]$, then the point $(a + 1, b)$ is not incident to the line L_a since $a + 1 \neq 0$. Finally, the point P is incident to the line L_a , the point P_{a+1} is not incident to the line $[0, -1]$, and the point P_1 is incident

to the line L , and the point $(0, 0)$ is not incident to the line L_{-1} . Therefore, the coloring is harmonious and $h(G) \leq q^2 + q + 2$. \square

Since a linear space \mathcal{S} is an incidence structure satisfying the following axioms:

1. Any two distinct points determine a line
2. Any line is incident to at least two distinct points
3. \mathcal{S} contains at least two distinct lines

using the same techniques as in Theorem 10, we can give bounds on the harmonious number in the incidence graph of the linear spaces.

Examples of linear spaces are the projective planes, the affine planes, and the 2 - (n, k) -designs. Clearly, the incidence graph of a linear space has diameter at most 4.

Now, we give a bound for the harmonious chromatic number of the incidence graph of a linear space related to the Erdős-Faber-Lovász Conjecture, see [1, 6, 7].

A *line coloring* of \mathcal{S} with k colors is an assignment of the lines of \mathcal{S} to a set of k colors. A line coloring of \mathcal{S} is called *proper* if any two intersecting lines have different colors. The *chromatic index* $\chi'(\mathcal{S})$ of \mathcal{S} is the smallest k such that there exists a proper line coloring of \mathcal{S} with k colors. Erdős, Faber and Lovász conjectured that the chromatic index of any finite linear space \mathcal{S} cannot exceed the number of its points, so if \mathcal{S} has n points then

$$\chi'(\mathcal{S}) \leq n.$$

Theorem 11. *Let \mathcal{S} be a linear space of n points, then its incidence graph G has the following bounds for $h(G)$,*

$$n \leq h(G) \leq n + \chi'(\mathcal{S}).$$

Moreover, if the Erdős-Faber-Lovász Conjecture is true for \mathcal{S} , then $h(G) \leq 2n$.

Proof. The incidence graph G of a linear space with point set \mathcal{P} has $h(G) \geq |\mathcal{P}| = n$ by axiom 1. To give the upper bound, we assign a coloring via a line coloring with $\chi'(\mathcal{S})$ colors and assign a different color to each vertex and the result follows because any two lines with the same color are at distance 3 or 4 in G . \square

Let n, b, k and r be positive integers with $n > 1$. Let $D = (P, B, I)$ be a triple consisting of a set P of n distinct objects, called points of D , a set B of b distinct objects, called blocks of D ($P \cap B = \emptyset$), and an incidence relation I , a subset of $P \times B$. We say that v is incident to u if exactly one of the ordered pairs (u, v) and (v, u) is in I ; then v is incident to u if and only if u is incident to v . D is called a 2 - (n, b, k, r) *block design* (for short, 2 - (n, b, k, r) *design*) if it satisfies the following axioms.

1. Each block of D is incident to exactly k distinct points of D .
2. Each point of D is incident to exactly r distinct blocks of D .
3. If u and v are distinct points of D , then there are exactly one block of D incident to both u and v .

A 2 - (n, b, k, r) design is called a balanced incomplete block design BIBD; it is called an (n, k) -design, too, since the parameters of a 2 - (n, b, k, r, λ) design are not all independent. The two equations connecting them are $nr = bk$ and $r(k - 1) = (n - 1)$. For a detailed introduction to block designs we refer to [2].

A design is *resolvable* if its blocks can be partitioned into r sets so that b/r blocks of each part are point-disjoint and each part is called a *parallel class*. Therefore, Theorem 11 implies that the incidence graph G of a resolvable design D satisfies $h(G) \leq n + \frac{n-1}{k-1}$.

When $k = 2$, the (n, k) -design is isomorphic to the complete graph K_n and it is resolvable when n is even. The following theorem improves the previous bounds.

Theorem 12. *Let G be the incidence graph of the complete graph K_n . Then*

$$\frac{3}{2}n \leq h(G) \leq \frac{5}{3}n + \theta(1).$$

Proof. For the lower bound, let G be the incidence graph of the complete graph K_n , with $V(K_n) = \{x_1, \dots, x_n\}$ and let $\varsigma : E(K_n) \cup V(K_n) \rightarrow \{c_1, \dots, c_{h(G)}\}$ be a harmonious coloring.

By Theorem 11, $n \leq h(G)$. For each $x_i \in V(K_n)$, let $c_i = \varsigma(x_i)$. For each $i \in \{1, \dots, h(G)\}$, let $e_i = \{e \in E(K_n) : \varsigma(e) = c_i\}$. Observe that each e_i induces a matching in K_n , $\sum_{i=1}^{h(G)} |e_i| = \binom{n}{2}$ and since ς is a harmonious coloring, for every $x_i \in V(K_n)$, x_i is not incident to an edge of e_i .

Let G^* be the spanning subgraph of K_n such that for each $xy \in E(G^*)$, $\varsigma(xy) \in \{c_1, \dots, c_n\}$. On the one hand, $e(G^*) = \sum_{i=1}^n |e_i| = \frac{1}{2} \sum_{i=1}^n \deg_{G^*}(x_i)$. On the other hand, for each $x_i \in V(G^*)$, if there is $x_i x_j \in E(G^*)$ such that $\varsigma(x_i x_j) = c_k$, then x_k is not adjacent to an edge of e_i . Thus, $|e_i| \leq \frac{n-1-\deg_{G^*}(x_i)}{2}$.

Therefore $\frac{1}{2} \sum_{i=1}^n \deg_{G^*}(x_i) = \sum_{i=1}^n |e_i| \leq \sum_{i=1}^n \frac{n-1-\deg_{G^*}(x_i)}{2}$ which implies that $\sum_{i=1}^n \deg_{G^*}(x_i) \leq \binom{n}{2}$. Thus, $e(G^*) \leq \frac{\binom{n}{2}}{2}$ and therefore $\sum_{j=n+1}^{h(G)} |e_j| \geq \frac{\binom{n}{2}}{2}$. Since, for each $i \in \{1, \dots, h(G)\}$, $|e_i| \leq \lfloor \frac{n}{2} \rfloor$ it follows that $h(G) - n \geq \frac{\binom{n}{2}}{2 \lfloor \frac{n}{2} \rfloor} \geq \lceil \frac{n}{2} \rceil$ and from here the result follows.

For the upper bound, let G be the incidence graph of the complete graph K_n and suppose $n = 6m - 3$ for some integer m . Let $K_{n+3} \cong K_{6m}$ be the complete graph with $V(K_{6m}) = \{x_1, \dots, x_{6m}\}$; and let $V_0 = \{x_1, \dots, x_{2m}\}$, $V_1 = \{x_{2m+1}, \dots, x_{4m}\}$ and $V_2 = \{x_{4m+1}, \dots, x_{6m}\}$. Let $\varsigma : E(K_{6m}) \cup V(K_{6m}) \rightarrow \{c_1, \dots, c_{10m}\}$ be the coloring defined as follows:

1. For each $x_i \in V(K_{6m})$, let $\varsigma(x_i) = i$.

For each $k \in \{0, 1, 2\}$, let $\{M_1^k, \dots, M_{2m-1}^k\}$ be a factorization of perfect matchings of the induced subgraph $K_{6m}[V_k]$.

2. For each $e \in M_t^j$, let $\varsigma(e) = 2mk^* + t$, with $k^* \equiv j + 1 \pmod{3}$.

Note that the colors $2m, 4m$ and $6m$ just appear in the vertices x_{2m}, x_{4m} and x_{6m} , respectively. Just remains to color the edges of a graph which is isomorphic to the complete tripartite graph $K_{2m, 2m, 2m}$. Let $\{M_1, \dots, M_{\chi'}\}$ be a decomposition of matchings of $K_{2m, 2m, 2m}$ where χ' is the chromatic index of $K_{2m, 2m, 2m}$, then $4m \leq \chi' \leq 4m + 1$ by Vizing's Theorem.

3. For each $e \in M_j$ let $\varsigma(e) = 6m + j$.

It is not hard to see that ς is a harmonious $(6m + 4m)$ -coloring of K_{6m} , and, by deleting the vertices x_{2m}, x_{4m} and x_{6m} we obtain a harmonious coloring with $10m - 3$ colors of $K_{6m-3} \cong K_n$, or $10m - 2$ colors if $K_{2m, 2m, 2m}$ is a graph of Class 2. Thus $h(G) \leq 10m - 2 = n + 4 \lceil \frac{n}{6} \rceil + 1$. For the remaining cases, by adding some suitable number l of vertices (obtaining a complete graph K of order multiple of 6 and a harmonious coloring of K) and by deleting those l vertices (and considering that we lose $\min\{k, 3\}$ colors) then

$$n + \lceil \frac{n}{2} \rceil \leq h(G) \leq n + 4 \lceil \frac{n}{6} \rceil + \theta(1)$$

and the result follows. \square

Next, we recall that a set $U \subset V(G)$ is a *packing set* of G if every pair of distinct vertices $u, v \in U$ satisfy $d(u, v) \geq 3$. The *packing number* $\rho(G)$ of G is the maximum cardinality of a packing set of G .

Theorem 13. *If G is a graph of order n , then*

$$\frac{n}{\rho(G)} \leq h(G) \leq n - \rho(G) + 1.$$

Proof. Suppose that $h(G) = k$ and let there be given a harmonious coloring of G with k colors and color classes V_1, V_2, \dots, V_k . Since $n = \sum_{i=1}^k |V_i| \leq k\rho(G)$, the lower bound follows.

Now, let U be a maximum packing set of vertices of G and assign the color 1 to each vertex of U . Assigning distinct colors different from 1 to each vertex of $V(G) - U$ produces a harmonious coloring of G . Hence $h(G) \leq |V(G) - U| + 1 = n - \rho(G) + 1$. \square

Observe that a harmonious homomorphism f of a graph G such that f identify a set of vertices in a maximum packing set produces a graph $f(G)$ of diameter at most 4.

4 On circulant graphs

Given a set $J \subseteq \{1, \dots, \lfloor \frac{n}{2} \rfloor\}$, the *circulant graph* $C_n(J)$ is defined as the graph with vertex set equal to \mathbb{Z}_n and $E(C_n(J)) = \{ij : |j - i| = l, l \in J\}$, we call J the set of *jumps* of $C_n(J)$. Complete graphs and cycles are examples of circulant graphs. Observe that the number of edges in $C_n(J)$ is $n|J|$ (except when $\lfloor \frac{n}{2} \rfloor \in J$ for n even).

In [4], Dębsty et al. proved that for a fixed set J , $h(C_n(J)) \geq \sqrt{2n|J|} + O(n^{0.25}(\log n)^{0.5})$. In other words, if the circulant graph has small regularity, then its harmonious chromatic number is close to the lower bound of Equation 1.

In Theorem 14 we present a family of circulant graphs with two jumps and a harmonious chromatic number equal to the lower bound of Equation 1. In Theorem 16, we give a family of circulant graphs of order n with regularity $\Omega(\sqrt{n})$ and a harmonious chromatic number equal to n , where n follows the asymptotic distribution of the prime power numbers.

Given a graph G , the graph G^{\boxtimes} is defined as follows, see [16]. If G has vertex set $V = \{w_1, w_2, \dots, w_n\}$, the graph G^{\boxtimes} has as vertex set two copies of V , namely, $\{u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n\}$ and two vertices x_i and y_j are adjacent if $w_i w_j$ is an edge of G , for the symbols $x, y \in \{u, v\}$. For instance, if $w_1 w_2$ is an edge of a graph G , the graph G^{\boxtimes} has the edges $u_1 u_2, v_1 v_2, u_1 v_2$ and $v_1 u_2$, see Figure 1.




Figure 1: An edge of G produces four edges in G^{\boxtimes} .

Theorem 14. *The circulant graph $C_{t(t-1)}(1, \frac{t(t-1)}{2} - 1)$ for $t \equiv -1 \pmod{4}$ has harmonious chromatic number equals the lower bound of Equation 1.*

Proof. The graph K_t^{\boxtimes} is isomorphic to a complete graph minus a perfect matching $K_{2t} - tK_2$. It has $\frac{(2t)(2t-1)}{2} - t$ edges, i.e., $4\frac{t(t-1)}{2}$ edges. If we take t odd, the complete graph K_t is an Eulerian graph and we can unfold its Eulerian tour $C_{\frac{t(t-1)}{2}}$ as $C_{\frac{t(t-1)}{2}}^{\boxtimes}$, see Figure 2 (left).

The edges between the two cycles $C_{\frac{t(t-1)}{2}}$ form a cycle C if $\frac{t(t-1)}{2}$ is odd, i.e., if $t \equiv -1 \pmod{4}$. Such cycle C has length $t(t-1)$. Figure 2 (right) shows the cycle C with the labels $w_0, w_1, \dots, w_{t(t-1)-1}$. Note that one cycle $C_{\frac{t(t-1)}{2}}$ (in the interior region) has labels w_{even} and the other cycle $C_{\frac{t(t-1)}{2}}$ (in the exterior region) has labels w_{odd} due to the fact that $C_{\frac{t(t-1)}{2}}^{\boxtimes}$ is isomorphic to the circulant graph $C_{t(t-1)}(1, \frac{t(t-1)}{2} - 1)$.

This circulant graph $C_{t(t-1)}(1, \frac{t(t-1)}{2} - 1)$ is colored with the $2t$ colors arising from the graph $K_{2t} - tK_2$ having an edge between every two color classes except for the color classes coming from the perfect matching tK_2 .




Figure 2: The graph $C_{\frac{t(t-1)}{2}}^{\triangleright}$.

Clearly, $h(C_{t(t-1)}(1, \frac{t(t-1)}{2} - 1)) \leq 2t$ because each class is independent and every two color classes have an edge between them except for t pairs. On the other hand,

$$\frac{1}{2} + \sqrt{\frac{1}{4} + 2(2t(t-1))} \leq h(C_{t(t-1)}(1, \frac{t(t-1)}{2} - 1)),$$

since $2t - 1 < \frac{1}{2} + \sqrt{4t^2 - 4t + \frac{1}{4}}$ because $t > 1$, the result follows. \square

Theorem 15. *If a graph G has size m and its harmonious number h equals the lower bound $\frac{1}{2} + \sqrt{\frac{1}{4} + 2m}$, then G^{\triangleright} has harmonious number $2h$.*

Proof. The graph G has size m equals to a triangular number and each two color classes has exactly one edge between them. Therefore, G^{\triangleright} has size $M = 4m$ and we can give a harmonious $2h$ -coloring in a natural way. Since $2h - 1 < \frac{1}{2} + \sqrt{\frac{1}{4} + 2M} = \frac{1}{2} + \sqrt{\frac{1}{4} + 2(4m)}$, the result follows. \square

A natural question is the following: Which are the circulant graphs of diameter two? The obvious answer is that they are those such that any number from 1 to n can be represented as the sum of two jumps, so we suppose that the question is rather when that can be done.

Graphs for which $\deg(u) + \deg(v) \geq n - 1$ for every two nonadjacent vertices u, v have diameter at least 2, thus circulant graphs which are r -regular of order n with $r \geq (n - 1)/2$ have harmonious number n . We give a construction of circulant graphs with harmonious number n , such that each vertex has degree $\theta(\sqrt{n})$.

A subset $D = \{d_1, \dots, d_k\}$ of an additive group Γ , of order n , is a (n, k) -difference set if for each $g \in \Gamma$, there is a unique pair of different elements $d_i, d_j \in D$ such that $g = d_i - d_j$. Therefore, if $D \subset \Gamma$ is a (n, k) -difference set and $g \in \Gamma$ is an arbitrary element, then $D + g := \{d_1 + g, d_2 + g, \dots, d_k + g\}$ is a (n, k) -difference set.

The parameters n and k of a (n, k) -difference set satisfy the necessary conditions $n - 1 = k(k - 1)$. It is conjectured that $q = k - 1$ must be a prime power, its hold for $q \leq 2,000,000$. Then $n = q^2 + q + 1$ and the construction of $(q^2 + q + 1, q + 1)$ -difference sets have an algebraic approach via the Galois fields of order q . See Table 1 for some examples of difference sets. See [12] for further reading.

\mathbb{Z}_n	Difference set	\mathbb{Z}_n	Difference set
\mathbb{Z}_7	1,2,4	\mathbb{Z}_{57}	1,2,4,14,33,37,44,53
\mathbb{Z}_{13}	1,2,5,7	\mathbb{Z}_{73}	1,2,4,8,16,32,37,55,64
\mathbb{Z}_{21}	1,4,5,10,12	\mathbb{Z}_{91}	1,2,4,10,28,50,57,62,78,82
\mathbb{Z}_{31}	1,2,4,9,13,19	\mathbb{Z}_{133}	1,2,4,13,21,35,39,82,89,95,105,110

Table 1: Some difference sets for \mathbb{Z}_n with small n .

Figure 3 shows an example of a circulant graph with diameter 2, where a difference set is used to define the set of jumps. Note that, by definition of circulant graphs and $12 \equiv 19 \pmod{31}$, we obtain $C_{31}(1, 2, 4, 9, 12, 13)$ for the different set $\{1, 2, 4, 9, 13, 19\}$.




Figure 3: The graph $C_{31}(1, 2, 4, 9, 12, 13)$, where only the jumps $0i$ for $i \in \{1, 2, 4, 9, 13, 19\}$ are shown.

Theorem 16. *Let q be a prime power, $n = q^2 + q + 1$, and let D be a difference set of \mathbb{Z}_n such that $0 \notin D$. Let $C_n(J)$ be a circulant graph with $J = \{i: i \in D \text{ if } i \leq n/2\} \cup \{n-i: i \in D \text{ if } i > n/2\}$, then $h(C_n(J)) = n$.*

Proof. The graph $C_n(J)$ is r -regular for $r = 2(q + 1) = 2\lceil\sqrt{n}\rceil = \theta(\sqrt{n})$ and its diameter is 2. To see this, take two non adjacent vertices u and v , then $l = u - v \notin J$, but there exist $i, j \in D$ such that $j - i = l$. Then u is adjacent to $w = u + j$ and w is adjacent to $w - i = v$. Therefore, the result follows. \square

5 Conclusions

Harmonious coloring was first proposed in 1982 [8], but still very little is known about it. For future work, we propose to study either the harmonious achromatic number defined in Section 2 or the greedy coloring via the harmonious homomorphisms. We define the *harmonious Grundy number* as the number of colors in the worst case using such greedy colorings.

Theorem 10 shows the harmonious chromatic number of the $(q + 1, 6)$ -cages, hence another possible problem is to estimate the harmonious chromatic number of the $(q + 1, 8)$ -cages, i.e., graphs that are the incidence graph of generalized quadrangles.

The smallest non-trivial generalized quadrangle is $GQ(2, 2)$, whose representation can be obtained through a 1-factorization of K_6 as follows, see [17]. If $V(K_6) = \{1, \dots, 6\}$, the points of $GQ(2, 2)$ are the edges of K_6




Figure 4: (Left) The harmonious chromatic number of the incidence graph of $GQ(2, 2)$ is 10. (Center) A color class with two points and a line produces other color classes via rotations of $\frac{2}{5}\pi$. (Right) A color class with two lines and a point produces other color classes via rotations of $\frac{2}{5}\pi$.

and each line has three points arising from a matching of K_6 . In Figure 4 (Left), $GQ(2, 2)$ is shown with a harmonious coloring of 10 colors where each color class has three elements: There are two types of color classes, the first one contains two points and a line (as in Figure 4 (Center)), while the second one contains two lines and a point (as in Figure 4 (Right)). Both types can be obtained if $GQ(2, 2)$ is drawn appropriately (as in Figure 4 (Left)) so that rotating $GQ(2, 2)$ through a $\frac{2}{5}\pi$ angle 5 times produces those color classes. Therefore, since the $(3, 8)$ -cage graph G is the incidence graph of $GQ(2, 2)$ and it has 45 edges, it requires at least 10 colors in any harmonious coloring, therefore $h(G) = 10$. As consequence, the achromatic number of G is also 10.

6 Statements and Declarations

This work was partially supported by PAPIIT-México under Project IN108121; CONACyT-México under projects 282280, A1-S-12891, 47510664; and PAIDI-México under Project PAIDI/007/21.

The authors have no relevant financial or non-financial interests to disclose.

All authors contributed to the study conception and design. The first draft of the manuscript was written by Gabriela Araujo-Pardo, Juan José Montellano-Ballesteros, Christian Rubio-Montiel, and Mika Olsen, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

References

- [1] G. Araujo-Pardo, Gy. Kiss, C. Rubio-Montiel, and A. Vázquez-Ávila. On line colorings of finite projective spaces. *Graphs Combin.*, 37(3):891–905, 2021.
- [2] T. Beth, D. Jungnickel, and H. Lenz. *Design theory. Vol. I*, volume 69 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 1999.
- [3] G. Chartrand and P. Zhang. *Chromatic graph theory*. Discrete Mathematics and its Applications. CRC Press, 2009.
- [4] M. Dębski, Z. Lonc, and P. Rzażewski. Achromatic and harmonious colorings of circulant graphs. *J. Graph Theory*, 87(1):18–34, 2018.

- [5] K. Edwards. The harmonious chromatic number and the achromatic number. In *Surveys in combinatorics, 1997 (London)*, volume 241 of *London Math. Soc. Lecture Note Ser.*, pages 13–47. Cambridge Univ. Press, Cambridge, 1997.
- [6] P. Erdős. Problems and results in graph theory and combinatorial analysis. In *Proceedings of the Fifth British Combinatorial Conference (Univ. Aberdeen, Aberdeen, 1975)*, pages 169–192. Congressus Numerantium, No. XV, 1976.
- [7] P. Erdős. On the combinatorial problems which I would most like to see solved. *Combinatorica*, 1(1):25–42, 1981.
- [8] O. Frank, F. Harary, and M. Plantholt. The line-distinguishing chromatic number of a graph. *Ars Combin.*, 14:241–252, 1982.
- [9] F. Harary, S. Hedetniemi, and G. Prins. An interpolation theorem for graphical homomorphisms. *Portugal. Math.*, 26:453–462, 1967.
- [10] J. W. P. Hirschfeld. *Projective geometries over finite fields*. The Clarendon Press Oxford University Press, New York, 1979. Oxford Mathematical Monographs.
- [11] J. E. Hopcroft and M. S. Krishnamoorthy. On the harmonious coloring of graphs. *SIAM J. Algebraic Discrete Methods*, 4(3):306–311, 1983.
- [12] D. Jungnickel, A. Pott, and K.W. Smith. Difference sets. In *Handbook of Combinatorial Designs*, pages 419–435. Chapman & Hall; CRC, 2007.
- [13] A. Kündrík. The harmonious chromatic number of a graph. In *Fourth Czechoslovakian Symposium on Combinatorics, Graphs and Complexity (Prachatice, 1990)*, volume 51 of *Ann. Discrete Math.*, pages 167–170. North-Holland, Amsterdam, 1992.
- [14] J. Mitchem. On the harmonious chromatic number of a graph. *Discrete Math.*, 74(1-2):151–157, 1989.
- [15] M. Olsen, C. Rubio-Montiel, and A. Silva-Ramírez. Zykov sums of digraphs with diachromatic number equal to their harmonious number. *Submitted*.
- [16] C. Rubio-Montiel. The 4-girth-thickness of the complete multipartite graph. *Electron. J. Graph Theory Appl. (EJGTA)*, 7(1):183–188, 2019.
- [17] E.W. Weisstein. Generalized Quadrangle. From *MathWorld*—A Wolfram Web Resource. <https://mathworld.wolfram.com/GeneralizedQuadrangle.html>. Accessed: 2022-03-04.