

Two-dimensional contiguous relations for linearization coefficients of orthogonal polynomials in the Askey-scheme

Howard S. Cohl^{*1} and Lisa Ritter^{*}

^{*} *Applied and Computational Mathematics Division, National Institute of Standards and Technology, Gaithersburg, MD 20899-8910, USA*

E-mail: howard.cohl@nist.gov, lisa.ritter@nist.gov

URL: <http://www.nist.gov/itl/math/msg/howard-s-cohl.cfm>

Received ???, in final form ???; Published online ???

doi:10.3842/JOURNAL.202*.***

Abstract. We produce two-dimensional contiguous relations for generalized hypergeometric functions by starting with linearization coefficients for some continuous generalized hypergeometric orthogonal polynomials in the Askey-scheme.

1 Introduction

In this paper we produce two-dimensional contiguous relations for linearization coefficients of hypergeometric orthogonal polynomials in the Askey-scheme. These contiguous relations are derived by using integrals of products of these orthogonal polynomials. These integrals are related to linearization coefficients for the polynomials. The idea for the two-dimensional contiguous relations goes back to a paper by Ismail, Kasraoui & Zeng (2013) [4]. In this paper we re-derive the general expression for the two-dimensional contiguous relations and then apply this relation to several specific examples, namely for the linearization of a product of two and three Gegenbauer polynomials, a product of two and three Hermite polynomials, linearization of two Jacobi polynomials and for a product of two unscaled and two scaled Laguerre polynomials.

2 Preliminaries

We adopt the following list conventions as follows. Within a list of items, we define

$$a + \left\{ \begin{matrix} x_1 \\ \vdots \\ x_n \end{matrix} \right\} := \{a + x_1, \dots, a + x_n\},$$

and when \pm is used within a list of values, we define $\pm a := \{a, -a\}$. Let $z \in \mathbb{C}$, $n, k \in \mathbb{N}_0$ unless otherwise stated. The definition that we use for the Pochhammer symbol is given by

$$(z)_n := (z)(z+1)\cdots(z+n-1), \quad (z)_0 := 1, \quad z \in \mathbb{C}, \quad (1)$$

$$(z_1, \dots, z_k)_n := (z_1)_n \cdots (z_k)_n. \quad (2)$$

We will also adopt the following compact notation for the minimum and maximum of any two integers, $m, n \in \mathbb{Z}$,

$$m \vee n := \max(m, n), \quad m \wedge n := \min(m, n).$$

¹Working remotely in Mission Viejo, California

Define the generalized hypergeometric series [2, Chapter 16]

$${}_rF_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; x \right) = \sum_{n=0}^{\infty} \frac{(a_1, \dots, a_r)_n}{(b_1, \dots, b_s)_n} \frac{x^n}{n!}, \quad (3)$$

and the Kampé de Fériet double hypergeometric series [8, (28)]

$$\begin{aligned} F_{l:m;n}^{p:q;k} \left(\begin{matrix} a_1, \dots, a_p : b_1, \dots, b_q; c_1, \dots, c_k \\ \alpha_1, \dots, \alpha_l : \beta_1, \dots, \beta_m; \gamma_1, \dots, \gamma_n \end{matrix}; x, y \right) \\ = \sum_{r,s=0}^{\infty} \frac{(a_1, \dots, a_p)_{r+s} (b_1, \dots, b_q)_r (c_1, \dots, c_k)_s}{(\alpha_1, \dots, \alpha_l)_{r+s} (\beta_1, \dots, \beta_m)_r (\gamma_1, \dots, \gamma_n)_s} \frac{x^r y^s}{r! s!}. \end{aligned} \quad (4)$$

3 Two-dimensional contiguous relations which correspond to the linearization of a product of orthogonal polynomials

In this section we derive the general relation for two-dimensional contiguous relations for linearization coefficients of unscaled orthogonal polynomials, namely Theorem 3.1 below. This theorem is similar to [4, Theorem 2.1], albeit for unscaled polynomials $p_n(x)$. In the remainder of this paper, we compute explicitly the linearization coefficients of the products of two and three orthogonal polynomials. From this we show examples of these contiguous relations for certain continuous hypergeometric orthogonal polynomials in the Askey scheme. However, in 7.6 we extend this theorem for scaled Laguerre polynomials. The general result presented in [4, Theorem 2.1] is for integrals of products of scaled orthogonal polynomials $p_n(\lambda x)$, for some $\lambda \in \mathbb{C}$.

Consider a sequence of continuous orthogonal polynomials $p_n(x)$, $m, n \in \mathbb{N}_0$, $x \in \mathbb{C}$, which satisfy the orthogonality relation

$$\int_{\mathcal{I}} p_m(x; \boldsymbol{\alpha}) p_n(x; \boldsymbol{\alpha}) d\mu = h_n(\boldsymbol{\alpha}) \delta_{m,n}, \quad (5)$$

where \mathcal{I} is the support of the measure μ and $\boldsymbol{\alpha}$ is a set of parameters. Orthogonal polynomials satisfy the following three-term recurrence relation, with the assumption $p_{-1}(x; \boldsymbol{\alpha}) := 0$,

$$p_{n+1}(x; \boldsymbol{\alpha}) = (A_n x + B_n) p_n(x; \boldsymbol{\alpha}) - C_n p_{n-1}(x; \boldsymbol{\alpha}). \quad (6)$$

The linearization coefficients $\mathbf{a}_{k,\mathbf{n}} := \mathbf{a}_{\mathbf{m}}$, where $\mathbf{n} := \{n_1, \dots, n_N\} \in \mathbb{N}_0^N$, $N \geq 2$, $k \in \mathbb{N}_0$, $\mathbf{m} := \{k\} \cup \mathbf{n}$, are defined using

$$p_{n_1}(x; \boldsymbol{\alpha}) \cdots p_{n_N}(x; \boldsymbol{\alpha}) = \sum_{k=0}^{n_1 + \cdots + n_N} \mathbf{a}_{\mathbf{m}} p_k(x; \boldsymbol{\alpha}). \quad (7)$$

Note that even though it is true that $k \in \{0, \dots, n_1 + \cdots + n_N\}$, it may be that k has a more restricted range depending on the specific orthogonal polynomials involved.

Now consider the integral over $N + 1$ orthogonal polynomials $P(\mathbf{m}; \boldsymbol{\alpha}) \in \mathbb{R}$, where $\mathbf{m} := \{n_1, \dots, n_{N+1}\}$ by

$$P(\mathbf{m}; \boldsymbol{\alpha}) := \int_{\mathcal{I}} p_{n_1}(x; \boldsymbol{\alpha}) \cdots p_{n_N}(x; \boldsymbol{\alpha}) p_{n_{N+1}}(x; \boldsymbol{\alpha}) d\mu. \quad (8)$$

We can formally see that integral is associated with the linearization of a product of N orthogonal polynomials as follows. Without loss of generality choose

$$n_1 \leq n_2 \leq \cdots \leq n_N \leq n_{N+1}. \quad (9)$$

Then after substituting (7) in (8) and using (5), one obtains

$$P(\mathbf{m}; \boldsymbol{\alpha}) = \int_{\mathcal{I}} \sum_{k=0}^{n_1 + \dots + n_N} \mathbf{a}_{\mathbf{m}} p_{n_{N+1}}(x; \boldsymbol{\alpha}) p_k(x; \boldsymbol{\alpha}) d\mu = h_{N+1}(\boldsymbol{\alpha}) \mathbf{a}_{\mathbf{m}}, \quad (10)$$

and since overlap in the orthogonality only occurs if $n_{N+1} \leq n_1 + \dots + n_N$, the integral $P(\mathbf{m}; \boldsymbol{\alpha})$ will vanish otherwise.

Now define

$$P_j^{\pm} := P_j^{\pm}(\mathbf{m}; \boldsymbol{\alpha}) := P(n_1, \dots, n_{j-1}, n_j \pm 1, n_{j+1}, \dots, n_{N+1}; \boldsymbol{\alpha}).$$

Theorem 3.1. *Let $N, j, k \in \mathbb{N}_0$, $1 \leq j < k \leq N + 1$, $n_j, n_k \in \mathbf{m}$, $x \in \mathcal{I}$, $A_{n_j}, B_{n_j}, C_{n_j}$ satisfy the three-term recurrence relation (6). Then the definite integral $P(\mathbf{m}; \boldsymbol{\alpha})$ corresponding to a product of $N + 1$ orthogonal polynomials defined by (8), satisfies the following sequence of $\binom{N+1}{2}$ contiguous relations*

$$(B_{n_j} A_{n_k} - A_{n_j} B_{n_k})P - A_{n_k} P_j^+ - C_{n_j} A_{n_k} P_j^- + A_{n_j} P_k^+ + C_{n_k} A_{n_j} P_k^- = 0.$$

Proof. Choose a quantum number $m := n_j \in \mathbf{m}$, and then evaluate P_j^+ using (8). Then use (6) to replace $p_{m+1}(x)$ with $(A_m x + B_m)p_m(x) - C_m p_{m-1}(x)$ in the integrand. Now choose a different quantum number $n := n_k \in \mathbf{m}$ and replace $x p_n(x)$ in the resulting expression with

$$x p_n(x) = \frac{1}{A_n} (p_{n+1} - B_n p_n(x) + C_n p_{n-1}). \quad (11)$$

Multiplying both sides of the resulting expression by A_n produces a five term two-dimensional contiguous relation for P involving P_j^{\pm} and P_k^{\pm} . Repeating this process for all $\binom{N+1}{2}$ unique combinations of quantum numbers in \mathbf{m} completes the proof. \blacksquare

Remark 3.2. *For a linearization of a product of N orthogonal polynomials, choose $j \neq k \in \{1, \dots, N\}$. The three-term recurrence relation (11) is symmetric under permutation of n_j and n_k . Hence each unique contiguous relation is the result of choosing two quantum numbers from $N + 1$ possibilities and therefore one will obtain a sequence of $\binom{n+1}{2}$ contiguous relations for the definite integral of a product of $N + 1$ orthogonal polynomials.*

In the remainder of the paper, we compute examples of these contiguous relations for certain continuous hypergeometric orthogonal polynomials in the Askey scheme.

4 Gegenbauer polynomials

The Gegenbauer (or ultraspherical) polynomials can be defined as [6, (9.8.19)]

$$C_n^{\lambda}(x) = \frac{(2\lambda)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n + 2\lambda \\ \lambda + \frac{1}{2} \end{matrix}; \frac{1-x}{2} \right), \quad (12)$$

where $n \in \mathbb{N}_0$, $\lambda \in \mathbb{C} \setminus \{0\}$, $x \in \mathbb{C}$. Gegenbauer polynomials are orthogonal on $x \in (-1, 1)$, with orthogonality relation [6, (9.8.20)]

$$\int_{-1}^1 C_m^{\lambda}(x) C_n^{\lambda}(x) (1-x^2)^{\lambda-\frac{1}{2}} dx = \frac{\pi \Gamma(2\lambda + n) \delta_{m,n}}{2^{2\lambda-1} (n+\lambda) n! \Gamma(\lambda)^2} =: h_n^{\lambda} \delta_{m,n}, \quad (13)$$

where $\lambda \in (-\frac{1}{2}, \infty) \setminus \{0\}$. The three-term recurrence relation for Gegenbauer polynomials is described via (6) with [2, Table 18.9.1]

$$A_n = \frac{2(n+\lambda)}{n+1}, \quad B_n = 0, \quad C_n = \frac{n+2\lambda-1}{n+1}. \quad (14)$$

4.1 Linearization of a product of two Gegenbauer polynomials

Gegenbauer polynomials have a linearization formula for a product of two Gegenbauer polynomials given by [2, (18.18.22)]

$$C_m^\lambda(x)C_n^\lambda(x) = \sum_{k=0}^m B_{k,m,n}^\lambda C_{m+n-2k}^\lambda(x), \quad (15)$$

where $m, n \in \mathbb{N}_0$, without loss of generality $n \geq m$, and

$$B_{k,m,n}^\lambda := \frac{(m+n+\lambda-2k)(m+n-2k)!(\lambda)_k(\lambda)_{m-k}(\lambda)_{n-k}(2\lambda)_{m+n-k}}{(m+n+\lambda-k)k!(m-k)!(n-k)!(\lambda)_{m+n-k}(2\lambda)_{m+n-2k}}. \quad (16)$$

Using (13), we see that (15) is equivalent to the following integral of a product of three Gegenbauer polynomials

$$C(k, m, n; \lambda) := \int_{-1}^1 C_m^\lambda(x)C_n^\lambda(x)C_{m+n-2k}^\lambda(x)(1-x^2)^{\lambda-\frac{1}{2}} dx = h_{m+n-2k}^\lambda B_{k,m,n}^\lambda. \quad (17)$$

Remark 4.1. Using the recurrence relations given by (14) with Theorem 3.1 one obtains the following contiguous relations:

$$(m+\lambda)(p+1)C_1^+ + (p+2\lambda-1)(m+\lambda)C_1^- = (p+\lambda)(m+1)C_2^+ + (p+\lambda)(m+2\lambda-1)C_2^-, \quad (18)$$

$$(n+\lambda)(m+1)C_2^+ + (m+2\lambda-1)(n+\lambda)C_2^- = (m+\lambda)(n+1)C_3^+ + (m+\lambda)(n+2\lambda-1)C_3^-, \quad (19)$$

$$(n+\lambda)(p+1)C_1^+ + (p+2\lambda-1)(n+\lambda)C_1^- = (p+\lambda)(n+1)C_3^+ + (p+\lambda)(n+2\lambda-1)C_3^-. \quad (20)$$

To use these with (17), we can alter it by taking $p = m + n - 2k$, to get the following:

$$C(p, m, n; \lambda) := h_p^\lambda B_{\frac{1}{2}(m+n-p),m,n}^\lambda. \quad (21)$$

For these contiguous relations (18)-(20) to be satisfied, there are conditions on the parameters. Namely, $p \in \{n-m, \dots, n+m\}$ and $m+n-p \pm 1$ must be even. These can be combined with (21), where the numbers 1, 2 and 3 correspond to the parameters p , m , and n respectively. This produces the following:

$$\begin{aligned} & \frac{(\lambda+m)(2\lambda+p)}{(\lambda+p+1)} B_{\frac{1}{2}(m+n-p-1),m,n}^\lambda + \frac{p(\lambda+m)}{(\lambda+p-1)} B_{\frac{1}{2}(m+n-p+1),m,n}^\lambda \\ &= (m+1)B_{\frac{1}{2}(m+n-p+1),m+1,n}^\lambda + (m+2\lambda-1)B_{\frac{1}{2}(m+n-p-1),m-1,n}^\lambda, \end{aligned} \quad (22)$$

$$\begin{aligned} & (n+\lambda)(m+1)B_{\frac{1}{2}(m+n-p+1),m+1,n}^\lambda + (m+2\lambda-1)(n+\lambda)B_{\frac{1}{2}(m+n-p-1),m-1,n}^\lambda \\ &= (m+\lambda)(n+1)B_{\frac{1}{2}(m+n-p+1),m,n+1}^\lambda + (m+\lambda)(n+2\lambda-1)B_{\frac{1}{2}(m+n-p-1),m,n-1}^\lambda, \end{aligned} \quad (23)$$

$$\begin{aligned} & \frac{(\lambda+n)(2\lambda+p)}{(\lambda+p+1)} B_{\frac{1}{2}(m+n-p-1),m,n}^\lambda + \frac{p(\lambda+n)}{(\lambda+p-1)} B_{\frac{1}{2}(m+n-p+1),m,n}^\lambda \\ &= (n+1)B_{\frac{1}{2}(m+n-p+1),m,n+1}^\lambda + (n+2\lambda-1)B_{\frac{1}{2}(m+n-p-1),m,n-1}^\lambda. \end{aligned} \quad (24)$$

Take for instance for (22) using the identities of (16) and (13). Then it reduces to

$$\begin{aligned}
& (m+\lambda)(p+1)C_1^+ + (p+2\lambda-1)(m+\lambda)C_1^- - (p+\lambda)(m+1)C_2^+ - (p+\lambda)(m+2\lambda-1)C_2^- \\
&= \frac{2p!(\lambda)_{\frac{1}{2}(n+m-p-1)}(\lambda)_{\frac{1}{2}(m+p-n-1)}(\lambda)_{\frac{1}{2}(n+p-m-1)}(2\lambda)_{\frac{1}{2}(n+m+p-1)}}{\Gamma(\frac{1}{2}(n+m-p+1))\Gamma(\frac{1}{2}(m-n+p+1))\Gamma(\frac{1}{2}(n-m+p+1))(\lambda)_{\frac{1}{2}(n+m+p-1)}(2\lambda)_{p-1}} \\
&\quad \times \left(\frac{(p+1)(\lambda+m)(2\lambda+m+p-n-1)(2\lambda+n+p-m-1)(4\lambda+n+m+p-1)}{(m+p-n+1)(n+p-m+1)(2\lambda+p-1)(2\lambda+m+n+p-1)(2\lambda+m+n+p+1)} \right. \\
&\quad - \frac{(m+1)(p+\lambda)(2\lambda+m+n-p-1)(2\lambda+m+p-n-1)(4\lambda+m+n+p-1)}{(m+n-p+1)(m+p-n+1)(2\lambda+p-1)(2\lambda+m+n+p-1)(2\lambda+m+n+p+1)} \\
&\quad - \frac{(\lambda+p)(2\lambda+m-1)(2\lambda+n+p-m-1)}{(n+p-m-1)(2\lambda+p-1)(2\lambda+m+n+p-1)} \\
&\quad \left. + \frac{(\lambda+m)(2\lambda+m+n-p-1)}{(m+n-p+1)(2\lambda+m+n+p-1)} \right). \tag{25}
\end{aligned}$$

Since the rational coefficient multiplying the factorials and Pochhammer symbols vanishes, we can see that this identity is trivially satisfied. So this is a clear validation of the contiguous relations implied by Theorem 3.1. Note that similar validations can be obtained by using (19), (20), which we leave to the reader.

4.2 Linearization of a product of three Gegenbauer polynomials

Now we present the linearization formula for a product of three Gegenbauer polynomials.

Theorem 4.2. Let $p, m, n \in \mathbb{N}_0$ and without loss of generality $p \leq m \leq n$, $\lambda \in (-\frac{1}{2}, \infty) \setminus \{0\}$, $x \in \mathbb{C}$. Then

$$C_p^\lambda(x)C_m^\lambda(x)C_n^\lambda(x) = \sum_{k=0}^{\lfloor \frac{p+m+n}{2} \rfloor} F_{k,p,m,n}^\lambda C_{p+m+n-2k}^\lambda(x), \tag{26}$$

where

$$F_{k,p,m,n}^\lambda := \begin{cases} D_{k,p,m,n}^\lambda, & \text{if } 0 \leq k \leq p-1, \\ E_{k,p,m,n}^\lambda, & \text{if } p \leq k \leq \lfloor \frac{p+m+n}{2} \rfloor, \end{cases} \tag{27}$$

$$\begin{aligned}
D_{k,p,m,n}^\lambda &:= \frac{(m+n)!(\lambda)_m(\lambda)_n(\lambda+m+n+p-2k)(m+n+p-2k)!(\lambda)_k(\lambda)_{p-k}(\lambda)_{m+n-k}(2\lambda)_{m+n+p-k}}{m!n!(\lambda)_{m+n}(\lambda+m+n+p-k)k!(p-k)!(m+n-k)!(\lambda)_{m+n+p-k}(2\lambda)_{m+n+p-2k}} \\
&\quad \times {}_{11}F_{10} \left(\begin{matrix} \{\lambda+p-k\}, \left\{ \begin{matrix} -k \\ -m \\ -n \\ -m-n+k \end{matrix} \right\}, \{-\lambda-m-n-p+k\}, \frac{-\lambda+2-m-n}{2}, \frac{-2\lambda+\{\frac{1}{2}\}-m-n}{2} \\ 1+p-k, \left\{ \begin{matrix} -\lambda+1-k \\ -\lambda+1-m \\ -\lambda+1-n \\ -\lambda+1-m-n+k \end{matrix} \right\}, \{-2\lambda+1-m-n-p+k\}, \frac{-\lambda-m-n}{2}, \frac{\{0\}-m-n}{2} \end{matrix} ; 1 \right), \tag{28}
\end{aligned}$$

$$\begin{aligned}
E_{k,p,m,n}^\lambda &:= \frac{(\lambda+m+n+p-2k)(m+n+2p-2k)!(\lambda)_p(\lambda)_{k-p}(\lambda)_{m+p-k}(\lambda)_{n+p-k}(\lambda)_{m+n+p-2k}(2\lambda)_{m+n+p-k}}{(\lambda+m+n+p-k)p!(k-p)!(m+p-k)!(n+p-k)!(\lambda)_{m+n+p-k}(\lambda)_{m+n+2p-2k}(2\lambda)_{m+n+p-2k}} \\
&\quad \times {}_{11}F_{10} \left(\begin{matrix} \{\lambda+k-p\}, \left\{ \begin{matrix} -p \\ -m-p+k \\ -n-p+k \\ -m-n-p+2k \end{matrix} \right\}, \{-\lambda-m-n-p+k\}, \frac{-\lambda+2k-2p-m-n+2}{2}, \frac{-2\lambda+\{\frac{1}{2}\}+2k-2p-m-n}{2} \\ 1+k-p, \left\{ \begin{matrix} -\lambda+1-p \\ -\lambda+1-m-p+k \\ -\lambda+1-n-p+k \\ -\lambda+1-m-n-p+2k \end{matrix} \right\}, \{-2\lambda+1-m-n-p+k\}, \frac{-\lambda+2k-2p-m-n}{2}, \frac{\{0\}+2k-2p-m-n}{2} \end{matrix} ; 1 \right). \tag{29}
\end{aligned}$$

Proof. Consider (26), where $F_{k,p,m,n}^\lambda$ is to be determined. Using (17), one obtains the following integral of a product of four Gegenbauer polynomials which is equivalent to the linearization of a product of three Gegenbauer polynomials, namely

$$F_{k,p,m,n}^\lambda = \frac{1}{h_{m+n+p-2k}^\lambda} \int_{-1}^1 C_p^\lambda(x) C_m^\lambda(x) C_n^\lambda(x) C_{p+m+n-2k}^\lambda(x) (1-x^2)^{\lambda-\frac{1}{2}} dx. \quad (30)$$

Now use (15) to write the product of Gegenbauer polynomials of degree m and n as a single sum over $l \in \mathbb{N}_0$, $0 \leq l \leq m$. This converts (30) into a single sum of an integral of a product of three Gegenbauer polynomials whose terms can be evaluated using (17). After avoiding the factor $l+p-k$ becoming negative (in which case $F_{k,p,m,n}^\lambda$ vanishes), we obtain

$$F_{k,p,m,n}^\lambda = \sum_{l=\max(0,k-p)}^m B_{l,m,n}^\lambda B_{k-l,p,m+n-2l}^\lambda,$$

which breaks the linearization formula into two regions depending on k , namely

$$C_p^\lambda(x) C_m^\lambda(x) C_n^\lambda(x) = \sum_{k=0}^{p-1} C_{p+m+n-2k}^\lambda(x) D_{k,p,l,m}^\lambda + \sum_{k=p}^{\lfloor \frac{p+m+n}{2} \rfloor} C_{p+m+n-2k}^\lambda(x) E_{k,p,l,m}^\lambda,$$

where

$$D_{k,p,m,n}^\lambda := \sum_{l=0}^m B_{l,m,n}^\lambda B_{k-l,p,m+n-2l}^\lambda, \quad E_{k,p,m,n}^\lambda := \sum_{l=0}^{p+m-k} B_{l+k-p,m,n}^\lambda B_{p-l,p,m+n+2p-2k-2l}^\lambda.$$

Factoring the products of linearization coefficients in terms of Pochhammer symbols and factorials completes the proof. \blacksquare

Remark 4.3. Setting $p = 0$ in (26) straightforwardly produces (15) since the first sum vanishes and the ${}_{11}F_{10}(1)$ in the second sum is unity.

Corollary 4.4. Let $p, m, n \in \mathbb{N}_0$ and without loss of generality $p \leq m \leq n$, $\lambda \in (-\frac{1}{2}, \infty) \setminus \{0\}$. Then

$$\frac{(2\lambda)_p (2\lambda)_m (2\lambda)_n}{p! m! n!} = \frac{(2\lambda)_{p+m+n}}{(p+m+n)!} \sum_{k=0}^{\lfloor \frac{p+m+n}{2} \rfloor} F_{k,p,m,n}^\lambda \frac{\left(\frac{-p-m-n}{2}\right)_k \left(\frac{-p-m-n+1}{2}\right)_k}{\left(\frac{-2\lambda-p-m-n+1}{2}\right)_k \left(\frac{-2\lambda-p-m-n+2}{2}\right)_k}. \quad (31)$$

Proof. Let $x = 1$ in (26) and [2, Table 18.6.1] $C_n^\lambda(1) = (2\lambda)_n/n!$. \blacksquare

Define the following integral of the product of four Gegenbauer polynomials

$$C(p, m, n, l; \lambda) := \int_{-1}^1 C_p^\lambda(x) C_m^\lambda(x) C_n^\lambda(x) C_l^\lambda(x) (1-x^2)^{\lambda-\frac{1}{2}} dx, \quad (32)$$

where $l = p + m + n - 2k$. We now present the contiguous relations for the integral of a product of four Gegenbauer polynomials.

Theorem 4.5. *Let $p, m, n \in \mathbb{N}_0$, and without loss of generality $p \leq m \leq n$, and $l \in \{0, \dots, p + m + n\}$ such that $p + m + n - l \pm 1$ is even. Then*

$$\begin{aligned} & (m + \lambda)(p + 1) F_{\frac{1}{2}(p+m+n-l+1), p+1, m, n} + (p + 2\lambda - 1)(m + \lambda) F_{\frac{1}{2}(p+m+n-l-1), p-1, m, n} \\ &= (p + \lambda)(m + 1) F_{\frac{1}{2}(p+m+n-l+1), p, m+1, n} + (p + \lambda)(m + 2\lambda - 1) F_{\frac{1}{2}(p+m+n-l-1), p, m-1, n}, \end{aligned} \quad (33)$$

$$\begin{aligned} & (n + \lambda)(p + 1) F_{\frac{1}{2}(p+m+n-l+1), p+1, m, n} + (p + 2\lambda - 1)(n + \lambda) F_{\frac{1}{2}(p+m+n-l-1), p-1, m, n} \\ &= (p + \lambda)(n + 1) F_{\frac{1}{2}(p+m+n-l+1), p, m, n+1} + (p + \lambda)(n + 2\lambda - 1) F_{\frac{1}{2}(p+m+n-l-1), p, m, n-1}, \end{aligned} \quad (34)$$

$$\begin{aligned} & (p + 1) F_{\frac{1}{2}(p+m+n-l+1), p+1, m, n} + (p + 2\lambda - 1) F_{\frac{1}{2}(p+m+n-l-1), p-1, m, n} \\ &= \frac{(2\lambda + l)(p + \lambda)}{(\lambda + l + 1)} F_{\frac{1}{2}(p+m+n-l-1), p, m, n} + \frac{l(p + \lambda)}{(\lambda + l - 1)} F_{\frac{1}{2}(p+m+n-l+1), p, m, n}, \end{aligned} \quad (35)$$

$$\begin{aligned} & (n + \lambda)(m + 1) F_{\frac{1}{2}(p+m+n-l+1), p, m+1, n} + (m + 2\lambda - 1)(n + \lambda) F_{\frac{1}{2}(p+m+n-l-1), p, m-1, n} \\ &= (m + \lambda)(n + 1) F_{\frac{1}{2}(p+m+n-l+1), p, m, n+1} + (m + \lambda)(n + 2\lambda - 1) F_{\frac{1}{2}(p+m+n-l-1), p, m, n-1}, \end{aligned} \quad (36)$$

$$\begin{aligned} & (m + 1) F_{\frac{1}{2}(p+m+n-l+1), p, m+1, n} + (m + 2\lambda - 1) F_{\frac{1}{2}(p+m+n-l-1), p, m-1, n} \\ &= \frac{(2\lambda + l)(m + \lambda)}{(\lambda + l + 1)} F_{\frac{1}{2}(p+m+n-l-1), p, m, n} + \frac{l(m + \lambda)}{(\lambda + l - 1)} F_{\frac{1}{2}(p+m+n-l+1), p, m, n}, \end{aligned} \quad (37)$$

$$\begin{aligned} & (n + 1) F_{\frac{1}{2}(p+m+n-l+1), p, m, n+1} + (n + 2\lambda - 1) F_{\frac{1}{2}(p+m+n-l-1), p, m, n-1} \\ &= \frac{(2\lambda + l)(n + \lambda)}{(\lambda + l + 1)} F_{\frac{1}{2}(p+m+n-l-1), p, m, n} + \frac{l(n + \lambda)}{(\lambda + l - 1)} F_{\frac{1}{2}(p+m+n-l+1), p, m, n}, \end{aligned} \quad (38)$$

where

$$F_{\frac{1}{2}(p+m+n-l\pm 1), p, m, n}^\lambda := \begin{cases} D_{\frac{1}{2}(p+m+n-l\pm 1), p, m, n}^\lambda, & \text{if } m + n - p + 2 \leq l \leq m + n + p, \\ E_{\frac{1}{2}(p+m+n-l\pm 1), p, m, n}^\lambda, & \text{if } 0 \leq l \leq m + n - p. \end{cases}$$

Proof. Aside from the contiguous relations (18)–(20), there are three more

$$(l + \lambda)(p + 1)C_1^+ + (p + 2\lambda - 1)(l + \lambda)C_1^- = (p + \lambda)(l + 1)C_4^+ + (p + \lambda)(l + 2\lambda - 1)C_4^-, \quad (39)$$

$$(l + \lambda)(m + 1)C_2^+ + (m + 2\lambda - 1)(l + \lambda)C_2^- = (m + \lambda)(l + 1)C_4^+ + (m + \lambda)(l + 2\lambda - 1)C_4^-, \quad (40)$$

$$(l + \lambda)(n + 1)C_3^+ + (n + 2\lambda - 1)(l + \lambda)C_3^- = (n + \lambda)(l + 1)C_4^+ + (n + \lambda)(l + 2\lambda - 1)C_4^-, \quad (41)$$

because we now have an integral of a product of four orthogonal polynomials and therefore there will be $\binom{4}{2} = 6$ contiguous relations. The parameters p, m, n and l are associated with the subscripts 1, 2, 3 and 4 respectively, and $l = p + m + n - 2k$. Applying this to the six contiguous relations (18)–(20), (39)–(41), and using Theorem 4.2 completes the proof. \blacksquare

5 Hermite polynomials

The Hermite polynomials can be defined as [6, (9.15.1)]

$$H_n(x) = (2x)^n {}_2F_0 \left(\begin{matrix} -\frac{1}{2}n, -\frac{1}{2}(n-1) \\ - \\ - \end{matrix}; -\frac{1}{x^2} \right), \quad (42)$$

where $n \in \mathbb{N}_0$, $x \in \mathbb{C}$. Hermite polynomials are orthogonal on $x \in (-\infty, \infty)$, with orthogonality relation [6, (9.15.2)]

$$\int_{-\infty}^{\infty} H_m(x) H_n(x) e^{-x^2} dx = \sqrt{\pi} 2^n n! \delta_{m,n} =: h_n \delta_{m,n}. \quad (43)$$

The recurrence relation (6) for Hermite polynomials is given through

$$A_n = 2, \quad B_n = 0, \quad C_n = 2n. \quad (44)$$

The definite integral of a product of $N \in \mathbb{N}_0$, $N \geq 2$, Hermite polynomials is defined by

$$\mathbf{H}(\mathbf{n}) := \int_{-\infty}^{\infty} H_{n_1}(x) \cdots H_{n_{N+1}}(x) e^{-x^2} dx,$$

where $\mathbf{n} := \{n_1, \dots, n_{N+1}\}$. Note that $\mathbf{H}(\mathbf{n})$ has a generating function given by [3, Exercise 4.11]

$$\exp\left(2 \sum_{1 \leq i < j \leq k} t_i t_j\right) = \frac{1}{\sqrt{\pi}} \sum_{n_1, \dots, n_k=0}^{\infty} \frac{t_1^{n_1} \cdots t_k^{n_k}}{n_1! \cdots n_k!} \mathbf{H}(\mathbf{n}). \quad (45)$$

5.1 Linearization of a product of two Hermite polynomials

The Hermite polynomials (42) have a linearization formula given by [2, (18.18.22)]

$$H_m(x)H_n(x) = \sum_{k=0}^m \mathbf{b}_{k,m,n} H_{m+n-2k}(x), \quad (46)$$

where $m, n \in \mathbb{N}_0$, without loss of generality $n \geq m$, and

$$\mathbf{b}_{k,m,n} := \frac{2^k m! n!}{k!(m-k)!(n-k)!}.$$

The definite integral corresponding to the linearization of a product of two Hermite polynomials is given by

$$\mathbf{H}(p, m, n) := \int_{-\infty}^{\infty} H_p(x)H_m(x)H_n(x) e^{-x^2} dx. \quad (47)$$

Using (46) and orthogonality (43), we see that $\mathbf{H}(p, m, n)$ is given by

$$\mathbf{H}(p, m, n) = \begin{cases} 0, & \text{if } p > n+m \text{ or } m > n+p \text{ or } n > m+p \\ & \text{or } ((p+m+n) \bmod 2) = 1, \\ \frac{\sqrt{\pi} m! n! p! 2^{\lfloor \frac{p+m+n}{2} \rfloor}}{\lfloor \frac{m+n-p}{2} \rfloor! \lfloor \frac{m+p-n}{2} \rfloor! \lfloor \frac{n+p-m}{2} \rfloor!}, & \text{otherwise.} \end{cases} \quad (48)$$

Corollary 5.1. *Let $|t_1|, |t_2|, |t_3| < 1$. Then $\mathbf{H}(p, m, n)$ has the following multilinear generating function (45), given by*

$$\exp(t_1 t_2 + t_1 t_3 + t_2 t_3) = \sum_{n,m=0}^{\infty} \sum_{p=0}^{m \wedge n} \frac{2^{p+m \vee n} t_1^{2p+|n-m|} t_2^m t_3^n}{(m \wedge n - p)! (p+|n-m|)! p!}.$$

Proof. Starting with (45), (48), for a fixed $m, n \in \mathbb{N}_0$, p is non-vanishing for $|n-m| \leq p \leq n+m$. Shifting the p -index by $|n-m|$ and scaling p by a power of two to remove the remaining vanishing values of $\mathbf{H}(p, n, m)$ completes the proof. \blacksquare

Remark 5.2. *Let $x \in \mathbb{C}$, $p, m, n \in \mathbb{N}_0$, and without loss of generality assume $m \leq n$. Further, let $p \in \{n-m, \dots, n+m\}$ and $p+m+n \pm 1$ even. Combining Theorem 3.1 with the coefficients*

of the recurrence relation given by (44) yields three contiguous relations for the integral of the product of three Hermite polynomials:

$$H_2^+ - H_1^+ - 2pH_1^- + 2mH_2^- = 0, \quad (49)$$

$$H_3^+ - H_1^+ - 2pH_1^- + 2nH_3^- = 0, \quad (50)$$

$$H_3^+ - H_2^+ - 2mH_2^- + 2nH_3^- = 0. \quad (51)$$

Using (48) with (49) the following identity is obtained:

$$\begin{aligned} & (m+1) \left[\frac{m+n-p-1}{2} \right]! \left[\frac{n+p-m+1}{2} \right]! \left[\frac{m+p-n-1}{2} \right]! \\ & - (p+1) \left[\frac{m+n-p+1}{2} \right]! \left[\frac{n+p-m-1}{2} \right]! \left[\frac{m+p-n-1}{2} \right]! \\ & - \left[\frac{m+p-n+1}{2} \right]! \left[\frac{m+n-p-1}{2} \right]! \left[\frac{n+p-m+1}{2} \right]! \\ & + \left[\frac{m+n-p+1}{2} \right]! \left[\frac{m+p-n+1}{2} \right]! \left[\frac{n+p-m-1}{2} \right]! = 0. \end{aligned} \quad (52)$$

This can be rearranged to the following

$$\begin{aligned} & \left(\frac{n+p-m+1}{2} \left((m+1) - \frac{m+p-n+1}{2} \right) + \frac{m+n-p+1}{2} \left(-(p+1) + \frac{m+p-n+1}{2} \right) \right) \\ & \times \left[\frac{m+n-p-1}{2} \right]! \left[\frac{n+p-m-1}{2} \right]! \left[\frac{m+p-n-1}{2} \right]! = 0. \end{aligned} \quad (53)$$

Since the coefficient multiplying the factorials vanishes, we can see that the identity is trivially satisfied. So this is a clear validation of the contiguous relations implied by Theorem 3.1. Note that similar validations can be obtained by using (50), (51), which we leave to the reader.

5.2 Linearization of a product of three Hermite polynomials

Theorem 5.3. Let $p, m, n \in \mathbb{N}_0$ and without loss of generality $p \leq m \leq n$, $x \in \mathbb{C}$. Then

$$H_p(x)H_m(x)H_n(x) = \sum_{k=0}^{\lfloor \frac{p+m+n}{2} \rfloor} f_{k,p,m,n} H_{p+m+n-2k}(x), \quad (54)$$

where

$$f_{k,p,m,n} := \begin{cases} d_{k,p,m,n}, & \text{if } 0 \leq k \leq p-1, \\ e_{k,p,m,n}, & \text{if } p \leq k \leq \lfloor \frac{p+m+n}{2} \rfloor, \end{cases} \quad (55)$$

$$d_{k,p,m,n} := \frac{(m+n)!p!2^k}{k!(p-k)!(m+n-k)!} {}_4F_3 \left(\begin{matrix} -k, -m, -n, -m-n+k, \\ 1+p-k, \frac{\{0\}-m-n}{2} \end{matrix}; \frac{1}{4} \right), \quad (56)$$

$$\begin{aligned} e_{k,p,m,n} &:= \frac{m!n!(m+n+2p-2k)!2^k}{(k-p)!(m+p-k)!(n+p-k)!(p+m+n-2k)!} \\ & \times {}_4F_3 \left(\begin{matrix} -p, -m-p+k, -n-p+k, -m-n-p+2k, \\ 1+k-p, \frac{\{0\}+2k-2p-m-n}{2} \end{matrix}; \frac{1}{4} \right). \end{aligned} \quad (57)$$

Proof. Using (cf. [6, (9.8.34)])

$$H_n(x) = n! \lim_{\alpha \rightarrow \infty} \alpha^{-\frac{n}{2}} C_n^\alpha \left(\frac{x}{\sqrt{\alpha}} \right)$$

four times in (26), and [2, (5.11.12)], the result follows. \blacksquare

Remark 5.4. Setting $p = 0$ in (54) straightforwardly produces (46) since the first sum vanishes and the ${}_4F_3(\frac{1}{4})$ in the second sum is unity.

Define the following definite integral of a product of four Hermite polynomials

$$H(k, p, m, n) := \int_{-\infty}^{\infty} H_k(x) H_p(x) H_m(x) H_n(x) e^{-x^2} dx.$$

Using orthogonality and the linearization relation (54) one produces the following corollary.

Corollary 5.5. Let $k, p, m, n \in \mathbb{N}_0$. Then

$$H(k, p, m, n) = \begin{cases} E_{k,p,m,n}, & \text{if } 0 \leq k \leq m + n - p, \\ D_{k,p,m,n}, & \text{if } m + n - p + 2 \leq k \leq m + n + p, \\ 0, & \text{otherwise,} \end{cases} \quad (58)$$

where

$$E_{k,p,m,n} := \frac{\sqrt{\pi} m! n! (p+k)! 2^{\lfloor \frac{k+p+m+n}{2} \rfloor}}{\lfloor \frac{m+n-p-k}{2} \rfloor! \lfloor \frac{k+p+n-m}{2} \rfloor! \lfloor \frac{k+p+m-n}{2} \rfloor!} {}_4F_3 \left(\begin{matrix} -k, -p, \lfloor \frac{n-m-p-k}{2} \rfloor, \lfloor \frac{m-n-p-k}{2} \rfloor \\ 1 + \lfloor \frac{m+n-p-k}{2} \rfloor, \frac{\{0\}}{2} - k - p \end{matrix} ; \frac{1}{4} \right),$$

$$D_{k,p,m,n} := \frac{\sqrt{\pi} k! p! (m+n)! 2^{\lfloor \frac{p+k+m+n}{2} \rfloor}}{\lfloor \frac{k+p-m-n}{2} \rfloor! \lfloor \frac{m+n+p-k}{2} \rfloor! \lfloor \frac{m+n+k-p}{2} \rfloor!} {}_4F_3 \left(\begin{matrix} -m, -n, \lfloor \frac{p-k-m-n}{2} \rfloor, \lfloor \frac{k-p-m-n}{2} \rfloor \\ 1 + \lfloor \frac{p+k-m-n}{2} \rfloor, \frac{\{0\}}{2} - m - n \end{matrix} ; \frac{1}{4} \right).$$

Proof. Start with Theorem 5.3 and then integrate both sides using the property of orthogonality for Hermite polynomials (43). This leads to the following integral for the linearization coefficients, namely

$$f_{k,p,m,n} = \frac{1}{h_{m+n+p-2k}} \int_{-\infty}^{\infty} H_p(x) H_m(x) H_n(x) H_{p+m+n-2k}(x) e^{-x^2} dx. \quad (59)$$

This allows us to write the integral of a product of four Hermite polynomials as

$$H(p, m, n, l) = h_l f_{\frac{1}{2}(p+m+n-l), p, m, n}. \quad (60)$$

\blacksquare

Theorem 5.6. Let $p, m, n \in \mathbb{N}_0$ and without loss of generality $p \leq m \leq n$, $k \in \{0, \dots, p+m+n\}$. Then the contiguous relations for the integral for a product of four Hermite polynomials are given by

$$F_{k,p,m+1,n} - F_{k,p+1,m,n} + 2mF_{k,p,m-1,n} - 2pF_{k,p-1,m,n} = 0, \quad (61)$$

$$F_{k,p,m,n+1} - F_{k,p+1,m,n} + 2nF_{k,p,m,n-1} - 2pF_{k,p-1,m,n} = 0, \quad (62)$$

$$F_{k+1,p,m,n} - F_{k,p+1,m,n} + 2kF_{k-1,p,m,n} - 2pF_{k,p-1,m,n} = 0, \quad (63)$$

$$F_{k,p,m,n+1} - F_{k,p,m+1,n} + 2nF_{k,p,m,n-1} - 2mF_{k,p,m-1,n} = 0, \quad (64)$$

$$F_{k+1,p,m,n} - F_{k,p,m+1,n} + 2kF_{k-1,p,m,n} - 2mF_{k,p,m-1,n} = 0, \quad (65)$$

$$F_{k+1,p,m+1,n} - F_{k,p,m,n+1} + 2kF_{k-1,p,m,n} - 2nF_{k,p,m,n-1} = 0, \quad (66)$$

where

$$F_{k,p,m,n} = \begin{cases} E_{k,p,m,n}, & \text{if } 0 \leq k \leq m+n-p, \\ D_{k,p,m,n}, & \text{if } m+n-p+2 \leq k \leq m+n+p, \\ 0, & \text{otherwise,} \end{cases} \quad (67)$$

Proof. In addition to the contiguous functions shown in Remark 5.2, there are three additional ones:

$$H_2^+ - H_1^+ - 2pH_1^- + 2mH_2^- = 0, \quad (68)$$

$$H_3^+ - H_1^+ - 2pH_1^- + 2nH_3^- = 0, \quad (69)$$

$$H_3^+ - H_2^+ - 2mH_2^- + 2nH_3^- = 0. \quad (70)$$

Combining these with (58), the contiguous relations of the theorem are obtained. \blacksquare

6 Jacobi polynomials

The Jacobi polynomials can be defined as [2, (18.5.7)]

$$P_n^{(\alpha,\beta)} = \frac{(\alpha+1)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{1-x}{2} \right). \quad (71)$$

where $n \in \mathbb{N}_0$, $\alpha, \beta > -1$, $x \in \mathbb{C}$. Jacobi polynomials are orthogonal on $x \in (-1, 1)$, with the orthogonality relation [2, (18.3)]

$$\int_{-1}^1 P_m^{(\alpha,\beta)}(x) P_n^{(\alpha,\beta)}(x) (1-x)^\alpha (1+x)^\beta dx = \frac{2^{\alpha+\beta+1} \Gamma(n+\alpha+1) \Gamma(n+\beta+1)}{(2n+\alpha+\beta+1) \Gamma(n+\alpha+\beta+1) n!} \delta_{n,m} =: h_n \delta_{n,m}. \quad (72)$$

The coefficients for the three-term recurrence relation for the Jacobi polynomials are given by [2, (18.9.2)]

$$A_n := \frac{(2n+\alpha+\beta+1)(2n+\alpha+\beta+2)}{2(n+1)(n+\alpha+\beta+1)}, \quad (73)$$

$$B_n := \frac{(\alpha^2 - \beta^2)(2n+\alpha+\beta+1)}{2(n+1)(n+\alpha+\beta+1)(2n+\alpha+\beta)}, \quad (74)$$

$$C_n := \frac{(n+\alpha)(n+\beta)(2n+\alpha+\beta+2)}{(n+1)(n+\alpha+\beta+1)(2n+\alpha+\beta)}. \quad (75)$$

6.1 Linearization of a product of two Jacobi polynomials

The linearization formula for Jacobi polynomials was given in a fundamental work by Rahman [7, (1.9)]. Rahman expressed the linearization coefficient in terms of a ${}_9F_8(1)$. His result was given in terms of some other variables $s = n - m$, $j = k - n + m$, and also contained a typographical error. The typographical error was that the term $(2s - 2n - \alpha - \beta)$ in Rahman's original publication should have been written as $(2s - 2n - \alpha - \beta)_j$. (Note that in [5, (2.1.1)], it was realized that Rahman's result contained a typographical error.) The corrected and further simplified version of Rahman's result is given as follows.

Theorem 6.1. *Rahman (1981).* Let $m, n \in \mathbb{N}_0$ and without loss of generality, $n \geq m$, $\alpha, \beta \in \mathbb{C}$. Then

$$P_m^{(\alpha, \beta)}(x)P_n^{(\alpha, \beta)}(x) = \sum_{k=0}^{2m} a_{k, m, n}^{\alpha, \beta} P_{k+n-m}^{(\alpha, \beta)}(x), \quad (76)$$

where

$$\begin{aligned} a_{k, m, n}^{\alpha, \beta} := & \frac{(\alpha + 1, \beta + 1)_n (\alpha + \beta + 1)_{2n-2m} (\alpha + \beta + 1)_{2m} (\alpha + \beta + 1 + 2n - 2m + 2k)}{m! (\alpha + \beta + 1)_m (\alpha + 1, \beta + 1)_{n-m} (\alpha + \beta + 2)_{2n} (\alpha + \beta + 1)} \\ & \times \frac{(n - m + 1, \alpha + \beta + 2n - 2m + 1, 2\alpha + 2\beta + 2n + 2, -2m, \alpha - \beta)_k}{k! (2\beta + 2n - 2m + 2, \alpha + n - m + 1, \alpha + \beta + 2n + 2, -\alpha - \beta - 2m)_k} \\ & \times {}_9F_8 \left(\begin{matrix} \beta + n - m + \frac{1}{2}, \frac{\beta + n - m + \frac{5}{2}}{2}, \beta + \frac{1}{2}, \beta + n + 1, -\alpha - m, \frac{\alpha + \beta + k + \frac{1}{2}}{2} + n - m, \frac{-k + \frac{0}{1}}{2} \\ \frac{\beta + n - m + \frac{1}{2}}{2}, n - m + 1, \frac{1}{2} - m, \alpha + \beta + n + \frac{3}{2}, \frac{\beta - \alpha - k + \frac{1}{2}}{2}, \frac{k + \frac{2}{3}}{2} + \beta + n - m \end{matrix} ; 1 \right). \end{aligned} \quad (77)$$

Proof. See Rahman (1981) [7, (1.9)] and [1, (3.3)] for description of correction. \blacksquare

Corollary 6.2. Let $l, m, n \in \mathbb{N}_0$ and without loss of generality $m \leq n$ and $l \in \{n - m, \dots, n + m\}$ using the linearization formula for Jacobi polynomials, it is possible to write the integral of the product of three Jacobi polynomials as follows:

$$P(l, m, n; \alpha, \beta) := \int_{-1}^1 P_m^{(\alpha, \beta)}(x) P_n^{(\alpha, \beta)}(x) P_l^{(\alpha, \beta)}(x) (1-x)^\alpha (1+x)^\beta dx = a_{l+m-n, n, m}^{\alpha, \beta} h_l, \quad (78)$$

where h_l is defined in (72).

Theorem 6.3. Let $l, m, n \in \mathbb{N}_0$ and without loss of generality $m \leq n$ and $l \in \{n - m, \dots, n + m\}$

$$\begin{aligned} & \frac{2(\alpha - \beta)(\alpha + \beta)(m - n)(\alpha + \beta + m + n + 1)(\alpha + \beta + 2n + 1)}{(\alpha + \beta + 2m)(\alpha + \beta + 2n)} a_{l+m-n, n, m}^{\alpha, \beta} \\ & + \frac{(\alpha + \beta + m + 1)(m + 1)(\alpha + \beta + 2n + 1)(\alpha + \beta + 2n + 2)}{(\alpha + \beta + 2m + 1)} a_{l+m-n+1, n, m+1}^{\alpha, \beta} \\ & + \frac{(\alpha + m)(\beta + m)(\alpha + \beta + 2m + 2)(\alpha + \beta + 2n + 1)(\alpha + \beta + 2n + 2)}{(\alpha + \beta + 2m)(\alpha + \beta + 2m + 1)} a_{l+m-n-1, n, m-1}^{\alpha, \beta} \\ & - (\alpha + \beta + 2m + 2)(\alpha + \beta + n + 1)(n + 1) a_{l+m-n-1, n+1, m}^{\alpha, \beta} \\ & - \frac{(\alpha + \beta + 2m + 2)(\alpha + n)(\beta + n)(\alpha + \beta + 2n + 2)}{(\alpha + \beta + 2n)} a_{l+m-n+1, n-1, m}^{\alpha, \beta} = 0, \end{aligned} \quad (79)$$

$$\begin{aligned} & \frac{2(\alpha - \beta)(\alpha + \beta)(l - n)(\alpha + \beta + n + l + 1)}{(\alpha + \beta + 2l)(\alpha + \beta + 2n)} a_{l+m-n, n, m}^{\alpha, \beta} \\ & - \frac{(\alpha + \beta + 2l + 2)(\alpha + \beta + n + 1)(n + 1)}{(\alpha + \beta + 2n + 1)} a_{l+m-n-1, n+1, m}^{\alpha, \beta} \\ & - \frac{(\alpha + n)(\beta + n)(\alpha + \beta + 2l + 2)(\alpha + \beta + 2n + 2)}{(\alpha + \beta + 2n + 1)(\alpha + \beta + 2n)} a_{l+m-n+1, n-1, m}^{\alpha, \beta} \\ & + \frac{(\alpha + \beta + 2n + 2)(\alpha + l + 1)(\beta + l + 1)(\alpha + \beta + 2l + 1)}{(\alpha + \beta + 2l + 3)} a_{l+m-n+1, n, m}^{\alpha, \beta} \\ & + \frac{l(\alpha + \beta + l)(\alpha + \beta + 2l + 2)(\alpha + \beta + 2n + 2)}{(\alpha + \beta + 2l - 1)(\alpha + \beta + 2l)} a_{l+m-n-1, n, m}^{\alpha, \beta} = 0, \end{aligned} \quad (80)$$

$$\begin{aligned}
& \frac{2(\alpha - \beta)(\alpha + \beta)(l - m)(\alpha + \beta + m + l + 1)}{(\alpha + \beta + 2l)(\alpha + \beta + 2l + 2)(\alpha + \beta + 2m)} a_{l+m-n,n,m}^{\alpha,\beta} \\
& - \frac{(m + 1)(\alpha + \beta + m + 1)}{(\alpha + \beta + 2m + 1)} a_{l+m-n+1,n,m+1}^{\alpha,\beta} \\
& - \frac{(\alpha + m)(\beta + m)(\alpha + \beta + 2m + 2)}{(\alpha + \beta + 2m)(\alpha + \beta + 2m + 1)} a_{l+m-n-1,n,m-1}^{\alpha,\beta} \\
& + \frac{(\alpha + l + 1)(\beta + l + 1)(\alpha + \beta + 2m + 2)}{(\alpha + \beta + 2l + 2)(\alpha + \beta + 2l + 3)} a_{l+m-n+1,n,m}^{\alpha,\beta} \\
& + \frac{l(\alpha + \beta + l)(\alpha + \beta + 2m + 2)}{(\alpha + \beta + 2l - 1)(\alpha + \beta + 2l)} a_{l+m-n-1,n,m}^{\alpha,\beta} = 0. \tag{81}
\end{aligned}$$

Proof. To determine the contiguous relations, the coefficients of the three-term recurrence relation defined in (73) are used. Combining these coefficients with the general form of the contiguous relation of Theorem 3.1 results in

$$\begin{aligned}
& \frac{(\alpha - \beta)(\alpha + \beta)(\alpha + \beta + 2n_j)(n_j - n_k)(\alpha + \beta + n_j + n_k + 1)(\alpha + \beta + 2n_j + 1)}{(n_j + 1)(\alpha + \beta + n_j + 1)(\alpha + \beta + 2n_j)(n_k + 1)(\alpha + \beta + n_k + 1)(\alpha + \beta + 2n_k)} P \\
& + \frac{(\alpha + \beta + 2n_k + 1)(\alpha + \beta + 2n_k + 2)}{2(n_k + 1)(\alpha + \beta + n_k + 1)} P_{n_j}^+ \\
& + \frac{(\alpha + n_j)(\beta + n_j)(\alpha + \beta + 2n_j + 2)(\alpha + \beta + 2n_k + 1)(\alpha + \beta + 2n_k + 2)}{2(n_j + 1)(\alpha + \beta + n_k + 1)(\alpha + \beta + 2n_j)(n_k + 1)(\alpha + \beta + n_k + 1)} P_{n_j}^- \\
& = \frac{(\alpha + \beta + 2n_j + 1)(\alpha + \beta + 2n_j + 2)(\alpha + n_k)(\beta + n_k)(\alpha + \beta + 2n_k + 2)}{2(n_j + 1)(\alpha + \beta + n_j + 1)(n_k + 1)(\alpha + \beta + n_k + 1)(\alpha + \beta + 2n_k)} P_{n_k}^+ \\
& + \frac{(\alpha + \beta + 2n_j + 1)(\alpha + \beta + 2n_j + 2)}{2(n_j + 1)(\alpha + \beta + n_j + 1)} P_{n_k}^-.
\end{aligned}$$

Then using Corollary 6.2 with the definition of the coefficient $a_{k,n,m}^{\alpha,\beta}$ in Theorem 6.1 completes the proof. \blacksquare

7 Laguerre polynomials

The Laguerre polynomials can be defined, $x \in \mathbb{C}$, $\alpha > -1$, [6, Section 9.12]

$$L_n^\alpha(x) = \frac{(\alpha + 1)_n}{n!} {}_1F_1\left(\begin{matrix} -n \\ \alpha + 1 \end{matrix}; x\right),$$

with orthogonality relation

$$\int_0^\infty L_m^\alpha(x) L_n^\alpha(x) x^\alpha e^{-x} dx = \frac{\Gamma(\alpha + 1 + n)}{n!} \delta_{m,n} =: h_m \delta_{m,n}.$$

For Laguerre polynomials, one has (6),

$$A_n := \frac{-1}{n + 1}, \quad B_n := \frac{2n + \alpha + 1}{n + 1}, \quad C_n := \frac{n + \alpha}{n + 1}. \tag{82}$$

Define $n_1, \dots, n_{N+1} \in \mathbb{N}_0$, $\Re\alpha > -1$,

$$\mathbf{L}(\mathbf{n}; \alpha) := \int_0^\infty L_{n_1}^\alpha(x) \cdots L_{n_{N+1}}^\alpha(x) x^\alpha e^{-x} dx. \tag{83}$$

Using the generating function for Laguerre polynomials [6, (9.12.10)]

$$(1-t)^{-\alpha-1} \exp\left(-\frac{xt}{1-t}\right) = \sum_{n=0}^{\infty} L_n^\alpha(x) t^n,$$

one finds that (83) has the following generating function [3, (9.3.7)]

$$\sum_{n_1, \dots, n_k=0}^{\infty} \mathbf{L}(\mathbf{n}; \alpha) t_1^{n_1} \cdots t_k^{n_k} = \Gamma(\alpha+1) ((1-t_1) \cdots (1-t_k))^{-\alpha+1} \left(1 + \sum_{j=1}^k \frac{t_j}{1-t_j}\right)^{-\alpha-1}. \quad (84)$$

7.1 Linearization of a product of two Laguerre polynomials

Theorem 7.1. *Let $x \in \mathbb{C}$, $\alpha \in \mathbb{C}$, $n, m \in \mathbb{N}_0$, and without loss of generality $n \geq m$. Then*

$$\begin{aligned} L_m^\alpha(x) L_n^\alpha(x) &= \frac{2^{2m} \left(\frac{1}{2}\right)_m (\alpha+1)_n}{(n-m)!} \\ &\times \sum_{k=n-m}^{n+m} \frac{(-1)^{k+n+m} k!}{(\alpha+1)_k (m+n-k)! (k-n+m)!} {}_3F_2 \left(\begin{matrix} -m-\alpha, \frac{n-m-k+\{0\}}{2} \\ n-m+1, \frac{1}{2}-m \end{matrix}; 1 \right) L_k^\alpha(x). \end{aligned}$$

Proof. Let $x = 1 - 2x\beta^{-1}$ in (76). Applying the relevant limiting relation [6, (9.8.16)]

$$\lim_{\beta \rightarrow \infty} P_n^{(\alpha, \beta)}(1 - 2x\beta^{-1}) = L_n^\alpha(x), \quad (85)$$

produces

$$\lim_{\beta \rightarrow \infty} P_m^{(\alpha, \beta)}(1 - 2x\beta^{-1}) P_n^{(\alpha, \beta)}(1 - 2x\beta^{-1}) = \lim_{\beta \rightarrow \infty} \sum_{k=n-m}^{n+m} P_k^{(\alpha, \beta)}(1 - 2x\beta^{-1}) \mathbf{a}_{k, m, n}^{\alpha, \beta}.$$

One also has the following useful asymptotic result. Let $k \in \mathbb{N}_0$, $b \in \mathbb{C}$, $c \in \mathbb{C} \setminus \{0\}$. Then

$$\lim_{a \rightarrow \infty} \frac{(ca+b)_k}{a^k} = c^k. \quad (86)$$

Using (86) to evaluate the limit on the right-hand side completes the proof. \blacksquare

Define $p, m, n \in \mathbb{N}_0$, $\alpha > -1$,

$$\mathbf{L}_{p, m, n}^\alpha := \int_0^\infty L_p^\alpha(x) L_m^\alpha(x) L_n^\alpha(x) x^\alpha e^{-x} dx,$$

and using (10), (82), we have for $|n-m| \leq p \leq m+n$,

$$\mathbf{L}_{p, m, n}^\alpha = \frac{\Gamma(\alpha+1+m \vee n) (-1)^{p+n+m} 2^{2(m \wedge n)} \left(\frac{1}{2}\right)_{m \wedge n}}{|n-m|! (m+n-p)! (p-|n-m|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m \wedge n, \frac{|n-m|-p+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m \wedge n \end{matrix}; 1 \right). \quad (87)$$

Hence we have the following generating function (84)

$$\begin{aligned} &\sum_{n, m=0}^{\infty} \frac{t^m t_n 2^{2(m \wedge n)} \left(\frac{1}{2}\right)_{m \wedge n} (\alpha+1)_{m \vee n}}{|n-m|!} \sum_{p=0}^{2(m \wedge n)} \frac{(-1)^p t_p^{p+|n-m|}}{(2(m \wedge n) - p)! p!} {}_3F_2 \left(\begin{matrix} -\alpha-m \wedge n, \frac{-p+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m \wedge n \end{matrix}; 1 \right) \\ &= ((1-t_p)(1-t_m)(1-t_n) + t_p(1-t_m)(1-t_n) + t_m(1-t_p)(1-t_n) + t_n(1-t_p)(1-t_m))^{-\alpha-1}. \end{aligned} \quad (88)$$

Theorem 7.2. *Let $p, m, n \in \mathbb{N}_0, a > 1$ such that $|n - m| \leq p \leq m + n$. Then, one has three contiguous relations for the product of the three Laguerre polynomials given by*

$$\begin{aligned}
& \frac{(m+\alpha)(m+n-p)\Gamma(\alpha+1+(m-1)\vee n)2^{2((m-1)\wedge n)}\left(\frac{1}{2}\right)_{(m-1)\wedge n}}{|n-m+1|!(p-|n-m+1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-(m-1)\wedge n, \frac{|n-m+1|-p+\{0\}}{2} \\ 1+|n-m+1|, \frac{1}{2}-(m+1)\wedge n \end{matrix} ; 1 \right) \\
& - \frac{(p+1)(m+n-p)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{|n-m|!(p-|n-m|+1)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+1+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) \\
& + \frac{(m+1)\Gamma(\alpha+1+(m+1)\vee n)2^{2((m+1)\wedge n)}\left(\frac{1}{2}\right)_{(m+1)\wedge n}}{(m+n-p+1)|n-m-1|!(p-|n-m-1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-(m+1)\wedge n, \frac{|n-m-1|-p+\{0\}}{2} \\ 1+|n-m-1|, \frac{1}{2}-(m+1)\wedge n \end{matrix} ; 1 \right) \\
& - \frac{(p+\alpha)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{(m+n-p+1)|n-m|!(p-|n-m-1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+1+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) \\
& - \frac{2(p-m)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{|n-m|!(p-|n-m|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) = 0, \tag{89}
\end{aligned}$$

$$\begin{aligned}
& \frac{(n+\alpha)(m+n-p)\Gamma(\alpha+1+m\vee(n-1))4^{m\wedge(n-1)}\left(\frac{1}{2}\right)_{m\wedge(n-1)}}{|n-m-1|!(p-|n-m-1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge(n-1), \frac{|n-m-1|-p+\{0\}}{2} \\ 1+|n-m-1|, \frac{1}{2}-m\wedge(n-1) \end{matrix} ; 1 \right) \\
& - \frac{(p+1)(m+n-p)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{|n-m|!(p-|n-m|+1)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+1+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) \\
& + \frac{(n+1)\Gamma(\alpha+1+m\vee(n+1))2^{2(m\wedge(n+1))}\left(\frac{1}{2}\right)_{m\wedge(n+1)}}{(m+n-p)|n-m+1|!(p-|n-m+1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge(n+1), \frac{|n-m+1|-p+\{0\}}{2} \\ 1+|n-m+1|, \frac{1}{2}-m\wedge(n+1) \end{matrix} ; 1 \right) \\
& - \frac{(p+\alpha)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{(m+n-p)|n-m|!(p-|n-m|+1)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+1+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) \\
& - \frac{2(p-n)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{|n-m|!(p-|n-m|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) = 0, \tag{90}
\end{aligned}$$

$$\begin{aligned}
& \frac{(n+\alpha)(m+n-p)\Gamma(\alpha+1+(m-1)\vee n)4^{((m-1)\wedge n)}\left(\frac{1}{2}\right)_{(m-1)\wedge n}}{|n-m+1|!(p-|n-m+1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-(m-1)\wedge n, \frac{|n-m+1|-p+\{0\}}{2} \\ 1+|n-m+1|, \frac{1}{2}-(m+1)\wedge n \end{matrix} ; 1 \right) \\
& - \frac{(m+\alpha)(m+n-p)\Gamma(\alpha+1+m\vee(n-1))4^{(m\wedge(n-1))}\left(\frac{1}{2}\right)_{m\wedge(n-1)}}{|n-m-1|!(p-|n-m-1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge(n-1), \frac{|n-m-1|-p+\{0\}}{2} \\ 1+|n-m-1|, \frac{1}{2}-m\wedge(n-1) \end{matrix} ; 1 \right) \\
& - \frac{(m+1)\Gamma(\alpha+1+(m+1)\vee n)2^{2((m+1)\wedge n)}\left(\frac{1}{2}\right)_{(m+1)\wedge n}}{(m+n-p+1)|n-m-1|!(p-|n-m+1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-(m+1)\wedge n, \frac{|n-m-1|-p+\{0\}}{2} \\ 1+|n-m-1|, \frac{1}{2}-(m+1)\wedge n \end{matrix} ; 1 \right) \\
& + \frac{(n+1)\Gamma(\alpha+1+m\vee(n+1))2^{2(m\wedge(n+1))}\left(\frac{1}{2}\right)_{m\wedge(n+1)}}{(m+n-p+1)|n-m+1|!(p-|n-m+1|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge(n+1), \frac{|n-m+1|-p+\{0\}}{2} \\ 1+|n-m+1|, \frac{1}{2}-m\wedge(n+1) \end{matrix} ; 1 \right) \\
& - \frac{2(m-n)\Gamma(\alpha+1+m\vee n)2^{2(m\wedge n)}\left(\frac{1}{2}\right)_{m\wedge n}}{|n-m|!(p-|n-m|)!} {}_3F_2 \left(\begin{matrix} -\alpha-m\wedge n, \frac{|n-m|-p+\{0\}}{2} \\ 1+|n-m|, \frac{1}{2}-m\wedge n \end{matrix} ; 1 \right) = 0. \tag{91}
\end{aligned}$$

Proof. Using Theorem 3.1, one has the following three contiguous relations:

$$(p+1)L_1^+ - (m+1)L_2^+ = (m+\alpha)L_2^- - (p+\alpha)L_1^- + 2(p-m)L, \quad (92)$$

$$(p+1)L_1^+ - (n+1)L_3^+ = (n+\alpha)L_3^- - (p+\alpha)L_1^- + 2(p-n)L, \quad (93)$$

$$(m+1)L_2^+ - (n+1)L_3^+ = (n+\alpha)L_3^- - (m+\alpha)L_2^- + 2(m-n)L. \quad (94)$$

Using the linearization of a product of two Laguerre polynomials (expressed as an integral of a product of three Laguerre polynomials) written as a hypergeometric function ${}_3F_2(1)$ (87) and substituting it into the above equations completes the proof. \blacksquare

7.2 Linearization of a product of two scaled Laguerre polynomials

Consider the integral associated with linearization coefficients for a product of two scaled Laguerre polynomials

$$\mathcal{L}_{p,m,n}^\alpha(a,b) := \int_0^\infty L_p^\alpha(x)L_m^\alpha(ax)L_n^\alpha(bx)x^\alpha e^{-x} dx,$$

where $\Re\alpha > -1$ and $\mathcal{L}_{p,n,n}^\alpha(a,b) = 0$ if $p \geq n+m+1$.

Remark 7.3. To see that $\Re\alpha > -1$ is true, the above integral can be written out in terms of hypergeometric series to become

$$\begin{aligned} \mathcal{L}_{p,m,n}^\alpha(a,b) &= \frac{(\alpha+1)_p(\alpha+1)_m(\alpha+1)_n}{p!m!n!} \\ &\quad \times \sum_{s=0}^p \sum_{k=0}^m \sum_{l=0}^n \frac{(-p)_s(-m)_k(-n)_l a^k b^l}{(\alpha+1)_s(\alpha+1)_k(\alpha+1)_l s!k!l!} \int_0^\infty x^{s+k+l+\alpha} e^{-x} dx. \end{aligned}$$

The integral then becomes $\Gamma(\alpha+1+s+k+l)$, where it is required that $\Re(\alpha+s+k+l) > -1$. Since this must be true for all values that s , k , and l take in the summation, this requirement takes the form $\Re\alpha > -1$.

We now present a theorem which describes double sum representations for this integral.

Theorem 7.4. Let $p, m, n \in \mathbb{N}_0$, $a, b > 0$, $\alpha \in \mathbb{C} \setminus -\mathbb{N}$. Then

$$\begin{aligned} \mathcal{L}_{p,m,n}^\alpha(a,b) &= \frac{\Gamma(\alpha+1)(\alpha+1)_n(-\alpha-p)_m}{p!m!(n+m-p)!} \left(-\frac{a}{b}\right)^m b^p \\ &\quad \times \sum_{k=0}^\infty \sum_{l=0}^\infty \frac{(-m)_k(-\alpha-m)_k(p+1)_l(\alpha+1+p)_l(p-n-m)_{k+l}}{(\alpha+p-m+1)_{k+l}(p+1)_{k+l-m}k!l!} \left(-\frac{b}{a}\right)^k b^l, \quad (95) \end{aligned}$$

Proof. By inserting the definition of the Laguerre polynomials, this integral can be converted to a triple sum, which can be evaluated using the integral definition of the gamma function, namely

$$\begin{aligned} \mathcal{L}_{p,m,n}^\alpha(a,b) &= \frac{(\alpha+1)_p(\alpha+1)_m(\alpha+1)_n}{p!m!n!} \\ &\quad \times \sum_{s=0}^p \sum_{k=0}^m \sum_{l=0}^n \frac{(-p)_s(-m)_k(-n)_l \Gamma(\alpha+1+s+k+l) a^k b^l}{(\alpha+1)_s(\alpha+1)_k(\alpha+1)_l s!k!l!}. \end{aligned}$$

By re-writing $\Gamma(\alpha+1+k+l+s) = \Gamma(\alpha+1)(\alpha+1)_{k+l}(\alpha+1+k+l)_s$, one can write the sum over s as a terminating Gauss hypergeometric series at unity. The hypergeometric series can be

evaluated using the Chu-Vandermonde identity [2, (15.4.24)]. This converts the triple sum into a double sum which produces

$$\begin{aligned} \mathcal{L}_{p,m,n}^\alpha(a,b) &= \frac{\Gamma(\alpha+1+m)\Gamma(\alpha+1+n)}{p!m!\Gamma(\alpha+1)} \\ &\times \sum_{k=0}^m \sum_{l=0}^{n-p+k} \frac{(-1)^{l+k}(-m)_k(p+1)_l(\alpha+1+p)_l a^k b^{p+l-k}}{(\alpha+1)_k(p+1)_{l-k}(\alpha+1+p)_{l-k}(n-p+k-l)!k!l!}. \end{aligned} \quad (96)$$

In order to extend the sum indices to infinity, one can reverse the order of the k index by setting $k \mapsto m-k$. Performing a series of standard manipulations for Pochhammer symbols produces the final form which completes the proof. \blacksquare

An alternative form of the integral $\mathcal{L}_{p,m,n}^\alpha(a,b)$ follows by utilizing the identity in Theorem 7.4.

$$(p+1)_{r-m} = \frac{(p-m)!}{p!} (p-m+1)_r, \quad (97)$$

which is valid for $p \geq m$.

Corollary 7.5. *Let $p, m, n \in \mathbb{N}_0$, $p \geq m$, $a, b > 0$, $\Re\alpha > -1$. Then*

$$\mathcal{L}_{p,m,n}^\alpha(a,b) = \frac{\Gamma(\alpha+1+n)(-\alpha-p)_m}{m!(p-m)!(n+m-p)!} \left(-\frac{a}{b}\right)^m b^p Z_{p,m,n}^\alpha(a,b), \quad (98)$$

where

$$Z_{p,m,n}^\alpha(a,b) = \sum_{l,k=0}^{\infty} \frac{(p-n-m)_{k+l}(-m, -\alpha-m)_k(p+1, \alpha+1+p)_l}{(p-m+1, \alpha+p-m+1)_{k+l} k!l!} \left(-\frac{b}{a}\right)^k b^l \quad (99)$$

$$= F_{2;0;0}^{1;2;2} \left(\begin{matrix} p-n-m : -m, -\alpha-m; p+1, \alpha+p+1 \\ p-m+1, \alpha+p-m+1 : -; - \end{matrix} ; -\frac{b}{a}, b \right), \quad (100)$$

where $F_{2;0;0}^{1;2;2}$ is terminating Kampé de Fériet double hypergeometric series (4).

Proof. The terminating double hypergeometric Kampé de Fériet form (4) of the integral $\mathcal{L}_{p,m,n}^\alpha(a,b)$ in Corollary 7.5 follows from Theorem 7.4 by utilizing the identity (97) ($p \geq m$) with $r = l+l$ and (4). This completes the proof. \blacksquare

Theorem 7.6. *Let $p, m, n \in \mathbb{N}_0$, $a, b, > 0$, $\alpha > -1$. Then the contiguous relations for the integral of the product of three Laguerre polynomials, which two are scaled is the following:*

$$a(p+\alpha)\mathcal{L}_1^- - (m+\alpha)\mathcal{L}_2^- + (2m+\alpha+1)\mathcal{L} - a(2p+\alpha+1)\mathcal{L} - (m+1)\mathcal{L}_2^+ + a(p+1)\mathcal{L}_1^+ = 0, \quad (101)$$

$$b(p+\alpha)\mathcal{L}_1^- - (n+\alpha)\mathcal{L}_3^- + (2n+\alpha+1)\mathcal{L} - b(2p+\alpha+1)\mathcal{L} - (n+1)\mathcal{L}_3^+ + b(p+1)\mathcal{L}_1^+ = 0, \quad (102)$$

$$b(m+\alpha)\mathcal{L}_2^- - a(n+\alpha)\mathcal{L}_3^- + a(2n+\alpha+1)\mathcal{L} - b(2m+\alpha+1)\mathcal{L} - a(n+1)\mathcal{L}_3^+ + b(m+1)\mathcal{L}_2^+ = 0, \quad (103)$$

where

$$\mathcal{L} := \mathcal{L}(p, m, n; \alpha; a, b), \quad (104)$$

$$\mathcal{L}_1^\pm := \mathcal{L}(p \pm 1, m, n; \alpha; a, b), \quad (105)$$

$$\mathcal{L}_2^\pm := \mathcal{L}(p, m \pm 1, n; \alpha; a, b), \quad (106)$$

$$\mathcal{L}_3^\pm := \mathcal{L}(p, m, n \pm 1; \alpha; a, b). \quad (107)$$

Proof. Again Theorem 3.1 will be used with the coefficients of the three-term recurrence relations of Laguerre polynomials. However, because we are now working with scaled Laguerre polynomials, one of these coefficients is now modified to the following:

$$A_n = -\frac{a}{n+1}. \quad (108)$$

Where a is the scaling factor of the Laguerre polynomial $L_n^\alpha(ax)$. The other two coefficients seen in (82) are unchanged. This can be found by inserting the scaled Laguerre polynomial into the three-term recurrence relation and solving for the coefficients. ■

Acknowledgements

We would like to thank Roberto S. Costas-Santos for valuable discussions.

References

- [1] R. A. Askey, R. K. Beatson, T. S. Chihara, H. S. Cohl, C. F. Dunkl, C. Koutschan, S. Olver, Y. Xu, W. zu Castell, and W. Zudilin. Report from the Open Problems session at OPSFA13. *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA), Special Issue on Orthogonal Polynomials, Special Functions and Applications*, 12:Paper 071, 12, 2016.
- [2] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.1.5 of 2022-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [3] M. E. H. Ismail. *Classical and Quantum Orthogonal Polynomials in One Variable*, volume 98 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2009. With two chapters by Walter Van Assche, With a foreword by Richard A. Askey, Corrected reprint of the 2005 original.
- [4] M. E. H. Ismail, A. Kasraoui, and J. Zeng. Separation of variables and combinatorics of linearization coefficients of orthogonal polynomials. *Journal of Combinatorial Theory, Series A*, 120(3):561–599, 2013.
- [5] A. J. E. M. Janssen, J. J. M. Braat, and P. Dirksen. On the Computation of the Nijboer-Zernike Aberration Integrals at Arbitrary Defocus. *Journal of Modern Optics*, 51(5):687–703, 2004.
- [6] R. Koekoek, P. A. Lesky, and R. F. Swarttouw. *Hypergeometric orthogonal polynomials and their q -analogues*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2010. With a foreword by Tom H. Koornwinder.
- [7] M. Rahman. A nonnegative representation of the linearization coefficients of the product of Jacobi polynomials. *Canadian Journal of Mathematics. Journal Canadien de Mathématiques*, 33(4):915–928, 1981.
- [8] H. M. Srivastava and Per W. Karlsson. *Multiple Gaussian hypergeometric series*. Ellis Horwood Series: Mathematics and its Applications. Ellis Horwood Ltd., Chichester, 1985.