

Norm Inequalities Related to Heinz and Logarithmic Means

Guanghua Shi

School of Mathematical Sciences, Yangzhou University, Yangzhou, Jiangsu, China
sghkanting@163.com

June 14, 2022

Abstract

In this paper, we got some refinements of the norm inequalities related to the Heinz mean and logarithmic mean.

AMS classification: 47A63; 94A17

Keywords: Heinz mean, Logarithmic mean, Positive function, Unitarily invariant norm

1 Introduction

There are several means that interpolate between the geometric and arithmetic means. For instance, the Heinz mean $H_t(a, b)$, defined by

$$H_t(a, b) = \frac{a^{1-t}b^t + a^tb^{1-t}}{2} \quad \text{for } 0 \leq t \leq 1.$$

In 1993, Bhatia-Davis [2] obtained that if A, B and X are $n \times n$ matrices with A, B positive semidefinite, then for every unitarily invariant norm $\|\cdot\|$,

$$\left\| \left\| A^{\frac{1}{2}} X B^{\frac{1}{2}} \right\| \right\| \leq \frac{1}{2} \left\| \left\| A^{1-t} X B^t + A^t X B^{1-t} \right\| \right\| \leq \frac{1}{2} \left\| \left\| AX + XB \right\| \right\|. \quad (1.1)$$

The logarithmic mean $L(a, b)$, defined by

$$L(a, b) = \frac{a - b}{\log a - \log b} = \int_0^1 a^t b^{1-t} dt,$$

also interpolates the geometric and arithmetic means. In 1999, Hiai-Kosaki [5] proved the following inequality

$$\left\| \left\| A^{\frac{1}{2}} X B^{\frac{1}{2}} \right\| \right\| \leq \left\| \left\| \int_0^1 A^v X B^{1-v} dv \right\| \right\| \leq \frac{1}{2} \left\| \left\| AX + XB \right\| \right\|. \quad (1.2)$$

Moreover, in 2006, Drissi [4] proved that the following Heinz-logarithmic inequality

$$\|A^{1-t}XB^t + A^tXB^{1-t}\| \leq 2 \left\| \int_0^1 A^vXB^{1-v}dv \right\| \quad (1.3)$$

holds for $\frac{1}{4} \leq t \leq \frac{3}{4}$.

A complex-valued function φ on \mathbb{R} is said to be positive definite if the matrix $[\varphi(x_i - x_j)]$ is positive semidefinite for all choices of real numbers x_1, x_2, \dots, x_n , and $n = 1, 2, \dots$. Set $M(a, b)$ and $N(a, b)$ are two symmetric homogeneous means on $(0, \infty) \times (0, \infty)$. M is said to strongly dominate N , denoted by $M \ll N$, if and only if the matrix

$$\left[\frac{M(\lambda_i, \lambda_j)}{N(\lambda_i, \lambda_j)} \right]_{i,j=1,\dots,n}$$

is positive semidefinite for any size n and $\lambda_1, \dots, \lambda_n > 0$. Drissi [4] proved that for $a, b \geq 0$, $H_t(a, b) \ll L(a, b)$ if and only if $\frac{1}{4} \leq t \leq \frac{3}{4}$. In general, the inequality $M \ll N$ could be stronger than the Löwner's order inequality $M \leq N$.

For more operator or norm inequalities related to the Heinz mean and logarithmic mean we refer the readers to [7–9] and the references therein.

2 Main results

Lemma 2.1. *For $\sinh x$ and $\cosh x$, we have*

(i) *If $|\beta| > |\alpha| > 0$, then the function $\frac{\cosh \alpha x}{\cosh \beta x}$ is positive definite.*

(ii) *If $|\beta| > |\alpha| > 0$ with α, β the same sign, then $\frac{\sinh \alpha x}{\sinh \beta x}$ is positive definite.*

(iii) *If $\beta > 0$ and $|\alpha| < \beta/2$, then $\frac{\beta x \cosh \alpha x}{\sinh \beta x}$ is positive definite.*

Proof. We follow a similar argument as in Chapter 5 of [1]. From the product representations in p. 147-148 of [1],

$$\frac{\sinh x}{x} = \prod_{k=1}^{\infty} \left(1 + \frac{x^2}{k^2\pi^2} \right), \quad \cosh x = \prod_{k=0}^{\infty} \left(1 + \frac{4x^2}{(2k+1)^2\pi^2} \right), \quad (2.1)$$

we have

$$\frac{\sinh \alpha x}{\sinh \beta x} = \frac{\alpha}{\beta} \prod_{k=1}^{\infty} \frac{1 + \alpha^2 x^2 / k^2 \pi^2}{1 + \beta^2 x^2 / k^2 \pi^2}, \quad \frac{\cosh \alpha x}{\cosh \beta x} = \prod_{k=0}^{\infty} \frac{1 + 4\alpha^2 x^2 / (2k+1)^2 \pi^2}{1 + 4\beta^2 x^2 / (2k+1)^2 \pi^2}. \quad (2.2)$$

Each factor in the product is of the form

$$\frac{1 + a^2 x^2}{1 + b^2 x^2} = \frac{a^2}{b^2} + \frac{1 - a^2/b^2}{1 + b^2 x^2}, \quad b^2 > a^2.$$

Since $1/(1+b^2x^2)$ is positive definite [1, 5.2.7], it follows that the functions $\cosh \alpha x / \cosh \beta x$ is positive definite for $|\beta| > |\alpha| > 0$, and $\sinh \alpha x / \sinh \beta x$ is positive definite for $|\beta| > |\alpha| > 0$ with α, β the same sign.

Since

$$\frac{\beta x \cosh \alpha x}{\sinh \beta x} = \frac{\frac{\beta}{2} x}{\sinh \frac{\beta}{2} x} \cdot \frac{\cosh \alpha x}{\cosh \frac{\beta}{2} x}$$

and $x/\sinh x$ is positive definite [1, 5.2.9], it follows from the above argument that when $\frac{\beta}{2} > |\alpha|$ the function $\frac{\beta x \cosh \alpha x}{\sinh \beta x}$ is positive definite. \square

Now, we define

$$L_s(a, b) = \frac{a^{1-s} b^s - a^s b^{1-s}}{(1-2s)(\log a - \log b)} = \frac{1}{1-2s} \int_s^{1-s} a^v b^{1-v} dv$$

for $a, b > 0$, and $0 \leq s < \frac{1}{2}$. When $s = 0$, it is the logarithmic mean. So we can call it the generalized logarithmic mean. And we also have $\lim_{s \rightarrow \frac{1}{2}} L_s(a, b) = a^{\frac{1}{2}} b^{\frac{1}{2}}$.

Theorem 2.2. *For Heinz mean and the generalized logarithmic mean, we have*

(i) *If $0 \leq s < 1/2$, and $|1 - 2t| < \frac{1-2s}{2}$, then $H_t(a, b) \ll L_s(a, b)$.*

(ii) *If $|1 - 2t| < |1 - 2s|$, then $H_t(a, b) \ll H_s(a, b)$.*

(iii) *If $0 \leq s < t < 1/2$, or $1 \geq s > t > 1/2$, then $L_t(a, b) \ll L_s(a, b)$.*

Proof. By definition, $H_t(a, b) \ll L_s(a, b)$ if

$$[y_{i,j}] = \left[\frac{H_t(\lambda_i, \lambda_j)}{L_s(\lambda_i, \lambda_j)} \right]_{i,j=1,\dots,n}$$

is positive semidefinite. Set $\lambda_i = e^{x_i}$ and $\lambda_j = e^{x_j}$, with $x_i, x_j \in \mathbb{R}$. Then

$$y_{i,j} = (1-2s) \frac{\left(\frac{x_i-x_j}{2}\right) \left(e^{(1-2t)\frac{x_i-x_j}{2}} + e^{(1-2t)\frac{x_j-x_i}{2}} \right)}{e^{(1-2s)\frac{x_i-x_j}{2}} - e^{(1-2s)\frac{x_j-x_i}{2}}}$$

Thus the matrix $[y_{i,j}]$ is congruent to one with entries

$$\frac{\beta \left(\frac{x_i-x_j}{2}\right) \cosh\left(\alpha \left(\frac{x_i-x_j}{2}\right)\right)}{\sinh\left(\beta \left(\frac{x_i-x_j}{2}\right)\right)},$$

where $\alpha = 1 - 2t$, $\beta = 1 - 2s$. Hence $[y_{i,j}]$ is positive semidefinite if and only if the function $\frac{\beta x \cosh \alpha x}{\sinh \beta x}$ is positive definite, which by lemma 2.1 is correct.

Similarly, we have $H_t(a, b) \ll H_s(a, b)$ if $\frac{\cosh \alpha x}{\cosh \beta x}$ is positive definite, and $L_t(a, b) \ll L_s(a, b)$ if $\frac{\sinh \alpha x}{\sinh \beta x}$ is positive definite. \square

Theorem 2.3. *Let A, B be any positive matrices. Then for any matrix X and for $0 \leq s < 1/2$ and $|1 - 2t| < (1 - 2s)/2$, we have*

$$\| \| A^{1-t} X B^t + A^t X B^{1-t} \| \| \leq \frac{2}{1-2s} \left\| \int_s^{1-s} A^v X B^{1-v} dv \right\|. \quad (2.3)$$

Proof. Firstly, we assume $A = B$. Since $\|\cdot\|$ is unitarily invariant, we may suppose A is diagonal with entries $\lambda_1, \dots, \lambda_n$. Then we have

$$A^{1-t}XA^t + A^tXA^{1-t} = Y \circ \left(\int_s^{1-s} A^v X A^{1-v} dv \right),$$

where Y is the matrix with entries

$$y_{i,j} = \frac{\lambda_i^t \lambda_j^{1-t} + \lambda_i^{1-t} \lambda_j^t}{\frac{\lambda_i^{1-s} \lambda_j^s - \lambda_i^s \lambda_j^{1-s}}{\log \lambda_i - \log \lambda_j}} = \frac{2H_t(\lambda_i, \lambda_j)}{(1-2s)L_s(\lambda_i, \lambda_j)}$$

A well-known result related to the Schur multiplier norm [6, Theorem 5.5.18, 5.5.19] says that if Y is any positive semidefinite matrix, then for all matrix X ,

$$\|Y \circ X\| \leq \max_i y_{ii} \|X\| \quad (2.4)$$

for every unitarily invariant norm. By Theorem 2.2, Y is positive semidefinite. Applying (2.4), we have

$$\|A^{1-t}XA^t + A^tXA^{1-t}\| \leq \frac{2}{1-2s} \left\| \int_s^{1-s} A^v X A^{1-v} dv \right\|. \quad (2.5)$$

Now replacing A and X in the inequality (2.5) by the 2 by 2 matrices $\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$ and $\begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}$. This gives the desired inequality (2.3). \square

When $s = 0$, we get Drissi's result (1.3). Moreover, when $s = 0, t = 1/2$, we get the first inequality of (1.2).

Theorem 2.4. *Let A, B be any positive matrices. Then for any matrix X and for $0 \leq s < 1/2$, we have*

$$\left\| \int_s^{1-s} A^v X B^{1-v} dv \right\| \leq \frac{1-2s}{2} \|A^{1-s} X B^s + A^s X B^{1-s}\|. \quad (2.6)$$

Proof. Suppose A is diagonal with entries $\lambda_1, \dots, \lambda_n$. Then we have

$$\int_s^{1-s} A^v X A^{1-v} dv = Y \circ (A^{1-s} X A^s + A^s X A^{1-s}),$$

where Y is the matrix with entries

$$y_{i,j} = \frac{\frac{\lambda_i^{1-s} \lambda_j^s - \lambda_i^s \lambda_j^{1-s}}{\log \lambda_i - \log \lambda_j}}{\lambda_i^s \lambda_j^{1-s} + \lambda_i^{1-s} \lambda_j^s} = \frac{(1-2s)L_s(\lambda_i, \lambda_j)}{2H_s(\lambda_i, \lambda_j)}.$$

By a similar argument as in the proof of Theorem 2.2, we know that Y is positive definite if and only if

$$\frac{\beta}{2} \frac{\sinh \beta x}{\beta x \cosh \beta x} = \frac{\beta}{2} \frac{\tanh \beta x}{\beta x}$$

is positive definite for $\beta = 1 - 2s > 0$, because $\tanh x/x$ is positive definite (see Bhatia [1, 5.2.11]). Thus Applying (2.4) we have

$$\left\| \int_s^{1-s} A^v X A^{1-v} dv \right\| \leq \frac{1-2s}{2} \left\| A^{1-s} X A^s + A^s X A^{1-s} \right\|. \quad (2.7)$$

Hence the desired result follows. \square

When $s = 0$, we get the second inequality of (1.2).

Theorem 2.5. *Let A, B be any positive matrices. Then for any matrix X and for $|1-2t| < |1-2s|$ with $1-2t, 1-2s$ the same sign, we have*

$$\left\| A^{1-t} X B^t - A^t X B^{1-t} \right\| \leq \left| \frac{1-2t}{1-2s} \right| \left\| A^{1-s} X B^s - A^s X B^{1-s} \right\|. \quad (2.8)$$

Proof. Suppose A is diagonal with entries $\lambda_1, \dots, \lambda_n$. Then we have

$$A^{1-t} X A^t - A^t X A^{1-t} = Y \circ (A^{1-s} X A^s - A^s X A^{1-s}),$$

where Y is the matrix with entries

$$y_{i,j} = \frac{\lambda_i^{1-t} \lambda_j^t - \lambda_i^t \lambda_j^{1-t}}{\lambda_i^{1-s} \lambda_j^s - \lambda_i^s \lambda_j^{1-s}}$$

Put $\lambda_i = e^{x_i}$ and $\lambda_j = e^{x_j}$, with $x_i, x_j \in \mathbb{R}$. Then Y is congruent to the matrix with entries

$$\frac{\sinh(\alpha \frac{x_i - x_j}{2})}{\sinh(\beta \frac{x_i - x_j}{2})}$$

where $\alpha = 1 - 2t, \beta = 1 - 2s$. Since $(\sinh \alpha x)/(\sinh \beta x)$ is positive for $|\beta| > |\alpha| > 0$ with α, β the same sign, it follows that Y is positive definite. Applying the inequality (2.4), we have

$$\left\| A^{1-t} X A^t - A^t X A^{1-t} \right\| \leq \left| \frac{1-2t}{1-2s} \right| \left\| A^{1-s} X A^s - A^s X A^{1-s} \right\|. \quad (2.9)$$

Hence the desired result follows. \square

Set $s = 0$ and $s = 1$, and combine the conclusions, we have the following inequality proved by Bhatia-Davis [3],

Corollary 2.6. *Let A, B be any positive matrices. Then for any matrix X and for $0 \leq t \leq 1$ we have*

$$\left\| A^{1-t} X B^t - A^t X B^{1-t} \right\| \leq |1-2t| \left\| AX - XB \right\|. \quad (2.10)$$

Acknowledgments

The author acknowledges support from National Natural Science Foundation of China, Grant No: 12001477, and the Natural Science Foundation of Jiangsu Province for Youth, Grant No: BK20190874.

References

- [1] R. Bhatia, Positive Definite Matrices. Princeton University Press, Princeton (2007).
- [2] R. Bhatia and C. Davis, More matrix forms of the arithmetic-geometric mean inequality, *SIAM J. Matrix Anal. Appl.*, 14 (1993), 132-136.
- [3] R. Bhatia and C. Davis, A Cauchy-Schwarz inequality for operators with applications, *Linear Algebra Appl.*, 223/224 (1995), 119-129.
- [4] D. Drissi, Sharp inequalities for some operator means, *SIAM J. Matrix Anal. Appl.*, 28(3) (2006), 822-828.
- [5] F. Hiai and H. Kosaki, Means for matrices and comparison of their norms, *Indiana Univ. Math. J.*, 48 (1999), 899-936.
- [6] R. A. Horn and C. R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, UK, 1991.
- [7] Y. Kapil and M. Singh, Contractive maps on operator ideals and norm inequalities. *Linear Algebra Appl.*, 459 (2014), 475-492.
- [8] H. Kosaki, Arithmetic-geometric mean and related inequalities for operators, *J. Funct. Anal.*, 156 (1998), 429-451.
- [9] X. Zhan, Inequalities for unitarily invariant norms, *SIAM J. Matrix Anal. Appl.*, 20 (1999), 466-470.