

THE DEGENERATE PRINCIPAL SERIES REPRESENTATIONS OF EXCEPTIONAL GROUPS OF TYPE E_8 OVER p -ADIC FIELDS

HEZI HALAWI¹ AND AVNER SEGAL²

ABSTRACT. In this paper, we study the reducibility of degenerate principal series of the simple, simply-connected exceptional group of type E_8 . Furthermore, we calculate the maximal semi-simple subrepresentation and quotient of these representations for almost all cases.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. The Main Theorem	12
4. Exceptional Cases	21
Appendix A. Branching Rules Database	38
References	45

1. Introduction

This paper is the final part in our project of studying the degenerate principal series of exceptional groups of type E_n . This paper is about E_8 , which, as often noted by David Kazhdan, is the smallest, split, simple, simply-connected, adjoint and simply laced group. In fact, this essentially completes the study of degenerate principal series of simple p -adic groups up to isogeny.

More precisely, let F be a non-Archimedean local field and let G denote the split simple group of type E_8 . For a maximal parabolic subgroup P of G with a Levi subgroup M and a 1-dimensional representation Ω of M , we consider the following two questions:

- Is the normalized parabolic induction $Ind_P^G(\Omega)$ irreducible?

2010 *Mathematics Subject Classification.* 22E50, 20G41, 20G05.

- If $\text{Ind}_P^G(\Omega)$ is reducible, what is the length of its maximal semi-simple subrepresentation and quotient?

We completely answer the first one in Theorem 3.1, and almost completely answer the second. In fact, there are only two pairs (P, Ω) (out of hundreds of cases) in which we were only able to show that the maximal semi-simple subrepresentation is of length at most 2. In both of these cases, we show that the irreducible spherical subquotient is a subrepresentation and describe the other possible irreducible subrepresentation in terms of its Langlands data. Further, we describe a decisive test to determine the length of the maximal semi-simple quotient for each of these cases, which would hopefully could be realized when stronger computing machines would be more commonly available. These two cases are detailed in Subsection 4.2.

In order to answer the above questions, we use the algorithm described in [HSa, Section 3] and [HSb, Section 3]. This provides an answer to both questions for almost all pairs (P, Ω) . For the remaining cases, not determined by the algorithm, further study is performed in Section 4. This project uses a script implemented in the Sagemath environment [TheYY].

The study of local degenerate principles series is useful for various reasons. One of which is the study of the degenerate residual spectrum of the adelic group. The study of the degenerate residual spectrum of the simple group of type F_4 is the topic of the first author's PhD dissertation and relies on the study of the local degenerate principal series of F_4 , performed in [CJ10]. The study of degenerate residual representations of $Spin_8$, with P being the Heisenberg parabolic subgroup, was performed by the second author in [Seg18, Seg19] and used to study exceptional θ -lifts in [GS].

The study of the degenerate residual spectrum of the adelic groups of type E_n is a work in progress as a joint project, [HSc], of both authors.

This paper is structured as follows:

- In Section 2, we recall basic notations and properties from representation theory of p -adic groups, we recall the algorithm described in [HSa] and [HSb] and we recall basic data on the exceptional group of type E_8 .
- In Section 3 we state our main theorem, Theorem 3.1, and list all cases which can be resolved using our algorithm. This algorithm constitutes of reducibility and irreducibility test as well as some test to check if the representation admits a unique irreducible subrepresentation.

- In Section 4 we go over the exceptional cases which could not have been fully resolved by our algorithm. We resolve most of these cases completely and make some progress towards the resolution of the remaining two cases.

Finally, we wish to address a question which was broached to us following [HSa] and [HSb]. The reducibility of a non-unitary degenerate principal series representation can be determined by the local Shahidi coefficients which would seem to make our algorithm obsolete. However, the algorithm, presented in Subsection 2.2, is useful for various other reasons such as:

- It allows us to determine the reducibility for unitary cases too.
- While local Shahidi coefficients can inform us regarding the reducibility of non-unitary degenerate principal series, the output of the reducibility test is also useful when studying global phenomena such as the Siegel-Weil identity (and indeed this data is used in [HSc]).
- Data from the reducibility and irreducibility tests is useful for studying the structure of reducible degenerate principal series and in particular for studying its socle and cosocle.

We also wish to point out that comparing the lists of non-unitary reducible degenerate principal series determined by our method with that given by Shahidi's method, was useful for debugging purposes. Indeed, as in the E_6 and E_7 case, our algorithm was decisive for all non-unitary cases. The only cases where the algorithm was unable to determine the reducibility of the degenerate principal series were unitary ones.

Acknowledgments. The second author was partially supported by grants 421/17 and 259/14 from the Israel Science Foundation as well as by the Junior Researcher Grant of Shamoon College of Engineering (SCE).

2. Preliminaries

This section has three parts. In the first part, we fix notations for this paper. This part is organized as an enumerated list in order to make the look up of notations easier.

In the second part we recall the algorithm which we implemented for the study of the degenerate principal series of E_8 .

Since the algorithm has been described in detail in [HSa, Section 3] and [HSb, Section 3], we skip much of the details and describe only the broad strokes of it, while keeping all details which are required for Section 3 and Section 4.

In the third part of this section we introduce the split group of type E_8 .

2.1. Groups, Characters and Representations

In this subsection, we fix notations and recall basic facts about the groups, characters and representations involved in this paper. For a more detailed discussion, the reader is encouraged to consider [HSa, Section 2] and [HSb, Section 2].

2.1.1. Groups

- (1) Let F be a non-Archimedean local field with norm $|\cdot|$.
- (2) Let G denote the F -points of a split simply-connected reductive group.
- (3) Let T be a maximal split torus of G .
- (4) Let B be a Borel subgroup of G such that $T \subset B$.
- (5) Let Φ_G denote the roots of G with respect to T and let $\Phi_G^+ \subset \Phi_G$ denote the positive roots of G with respect to B .
- (6) Let $\Delta_G = \{\alpha_1, \dots, \alpha_n\}$ be the set of simple roots of Φ_G with respect to B .
- (7) Let $n = |\Delta_G| = \dim_F(T)$ denote the rank of G .
- (8) Let $\Phi_G^\vee = \{\alpha^\vee \mid \alpha \in \Phi_G\}$ denote the set of coroots of G with respect to T .
- (9) We use $\langle \cdot, \cdot \rangle$ to denote the usual pairing between characters and co-characters of T .
- (10) Let $\omega_{\alpha_1}, \dots, \omega_{\alpha_n}$ denote the fundamental weights of T which satisfy

$$\langle \omega_{\alpha_i}, \alpha_j^\vee \rangle = \delta_{i,j}.$$

- (11) Let $W = \langle s_i \mid 1 \leq i \leq n \rangle$ denote the Weyl group of G with respect to T , generated by the simple reflections s_i associated with the simple roots α_i .
- (12) For $\Theta \subset \Delta_G$, let $P_\Theta = \langle B, s_i \mid \alpha_i \in \Theta \rangle = M_\Theta \cdot U_\Theta$ be the standard parabolic subgroup of G associated to Θ . We denote its Levi subgroup by M_Θ and its unipotent radical by U_Θ .
- (13) Let Φ_{M_Θ} , $\Phi_{M_\Theta}^+$ and Δ_{M_Θ} denote the roots, positive roots and simple roots of M_Θ with respect to T and $B \cap M_\Theta$ respectively.
- (14) Let $W_{M_\Theta} = \langle s_i \mid \alpha_i \in \Theta \rangle$ denote the Weyl group of M_Θ with respect to T .

- (15) Let $P_i = P_{\Delta_G \setminus \{\alpha_i\}}$ and $M_i = M_{\Delta_G \setminus \{\alpha_i\}}$ denote a maximal (proper) standard parabolic subgroup of G and its Levi subgroup.
- (16) For maximal standard Levi subgroups M_i and M_j of G , we write $M_{i,j} = M_i \cap M_j$.
- (17) We denote the rank 1 Levi subgroups by $L_i = M_{\{\alpha_i\}}$.

2.1.2. Characters

- (1) Let $\mathbf{X}(G) = \{\Omega : G \rightarrow \mathbb{C}^\times\}$ denote the complex manifold of continuous characters of G , we use additive notations for this group, that is

$$(\Omega_1 + \Omega_2)(g) = \Omega_1(g) \cdot \Omega_2(g).$$

We usually use the letter Ω to denote elements in $\mathbf{X}(M)$, for a non-minimal Levi subgroup M of G , while using λ to denote an element of $\mathbf{X}(T)$. Also, note that W acts on $\mathbf{X}(T)$ via its action on T .

- (2) We denote the set of unramified elements $\Omega \in \mathbf{X}(T)$ by $\mathbf{X}^{un}(T)$.
- (3) Let $\mathbf{1}_G \in \mathbf{X}(G)$ denote the trivial character of G .
- (4) We say that $\chi \in \mathbf{X}(T)$ has finite order if there exists $k \in \mathbb{N}$ such that $\chi^k = \mathbf{1}_T$. The order of χ , denoted by $ord(\chi)$, is the minimal $k \in \mathbb{N}$ such that $\chi^k = \mathbf{1}_T$.

In particular, every element $\Omega \in \mathbf{X}(F^\times)$ can be written as $\Omega = s + \chi$, where $s \in \mathbb{C}$ and χ is of finite order. Namely,

$$\Omega(x) = \chi(x) |x|^s \quad \forall x \in F^\times.$$

It holds that $\mathbf{X}^{un}(F^\times)$ can be described by all characters of the forms $|x|^s$ for some $s \in \mathbb{C}$.

- (5) We write $\text{Re}(\Omega)$ for the character

$$\text{Re}(\Omega)(x) = |x|^{\text{Re}(s)}.$$

- (6) We say that $\lambda \in \mathbf{X}(T)$ is **anti-dominant** if

$$\text{Re}(\langle \lambda, \alpha_i^\vee \rangle) \leq 0 \quad \forall 1 \leq i \leq n.$$

Note that every W_G -orbit in $\mathbf{X}(T)$ contains at least one anti-dominant element and all anti-dominant elements in the same W_G -orbit have an equal real part. As a convention, we denote an anti-dominant element by $\lambda_{a.d.}$.

- (7) Let $\Omega_{i,s,\chi}$ denote the character of a maximal Levi subgroup M_i of G associated with $(s + \chi) \circ \omega_{\alpha_i}$, where $s \in \mathbb{C}$ and $\chi \in \mathbf{X}(F^\times)$ is of finite order. Note that if G is simple, then any element in $\mathbf{X}(M_i)$ can be written this way.

2.1.3. Representations

- (1) Let $Rep(G)$ denote the category of admissible representations of G .
(2) As above, $\mathbf{1}_G$ denotes the trivial representation of G .
(3) Let $i_M^G : Rep(M) \rightarrow Rep(G)$ and $r_M^G : Rep(G) \rightarrow Rep(M)$ denote the functors of normalized parabolic induction and Jacquet functor, adjunct by the Frobenius reciprocity:

$$(2.1) \quad \text{Hom}_G(\pi, i_M^G \sigma) \cong \text{Hom}_M(r_M^G \pi, \sigma).$$

- (4) For $\pi, \sigma \in Rep(G)$, such that σ is irreducible, let $mult(\sigma, \pi)$ denote the multiplicity of σ in the Jordan-Hölder series of π .
(5) Let $\mathfrak{R}(G)$ denote the Grothendieck ring of $Rep(G)$ and let $[\pi]$ denote the image of $\pi \in Rep(G)$ in $\mathfrak{R}(G)$. Recall that $\mathfrak{R}(G)$ admits a partial order such that $\pi_1 \leq \pi_2$ if $mult(\sigma, \pi_1) \leq mult(\sigma, \pi_2)$ for every irreducible $\sigma \in Rep(G)$.
(6) We remind the reader that, for Levi subgroups L and M of G and $\sigma \in Rep(M)$, the composition $[r_L^G i_M^G \sigma]$ is given by the *geometric lemma* ([BZ77, Lemma 2.12], [Cas74, Theorem 6.3.6]):

$$(2.2) \quad [r_L^G i_M^G \sigma] = \sum_{w \in W^{M,L}} [i_{L'}^L \circ w \circ r_{M'}^M \sigma],$$

where:

- $W^{M,L} = \{w \in W \mid w(\Phi_M^+) \subseteq \Phi_G^+, w^{-1}(\Phi_L^+) \subseteq \Phi_G^+\}$ is the set of shortest representatives in W of the double coset space $W_L \backslash W_G / W_M$.
 - For $w \in W^{M,L}$ we write $M' = M \cap w^{-1}Lw$ and $L' = wMw^{-1} \cap L$.
- (7) For $\pi \in Rep(G)$, we write $[r_T^G \pi] = \sum_{i=1}^l n_i \times [\lambda_i]$ for certain $\lambda_i \in \mathbf{X}(T)$ such that $mult(\lambda_i, \pi) = n_i > 0$. Since $\dim_{\mathbb{C}}(r_T^G \pi)$ is finite, there are only finitely many such λ_i . We call such λ_i **the exponents of π** .
(8) The representations $\pi = i_{M_i}^G(\Omega_{M_i,s,\chi})$ are called **degenerate principal series**. The exponent $\lambda_0 = r_T^{M_i}(\Omega_{M_i,s,\chi})$ is called the **initial exponent** of π .

- (9) We say that $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$ is **regular** if $\text{Stab}_W(\lambda_0) = \{1\}$, where $\lambda_0 = r_T^{M_i}(\Omega_{M_i, s, \chi})$.
- (10) Let w_0 denote the longest element in $W^{M_i, T}$. It holds that $w_0 \cdot (i_{M_i}^G(\Omega_{M_i, s, \chi})) = i_{M_j}^G(\Omega_{M_j, -s, \bar{\chi}})$, where $\bar{\chi}$ is the complex conjugate of χ and $M_j = w_0 M_i w_0^{-1}$. We note that $M_j = M_i$, except when G is of type A_n , D_{2n+1} or E_6 . We call $i_{M_j}^G(\Omega_{M_j, -s, \bar{\chi}})$ the **invert representation** of $i_{M_i}^G(\Omega_{M_i, s, \chi})$. We note that the invert representation has the same irreducible constituents but in an "inverted order". That is, $i_{M_j}^G(\Omega_{M_j, -s, \bar{\chi}})$ admits a Jordan-Hölder series whose irreducible quotients appear an inverted order than that of $i_{M_i}^G(\Omega_{M_i, s, \chi})$. When $\chi = \mathbf{1}$, the invert was defined in [Jan95, Remark 2.2.5] as a variation of the Iwahori-Matsumoto involution.

2.2. The Algorithm

We now recall parts of the algorithm used by us to study the degenerate principal series $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$. For a more detailed account, the reader is encouraged to consider [HSa, Section 3] and [HSb, Section 3].

We identify the data defining π by a triple of numbers $[i, s, \text{ord}(\chi)]$. In particular, the reducibility and lengths of the maximal semi-simple subrepresentation and quotient depend only on this triple (and are uniform among χ s with the same order).

As explained in [HSa, Remark 3.1], we may assume, without loss of generality, that $s \in \mathbb{R}$. Furthermore, it is enough to consider only the cases where $s \leq 0$ since the invert representation of π admits a Jordan-Hölder series with same irreducible quotients appearing but in inverted order.

The algorithm has the following parts:

- (1) Determine all non-regular such π - there is a finite number of such cases!
- (2) Determine all reducible regular π - there are only finitely many such cases!
- (3) Apply reducibility tests to non-regular π (these may be inconclusive).
- (4) Apply an irreducibility test to non-regular π (this too may be inconclusive). This test uses the so called *branching rule calculation* introduced in Subsection 2.2.2.
- (5) Determine if π admits a unique irreducible subrepresentation. As will be explained in Subsection 2.2.3, for $s < 0$, π admits a unique irreducible quotient.

The treatment of the regular cases is standard and is outlined in [HSa, Subsection 3.1]. In what follows, we describe tools used mostly for dealing with the non-regular case.

2.2.1. Reducibility Tests

In order to prove that a representation $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$ is reducible, we provide another $\pi' \in \text{Rep}(G)$ such that π and π' share a common irreducible constituent, while $\pi \neq \pi'$. Usually, it is enough to consider $\pi' = i_{M_j}^G(\Omega_{M_j, t, \chi^l})$, where l is a totative¹ of $\text{ord}(\chi)$. In other cases, one takes $\pi' = i_{M_{j_1, j_2}}^G \Omega_{s_1, s_2, \chi, k_1, k_2}$, where $\Omega_{s_1, s_2, \chi, k_1, k_2} \in \mathbf{X}(M_{j_1, j_2})$ is associated with

$$(2.3) \quad (s_1 + \chi^{k_1}) \circ \omega_{\alpha_{j_1}} + (s_2 + \chi^{k_2}) \circ \omega_{\alpha_{j_2}},$$

such that at least one of k_1 and k_2 is a totative of $\text{ord}(\chi)$.

In particular, one checks that:

- π and π' share a common anti-dominant exponent and
- $r_T^G \pi \not\leq r_T^G \pi'$.

Which guarantees that π is reducible and shares a common irreducible subquotient with π' (possibly π' itself). This is done via Tadić's criterion (See [Tad98, Lemma 3.1] and [HSb, Section 3B] for more information).

2.2.2. Irreducibility Tests - Branching Rule Calculations

We test the irreducibility of $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$ using a method we call *branching rule calculations*. This method is also used, in some cases, to determine the number of irreducible subrepresentations of π .

Let

$$\mathcal{S} = \{f : \mathbf{X}(T) \rightarrow \mathbb{N} \mid f \text{ has a finite support}\}.$$

and note that \mathcal{S} is endowed with a natural partial order. For any $\sigma \in \text{Rep}(G)$, let $f_\sigma \in \mathcal{S}$ be defined by

$$f_\sigma(\lambda) = \text{mult}(\lambda, r_T^G \sigma).$$

In particular, for $\sigma = \pi$, f_π can be calculated directly using Equation (2.2).

A sequence of functions $\{f_j\}_{j=0}^k$ is called a **unital σ -dominated sequence** if it satisfies the condition $f_0 \leq f_1 \leq \dots \leq f_k \leq f_\sigma$ and $f_0 = \delta_\lambda$ for some $\lambda \leq r_T^G \sigma$. Here δ_λ denotes the

¹That is, an integer $0 < l \leq \text{ord}(\chi)$ which is coprime with $\text{ord}(\chi)$

Kronecker delta function given by

$$\delta_\lambda(\lambda') = \begin{cases} 1, & \lambda' = \lambda \\ 0, & \lambda' \neq \lambda \end{cases}.$$

For a given $\lambda \leq r_T^G \sigma$, we describe a recipe of constructing a σ -dominated sequence $\{f_j\}_{j=0}^k$ such that $f_0 = \delta_\lambda$.

The construction of the sequence $\{f_j\}_{j=0}^k$ is done using the following recursive process:

- (1) Set $f_0 = \delta_\lambda$.
- (2) Assume that $f_0 < \dots < f_l$ were determined. We choose $\lambda' \in \text{supp}(f_l)$ and a Levi subgroup L of G such that M admits a unique irreducible representation τ such that $\lambda' \leq r_T^L \tau$.
- (3) For a choice of (λ', L, τ) as above, for any $\mu \in \mathbf{X}(T)$, let

$$g(\mu) = \max \left\{ f_l(\mu), \left\lceil \frac{f_l(\lambda')}{\text{mult}(\lambda', r_T^L \tau)} \right\rceil \cdot \text{mult}(\mu, r_T^L \tau) \right\}.$$

- (4) If $f_l < g_{\lambda', L, \tau}$ for some choice of (λ', L, τ) , set $f_{l+1} = g$ and go back to step (2). Otherwise, we take $k = l$ and the process terminates.

This process is later referred to as a **branching rule calculations**.

Remark 2.1. The uniqueness condition on τ in item (2) can be slightly relaxed. In fact, one actually only needs that $[r_T^L \tau]$ be unique. Also, we note that one can apply this process to any function $f_0 \leq f_\sigma$ and not only delta functions, in this case we call the resulting sequence a **non-unital σ -dominated sequence**.

A more detailed account of this recipe can be found in [HSa, Subsection 3.3] and [HSb, Subsection 3.3]. Furthermore, a pair of explicit examples of such calculations, in the case of $SL_4(F)$, can be found in [HSb, Appendix C] and an explicit example of such calculations in the case of the group E_6 can be found in [HSa, Appendix B].

The calculation of the sequence $f_0 \leq f_1 \leq \dots \leq f_k \leq f_\sigma$ as above relies on a database of irreducible representations of standard Levi subgroups of G , the database used by us can be found in Appendix A.

The terminal function f_k in this sequence provides a lower bound to the multiplicities of exponents appearing in $r_T^G \sigma$. Throughout this manuscript, we refer to this process as a *branching rule calculation* (associated to λ).

In particular, for $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$, let $\lambda_{a.d.}$ denote an anti-dominant exponent of π and let π_0 denote an irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$. We construct a sequence $f_0 \leq f_1 \leq \dots \leq f_k \leq f_{\pi_0}$ as above. If it holds that $f = f_{\pi}$, then it follows that $r_T^G \pi = r_T^G \pi_0$ and since all subquotients of π have their cuspidal support along B , it follows that $\pi = \pi_0$. In particular, π is irreducible.

In fact, in order to prove that π is irreducible, it is enough to show that

$$\begin{aligned} \text{mult}(\lambda_0, r_T^G \pi_0) &= \text{mult}(\lambda_0, r_T^G \pi) \\ \text{mult}(\lambda_1, r_T^G \pi_0) &= \text{mult}(\lambda_1, r_T^G \pi), \end{aligned}$$

where λ_0 is the initial exponent of π and λ_1 is its terminal exponent, namely the initial exponent of the invert of π . This would imply that π is both its unique irreducible subrepresentation and quotient and hence it is irreducible.

2.2.3. Irreducible Subrepresentations

Bellow, we describe a few methods that can be realized as part of the algorithm, in order to determine the length of the maximal semi-simple subrepresentation of the degenerate principal series representation $\pi = i_{M_i, s, \chi}^G$.

- (1) If $\text{mult}(\lambda_0, r_T^G \pi) = 1$, where λ_0 is the initial exponent of π , then π admits a unique irreducible subrepresentation.
- (2) Let $\lambda_{a.d.}$ denote an anti-dominant exponent of π and let π_0 denote an irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$. If

$$(2.4) \quad \text{mult}(\lambda_0, r_T^G \pi_0) = \text{mult}(\lambda_0, r_T^G \pi),$$

then π_0 is the unique irreducible subrepresentation of π .

The condition in Equation (2.4) can, in many cases, be verified using branching rule calculations. Also, this argument seems to be relevant only when $s \leq 0$, since otherwise π would be irreducible (and thus would not be the subject of investigation for this part of the algorithm).

- (3) Given a unitary degenerate principal series $\pi = i_{M_i}^G(\Omega_{M_i, 0, \chi})$, then π is semi-simple of length at most 2 by [Ban02, Lemma 5.2]. In particular, if it is reducible, it is of length 2.

We note that, in a number of cases, the length of the semi-simple subrepresentation cannot be determined by these methods. This can happen for cases where the length

of the semi-simple subrepresentation can be either 1 or more. These cases are listed in Theorem 3.1 and are dealt with in Section 4 using various other methods.

Remark 2.2. For $s > 0$, the multiplicity of the initial exponent always appears in $r_T^G \pi$ with multiplicity 1 and thus π admits a unique irreducible subrepresentation. Conversely, this is why π always admits a unique irreducible quotient when $s < 0$.


2.3. The Exceptional Group of Type E_8

Let G be the split, semi-simple, simply-connected group of type E_8 . In this subsection we describe the structure of G . We fix a Borel subgroup B and a maximal split torus $T \subset B$ with notations as in Subsection 2.1. The set of roots, Φ_G , contains 240 roots. The group G is generated by symbols

$$\{x_\alpha(r) : \alpha \in \Phi_G, r \in F\}$$

subject to the Chevalley relations as in [Ste68, Section 6].

We label the simple roots Δ_G and the Dynkin diagram of G using the Bourbaki labelling:



Recall that for $\Theta \subset \Delta_G$ we denote by M_Θ the standard Levi subgroup of G such that $\Delta_M = \Theta$. We let M_i denote the Levi subgroup of the maximal parabolic subgroup $P_i = P_{\Delta_G \setminus \{\alpha_i\}}$.

Lemma 2.3. *Under these notations, it holds that:*

- (1) $M_1 \cong \left\{ g \in GSpin_{14}(F) \mid \det(g) \in (F^\times)^2 \right\}$.
- (2) $M_2 \cong GL_8(F)$.
- (3) $M_3 \cong \{(g_1, g_2) \in GL_2(F) \times GL_7(F) \mid \det(g_1) = \det(g_2)\}$.
- (4) $M_4 \cong \{(g_1, g_2, g_3) \in GL_3(F) \times GL_2(F) \times GL_5(F) \mid \det(g_1) = \det(g_2) = \det(g_3)\}$.
- (5) $M_5 \cong \{(g_1, g_2) \in GL_5(F) \times GL_4(F) \mid \det(g_1) = \det(g_2)\}$.
- (6) $M_6 \cong \left\{ (g_1, g_2) \in GSpin_{10}(F) \times GL_3(F) \mid \det(g_1) = \det(g_2) \in (F^\times)^2 \right\}$.
- (7) $M_7 \cong \{(g_1, g_2) \in GE_6(F) \times GL_2(F) \mid \det(g_1) = \det(g_2)\}$.
- (8) $M_8 \cong GE_7(F)$.

Where \det denotes the similitude factors on the relevant groups (in particular, the similitude factor on GL_n is the usual determinant).

We record here, for $1 \leq i \leq 8$, the cardinality of $W^{M_i, T}$, the set of shortest representatives of W_G/W_{M_i}

i	1	2	3	4	5	6	7	8
$ W^{M_i, T} $	2,160	17,280	69,120	483,840	241,920	60,480	6,720	240

We also mention that $|W_G| = 696,729,600$. Every $\lambda \in \mathbf{X}(T)$ is of the form

$$\lambda = \sum_{i=1}^8 \Omega_i \circ \bar{\omega}_{\alpha_i}.$$

As a shorthand, we will write

$$\left(\begin{array}{cccccccc} & & \Omega_2 & & & & & \\ \Omega_1 & \Omega_3 & \Omega_4 & \Omega_5 & \Omega_6 & \Omega_7 & \Omega_8 & \end{array} \right) = \sum_{i=1}^8 \Omega_i \circ \bar{\omega}_{\alpha_i}.$$

3. The Main Theorem

Theorem 3.1. *Let $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$ with $s \leq 0$ and let $k = \text{ord}(\chi)$.*

(1) *The following tables Tables 1-8 below lists all triples $[i, s, k]$ such that π is either non-regular or reducible. In particular, for each triple $[i, s, k]$ the entry in the i th table for this value of s and k will be*

- **irr.** for non-regular and irreducible π .
- **red.** for non-regular and reducible π .
- **red.*** for regular and reducible π .

For any triple $[i, s, k]$, not appearing in the tables, the degenerate principal series $i_{M_i}^G(\Omega_{M_i, s, \chi})$, with $\text{ord}(\chi) = k$, is regular and irreducible.

(2) *All $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$ admit a unique irreducible subrepresentation, with the exception of:*

- (a) *$[i, s, k]$ is one of $[1, -5/2, 1]$, $[3, -1/2, 2]$ $[6, 0, 1]$, $[6, 0, 2]$ and $[7, -3/2, 1]$. In these cases, the representation π admits a maximal semi-simple subrepresentation of length 2.*
- (b) *$[i, s, k]$ is one of $[2, -1/2, 1]$ and $[5, -1/2, 1]$, in which case the length of the maximal semi-simple subrepresentation of π is at most 2.*

(1) For $P = P_1$

s	$-\frac{23}{2}$	$-\frac{21}{2}$	$-\frac{19}{2}$	$-\frac{17}{2}$	$-\frac{15}{2}$	$-\frac{13}{2}$	$-\frac{11}{2}$	$-\frac{9}{2}$	$-\frac{7}{2}$	-3	$-\frac{5}{2}$	-2	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	0	
$ord(\chi)$																	
1	<i>red.*</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	
2	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>

TABLE 1. P_1 -Reducibility Points

(2) For $P = P_2$

s	$-\frac{17}{2}$	$-\frac{15}{2}$	$-\frac{13}{2}$	$-\frac{11}{2}$	$-\frac{9}{2}$	$-\frac{7}{2}$	-3	$-\frac{5}{2}$	-2	$-\frac{3}{2}$	$-\frac{7}{6}$	-1	$-\frac{5}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	0
$ord(\chi)$																
1	<i>red.*</i>	<i>irr.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>
2	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	\times	<i>irr.</i>	\times	<i>red.</i>	\times	<i>irr.</i>
3	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	\times	<i>irr.</i>	<i>irr.</i>	\times

TABLE 2. P_2 -Reducibility Points

(3) For $P = P_3$

s	$-\frac{13}{2}$	$-\frac{11}{2}$	$-\frac{9}{2}$	$-\frac{7}{2}$	-3	$-\frac{5}{2}$	-2	$-\frac{3}{2}$	$-\frac{7}{6}$	-1	$-\frac{5}{6}$	$-\frac{3}{4}$	$-\frac{1}{2}$	$-\frac{1}{4}$	$-\frac{1}{6}$	0
$ord(\chi)$																
1	<i>red.*</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>

2	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	\times	<i>red.</i>	\times	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	\times	<i>irr.</i>
3	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>red.</i>	\times	<i>irr.</i>	\times	<i>irr.</i>	\times	<i>irr.</i>	\times
4	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	\times	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	\times	<i>irr.</i>

TABLE 3. P_3 -Reducibility Points

(4) For $P = P_4$

$ord(\chi) \backslash s$	$-\frac{9}{2}$	$-\frac{7}{2}$	$-\frac{5}{2}$	-2	$-\frac{3}{2}$	$-\frac{7}{6}$	-1	$-\frac{5}{6}$	$-\frac{3}{4}$	$-\frac{1}{2}$	$-\frac{1}{3}$	$-\frac{3}{10}$	$-\frac{1}{4}$	$-\frac{1}{6}$	$-\frac{1}{10}$	0
1	<i>red.*</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>
2	\times	\times	<i>red.*</i>	<i>red.</i>	<i>red.</i>	\times	<i>red.</i>	\times	<i>red.</i>	<i>red.</i>	<i>irr.</i>	\times	<i>irr.</i>	<i>irr.</i>	\times	<i>irr.</i>
3	\times	\times	\times	\times	<i>red.*</i>	<i>red.</i>	\times	<i>red.</i>	\times	<i>red.</i>	<i>irr.</i>	\times	\times	<i>irr.</i>	\times	<i>irr.</i>
4	\times	\times	\times	\times	\times	\times	<i>red.*</i>	\times	<i>red.</i>	<i>red.</i>	\times	\times	<i>irr.</i>	\times	\times	<i>irr.</i>
5	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	\times	<i>red.</i>	\times	\times	<i>irr.</i>	\times
6	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	\times	\times	<i>irr.</i>	\times	<i>irr.</i>

TABLE 4. P_4 -Reducibility Points

(5) For $P = P_5$

DEGENERATE PRINCIPAL SERIES OF D_2

$ord(\chi) \backslash s$	$-\frac{11}{2}$	$-\frac{9}{2}$	$-\frac{7}{2}$	$-\frac{5}{2}$	-2	$-\frac{3}{2}$	$-\frac{7}{6}$	-1	$-\frac{5}{6}$	$-\frac{3}{4}$	$-\frac{1}{2}$	$-\frac{3}{10}$	$-\frac{1}{4}$	$-\frac{1}{6}$	$-\frac{1}{10}$	0
1	<i>red.*</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>
2	\times	\times	\times	<i>red.*</i>	<i>red.</i>	<i>red.</i>	\times	<i>red.</i>	\times	<i>irr.</i>	<i>red.</i>	\times	<i>irr.</i>	\times	\times	<i>irr.</i>
3	\times	\times	\times	\times	\times	<i>red.*</i>	<i>red.</i>	\times	<i>red.</i>	\times	<i>red.</i>	\times	\times	<i>irr.</i>	\times	\times
4	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	\times	<i>irr.</i>	<i>red.</i>	\times	<i>irr.</i>	\times	\times	<i>irr.</i>
5	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	\times	\times	<i>irr.</i>	\times

TABLE 5. P_5 -Reducibility Points

(6) For $P = P_6$

$ord(\chi) \backslash s$	-7	-6	-5	-4	-3	$-\frac{5}{2}$	-2	$-\frac{5}{3}$	$-\frac{3}{2}$	$-\frac{4}{3}$	-1	$-\frac{2}{3}$	$-\frac{1}{2}$	$-\frac{1}{3}$	$-\frac{1}{4}$	0
1	<i>red.*</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>
2	\times	\times	\times	\times	<i>red.*</i>	<i>red.</i>	<i>red.</i>	\times	<i>irr.</i>	\times	<i>red.</i>	\times	<i>red.</i>	\times	<i>irr.</i>	<i>red.</i>
3	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	\times	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	\times	<i>irr.</i>	\times	<i>irr.</i>
4	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	\times	<i>irr.</i>	<i>irr.</i>

TABLE 6. P_6 -Reducibility Points

(7) For $P = P_7$

$ord(\chi) \backslash s$	$-\frac{19}{2}$	$-\frac{17}{2}$	$-\frac{15}{2}$	$-\frac{13}{2}$	$-\frac{11}{2}$	$-\frac{9}{2}$	-4	$-\frac{7}{2}$	-3	$-\frac{5}{2}$	-2	$-\frac{3}{2}$	-1	$-\frac{1}{2}$	$-\frac{1}{6}$	0
1	<i>red.*</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>
2	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	\times	<i>irr.</i>
3	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>	<i>irr.</i>

TABLE 7. P_7 -Reducibility Points

(8) For $P = P_8$

$ord(\chi) \backslash s$	$-\frac{29}{2}$	$-\frac{27}{2}$	$-\frac{25}{2}$	$-\frac{23}{2}$	$-\frac{21}{2}$	$-\frac{19}{2}$	$-\frac{17}{2}$	$-\frac{15}{2}$	$-\frac{13}{2}$	$-\frac{11}{2}$	$-\frac{9}{2}$	$-\frac{7}{2}$	$-\frac{5}{2}$	$-\frac{3}{2}$	$-\frac{1}{2}$	0
1	<i>red.*</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>irr.</i>	<i>red.</i>	<i>irr.</i>
2	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	\times	<i>red.*</i>
																<i>irr.</i>

TABLE 8. P_8 -Reducibility Points

Remark 3.2. According to [GS05], the minimal representation of E_8 is the unique irreducible subrepresentation of $[8, -19/2, 1]$ (Note that P_8 is the Heisenberg parabolic subgroup of E_8). From the data provided by Tadić's reducibility criterion, it follows that the minimal representation is also a subquotient of the following cases: $[1, -17/2, 1]$, $[2, -13/2, 1]$ and $[4, -7/2, 1]$. Indeed, it is isomorphic to their unique irreducible subrepresentations.

Proof. We separate the proof into four parts: proof of reducibility (for all reducible π), proof of irreducibility (for most irreducible cases), proof of unique irreducible subrepresentation (for most cases) and exceptional cases. The last part is dealt with in Section 4 while the remainder of this section is devoted to the first three parts (which use the algorithm from Subsection 2.2).

3.1. Reducibility

For most reducible cases, it is enough to find $\pi' \neq \pi$ which shares an irreducible subquotient with π . As explained in Subsection 2.2.1, for most cases, one can find $\pi' = i_{M_i}^G(\Omega_{M_j, t, \chi^l})$ which satisfy the required conditions, while in others one needs to find π' of the form $\pi' = i_{M_{j_1, j_2}}^G \Omega_{s_1, s_2, \chi, k_1, k_2}$, where $\Omega_{s_1, s_2, \chi, k_1, k_2}$ is given by Equation (2.3).

In the following tables we list, for each reducible $[i, s, ord(\chi)]$ a triple $[j, t, ord(\chi^l)] = [j, t, ord(\chi)]$ or $[[j_1, j_2], [s_1, s_2], [k_1, k_2]]$ which provides a representation $\pi' \neq \pi$ sharing a common irreducible subquotient with π :

- For $P = P_1$






$s \backslash ord(\chi)$	1	2
$-\frac{17}{2}$	$[8, -\frac{19}{2}, 1]$	
$-\frac{13}{2}$	$[8, -\frac{11}{2}, 1]$	
$-\frac{11}{2}$	$[8, -\frac{5}{2}, 1]$	
$-\frac{7}{2}$	$[[1, 8], [-\frac{7}{2}, -2], [0, 0]]$	
$-\frac{5}{2}$	$[3, -\frac{5}{2}, 1]$	
$-\frac{1}{2}$	$[7, -\frac{3}{2}, 1]$	$[2, -\frac{5}{2}, 2]$

TABLE 9. Data for the proof of the reducibility of $i_{M_1}^G(\Omega_{M_1, s, \chi})$

- For $P = P_2$

$s \backslash ord(\chi)$	1	2
$-\frac{13}{2}$	$[8, -\frac{19}{2}, 1]$	
$-\frac{11}{2}$	$[8, -\frac{13}{2}, 1]$	
$-\frac{9}{2}$	$[8, -\frac{3}{2}, 1]$	
$-\frac{7}{2}$	$[1, -\frac{7}{2}, 1]$	
$-\frac{5}{2}$	$[7, -\frac{3}{2}, 1]$	$[1, -\frac{1}{2}, 2]$
-2		
$-\frac{3}{2}$	$[[1, 8], [\frac{1}{2}, -2], [0, 0]]$	$[[2, 8], [-2, 0], [1, 1]]$
$-\frac{1}{2}$	$[6, -1, 1]$	$[6, -1, 2]$

 TABLE 10. Data for the proof of the reducibility of $i_{M_2}^G(\Omega_{M_2, s, \chi})$

- For $P = P_3$

$s \backslash ord(\chi)$	1	2	3
$-\frac{11}{2}$	$[1, -\frac{19}{2}, 1]$		
$-\frac{9}{2}$	$[8, -\frac{15}{2}, 1]$		
$-\frac{7}{2}$	$[7, -\frac{9}{2}, 1]$		
$-\frac{5}{2}$	$[7, -\frac{5}{2}, 1]$	$[7, -\frac{5}{2}, 2]$	
-2	$[7, -1, 1]$	$[7, -1, 2]$	
$-\frac{3}{2}$	$[2, -\frac{3}{2}, 1]$	$[[6, 7], [1, -\frac{7}{2}], [0, 1]]$	
$-\frac{7}{6}$	$[2, -\frac{5}{6}, 1]$		$[[4, 5], [\frac{3}{2}, -\frac{19}{6}], [0, 1]]$
-1	$[[6, 8], [-1, 0], [0, 0]]$	$[[6, 7], [-3, 4], [0, 1]]$	
$-\frac{1}{2}$	$[6, 0, 1]$	$[[2, 3], [-\frac{5}{2}, 1], [1, 1]]$	

 TABLE 11. Data for the proof of the reducibility of $i_{M_3}^G(\Omega_{M_3, s, \chi})$

- For $P = P_4$
-

$s \backslash ord(\chi)$	1	2	3	4	5
$-\frac{7}{2}$	$[8, -\frac{19}{2}, 1]$				
$-\frac{5}{2}$	$[7, -\frac{9}{2}, 1]$				
-2	$[7, -3, 1]$	$[7, -3, 2]$			
$-\frac{3}{2}$	$[6, -2, 1]$	$[7, -\frac{1}{2}, 2]$			
$-\frac{7}{6}$	$[6, -\frac{4}{3}, 1]$		$[[1, 6], [6, -\frac{10}{3}], [0, 2]]$		
-1	$[3, -1, 1]$	$[3, -1, 2]$			
$-\frac{5}{6}$	$[6, -\frac{1}{3}, 1]$		$[6, -\frac{1}{3}, 3]$		
$-\frac{3}{4}$	$[3, -\frac{1}{4}, 1]$	$[3, -\frac{1}{4}, 2]$		$[[5, 6], [-\frac{11}{4}, \frac{17}{4}], [3, 3]]$	
$-\frac{1}{2}$	$[5, -\frac{1}{2}, 1]$	$[[3, 8], [-\frac{1}{2}, \frac{3}{2}], [1, 0]]$	$[[3, 8], [-\frac{1}{2}, \frac{3}{2}], [1, 1]]$	$[[2, 6], [-1, 0], [3, 3]]$	
$-\frac{3}{10}$	$[5, -\frac{1}{10}, 1]$				$[[2, 5], [-\frac{11}{10}, \frac{1}{5}], [2, 1]]$

 TABLE 12. Data for the proof of the reducibility of $i_{M_4}^G(\Omega_{M_4, s, \chi})$

- For $P = P_5$

$s \backslash ord(\chi)$	1	2	3	4
$-\frac{9}{2}$	$[8, -\frac{21}{2}, 1]$			
$-\frac{7}{2}$	$[7, -\frac{11}{2}, 1]$			
$-\frac{5}{2}$	$[6, -3, 1]$			
-2	$[1, -1, 1]$	$[1, -1, 2]$		
$-\frac{3}{2}$	$[2, -\frac{3}{2}, 1]$	$[2, -\frac{3}{2}, 2]$		
$-\frac{7}{6}$	$[2, -\frac{1}{6}, 1]$		$[2, -\frac{1}{6}, 3]$	
-1	$[6, -\frac{1}{2}, 1]$	$[6, -\frac{1}{2}, 2]$		
$-\frac{5}{6}$	$[3, -\frac{1}{6}, 1]$		$[[2, 3], [-\frac{7}{6}, 0], [2, 0]]$	
$-\frac{1}{2}$	$[[3, 8], [-\frac{1}{2}, \frac{3}{2}], [0, 0]]$	$[[3, 7], [-\frac{3}{2}, \frac{3}{2}], [1, 1]]$	$[[3, 8], [-\frac{1}{2}, \frac{3}{2}], [1, 0]]$	$[[2, 5], [\frac{1}{2}, -1], [3, 3]]$

 TABLE 13. Data for the proof of the reducibility of $i_{M_5}^G(\Omega_{M_5, s, \chi})$

- For $P = P_6$

$s \backslash ord(\chi)$	1	2	3
-6	$[8, -\frac{23}{2}, 1]$		
-5	$[7, -\frac{13}{2}, 1]$		

-4	$[8, -\frac{7}{2}, 1]$		
-3	$[1, -\frac{7}{2}, 1]$		
$-\frac{5}{2}$	$[1, -2, 1]$	$[1, -2, 2]$	
-2	$[7, -\frac{1}{2}, 1]$	$[[2, 3], [-\frac{3}{2}, -1], [1, 0]]$	
-1	$[2, -\frac{1}{2}, 1]$	$[2, -\frac{1}{2}, 2]$	$[[1, 7], [-2, \frac{5}{2}], [1, 1]]$
$-\frac{1}{2}$	$[[2, 8], [-\frac{1}{2}, -1], [0, 0]]$	$[[1, 2], [1, -\frac{3}{2}], [0, 1]]$	
0	$[3, -\frac{1}{2}, 1]$	$[[3, 7], [-\frac{3}{2}, \frac{1}{2}], [1, 1]]$	

 TABLE 14. Data for the proof of the reducibility of $i_{M_6}^G(\Omega_{M_6, s, \chi})$

- For $P = P_7$

$s \backslash ord(\chi)$	1	2
$-\frac{17}{2}$	$[8, -\frac{25}{2}, 1]$	
$-\frac{11}{2}$	$[8, -\frac{11}{2}, 1]$	
$-\frac{9}{2}$	$[8, -\frac{1}{2}, 1]$	
$-\frac{5}{2}$	$[1, -\frac{5}{2}, 1]$	$[3, -\frac{5}{2}, 2]$
$-\frac{3}{2}$	$[1, -\frac{1}{2}, 1]$	
$-\frac{1}{2}$	$[[6, 8], [-\frac{7}{2}, \frac{15}{2}], [0, 0]]$	$[[3, 8], [-\frac{5}{2}, \frac{3}{2}], [1, 0]]$

 TABLE 15. Data for the proof of the reducibility of $i_{M_7}^G(\Omega_{M_7, s, \chi})$

- For $P = P_8$

$s \backslash ord(\chi)$	1
$-\frac{19}{2}$	$[1, -\frac{17}{2}, 1]$
$-\frac{11}{2}$	$[1, -\frac{13}{2}, 1]$
$-\frac{1}{2}$	$[[1, 8], [-\frac{11}{2}, -1], [0, 0]]$

 TABLE 16. Data for the proof of the reducibility of $i_{M_8}^G(\Omega_{M_8, s, \chi})$

3.2. Irreducibility

For most irreducible non-regular $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$, it is possible to prove irreducibility using the branching rules calculation, as explained in Subsection 2.2.2. The only exceptions were those of the cases $[4, 0, 1]$ and $[4, 0, 2]$, which will be dealt with in Section 4.

3.3. Unique Irreducible Subrepresentation

Here too, for most reducible non-regular $\pi = i_{M_i}^G(\Omega_{M_i, s, \chi})$, one can use the algorithm in Subsection 2.2.3 to determine the length of the maximal semi-simple subrepresentation. There are, however, two kinds of exceptions:

- π with unique irreducible subrepresentation which could not have been determined using the algorithm: $[2, -5/2, 1]$, $[3, -3/2, 1]$, $[3, -3/2, 2]$, $[4, -1/2, 1]$, $[4, -1/2, 2]$, $[4, -3/2, 1]$, $[5, -3/2, 1]$, $[6, -2, 1]$ and $[6, -2, 2]$.
- Non-unitary π with a maximal semi-simple subrepresentation of length 2: $[1, -5/2, 1]$, $[3, -1/2, 2]$ and $[7, -3/2, 1]$.

Also, there are the cases $[2, -1/2, 1]$ and $[5, -1/2, 1]$ where we are able to show that the maximal semi-simple subrepresentation is of length at most 2. In these cases we further show that the irreducible spherical subquotient is a subrepresentation and describe the other candidate subrepresentation in terms of its Langlands data. \square

4. Exceptional Cases

In this section, we finish the proof of Theorem 3.1 by resolving the cases not solved by the algorithm of Subsection 2.2.

4.1. Fully Resolved Cases

In this subsection, we deal with all cases listed in Subsection 3.2 and Subsection 3.3 except for $[2, -1/2, 1]$ and $[5, -1/2, 1]$. We organize the discussion according to similarities between the arguments used for the various cases, such as:

- Using a chain of isomorphisms to embed π in a parabolic induction that admits a unique irreducible subrepresentation in common with π .

$[4, -1/2, 2]$	$\begin{pmatrix} \chi & & & & & & & \\ \chi & \chi & \chi & -1 & \chi & \chi & 0 & \end{pmatrix}$	$\{1, 2, 3, 4, 6, 7, 8\}$	$SL_5 \times SL_4$	1
$[5, -1/2, 2]$	$\begin{pmatrix} \chi & & & & & & & \\ \chi & \chi & \chi & -1 & \chi & \chi & \chi & \end{pmatrix}$	$\{1, 2, 3, 4, 6, 7, 8\}$	$SL_5 \times SL_4$	1
$[6, -2, 2]$	$\begin{pmatrix} \chi & & & & & & & \\ \chi & \chi & -1 & 0 & \chi & -1 & -1 & \end{pmatrix}$	$\{1, 2, 3, 5, 6\}$	$SL_3 \times SL_3 \times SL_2$	2

TABLE 17. Data for the proof of Lemma 4.1.

In the cases $[3, -1/2, 2]$, $[4, -1/2, 2]$ and $[5, -1/2, 2]$ the representation $(i_T^L \lambda_{a.d.}) \Big|_{L^{der}}$ is irreducible and hence, so is $i_T^L \lambda_{a.d.}$. In the other two cases, $[3, -3/2, 2]$ and $[6, -2, 2]$, the representation $(i_T^L \lambda_{a.d.}) \Big|_{L^{der}}$ has length 2. Since L is generated by L^{der} and T , the reducibility of $i_T^L \lambda_{a.d.}$ is determined by the action of T on the pieces of $(i_T^L \lambda_{a.d.}) \Big|_{L^{der}}$. Indeed, it follows that $i_T^L \lambda_{a.d.}$ is irreducible (see [HSb, Proposition 4.7(3)] for more details).

We note that

$$\operatorname{Re}(\langle \lambda, \alpha^\vee \rangle) < 0 \quad \forall \alpha \notin \Delta_\lambda.$$

Hence, by Langlands' unique irreducible subrepresentation theorem (see [Jan98, Section 1]),

$$i_T^G \lambda_{a.d.} \cong i_L^G (i_T^L \lambda_{a.d.})$$

admits a unique irreducible subrepresentation π_0 which appears in $i_T^G \lambda_{a.d.}$ with multiplicity 1. We now argue that

$$\operatorname{mult}(\lambda_{a.d.}, r_T^G \pi_0) = \operatorname{mult}(\lambda_{a.d.}, r_T^G (i_T^G \lambda_{a.d.})).$$

Assume to the contrary that there exists a subquotient $\tau \neq \pi_0$ of $i_T^G \lambda_{a.d.}$ such that $\lambda_{a.d.} \leq r_T^G(\tau)$. By a central character argument, see [HSa, Lemma. 3.12], it holds that $\tau \hookrightarrow i_T^G \lambda_{a.d.}$. Since π_0 is the unique irreducible subrepresentation of $i_T^G \lambda_{a.d.}$, it follows that $\tau \cong \pi_0$, contradicting the multiplicity 1 property of π_0 . \square

Remark 4.2. If one chooses a different anti-dominant element $\lambda'_{a.d.}$ in the orbit of $\lambda_{a.d.}$ in Table 17, they would have the same real part. Thus $\Theta \subset \Delta$ and L^{der} are invariant under the choice of representative of the orbit. Further more, $(i_T^L \lambda'_{a.d.}) \Big|_{L^{der}}$ is isomorphic to $(i_T^L \lambda_{a.d.}) \Big|_{L^{der}}$

Proposition 4.3. *In the following cases, $[i, s, ord(\chi)]$, the representation $\pi = i_{M_i}^G \Omega_{i,s,\chi}$ admits a unique irreducible subrepresentation:*

$$[3, -3/2, 2], [6, -2, 2]$$

Proof. Let $\lambda_{a.d.}$ be an anti-dominant exponent of π and π_0 be the unique irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$ as in Table 17 and Lemma 4.1. One checks, using a branching rule calculation, that in these cases

$$mult(\lambda_{a.d.}, r_T^G \pi_0) = |\text{Stab}_W(\lambda_{a.d.})| \Rightarrow mult(\lambda_0, r_T^G \pi_0) = mult(\lambda_0, r_T^G \pi) = 2.$$

From which the claim follows. Here,

$$|\text{Stab}_W(\lambda_{a.d.})| = \begin{cases} 8, & [i, s, ord(\chi)] = [3, -3/2, 2] \\ 4, & [i, s, ord(\chi)] = [6, -2, 2]. \end{cases}$$

□

In the following, we make a repeated use of a certain inclusion argument. It is thus convenient to state it in general at this point. For that matter, let $\Omega_0 = \Omega_{i,s,\chi}$ and $\pi = i_{M_i}^G \Omega_0$. For $j \neq i$ let $M_{i,j} = M_i \cap M_j$ and $\Omega_1 = r_{M_{i,j}}^{M_i} \Omega_0$. By Frobenius reciprocity,

$$\Omega_0 \hookrightarrow i_{M_{i,j}}^{M_i} \Omega_1.$$

and hence, by induction in stages,

$$(4.1) \quad \pi = i_{M_i}^G \Omega_0 \hookrightarrow i_{M_i}^G \left(i_{M_{i,j}}^{M_i} \Omega_1 \right) \cong i_{M_{i,j}}^G \Omega_1 \cong i_{M_j}^G \left(i_{M_{i,j}}^{M_j} \Omega_1 \right).$$

Proposition 4.4. *In the following cases, $[i, s, ord(\chi)]$, the representation $\pi = i_{M_i}^G \Omega_{i,s,\chi}$ admits a unique irreducible subrepresentation:*

$$[2, -5/2, 1], [3, -3/2, 1], [4, -1/2, 1], [4, -3/2, 1], [5, -1/2, 2], [5, -3/2, 1], [6, -2, 1].$$

Proof. Since the proof is similar for all cases, we start by outlining it and conclude by listing the relevant data for each of the cases in Table 18.

For $[i, s, ord(\chi)]$ as above, let $\lambda_0 = r_T^G \Omega_{i,s,\chi}$. For $j \neq i$ as in Table 18, let $M_{i,j} = M_i \cap M_j$ and $\Omega_1 = r_{M_{i,j}}^{M_i} \Omega_0$. It holds that

$$\lambda_0 = r_T^{M_i} \Omega = r_T^{M_{i,j}} \Omega_1.$$

Assume that $i_{M_{i,j}}^{M_j} \Omega_1$ is an irreducible representation of M_j , this can be verified by a branching rule calculation in M_j or by referring to previous knowledge on the representation theory of M_j as explained bellow in Remark 4.5. Let λ_1 be an M_j -anti-dominant exponent of $i_{M_{i,j}}^{M_j} \Omega_1$, that is $\lambda_1 \leq r_T^{M_j} \left(i_{M_{i,j}}^{M_j} \Omega_1 \right)$ and

$$(4.2) \quad \operatorname{Re}(\langle \lambda_1, \alpha_l^\vee \rangle) \leq 0 \quad \forall l \neq j.$$

By a central character argument, see [HSa, Lemma. 3.12], it holds that

$$i_{M_{i,j}}^{M_j} \Omega_1 \hookrightarrow i_T^{M_j} \lambda_1.$$

Hence, by Equation (4.1) and induction in stages,

$$\pi \hookrightarrow i_{M_j}^G \left(i_{M_{i,j}}^{M_j} \Omega_1 \right) \hookrightarrow i_T^G \lambda_1.$$

By Frobenius reciprocity, any irreducible subrepresentation τ of $i_T^G \lambda_1$ must satisfy

$$\lambda_1 \leq r_T^G \tau.$$

On the other hand, let $\lambda_{a.d.}$ be an anti-dominant exponent of π and let π_0 denote the unique irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$. If $\operatorname{ord}(\chi) = 1$ the existence of π_0 is automatic, and if $\operatorname{ord}(\chi) = 2$, it follows from Lemma 4.1.

In each of these cases, a branching rule calculation implies that

$$\operatorname{mult}(\lambda_1, r_T^G \pi) = \operatorname{mult}(\lambda_1, r_T^G \pi_0).$$

It follows that π_0 is the unique irreducible subquotient of π which admits λ_1 as an exponent.

In other words, π_0 is the unique irreducible subquotient of π such that

$$\pi_0 \hookrightarrow i_T^G \lambda_1.$$

On the other hand, since

$$\operatorname{mult}(\lambda_{a.d.}, r_T^G \pi) = \operatorname{mult}(\lambda_{a.d.}, r_T^G i_T^G \lambda_1) = \operatorname{mult}(\lambda_{a.d.}, r_T^G \pi_0),$$

it follows that π_0 appears in $i_T^G \lambda_1$, and in π , with multiplicity 1. We conclude that π_0 is the unique irreducible subrepresentation of π .

i	s	$\operatorname{ord}(\chi)$	j	λ_0	λ_1	$\lambda_{a.d.}$
2	$-\frac{5}{2}$	1	1	$\begin{pmatrix} 5 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} -1 \\ 2 & -1 & 0 & -1 & 0 & -1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 \\ -1 & 0 & -1 & 0 & 0 & -1 & 0 \end{pmatrix}$

3	$-\frac{3}{2}$	1	7	$\begin{pmatrix} -1 \\ -1 & 4 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & -1 & 1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & 0 \end{pmatrix}$
4	$-\frac{1}{2}$	1	8	$\begin{pmatrix} -1 \\ -1 & -1 & 3 & -1 & -1 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \end{pmatrix}$
4	$-\frac{3}{2}$	1	6	$\begin{pmatrix} -1 \\ -1 & -1 & 2 & -1 & -1 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & -1 \end{pmatrix}$
5	$-\frac{1}{2}$	2	8	$\begin{pmatrix} -1 \\ -1 & -1 & -1 & 4+\chi & -1 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} \chi \\ \chi & \chi & -1+\chi & \chi & \chi & \chi & 1+\chi \end{pmatrix}$	$\begin{pmatrix} \chi \\ \chi & \chi & \chi & -1 & \chi & \chi & \chi \end{pmatrix}$
5	$-\frac{3}{2}$	1	7	$\begin{pmatrix} -1 \\ -1 & -1 & -1 & 3 & -1 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & -1 & 1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & 0 \end{pmatrix}$
6	-2	1	8	$\begin{pmatrix} -1 \\ -1 & -1 & -1 & -1 & 4 & -1 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & -1 & -1 & 2 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & -1 \end{pmatrix}$

TABLE 18. Data for the proof of Proposition 4.4.

For the case $[5, -1/2, 2]$, χ denotes in Table 18 a character of F^\times of order 2.

□

Remark 4.5. The above argument, as well as other arguments bellow, used the fact that $\sigma = \iota_{M_{i,j}}^{M_j} \Omega_1$ is an irreducible representation of M_j . One way to prove the irreducibility of σ is using branching rule calculations but it can also be inferred from previous works. We wish to allow for a common reference for following arguments and thus we treat all values of j and not only 1, 6, 7 and 8.

More precisely, σ is a degenerate principal series of M_j and if its restriction $\sigma|_{M_j^{der}}$ to M_j^{der} is irreducible, then so is σ . For different values of j , one can find the list of reducible and irreducible degenerate principal series of M_j^{der} in the following sources:

- [BJ03] for $j = 1$.
- [Tad92] for $j = 2, 3, 4, 5$
- [BJ03] and [Tad92] for $j = 6$.
- [HSa, HSb] for $j = 7, 8$.

It should be noted here that [BJ03] deals only with representations of orthogonal groups, while $M_1^{der} = Spin_{14}$ and $M_6^{der} = Spin_{10} \times SL_2$. That is, in order to prove irreducibility, one should also take the isogeny map into account, using [Tad92].

A slight modification of the argument made in Proposition 4.4 can be used to prove:

Proposition 4.6. *The representation $\pi = i_{M_i}^G \Omega_{4, -\frac{1}{2}, \chi}$, where χ has order 2, admits a unique irreducible subrepresentation.*

Proof. Let $\Omega_0 = \Omega_{4, -\frac{1}{2}, \chi}$ and let

$$\lambda_0 = r_T^{M_4} \Omega_0 = \begin{pmatrix} & & & -1 & & & & \\ -1 & -1 & 3 + \chi & -1 & -1 & -1 & -1 & \end{pmatrix}.$$

We fix an anti-dominant exponent

$$\lambda_{a.d.} = \begin{pmatrix} & & & \chi & & & & \\ \chi & \chi & \chi & -1 & \chi & \chi & 0 & \end{pmatrix}$$

of π . Let $\Omega_1 = r_{M_{3,4}}^{M_4} \Omega_0$. By Equation (4.1),

$$\pi \hookrightarrow i_{M_3}^G \left(i_{M_{3,4}}^{M_3} \Omega_1 \right).$$

The representation $i_{M_{3,4}}^{M_3} \Omega_1$ of M_3 is an irreducible degenerate principal series of M_3 . This can be verified by a branching rule calculation in M_7 or by a similar argument to that of Remark 4.5. Furthermore, it holds that $i_{M_{3,4}}^{M_3} \Omega_1 \cong i_{M_{3,7}}^{M_3} \Omega_2$, where

$$r_T^{M_{3,7}} \Omega_2 = \begin{pmatrix} & & & -1 & & & & \\ -1 & 3 & -1 & -1 & -1 & 2 + \chi & -1 & \end{pmatrix}$$

and $i_{M_{3,7}}^{M_3} \Omega_2$ is the invert of the irreducible $i_{M_{3,4}}^{M_3} \Omega_1$. It follows, by induction in stages, that

$$(4.3) \quad \pi \hookrightarrow i_{M_3}^G \left(i_{M_{3,7}}^{M_3} \Omega_2 \right) \cong i_{M_7}^G \left(i_{M_{3,7}}^{M_7} \Omega_2 \right).$$

The M_7 -anti-dominant exponent of $i_{M_{3,7}}^{M_7} \Omega_2$ (see Equation (4.2)) is given by

$$\lambda_1 = \begin{pmatrix} & & & 0 & & & & \\ -1 & 0 & -1 & 0 & 0 & 2 + \chi & -1 & \end{pmatrix}.$$

By a central character argument, see [HSa, Lemma. 3.12],

$$i_{M_{3,7}}^{M_7} \Omega_1 \hookrightarrow i_T^{M_7} \lambda_1.$$

and hence

$$\pi \hookrightarrow i_T^G \lambda_1.$$

On the other hand, for the unique irreducible subrepresentation π_0 of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$ as in Lemma 4.1, a branching rule calculation shows that

$$\text{mult}(\lambda_1, r_T^G \pi) = \text{mult}(\lambda_1, r_T^G \pi_0) = 48.$$

Arguing as in Proposition 4.4, it follows that π admits a unique irreducible subrepresentation. \square

Proposition 4.7. *The representation $\pi = i_{M_4}^G \Omega_{4,0,\chi}$, where $\chi^2 = 1$, is irreducible.*

Proof. Let $\Omega_0 = \Omega_{4,0,\chi}$ and let $\Omega_1 = r_{M_{2,4}}^{M_4} \Omega_0$. By Equation (4.1),

$$\pi \hookrightarrow i_{M_2}^G \left(i_{M_{2,4}}^{M_2} \Omega_1 \right).$$

In what follows, we consider a sequence of isomorphisms, similar to the one in Equation (4.3). That is, we wish to write a sequence of isomorphisms:

$$(4.4) \quad i_{M_{i_k}}^G \left(i_{M_{i_k, j_k}}^{M_{i_k}} \Omega_k \right) \cong i_{M_{j_{k+1}}}^G \left(i_{M_{i_{k+1}, j_{k+1}}}^{M_{j_{k+1}}} \Omega_{k+1} \right) \cong i_{M_{i_{k+1}}}^G \left(i_{M_{i_{k+1}, j_{k+1}}}^{M_{i_{k+1}}} \Omega_{k+1} \right),$$

where

- $j_{k+1} = i_k$.
- Ω_k is a 1-dimensional representation of M_{i_k, j_k} .
- Ω_{k+1} is a 1-dimensional representation of $M_{i_k, j_{k+1}}$.
- $i_{M_{i_k, j_k}}^{M_{i_k}} \Omega_k$ is an irreducible representation of M_{i_k} .
- $i_{M_{j_{k+1}, i_{k+1}}}^{M_{j_{k+1}}} \Omega_{k+1}$ is the invert representation of $i_{M_{i_k, j_k}}^{M_{i_k}} \Omega_k$.

The sequence in Equation (4.4) relies on an isomorphism

$$(4.5) \quad i_{M_{i_k, j_k}}^{M_{i_k}} \Omega_k \cong i_{M_{j_{k+1}, i_{k+1}}}^{M_{j_{k+1}}} \Omega_{k+1}$$

of irreducible representations for each k . The irreducibility of each of the $i_{M_{i_k, j_k}}^{M_{i_k}} \Omega_k$ follows from a branching rule calculation in M_{i_k} and it can also be inferred from the references in Remark 4.5.

We summarize this data for the sequence of isomorphisms in the following table

k	i_k	j_k	$r_T^{M_{i_k, j_k}} \Omega_k$	i_{k+1}	$r_T^{M_{j_{k+1}, i_{k+1}}} \Omega_{k+1}$
1	2	4	$\begin{pmatrix} -1 \\ -1 & -1 & \frac{7}{2} + \chi & -1 & -1 & -1 & -1 \end{pmatrix}$	6	$\begin{pmatrix} \frac{7}{2} + \chi \\ -1 & -1 & -1 & -1 & \frac{5}{2} + \chi & -1 & -1 \end{pmatrix}$
2	6	2	$\begin{pmatrix} \frac{7}{2} + \chi \\ -1 & -1 & -1 & -1 & \frac{5}{2} + \chi & -1 & -1 \end{pmatrix}$	5	$\begin{pmatrix} -1 \\ -1 & -1 & -1 & \frac{5}{2} + \chi & \frac{3}{2} + \chi & -1 & -1 \end{pmatrix}$

3	5	6	$\begin{pmatrix} & -1 \\ -1 & -1 & -1 & \frac{5}{2} + \chi & \frac{3}{2} + \chi & -1 & -1 \end{pmatrix}$	8	$\begin{pmatrix} & -1 \\ -1 & -1 & -1 & 4 & -1 & -1 & \frac{1}{2} + \chi \end{pmatrix}$
---	---	---	--	---	---

TABLE 19. Data for the proof of Proposition 4.7.

Namely, the following sequence of isomorphisms hold

$$(4.6) \quad \begin{aligned} i_{M_2}^G \left(i_{M_{2,4}}^{M_2} \Omega_1 \right) &\cong i_{M_2}^G \left(i_{M_{2,6}}^{M_2} \Omega_1 \right) \cong i_{M_6}^G \left(i_{M_{2,6}}^{M_6} \Omega_1 \right) \\ &\cong i_{M_6}^G \left(i_{M_{5,6}}^{M_6} \Omega_1 \right) \cong i_{M_5}^G \left(i_{M_{5,6}}^{M_5} \Omega_1 \right) \cong i_{M_8}^G \left(i_{M_{5,8}}^{M_8} \Omega_4 \right). \end{aligned}$$

In each step we use either induction in stages or invert a representation. That is:

- $i_{M_2}^G \left(i_{M_{2,4}}^{M_2} \Omega_1 \right) \cong i_{M_2}^G \left(i_{M_{2,6}}^{M_2} \Omega_1 \right)$ since $i_{M_{2,4}}^{M_2} \Omega_1$ and $i_{M_{2,6}}^{M_2} \Omega_1$ are two irreducible degenerate principal series representations of M_2 which invert to each other.
- $i_{M_2}^G \left(i_{M_{2,6}}^{M_2} \Omega_1 \right) \cong i_{M_6}^G \left(i_{M_{2,6}}^{M_6} \Omega_1 \right)$ by induction in stages.
- etc.

It follows that

$$\pi \hookrightarrow i_{M_8}^G \left(i_{M_{5,8}}^{M_8} \Omega_4 \right).$$

Let

$$\lambda_1 = \begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & \frac{5}{2} + \chi \end{pmatrix}$$

be the M_8 -anti-dominant exponent of $i_{M_8}^G \left(i_{M_{5,8}}^{M_8} \Omega_4 \right)$. By a central character argument, see [HSa, Lemma. 3.12], it follows that

$$\pi \hookrightarrow i_{M_8}^G \left(i_{M_{5,8}}^{M_8} \Omega_4 \right) \hookrightarrow i_T^G \lambda_1.$$

Let

$$\lambda_{a.d.} = \begin{pmatrix} 0 \\ 0 & -\frac{1}{2} + \chi & 0 & 0 & -\frac{1}{2} + \chi & 0 & 0 \end{pmatrix}$$

denote the anti-dominant exponent of π and let π_0 denote the unique (due to [HSb, Lemma A.1]) subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$. A branching rule calculation yields that

$$\text{mult}(\lambda_1, r_T^G \pi_0) = 288 = \text{mult}(\lambda_1, r_T^G \pi)$$

Hence, π_0 is the unique subquotient of π which is a subrepresentation of $i_T^G \lambda_1$. Thus, π_0 is the unique irreducible subrepresentation of π . Since π is semi-simple, it follows that it is irreducible. \square

Proposition 4.8. *In the following cases, $[i, s, \text{ord}(\chi)]$, the maximal semi-simple subrepresentation of $\pi = i_{M_i}^G \Omega_{i,s,\chi}$ has length 2:*

$$[1, -5/2, 1], [3, -1/2, 2].$$

Proof. We prove the two cases using a similar argument and so we deal with them simultaneously and point out where the arguments diverge. Let $\Omega_0 = \Omega_{i,s,\chi}$ and $\pi = i_{M_i}^G \Omega_0$. We point out that

$$\text{mult}(\lambda_0, r_T^G \pi) = 2,$$

where $\lambda_0 = r_T^{M_i} \Omega_0$. Hence, π admits a maximal semi-simple subrepresentation of length at most 2.

For each case, we write a similar sequence of isomorphisms as in Equation (4.4) (see also Equation (4.6) and Remark 4.5), summarized in the following tables (analog to Table 19):

- Fix a character χ of order 2. In the case $[3, -1/2, 2]$:

k	i_k	j_k	$r_T^{M_{i_k \cdot j_k}} \Omega_k$	i_{k+1}	$r_T^{M_{j_{k+1}, i_{k+1}}} \Omega_{k+1}$
1	1	3	$\begin{pmatrix} -1 \\ -1 & 5+\chi & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$	2	$\begin{pmatrix} 5+\chi \\ 2+\chi & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$
2	2	1	$\begin{pmatrix} 5+\chi \\ 2+\chi & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$	8	$\begin{pmatrix} 5 \\ -1 & -1 & -1 & -1 & -1 & -1 & 4+\chi \end{pmatrix}$

TABLE 20. Data for the proof of Proposition 4.8 in the case $[3, -1/2, 2]$.

- In the case $[1, -5/2, 1]$:

k	i_k	j_k	$r_T^{M_{i_k \cdot j_k}} \Omega_k$	i_{k+1}	$r_T^{M_{j_{k+1}, i_{k+1}}} \Omega_{k+1}$
1	2	1	$\begin{pmatrix} -1 \\ 8 & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix}$	8	$\begin{pmatrix} 5 \\ -1 & -1 & -1 & -1 & -1 & -1 & -2 \end{pmatrix}$

TABLE 21. Data for the proof of Proposition 4.8 in the case $[1, -5/2, 1]$.

In order to match notations with the previous case, we write $\Omega_3 = \Omega_2$ for this case.

The conclusion of the above discussion is that in both cases it holds that

$$\pi \hookrightarrow i_{M_{2,8}}^G \Omega_3 \cong i_{M_8}^G \left(i_{M_{2,8}}^{M_8} \Omega_3 \right).$$

The representation $i_{M_{2,8}}^{M_8} \Omega_3$ is reducible and admits a maximal semi-simple subrepresentation of length 2 (this is the case denoted by $[2, -1, 1]$ in [HSb]). It follows that $i_{M_{2,8}}^G \Omega_3$ admits a maximal semi-simple subrepresentation of length at least 2.

On the other hand,

$$\text{mult} \left(\lambda_1, r_T^G \left(i_{M_{2,8}}^G \Omega_3 \right) \right) = \text{mult} \left(\lambda_1, r_T^G \pi \right) = 2,$$

where $\lambda_1 = r_T^{M_{2,8}} \Omega_3$. Hence, $i_{M_{2,8}}^G \Omega_3$ admits a maximal semi-simple subrepresentation of length precisely 2 and both of these subrepresentation intersect π , from which the claim follows. \square

The following is a slight variation of the previous argument.

Proposition 4.9. *The representation $\pi = i_{M_7}^G \Omega_{7,-3/2}$ admits a maximal semi-simple subrepresentation of length 2.*

Proof. Let $\Omega_0 = \Omega_{7,-3/2}$ and $\lambda_0 = r_T^G \Omega_0$. We note that

$$\text{mult} \left(\lambda_0, r_T^G \pi \right) = 2.$$

Hence, π admits a maximal semi-simple subrepresentation of length at most 2.

We write a similar sequence of isomorphisms to that of Equation (4.4) (see also Equation (4.6) and Remark 4.5), summarized in the following table (analog to Table 19):

k	i_k	j_k	$r_T^{M_{i_k, j_k}} \Omega_k$	i_{k+1}	$r_T^{M_{j_{k+1}, i_{k+1}}} \Omega_{k+1}$
1	3	7	$\begin{pmatrix} -1 \\ -1 & -1 & -1 & -1 & -1 & 7 & -1 \end{pmatrix}$	4	$\begin{pmatrix} -1 \\ -1 & 5 & -2 & -1 & -1 & -1 & -1 \end{pmatrix}$
2	4	3	$\begin{pmatrix} -1 \\ -1 & 5 & -2 & -1 & -1 & -1 & -1 \end{pmatrix}$	1	$\begin{pmatrix} -1 \\ -4 & -1 & 3 & -1 & -1 & -1 & -1 \end{pmatrix}$

TABLE 22. Data for the proof of Proposition 4.9.

It follows that

$$\pi \hookrightarrow i_{M_1}^G \left(i_{M_{1,4}}^{M_1} \Omega_3 \right).$$

Since $i_{M_{1,4}}^{M_1} \Omega_3$ is a reducible unitary degenerate principal series of M_1 , it follows that $i_{M_{1,4}}^{M_1} \Omega_3 = \sigma_0 \oplus \sigma_1$, with $\lambda_{a.d.} \leq r_T^{M_1} \sigma_0$ and $\lambda_{a.d.} \not\leq r_T^{M_1} \sigma_1$, where

$$\lambda_{a.d.} = \begin{pmatrix} 0 \\ -1 & 0 & -1 & 0 & 0 & -1 & 0 \end{pmatrix}.$$

On the other hand, $\lambda_1 \leq r_T^{M_1} \sigma_0, r_T^{M_1} \sigma_1$, where $\lambda_1 = r_T^{M_{1,4}} \Omega_3$. Using Equation (2.2), one checks that

$$\text{mult} \left(\lambda_1, r_T^G i_{M_{1,4}}^G \Omega_3 \right) = \text{mult} \left(\lambda_1, r_T^G \pi \right) = 2.$$

On the other hand,

$$\text{mult} \left(\lambda_1, r_T^G i_{M_1}^G \sigma_0 \right) = \text{mult} \left(\lambda_1, r_T^G i_{M_1}^G \sigma_1 \right) = 1.$$

It follows that $i_{M_1}^G \sigma_0$ and $i_{M_1}^G \sigma_1$ each contains a unique irreducible subrepresentation, π_0 and π_1 , both of which intersect π . Thus,

$$\pi_0 \oplus \pi_1 \hookrightarrow \pi.$$

□

4.2. Unresolved Cases

We finish with two cases we were unable to fully resolve, these are the cases of $\pi = i_{M_2}^G \Omega_{2, -\frac{1}{2}}$ and $\pi = i_{M_5}^G \Omega_{5, -\frac{1}{2}}$. In both cases we show that π admits a maximal semi-simple subrepresentation of length 1 or 2 and that the spherical subquotient of π is a subrepresentation. We were, however, unable to determine the precise length of the socle. We do, however, outline computational methods to determine this in the future, when stronger computers are readily available.

Let $\pi = i_{M_i}^G (\Omega_{M_i, s, \chi})$ with $s < 0$. Also, let $\lambda_{a.d.}$ be an anti-dominant exponent of π and let π_0 be an irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$. It seems to be the case that π_0 is a subrepresentation of π automatically. However, we were unable to find a general proof for that. While this usually follow from a simple branching rule calculation, this line of argument was insufficient in the following case and the claim requires a more circumvent proof.

Lemma 4.10. *Let $\pi = i_{M_2}^G \Omega_{2, -\frac{1}{2}}$ and let π_0 denote the unique irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$, where*

$$\lambda_{a.d.} = \begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

Then, $\pi_0 \hookrightarrow \pi$.

Proof. Since $\lambda_{a.d.}$ is the anti-dominant exponent of π , the uniqueness of π_0 follows from [HSb, Lemma A.1]. In fact, it follows that π_0 appears in $i_T^G \lambda_{a.d.}$ with multiplicity 1.

We fix the following exponents of π

$$\begin{aligned} \lambda_0 &= \begin{pmatrix} 7 \\ -1 & -1 & -1 & -1 & -1 & -1 & -1 \end{pmatrix} \\ \lambda_1 &= \begin{pmatrix} -1 \\ -1 & -1 & -1 & -1 & 5 & -1 & -1 \end{pmatrix} \\ \lambda_2 &= \begin{pmatrix} -1 \\ -1 & -1 & -2 & 1 & 4 & -1 & -1 \end{pmatrix}, \end{aligned}$$

where λ_0 is its initial exponent.

By a branching rule calculation, we have

$$\text{mult}(\lambda_1, r_T^G \pi) = \text{mult}(\lambda_1, r_T^G \pi_0) = 2.$$

On the other hand, by a central character argument (see [HSa, Lemma. 3.12] for details) it holds that $\pi_0 \hookrightarrow i_T^G \lambda_1$. We consider the normalized intertwining operator (see [HSb, Subsection 2D] for a short account on these operators and their properties)

$$i_T^G \lambda_1 \xrightarrow{N_w} i_T^G \lambda_0,$$

where

$$w = s_5 s_4 s_3 s_2 s_4 s_5 s_1 s_3 s_4 s_2.$$

Note that it decomposes as $N_w(\lambda_1) = N_{w'}(\lambda_2) \circ N_{s_5}(\lambda_1)$, where

$$w' = s_4 s_3 s_2 s_4 s_5 s_1 s_3 s_4 s_2$$

and that $N_{w'}(\lambda_2)$ is an isomorphism between $i_T^G \lambda_2$ and $i_T^G \lambda_0$.

On the other hand, the operator

$$i_T^G \lambda_1 \xrightarrow{N_{s_5}} i_T^G \lambda_2,$$

is not an isomorphism. However, Equation (2.2) implies that

$$\text{mult}(\lambda_{a.d.}, r_T^G \ker(N_{s_5}(\lambda_1))) = 0.$$

It follows that π_0 is not contained in the kernel of $N_{s_5}(\lambda_1)$ or $N_w(\lambda_1)$ and hence π_0 is a subrepresentation of $i_T^G \lambda_0$. In particular, since π_0 appears in $i_T^G \lambda_0$ with multiplicity 1, it is also a subrepresentation of π . □

Remark 4.11. By Equation (2.2), $\text{mult}(\lambda_0, r_T^G \pi) = 2$. Hence, the length of the socle of π is either 1 or 2. We point out that, by Table 14, π_0 is also an irreducible subquotient of $i_{M_6}^G \Omega_{6,-1}$. It follows from Equation (2.2) that $\text{mult}(\lambda_0, r_T^G i_{M_6}^G \Omega_{6,-1}) = 1$ and hence also $\text{mult}(\lambda_0, r_T^G \pi_0) = 1$, while on the other hand $\text{mult}(\lambda_0, r_T^G \pi) = 2$. Hence, π admits an irreducible subquotient $\pi_1 \neq \pi_0$ of π such that $\lambda_0 \leq r_T^G \pi_1$. It remains to determine whether π_1 is a subrepresentation of π or not. Namely, whether the length of the maximal semi-simple subrepresentation of π is 1 or 2.

It seems that the methods used above are futile for this case. One could technically use the method of [HSb, Proposition 4.9] to determine whether π_1 is a subrepresentation of π , or not. The method described there has two steps:

- Find a standard intertwining operator $N_w(\lambda_0)$, such that the kernel of the operator is composed of the irreducible subrepresentations of π other than π_0 .
- Calculate the dimension d of the space of Iwahori-fixed vectors in the kernel of $N_w(\lambda_0)$. If $d = 0$, then π_0 is the unique irreducible subrepresentation and if $d > 0$ then π admits a socle of length at least 2.

This is, however, beyond the capabilities of the computers available to us as the minimal Weyl word that could be used for this is

$$w = s_5 s_4 s_3 s_2 s_4 s_5 s_1 s_3 s_4 s_2.$$

which is of length 10. The issue is that the required computing time grows exponentially with the length of the Weyl word and the cardinalities of relevant Weyl groups and their coset spaces.

Bellow, we suggest a slightly simpler method to determine this in hope that it could be used in the near future. In the discussion, we give further indications on the properties of π_1 .

Let $\pi, \pi_0, \lambda_0, \lambda_1$ and $\lambda_{a.d.}$ be as in Lemma 4.10 and let w be as above.

Let $\Theta_0 = \{1, 2, 3, 4, 5\}$. Since $w \in W_{\Theta_0}$, the action of the intertwining operator $N_w(\lambda_0)$ factors through any of M_6 , M_7 or M_8 . We will describe a unified argument for a calculation that can be performed for each choice of M_j , with $j \in \{6, 7, 8\}$, to factor through. We will then explain why factorizing through M_8 seems to be the most efficient.

Write $\Omega_0 = \Omega_{2, -\frac{1}{2}, 1}$ and $\Omega_1 = r_{M_2}^{M_2, j}(\Omega_0)$.

We now recall, from [HSa, HSB, BJ03], that

$$\text{length}\left(i_{M_2, j}^{M_j} \Omega_1\right) = \begin{cases} 2, & j = 6, 7 \\ 3, & j = 8. \end{cases}$$

More precisely:

- If $j = 6$ or $j = 7$, $\tau = i_{M_2, j}^{M_j} \Omega_1$ is of length 2 and one may write a non-splitting exact sequence

$$\tau_0 \hookrightarrow \tau \twoheadrightarrow \tau_1,$$

where τ_0 and τ_1 are irreducible and τ_1 is spherical. In the case $j = 6$, the quotient τ_1 is in-fact 1-dimensional.

- If $j = 8$, $\tau = i_{M_2, 8}^{M_8} \Omega_1$ is of length 3 and one may write a non-splitting exact sequence

$$\tau_0 \hookrightarrow \tau \twoheadrightarrow \tau_1 \oplus \tau_{-1},$$

where τ_0 , τ_{-1} and τ_1 are irreducible and τ_1 is spherical.

Since

$$\text{mult}\left(\lambda_1, r_T^{M_j} \tau\right) = \text{mult}\left(\lambda_1, r_T^{M_j} \tau_1\right) = 1,$$

we conclude that the kernel and image of the intertwining operator $N_w^{M_j}(\lambda_0)$ is given by:

$$I = \text{Im}\left(N_w^{M_j}(\lambda_0)\right) = \tau_1, \quad K = \ker\left(N_w^{M_j}(\lambda_0)\right) = \begin{cases} \tau_0, & j = 6, 7 \\ \tau_0 + \tau_{-1}, & j = 8. \end{cases}$$

Hence, it holds that

$$i_{M_j}^G(K) \hookrightarrow i_{M_2, j}^G(\Omega_1) \twoheadrightarrow i_{M_j}^G(I).$$

On the other hand, since

$$\text{mult}\left(\lambda_1, r_T^G \pi_0\right) = \text{mult}\left(\lambda_1, r_T^G i_{M_2, j}^G(\Omega_1)\right) = \text{mult}\left(\lambda_1, r_T^G \pi\right) = 2,$$

it follows that π_0 is the unique irreducible subrepresentation of $i_{M_j}^G(I)$.

It follows from the above discussion that π admits a unique irreducible subrepresentation if and only if $\pi \cap i_{M_j}^G(K) = 0$.

In order to determine if the intersection is indeed 0, we consider

$$\dim_{\mathbb{C}} \left[\pi^{\mathcal{J}} \cap i_{M_j}^G(K)^{\mathcal{J}} \right],$$

where \mathcal{J} denotes the Iwahori-Hecke algebra of G . That is, we calculate the intersection of the 17,280-dimensional space $\pi^{\mathcal{J}}$ with $i_{M_j}^G(K)^{\mathcal{J}}$ whose dimension depends on j and is given by

$$\dim_{\mathbb{C}} \left(i_{M_j}^G(K)^{\mathcal{J}} \right) = \left[\dim_{\mathbb{C}} \left(r_T^{M_j}(K) \right) \right] \times |W^{M_j, T}|.$$

We collect relevant data in the following table:

j	6	7	8
$\dim_{\mathbb{C}} r_T^{M_j}(K)$	1919	15	120=105+15
$ W^{M_j, T} $	60,480	6,720	240
$\dim_{\mathbb{C}} \left(i_{M_j}^G(K)^{\mathcal{J}} \right)$	116,061,120	100,800	28,800

Thus, while the choice of $j = 6$ seems more intuitive, it seems like using $j = 8$ would be more efficient.

We now turn to deal with the other unresolved case, $\pi = i_{M_5}^G \Omega_{5, -\frac{1}{2}}$. In this case, a branching rule calculation shows that the irreducible spherical subquotient π_0 of π is a subrepresentation. However, from Equation (2.2) we have $\text{mult}(\lambda_0, r_T^G \pi) = 30$, where

$$\lambda_0 = \begin{pmatrix} -1 \\ -1 & -1 & -1 & 4 & -1 & -1 & -1 \end{pmatrix}$$

is the initial exponent of π . Thus, it is not immediately clear that the length of the socle of π is at most 2.

Lemma 4.12. *The representation $\pi = i_{M_5}^G \Omega_{5, -\frac{1}{2}}$ admits a maximal semi-simple subrepresentation of length at most 2.*

Proof. We consider the following exponents of π :

$$\begin{aligned} \lambda_0 &= \begin{pmatrix} -1 \\ -1 & -1 & -1 & 4 & -1 & -1 & -1 \end{pmatrix}, \\ \lambda_1 &= \begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 1 \end{pmatrix} \\ \lambda_{a.d.} &= \begin{pmatrix} 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \end{pmatrix}. \end{aligned}$$

Here, λ_0 is the initial exponent of π , $\lambda_{a.d.}$ is its anti-dominant exponent. Let π_0 denote the unique irreducible subquotient of π such that $\lambda_{a.d.} \leq r_T^G \pi_0$.

Since $\text{mult}(\lambda_0, r_T^G \pi) = 30$ and since a branching rule calculation yielded only $\text{mult}(\lambda_0, r_T^G \pi_0) \geq 4$, the claim is not immediate. However, this does imply that $\pi_0 \hookrightarrow \pi$.

On the other hand, [HSb, Lemma A.1] implies that for every $\sigma \in \text{Rep}(G)$, it holds that

$$288 \mid \text{mult}(\lambda_1, r_T^G \sigma).$$

Let $\Omega_1 = r_{M_{5,8}}^{M_5} \Omega_{5,-1\frac{1}{2}}$ and note that $i_{M_{5,8}}^{M_8} \Omega_1$ is irreducible. Hence, reasoning as in Proposition 4.4, it follows that

$$\pi \hookrightarrow i_{M_8}^G \left(i_{M_{5,8}}^{M_8} \Omega_1 \right) \hookrightarrow i_T^G \lambda_1$$

as $\lambda_1 = r_T^{M_{5,8}} \Omega_1$. By Equation (2.2),

$$\text{mult}(\lambda_1, r_T^G \pi) = 864$$

and on the other hand, a branching rule calculation yields,

$$\text{mult}(\lambda_1, r_T^G \pi_0) \geq 576.$$

Hence, π could have at most one more subquotient π_1 such that $\pi_1 \hookrightarrow i_T^G \lambda_1$. Thus, the length of the maximal semi-simple subrepresentation of π is at most 2. \square

Remark 4.13. Attempting to follow the methods of [HSb, Proposition 4.9] in order to determine the length of the socle of π is, too, beyond the capabilities of current available computers and even more so. The exponent λ “closest” to λ_0 such that a branching rule calculation guarantees that $\text{mult}(\lambda, r_T^G \pi_0) = \text{mult}(\lambda, r_T^G \pi)$ is $\lambda = \lambda_{a.d.}$ and the shortest Weyl element w such that $w \cdot \lambda_0 = \lambda$ is

$$w = s_5 s_6 s_7 s_8 s_4 s_3 s_2 s_4 s_5 s_6 s_7 s_4 s_1 s_3 s_2 s_4 s_5 s_6 s_4 s_1 s_3 s_2 s_4 s_5$$

which is of length 24 and thus, the associated intertwining operator cannot be realistically generated in currently available computers. Also, since this word contains all 8 generators of W , the associated operator does not factor via a Levi subgroup and the method suggested in Remark 4.11 is not applicable here.

Here, we are able to show that the same calculation can be performed with a Weyl element of length 21. While this is an improvement, this calculation still seems to be unfeasible.

As noted above,

$$\pi \hookrightarrow i_T^G \lambda_1 = i_{M_4}^G \left(i_T^{M_4} \lambda_1 \right).$$

We also note that, by [HSb, Lemma A.4 and Equation (OR)], $i_T^{M_4} \lambda_1$ is non-semi-simple of length 2, we write

$$\sigma_1 \hookrightarrow i_T^{M_4} \lambda_1 \twoheadrightarrow \sigma_0,$$

where σ_0 is spherical.

Let

$$\lambda_2 = \begin{pmatrix} 0 \\ 0 & 0 & -1 & 0 & 0 & -1 & 1 \end{pmatrix}.$$

We show that $i_T^{M_4} \lambda_2 = \sigma_0 \oplus \sigma_1$ and that π_i is the unique irreducible subrepresentation of $i_{M_4}^G \sigma_i$.

Indeed, let Ω_2 denote the 1-dimensional representation of L_5 such that $r_T^{L_5} \Omega_2 = \lambda_2$. By [NSS, Corollary 4.4], both $i_{L_5}^{M_4} \Omega_2$ and $i_{L_5}^{M_4} (\Omega_2 \otimes St_{L_5})$ admit a unique irreducible subrepresentation. Similarly, so do $i_{L_5}^G \Omega_2$ and $i_{L_5}^G (\Omega_2 \otimes St_{L_5})$.

On the other hand, since

$$i_T^{L_5} \lambda_2 = \Omega_2 \oplus (\Omega_2 \otimes St_{L_5}),$$

it follows that $\sigma_0 = i_T^{L_5} \Omega_2$ and $\sigma_1 = i_T^{L_5} (\Omega_2 \otimes St_{L_5})$. We thus conclude that π_i is the unique irreducible subrepresentation of $i_{M_4}^G \sigma_i$.

We now note that $\sigma_1 = \ker \left(N_{s_8(\lambda_1)}^{M_4} \right)$. On the other hand, $\pi \hookrightarrow i_T^G \lambda_1$. If π_1 is a subrepresentation of π , then the kernel of $N_{w'}(\lambda_0) \Big|_{\pi}$ is non-trivial, where

$$w' = s_8 s_4 s_3 s_2 s_4 s_5 s_6 s_7 s_4 s_1 s_3 s_2 s_4 s_5 s_6 s_4 s_1 s_3 s_2 s_4 s_5.$$

Here, too, all 8 generators of W appear and thus, the method suggested in Remark 4.11 cannot be applied to this case.

Appendix A. Branching Rules Database

In this section, we retain the notations of Subsection 2.1 and list data on irreducible representations of Levi subgroups which is used for the branching rule calculations performed for this paper.

That is, we wish to list triples of (λ, M, τ) where λ is a character of T , M is a Levi subgroup of G and τ is the unique irreducible representation of M such that $\lambda \leq r_T^M \tau$.

We start, by a general rule. For a character λ of T , set

$$\Theta_\lambda = \{ \alpha \in \Delta_L \mid \langle \lambda, \alpha^\vee \rangle = 0 \}.$$

It holds that $M = M_{\Theta_\lambda}$ admits a unique irreducible representation τ such that $\lambda \leq r_T^M \tau$, as shown in [NSS, Lemma 2.3], which satisfy

$$\left[r_T^{M_{\Theta_\lambda}} \tau \right] = \left| W_{M_{\Theta_\lambda}} \right| \times [\lambda].$$

This can be decoded by the *Orthogonality Rule* which states that, for any irreducible representation σ of G it holds that

$$\lambda \leq \sigma \Rightarrow \left| W_{M_{\Theta_\lambda}} \right| \times [\lambda] \leq r_T^G \sigma.$$

For other branching rules however, instead of listing the data by Levi subgroups M of G , it is more convenient to list them by simple factors L of the derived subgroups M^{der} . Moreover, we recall that the uniqueness of τ is not required, only the uniqueness of $[r_T^L \tau]$ is. Thus we list the semi-simplified Jacquet functors $[r_T^L \tau]$ of irreducible representations. Finally, it would be more convenient to write these in the “intrinsic coordinates” of a maximal split torus of L instead of those of the torus of G .

That is, we list triples of data $(L, \lambda, [r_T^G \tau])$, where:


- L is a simple, split and simply-connected p -adic group (whose Dynkin diagram is a sub-Dynkin diagram of E_8), by abuse of notations we fix a maximal split torus T of L .
- λ is a character of T .
- $[r_T^L \tau]$ is the unique semi-simplified Jacquet functor of an irreducible representation τ of L such that $\lambda \leq r_T^G \tau$.

We separate the list of branching rules by the type of L . Most of these rules are listed in [HSb, Appendix A] while the rest can be deduced from the results of [HSa, HSb, BJ03, Jan96, Jan93, BZ76].

For convenience of applying the branching rules, we list the elements appearing in $[r_T^L \tau]$ using the action of the Weyl group $W = W_L$ of L on the characters of T . Also, we point out the each rules can be written with several variations, either due to automorphisms of the Dynkin diagra (if L is of type A_n , D_n or E_6) but also due to the Aubert involution (see [Ban02]), **we list only one variation of each rule.**

A.1. L of type A_n

Let L be a simple group of type A_n . We think of it as a simple factor in the derived group M^{der} of a Levi subgroup M of G . We fix a labeling for the Dynkin diagram of type A_n and use this labeling to formulate the branching rules arising from L of type A_n instead of the labeling inherited from that of the Dynkin diagram of type E_8 given in Subsection 2.3. This labeling is given by



Bellow, we list the branching rules arising from L of type A_n which were implemented by us in the context of this paper, this list is by no means exhaustive and there are many branching rules arising from other irreducible representations of groups of type A_n :

- If L of type A_1 , it admits a unique simple root β_1 . For λ such that $\langle \lambda, \beta_1^\vee \rangle \neq \pm 1$, there exist a unique irreducible representation τ such that $\lambda \leq r_T^L \tau$. In fact, it holds that

$$[r_T^L(\tau)] = [\lambda] + [s_{\beta_1} \cdot \lambda].$$

This could be encoded as follows

$$(A.1) \quad \lambda \leq r_T^L(\tau), \langle \lambda, \beta_1^\vee \rangle \neq \pm 1 \implies [\lambda] + [s_{\beta_1} \cdot \lambda] \leq [r_T^L(\tau)].$$

In what follows, branching rules will be encoded in this fashion.

- For group L of type A_n with $n > 2$, we have the following rule:

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = \pm 1, \\ \langle \lambda, \beta_k^\vee \rangle = 0 \quad \forall 2 \leq k \leq n \end{array} \right. \implies [r_T^L(\tau)] = \sum_{w \in W^{L,T}} (n - l(w)) \cdot (n - 1)! [w \cdot \lambda],$$

where $M = M_{\{\beta_2, \dots, \beta_n\}}$.

For example, in type A_2 , this rule can be written as

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = \pm 1, \\ \langle \lambda, \beta_2^\vee \rangle = 0 \end{array} \right. \implies 2 \times [\lambda] + [s_{\beta_1} \cdot \lambda] \leq [r_T^L(\tau)].$$

In type A_3 , this rule can be written as

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = 1, \\ \langle \lambda, \beta_2^\vee \rangle = \langle \lambda, \beta_3^\vee \rangle = 0 \end{array} \right. \implies 6 \times [\lambda] + 4 \times [s_{\beta_1} \cdot \lambda] + 2 \times [s_{\beta_2} s_{\beta_1} \cdot \lambda] \leq [r_T^L(\tau)]$$

- For L of type A_3 we have the following two rules:

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = 1, \\ \langle \lambda, \beta_2^\vee \rangle = 0, \\ \langle \lambda, \beta_3^\vee \rangle = -1 \end{array} \right. \implies 2 \times [\lambda] + [s_{\beta_1} \cdot \lambda] + [s_{\beta_3} \cdot \lambda] + 2 \times [s_{\beta_1} s_{\beta_3} \cdot \lambda] \leq [r_T^L(\tau)]$$

and

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = 1, \\ \langle \lambda, \beta_2^\vee \rangle = 0, \\ \langle \lambda, \beta_3^\vee \rangle = 1 \end{array} \right. \implies 2 \times [\lambda] + [s_{\beta_1} \cdot \lambda] + [s_{\beta_3} \cdot \lambda] \leq [r_T^L(\tau)].$$

- In type A_4 , we have the following two rules:

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_2^\vee \rangle = 1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 1, 3, 4 \end{array} \right. \implies 12 \times [\lambda] + 8 \times [s_{\beta_2} \cdot \lambda] + 4 \times [s_{\beta_3} s_{\beta_2} \cdot \lambda] + 4 \times [s_{\beta_1} s_{\beta_2} \cdot \lambda] + 2 \times [s_{\beta_1} s_{\beta_3} s_{\beta_2} \cdot \lambda] \leq [r_T^L(\tau)].$$

and

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = 1, \\ \langle \lambda, \beta_4^\vee \rangle = -1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 2, 3 \end{array} \right. \implies 6 \times [\lambda] + 4 \times [s_{\beta_1} \cdot \lambda] + 4 \times [s_{\beta_4} \cdot \lambda] + 4 \times [s_{\beta_1} s_{\beta_4} \cdot \lambda] + 2 \times [s_{\beta_3} s_{\beta_4} \cdot \lambda] \\ + 2 \times [s_{\beta_2} s_{\beta_1} \cdot \lambda] + 4 \times [s_{\beta_4} s_{\beta_2} s_{\beta_1} \cdot \lambda] + 4 \times [s_{\beta_1} s_{\beta_3} s_{\beta_4} \cdot \lambda] \leq [r_T^L(\tau)].$$

- In type A_5 we have the following two rules:

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_3^\vee \rangle = 1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 1, 2, 4, 5 \end{array} \right.$$

$$\implies 36 \times [\lambda] + 24 \times [s_{\beta_3} \cdot \lambda] + 12 \times [s_{\beta_2} s_{\beta_3} \cdot \lambda] + 12 \times [s_{\beta_4} s_{\beta_3} \cdot \lambda] + 6 \times [s_{\beta_2} s_{\beta_4} s_{\beta_3} \cdot \lambda] \leq [r_T^L(\tau)].$$


and

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = 1, \\ \langle \lambda, \beta_4^\vee \rangle = -1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 2, 3, 5 \end{array} \right.$$

$$\begin{aligned} \implies & 12 \times [\lambda] + 12 \times [s_{\beta_5} s_{\beta_4} s_{\beta_2} s_{\beta_1} \cdot \lambda] + 8 \times [s_{\beta_1} \cdot \lambda] + 8 \times [s_{\beta_4} \cdot \lambda] + 8 \times [s_{\beta_1} s_{\beta_4} \cdot \lambda] \\ & + 8 \times [s_{\beta_4} s_{\beta_2} s_{\beta_1} \cdot \lambda] + 8 \times [s_{\beta_1} s_{\beta_3} s_{\beta_4} \cdot \lambda] + 8 \times [s_{\beta_5} s_{\beta_4} s_{\beta_1} \cdot \lambda] + 4 \times [s_{\beta_3} s_{\beta_4} \cdot \lambda] \\ & + 4 \times [s_{\beta_5} s_{\beta_4} \cdot \lambda] + 4 \times [s_{\beta_5} s_{\beta_1} \cdot \lambda] + 4 \times [s_{\beta_5} s_{\beta_1} s_{\beta_3} s_{\beta_4} \cdot \lambda] + 2 \times [s_{\beta_5} s_{\beta_3} s_{\beta_4} \cdot \lambda] \leq [r_T^L(\tau)]. \end{aligned}$$

A.2. L of type D_n

We fix the following labeling of the Dynkin diagram of type D_n .



If L is of type D_n we encode the branching rules in a similar fashion to that of type A_n .

- For L of type D_4 , we have the following two rules:

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_2^\vee \rangle = 1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 1, 3, 4 \end{array} \right.$$

$$\begin{aligned} \implies & 8 \times [\lambda] + 5 \times [s_{\beta_2} \cdot \lambda] + 2 \times [s_{\beta_1} s_{\beta_2} \cdot \lambda] + 2 \times [s_{\beta_3} s_{\beta_2} \cdot \lambda] + 2 \times [s_{\beta_4} s_{\beta_2} \cdot \lambda] \\ & + [s_{\beta_1} s_{\beta_3} s_{\beta_2} \cdot \lambda] + [s_{\beta_1} s_{\beta_4} s_{\beta_2} \cdot \lambda] + [s_{\beta_3} s_{\beta_4} s_{\beta_2} \cdot \lambda] \leq [r_T^L(\tau)]. \end{aligned}$$

and

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_1^\vee \rangle = 1, \\ \langle \lambda, \beta_4^\vee \rangle = -1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 2, 3 \end{array} \right.$$

$$\implies 12 \times [\lambda] + 8 \times [s_{\beta_1} \cdot \lambda] + 8 \times [s_{\beta_4} \cdot \lambda] + 12 \times [s_{\beta_1} s_{\beta_4} \cdot \lambda] \\ + 4 \times [s_{\beta_2} s_{\beta_4} \cdot \lambda] + 4 \times [s_{\beta_2} s_{\beta_1} \cdot \lambda] \leq [r_T^L(\tau)].$$

- For L of type D_5 , we have the following two rules:

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_5^\vee \rangle = 1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 1, 2, 3, 4 \end{array} \right.$$

$$\implies 120 \times [\lambda] + 96 \times [s_{\beta_5} \cdot \lambda] + 72 \times [s_{\beta_3} s_{\beta_5} \cdot \lambda] + 48 \times [s_{\beta_2} s_{\beta_3} s_{\beta_5} \cdot \lambda] \\ + 48 \times [s_{\beta_4} s_{\beta_3} s_{\beta_5} \cdot \lambda] + 32 \times [s_{\beta_4} s_{\beta_2} s_{\beta_3} s_{\beta_5} \cdot \lambda] + 24 \times [s_{\beta_1} s_{\beta_2} s_{\beta_3} s_{\beta_5} \cdot \lambda] \\ + 16 \times [s_{\beta_3} s_{\beta_2} s_{\beta_4} s_{\beta_3} s_{\beta_5} \cdot \lambda] + 16 \times [s_{\beta_1} s_{\beta_2} s_{\beta_4} s_{\beta_3} s_{\beta_5} \cdot \lambda] + 8 \times [s_{\beta_3} s_{\beta_1} s_{\beta_2} s_{\beta_4} s_{\beta_3} s_{\beta_5} \cdot \lambda] \leq [r_T^L(\tau)].$$

and

$$\left\{ \begin{array}{l} \lambda \leq r_T^L(\tau), \\ \langle \lambda, \beta_3^\vee \rangle = 1, \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 1, 2, 4, 5 \end{array} \right.$$

$$\implies 24 \times [\lambda] + 16 \times [s_{\beta_3} \cdot \lambda] + 8 \times [s_{\beta_2} s_{\beta_3} \cdot \lambda] + 8 \times [s_{\beta_4} s_{\beta_3} \cdot \lambda] \\ + 8 \times [s_{\beta_5} s_{\beta_3} \cdot \lambda] + 4 \times [s_{\beta_4} s_{\beta_5} s_{\beta_3} \cdot \lambda] + 4 \times [s_{\beta_2} s_{\beta_5} s_{\beta_3} \cdot \lambda] \\ + 4 \times [s_{\beta_2} s_{\beta_4} s_{\beta_3} \cdot \lambda] + 2 \times [s_{\beta_4} s_{\beta_2} s_{\beta_5} s_{\beta_3} \cdot \lambda] + 2 \times [s_{\beta_3} s_{\beta_4} s_{\beta_2} s_{\beta_5} s_{\beta_3} \cdot \lambda] \leq [r_T^L(\tau)].$$

- For L of type D_6 , we use one branching rule. For this rule, however, the list of exponents contains 30 different exponents of various multiplicities. Thus, we give a rough explanation on how to determine the Jacquet functor of τ in the relevant case, see [HSb, Lemma A.6] for more details. First, we consider the degenerate principal series representation σ of L corresponding to the notation $[D_6, 3, 0, 1]$. This is a direct sum of two irreducible representations, one spherical and the other is not. We consider the spherical irreducible constituent τ , this is the unique irreducible

representation of L such that

$$\begin{cases} \lambda \leq r_T^L \tau \\ \langle \lambda, \beta_1^\vee \rangle = \langle \lambda, \beta_4^\vee \rangle = -1 \\ \langle \lambda, \beta_k^\vee \rangle = 0, \quad k = 2, 3, 5, 6. \end{cases}$$

Let τ' denote the non-spherical constituent of σ . Using the method described in [HSb, Appendix B], one can show that

$$\begin{aligned} \dim_{\mathbb{C}}(r_T^L \tau) &= 155 \\ \dim_{\mathbb{C}}(r_T^L \tau') &= 5 \\ \dim_{\mathbb{C}}(r_T^L \sigma) &= 160. \end{aligned}$$

The initial exponent of σ is given by

$$\lambda_0 = \begin{bmatrix} & & & -1 & & \\ & & & & & \\ -1 & -1 & 3 & -1 & -1 & \end{bmatrix}.$$

Applying Equation (A.1), together with the above, yields


$$[r_T^L \tau'] = [\lambda_0] + [s_{\beta_3} \cdot \lambda_0] + [s_{\beta_2} s_{\beta_3} \cdot \lambda_0] + [s_{\beta_4} s_{\beta_3} \cdot \lambda_0] + [s_{\beta_2} s_{\beta_4} s_{\beta_3} \cdot \lambda_0].$$

One then computes $[r_T^L \sigma]$ using Equation (2.2) and the structure of $[r_T^L \tau]$ follows by reducing the multiplicity of these 5 exponents by 1.

A.3. L of type E_n


In the branching rules for L of type E_n , the list of exponents is long. Thus, we only list the irreducible degenerate principal series of L from which we derive the branching rules used by us. One then needs to compute $[r_T^L \tau]$ using Equation (2.2) in order to determine the branching rule explicitly.

We fix the following labeling of the Dynkin diagram of group of type E_6 .



The irreducible degenerate principal series of L of type E_6 whose Jacquet functor $[r_T^L \tau]$ contribute additional information to the one given previously are: $[E_6, 5, -\frac{1}{2}, 1]$ and $[E_6, 3, -\frac{1}{2}, 1]$.

We fix the following labeling of the Dynkin diagram of group of type E_7 .



The irreducible degenerate principal series of L whose Jacquet functor $[r_T^L \tau]$ contribute additional information to the one given previously are: $[E_7, 4, 0, 1]$, $[E_7, 5, 0, 1]$ and $[E_7, 5, -\frac{3}{2}, 1]$.

References

- [Ban02] Dubravka Ban. The Aubert involution and R-groups. *Ann. Sci. École Norm. Sup. (4)*, 35(5):673–693, 2002.
- [BJ03] Dubravka Ban and Chris Jantzen. Degenerate principal series for even-orthogonal groups. *Represent. Theory*, 7:440–480 (electronic), 2003.
- [BZ76] I. N. Bernšteĭn and A. V. Zelevinskiĭ. Representations of the group $GL(n, F)$, where F is a local non-Archimedean field. *Uspehi Mat. Nauk*, 31(3(189)):5–70, 1976.
- [BZ77] I. N. Bernstein and A. V. Zelevinsky. Induced representations of reductive p -adic groups. I. *Ann. Sci. École Norm. Sup. (4)*, 10(4):441–472, 1977.
- [Cas74] W. Casselman. Introduction to the theory of admissible representations of p -adic reductive groups. <http://www.math.ubc.ca/~cass/research/pdf/p-adic-book.pdf>, 1974.
- [CJ10] Seungil Choi and Chris Jantzen. Degenerate principal series for the exceptional p -adic groups of type F_4 . *J. Lie Theory*, 20(4):785–806, 2010.
- [GK81] S. S. Gelbart and A. W. Knap. Irreducible constituents of principal series of $SL_n(k)$. *Duke Math. J.*, 48(2):313–326, 1981.
- [GS] Nadya Gurevich and Avner Segal. Poles of the standard \mathcal{L} -function of G_2 and the rallis-schiffmann lift. *Preprint*.
- [GS05] Wee Teck Gan and Gordan Savin. On minimal representations definitions and properties. *Represent. Theory*, 9:46–93, 2005.
- [HSa] Hezi Halawi and Avner Segal. The degenerate principal series representations of exceptional groups of type e_6 over p -adic fields. *To appear in the Israel Journal of Mathematics*.
- [HSb] Hezi Halawi and Avner Segal. The degenerate principal series representations of exceptional groups of type e_7 over p -adic fields. *Preprint*.
- [HSc] Hezi Halawi and Avner Segal. Poles, residues and Siegel-Weil identities of degenerate Eisenstein series on split exceptional groups of type e_n . *Unpublished*.

- [Jan93] Chris Jantzen. Degenerate principal series for symplectic groups. *Mem. Amer. Math. Soc.*, 102(488):xiv+111, 1993.
- [Jan95] Chris Jantzen. On the Iwahori-Matsumoto involution and applications. *Ann. Sci. École Norm. Sup. (4)*, 28(5):527–547, 1995.
- [Jan96] Chris Jantzen. Degenerate principal series for symplectic and odd-orthogonal groups. *Mem. Amer. Math. Soc.*, 124(590):viii+100, 1996.
- [Jan98] Chris Jantzen. Some remarks on degenerate principal series. *Pacific J. Math.*, 186(1):67–87, 1998.
- [NSS] Taeuk Nam, Avner Segal, and Lior Silberman. Singularities of intertwining operators and certain decompositions of principal series representations. *preprint*, <https://arxiv.org/abs/1811.00803>.
- [Seg18] Avner Segal. The degenerate Eisenstein series attached to the Heisenberg parabolic subgroups of quasi-split forms of $Spin_8$. *Trans. Amer. Math. Soc.*, 370(8):5983–6039, 2018.
- [Seg19] Avner Segal. The degenerate residual spectrum of quasi-split forms of $Spin_8$ associated to the Heisenberg parabolic subgroup. *Trans. Amer. Math. Soc.*, 372(9):6703–6754, 2019.
- [Ste68] Robert Steinberg. *Lectures on Chevalley groups*. Yale University, New Haven, Conn., 1968. Notes prepared by John Faulkner and Robert Wilson.
- [Tad92] Marko Tadić. Notes on representations of non-Archimedean $SL(n)$. *Pacific J. Math.*, 152(2):375–396, 1992.
- [Tad98] Marko Tadić. On reducibility of parabolic induction. *Israel J. Math.*, 107:29–91, 1998.
- [TheYY] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version x.y.z)*, YYYY. <http://www.sagemath.org>.

¹ SCHOOL OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, POB 653, BE'ER SHEVA 84105, ISRAEL

² MATHEMATICS DEPARTMENT, SHAMOON COLLEGE OF ENGINEERING, 56 BIALIK ST., BEER-SHEVA 84100, ISRAEL

Email address: halawi@post.bgu.ac.il, avnerse@sce.ac.il