

# CURVILINEARITY AND ORTHOGONALITY

VLADIMIR S. CHELYSHKOV\*

## Abstract.

We introduce sequences of functions orthogonal on a finite interval: proper orthogonal rational functions, orthogonal exponential functions, and orthogonal logarithmic functions.

**Key words.** Altered Legendre polynomials, orthogonal rational functions, orthogonal exponential functions, orthogonal logarithmic functions.

**AMS subject classifications.** 41A10, 65D15, 65L60

**1. Introduction.** General properties of orthogonal rational functions were studied in [2, 13]. Specific examples of orthogonal rational functions on the semi-axis were given in [1, 9, 14]. Alternative orthogonal rational functions on a half-line which provide term-by-term increasing rate of decay at infinity were introduced in [7].

Orthogonal exponential functions on the semi-axis were presented in [10]. Alternative orthogonal exponential functions on the semi-axis with term-by-term increasing rate of decay at infinity were developed in [4, 5, 6].

Standard theory of orthogonal polynomials [11] includes statement that zeros of the polynomials are distributed in an orthogonality interval. Alternative orthogonal polynomials [5, 6] have zeros both inside and at one end of an interval. Structured orthogonal polynomials [8] admit zeros inside and at both ends of an interval. One can also introduce orthogonal polynomials that have prescribed zeros outside of an orthogonality interval. Being not of specific interest by itself, such altered orthogonal polynomials may originate orthogonal rational functions and other orthogonal functions on an interval.

In this preliminary notice, we alter Legendre polynomials [12] and use transforms for independent variable  $x$  on the  $x$ -axis to compose sequences of proper orthogonal rational functions on an interval, as well as sequences orthogonal exponential functions and orthogonal logarithmic functions on an interval.

**2. Altered Legendre polynomials.** Let  $\tilde{P}_n(x)$  be shifted Legendre polynomials. Making use of  $\tilde{P}_n(x)$ , we introduce polynomials

$$A_n(x) = \frac{x+1}{2} \tilde{P}_{n-1}(x), \quad x \in [-1, 1], \quad n = 1, 2, \dots \quad (2.1)$$

that are orthogonal

$$\int_0^1 \frac{A_m(x)A_n(x)}{(x+1)^2} dx = \frac{\delta_{mn}}{4(2n-1)}$$

and have a fixed real zero  $x = -1$  outside the interval of orthogonality. Properties of  $A_n(x)$  can be easily derived from definition (2.1) and properties of Legendre polynomials. Altered Legendre polynomials  $A_n(x)$  form a basis in  $L^2[0, 1]$ , and a polynomial  $\tilde{P}_k(x)$  can be approximated by the sequence  $A_n(x)$  on the interval. In particular, one can find

$$\tilde{P}_0(x) = 1 \approx 2 \ln 2 \cdot A_1(x) - 6(3 \ln 2 - 2)A_2(x) + 10(13 \ln 2 - 9)A_3(x) - \dots \quad (2.2)$$

Expansion (2.2) provides fast convergence for  $x \in [0, 1]$  and shows good results for extrapolation outside of the interval.

\*Oglethorpe University

**3. Proper orthogonal rational functions on an interval.** Making use of Cayley transform<sup>1</sup> [3]

$$x \rightarrow \frac{1-x}{1+x}$$

that maps points  $\{-1, 0, 1\}$  to points  $\{\infty, 1, 0\}$  on the  $x$ -axis, we define proper Legendre rational functions as a composition

$$R_n(x) := (-1)^{n-1} A_n \left( \frac{1-x}{1+x} \right), \quad x \in [0, \infty), \quad n = 1, 2, \dots .$$

Functions  $R_n(x)$  are orthogonal on the interval  $[0, 1]$ , and

$$\int_0^1 R_m(x) R_n(x) dx = \frac{\delta_{mn}}{2(2n-1)}.$$

It may be noted that function

$$f(x) = \frac{1+x}{1-x}$$

is an involution, so function  $f^{-1}(x)$  provides the same composition as  $f(x)$  does.




FIG. 3.1.  $R_n(x)$ ,  $n = 1 - 6$ .

We find

$$R_1(x) = \frac{1}{x+1},$$

$$R_2(x) = \frac{1-3x}{(x+1)^2},$$

$$R_3(x) = \frac{13x^2 - 10x + 1}{(x+1)^3},$$

$$R_4(x) = \frac{(1-3x)(21x^2 - 18x + 1)}{(x+1)^4},$$

<sup>1</sup>The transform permutes elements  $\{-1, 0, 1, \infty\}$  in sequence.

$$R_5(x) = \frac{321x^4 - 516x^3 + 246x^2 - 36x + 1}{(1+x)^5},$$

$$R_6(x) = \frac{(1-3x)(1-52x+454x^2-948x^3+561x^4)}{(1+x)^6},$$

and

$$\lim_{x \rightarrow \infty} R_n(x) = 0.$$

From properties of Legendre polynomials it follows that  $R_n(x)$  with even indices have a zero at  $x = 1/3$ .

Functions  $R_n(x)$  form a basis in  $L^2[0, 1]$ . In the complex  $z$ -plain,  $R_n(z)$  has one  $n$ -degree pole at  $z = -1$ .

**4. Orthogonal exponential functions on an interval.** Making use of a transform of exponential type

$$x \rightarrow 2^{1-x} - 1$$

that maps points  $\{-1, 0, 1\}$  to points  $\{\infty, 1, 0\}$  on the  $x$ -axis, we compose a sequence of Legendre exponential functions

$$E_n(x) := (-1)^{n-1} A_n(2^{1-x} - 1), \quad x \in [0, \infty), \quad n = 1, 2, \dots$$

that obey orthogonality relation

$$\int_0^1 2^x E_m(x) E_n(x) dx = \frac{\delta_{mn}}{2 \ln 2 (2n-1)}.$$

For convenience, we introduce polynomial functions  $F_n(t)$  and specify  $E_n(x)$  as compositions of  $F_n(t)$  and an exponential function  $t(x) = 2^{-x}$ , that is,

$$E_n(x) \equiv F_n(t)_{t=2^{-x}} \quad n = 1, 2, \dots$$

We find




FIG. 4.1.  $E_n(x)$ ,  $n = 1 - 6$ .

$$F_1(t) = t,$$

$$F_2(t) = t(4t - 3),$$

$$F_3(t) = t(24t^2 - 36t + 13),$$

$$F_4(t) = t(4t - 3)(40t^2 - 60t + 21),$$

$$F_5(t) = t(1120t^4 - 3360t^3 + 3720t^2 - 1800t + 321),$$

$$F_6(t) = t(4t - 3)(2016t^4 - 6048t^3 + 6664t^2 - 3192t + 561),$$

and

$$\lim_{x \rightarrow \infty} E_n(x) = 0.$$

It appears that  $E_n(x)$  with even indices have a zero at

$$x = 2 - \ln 3 / \ln 2 = 0.415037499\dots \quad (4.1)$$

Functions  $E_n(x)$  form a basis in  $L^2[0, 1]$ .

**5. Orthogonal logarithmic functions on an interval.** A function

$$g(x) = 2^{1-x} - 1$$

is not an involution, but it is a near involution on the interval  $[0, 1]$ . Inverse of  $g(x)$  is the function

$$g^{-1}(x) = 1 - \log_2(1 + x),$$

and one may find that

$$\left( \int_0^1 (g(x) - g^{-1}(x))^2 dx \right)^{1/2} = 0.002719224\dots$$

Also,

$$g(x) = g^{-1}(x) \quad \text{at} \quad x = 0, \quad x = 0.456999559\dots, \quad \text{and} \quad x = 1.$$

Making use of a transform of logarithmic type

$$x \rightarrow 1 - \log_2(1 + x)$$

that maps the points  $\{-1, 0, 1\}$  to  $\{3, 1, 0\}$  on the  $x$ -axis, we define Legendre logarithmic functions

$$L_n(x) := (-1)^{n-1} A_n(1 - \log_2(1 + x)), \quad x \in [0, 3], \quad n = 1, 2, \dots$$

Functions  $L_n(x)$  are orthogonal on the interval  $[0, 1]$ , and

$$\int_0^1 \frac{1}{\frac{1+x}{4} \log_2^2 \frac{1+x}{4}} L_m(x) L_n(x) dx = \frac{\delta_{mn}}{1/\ln 2(2n-1)}.$$

Also, we introduce polynomial functions  $G_n(t)$  and identify  $L_n(x)$  as compositions of  $G_n(t)$  and a logarithmic function  $t(x) = \log_2(1 + x)$ ,

$$L_n(x) \equiv G_n(t)_{t=\log_2(1+x)} \quad n = 1, 2, \dots$$

We find

$$G_n(t) = (-1)^{n-1} \left(1 - \frac{1}{2}t\right) \tilde{P}_{n-1}(t),$$

$$L_n(3) = 0.$$

It appears that  $L_n(x)$  with even indices have a zero at

$$x = \sqrt{2} - 1 = 0.414213562\dots \quad (5.1)$$

Numbers in (4.1) and (5.1) are “almost the same”. Actually, the graphs of the functions




FIG. 5.1.  $R_4(x)$  (green),  $E_4(x)$  (yellow),  $L_4(x)$  (red).

$E_n(x)$  and  $L_n(x)$  are very close on the interval of orthogonality, but they differ outside of the interval.

Functions  $L_n(x)$  form a basis in  $L^2[0, 1]$ .

**6. Conjecture.** Calculations show that the following statement may take place. Let  $r_{nk}$ ,  $e_{nk}$ , and  $l_{nk}$  be abscissas of local extrema for functions  $R_n(x)$ ,  $E_n(x)$ , and  $L_n(x)$ . Then  $R_n(r_{nk}) = E_n(e_{nk}) = L_n(l_{nk})$ .

**7. Conclusion.** Altered Legendre polynomials  $A_n(x)$  originate three sequences of orthogonal functions,  $R_n(x)$ ,  $E_n(x)$ , and  $L_n(x)$ .

The sequences are bases for approximation in  $L^2[0, 1]$  space, and they may provide spectral convergence properties. One may notice that a linear function  $s(x) = ax + b$  cannot be represented exactly by a finite linear combination of any of the three sequences, and the bases do not provide options for linear approximation.

Similar compositions for altered alternative orthogonal polynomials can be considered.

Also, one may extend definition of  $A_n(x)$  to a three-parametric family of functions

$${}_\gamma A_n^{(\alpha, \beta)}(x) = ((x + 1)/2)^\gamma \tilde{P}_{n-1}^{(\alpha, \beta)}(x), \quad \gamma > 0, \quad n > 0,$$

where  $\tilde{P}_{n-1}^{(\alpha, \beta)}(x)$  are shifted Jacobi polynomials.

#### REFERENCES

- [1] J.P. BOYD, *Orthogonal rational functions on a semi-infinite interval*, J. Comput. Phys., Vol. 70, No 1, (1987), pp. 63-88.
- [2] A. BULTHEEL, P. GONZÁLES VERA, E. HENDRIKSEN, O. NJÅSTAD, *Orthogonal rational functions*, Cambridge Monographs on Applied and Computational Mathematics, Vol. 5, Cambridge University Press, Cambridge, 1999.

- [3] A. CAYLEY, *Sur quelques propriétés des déterminants gauches*, Journal für die reine und undewandte Mathematik, 32: pp. 119-123, 1846.
- [4] V. S. CHELYSHKOV, *Sequences of exponential polynomials orthogonal on the semi-axis*, Dokl. Akad. Nauk Ukr. SSR, Ser. A, No 1, pp. 14–17, 1987.
- [5] V. S. CHELYSHKOV, *Alternative orthogonal polynomials and quadratures*, Electron. Trans. Numer. Anal., Vol. 25, pp. 17-26, 2006.
- [6] V. S. CHELYSHKOV, *Alternative Jacobi polynomials and orthogonal exponentials*, arXiv:1105.1838, 11pp., 2011.
- [7] V. CHELYSHKOV, *Alternative orthogonal rational functions on a half line*, arXiv:1504.05248, 8 pp., 2015.
- [8] V. S. CHELYSHKOV, *Spectral shape preserving approximation*, arXiv:1801.09282, 15pp., 2020.
- [9] GUO, BEN-YU; SHEN, JIE; WANG, ZHONG-QING, *Chebyshev rational spectral and pseudospectral methods on a semi-infinite interval*, Int. J. Numer. Methods Eng. 53 (1), pp. 65–84, 2002.
- [10] O. JAROCH, *Approximation by exponential functions*, Aplikace Matematiky, Vol. 7, No 4, pp. 249 – 264, 1962 (in Czech).
- [11] T.N. KOORWINDER, R. WONG, R. KOEKOEK, R.F. SWARTTOUW, *Orthogonal polynomials*, in NIST Handbook of Mathematical Functions. Cambridge University Press, 2010. (dlmf.nist.gov).
- [12] A.-M. LEGENDRE, *Recherches sur l'attraction des sphéroïdes homogènes*, Mémoires de Mathématiques. et de Physique, présentes à l'Académie Royale des Sciences, par divers savans, et las dans ses Assemblées. Vol. X. Paris, pp. 411-435, 1785 (1782).
- [13] J. VAN DEUN, A. BULTHEEL, *Orthogonal rational functions and quadrature on an interval*, Journal of computational and applied mathematics, 153: pp. 487-495, 2003.
- [14] WANG ZHONG-QING, GUO BEN-YU, *A mixed spectral method for incompressible fluid flow in an infinite strip*, Journal of Computational and Applied Mathematics, Volume 24, N.3, pp. 343-364, 2005.