

Algebraically closed σ -fields

Masood Aryapoor

Division of Mathematics and Physics

Mälardalen University

Hamngatan 15, 632 17, Eskilstuna, Sweden

Abstract

The concept of a skew root of a skew polynomial is used to introduce notions of algebraic closedness for σ -fields, that is, a field equipped with an endomorphism. It is shown that every σ -field can be embedded in algebraically closed σ -fields of different types.

1 Introduction

Skew polynomials were first studied by Ore [23]. Since Ore's article was published in 1933, skew polynomials have been the subject of study by many authors. In [16], Lam introduced the notion of a skew root of a skew polynomial in order to study a generalization of the Vandermonde matrix (see also [14]). This notion allows us to define a concept of algebraicity which has been used by Lam and Leroy to formulate and solve some problems regarding skew polynomials, see [15, 17, 18]. The main goal of this paper is to study this concept of algebraicity in the context of field extensions.

The paper is organized as follows. Section 2 deals with preliminaries and elementary problems such as interpolation for skew polynomials and Vieta's formula for skew polynomials. In Section 3, we study field extensions from the perspective of skew polynomials. It is shown that for any nonconstant skew polynomial, there is an extension field in which the given skew polynomial has a skew root. Section 4 deals with the main subject of this paper, namely, algebraically closed σ -fields. It turns out that there are different notions of "algebraic closedness" in the context of skew roots, some of which are introduced in this section, and moreover, it is shown that they exist. Finally, in Section 5, we discuss the notion of a partial fraction decomposition for skew polynomials. It is shown that an arbitrary "skew rational functions" over an algebraically closed σ -field has a relatively simple partial fraction decomposition.

We conclude the introduction with some remarks. In this paper, all rings are assumed to be associative and contain an identity element. In general, a

skew polynomial is an element of a skew polynomial ring $R[T; \sigma, \delta]$, where R is a (not necessarily commutative) ring, $\sigma : R \rightarrow R$ is an endomorphism and $\delta : R \rightarrow R$ is a σ -derivation. However, in this paper, we work exclusively with skew polynomial rings $R[T; \sigma, \delta]$ for which R is a commutative field and δ is the zero map. Given a set X , we denote the identity map of X by 1_X . For a subset X of a ring R , the notation X^* stands for $X \setminus \{0\}$.

2 Skew polynomials and skew roots

This section is devoted to preliminaries and some elementary results regarding skew polynomials and skew roots. We begin by reviewing basic facts about the ring of skew polynomials over a field, see [7, Chapter 2] for more details. Let K be a (commutative) field and $\sigma : K \rightarrow K$ be an endomorphism. The pair (K, σ) is called a σ -field. A (nonzero) skew polynomial $P(T)$ over K is a "polynomial" of the form

$$P(T) = a_0 + a_1T + \cdots + a_nT^n,$$

where $n \geq 0$ and $a_0, \dots, a_n \in K$ with $a_n \neq 0$. The number n , denoted by $\deg P$, is called the degree of $P(T)$. We usually use the word "polynomial" instead of the expression "skew polynomial" if there is no risk of confusion. The set of all skew polynomials over K is denoted by $K[T; \sigma]$. This set has a ring structure which resembles the ring of ordinary polynomials. A crucial difference is that $Ta = \sigma(a)T$ for all $a \in K$. It is known that Euclidean algorithm for right division holds $K[T; \sigma]$, a fact which has a number of consequences: $K[T; \sigma]$ is a left principal ideal domain; $K[T; \sigma]$ is a left Ore domain whose field of fractions is denoted by $K(T; \sigma)$. Elements of $K(T; \sigma)$ can be expressed as $P(T)^{-1}Q(T)$ where $0 \neq P(T), Q(T) \in K[T; \sigma]$. Besides the rings $K[T; \sigma]$ and $K(T; \sigma)$, one has the ring of "skew power series" $K[[T; \sigma]]$, and if σ is an automorphism, the skew field $K((T; \sigma))$ of Laurent series over (K, σ) whose elements are of the form

$$a_0 + a_1T + \cdots + a_nT^n + \cdots,$$

and

$$a_{-m}T^{-m} + \cdots + a_{-1}T^{-1} + a_0 + a_1T + \cdots + a_nT^n + \cdots,$$

respectively. We have the following inclusions:

$$\begin{array}{ccc} K[T; \sigma] & \subset & K[[T; \sigma]] \\ \cap & & \cap \\ K(T; \sigma) & \subset & K((T; \sigma)) \end{array}$$

If $\sigma : K \rightarrow K$ is an automorphism, then $K[T; \sigma]$ is a principal ideal domain, that is, it is both a left and a right principal ideal domain. Moreover, any $P(T) \in K[T; \sigma]$ can be written as a "right polynomial":

$$P(T) = b_0 + Tb_1 + \cdots + T^nb_n,$$

where $b_0, \dots, b_n \in K$. In fact, if $P(T)$, as a left polynomial, has the expression

$$P(T) = a_0 + a_1T + \cdots + a_nT^n,$$

then $P(T)$, as a right polynomial, has the expression

$$P(T) = a_0 + T\sigma^{-1}(a_1) + \cdots + T^n\sigma^{-n}(a_n).$$

As an example of a ring of skew polynomials, we have the ring $\mathbb{C}[T; \bar{\cdot}]$ of skew polynomials over \mathbb{C} , where the endomorphism $\bar{\cdot} : \mathbb{C} \rightarrow \mathbb{C}$ is the familiar map sending a complex number to its complex conjugate.

2.1 Preliminaries

In this subsection, we review some known facts regarding roots of skew polynomials. The main references for this part are the papers [16, 14]. Throughout this subsection, (K, σ) is a fixed σ -field.

An element $a \in K$ is called a σ -conjugate of $b \in K$ over a subset X of K if there exists $0 \neq x \in X$ such that $a = \sigma(x)bx^{-1}$. We drop the phrase "over X " when $X = K$. It is easy to see that the notion of σ -conjugacy over a subgroup X of the multiplicative group K^* is an equivalence relation in which case the σ -conjugacy class containing $a \in K$ is denoted by $C(a, X)$. When $X = K^*$, we write $C(a)$ instead of $C(a, X)$.

The maps $N_i : K \rightarrow K$, where $i \geq 0$, are defined recursively as follows

$$N_i(a) = \begin{cases} 1 & \text{if } i = 0, \\ \sigma(N_{i-1}(a))a & \text{if } i > 0. \end{cases}$$

Clearly, $N_i(a) = \sigma^{i-1}(a) \cdots \sigma(a)a$ for all $i \geq 1$. It is easy to verify the following identities:

$$N_{i+j}(a) = \sigma^i(N_j(a))N_i(a), \text{ for all } a \in K \text{ and } i, j \geq 0, \quad (2.1)$$

$$N_i(\sigma(a)ba^{-1}) = \sigma^i(a)N_i(b)a^{-1}, \text{ for all } a \in K^*, b \in K \text{ and } i \geq 0. \quad (2.2)$$

Note that $N_i(ab) = N_i(a)N_i(b)$ for all $a, b \in K$. Furthermore, if $c \in K$ is fixed by σ , then $N_i(c) = c^i$ for all $i \geq 0$.

A polynomial $P(T) = \sum_{i=0}^n a_iT^i \in K[T; \sigma]$ can be regarded as a function $P : K \rightarrow K$ as follows:

$$P(a) := \sum_{i=0}^n a_iN_i(a).$$

Note that $P(c) = \sum_{i=0}^n a_ic^i$ for all c in the fixed field of σ , that is, the field $\{c \in K \mid \sigma(c) = c\}$. The element $P(a)$ is called the value of P at a . This definition is justified by the following lemma:

Lemma 2.1. For every $P(T) \in K[T; \sigma]$ and $a \in K$, there exists a unique polynomial $Q(T) \in K[T; \sigma]$ such that

$$P(T) = Q(T)(T - a) + P(a).$$

A simple consequence of this lemma is the following:

$$P(a) = 0 \iff P(T) \in K[T; \sigma](T - a).$$

A polynomial $P(T) \in K[T; \sigma]$ is said to vanish on a subset S of K if $P(a) = 0$ for all $a \in S$. In particular, if $P(a) = 0$, where $a \in K$, the element a is called a (right skew) root of the polynomial $P(T) \in K[T; \sigma]$.

The following formula is known as the product formula:

Lemma 2.2. For all $P(T), Q(T) \in K[T; \sigma]$, we have

$$(PQ)(a) = \begin{cases} 0 & \text{if } Q(a) = 0, \\ P(\sigma(Q(a)) a Q(a)^{-1}) Q(a) & \text{if } Q(a) \neq 0, \end{cases}$$

where $(PQ)(T) = P(T)Q(T)$, as elements of $K[T; \sigma]$.

One consequence of these lemmas is that a skew polynomial of degree n has roots in at most n different σ -conjugacy classes.

A subset S of K is called σ -algebraic (or simply algebraic when there is no risk of confusion) if there exists a nonzero $P(T) \in K[T; \sigma]$ which vanishes on S . One can show that every finite subset of K is σ -algebraic. Moreover, for any σ -algebraic subset S of K , there exists a unique monic polynomial $P_S(T) \in K[T; \sigma]$ which vanishes on S , and any skew polynomial, vanishing on S , belongs to the left ideal $K[T; \sigma]P_S(T)$. The polynomial $P_S(T)$ is called the minimal (skew) polynomial of S and its degree, denoted by $rk(S)$, is called the rank of S . If S is not σ -algebraic, its rank is defined to be ∞ . An element $a \in K$ is called P-dependent on a σ -algebraic subset S of K if $P_S(a) = 0$. A subset S of K is called P-independent if no $a \in S$ is P-dependent on $S \setminus \{a\}$.

Given $a_1, \dots, a_n \in K$, the σ -Vandermonde matrices $V^\sigma(a_1, \dots, a_n) \in K^{n \times n}$ and $V_\infty^\sigma(a_1, \dots, a_n) \in K^{\infty \times n}$ are defined as follows

$$V^\sigma(a_1, \dots, a_n) = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ N_1(a_1) & N_1(a_2) & \cdots & N_1(a_n) \\ N_2(a_1) & N_2(a_2) & \cdots & N_2(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_{n-1}(a_1) & N_{n-1}(a_2) & \cdots & N_{n-1}(a_n) \end{pmatrix},$$

$$V_\infty^\sigma(a_1, \dots, a_n) = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ N_1(a_1) & N_1(a_2) & \cdots & N_1(a_n) \\ N_2(a_1) & N_2(a_2) & \cdots & N_2(a_n) \\ N_3(a_1) & N_3(a_2) & \cdots & N_3(a_n) \\ \vdots & \vdots & \cdots & \vdots \end{pmatrix}$$

The proof of the following result can be found in [16].

Theorem 2.3. *Let $a_1, \dots, a_n \in K$ be arbitrary.*

(a) *The σ -Vandermonde matrices $V^\sigma(a_1, \dots, a_n)$ and $V_\infty^\sigma(a_1, \dots, a_n)$ have equal rank, and their rank is equal to the rank of the subset $\{a_1, \dots, a_n\}$ of K . In particular, $\{a_1, \dots, a_n\}$ is P -independent iff $V^\sigma(a_1, \dots, a_n)$ is invertible.*

(b) *If $\sum_{i=1}^n b_i N_j(a_i) = 0$ for $j = 0, 1, \dots, n-1$, then $\sum_{i=1}^n b_i N_j(a_i) = 0$ for all $j \geq 0$.*

We remark that the conclusion of Part (b) holds for any skew polynomial, that is, $\sum_{i=1}^n b_i P(a_i) = 0$ for all $P(T) \in K[T; \sigma]$.

Theorem 2.3 can be used to analyze the concept of P -dependence in terms of linear dependence: Consider the K -vector space K^∞ of "infinite" row vectors over K , that is,

$$K^\infty = \{(a_0, a_1, a_2, \dots) \mid a_0, a_1, \dots \in K\}.$$

Let $N : K^* \rightarrow K^\infty$ be the map

$$N(a) = (N_0(a), N_1(a), N_2(a), \dots).$$

Using this map, we can reformulate Theorem 2.3:

Proposition 2.4. *1) A set $S \subset K^*$ is σ -algebraic over K iff the vector subspace N_S of K^∞ which is generated by all $N(a)$, where $a \in S$, is finite-dimensional. Moreover, we have $\text{rk}(S) = \dim N_S$.*

2) Elements $a_1, \dots, a_n \in K^$ are P -independent iff the vectors $N(a_1), \dots, N(a_n)$ are linearly independent over K . In particular, an element $a \in K^*$ is P -dependent on a subset S of K^* iff the row vector $N(a)$ belongs to the vector space N_S .*

The vector space K^∞ can be considered as a (commutative) K -algebra using the multiplication

$$(a_0, a_1, a_2, \dots)(b_0, b_1, b_2, \dots) = (a_0 b_0, a_1 b_1, a_2 b_2, \dots).$$

Since each $N_i : K \rightarrow K$ satisfies the identity

$$N_i(ab) = N_i(a)N_i(b), \text{ where } a, b \in K,$$

we see that N_{K^*} is a subalgebra of the K -algebra K^∞ . More generally, N_S is a subalgebra of the K -algebra K^∞ if $S \subset K^*$ is a multiplicative subset of K , that is, $1 \in S$, and $ab \in S$ when $a, b \in S$.

2.2 Left and right roots

In this part, we assume that $\sigma : K \rightarrow K$ is a field automorphism. As mentioned at the beginning of this section, elements of $K[T; \sigma]$ can also be considered as right polynomials. As a consequence, one has the "left" analogues of the concepts and results of the preceding subsection. More precisely, any polynomial $P(T) = \sum_{i=0}^n a_i T^i \in K[T; \sigma]$ gives rise to a left evaluation map $P_l : K \rightarrow K$ defined by

$$P_l(a) = \sum_{i=0}^n \sigma^{-i}(a_i) N_{-i}(a),$$

where $N_{-i}(a) = \sigma^{-i+1}(a) \cdots \sigma^{-1}(a)a$ for $a \in K$ and $i \geq 1$. Moreover, the "left" version of Lemma 3.8 holds. More precisely, for every polynomial $P(T) \in K[T; \sigma]$ and $a \in K$, there exists a unique polynomial $Q(T) \in K[T; \sigma]$ such that

$$P(T) = (T - a)Q(T) + P_l(a).$$

In particular, $P(T) \in (T - a)K[T; \sigma]$ iff $P_l(a) = 0$ in which case a is called a left (skew) root of $P(T)$. We leave it to the reader to define and prove the left analogues of the concepts and results of the preceding subsection. We shall use the prefixes "left" and "right" to distinguish between the concepts defined using P_l and the ones presented in the preceding subsection. However, we usually drop the prefix "right" when there is no risk for confusion. The following proposition gives a relationship between the concepts of left P-dependence and right P-dependence.

Proposition 2.5. *For every $a_1, \dots, a_n \in K^*$, the following are equivalent:*

- (1) *The right rank of the set $\{a_1, \dots, a_n\}$ is less than n , or equivalently, the elements a_1, \dots, a_n are right P-dependent.*
- (2) *There exist $b_1, \dots, b_n \in K$, not all equal to zero, such that*

$$b_1(1 - a_1 T)^{-1} + \cdots + b_n(1 - a_n T)^{-1} = 0.$$

- (3) *There exist $b_1, \dots, b_n \in K$, not all equal to zero, such that*

$$b_1(1 - a_1 T)^{-1} + \cdots + b_n(1 - a_n T)^{-1} \in K[T; \sigma].$$

- (4) *The left rank of the set $\{\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_n^{-1})\}$ is less than n , or equivalently, the elements $\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_n^{-1})$ are left P-dependent.*

Proof. The equivalence of (1) and (2) follows from Theorem 2.3 and the fact that $(1 - aT)^{-1}$, as an element of $K[[T; \sigma]]$, is the series

$$(1 - aT)^{-1} = \sum_{i=0}^{\infty} N_i(a) T^i.$$

Clearly, (2) implies (3). To prove (3) \implies (4), let there be $b_1, \dots, b_n \in K$, not all equal to zero, such that

$$b_1(1 - a_1T)^{-1} + \dots + b_n(1 - a_nT)^{-1} = P(T) \in K[T; \sigma].$$

Without loss of generality, we may assume that $b_n \neq 0$. This equality can be rewritten as follows

$$b_1\sigma^{-1}(a_1^{-1})(\sigma^{-1}(a_1^{-1}) - T)^{-1} + \dots + b_n\sigma^{-1}(a_n^{-1})(\sigma^{-1}(a_n^{-1}) - T)^{-1} = P(T).$$

Let $Q(T) \in K[T; \sigma]$ have $\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_n^{-1})$ as left zeros, that is,

$$Q_l(\sigma^{-1}(a_i^{-1})) = 0,$$

for $1 \leq i < n$. It follows that

$$\begin{aligned} b_n\sigma^{-1}(a_n^{-1})(\sigma^{-1}(a_n^{-1}) - T)^{-1}Q(T) &= P(T)Q(T) \\ &+ \sum_{i=1}^{n-1} b_i\sigma^{-1}(a_i^{-1})(T - \sigma^{-1}(a_i^{-1}))^{-1}Q(T) \end{aligned}$$

belongs to $K[T; \sigma]$. Therefore, we see that $Q_l(\sigma^{-1}(a_n^{-1})) = 0$. Therefore, we conclude that $\sigma^{-1}(a_n^{-1})$ is left P-dependent on the set $\{\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_{n-1}^{-1})\}$. In particular, the left rank of the set $\{\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_n^{-1})\}$ is less than n . This completes the proof of (3) \implies (4). It remains to prove (4) \implies (2). Let $\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_n^{-1})$ be left P-dependent. Without loss of generality, we may assume that $\sigma^{-1}(a_n^{-1})$ is P-dependent on the set $\{\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_{n-1}^{-1})\}$. Let $Q(T) \in K[T; \sigma]$ be the minimal left polynomial of $\{\sigma^{-1}(a_1^{-1}), \dots, \sigma^{-1}(a_{n-1}^{-1})\}$. Note that the degree of $Q(T)$ is less than n . This fact, together with the fact that

$$(\sigma^{-1}(a_n^{-1}) - T)^{-1}Q(T) \in K[T; \sigma],$$

for all $1 \leq i \leq n$, imply that there are $b_1, \dots, b_n \in K$, not all equal to zero, such that

$$b_1(\sigma^{-1}(a_1^{-1}) - T)^{-1}Q(T) + \dots + b_n(\sigma^{-1}(a_n^{-1}) - T)^{-1}Q(T) = 0.$$

Since $K(T; \sigma)$ is a skew field, it follows that

$$b_1(\sigma^{-1}(a_1^{-1}) - T)^{-1} + \dots + b_n(\sigma^{-1}(a_n^{-1}) - T)^{-1} = 0,$$

or equivalently,

$$b_1\sigma^{-1}(a_1)(1 - a_1T)^{-1} + \dots + b_n\sigma^{-1}(a_n)(1 - a_nT)^{-1} = 0,$$

and we are done. \square

As a consequence of this proposition, we have

Corollary 2.6. *A set $S \subset K^*$ is right σ -algebraic iff the set*

$$\widehat{S} = \{\sigma^{-1}(a^{-1}) \mid a \in S\},$$

is left σ -algebraic. Furthermore, the right rank of an arbitrary subset S of K^ is equal to the left rank of \widehat{S} .*

In the case when σ is an involution, that is, $\sigma^2 = 1_K$, we have

$$\widehat{S} = \{\sigma(a^{-1}) \mid a \in S\}.$$

As an example, a set $S \subset \mathbb{C}^*$ is right σ -algebraic iff the set

$$\widehat{S} = \left\{ \frac{1}{\bar{c}} \mid c \in S \right\},$$

is left σ -algebraic. Here, $\sigma : \mathbb{C} \rightarrow \mathbb{C}$ is the complex conjugation map.

Although Proposition 2.5 gives a relationship between left and right ranks, it does not provide any relationship between minimal left and right polynomials. In order to derive such a relationship, we need the following result which deals with anti-automorphisms of rings of skew polynomials. Recall that a map $\alpha : R \rightarrow R$, where R is a ring, is called an anti-automorphism, if it is an additive group isomorphism, $\alpha(1) = 1$, and $\alpha(ab) = \alpha(b)\alpha(a)$ for all $a, b \in R$. In the following proposition, we give a characterization of anti-automorphisms of $K[T; \sigma]$. For the case of automorphisms of skew polynomial rings over (not necessarily commutative) rings, see [5].

Proposition 2.7. *Suppose that $\sigma : K \rightarrow K$ is an automorphism not equal to the identity map 1_K . Let $\alpha : K[T; \sigma] \rightarrow K[T; \sigma]$ be a map. If α is an anti-automorphism, then the following hold:*

- (1) $\alpha(K) = K$, and the restriction α_0 of α to K is an automorphism of K .
- (2) There exists an element $a_0 \in K^*$ such that $\alpha(T) = a_0T$.
- (3) $\sigma\alpha_0\sigma = \alpha_0$.

Conversely, if $\alpha_0 : K \rightarrow K$ is an automorphism of K which satisfies Property (3), then, for every $a_0 \in K^$, there exists a unique anti-automorphism $\alpha : K[T; \sigma] \rightarrow K[T; \sigma]$, with $\alpha(T) = a_0T$, whose restriction to K is α_0 .*

Proof. Let $\alpha : K[T; \sigma] \rightarrow K[T; \sigma]$ be an anti-automorphism. First, we prove (1). Since the set of units of $K[T; \sigma]$ is K^* , we see that $\alpha(K) \subset K$. Similarly, we have $\alpha^{-1}(K) \subset K$ since α^{-1} is an anti-automorphism of $K[T; \sigma]$. It follows that $\alpha(K) = K$. Clearly, $\alpha_0 : K \rightarrow K$ is an automorphism of K since K is commutative. This completes the proof of (1). A simple argument using the degree function reveals that $\alpha(T) = a_0T + b_0$ for some $a_0, b_0 \in K$ with $a_0 \neq 0$. For $a \in K$, we have

$$Ta = \sigma(a)T \implies \alpha(Ta) = \alpha(\sigma(a)T) \implies \alpha_0(a)\alpha(T) = \alpha(T)\alpha_0(\sigma(a)),$$

which in turn implies that

$$\alpha_0(a)a_0T + \alpha_0(a)b_0 = a_0T\alpha_0(\sigma(a)) + b_0\alpha_0(\sigma(a)) = a_0\sigma(\alpha_0(\sigma(a)))T + b_0\alpha_0(\sigma(a)).$$

It follows that

$$\forall a \in K, \alpha_0(a)a_0 = a_0\sigma(\alpha_0(\sigma(a))) \text{ and } \alpha_0(a)b_0 = b_0\alpha_0(\sigma(a)).$$

The first relation proves (3) because $a_0 \neq 0$. If $b_0 \neq 0$, then α_0 must be equal to $\alpha_0\sigma$ which implies that $\sigma = 1_K$ since α is a bijection. This contradicts the assumption that $\sigma \neq 1_K$. This completes the proof of (2).

To prove the second statement, let $\alpha_0 : K \rightarrow K$ be an automorphism satisfying (1) and (3). Given $a_0 \in K^*$, we define $\alpha : K[T; \alpha] \rightarrow K[T; \alpha]$, as follows

$$\alpha\left(\sum_{i=0}^n b_i T^i\right) = \sum_{i=0}^n \sigma^i(\alpha_0(b_i)) N_i(a_0) T^i.$$

It is easy to see that α is an additive group isomorphism. Since $\alpha(1) = 1$, $\alpha(T) = a_0T$, and the restriction of α to K is α_0 , we only need to prove that

$$\alpha(P(T)Q(T)) = \alpha(Q(T))\alpha(P(T)),$$

for all $P(T), Q(T) \in K[T; \sigma]$. By the additivity of α , it is enough to show that

$$\alpha((aT^m)(bT^n)) = \alpha(bT^n)\alpha(aT^m),$$

for all $a, b \in K$ and $m, n \geq 0$. We have

$$\alpha((aT^m)(bT^n)) = \alpha(a\sigma^m(b)T^{m+n}) = \sigma^{m+n}(\alpha_0(a\sigma^m(b))) N_{m+n}(a_0) T^{m+n},$$

and

$$\begin{aligned} \alpha(bT^n)\alpha(aT^m) &= \sigma^n(\alpha_0(b)) N_n(a_0) T^n \sigma^m(\alpha_0(a)) N_m(a_0) T^m \\ &= \sigma^n(\alpha_0(b)) N_n(a_0) \sigma^n(\sigma^m(\alpha_0(a))) N_m(a_0) T^{m+n}. \end{aligned}$$

Therefore, we need to show that

$$\sigma^{m+n}(\alpha_0(a\sigma^m(b))) N_{m+n}(a_0) = \sigma^n(\alpha_0(b)) N_n(a_0) \sigma^n(\sigma^m(\alpha_0(a))) N_m(a_0),$$

for all $a, b \in K$ and $m, n \geq 0$. Using Equation 2.1 and the fact that σ and α_0 are ring homomorphisms, we can simplify this equation to obtain

$$\sigma^{m+n}(\alpha_0(\sigma^m(b))) = \sigma^n(\alpha_0(b)).$$

It is now easy to use $\sigma\alpha_0\sigma = \alpha_0$ to show that this relation holds for all $b \in K$ and $m, n \geq 0$. The uniqueness part follows from the fact that K and T generate $K[T; \sigma]$ as a ring. This completes the proof of the second statement. \square

Let $AAut(K, \sigma)$ denote the set of all anti-automorphisms of $K[T; \sigma]$ and $Aut_\sigma(K)$ denote the set of all automorphism $\alpha_0 : K \rightarrow K$ such that $\sigma\alpha_0\sigma = \alpha_0$. Using Proposition 2.7, we obtain a 1-1 correspondence between $AAut(K, \sigma)$ and the set $Aut_\sigma(K) \times K^*$, given by $\alpha \mapsto (\alpha|_K, \alpha(T)T^{-1})$. It may happen that $Aut_\sigma(K)$ is empty, in which case $AAut(K, \sigma)$ is empty as well. As an example, the reader can verify that if K is any cyclic extension of the field of rational numbers, and σ is a generator of the Galois group of K over \mathbb{Q} , then $AAut(K, \sigma)$ is nonempty iff $\sigma^2 = 1_K$ (in the case $\sigma^2 = 1_K$, we have $1_K \in Aut_\sigma(K)$). If $AAut(K, \sigma)$ is nonempty, then we have the following result concerning left and right roots:

Proposition 2.8. *Let $\alpha \in AAut(K, \sigma)$ satisfy $\alpha(T) = T$, and set $\alpha_0 = \alpha|_K$. Then, the following hold:*

- (1) *$a \in K$ is a left root of a polynomial $P(T) \in K[T; \sigma]$ iff $\alpha_0(a)$ is a right root of $\alpha(P(T))$.*
- (2) *A polynomial $P(T) \in K[T; \sigma]$ is the minimal left polynomial of a subset of K iff $\alpha(P(T))$ is the minimal right polynomial of a subset of K .*
- (3) *The assignment $S \mapsto \alpha(S)$ establishes a 1-1 correspondence between the set of all left σ -algebraic subsets of K and the set of all right σ -algebraic subsets of K . Moreover, the left rank of S is equal to the right rank of $\alpha(S)$.*

Proof. (1) If a is a left root of $P(T) \in K[T; \sigma]$, then

$$P(T) = (T - a)Q(T),$$

for some $Q(T) \in K[T; \sigma]$. Applying α to this relation gives

$$\alpha(P(T)) = \alpha((T - a)Q(T)) = \alpha(Q(T))\alpha(T - a) = \alpha(Q(T))(T - \alpha_0(a)),$$

showing that $\alpha_0(a)$ is a right root of $\alpha(P(T))$. The converse is proved in a similar fashion. Now, Parts (2) and (3) follow easily from (1). \square

In the case when σ is an involution, the identity map of $K[T; \sigma]$ belongs to $AAut(K, \sigma)$. In this case, an application of this proposition gives the following: (1) a is a left root of $\sum_{i=0}^n a_i T^i$ iff a is a right root of $\sum_{i=0}^n T^i a_i$; (2) $S \subset K$ is left σ -algebraic iff it is right σ -algebraic, and moreover, the left rank of S is equal to its right rank.

2.3 Values of skew polynomials

In this subsection, we present some elementary results concerning values of skew polynomials. As opposed to ordinary polynomials in an indeterminate over commutative fields, nonzero skew polynomials over commutative fields may have infinitely many roots. For example, the polynomial

$$T^2 - 1 \in \mathbb{C}[T; \bar{\cdot}],$$

vanishes on the unit circle $\{z \in \mathbb{C} \mid |z| = 1\}$. However, we have the following result, which is well-known for ordinary polynomials over commutative fields. The result also holds for polynomials in a central indeterminate over a skew field (see [13, Theorem 16.7]).

Proposition 2.9. *Let (K, σ) be an infinite σ -field and $P(T) \in K[T; \sigma]$. If $P(a) = 0$ for all $a \in K$, then $P(T) = 0$.*

Proof. Let F be the fixed field of σ . Suppose, on the contrary, that a skew polynomial $P(T) = \sum_{i=0}^n a_i T^i \neq 0$ vanishes on K . Then, all the σ -conjugacy classes of K must be σ -algebraic. It follows from [18, Theorem 5.17] that the degree $[K : F]$ of the extension K/F is finite. Therefore, F must be an infinite field. Since $P(c) = \sum_{i=0}^n a_i c^i$, where $c \in F$, the nonzero ordinary polynomial $\sum_{i=0}^n a_i x^i \in K[x]$, must have infinitely many zeros over the commutative field K , a contradiction. \square

This theorem does not hold for finite σ -fields: Since any finite subset of a σ -field is σ -algebraic, we see that there is a nonzero polynomial $P(T) \in K[T; \sigma]$ vanishing on K if (K, σ) is a finite σ -field, in which case, the nonconstant polynomial $P(T) + 1$ has no roots in K . We note in passing that if (K, σ) is a finite σ -field, then the set of all $P(T) \in K[T; \sigma]$ which vanish on K is a nonzero (two-sided) ideal of $K[T; \sigma]$.

Next, we prove a version of the Bray–Whaples theorem for skew polynomials, see [4] for the result which deals with polynomials in a central indeterminate over a skew field.

Proposition 2.10. *Let $a_1, \dots, a_n \in K$ be mutually nonconjugate over K . Then, there is a unique monic skew polynomial $P(T) \in K[T; \sigma]$ of degree n whose roots are exactly a_1, \dots, a_n . Moreover, if*

$$Q(a_1) = \dots = Q(a_n) = 0,$$

where $Q(T) \in K[T; \sigma]$, then $Q(T) \in K[T; \sigma]P(T)$.

Proof. Let $P(T) \in K[T; \sigma]$ be the minimal skew polynomial of $\{a_1, \dots, a_n\}$. Since a_1, \dots, a_n are mutually nonconjugate, the degree of $P(T)$ must be n . Therefore, it remains to show that $P(T)$ has no roots other than a_1, \dots, a_n . This holds trivially if $n = 1$. So, let $n > 1$. Suppose, on the contrary, that there exists $b \in K$ not equal to a_1, \dots, a_n such that $P(b) = 0$. By Lemma 2.1, we have

$$P(T) = Q(T)(T - b),$$

for some $Q(T) \in K[T; \sigma]$. Since $b \neq a_i$ for all i , it follows from Lemma 2.2 that the skew polynomial $Q(T)$ has the following roots

$$c_i = \sigma(a_i - b)a_i(a_i - b)^{-1} \text{ where } i = 1, \dots, n.$$

Clearly, each c_i is conjugate to a_i . Therefore, the roots c_1, \dots, c_n of $Q(T)$ are mutually nonconjugate, contradicting the fact that a skew polynomial of degree less than n cannot have roots in n different conjugacy classes. The second statement follows from the fact that $P(T)$ is the minimal polynomial of $\{a_1, \dots, a_n\}$. \square

In general, the classical Lagrange interpolation does not work for skew polynomials since an element $a \in K$ can be P-dependent on a subset not containing a (see the discussion after Proposition 2.11). However, we have the following version of interpolation for skew polynomials, which can be regarded as a generalization of Proposition 2.10. The problem of interpolation for polynomials in a central indeterminate over a skew field is treated in [3].

Proposition 2.11. *Let $S = \{a_1, \dots, a_n\} \subset K$ be P-independent. For any $b_1, \dots, b_n \in K$, there exists a unique skew polynomial $P(T) \in K[T; \sigma]$ of degree less than $n = rk(S)$ such that $P(a_i) = b_i$ for all i . Moreover, a polynomial $Q(T) \in K[T; \sigma]$ satisfies $Q(a_i) = b_i$ for all i , iff*

$$Q(T) \in P(T) + K[T; \sigma]P_S(T).$$

Proof. We note that $rk(S) = n$ by [16, Lemma 12]. Let $P_i(T) \in K[T; \sigma]$ be the minimal skew polynomial of $S \setminus \{a_i\}$. The degree of each $P_i(T)$ is $rk(S) - 1 = n - 1$. Since a_1, \dots, a_n are P-independent, $P_i(a_i)$ cannot be zero. It is easy to verify that the polynomial

$$P(T) = \sum_{i=1}^n b_i P_i(a_i)^{-1} P_i(T),$$

satisfies the desired property. The uniqueness of $P(T)$ follows from the fact that the degree of each $P_i(T)$ is less than $rk(S)$. \square

A consequence of this proposition is that if $a \in K$ is P-dependent on a P-independent set $S = \{a_1, \dots, a_n\}$, then for any $Q(T) \in K[T; \sigma]$, the value of $Q(a)$ is completely determined by the values $Q(a_1), \dots, Q(a_n)$. In fact, we have

$$Q(a) = \sum_{i=1}^n Q(a_i) P_i(a_i)^{-1} P_i(a),$$

where $P_i(T)$ is the minimal skew polynomial of $S \setminus \{a_i\}$. More generally, one can easily show that the value of $Q(a)$ is completely determined by the values $Q(a), a \in S$, if a is P-dependent on a (not necessarily P-independent) σ -algebraic set $S \subset K$.

2.4 Vieta's formulas for skew polynomials

The classical formulas of Vieta express the coefficients of an ordinary polynomial over a commutative field in terms of its roots. Here, we present analogous formulas for skew polynomials. The formulas in the second part of the following result can be regarded as analogues of Vieta's formulas (see [10], [11, Section 7.2] and [12] for a more general discussion).

Theorem 2.12. *Let $S = \{a_1, \dots, a_n\}$ be P -independent with minimal polynomial $P_S(T) = \sum_{i=0}^n b_i T^i \in K[T; \sigma]$.*

(a) *For every $a \in K$, we have*

$$P_S(a) = \frac{\det(V^\sigma(a_1, \dots, a_n, a))}{\det(V^\sigma(a_1, \dots, a_n))}.$$

(b) *For $i = 0, \dots, n$,*

$$b_i = \frac{(-1)^{n+i}}{\det(V^\sigma(a_1, \dots, a_n))} \begin{vmatrix} 1 & 1 & \cdots & 1 \\ N_1(a_1) & N_1(a_2) & \cdots & N_1(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_{i-1}(a_1) & N_{i-1}(a_2) & \cdots & N_{i-1}(a_n) \\ N_{i+1}(a_1) & N_{i+1}(a_2) & \cdots & N_{i+1}(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_n(a_1) & N_n(a_2) & \cdots & N_n(a_n) \end{vmatrix} \quad (2.3)$$

Moreover, $b_0 = (-1)^n \frac{\sigma(\det(V^\sigma(a_1, \dots, a_n)))}{\det(V^\sigma(a_1, \dots, a_n))} a_1 a_2 \cdots a_n$.

Proof. Using cofactor expansion along the last column of the matrix $V^\sigma(a_1, \dots, a_n, a)$, we obtain

$$\det(V^\sigma(a_1, \dots, a_n, a)) = \sum_{i=0}^n c_i N_i(a),$$

where

$$c_i = (-1)^{n+i} \begin{vmatrix} 1 & 1 & \cdots & 1 \\ N_1(a_1) & N_1(a_2) & \cdots & N_1(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_{i-1}(a_1) & N_{i-1}(a_2) & \cdots & N_{i-1}(a_n) \\ N_{i+1}(a_1) & N_{i+1}(a_2) & \cdots & N_{i+1}(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_n(a_1) & N_n(a_2) & \cdots & N_n(a_n) \end{vmatrix}.$$

Setting $Q(T) = \sum_{i=0}^n c_i T^i$, we have

$$Q(a) = \sum_{i=0}^n c_i N_i(a) = \det(V^\sigma(a_1, \dots, a_n, a)),$$

for all $a \in K$. By Part (a) of Theorem 2.3, $Q(a_i) = 0$ for all $i = 1, \dots, n$. It follows that

$$Q(T) \in K[T; \sigma]P_S(T).$$

Since $P_S(T)$ and $Q(T)$ have the same degree, and $c_n = \det(V^\sigma(a_1, \dots, a_n))$, we conclude that

$$Q(T) = V^\sigma(a_1, \dots, a_n)P_S(T),$$

and

$$P_S(a) = \frac{\det(V^\sigma(a_1, \dots, a_n, a))}{\det(V^\sigma(a_1, \dots, a_n))},$$

for all $a \in K$.

(b) Formula 2.3 follows from (a). To prove the last formula, we set $i = 0$ in Formula 2.3 and use Identity 2.1. We see that $(-1)^n b_0 \det(V^\sigma(a_1, \dots, a_n))$ is equal to

$$\begin{aligned} & \begin{vmatrix} N_1(a_1) & N_1(a_2) & \cdots & N_1(a_n) \\ N_2(a_1) & N_2(a_2) & \cdots & N_2(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_n(a_1) & N_n(a_2) & \cdots & N_n(a_n) \end{vmatrix} \\ = & \begin{vmatrix} a_1 & a_2 & \cdots & a_n \\ a_1\sigma(N_1(a_1)) & a_2\sigma(N_1(a_2)) & \cdots & a_n\sigma(N_1(a_n)) \\ a_1\sigma(N_2(a_1)) & a_2\sigma(N_2(a_2)) & \cdots & a_n\sigma(N_2(a_n)) \\ \vdots & \vdots & \cdots & \vdots \\ a_1\sigma(N_{n-1}(a_1)) & a_2\sigma(N_{n-1}(a_2)) & \cdots & a_n\sigma(N_{n-1}(a_n)) \end{vmatrix} \\ = & \begin{vmatrix} 1 & 1 & \cdots & 1 \\ \sigma(N_1(a_1)) & \sigma(N_1(a_2)) & \cdots & \sigma(N_1(a_n)) \\ \sigma(N_2(a_1)) & \sigma(N_2(a_2)) & \cdots & \sigma(N_2(a_n)) \\ \vdots & \vdots & \cdots & \vdots \\ \sigma(N_{n-1}(a_1)) & \sigma(N_{n-1}(a_2)) & \cdots & \sigma(N_{n-1}(a_n)) \end{vmatrix} a_1 a_2 \cdots a_n \\ = & \sigma \left(\begin{vmatrix} 1 & 1 & \cdots & 1 \\ N_1(a_1) & N_1(a_2) & \cdots & N_1(a_n) \\ N_2(a_1) & N_2(a_2) & \cdots & N_2(a_n) \\ \vdots & \vdots & \cdots & \vdots \\ N_{n-1}(a_1) & N_{n-1}(a_2) & \cdots & N_{n-1}(a_n) \end{vmatrix} \right) a_1 a_2 \cdots a_n \\ = & \sigma(\det(V^\sigma(a_1, \dots, a_n))) a_1 a_2 \cdots a_n, \end{aligned}$$

which proves the desired formula. \square

We note that Part (a) of this theorem provides a method for finding the minimal polynomial of any finite P-independent set. In particular, one can apply this method to find the polynomials $P_i(T)$'s used in the proof of Proposition 2.11.

3 σ -Extensions

This section deals with the notion of σ -algebraicity in the context of σ -extensions. Since it is more convenient to use the language of difference rings, we begin by reviewing basic facts about difference rings. For a detailed account of the theory of difference rings, we refer the reader to the book [20].

By a difference ring, we mean a pair (R, σ_R) where R is a commutative ring and $\sigma_R : R \rightarrow R$ is an injective endomorphism. The ring R is called the underlying ring of the difference ring (R, σ) and the endomorphism σ_R is called its structure map. A difference ring is also called a σ -ring, a term we mainly use in this paper. By abuse of notation, we often write σ in place of σ_R if there is no risk of confusion. Note that any ring equipped with its identity map gives rise to a σ -ring which we call a trivial σ -ring. In what follows, we present some definitions and facts regarding difference rings which are relevant to this work.

- (1) A σ -ring is called a σ -field (resp., σ -domain) if its underlying ring is a field (resp., an integral domain).
- (2) By a σ -ideal of a σ -ring (R, σ) , we mean an ideal I of R which satisfies

$$\forall r \in R, r \in I \iff \sigma(r) \in I.$$

For any σ -ideal I of a σ -ring (R, σ) , we can consider a σ -ring structure on R/I defined by $r + I \mapsto \sigma(r) + I$, which is called the quotient of (R, σ) modulo I , and denoted by $(R/I, \sigma)$.

- (3) A σ -homomorphism $\phi : (R, \sigma_R) \rightarrow (S, \sigma_S)$ between σ -rings is a ring homomorphism of $\phi : R \rightarrow S$ between the underlying rings which satisfies $\phi \sigma_R = \sigma_S \phi$. It is easy to see that σ -rings and σ -homomorphisms form a category. If I is a σ -ideal of (R, σ) , then the quotient map $q : R \rightarrow R/I$ gives rise to a σ -homomorphism $q : (R, \sigma) \rightarrow (R/I, \sigma)$. If $\phi : (R, \sigma_R) \rightarrow (S, \sigma_S)$ is a σ -homomorphism, then $\ker \phi$ is a σ -ideal of (R, σ) , and the map $R/\ker \phi \rightarrow S$ is an (injective) σ -homomorphism.

- (4) By a σ -subring of a σ -ring (R, σ) , we mean a subring S of R which satisfies $\sigma(S) \subset S$, in which case, the inclusion map becomes a σ -homomorphism $i : (S, \sigma) \rightarrow (R, \sigma)$. In particular, if both S and R are fields, (S, σ) is called a σ -subfield of (R, σ) , and (R, σ) is called an *extension σ -field* (or simply *σ -extension*) of (S, σ) .

- (5) Let (S, σ) be a σ -subring of a σ -ring (R, σ) . For a subset X of (R, σ) , the intersection of all σ -subrings of (R, σ) containing $S \cup X$, is itself a σ -subring of (R, σ) which is called the σ -subring of (R, σ) generated by X over (S, σ) , and denoted by $S[X]_\sigma$. In case, $X = \{x_1, \dots, x_n\}$ is finite, we denote $S[X]_\sigma$ by $S[x_1, \dots, x_n]_\sigma$.

- (6) Let (K, σ) be a σ -subfield of a σ -field (L, σ) . For a subset X of (L, σ) , the intersection of all σ -subfields of (L, σ) containing $K \cup X$ is itself a σ -field of (L, σ) which is called the σ -subfield of (L, σ) generated by X over (K, σ) , and denoted by $K\langle X \rangle$. In case, $X = \{x_1, \dots, x_n\}$ is finite, we denote $K\langle X \rangle$

by $K\langle x_1, \dots, x_n \rangle$.

(7) A σ -ring (R, σ) is called *inversive* if $\sigma : R \rightarrow R$ is onto. It is known that for any σ -ring (R, σ) , there exists a σ -ring (U, σ) containing (R, σ) as a σ -subring such that any σ -homomorphism $\psi : (R, \sigma) \rightarrow (S, \sigma_S)$, where (S, σ_S) is inversive, has a unique extension to a σ -homomorphism $(U, \sigma) \rightarrow (S, \sigma_S)$. The σ -ring (U, σ) , called a *universal inversive closure* of (R, σ) , is unique up to σ -isomorphism. Moreover, (U, σ) has the following property:

$$\forall u \in U, \exists n \geq 1, \sigma^n(u) \in R.$$

Also, (U, σ) is a σ -field (resp., σ -domain) if (R, σ) is a σ -field (resp., σ -domain).

(8) A multiplicative σ -subset of a σ -ring (R, σ) is a multiplicative subset S of R which satisfies

$$\forall r \in R, (r \in S \implies \sigma(r) \in S).$$

Given a multiplicative σ -subset of a σ -domain (R, σ) , the map

$$rs^{-1} \mapsto \sigma(r)\sigma(s)^{-1},$$

is a well-defined map on the localization ring $S^{-1}R$, turning $S^{-1}R$ into a σ -domain. Moreover, the ring homomorphism

$$R \rightarrow S^{-1}R, r \mapsto r1^{-1},$$

is a σ -homomorphism. In particular, if (R, σ) is a σ -domain, then the quotient field $\text{Frac}(R)$ of R acquires a σ -field structure for which the inclusion map $R \rightarrow \text{Frac}(R)$ is a σ -homomorphism. The resulting σ -field $(\text{Frac}(R), \sigma)$ is called the σ -field of fractions of (R, σ) .

(9) Let (R, σ) be a σ -field. Consider the ring

$$R\{x\} := R[x(0), x(1), x(2), \dots],$$

of polynomials over R in (commuting) indeterminates $x(0), x(1), \dots$. There is a unique endomorphism $\sigma : R\{x\} \rightarrow R\{x\}$ which extends $\sigma : R \rightarrow R$ and satisfies $\sigma(x(i)) = x(i+1)$ for $i \geq 0$. Clearly, $\sigma : R\{x\} \rightarrow R\{x\}$ is injective. Therefore, the pair $(R\{x\}, \sigma)$ defines a σ -ring which includes (R, σ) as a σ -subring. The σ -ring $(R\{x\}, \sigma)$ is called the *ring of difference polynomials* in the difference indeterminate x over (R, σ) . It has the following familiar property: For any σ -homomorphism $\phi : (R, \sigma) \rightarrow (S, \sigma_S)$ and any element $s \in S$, there exists a unique σ -homomorphism $\phi : (R\{x\}, \sigma) \rightarrow (S, \sigma_S)$ which extends $\phi : (R, \sigma) \rightarrow (S, \sigma_S)$ and sends $x(0)$ to s .

3.1 σ -Algebraic elements in σ -extensions

In this subsection, we let (L, σ) be an extension σ -field of a σ -field (K, σ) . An element $a \in L$ is called σ -algebraic over K (or algebraic over (K, σ)) if a is a root of a polynomial $P(T) \in K[T; \sigma]$. The following lemma gives a criterion for an element $a \in L$ to be algebraic over (K, σ) .

Lemma 3.1. *For any element $a \in L$, the following are equivalent:*

- (1) a is σ -algebraic over K .
- (2) The K -vector subspace of L generated by the elements $N_i(a)$, $i \geq 0$, is finite-dimensional.
- (3) There exists a finite-dimensional K -vector subspace V of L such that $1 \in V$ and $a\sigma(V) \subset V$.

Proof. (1) \implies (2): Let $a \in L$ be σ -algebraic over K . Then, we have

$$c_0 + c_1 N_1(a) + \cdots + c_n N_n(a) = 0,$$

for some $c_0, \dots, c_n \in K$ with $c_n \neq 0$. Applying σ to this equation and then multiplying with a , we obtain

$$\sigma(c_0)a + \sigma(c_1)a\sigma(N_1(a)) + \cdots + \sigma(c_n)a\sigma(N_n(a)) = 0.$$

Since $a\sigma(N_i(a)) = N_{i+1}(a)$, this equation can be rewritten as

$$\sigma(c_0)N_1(a) + \sigma(c_1)N_2(a) + \cdots + \sigma(c_n)N_{n+1}(a) = 0,$$

proving that $N_{n+1}(a)$ is a K -linear combination of the elements

$$N_0(a), N_1(a), \dots, N_n(a).$$

Inductively, one can see that this holds for all $N_i(a)$, where $i \geq n + 1$. In particular, the K -vector subspace of L generated by the elements $N_i(a)$, $i \geq 0$, is finite-dimensional.

(2) \implies (3): We show that the finite-dimensional K -vector subspace V of L generated by $N_i(a)$, $i \geq 0$, satisfies the desired conditions. Clearly, $1 = N_0(a) \in V$. Using the formula $N_{i+1}(a) = a\sigma(N_i(a))$, one can easily see that $a\sigma(V) \subset V$, and we are done.

(3) \implies (1): Let V be a finite-dimensional K -vector subspace of L which contains 1 and satisfies $a\sigma(V) \subset V$. Let n be the dimension of V . Since $1 \in V$ and $a\sigma(V) \subset V$, the identity $N_{i+1}(a) = a\sigma(N_i(a))$ implies that $N_i(a) \in V$ for all $i \geq 0$. In particular, $N_0(a), \dots, N_n(a)$ are linearly dependent over K , that is, a is σ -algebraic over K . \square

Now, we give some applications of this lemma.

Proposition 3.2. *Let $b \in L$. There is $x \in L^*$ such that $\sigma(x)bx^{-1}$ is algebraic over (K, σ) iff there exists a nonzero finite-dimensional K -vector subspace W of L such that $b\sigma(W) \subset W$.*

Proof. Let $a = \sigma(x)bx^{-1}$ be algebraic over (K, σ) for some $0 \neq x \in L$. By Lemma 3.1, there exists a finite-dimensional K -vector subspace V of L such that $1 \in V$ and $a\sigma(V) \subset V$. We have

$$\sigma(x)bx^{-1}\sigma(V) \subset V \implies b\sigma(xV) \subset xV.$$

Therefore, the finite-dimensional K -vector space $W = xV$ satisfies the desired condition.

Conversely, let $b\sigma(W) \subset W$ for some finite-dimensional K -vector subspace W of L . Since W is nonzero, W contains a nonzero element x . Setting $V = x^{-1}W$, we have

$$(\sigma(x)bx^{-1})V = (\sigma(x)bx^{-1})\sigma(x^{-1}W) = x^{-1}b\sigma(W) \subset x^{-1}W = V.$$

Since $1 \in V = x^{-1}W$, Lemma 3.1 implies that $\sigma(x)bx^{-1}$ is algebraic over (K, σ) and we are done. \square

Proposition 3.3. *If $a, b \in L$ are σ -algebraic over K , then $\sigma(a)$ and ab are σ -algebraic over K too.*

Proof. By Lemma 3.1, there exist finite-dimensional K -vector subspaces V and W of L such that $1 \in V \cap W$, $a\sigma(V) \subset V$ and $a\sigma(W) \subset W$. It is easy to see that the set

$$VW = \{xy \mid x \in V, y \in W\},$$

is a finite-dimensional K -vector subspace of L . Clearly, we have $1 \in VW$. Moreover,

$$(ab)\sigma(VW) = (a\sigma(V))(b\sigma(W)) \subset VW.$$

It follows from Lemma 3.1 that ab is σ -algebraic over K . To prove that $\sigma(a)$ is σ -algebraic over K , let $x_1 = 1, x_2, \dots, x_n$ be a basis for the vector space V . Consider the vector space

$$V' = K\sigma(x_1) + K\sigma(x_2) + \dots + K\sigma(x_n).$$

Clearly, V' is finite-dimensional over K and $1 \in V'$. Since $\sigma(V) \subset V'$, we see that

$$\sigma(a)\sigma(V') = \sigma(aV') = \sigma(K(a\sigma(x_1)) + \dots + K(a\sigma(x_n))) \subset \sigma(V) \subset V'.$$

By Lemma 3.1, a is σ -algebraic over K . \square

Proposition 3.4. *Let $a \in L$ be σ -algebraic over K and V be the K -vector subspace of L generated by $N_i(a)$, where $i \geq 0$. Then any $b \in L$ which is σ -conjugate to a over V is σ -algebraic over K .*

Proof. By Lemma 3.1, V is finite-dimensional over K , and $a\sigma(V) \subset V$. Let $b \in L$ be σ -conjugate to a over V . Then, there exists $x \in V^*$ such that $b = \sigma(x)ax^{-1}$. Consider the vector space

$$x^{-1}V = \{x^{-1}y \mid y \in V\}.$$

Clearly, V is a finite-dimensional vector space over K . Since, $x \in V$, we have $1 \in V$. Moreover, we have

$$b\sigma(x^{-1}V) = \sigma(x)ax^{-1}\sigma(x^{-1}V) = x^{-1}a\sigma(V) \subset x^{-1}V.$$

By Lemma 3.1, b must be σ -algebraic over K . □

The sum of two σ -algebraic elements need not be σ -algebraic, as the following example shows.

Example 3.1. Consider the σ -field $(\mathbb{Q}(x), \sigma)$, where σ is the unique automorphism of $\mathbb{Q}(x)$ determined by $\sigma(x) = x^{-1}$. Considering the σ -field $(\mathbb{Q}(x), \sigma)$ as an σ -extension of the trivial σ -field $(\mathbb{Q}, 1_{\mathbb{Q}})$, we see that x is σ -algebraic over \mathbb{Q} because it is a root of the polynomial $T^2 - 1 \in \mathbb{Q}[T; 1_{\mathbb{Q}}]$. It is easy to show that

$$N_i(x+1) = \frac{(x+1)^i}{x^{\lfloor \frac{i}{2} \rfloor}},$$

for all $i \geq 0$. In particular, the elements $N_i(x+1)$, $i \geq 0$, generate an infinite-dimensional \mathbb{Q} -vector subspace of $\mathbb{Q}(x)$. Using Lemma 3.1, we conclude that, the element $x+1$, which is the sum of the σ -algebraic elements x and 1 , is not σ -algebraic over $(\mathbb{Q}, 1_{\mathbb{Q}})$.

Using Lemma 2.2 and the fact that $K[T; \sigma]$ is a left principal ideal domain, we see that if $a \in L$ is algebraic over (K, σ) , then there exists a unique monic skew polynomial $P_a(T) \in K[T; \sigma]$ such that

$$\{Q(T) \in K[T; \sigma] \mid Q(a) = 0\} = K[T; \sigma]P_a(T).$$

The skew polynomial $P_a(T)$ is called the *minimal (skew) polynomial* of a over (K, σ) . In general, minimal polynomials need not be irreducible. For example, the minimal polynomial of $i \in \mathbb{C}$ in the σ -extension $(\mathbb{C}, -)$ of the σ -field $(\mathbb{R}, 1_{\mathbb{R}})$ is $T^2 - 1$ which is not irreducible in $\mathbb{R}[T; 1_{\mathbb{R}}]$. However, we can show that every element has a σ -conjugate whose minimal polynomial is irreducible. Here are the details: As before, let (L, σ) be an extension σ -field of (K, σ) and $a \in L$ be σ -algebraic over K . Assume that the minimal polynomial $P(T) = P_a(T)$ of a over (K, σ) is reducible. It follows that there exists $Q(T), R(T) \in K[T; \sigma]$ of degree > 0 such that

$$P(T) = Q(T)R(T).$$

Since $R(a) \neq 0$, Lemma 2.2 implies that $Q(a_1) = 0$ where

$$a_1 = \sigma(R(a)) a R(a)^{-1} \in K\langle a \rangle.$$

Here, $K\langle a \rangle$ is the σ -subfield of L which is generated by K and a . In particular, a_1 is σ -conjugate to a over $K\langle a \rangle$, and a_1 is σ -algebraic over K . Let $P_1(T) \in K[T; \sigma]$ be the minimal skew polynomial of a_1 over K . Note that $\deg P_1 < \deg P$ and $Q(T) = Q_1(T)P_1(T)$ for some $Q_1(T) \in K[T; \sigma]$. Continuing this process, we obtain a sequence of elements a_1, a_2, \dots such that $a_{i+1} \in K\langle a_i \rangle$ is σ -conjugate to a_i over $K\langle a_i \rangle$, and a_{i+1} is σ -algebraic over K with minimal skew polynomial $P_{i+1}(T) \in K[T; \sigma]$. Furthermore,

$$\deg P > \deg P_1 > \deg P_2 > \dots$$

Therefore, after finitely many steps, we obtain an element a_i whose minimal skew polynomial is irreducible over (K, σ) . Thus we have proved the following

Proposition 3.5. *Let $a \in (L, \sigma)$ be σ -algebraic over (K, σ) . Then, there exists $b \in K\langle a \rangle$ such that (1) b is σ -conjugate to a over $K\langle a \rangle$, (2) b is σ -algebraic over (K, σ) and the minimal skew polynomial of b over K is irreducible, and (3) $P_a(T) = Q(T)P_b(T)R(T)$ for some $Q(T), R(T) \in K[T; \sigma]$.*

A σ -extension (L, σ) of a σ -field (K, σ) is called σ -algebraic if, as a σ -field, (L, σ) is generated by a family of its elements each of which is σ -algebraic over K . It is called σ -simple if $L = K\langle a \rangle$, where a is σ -algebraic over K . Following [26], we say that two elements $a \in L_1$ and $b \in L_2$ in σ -extensions (L_1, σ) and (L_2, σ) of (K, σ) have the same (K, σ) -type (or simply K -type) if there is an isomorphism $K\langle a \rangle \rightarrow K\langle b \rangle$ of σ -fields over (K, σ) which sends a to b . The proof of the following lemma is left to the reader.

Lemma 3.6. (1) *Let a, b have the same K -type. Then, a is σ -algebraic over K iff b is σ -algebraic over K .*

(2) *If a and b are σ -algebraic of the same K -type, then they have the same minimal polynomial over (K, σ) .*

In contrast to ordinary field extensions, the converse of the second statement in this lemma does not hold in the case of σ -algebraic elements. More precisely, it may happen that two σ -algebraic elements have the same minimal polynomial, but are not of the same type. As an example, the reader can verify that the element i in the σ -extension $(\mathbb{C}, -)$ of the trivial σ -field $(\mathbb{R}, 1_{\mathbb{R}})$, and the element x in the σ -extension $(\mathbb{R}(x), \sigma)$, where $\sigma(x) = 1/x$, of the trivial σ -field $(\mathbb{R}, 1_{\mathbb{R}})$ have the same minimal polynomial $T^2 - 1$ over $(\mathbb{R}, 1_{\mathbb{R}})$. However, i and x are not of the same \mathbb{R} -type.

We conclude this subsection with a general observation. Let (L, σ) be a σ -extension of (K, σ) . For an arbitrary element $a \in L$, there is a unique σ -homomorphism

$$(K\langle x \rangle, \sigma) \rightarrow K[a]_{\sigma},$$

which acts as the identity map on K and sends x to a . Clearly, this map is onto. So, letting J_a be the kernel of this map, we obtain a σ -isomorphism

$$\psi_a : \left(\frac{K\{x\}}{J_a}, \sigma \right) \rightarrow K[a]_\sigma,$$

which through localization gives rise to a σ -isomorphism of σ -fields:

$$\phi_a : \left(\text{Frac}\left(\frac{K\{x\}}{J_a}\right), \sigma \right) \rightarrow K\langle a \rangle.$$

Keeping the above notations, we prove the following:

Proposition 3.7. *Let (L_1, σ_1) and (L_2, σ_2) be σ -extensions of (K, σ) . Then, $a \in L_1$ and $b \in L_2$ are of the same K -type iff $J_a = J_b$.*

Proof. If $J_a = J_b$, then $\phi_b \phi_a^{-1} : K\langle a \rangle \rightarrow K\langle b \rangle$ is a σ -isomorphism which acts as the identity map on K and sends a to b . Therefore, a and b are of the same K -type. Conversely, let a and b have the same K -type. Then, there is a σ -isomorphism $\phi : K\langle a \rangle \rightarrow K\langle b \rangle$ which acts as the identity map on K and sends a to b . Then, the σ -isomorphism

$$\phi_b^{-1} \phi \phi_a : \left(\text{Frac}\left(\frac{K\{x\}}{J_a}\right), \sigma \right) \rightarrow \left(\text{Frac}\left(\frac{K\{x\}}{J_b}\right), \sigma \right),$$

acts as the identity map on K and sends $x + J_a$ to $x + J_b$, from which it follows that $J_a = J_b$. \square

3.2 Minimal polynomials

In this part, we turn to the question of deciding when a given skew polynomial can be a minimal polynomial. In what follows, let (K, σ) be a σ -field, and

$$P(T) = T^{n+1} - c_n T^n - \cdots - c_1 T - c_0 \in K[T; \sigma],$$

be a (monic) polynomial of degree $n+1 > 0$. Although, most of the results of this part hold if $P(T) \notin T K[T; \sigma]$, we assume, for simplicity, that $P(0) \neq 0$, that is, $c_0 \neq 0$ (see Remark 3.1).

Consider the ring $K[y_0, \dots, y_{n-1}]$ of polynomials over K in n (commuting) indeterminates, and let V_n be the set of all $Y \in K[y_0, \dots, y_{n-1}]$ of the form

$$Y = d_0 + d_1 y_0 + d_2 y_0 y_1 + \cdots + d_{n-1} y_0 y_1 \cdots y_{n-2} + d_n y_0 y_1 \cdots y_{n-1},$$

where $d_0, \dots, d_n \in K$. Clearly, V_n is an $(n+1)$ -dimensional K -vector subspace of $K[y_0, \dots, y_{n-1}]$. First, we define a map $L : V_n \rightarrow V_n$ as follows: For

$$Y = d_0 + d_1 y_0 + \cdots + d_n y_0 y_1 \cdots y_{n-1} \in V_n,$$

we set

$$L(Y) = \sigma(d_n)c_0 + (\sigma(d_0) + \sigma(d_n)c_1)y_0 + \cdots + (\sigma(d_{n-1}) + \sigma(d_n)c_n)y_0y_1 \cdots y_{n-1}.$$

Next, we define a ring homomorphism $\phi : K[y_0, \dots, y_{n-1}] \rightarrow K(y_0, \dots, y_{n-1})$ as follows: By the universal property of $K[y_0, \dots, y_{n-1}]$, there exists a unique ring homomorphism

$$\phi : K[y_0, \dots, y_{n-1}] \rightarrow K(y_0, \dots, y_{n-1}),$$

such that $\phi(a) = \sigma(a)$, for all $a \in K$, and

$$\phi(y_i) = \begin{cases} y_{i+1} & i = 0, 1, \dots, n-2, \\ \frac{L(y_0y_1 \cdots y_{n-1})}{y_0y_1 \cdots y_{n-1}} & i = n-1. \end{cases}$$

The following lemma gives some properties of the maps L and ϕ .

- Lemma 3.8.** (1) For $Y \in V_n$, $L(Y) = 0$ iff $Y = 0$.
(2) The ring homomorphism ϕ is injective.
(3) For all $Y \in V_n$ and $i > 0$, $L^i(Y) = y_0\phi(y_0) \cdots \phi^{i-1}(y_0)\phi^i(Y)$.

Proof. (1) Clearly, $L(0) = 0$. Assume that $L(Y) = 0$ for some nonzero

$$Y = d_0 + d_1y_0 + \cdots + d_ny_0y_1 \cdots y_{n-1} \in V_n.$$

If $d_n = 0$, then $L(Y) = 0$ implies that

$$\sigma(d_0) = \sigma(d_1) = \cdots = \sigma(d_{n-1}) = 0 \implies Y = 0.$$

If $d_n \neq 0$, then $\sigma(d_n)c_0 = 0$ implies that $c_0 = 0$, a contradiction. This completes the proof of (1).

(2) Suppose, on the contrary, that ϕ is not injective. Then, there exist ordinary polynomials

$$f_0(y_0, \dots, y_{n-2}), \dots, f_m(y_0, \dots, y_{n-2}) \in K[y_0, \dots, y_{n-2}],$$

such that $f_0f_m \neq 0$ and $\phi(\sum_{i=0}^m f_i(y_0, \dots, y_{n-2})y_{n-1}^i) = 0$. Computing the left-hand side using the definition of ϕ , we obtain

$$\sum_{i=0}^m g_i(y_1, \dots, y_{n-1}) \left(\frac{L(y_0y_1 \cdots y_{n-1})}{y_0y_1 \cdots y_{n-1}} \right)^i = 0, \quad (3.1)$$

where $g_0, \dots, g_m \in K[y_0, \dots, y_{n-2}]$, with $g_0g_m \neq 0$. Writing

$$\begin{aligned} \frac{L(y_0y_1 \cdots y_{n-1})}{y_0y_1 \cdots y_{n-1}} &= \frac{c_0 + c_1y_0 + \cdots + c_{n-1}y_0y_1 \cdots y_{n-2} + c_ny_0y_1 \cdots y_{n-1}}{y_0y_1 \cdots y_{n-1}} \\ &= c_0y_0^{-1}y_1^{-1} \cdots y_{n-1}^{-1} + c_1y_1^{-1} \cdots y_{n-1}^{-1} + \cdots + c_{n-1}y_{n-1}^{-1} + c_n \end{aligned}$$

and expanding the left-hand side of Equation 3.1 as a polynomial in y_0^{-1} , we see that the highest coefficient of y_0^{-1} in the left-hand side of the equation is equal to

$$(c_0 y_1^{-1} \cdots y_{n-1}^{-1})^m g_m(y_1, \dots, y_{n-1}) \neq 0,$$

a contradiction. This completes the proof of (2).

(3) We first note that the homomorphism ϕ can be extended to an endomorphism $\phi : K(y_0, \dots, y_{n-1}) \rightarrow K(y_0, \dots, y_{n-1})$, by Part (2). In particular, the formula given in (3) is well-defined. Given

$$Y = d_0 + d_1 y_0 + d_2 y_0 y_1 + \cdots + d_n y_0 y_1 \cdots y_{n-1} \in V_n,$$

we write

$$\begin{aligned} \phi(Y) &= \sigma(d_0) + \sigma(d_1) y_1 + \sigma(d_2) y_1 y_2 + \cdots + \sigma(d_n) y_1 y_2 \cdots y_{n-1} y_n \\ &= \sigma(d_0) + \sigma(d_1) y_1 + \sigma(d_2) y_1 y_2 + \cdots + \sigma(d_{n-1}) y_1 y_2 \cdots y_{n-1} \\ &\quad + \sigma(d_n) \frac{c_0 + c_1 y_0 + \cdots + c_n y_0 y_1 \cdots y_{n-1}}{y_0} \end{aligned}$$

This gives

$$\begin{aligned} y_0 \phi(Y) &= \sigma(d_0) y_0 + \sigma(d_1) y_0 y_1 + \cdots + \sigma(d_{n-1}) y_0 y_1 y_2 \cdots y_{n-1} \\ &\quad + \sigma(d_n) (c_0 + c_1 y_0 + c_2 y_1 y_2 + \cdots + c_n y_0 y_1 \cdots y_{n-1}) \\ &= \sigma(d_n) c_0 + (\sigma(d_0) + \sigma(d_n) c_1) y_0 + \cdots + (\sigma(d_{n-1}) + \sigma(d_n) c_n) y_0 y_1 \cdots y_{n-1} \\ &= L(Y) \end{aligned}$$

proving (3) for $i = 1$. Now, a simple induction yields the desired formula. \square

Let S be the multiplicative subset of $K[y_0, \dots, y_{n-1}]$ which is generated by all the nonzero elements of V_n . It follows from Lemma 3.8 that ϕ , through localization, gives rise to a ring homomorphism

$$\sigma_P : S^{-1}K[y_0, \dots, y_{n-1}] \rightarrow S^{-1}K[y_0, \dots, y_{n-1}], \quad \sigma_P(rs^{-1}) = \phi(r)\phi(s)^{-1},$$

where $r \in K[y_0, \dots, y_{n-1}]$ and $s \in S$. Using Part (2) of Lemma 3.8, we see that σ_P is injective. Therefore, the pair

$$(K[P], \sigma_P) := (S^{-1}K[y_0, \dots, y_{n-1}], \sigma_P),$$

is a σ -domain. The σ -field of fractions of $(K[P], \sigma_P)$ is denoted by $(K(P), \sigma_P)$. Note that $(K(P), \sigma_P)$ is a σ -extension of (K, σ) . Now, we show that $P(T)$ is the minimal polynomial of an element in $(K(P), \sigma_P)$.

Proposition 3.9. *The element $y_0 \in K(P)$ is a nonzero root of $P(T)$ in the σ -extension $(K(P), \sigma)$ of (K, σ) . Moreover, the minimal skew polynomial of $y_0 \in K(P)$ over (K, σ) is $P(T)$.*

Proof. The fact that $y_0 \in K(P)$ is a root of $P(T)$ follows from the fact that

$$P(y_0) = y_0 y_1 \cdots y_{n-1} \frac{L(y_0 y_1 \cdots y_{n-1})}{y_0 y_1 \cdots y_{n-1}} - c_n y_0 y_1 \cdots y_{n-1} - \cdots - c_1 y_0 - c_0 = 0.$$

If y_0 is a root of a skew polynomial

$$d_0 + d_1 T + \cdots + d_m T^m \in K[T; \sigma],$$

where $m < \deg P$, then

$$0 = d_0 + d_1 y_0 + d_2 y_0 y_1 + \cdots + d_m y_0 y_1 \cdots y_{m-1},$$

which implies $d_0 = \dots = d_{m-1} = 0$, since y_0, \dots, y_{n-1} are algebraically independent over K . This completes the proof. \square

Next, we show that the σ -domain $(K[P], \sigma_P)$ has the following universal property:

Theorem 3.10. *An element $0 \neq a$ in a σ -extension (L, σ) of (K, σ) has $P(T)$ as its minimal polynomial over (K, σ) iff there exists a σ -homomorphism*

$$\psi_a : (K[P], \sigma_P) \rightarrow (L, \sigma),$$

over (K, σ) , such that $\psi_a(y_0) = a$. Moreover, such a σ -homomorphism is unique.

Proof. First, suppose that $P(T)$ is the minimal polynomial of $a \in L$. By the universal property of $K[y_0, \dots, y_{n-1}]$, there exists a unique K -algebra homomorphism

$$\psi : K[y_0, \dots, y_{n-1}] \rightarrow L$$

such that $\psi(y_i) = \sigma^i(a)$ for $i = 0, 1, \dots, n-1$. The condition $P(a) = 0$ gives

$$\sigma^n(a) \cdots \sigma(a) a = c_0 + c_1 a + c_2 a \sigma(a) + \cdots + c_n a \sigma(a) \sigma^2(a) \cdots \sigma^{n-1}(a),$$

from which it follows that

$$\sigma^n(a) \psi(y_{n-1}) \cdots \psi(y_0) = \psi(L(y_0, \dots, y_{n-1})).$$

Localizing ψ at $L(y_0, \dots, y_{n-1})$, we obtain a K -algebra homomorphism

$$\psi_1 : K[y_0, \dots, y_{n-1}][L(y_0, \dots, y_{n-1})^{-1}] \rightarrow L,$$

such that $\psi_1\left(\frac{L(y_0 y_1 \cdots y_{n-1})}{y_0 y_1 \cdots y_{n-1}}\right) = \sigma^n(a)$. For any

$$0 \neq r = d_0 + d_1 y_0 + d_2 y_0 y_1 + \cdots + d_n y_0 y_1 \cdots y_{n-1} \in S,$$

we have

$$\psi_1(r) = d_0 + d_1 a + d_2 a \sigma(a) + \cdots + d_n a \sigma(a) \cdots \sigma^{n-1}(a) \neq 0,$$

since the minimal polynomial of a has degree $n + 1$. Therefore, we can localize ψ_1 at S to obtain a K -algebra homomorphism

$$\psi_a : S^{-1}K[y_0, \dots, y_{n-1}] \rightarrow L.$$

Since $\psi_a(\sigma_P(y_i)) = \sigma(\psi_a(y_i))$ for $i = 0, 1, \dots, n-1$, one can easily verify that ψ_a is a σ -homomorphism satisfying $\psi_a(y_0) = a$. Conversely, suppose that

$$\psi_0 : (K[P], \sigma_P) \rightarrow (L, \sigma),$$

is a σ -homomorphism over (K, σ) which satisfies $\psi_0(y_0) = a$. We have $P(a) = 0$ since ψ_0 is a σ -homomorphism, and $P(y_0) = 0$ by Proposition 3.9. The fact that the image of any element in S is invertible in L implies that $P(T)$ is the minimal polynomial of a . The uniqueness part follows from the fact that any σ -homomorphism on $(K[P], \sigma_P)$ is completely determined by its value at y_0 and its restriction to K . \square

Now, we give some applications of this theorem and Proposition 3.9. The following corollary is an immediate consequence of Proposition 3.9.

Corollary 3.11. *Any nonconstant monic polynomial $Q(T) \in K[T; \sigma]$ satisfying $Q(0) \neq 0$ is the minimal polynomial of a (nonzero) element in some σ -extension of (K, σ) .*

The following proposition, the proof of which is left to the reader, explains the connection between K -types and prime σ -ideals.

Proposition 3.12. *Let $Q(T) \in K[T; \sigma]$ be a nonconstant monic polynomial $Q(T) \in K[T; \sigma]$ satisfying $Q(0) \neq 0$. For every prime σ -ideal I of $(K[Q], \sigma_Q)$, the polynomial $Q(T)$ is the minimal polynomial of the element $y_0 + I$ in the σ -field $(\text{Frac}(K[Q]/I), \sigma)$ of fractions of $(K[Q]/I, \sigma)$. Moreover, the following holds:*

- (1) $(\text{Frac}(K[Q]/I), \sigma) = K\langle y_0 + I \rangle$,
- (2) For any element $0 \neq a \in L$ in a σ -extension of (K, σ) whose minimal polynomial over K is $Q(T)$, there is a unique prime σ -ideal I_a of $(K[Q], \sigma_Q)$ such that $y_0 + I_a$ has the same K -type as a .

The assignment $a \mapsto I_a$, given in this proposition, establishes a 1-1 correspondence between distinct K -types with a fixed minimal polynomial $Q(T) \in K[T; \sigma]$, and the set of prime σ -ideals of $(K[Q], \sigma_Q)$. We study this correspondence more closely. Let $Q(T) \in K[T; \sigma]$ be a nonconstant monic polynomial $Q(T) \in K[T; \sigma]$ satisfying $Q(0) \neq 0$. Let $\mathcal{P}(Q)$ be the set of all prime σ -ideals of $(K[Q], \sigma_Q)$. Let $\mathcal{L}(Q)$ be the class of all elements in σ -extensions of (K, σ) whose minimal polynomial is $Q(T)$. Consider the equivalence relation \sim on $\mathcal{L}(Q)$ for which $a \sim b$ iff a and b have the same K -type. Let $\mathcal{T}(Q)$ be the class of equivalence classes of \sim in $\mathcal{L}(Q)$. Then, the assignment $I \mapsto [y_0 + I]$, where $[a]$ denotes the equivalence class of $a \in \mathcal{L}(Q)$,

establishes a 1-1 correspondence between $\mathcal{P}(Q)$ and $\mathcal{T}(Q)$. In particular, $\mathcal{T}(Q)$ is a set, called the set of K -types of $Q(T)$. Using this correspondence, we can define a relation \leq on $\mathcal{T}(Q)$: Given $[a], [b] \in \mathcal{L}(Q)$, we write $[a] \leq [b]$ if $I_a \subset I_b$. There is a unique minimal element in $\mathcal{T}(Q)$ which corresponds to the prime σ -ideal (0) of $(K[Q], s_Q)$. Also, using Zorn's lemma, one can show that for any $[a] \in \mathcal{T}(Q)$, there exists a maximal element $[b] \in \mathcal{T}(Q)$ such that $[a] \leq [b]$. We note that the set $\mathcal{P}(Q)$ is a subset of the affine scheme $\text{Spec}(K[y_0, \dots, y_{n-1}])$, where $n = \deg Q - 1$. We conclude this part with an example and a remark:

Example 3.2. *Let K be an arbitrary (commutative) field. Consider the trivial σ -field $(K, 1_K)$. Let $P(T) \in K[T; 1_K]$ be the polynomial*

$$P(T) = T^2 - 1.$$

One can easily check that $K[P] = S^{-1}K[y_0]$, where S is the multiplicative subset of $K[y_0]$ which is generated by $y_0 - a$, where $a \in K$. Moreover, we have $\sigma_P(y_0) = y_0^{-1}$. Since every prime ideal of $K[P]$, apart from the zero ideal, is of the form $K[P]f(y_0)$ where $f(y_0) \in K[y_0]$ is an irreducible polynomial of degree > 1 , it is easy to see that a prime ideal $K[P]f(y_0)$ of $K[P]$ is a σ -ideal iff there exists an element $a \in K$ such that

$$y_0^n f\left(\frac{1}{y_0}\right) = af(y_0),$$

where n is the degree of f . In other words, $K[P]f(y_0)$ is a prime σ -ideal iff $f_0(y)$ is irreducible and palindromic of degree > 1 . Therefore, there is a 1-1 correspondence between the set of monic palindromic irreducible polynomials $f(y_0) \in K[y_0]$ of degree 1 and the set $\mathcal{P}(T^2 - 1) \setminus \{(0)\}$. In particular, if K is an algebraically closed field, we have $\mathcal{P}(T^2 - 1) = \{(0)\}$. We note that the σ -extension corresponding to the zero ideal (0) is $(K(y_0), \sigma)$ with $\sigma(y_0) = y_0^{-1}$.

Remark 3.1. *Results similar to the ones presented in this part hold for any monic polynomial not contained in $TK[T; \sigma]$. In fact, one can show that a monic polynomial $P(T) \in K[T; \sigma]$ is the minimal polynomial of a (nonzero) element in some σ -extension of (K, σ) iff $P(T) \notin TK[T; \sigma]$. Also, a skew polynomial $P(T) \in K[T; \sigma]$ has a nonzero root in some σ -extension of (K, σ) iff $P(T)$ is not of the form aT^m where $a \in K^*$ and $m \geq 0$. Note that the condition $P(T) \notin TK[T; \sigma]$ is equivalent to the condition $P(0) \neq 0$ provided that (K, σ) is inversive.*

4 Algebraically closed σ -fields

As is well-known, every (commutative) field has an algebraic closure which is unique up to isomorphism. This property of fields is based on the following two facts:

F1 Any irreducible polynomial over a field has a root in some extension of the field.

F2 Any polynomial over a field has a unique root up to isomorphism.

Thanks to F1, one can construct an algebraic closure of a given field by successively "adjoining" roots of irreducible polynomials. The uniqueness of algebraic closure for fields is a consequence of F2.

More generally, the notion of root can be defined for other algebraic objects, using which one can define the corresponding concept of algebraic closedness. Generally speaking, if F1 holds for a particular algebraic object, then one can construct algebraic closures for the algebraic object under study by adjoining roots "one after another". However, if F2 does not hold, there may exist several notions of algebraic closedness. We refer the reader to [9, 22] for the case of skew fields, and [21, 25] for the case of groups, (see also [24] for a model-theoretic approach to this problem). In this section, we investigate the question of algebraic closedness for σ -fields. Note that F1 holds for skew roots, see Proposition 3.9. However, F2 does not hold for skew roots, see Proposition 3.12. The lack of F2 for σ -fields gives rise to different notions of algebraic closedness for σ -fields.

4.1 Some notions of algebraically closed σ -fields

The simplest notion of algebraic closedness for σ -field is the following: A σ -field (L, σ) is called *1-algebraically closed* if every nonconstant polynomial in $L[T; \sigma]$ has a root in L . In this part, we show that every σ -field has a σ -extension which is 1-algebraically closed. To prove this result, we implement a classical method used to prove the existence of algebraic closures for ordinary fields, see [19, Theorem 2.5].

Theorem 4.1. *Every σ -field can be embedded in a 1-algebraically closed σ -field.*

Proof. Let (K, σ) be an arbitrary σ -field. The idea is to adjoin skew roots of irreducible skew polynomials until there are no irreducible polynomials of degree > 1 left to deal with. This will be done in a series of steps as described below:

Selecting and ordering polynomials (Step 1): Let D be the set of all monic irreducible polynomials in $K[T; \sigma]$ of degree > 1 which do not have a root in K . Consider a total order on $D_1 = \{1\} \cup D$ which turns D_1 into a well-ordered set with 1 being its least element. Let α_0 be the ordinal of D_1 . Given an ordinal $\alpha \leq \alpha_0$, we denote the α -th element of D_1 by P_α . Note that any polynomial in D has a nonzero constant term since otherwise it would not be irreducible.

Adjoining roots (Step 2): Using transfinite induction, we define a sequence (K_α, σ) , where $\alpha \leq \alpha_0$ is an ordinal, of σ -fields as follows: we set

$(K_0, \sigma) = (K, \sigma)$ and $K_\alpha = (\cup_{\beta < \alpha} K_\beta)(P_\alpha)$, see Proposition 3.9. We note that (K_α, σ) is a σ -extension of (K_β, σ) if $\beta \leq \alpha \leq \alpha_0$. Now, we consider the σ -field $(K^{(1)}, \sigma) = (K_{\alpha_0}, \sigma)$. Clearly, $(K^{(1)}, \sigma)$ is a σ -extension of (K, σ) . By construction, every $P(T) \in D$ has a root in $K^{(1)}$. Therefore, any irreducible nonconstant polynomial in $K[T; \sigma]$ has a root in $K^{(1)}$.

Induction (Step 3): Applying Steps 1 and 2 to $(K^{(1)}, \sigma)$ gives rise to a σ -extension $(K^{(2)}, \sigma)$ of $(K^{(1)}, \sigma)$ in which any nonconstant polynomial belonging to $K^{(1)}[T; \sigma]$ has a root. Inductively, we obtain a sequence of σ -fields

$$(K^{(1)}, \sigma) \subset (K^{(2)}, \sigma) \subset (K^{(3)}, \sigma) \subset \dots,$$

such that any nonconstant polynomial belonging to $K^{(n)}[T; \sigma]$ has a root in $(K^{(n+1)}, \sigma)$. Finally, we consider the σ -field

$$(K^\infty, \sigma) = \cup_{n=1}^\infty (K^{(n)}, \sigma).$$

Any skew polynomial $P(T) \in K^\infty[T; \sigma]$ belongs to some $K^{(n)}[T; \sigma]$. It follows that if $P(T)$ has a root in $(K^{(n+1)}, \sigma)$, hence in (K^∞, σ) . Therefore, (K^∞, σ) is 1-algebraically closed and contains (K, σ) as a σ -subfield. This completes the proof of the theorem. \square

A 1-algebraically closed σ -field is only required to contain at least one root of every nonconstant skew polynomial. The requirement that any skew polynomial have more than one root, gives rise to the following notion of algebraic closedness: Let c be a cardinal number. A σ -field (L, σ) is called *c-algebraically closed* if any skew polynomial over (L, σ) , of degree > 1 and with a nonzero constant term, has at least c distinct roots in L . Clearly, if $c \leq c'$ are two cardinals, then any c' -algebraically closed σ -field is c -algebraically closed.

Theorem 4.2. *Let c be a cardinal number. Any σ -field can be embedded in a c -algebraically closed σ -field.*

Proof. The proof is similar to that of Theorem 4.1. Here is an outline of the proof. The details are left to the reader to verify.

Selecting and ordering polynomials (Step 1): Let D be the set of all monic polynomials in $K[T; \sigma]$ which are of degree > 1 and have a nonzero constant term. Consider a total order on $D_1 = \{1\} \cup D$ which turns D_1 into a well-ordered set with 1 being its least element. Let α_0 be the ordinal of D_1 . Given an ordinal $\alpha \leq \alpha_0$, we denote the α -th element of D_1 by P_α .

Adjoining roots (Step 2): Using transfinite induction, we define a sequence (K_α, σ) , where $\alpha \leq \alpha_0$ is an ordinal, of σ -fields as follows: We set $(K_0, \sigma) = (K, \sigma)$; K_α is obtained from $\cup_{\beta < \alpha} K_\beta$ by adjoining c copies of roots of P_α using Proposition 3.9. In particular, $(K^{(1)}, \sigma) = (K_{\alpha_0}, \sigma)$ is a σ -extension of (K, σ) , and, by construction, every $P(T) \in D$ has at least c distinct roots in $K^{(1)}$.

Induction (Step 3): Using Steps 1, 2 and induction, we can construct a sequence of σ -fields

$$(K^{(1)}, \sigma) \subset (K^{(2)}, \sigma) \subset (K^{(3)}, \sigma) \subset \dots,$$

such that any polynomial belonging to $K^{(n)}[T; \sigma]$, of degree > 1 and with a nonzero constant term, has a root in $(K^{(n+1)}, \sigma)$. One can see that the σ -field

$$(K^\infty, \sigma) = \cup_{n=1}^\infty (K^{(n)}, \sigma).$$

is c -algebraically closed and contains (K, σ) as a σ -subfield. \square

σ -field is c -algebraically closed, it is only enough to check the condition for polynomials of degree 2.

In the proof of Theorem 4.2, we have used Proposition 3.9 as a consequence of which most roots would be of the same type. Taking into account that skew polynomials can have roots of different types, we introduce the following definition: A σ -field (L, σ) is called *type-algebraically closed* if for any element a in some σ -extension of (L, σ) with minimal polynomial $P(T) \in L[T; \sigma]$ over (L, σ) , there is $b \in L$ whose minimal polynomial over the σ -subfield K of (L, σ) generated by the coefficients of $P(T)$ is the polynomial $P(T)$, and a and b have the same K -type. Using Proposition 3.12 and modifying the steps in the proof Theorem 4.1, one can prove the existence of type-algebraically closed σ -fields. More precisely, we have the following result the proof of which is left to the reader.

Theorem 4.3. *Any σ -field can be embedded in a type-algebraically closed σ -field.*

One can define a "hybrid version" of algebraic closedness in which every polynomial has at least a prescribed number of distinct roots of every possible type. We leave it to the reader to formulate and prove the existence of such algebraically closed σ -fields. We conclude this part with an observation regarding polynomial factorization in algebraically closed σ -fields.

Proposition 4.4. *(a) Let (K, σ) be a 1-algebraically closed σ -field. Then, any nonzero skew polynomial $P(T) \in K[T; \sigma]$ splits completely, that is, there exist $a, a_1, \dots, a_n \in K^*$ and $m \geq 0$ such that*

$$P(T) = a(T - a_n) \cdots (T - a_2)(T - a_1)T^m.$$

(b) Let (K, σ) be \aleph_0 -algebraically closed. Then, any $P(T) \in K[T; \sigma]$ can be written as

$$P(T) = a(T - a_n) \cdots (T - a_2)(T - a_1)T^m,$$

where $a \in K^$, $a_1, \dots, a_n \in K^*$ are distinct, and $m \geq 0$.*

Proof. Let $P(T) \in K[T; \sigma]$. There exists $m \geq 0$ and $Q(T) \in K[T; \sigma]$ such that $Q(T)$ has a nonzero constant term and $P(T) = Q(T)T^m$. If $Q(T)$ is a constant, we are done. Otherwise, there exists $a_1 \in K$ such that $Q(a_1) = 0$ because (K, σ) is 1-algebraically closed. By Lemma 2.1, there exists $Q_1(T) \in K[T; \sigma]$ such that

$$Q(a) = Q_1(a)(T - a_1).$$

Clearly, $Q_1(T)$ has a nonzero constant term. Now, we prove (a) and (b) as follows:

(a) Using induction on the degree of P , one can obtain the desired factorization.

(b) If $Q_1(T)$ is a constant, we are done. Otherwise, we can find a root $a_1 \neq a_2 \in K$ since K is \aleph_0 -algebraically closed. A simple induction on the degree of $Q(T)$ completes the proof. \square

4.2 Conjugacy classes in algebraically closed σ -fields

In order to determine whether a σ -field is c -algebraically closed for $c > 1$, it is enough to study polynomials of degree two. More precisely, we have the following:

Proposition 4.5. *Let $c > 1$ be a cardinal number. A σ -field (L, σ) is c -algebraically closed iff it is 1-algebraically closed and every skew polynomial $P(T) \in L[T; \sigma]$ of degree 2 whose constant term is nonzero has at least c distinct roots in L .*

Proof. To prove the "if" direction, let $P(T) \in L[T; \sigma]$ be an arbitrary polynomial satisfying $\deg P > 1$ and $P(0) \neq 0$. Since (L, σ) is 1-algebraically closed, it follows from Proposition 4.4, that there are skew polynomials $P_1(T), P_2(T) \in L[T; \sigma]$ such that $\deg P_2 = 2$ and $P(T) = P_1(T)P_2(T)$. Since any root of $P_2(T)$ is a root of $P_1(T)$, we see that $P(T)$ has at least c distinct roots in L . The other direction is trivial. \square

To present the next result, we need to review some facts regarding roots of skew polynomials in a fixed conjugacy class. For more details see [14, 18]. Let (K, σ) be a σ -field with fixed field F . Let $P(T) = \sum_{i=0}^n a_i T^i \in K[T; \sigma]$. An element $\sigma(x)x^{-1}$, where $x \in K^*$, is a root of $P(T)$ iff

$$P(\sigma(x)x^{-1}) = 0 \iff \sum_{i=0}^n a_i \sigma^i(x)x^{-1} = 0 \iff \sum_{i=0}^n a_i \sigma^i(x) = 0.$$

The set $E(P, 1)$ of all $x \in K$ for which $\sum_{i=0}^n a_i \sigma^i(x) = 0$ is a vector space over F . One can see that the assignment $[x] \mapsto \sigma(x)x^{-1}$ establishes a bijection between the projective space $\mathbb{P}(E(P, 1))$ over F and the set of roots

of $P(T)$ in the conjugacy class $C(1)$ containing 1. In particular, if F and $\dim_F E(P, 1)$ are finite, the number of roots of $P(T)$ in $C(1)$ is equal to

$$\frac{|F|^{\dim_F E(P,1)} - 1}{|F| - 1}.$$

Theorem 4.6. *Let (K, σ) be a 2-algebraically closed σ -field with fixed field F . Then, K is 3-algebraically closed iff K has exactly two distinct σ -conjugacy classes. Moreover, K is 3-algebraically closed, then (K, σ) is $(1 + |F|)$ -algebraically closed. In particular, if K is 3-algebraically closed and F is infinite, (K, σ) is \aleph_0 -algebraically closed.*

Proof. Generally speaking, any σ -field has at least two distinct σ -conjugacy classes: the trivial conjugacy class $\{0\}$ and the conjugacy class containing 1. If (K, σ) has a nonzero element a which is not conjugate to 1, then Proposition 2.10 implies that there is a skew polynomial of degree two whose roots in K are exactly 1 and a . Therefore, if K has at least three distinct conjugacy classes, then it cannot be 3-algebraically closed. To prove the second statement, we assume that K has exactly two distinct σ -conjugacy classes. Let $P(T) \in K[T; \sigma]$ be a polynomial of degree > 1 such that $P(0) \neq 0$. Since (K, σ) has only one nontrivial conjugacy classes, we see that the cardinality of the set of roots of $P(T)$ is equal to the cardinality of the projective space $\mathbb{P}(E(P, 1))$. Since (K, σ) is 2-algebraically closed, we must have $\dim_F E(P, 1) > 1$ from which it follows that the cardinality of $\mathbb{P}(E(P, 1))$ is at least $\geq 1 + |F|$, and we are done. \square

We remark that a σ -field has exactly two distinct σ -conjugacy classes iff the map $x \mapsto \sigma(x)x^{-1}$ is a bijection of K^* . As an application, we have the following:

Corollary 4.7. *Let $n > 2$ be a natural number. If (K, σ) is an n -algebraically closed σ -field which is not $(n+1)$ -algebraically closed, then the characteristic p of K is nonzero, and moreover, n is of the form*

$$\frac{p^{mk} - 1}{p^k - 1},$$

for some natural numbers m, k with $m > 1$.

Proof. Since (K, σ) is 3-algebraically closed, it follows from Theorem 4.6 that the fixed field F of K is finite. In particular, the characteristic p of K is nonzero, and $|F| = p^k$ for some natural number k . Since (K, σ) is not $(n + 1)$ -algebraically closed, there exists a nonconstant polynomial $P(T) \in K[T; \sigma^n]$ of degree > 1 such that $P(0) \neq 0$ and $P(T)$ has exactly n distinct roots. Since the number of roots of $P(T)$ is equal to

$$\frac{|F|^{\dim_F E(P,1)} - 1}{|F| - 1} = \frac{p^{k \dim_F E(P,1)} - 1}{p^k - 1},$$

the result follows. \square

Another fact regarding algebraically closed σ -fields is given in the following proposition.

Proposition 4.8. *Let (K, σ) be a nontrivial σ -field, that is, $\sigma \neq 1_K$. If (K, σ) is 1-algebraically closed, then σ is of infinite order.*

Proof. Assume on the contrary that σ is of finite order, say order n . Consider the skew polynomial $P_a(T) = T^n - a \in K[T; \sigma]$ where $a \in K$. An element $x \in K$ is a root of $P_a(T)$ iff

$$N_n(x) = \sigma^{n-1}(x) \cdots \sigma(x)x = a.$$

Since $\sigma^n = 1_K$, we see that $N_n(x) = N_{K/F}(x)$ is the norm of x over the fixed field F of K . In particular, we must have $a \in F$ if $P_a(T)$ has a root in K . Since (K, σ) is 1-algebraically closed, we conclude that $K = F$, contradicting the assumption that $\sigma \neq 1_K$. \square

In the next section, we give examples of n -algebraically closed σ -fields.

4.3 Examples of algebraically closed σ -fields

We begin with a trivial example. Let K be an (ordinary) algebraically closed field. It is easy to see that the trivial σ -field $(K, 1_K)$ is a 1-algebraically closed σ -field. However, it is not 2-algebraically closed because the skew polynomial $T^2 - 2T + 1 \in K[T; 1_K]$ has exactly one root in K .

To give a nontrivial example, we let K be an (ordinary) algebraically closed field of characteristic $p > 0$. Let $\sigma : K \rightarrow K$ be the Frobenius homomorphism, that is, $\sigma(a) = a^p$ for all $a \in K$. Given a natural number n , we obtain the σ -field (K, σ^n) . It is known that the fixed field of (K, σ^n) is the finite field \mathbb{F}_{p^n} with p^n elements. Working over (K, σ^n) , one can easily check that

$$N_i(a) = a^{\frac{p^{in}-1}{p^n-1}},$$

for all $a \in K$ and natural numbers $i \geq 1$. Therefore, given a skew polynomial $P(T) = \sum_{i=0}^m a_i T^i \in K[T; \sigma^n]$ and $a \in K$, we obtain

$$P(a) = \sum_{i=0}^m a_i N_i(a) = \sum_{i=0}^m a_i a^{\frac{p^{in}-1}{p^n-1}}.$$

Setting

$$f(x) = \sum_{i=0}^m a_i x^{\frac{p^{in}-1}{p^n-1}} \in K[x],$$

we see that $P(a) = f(a)$ for all $a \in K$. Since K is algebraically closed, we conclude that (K, σ^n) is 1-algebraically closed. In fact, more can be said. Assume that $P(0) \neq 0$. Then, $f(x)$ is separable. Since the degree of f is

$$\frac{p^{nm} - 1}{p^n - 1}$$

where m is the degree of $P(T)$, we see that the number of distinct roots of the skew polynomial $P(T)$ is

$$\frac{p^{nm} - 1}{p^n - 1}.$$

In particular, we see that (K, σ^n) is $(p^n + 1)$ -algebraically closed, but not $(p^n + 2)$ -algebraically closed, since if $\deg P = 2$, then $P(T)$ has exactly $p^n + 1$ distinct roots.

4.4 c -Algebraically closed inversive σ -fields

In the preceding part, we proved the existence of various types of algebraically closed σ -field. Here, we prove the existence of c -algebraically closed inversive σ -fields. Recall that a σ -field (K, σ) is called inversive if σ is an automorphism.

Proposition 4.9. *Let c be a cardinal number and (K, σ) be c -algebraically closed σ -field. Then, the inversive closure of (K, σ) is c -algebraically closed.*

Proof. Let (L, σ) be the inversive closure of (K, σ) . Assume that a polynomial

$$P(T) = \sum_{i=0}^n a_i T^i \in L[T; \sigma],$$

satisfies $n > 1$ and $a_0 \neq 0$. Since (L, σ) is the inversive closure of (K, σ) , there exists $N \in \mathbb{N}$ such that $\sigma^N(a_i) \in K$ for all $i = 0, 1, \dots, n$. Then, the skew polynomial

$$Q(T) = \sum_{i=0}^n \sigma^N(a_i) T^i,$$

belongs to $K[T; \sigma]$. It is easy to see that if $a \in L$ is a root of $Q(T)$, then $\sigma^{-N}(a)$ is a root of $P(T)$. Since $Q(T)$ has at least c distinct roots in K , we conclude that $P(T)$ has at least c distinct roots in L . This completes the proof of the proposition. \square

As a consequence of this proposition, we have the following result:

Corollary 4.10. *Let c be a cardinal number. Any σ -field (K, σ) can be embedded in a c -algebraically closed inversive σ -field.*

Proof. By Theorem 4.2, (K, σ) can be embedded in a c -algebraically closed σ -field (L, σ) . Then, by Proposition 4.9, the inversive closure of (L, σ) is c -algebraically closed. Since the inversive closure of (L, σ) contains (K, σ) as a σ -subfield, the result follows. \square

5 Partial fraction decomposition

In this section, we present a version of a partial fraction decomposition for the field of fractions of the ring of skew polynomials. We begin with some standard definitions and facts regarding factorization in (not necessarily commutative) principal ideal domains, see [6, 8] for more details. Let R be a (not necessarily commutative) principal ideal domain (or PID for short). A nonunit $a \in R$ is called irreducible if $a = bc$, where $b, c \in R$, implies that b or c is a unit in R . Elements $a, b \in R$ are called similar if the left modules R/Ra and R/Rb are isomorphic, or equivalently, the right modules R/aR and R/bR are isomorphic. Elements $a, b \in R$ are called left (resp., right) co-prime if $aR + bR = R$ (reps., $Ra + Rb = R$). They are called co-prime if they are both left and right co-prime. An element $a \in R$ is called right decomposable if $a = b_1c_1 = b_2c_2$ for some nonunits $b_1, b_2, c_1, c_2 \in R$ such that $Rc_1 + Rc_2 = R$. An element is called right indecomposable if it is not right decomposable. The notion of left decomposable (indecomposable) is defined similarly. It is known that any PID R is a UFD, that is, every nonunit $a \in R$ can be written as

$$a = a_1 a_2 \cdots a_n,$$

where a_1, \dots, a_n are irreducible, and moreover, if $a = b_1 \cdots b_m$, where b_1, \dots, b_m are irreducible, then $m = n$, and there is a permutation $\tau \in S_n$ such that each b_i is similar to $a_{\tau(i)}$.

5.1 Partial fraction decomposition in a PID

In this subsection, R is a fixed PID. It is known that R is an Ore domain, hence has a field of fractions $F = \text{Frac}(R)$. A fraction $ab^{-1} \in F$, where $a, b \in R$, is called *reduced* if a and b are right co-prime, that is, $Ra + Rb = R$.

Lemma 5.1. *Any $x \in F$ has at least one reduced fraction $x = ab^{-1}$ which is unique up to multiplication by units. More precisely, if $ab^{-1} = a_1b_1^{-1}$ are two reduced fractions, then $a_1 = au$ and $b_1 = bu$ for some unit $u \in R$. More generally, if $ab^{-1} = a_1b_1^{-1}$ and ab^{-1} is reduced, then there exists $r \in R$ such that $a_1 = ar$ and $b_1 = br$*

Proof. Let $x = ab^{-1}$ be an arbitrary element of F . We have $Ra + Rb = Rc$ for some $c \in R$ because R is a PID. It follows that $a = a_1c$ and $b = b_1c$ for some $a_1, b_1 \in R$. Therefore, $Ra_1 + Rb_1 = R$ which implies that $x = a_1b_1^{-1}$ is

a reduced fraction. This proves the existence part. Now, let $ab^{-1} = a_1b_1^{-1}$, where ab^{-1} is reduced, that is, $xa + yb = 1$ for some $x, y \in R$. Since R is an Ore domain, $bR \cap b_1R \neq \{0\}$ from which it follows that there exist $c, c_1 \in R^*$ such that $bc = b_1c_1$. We have

$$ab^{-1} = a_1b_1^{-1} \implies (ab^{-1})(bc) = (a_1b_1^{-1})(b_1c_1) \implies ac = a_1c_1.$$

Therefore, we obtain

$$c = (xa + yb)c = xac + ybc = xa_1c_1 + yb_1c_1 = (xa_1 + yb_1)c_1.$$

Setting $r = xa_1 + yb_1$, we obtain

$$a_1c_1 = ac = arc_1 \implies a_1 = ar.$$

In a similar way, one can show that $b_1 = br$. This completes the proof of the lemma. \square

The following result can be regarded as a generalization of the method of partial fraction decomposition, see [19] for the case of a commutative principal ideal domain.

Theorem 5.2. *Every fraction of R can be written as a sum of fractions of the form ab^{-1} , where $a, b \in R$ with $0 \neq b$ being right indecomposable.*

Proof. Let S be the set of all fractions that cannot be written in the stated form. We need to show that S is empty. Assume, on the contrary, that S is nonempty. Consider the set I of all ideals of the form Rb where $ab^{-1} \in S$ for some nonunits $a \in R$ and $b \in R^*$. Since R is a PID, I has a maximal element Rb_0 with respect to inclusion. Then $ab_0^{-1} \in S$ for some $a \in R$. Note that b_0 cannot be right indecomposable since otherwise $ab_0^{-1} \notin S$. So, b_0 must be right decomposable, that is, we have $b_0 = b_1c_1 = b_2c_2$ for some $b_1, c_1, b_2, c_2 \in R$ such that $Rc_1 + Rc_2 = R$. In particular, there exist $a_1, a_2 \in R$ such that $a_1c_1 + a_2c_2 = a$. Moreover, $b_0R \subsetneq b_1R$ and $b_0R \subsetneq b_2R$ since c_1 and c_2 are non-units. Now, we have

$$a = a_1c_1 + a_2c_2 \implies ab_0^{-1} = a_1b_1^{-1} + a_2b_2^{-1}.$$

By the maximality of Rb_0 , neither $a_1b_1^{-1}$ nor $a_2b_2^{-1}$ can belong to S . Therefore, both $a_1b_1^{-1}$ and $a_2b_2^{-1}$ can be written in the desired form. Consequently, ab_0^{-1} which is the sum of these two fractions can be written in the desired form, a contradiction. This complete the proof. \square

5.2 Partial fraction decomposition in $K(T; \sigma)$

Let (K, σ) be an inversive σ -field. It is known that $K[T; \sigma]$ is a PID. Its field of fractions is denoted by $K(T; \sigma)$. In particular, we can apply Theorem 5.2 to $K[T; \sigma]$. In fact, we have the following stronger result in the case of $K(T; \sigma)$.

Theorem 5.3. *Every fraction $P(T)Q(T)^{-1} \in K(T; \sigma)$ can be written as*

$$P(T)Q(T)^{-1} = P_0(T) + \sum_{i=1}^m P_i(T)Q_i(T)^{-1},$$

where $P_0(T), P_1(T), Q_1(T), \dots, P_n(T), Q_n(T) \in K[T; \sigma]$, $\deg P_i < \deg Q_i$ for every i , each $Q_i(T)$ is right indecomposable, and each fraction $P_i(T)Q_i(T)^{-1}$ is reduced.

Proof. Let S be the set of all partial fraction decompositions

$$P(T)Q(T)^{-1} = P_0(T) + \sum_{i=1}^m P_i(T)Q_i(T)^{-1},$$

where $P_0(T), P_1(T), Q_1(T), \dots, P_m(T), Q_m(T) \in K[T; \sigma]$, and each $Q_i(T)$ is right indecomposable of degree > 0 . By Theorem 5.2, the set S is nonempty. Therefore, we can choose

$$P(T)Q(T)^{-1} = P_0(T) + \sum_{i=1}^m P_i(T)Q_i(T)^{-1}$$

in S for which $\sum_{i=1}^m \deg P_i$ is as small as possible. I claim that this decomposition satisfies the desired conditions. Assume, on the contrary, that $\deg P_i \geq \deg Q_i$ for some i , say $i = 1$. Using the Euclidean division algorithm in $K[T; \sigma]$, we write

$$P_1(T) = P'(T)Q_1(T) + R(T),$$

where $P'(T), R(T) \in K[T; \sigma]$ and $\deg R < \deg Q$. Then, the decomposition

$$P(T)Q(T)^{-1} = (P_0(T) + P'(T)) + \left(R(T)Q_1(T)^{-1} + \sum_{i=2}^m P_i(T)Q_i(T)^{-1} \right),$$

belongs to S , which is a contradiction since

$$\deg R + \sum_{i=2}^m \deg P_i < \deg Q + \sum_{i=2}^m \deg P_i \leq \sum_{i=1}^m \deg P_i.$$

This completes the proof of the claim. If some $P_i(T)Q_i(T)^{-1}$ is not reduced, we can replace $P_i(T)Q_i(T)^{-1}$ by a reduced fraction whose numerator has a smaller degree than the degree of $P_i(T)$, contradicting the choice of the partial fraction decomposition. This completes the proof. \square

Next, we show that a stronger result is valid in the case of 2-algebraically closed σ -fields.

Theorem 5.4. *Let (K, σ) be a 2-algebraically closed inversive σ -field. Any fraction $P(T)Q(T)^{-1} \in K(T; \sigma)$ has a partial fraction decomposition of the form*

$$P(T)Q(T)^{-1} = P_0(T) + \sum_{i=1}^m a_i T^{-i} + \sum_{i=1}^n b_i (1 - c_i T)^{-1},$$

where $P_0(T) \in K[T; \sigma]$, $a_1, \dots, a_m \in K$, $b_1, \dots, b_n, c_1, \dots, c_n \in K^*$, and c_1, \dots, c_n are (right) P -independent. Moreover, if

$$P(T)Q(T)^{-1} = P_1(T) + \sum_{i=1}^{m_1} a'_i T^{-i} + \sum_{i=1}^{n_1} b'_i (1 - c'_i T)^{-1},$$

is another such decomposition, then we have $P_0(T) = P_1(T)$, $m = m_1$, and $a_i = a'_i$ for all i .

Proof. First, we show that a nonconstant polynomial $P(T) \in K[T; \sigma]$ is right indecomposable iff $P(T) = aT^m$ for some $m > 0$ and $a \in K^*$, or $P(T) = a(1 - cT)$ for some $a, c \in K^*$. To prove the "only if" part of this statement, assume that $P(T) \in K[T; \sigma]$ is right indecomposable. There is $m \geq 0$ and $Q(T) \in K[T; \sigma]$ with a nonzero constant term such that $P(T) = Q(T)T^m$. Since (K, σ) is inversive, there exists $Q_1(T) \in K[T; \sigma]$ with a nonzero constant term such that

$$P(T) = Q(T)T^m = T^m Q_1(T).$$

Since $Q_1(T)$ and T^m are right co-prime, m must be zero or $Q_1(T)$ must be a constant. In the latter case, we have $P(T) = aT^m$ for some $a \in K^*$. In the former case, $Q_1(T)$ must be a linear skew polynomial, since otherwise the fact that (K, σ) is 2-algebraically closed implies that $Q_1(T)$ has at least two distinct roots, say $a_1 \neq a_2$. It follows that

$$Q(T) = R_1(T)(T - a_1) = R_2(T)(T - a_2),$$

for some polynomials $R_1(T), R_2(T) \in K[T; \sigma]$. Since $T - a_1$ and $T - a_2$ are right co-prime, $Q(T)$ must be right decomposable, a contradiction. This completes the proof of the "only if" part. Conversely, it is clear that any linear skew polynomial is right indecomposable. Since $T^m = P(T)Q(T)$ implies that $P(T) = aT^n$ and $Q(T) = (1/a)T^{m-n}$ for some $0 \neq n \neq m$, we see that T^m is right indecomposable. This completes the proof of the statement. Now, we can prove the theorem: It follows from the above statement and Theorem 5.3 that any $P(T)Q(T)^{-1} \in K(T; \sigma)$ has a partial fraction decomposition of the form

$$P(T)Q(T)^{-1} = P_0(T) + \sum_{i=1}^m a_i T^{-i} + \sum_{i=1}^n b_i (1 - c_i T)^{-1},$$

where $P_0(T) \in K[T; \sigma]$, $a_1, \dots, a_m, b_1, \dots, b_n \in K$, and the elements c_1, \dots, c_n belong to K^* . Using Proposition 2.5, it is now easy to show that there is such a partial fraction decomposition which, in addition, satisfies the requirement that c_1, \dots, c_n be P-independent.

To prove the uniqueness part, we first note that the summand $\sum_{i=1}^m a_i T^{-i}$ does not belong to $K[[T; \sigma]]^*$ while the other summands belong to $K[[T; \sigma]]$, from which the equality $\sum_{i=1}^m a_i T^{-i} = \sum_{i=1}^{m_1} a'_i T^{-i}$ follows. Finally, using the degree function on $K(T; \sigma)$, one can easily show that $P_1(T)$ must be equal to $P_0(T)$. \square

5.3 The twisted Hadamard algebra

We conclude this paper with an application for which we need to recall the definition of the twisted Hadamard algebra over a σ -field. For more details, see [1]. Let (K, σ) be an inversive σ -field. An element in $K[[T, \sigma]]$ is called a rational twisted series if it belongs to the subring $K(T; \sigma)$. A rational twisted series is called regular if its degree is negative. The set of all regular rational series over (K, σ) is denoted by $H(T; \sigma)$. In [1], it is shown that the set $H(T; \sigma)$ equipped with the addition operation inherited from $K[[T, \sigma]]$ and the Hadamard multiplication \odot , that is,

$$\left(\sum_{i=0}^{\infty} a_i T^i \right) \odot \left(\sum_{i=0}^{\infty} b_i T^i \right) = \sum_{i=0}^{\infty} a_i b_i T^i,$$

is a K -algebra, called the twisted Hadamard algebra over (K, σ) . Here, we give a characterization of the twisted Hadamard algebra in the case when (K, σ) is 2-algebraically closed. For a characterization of the ordinary Hadamard algebra in terms of exponential polynomials, see [2]. In the following theorem, we use the K -algebra N_{K^*} introduced in Subsection 2.1.

Theorem 5.5. *There exists a unique K -algebra homomorphism*

$$\alpha : N_{K^*} \rightarrow H(T; \sigma),$$

which satisfies $\alpha(N(a)) = (1 - aT)^{-1}$ for all $a \in K^$. Furthermore, α is injective, and, if K is a 2-algebraically closed inversive σ -field, α is an isomorphism of K -algebras.*

Proof. The vector space K^∞ , as defined in Subsection 2.1, is a K -algebra under pointwise multiplication. Consider the map $\beta : K^\infty \rightarrow K[[T; \sigma]]$ defined by

$$\beta(a_0, a_1, a_2, \dots) = \sum_{i=0}^{\infty} a_i T^i.$$

It is easy to see that β is a K -algebra homomorphism where $K[[T; \sigma]]$ is equipped with the Hadamard product. Let $\alpha : N_{K^*} \rightarrow K[[T; \sigma]]$ be the

restriction of β to N_{K^*} . Note that $(1 - aT)^{-1} \in H(T; \sigma)$, for all $a \in K^*$. Since N_{K^*} , as a vector space, is generated by $N(a)$'s, and $H(T; \sigma)$ is closed under the Hadamard product, we see that the image of α is contained in $H(T; \sigma)$. The uniqueness of α follows from the fact that the elements $N(a)$, $a \in K$, generate N_{K^*} as a vector space over K . Clearly, α is injective. The fact that α is surjective when K is a 2-algebraically closed inversive σ -field follows from Theorem 5.4. □

As a consequence of this result, we have the following:

Corollary 5.6. *Let $\sum_{i=0}^{\infty} a_i T^i$ be a rational twisted series over a 2-algebraically closed inversive σ -field K . Then, there exist element $b_1, \dots, b_n, c_1, \dots, c_n \in K^*$ such that*

$$a_i = \sum_{j=1}^n b_j N_i(c_j),$$

for all large enough i .

Proof. Since every rational twisted series is a sum of a skew polynomial and a regular series, the result follows from Theorem 5.5. □

Acknowledgements

I would like to thank A. Leroy for his valuable suggestions and comments which led to an improvement of the paper.

References

- [1] Masood Aryapoor. Rational twisted series. *arXiv preprint arXiv:2202.11131*, 2022.
- [2] Jean Berstel and Christophe Reutenauer. *Rational series and their languages*, volume 12. Springer-Verlag, 1988.
- [3] Vladimir Bolotnikov. Lagrange interpolation over division rings. *Communications in Algebra*, 48(9):4065–4084, 2020.
- [4] Una Bray and George Whaples. Polynomials with coefficients from a division ring. *Canadian Journal of Mathematics*, 35(3):509–515, 1983.
- [5] Chen-Lian Chuang. Automorphisms of ore extensions. *Israel Journal of Mathematics*, 197(1):437–452, 2013.
- [6] Paul M Cohn. Noncommutative unique factorization domains. *Transactions of the American Mathematical Society*, 109(2):313–331, 1963.

- [7] Paul M Cohn. *Skew Fields: Theory of General Division Rings*. Encyclopedia of Mathematics and its Applications, 57. Cambridge University Press, Cambridge, 1995.
- [8] Paul Moritz Cohn. Unique factorization domains. *The American Mathematical Monthly*, 80(1):1–18, 1973.
- [9] PM Cohn. Presentations of skew fields. i. existentially closed skew fields and the nullstellensatz. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 77, pages 7–19. Cambridge University Press, 1975.
- [10] Jonathan Delenclos and André Leroy. Noncommutative symmetric functions and w-polynomials. *Journal of Algebra and its Applications*, 6(05):815–837, 2007.
- [11] I.M. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V.S. Retakh, and J.Y. Thibon. Noncommutative symmetric functions. *Advances in Mathematics*, 112(2):218–348, 1995.
- [12] Israel Gelfand and Vladimir Retakh. Noncommutative vieta theorem and symmetric functions. In *The Gelfand Mathematical Seminars, 1993–1995*, pages 93–100. Springer, 1996.
- [13] Tsit-Yuen Lam. *A first course in noncommutative rings*, volume 131. Springer, 1991.
- [14] Tsit-Yuen Lam and André Leroy. Vandermonde and wronskian matrices over division rings. *Journal of Algebra*, 119(2):308–336, 1988.
- [15] Tsit Yuen Lam and André Leroy. Wedderburn polynomials over division rings, i. *Journal of Pure and Applied Algebra*, 186(1):43–76, 2004.
- [16] T.Y. Lam. A general theory of vandermonde matrices. *Exposition. Math.*, 4(3):193–215, 1986.
- [17] TY Lam and A Leroy. Hilbert 90 theorems over division rings. *Transactions of the American Mathematical Society*, 345(2):595–622, 1994.
- [18] TY Lam and André Leroy. Algebraic conjugacy classes and skew polynomial rings. In *Perspectives in ring theory*, pages 153–203. Springer, 1988.
- [19] Serge Lang. *Algebra*, volume 211. Springer Science & Business Media, 2012.
- [20] Alexander Levin. *Difference algebra*, volume 8. Springer Science & Business Media, 2008.

- [21] Bernhard Hermann Neumann. Adjunction of elements to groups. *Journal of the London Mathematical Society*, 1(1):4–11, 1943.
- [22] Ivan Niven. Equations in quaternions. *The American Mathematical Monthly*, 48(10):654–661, 1941.
- [23] Oystein Ore. Theory of non-commutative polynomials. *Annals of mathematics*, pages 480–508, 1933.
- [24] Abraham Robinson. On the notion of algebraic closedness for noncommutative groups and fields. *The Journal of Symbolic Logic*, 36(3):441–444, 1971.
- [25] William R Scott. Algebraically closed groups. *Proceedings of the American Mathematical Society*, 2(1):118–121, 1951.
- [26] Jan Treur. Separate zeros and galois extensions of skew fields. *Journal of Algebra*, 120(2):392–405, 1989.