

Existence and convergence of commutative mappings on some results of fixed point theory in a class of generalized non-expansive mappings

Gezahegn Anberber Tadesse Department of Mathematics, College of Computational and Natural Science, Addis Ababa University, Ethiopia.

Abstract

In this paper, we introduce a commutative mappings satisfying the class of generalized nonexpansive mappings which is wider than the class of mappings satisfying the condition (C), so called Condition $B_{\gamma,\mu}$. The results obtained in this paper extend and generalized non-expansive mappings and other results in this direction. Different properties and some fixed point results for the mappings are obtained here.

Keywords: Nonexpansive mappings, Condition (C), Condition $B_{\gamma,\mu}$, Commuting mappings, Fixed point.

1 Introduction

Fixed point theory and its application played an important role in many areas of applied science and solved many problems rising in engineering, mathematical economics and optimization. The generalization of non-expansive mappings and the study of related fixed point theorems with different practical applications in nonlinear functional analysis have found great importance during the recent decades [1, 3, 2, 4, 5, 13, 7, 8, 11, 9, 13, 14, 15, 16, 18, 21]. Several prominent authors [6, 10, 12, 9, 15, 16, 20] have contributed immensely in this field, and different new classes of mappings with interesting properties have been developed in this context.

Suzuki [20] introduce the concept of generalized nonexpansive mappings which is called Suzuki generalized nonexpansive mappings or the condition (C) and received some fixed point results and convergence results for such mapping in Banach space. Recently in 2018, Patir et al. [17] defined a class of generalized nonexpansive mappings on a nonempty subset C of a Banach space X which is wider than Suzuki's generalized nonexpansive mappings. Such type of mappings was called the class of mappings satisfying the condition $B_{\gamma,\mu}$ on C . (see [17]). For a nonempty bounded and convex subset C , every self-mapping T on C satisfying $B_{\gamma,\mu}$ condition has an almost fixed point sequence [8], but preserving their fixed point properties.

Having these inspiring background, we can ask:

Problem 1 *Is that extend the existence and converge theorem of the self mappings T of the condition $B_{\gamma,\mu}$ to the finite commutative mappings on this condition?*

To answer this, we can apply some facts related to the condition $B_{\gamma,\mu}$ on a non empty subset of a Banach space X .

2 Preliminaries

We now recall the following definitions, propositions and lemmas to be use the main result.

Definition 1 *Let C be a nonempty subset of a Banach space X . A mapping $T : C \rightarrow X$ is said to be **non-expansive** if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$.*

Definition 2 *For a nonempty subset C of a Banach space X , a mapping $T : C \rightarrow X$ is called **quasi-nonexpansive** if $\|Tx - z\| \leq \|x - z\|$ for all $x \in C$ and $z \in F(T)$ (where $F(T)$ denotes the set of all fixed points of T).*

Definition 3 ([20]) *For a nonempty subset C of a Banach space X , a mapping $T : C \rightarrow X$ is said to satisfy **the condition (C)** on C if $\frac{1}{2}\|x - Tx\| \leq \|x - y\|$ implies $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$.*

Remark 1 *Every nonexpansive mapping satisfies the condition (C) on C . But there are also some noncontinuous mappings satisfying the condition C , (see [20]).*

Definition 4 ([20]) *For a nonempty subset C of a Banach space X and $\lambda \in (0, 1)$, a mapping $T : C \rightarrow X$ is said to satisfy **(C_λ) -condition** on C if $\lambda\|x - Tx\| \leq \|x - y\|$ implies $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$.*

Definition 5 ([17]) Let C be a nonempty subset of a Banach space X . Let $\gamma \in [0, 1]$ and $\mu \in [0, \frac{1}{2}]$ such that $2\mu \leq \gamma$. A mapping $T : C \rightarrow X$ is said to satisfy **the condition** $B_{\gamma, \mu}$ on C if, for all x, y in C , $\gamma\|x - Tx\| \leq \|x - y\| + \mu\|y - Ty\|$ implies $\|Tx - Ty\| \leq (1 - \gamma)\|x - y\| + \mu(\|x - Ty\| + \|y - Tx\|)$.

Since, this class includes the class of nonexpansive mappings (for $\gamma = \mu = 0$).

Remark 2 Let C be a nonempty subset of a Banach space X . If a mapping satisfies the condition (C), then it will satisfy the condition $B_{\gamma, \mu}$ for $\gamma = \mu = 0$.

Definition 6 Let C be a nonempty closed convex and bounded subset of a Banach space X and a self mapping T on C is satisfies condition $B_{\gamma, \mu}$ on C , then there exists a sequence $\{x_n\}$ such that $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$. Such a sequence is called almost fixed point sequence for T .

In 2005, Suzuki ([19]) proved the mappings T_1 and T_2 are non-expansive mapping with $T_1 \circ T_2 = T_2 \circ T_1$ by **Lemma 3.1** have a common fixed point of T_1 and T_2 . In similar way to extend the prove of finite commutative mappings for some class of generalized non-expansive have a common fixed point.

The following are some basic properties of mappings which satisfy the condition $B_{\gamma, \mu}$ on C .

Proposition 1 ([17]) Let C be a nonempty subset of a Banach space X . Let $T : C \rightarrow C$ satisfy the condition $B_{\gamma, \mu}$ on C . Then, for all $x, y \in C$ and for $\theta \in [0, 1]$,

$$(i) \|Tx - T^2x\| \leq \|x - Tx\|,$$

(ii) at least one of the following ((a) and (b)) holds:

$$(a) \frac{\theta}{2}\|x - Tx\| \leq \|x - y\|,$$

$$(b) \frac{\theta}{2}\|Tx - T^2x\| \leq \|Tx - y\|.$$

The condition (a) implies $\|Tx - Ty\| \leq (1 - \frac{\theta}{2})\|x - y\| + \mu(\|x - Ty\| + \|y - Tx\|)$ and the condition (b) implies $\|T^2x - Ty\| \leq (1 - \frac{\theta}{2})\|Tx - y\| + \mu(\|Tx - Ty\| + \|y - T^2x\|)$.

(iii)

$$\begin{aligned} \|x - Ty\| &\leq (3 - \theta)\|x - Tx\| + (1 - \frac{\theta}{2})\|x - y\| + \mu(2\|x - Tx\| \\ &\quad + \|x - Ty\| + \|y - Tx\| + 2\|Tx - T^2x\|). \end{aligned}$$

Proposition 2 ([17]) Let C be a nonempty convex and bounded subset of a Banach space X and T be a self-mapping on C . We assume that T satisfies the condition $B_{\gamma, \mu}$ on C . For $x_0 \in C$, let a sequence $\{x_n\}$ in C be defined by;

$$x_{n+1} = \lambda Tx_n + (1 - \lambda)x_n, \tag{1}$$

where $\lambda \in [\gamma, 1) - \{0\}$, $n \in \mathbb{N} \cup \{0\}$. Then $\|Tx_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty$.

3 Main results

In this section, we prove our main results. That is, we prove the convergence and the existence theorem of fixed point of commuting mappings satisfying the condition $B_{\gamma, \mu}$. First, we give the following lemma, which will play an important role in the sequel.

Lemma 1 Let C be a nonempty closed convex subset of a Banach space X . Let T_1 and T_2 be mappings with $T_1 \circ T_2 = T_2 \circ T_1$ satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $(0, \frac{1}{2})$ converging to 0, and let $\{x_n\}$ be a sequence in C such that $\{x_n\}$ converges strongly to some $z \in C$. Then z is a common fixed point of T_1 and T_2 .

Proof. Since a sequence $\{x_n\}$ in C converges strongly to some $z \in C$. And by the condition $B_{\gamma, \mu}$ we get

$$\|T_1x_n - T_1z\| \leq (1 - \gamma)\|x_n - z\| + \mu\left(\|x_n - T_1z\| + \|z - T_1x_n\|\right). \tag{2}$$

Again by using (2)

$$\begin{aligned}
\|x_n - T_1 z\| &\leq \|x_n - T_1 x_n\| + \|T_1 x_n - T_1 z\| \\
&\leq \|x_n - T_1 x_n\| + (1 - \gamma)\|x_n - z\| + \mu\left(\|x_n - T_1 z\| + \|z - T_1 x_n\|\right) \\
&\leq \|x_n - T_1 x_n\| + (1 - \gamma)\|x_n - z\| + \mu\left(\|x_n - T_1 z\| + \|z - x_n\| + \|x_n - T_1 x_n\|\right).
\end{aligned}$$

It implies that

$$(1 - \mu)\|x_n - T_1 z\| \leq (1 + \mu)\|T_1 x_n - x_n\| + (1 - \gamma + \mu)\|x_n - z\|.$$

So taking $n \rightarrow \infty$ and using Proposition 2, we get

$$(1 - \mu)\|z - T_1 z\| \leq 0,$$

since $\mu \neq 1$. It implies that

$$T_1 z = z \quad (3)$$

This shows that z is a fixed point for T_1 . We note that

$$T_1 \circ T_2 z = T_2 \circ T_1 z = T_2 z. \quad (4)$$

We assume that z is not a fixed point of T_2 . Put

$$\epsilon = \frac{\|T_2 z - z\|}{3} > 0.$$

Then there exists $m \in \mathbb{N}$ such that

$$\|x_m - z\| < \epsilon, \quad \frac{\|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\|}{\alpha_m} < \epsilon.$$

Since,

$$\begin{aligned}
3\epsilon = \|T_2 z - z\| &\leq \|T_2 z - x_m\| + \|x_m - z\| \\
&< \|T_2 z - x_m\| + \epsilon \\
\Rightarrow 2\epsilon &< \|T_2 z - x_m\|.
\end{aligned}$$

So, by using (4) and Proposition 1, (i) for $\gamma = \frac{\theta}{2}, \theta \in [0, 1]$, we obtain, we obtain

$$\begin{aligned}
\|T_2 z - x_m\| &\leq \|T_2 z - (1 - \alpha_m)T_1 x_m - \alpha_m T_2 x_m\| + \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&\leq (1 - \alpha_m)\|T_2 z - T_1 x_m\| + \alpha_m\|T_2 z - T_2 x_m\| + \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&= (1 - \alpha_m)\|T_2 \circ T_1 z - T_1 x_m\| + \alpha_m\|T_2 z - T_2 x_m\| \\
&\quad + \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&\leq (1 - \alpha_m)\left[(1 - \gamma)\|T_2 z - x_m\| + \mu(\|T_2 z - T_1 x_m\| + \|x_m - T_2 z\|)\right] + \\
&\quad \alpha_m\left[(1 - \gamma)\|z - x_m\| + \mu(\|z - T_2 x_m\| + \|x_m - T_2 z\|)\right] + \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&\leq (1 - \alpha_m)\left[(1 - \gamma)\|T_2 z - x_m\| + \mu(\|T_2 z - T_1 x_m\| + \|x_m - T_2 z\|)\right] + \\
&\quad \alpha_m\left[(1 - \gamma)\|z - x_m\| + \mu\|z - x_m\| + \mu\|x_m - z\|\right] + \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&\leq (1 - \alpha_m)\left[\frac{1 - \gamma + \mu}{1 - \mu}\right]\|T_2 z - x_m\| + \alpha_m\|z - x_m\| + \\
&\quad \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&\leq (1 - \alpha_m)\|T_2 z - x_m\| + \alpha_m\|z - x_m\| + \|(1 - \alpha_m)T_1 x_m + \alpha_m T_2 x_m - x_m\| \\
&< (1 - \alpha_m)\|T_2 z - x_m\| + \alpha_m \epsilon + \alpha_m \epsilon \\
&= (1 - \alpha_m)\|T_2 z - x_m\| + 2\alpha_m \epsilon \\
&< (1 - \alpha_m)\|T_2 z - x_m\| + \alpha_m\|T_2 x_m - x_m\| \\
&= \|T_2 z - x_m\|,
\end{aligned}$$

since $\frac{1 - \gamma + \mu}{1 - \mu} \leq 1$, which is a contradiction. Hence, z is a common fixed point of T_1 and T_2 ■

Lemma 2 Let C be a nonempty closed convex and bounded subset of a Banach space X . Let T_1 and T_2 be mappings with $T_1 \circ T_2 = T_2 \circ T_1$ satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $[0, \frac{1}{2}]$ converging to 0. Define a sequence $\{x_n\}$ in C by $x_0 \in C$ and

$$x_{n+1} = \lambda(1 - \alpha_n)T_1x_n + \lambda\alpha_nT_2x_n + (1 - \lambda)x_n, \quad (5)$$

for $\lambda \in [\gamma, 1) - \{0\}$, $n = 0, 1, 2, \dots$. Then

$$\lim_{n \rightarrow \infty} \|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n - x_n\| = 0. \quad (6)$$

Proof. Since a sequence $\{x_n\}$ in C defined as (5) and assume that z is a common fixed point of T_1 and T_2 , then, by condition $B_{\gamma, \mu}$ we get

$$\begin{aligned} \|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n - x_n\| &\leq \lambda\|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n - x_n\| \\ &= \|x_{n+1} - x_n\| \\ &\leq \|x_{n+1} - x_n\| + \mu\|x_n - T_1x_n\|. \end{aligned}$$

Implies that

$$\begin{aligned} &\|(1 - \alpha_{n+1})T_1x_{n+1} + \alpha_{n+1}T_2x_{n+1} - T_1x_n\| \leq (1 - \alpha_{n+1})\|T_1x_{n+1} - T_1x_n\| + \alpha_{n+1}\|T_2x_{n+1} - T_1x_n\| \\ &\leq (1 - \alpha_{n+1})\left[\|T_1x_{n+1} - T_1z\| + \|T_1z - T_1x_n\|\right] + \alpha_{n+1}\left[\|T_2x_{n+1} - T_2z\| + \|T_2z - T_1x_n\|\right] \\ &\leq (1 - \alpha_{n+1})\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|x_{n+1} - T_1z\| + \|z - T_1x_{n+1}\|) + (1 - \gamma)\|x_n - z\| + \mu(\|x_n - T_1z\| + \|z - T_1x_n\|)\right] \\ &\quad + \alpha_{n+1}\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|x_{n+1} - T_2z\| + \|z - T_2x_{n+1}\|) + (1 - \gamma)\|x_n - z\| + \mu(\|x_n - T_2z\| + \|z - T_2x_n\|)\right] \\ &\leq (1 - \alpha_{n+1})\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|x_{n+1} - z\| + \|z - x_{n+1}\|) + (1 - \gamma)\|x_n - z\| + \mu(\|x_n - z\| + \|z - x_n\|)\right] + \\ &\quad \alpha_{n+1}\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|x_{n+1} - z\| + \|z - x_{n+1}\|) + (1 - \gamma)\|x_n - z\| + \mu(\|x_n - z\| + \|z - x_n\|)\right] \\ &\leq (1 - \alpha_{n+1})\left[\|x_{n+1} - z\| + \|x_n - z\|\right] + \alpha_{n+1}\left[\|x_{n+1} - z\| + \|x_n - z\|\right] \\ &= \|x_{n+1} - z\| + \|x_n - z\|. \end{aligned}$$

Now from the following we have that,

$$\|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n - x_n\| \leq (1 - \alpha_n)\|T_1x_n - x_n\| + \alpha_n\|T_2x_n - x_n\|.$$

Taking $n \rightarrow \infty$ on both side of the above inequality and using Proposition 2 we get

$$\lim_{n \rightarrow \infty} \|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n - x_n\| = 0.$$

For a nonempty compact convex subset C of X , we have the following fixed point result.

Theorem 1 Let C be a compact and convex subset of a Banach space X . Let T_1 and T_2 be mappings with $T_1 \circ T_2 = T_2 \circ T_1$ satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $[0, \frac{1}{2}]$ converging to 0. For $x_0 \in C$, let $\{x_n\}$ be a sequence in C as defined by (5). Then $\{x_n\}$ converges strongly to a common fixed point T_1 and T_2 .

Proof. Since C is compact, there exists a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ and $z \in C$ such that $\{x_{n_j}\}$ converges to $z \in C$ By (Lemma 1)

Now by Proposition 1 (ii), for $\gamma = \frac{\theta}{2}$, $\theta \in [0, 1]$,

$$\begin{aligned} \gamma\|x_{n_j} - T_1x_{n_j}\| &\leq \|x_{n_j} - z\| \\ \Rightarrow \gamma\|x_{n_j} - T_1x_{n_j}\| &\leq \|x_{n_j} - z\| + \mu\|z - T_1z\|. \end{aligned}$$

So by condition $B_{\gamma, \mu}$

$$\|T_1x_{n_j} - T_1z\| \leq (1 - \gamma)\|x_{n_j} - z\| + \mu(\|x_{n_j} - T_1z\| + \|z - T_1x_{n_j}\|) \quad (7)$$

Again, by using (7) we have

$$\begin{aligned}
\|x_{n_j} - T_1 z\| &\leq \|x_{n_j} - T_1 x_{n_j}\| + \|T_1 x_{n_j} - T_1 z\| \\
&\leq \|x_{n_j} - T_1 x_{n_j}\| + (1 - \gamma)\|x_{n_j} - z\| + \mu(\|x_{n_j} - T_1 z\| + \|z - T_1 x_{n_j}\|) \\
&\leq \|x_{n_j} - T_1 x_{n_j}\| + (1 - \gamma)\|x_{n_j} - z\| + \mu(\|x_{n_j} - T_1 z\| + \|z - x_{n_j}\| + \|x_{n_j} - T_1 x_{n_j}\|).
\end{aligned}$$

So taking $n_j \rightarrow \infty$ and using Proposition 2, we get

$$\begin{aligned}
(1 - \mu)\|z - T_1 z\| &\leq 0 \\
\Rightarrow Tz &= z,
\end{aligned}$$

since $\mu \neq 1$, shows that z is a fixed point T_1 . Similarly by applying Lemma 1, we have that $z = T_2 z$. Hence, z is a common fixed point of T_1 and T_2 .

Now by the condition $B_{\gamma, \mu}$ we have

$$\begin{aligned}
\|T_1 x_n - z\| &= \|T_1 x_n - T_1 z\| \\
&\leq (1 - \gamma)\|x_n - z\| + \mu(\|x_n - T_1 z\| + \|z - T_1 x_n\|) \\
&= (1 - \gamma)\|x_n - z\| + \mu\|x_n - z\| + \mu\|z - T_1 x_n\|.
\end{aligned}$$

Which implies that

$$\|T_1 x_n - z\| \leq \left(\frac{1 - \gamma + \mu}{1 - \mu}\right)\|x_n - z\| \leq \|x_n - z\|, \quad (8)$$

since $2\mu \leq \gamma$. Similarly we have that $\|T_2 x_n - z\| \leq \|x_n - z\|$. Now by using (5) and (8), we have that:

$$\begin{aligned}
\|x_{n+1} - z\| &= \|\lambda(1 - \alpha_n)T_1 x_n + \lambda\alpha_n T_2 x_n + (1 - \lambda)x_n - z\| \\
&\leq \lambda(1 - \alpha_n)\|T_1 x_n - T_1 z\| + \lambda\alpha_n\|T_2 x_n - T_2 z\| + (1 - \lambda)\|x_n - z\| \\
&\leq \lambda(1 - \alpha_n)\left[(1 - \gamma)\|x_n - z\| + \mu(\|x_n - T_1 z\| + \|z - T_1 x_n\|)\right] + \\
&\quad \lambda\alpha_n\left[(1 - \gamma)\|x_n - z\| + \mu(\|x_n - T_2 z\| + \|z - T_2 x_n\|)\right] + (1 - \lambda)\|x_n - z\| \\
&\leq \lambda(1 - \alpha_n)\left[(1 - \gamma)\|x_n - z\| + 2\mu\|x_n - z\|\right] + \\
&\quad \lambda\alpha_n\left[(1 - \gamma)\|x_n - z\| + 2\mu\|x_n - z\|\right] + (1 - \lambda)\|x_n - z\| \\
&\leq \lambda(1 - \alpha_n)\left[(1 - \gamma)\|x_n - z\| + \gamma\|x_n - z\|\right] + \\
&\quad \lambda\alpha_n\left[(1 - \gamma)\|x_n - z\| + \gamma\|x_n - z\|\right] + (1 - \lambda)\|x_n - z\| \\
&= \lambda(1 - \alpha_n)\|x_n - z\| + \lambda\alpha_n\|x_n - z\| + (1 - \lambda)\|x_n - z\| \\
&= \|x_n - z\|.
\end{aligned}$$

Thus, $\{\|x_n - z\|\}$ is a monotonically decreasing non-negative sequence of real numbers, and will converges to some real numbers, say u .

Now by using (??) we get,

$$\begin{aligned}
\|x_n - z\| = \|x_n - T_1 z\| &\leq \|x_n - T_1 x_n\| + (1 - \gamma)\|x_n - z\| + \mu\left(\|x_n - T_1 z\| + \|z - T_1 x_n\|\right) \\
&\leq \|x_n - T_1 x_n\| + (1 - \gamma)\|x_n - z\| + \mu\left(2\|x_n - z\| + \|x_n - T_1 x_n\|\right),
\end{aligned}$$

since $z = T_1 \circ T_2 z = T_2 \circ T_1 z = T_1 z = T_2 z$. Then it implies that

$$\|x_n - z\| \leq \|x_n - T_1 x_n\| + (1 - \gamma)\|x_n - z\| + \mu\left(2\|x_n - z\| + \|x_n - T_1 x_n\|\right) \quad (9)$$

Taking the limit as $n \rightarrow \infty$ of (9) and using Proposition 2 we get $u \leq (1 - \gamma)u + \mu(2u)$ implies that $(\gamma - 2\mu)u \leq 0$, which is possible only for $u = 0$, since $2\mu \leq \gamma$. Hence, $\{x_n\}$ converges strongly to the common fixed point of T_1 and T_2 . \blacksquare

Lemma 3 Let C be a nonempty closed convex subset of a Banach space X . Let T_1, T_2 and T_3 be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $(0, \frac{1}{2})$ converging to 0, and let $\{x_n\}$ be a sequence in C such that $\{x_n\}$ converges strongly to some $z \in C$. Then z a common fixed point of T_1, T_2 and T_3 .

Proof. Assume that a sequence $\{x_n\}$ in C converges strongly to some $z \in C$. Since by Lemma 1, we have that z is a common fixed point of T_1 and T_2 . And we note that;

$$T_1 \circ T_3 z = T_3 \circ T_1 z = T_3 z, \quad T_2 \circ T_3 z = T_3 \circ T_2 z = T_3 z. \quad (10)$$

We assume that z is not a fixed point of T_3 . Put

$$\epsilon = \frac{\|T_3 z - z\|}{3} > 0.$$

Then there exists $m \in \mathbb{N}$ such that

$$\|x_m - z\| < \epsilon, \quad \frac{\|(1 - \alpha_m - \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m + \alpha_m^2 T_2 x_m - x_m\|}{\alpha_m^2} < \epsilon,$$

since,

$$\begin{aligned} 3\epsilon = \|T_3 z - z\| &\leq \|T_3 z - x_m\| + \|x_m - z\| \\ &< \|T_3 z - x_m\| + \epsilon \\ \Rightarrow 2\epsilon &< \|T_3 z - x_m\|. \end{aligned}$$

So, by using Proposition 1 for $\gamma = \frac{\theta}{2}, \theta \in [0, 1]$ and (10), we obtain

$$\begin{aligned} \|T_3 z - x_m\| &\leq \|T_3 z - (1 - \alpha_m - \alpha_m^2)T_1 x_m - \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m\| \\ &\quad + \|(1 - \alpha_m - \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m + \alpha_m^2 T_3 x_m - x_m\| \\ &\leq (1 - \alpha_m - \alpha_m^2)\|T_3 z - T_1 x_m\| + \alpha_m \|T_3 z - T_2 x_m\| + \alpha_m^2 \|T_3 z - T_3 x_m\| \\ &\quad + \|(1 - \alpha_m + \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m - x_m\| \\ &= (1 - \alpha_m - \alpha_m^2)\|T_3 \circ T_1 z - T_1 x_m\| + \alpha_m \|T_3 \circ T_2 z - T_2 x_m\| + \alpha_m^2 \|T_3 z - T_3 x_m\| \\ &\quad + \|(1 - \alpha_m + \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m - x_m\| \\ &\leq (1 - \alpha_m - \alpha_m^2) \left[(1 - \gamma)\|T_3 z - x_m\| + \mu(\|T_3 z - x_m\|) + \|x_m - T_3 z\| \right] \\ &\quad + \alpha_m \left[(1 - \gamma)\|T_3 z - x_m\| + \mu(\|T_3 z - T_2 x_m\| + \|x_m - T_3 z\|) \right] + \alpha_m^2 \|T_3 z - T_3 x_m\| \\ &\quad + \|(1 - \alpha_m + \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m - x_m\| \\ &\leq (1 - \alpha_m - \alpha_m^2) \left[\frac{1 - \gamma + \mu}{1 - \mu} \|T_3 z - x_m\| + \alpha_m \left[\frac{1 - \gamma + \mu}{1 - \mu} \|T_3 z - x_m\| + \alpha_m^2 \|z - x_m\| \right] \right. \\ &\quad \left. + \|(1 - \alpha_m + \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m - x_m\| \right] \\ &\leq (1 - \alpha_m - \alpha_m^2)\|T_3 z - x_m\| + \alpha_m \|T_3 z - x_m\| + \alpha_m^2 \|z - x_m\| \\ &\quad + \|(1 - \alpha_m + \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m - x_m\| \quad (\text{since } \frac{1 - \gamma + \mu}{1 - \mu} \leq 1) \\ &= (1 - \alpha_m^2)\|T_3 z - x_m\| + \alpha_m^2 \|z - x_m\| \\ &\quad + \|(1 - \alpha_m + \alpha_m^2)T_1 x_m + \alpha_m T_2 x_m - \alpha_m^2 T_3 x_m - x_m\| \\ &< (1 - \alpha_m^2)\|T_3 z - x_m\| + \alpha_m^2 \epsilon + \alpha_m^2 \epsilon \\ &= (1 - \alpha_m^2)\|T_3 z - x_m\| + 2\alpha_m^2 \epsilon \\ &< (1 - \alpha_m^2)\|T_3 z - x_m\| + \alpha_m^2 \|T_3 z - x_m\| \\ &= \|T_3 z - x_m\|, \end{aligned}$$

which is a contradiction. Hence, z is a common fixed point of T_1, T_2 and T_3 ■

Lemma 4 Let C be a nonempty closed convex subset of a Banach space X . Let T_1, T_2 and T_3 be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $[0, \frac{1}{2}]$ converging to 0. Define a sequence $\{x_n\}$ in C by $x_0 \in C$ and

$$x_{n+1} = \lambda(1 - \alpha_n - \alpha_n^2)T_1x_n + \lambda\alpha_nT_2x_n + \lambda\alpha_n^2T_3x_n + (1 - \lambda)x_n, \quad (11)$$

for $\lambda \in [\gamma, 1) - \{0\}$, $n = 0, 1, 2, \dots$. Then

$$\lim_{n \rightarrow \infty} \|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n + \alpha_n^2T_3x_n - x_n\| = 0. \quad (12)$$

Proof. Since a sequence $\{x_n\}$ in C defined as (11) and assume that z is a common fixed point of T_1, T_2 and T_3 , then, by condition $B_{\gamma, \mu}$ we get

$$\begin{aligned} \gamma\|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n + \alpha_n^2T_3x_n - x_n\| &\leq \lambda\|(1 - \alpha_n)T_1x_n + \alpha_nT_2x_n + \alpha_n^2T_3x_n - x_n\| \\ &= \|x_{n+1} - x_n\| \\ &\leq \|x_{n+1} - x_n\| + \mu\|x_n - T_1x_n\|. \end{aligned}$$

Implies that

$$\begin{aligned} &\|(1 - \alpha_{n+1} - \alpha_{n+1}^2)T_1x_{n+1} + \alpha_{n+1}T_2x_{n+1} + \alpha_{n+1}^2T_3x_{n+1} - T_1x_n\| \\ &\leq (1 - \alpha_{n+1} - \alpha_{n+1}^2)\|T_1x_{n+1} - T_1x_n\| + \alpha_{n+1}\|T_2x_{n+1} - T_1x_n\| + \alpha_{n+1}^2\|T_3x_{n+1} - T_1x_n\| \\ &\leq (1 - \alpha_{n+1} - \alpha_{n+1}^2)\left[\|T_1x_{n+1} - T_1z\| + \|T_1z - T_1x_n\|\right] + \alpha_{n+1}\left[\|T_2x_{n+1} - T_2z\| + \|T_2z - T_1x_n\|\right] \\ &\quad + \alpha_{n+1}^2\left[\|T_3x_{n+1} - T_3z\| + \|T_3z - T_1x_n\|\right] \\ &\leq (1 - \alpha_{n+1} - \alpha_{n+1}^2)\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|z - T_1x_{n+1}\| + \|x_{n+1} - T_1z\|) + (1 - \gamma)\|x_n - z\| \right. \\ &\quad \left. + \mu(\|z - T_1x_n\| + \|x_n - T_1z\|)\right] + \alpha_{n+1}\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|z - T_2x_{n+1}\| + \|x_{n+1} - T_2z\|) + \right. \\ &\quad \left. (1 - \gamma)\|x_n - z\| + \mu(\|z - T_1x_n\| + \|x_n - T_2z\|)\right] + \alpha_{n+1}^2\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|z - T_3x_{n+1}\| + \|x_{n+1} - T_3z\|) + \right. \\ &\quad \left. (1 - \gamma)\|x_n - z\| + \mu(\|z - T_1x_n\| + \|x_n - T_3z\|)\right] \\ &\leq (1 - \alpha_{n+1} - \alpha_{n+1}^2)\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|z - x_{n+1}\| + \|x_{n+1} - z\|) + (1 - \gamma)\|x_n - z\| \right. \\ &\quad \left. + \mu(\|z - x_n\| + \|x_n - z\|)\right] + \alpha_{n+1}\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|z - x_{n+1}\| + \|x_{n+1} - z\|) + \right. \\ &\quad \left. (1 - \gamma)\|x_n - z\| + \mu(\|z - x_n\| + \|x_n - z\|)\right] + \alpha_{n+1}^2\left[(1 - \gamma)\|x_{n+1} - z\| + \mu(\|z - x_{n+1}\| + \|x_{n+1} - z\|) + \right. \\ &\quad \left. (1 - \gamma)\|x_n - z\| + \mu(\|z - x_n\| + \|x_n - z\|)\right] \\ &\leq (1 - \alpha_{n+1} - \alpha_{n+1}^2)\left[(1 - \gamma)\|x_{n+1} - z\| + \gamma\|z - x_{n+1}\| + (1 - \gamma)\|x_n - z\| + \gamma\|z - x_n\|\right] + \\ &\quad \alpha_{n+1}\left[(1 - \gamma)\|x_{n+1} - z\| + \gamma\|z - x_{n+1}\| + (1 - \gamma)\|x_n - z\| + \gamma\|z - x_n\|\right] + \\ &\quad \alpha_{n+1}^2\left[(1 - \gamma)\|x_{n+1} - z\| + \gamma\|z - x_{n+1}\| + (1 - \gamma)\|x_n - z\| + \gamma\|z - x_n\|\right] \\ &= (1 - \alpha_{n+1} - \alpha_{n+1}^2)\left[\|x_{n+1} - z\| + \|x_n - z\|\right] + \alpha_{n+1}\left[\|x_{n+1} - z\| + \|x_n - z\|\right] + \alpha_{n+1}^2\left[\|x_{n+1} - z\| + \|x_n - z\|\right] \\ &= \|x_{n+1} - z\| + \|x_n - z\| \end{aligned}$$

Now from the following we have that,

$$\|(1 - \alpha_n - \alpha_n^2)T_1x_n + \alpha_nT_2x_n + \alpha_n^2T_3x_n - x_n\| \leq (1 - \alpha_n - \alpha_n^2)\|T_1x_n - x_n\| + \alpha_n\|T_2x_n - x_n\| + \alpha_n^2\|T_3x_n - x_n\|.$$

Taking $n \rightarrow \infty$ on both side of the above inequality and using Proposition 2 we get

$$\lim_{n \rightarrow \infty} \|(1 - \alpha_n - \alpha_n^2)T_1x_n + \alpha_nT_2x_n + \alpha_n^2T_3x_n - x_n\| = 0.$$

Theorem 2 Let C be a compact and convex subset of a Banach space X . Let T_1, T_2 and T_3 be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $[0, \frac{1}{2}]$ converging to 0. For $x_0 \in C$, let $\{x_n\}$ be a sequence in C as defined by (11). Then $\{x_n\}$ converges strongly to a common fixed point T_1, T_2 and T_3 .

Proof. Since C is compact, there exists a subsequence $\{x_{n_j}\}$ of $\{x_n\}$ and $z \in C$ such that $\{x_{n_j}\}$ converges to $z \in C$ By (Lemma 1)

Now by Proposition 1 (ii), for $\gamma = \frac{\theta}{2}$, $\theta \in [0, 1]$,

$$\begin{aligned} \gamma \|x_{n_j} - T_1 x_{n_j}\| &\leq \|x_{n_j} - z\| \\ \Rightarrow \gamma \|x_{n_j} - T_1 x_{n_j}\| &\leq \|x_{n_j} - z\| + \mu \|z - T_1 z\|. \end{aligned}$$

So by condition $B_{\gamma, \mu}$ and by using (7) we have

$$\begin{aligned} \|x_{n_j} - T_1 z\| &\leq \|x_{n_j} - T_1 x_{n_j}\| + \|T_1 x_{n_j} - T_1 z\| \\ &\leq \|x_{n_j} - T_1 x_{n_j}\| + (1 - \gamma) \|x_{n_j} - z\| + \mu (\|x_{n_j} - T_1 z\| + \|z - T_1 x_{n_j}\|) \\ &\leq \|x_{n_j} - T_1 x_{n_j}\| + (1 - \gamma) \|x_{n_j} - z\| + \mu (\|x_{n_j} - T_1 z\| + \|z - x_{n_j}\| + \|x_{n_j} - T_1 x_{n_j}\|). \end{aligned}$$

So taking $n_j \rightarrow \infty$ and using Proposition 2, we get

$$\begin{aligned} (1 - \mu) \|z - T_1 z\| &\leq 0 \\ \Rightarrow Tz &= z, \end{aligned}$$

since $\mu \neq 1$, shows that z is a fixed point T_1 . Therefore, by applying Lemma 3 and Theorem 1, z is a common fixed point of T_1 and T_2 .

Then by using (11) and (8), we have the following:

$$\begin{aligned} \|x_{n+1} - z\| &= \|\lambda(1 - \alpha_n - \alpha_n^2)T_1 x_n + \lambda\alpha_n T_2 x_n + \lambda\alpha_n^2 T_3 x_n + (1 - \lambda)x_n - z\| \\ &\leq \lambda(1 - \alpha_n - \alpha_n^2)\|T_1 x_n - z\| + \lambda\alpha_n \|T_2 x_n - z\| + \lambda\alpha_n^2 \|T_3 x_n - z\| + (1 - \lambda)\|x_n - z\| \\ &\leq \lambda(1 - \alpha_n - \alpha_n^2) \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - T_1 z\| + \|z - T_1 x_n\|) \right] + \\ &\quad \lambda\alpha_n \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - T_2 z\| + \|z - T_2 x_n\|) \right] + \\ &\quad \lambda\alpha_n^2 \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - T_3 z\| + \|z - T_3 x_n\|) \right] + (1 - \lambda) \|x_n - z\| \\ &\leq \lambda(1 - \alpha_n - \alpha_n^2) \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - z\| + \|z - x_n\|) \right] + \\ &\quad \lambda\alpha_n \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - z\| + \|z - x_n\|) \right] + \\ &\quad \lambda\alpha_n^2 \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - z\| + \|z - x_n\|) \right] + (1 - \lambda) \|x_n - z\| \\ &\leq \lambda(1 - \alpha_n - \alpha_n^2) \|x_n - z\| + \lambda\alpha_n \|x_n - z\| + \lambda\alpha_n^2 \|x_n - z\| + (1 - \lambda) \|x_n - z\| \quad (\text{since } 2\mu \leq \gamma) \\ &= \|x_n - z\|. \end{aligned}$$

Thus, $\{\|x_n - z\|\}$ is a monotonically decreasing non-negative sequence of real numbers, and will converges to some real numbers. Therefore, by similar arguments of Lemma 1 the sequence $\{x_n\}$ converges strongly to the common fixed point of T_1, T_2 and T_3 . ■

Lemma 5 Let C be a nonempty closed convex subset of a Banach space X . Let $l \in \mathbb{N}$ with $l \geq 2$ and let T_1, T_2, \dots, T_l be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $(0, \frac{1}{2})$ converging to 0, and let $\{x_n\}$ be a sequence in C such that $\{x_n\}$ converges strongly to some $z \in C$. Then z a common fixed point of $T_1, T_2 \dots T_l$.

Proof. We will prove this lemma by induction. We have already proved the conclusion in the case $l = 2, 3$. Fix $l \in \mathbb{N}$, $l \geq 4$. We assume the conclusion holds for every integer less than l and grater than 1.

Let T_1, T_2, \dots, T_l be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Then by Lemma 3 and the induction assumption, we have that z is a common fixed point of T_1, T_2, \dots, T_{l-1} . And we note that;

$$T_k \circ T_l z = T_l \circ T_k z = T_l z, \text{ with } 1 \leq k < l, \forall k \in \mathbb{N}. \quad (13)$$

We assume that z is not a fixed point of T_m . Put

$$\epsilon = \frac{\|T_l z - z\|}{3} > 0.$$

Then there exists $m \in \mathbb{N}$ such that

$$\|x_m - z\| < \epsilon, \quad \frac{\|(1 - \sum_{k=1}^{l-1} \alpha_m^k)T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\|}{\alpha_m^{l-1}} < \epsilon.$$

Since

$$\begin{aligned} 3\epsilon = \|T_m z - z\| &\leq \|T_l z - x_m\| + \|x_m - z\| \\ &< \|T_l z - x_m\| + \epsilon \\ \Rightarrow 2\epsilon &< \|T_l z - x_m\|. \end{aligned}$$

So, we obtain the following

$$\begin{aligned}
\|T_l z - x_m\| &\leq \|T_l z - (1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m - \sum_{k=2}^l \alpha_m^{k-1} T_k x_m\| \\
&\quad + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&\leq (1 - \sum_{k=1}^{l-1} \alpha_m^k) \|T_l z - T_1 x_m\| + \sum_{k=2}^{l-1} \alpha_m^{k-1} \|T_l z - T_k x_m\| + \alpha_m^{l-1} \|T_l z - T_l x_m\| \\
&\quad + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&= (1 - \sum_{k=1}^{l-1} \alpha_m^k) \|T_l \circ T_k z - T_1 x_m\| + \sum_{k=2}^{l-1} \alpha_m^{k-1} \|T_l \circ T_k z - T_k x_m\| + \alpha_m^{l-1} \|T_l z - T_l x_m\| \\
&\quad + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&\leq (1 - \sum_{k=1}^{l-1} \alpha_m^k) \left((1 - \gamma) \|T_l z - x_m\| + \mu (\|T_l z - x_m\| + \|x_m - T_l z\|) \right) + \\
&\quad \sum_{k=2}^{l-1} \alpha_m^{k-1} \left((1 - \gamma) \|T_l z - x_m\| + \mu (\|T_k z - x_m\| + \|x_m - T_l z\|) \right) + \alpha_m^{l-1} \|z - x_m\| \\
&\quad + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&\leq (1 - \sum_{k=1}^{l-1} \alpha_m^k) \left[\frac{(1 - \gamma + \mu)}{1 - \mu} \right] \|T_l z - x_m\| + \sum_{k=2}^{l-1} \alpha_m^{k-1} \left[\frac{(1 - \gamma + \mu)}{1 - \mu} \right] \|T_l z - x_m\| + \alpha_m^{l-1} \|z - x_m\| \\
&\quad + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&\leq (1 - \sum_{k=1}^{l-1} \alpha_m^k) \|T_l z - x_m\| + \sum_{k=2}^{l-1} \alpha_m^{k-1} \|T_l z - x_m\| + \alpha_m^{l-1} \|z - x_m\| \\
&\quad + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&= \|T_l z - x_m\| + \alpha_m^{l-1} \|z - x_m\| + \|(1 - \sum_{k=1}^{l-1} \alpha_m^k) T_1 x_m + \sum_{k=2}^l \alpha_m^{k-1} T_k x_m - x_m\| \\
&< \|T_l z - x_m\| + \alpha_m^{l-1} \epsilon + \alpha_m^{l-1} \epsilon \\
&= (1 - \alpha_m^{l-1}) \|T_l z - x_m\| + 2\alpha_m^{l-1} \epsilon \\
&< (1 - \alpha_m^{l-1}) \|T_l z - x_m\| + \alpha_m^{l-1} \|T_l z - x_m\| \\
&= \|T_l z - x_m\|,
\end{aligned}$$

which is a contradiction. Hence, z is a common fixed point of T_1, T_2, \dots, T_l , $l \in \mathbb{N}$. ■

Lemma 6 Let C be a nonempty closed convex and bounded subset of a Banach space X . Let T_1, T_2, \dots, T_l , $l \in \mathbb{N}$ be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $[0, \frac{1}{2}]$ converging to 0. Define a sequence $\{x_n\}$ in C by $x_0 \in C$ and

$$x_{n+1} = \lambda \left(1 - \sum_{k=1}^{l-1} \alpha_n^k \right) T_1 x_n + \lambda \sum_{k=2}^l \alpha_n^{k-1} T_k x_n + (1 - \lambda) x_n, \quad (14)$$

for $\lambda \in [\gamma, 1) - \{0\}$, $n = 0, 1, 2, \dots$. Then

$$\lim_{n \rightarrow \infty} \left\| \left(1 - \sum_{k=1}^{l-1} \alpha_n^k\right) T_1 x_n + \sum_{k=2}^l \alpha_n^{k-1} T_k x_n - x_n \right\| = 0. \quad (15)$$

Proof. Since and a sequence $\{x_n\}$ in C defined as (14) and assume that z is a common fixed point of T_1, T_2, \dots, T_l then, by condition $B_{\gamma, \mu}$ we get

$$\begin{aligned} \gamma \left\| \left(1 - \sum_{k=1}^{l-1} \alpha_n^k\right) T_1 x_n + \sum_{k=2}^l \alpha_n^{k-1} T_k x_n - x_n \right\| &\leq \lambda \left\| \left(1 - \sum_{k=1}^{l-1} \alpha_n^k\right) T_1 x_n + \sum_{k=2}^l \alpha_n^{k-1} T_k x_n - x_n \right\| \\ &= \|x_{n+1} - x_n\| \\ &\leq \|x_{n+1} - x_n\| + \mu \|x_n - T_1 x_n\|. \end{aligned}$$

It implies that

$$\begin{aligned} &\left\| \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) T_1 x_{n+1} + \sum_{k=2}^{l-1} \alpha_{n+1}^{k-1} T_k x_{n+1} - T_1 x_n \right\| \\ &\leq \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) \|T_1 x_{n+1} - T_1 x_n\| + \sum_{k=2}^l \alpha_{n+1}^{k-1} \|T_k x_{n+1} - T_1 x_n\| + \\ &\leq \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) \left[\|T_1 x_{n+1} - T_1 z\| + \|T_1 - T_1 x_n\| \right] + \sum_{k=2}^l \alpha_{n+1}^{k-1} \left[\|T_k x_{n+1} - T_k z\| + \|T_k z - T_1 x_n\| \right] \\ &\leq \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) \left[(1 - \gamma) \|x_{n+1} - z\| + \mu (\|z - T_1 x_{n+1}\| + \|x_{n+1} - T_1 z\|) + (1 - \gamma) \|x_n - z\| + \right. \\ &\quad \left. \mu (\|z - T_1 x_n\| + \|x_n - T_1 z\|) \right] + \sum_{k=2}^l \alpha_{n+1}^{k-1} \left[(1 - \gamma) \|x_{n+1} - z\| + \mu (\|z - T_k x_{n+1}\| + \right. \\ &\quad \left. \|x_{n+1} - T_k z\|) + (1 - \gamma) \|x_n - z\| + \mu (\|z - T_1 x_n\| + \|x_n - T_k z\|) \right] \\ &\leq \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) \left[(1 - \gamma) \|x_{n+1} - z\| + \mu (\|z - x_{n+1}\| + \|x_{n+1} - z\|) + (1 - \gamma) \|x_n - z\| + \right. \\ &\quad \left. \mu (\|z - x_n\| + \|x_n - z\|) \right] + \sum_{k=2}^l \alpha_{n+1}^{k-1} \left[(1 - \gamma) \|x_{n+1} - z\| + \mu (\|z - x_{n+1}\| + \right. \\ &\quad \left. \|x_{n+1} - z\|) + (1 - \gamma) \|x_n - z\| + \mu (\|z - x_n\| + \|x_n - z\|) \right] \\ &\leq \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) \left[(1 - \gamma) \|x_{n+1} - z\| + \gamma \|x_{n+1} - z\| + (1 - \gamma) \|x_n - z\| + \gamma \|x_n - z\| \right] + \\ &\quad \sum_{k=2}^l \alpha_{n+1}^{k-1} \left[(1 - \gamma) \|x_{n+1} - z\| + \gamma \|x_{n+1} - z\| + (1 - \gamma) \|x_n - z\| + \gamma \|x_n - z\| \right] \\ &= \left(1 - \sum_{k=1}^{l-1} \alpha_{n+1}^k\right) \left[\|x_{n+1} - z\| + \|x_n - z\| \right] + \sum_{k=2}^l \alpha_{n+1}^{k-1} \left[\|x_{n+1} - z\| + \|x_n - z\| \right] \\ &= \|x_{n+1} - z\| + \|x_n - z\|. \end{aligned}$$

Then, we have

$$\left\| \left(1 - \sum_{k=1}^{l-1} \alpha_n^k\right) T_1 x_n + \sum_{k=2}^l \alpha_n^{k-1} T_k x_n - x_n \right\| \leq \left(1 - \sum_{k=1}^{l-1} \alpha_n^k\right) \|T_1 x_n - x_n\| + \sum_{k=2}^l \alpha_n^{k-1} \|T_k x_n - x_n\|$$

Now by taking $n \rightarrow \infty$ and applying Proposition 2, we get

$$\lim_{n \rightarrow \infty} \left\| \left(1 - \sum_{k=1}^{l-1} \alpha_n^k\right) T_1 x_n + \sum_{k=2}^l \alpha_n^{k-1} T_k x_n - x_n \right\| = 0.$$

Theorem 3 Let C be a compact and convex subset of a Banach space X . Let T_1, T_2, \dots, T_l be commuting mappings satisfying the condition $B_{\gamma, \mu}$ on C . Let $\{\alpha_n\}$ be a sequence in $[0, \frac{1}{2}]$ converging to 0. For $x_0 \in C$, let $\{x_n\}$ be a sequence in C as defined by (14). Then $\{x_n\}$ converges strongly to a common fixed point of T_1, T_2, \dots, T_l .

Proof. By using Theorem 1 and induction assumption, assume that z is a common fixed point of T_1, T_2, \dots, T_l . Then by using (14) and (8), we have

$$\begin{aligned}
\|x_{n+1} - z\| &= \left\| \lambda \left(1 - \sum_{k=1}^{l-1} \alpha_n^k \right) T_1 x_n + \lambda \sum_{k=2}^l \alpha_n^{k-1} T_k x_n + (1 - \lambda) x_n - z \right\| \\
&\leq \lambda \left(1 - \sum_{k=1}^{l-1} \alpha_n^k \right) \|T_1 x_n - z\| + \lambda \sum_{k=2}^l \alpha_n^{k-1} \|T_k x_n - z\| + (1 - \lambda) \|x_n - z\| \\
&\leq \lambda \left(1 - \sum_{k=1}^{l-1} \alpha_n^k \right) \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - T_1 z\| + \|z - T_1 x_n\|) \right] + \\
&\quad \lambda \sum_{k=2}^l \alpha_n^{k-1} \left[(1 - \gamma) \|x_n - z\| + \mu (\|x_n - T_k z\| + \|z - T_k x_n\|) \right] + (1 - \gamma) \|x_n - z\| \\
&\leq \lambda \left(1 - \sum_{k=1}^{l-1} \alpha_n^k \right) \left[\frac{1 - \gamma + \mu}{1 - \mu} \right] \|x_n - z\| + \lambda \sum_{k=2}^l \alpha_n^{k-1} \left[\frac{1 - \gamma + \mu}{1 - \mu} \right] \|x_n - z\| + (1 - \gamma) \|x_n - z\| \\
&\leq \lambda \left(1 - \sum_{k=1}^{l-1} \alpha_n^k \right) \|x_n - z\| + \lambda \sum_{k=2}^l \alpha_n^{k-1} \|x_n - z\| + (1 - \gamma) \|x_n - z\| \\
&= \lambda \|x_n - z\| + (1 - \gamma) \|x_n - z\| \\
&= \|x_n - z\|.
\end{aligned}$$

Thus, $\{\|x_n - z\|\}$ is a monotonically decrease non-negative sequence of real numbers, and will converge to some real number. Therefore, by Theorem 2 and induction assumption, the sequence $\{x_n\}$ converges strongly to the common fixed point of T_1, T_2, \dots, T_l . ■

4 Conclusion

In this paper, our purpose was to extend the classic approach of fixed point searching from [19], by taking the condition $B_{\gamma, \mu}$ and the process by properly using the finite number of operators in to its structure. In other words, we brought together both the Suzuki's strong nonexpansive mappings and the Patir et al. [17] condition $B_{\gamma, \mu}$ under the same iteration process. Under the resulted iteration process, we proved the existence of a common fixed point for finite number of commutative mapping satisfies the condition $B_{\gamma, \mu}$. We proves several strong convergence results and existence of the common fixed point for the sequence of approximation by those iteration.

References

- [1] H. Afshari, H. Aydi, and E. Karapinar. Existence of fixed points of set-valued mappings in b-metric spaces. *East Asian mathematical journal*, 32(3):319–332, 2016.
- [2] Ü. Aksoy, E. Karapinar, and İ. M Erhan. Fixed point theorems in complete modular metric spaces and an application to anti-periodic boundary value problems.
- [3] Ü. Aksoy, E. Karapinar, and İ. M Erhan. Fixed points of generalized alpha-admissible contractions on b-metric spaces with an application to boundary value problems.
- [4] B. Alqahtani, A. Fulga, and E. Karapinar. Non-unique fixed point results in extended b-metric space. *Mathematics*, 6(5):68, 2018.

- [5] A Betiuk-Pilarska and T D. Benavides. The fixed point property for some generalized nonexpansive mappings and renormings. *Journal of Mathematical Analysis and Applications*, 429(2):800–813, 2015.
- [6] Deepmala and R. P. Agarwal. Existence and uniqueness of solutions for certain functional equations and system of functional equations arising in dynamic programming. *Analele Universitatii” Ovidius” Constanta-Seria Matematica*, 24(1):3–28, 2016.
- [7] S Dhompongsa, W I., and A Kaewkhao. Edelstein’s method and fixed point theorems for some generalized nonexpansive mappings. *Journal of Mathematical Analysis and Applications*, 350(1):12–17, 2009.
- [8] J. García-Falset, E. Llorens-Fuster, and T. Suzuki. Fixed point theory for a class of generalized nonexpansive mappings. *Journal of mathematical analysis and applications*, 375(1):185–195, 2011.
- [9] M Gunaseelan and LN Mishra. Coupled fixed point theorems on complex partial metric space using different type of contractive conditions. *Scientific Publications of the State University of Novi Pazar Series A: Applied Mathematics, Informatics and Mechanics*, 11(2):117–123, 2019.
- [10] S Kanas and H. M. Srivastava. Linear operators associated with k-uniformly convex functions. *Integral Transforms and Special Functions*, 9(2):121–132, 2000.
- [11] M. A. Khamsi and A. R. Khan. On monotone nonexpansive mappings in l_1 $([0, 1])$ $l_{-1}([0, 1])$. *Fixed Point Theory and Applications*, 2015(1):1–5, 2015.
- [12] F. Lael and Z. Heidarpour. Fixed point theorems for a class of generalized nonexpansive mappings. *Fixed Point Theory and Applications*, 2016(1):1–7, 2016.
- [13] L. N. Mishra, S. K. Tiwari, and V. N. Mishra. Fixed point theorems for generalized weakly s-contractive mappings in partial metric spaces. *J. Appl. Anal. Comput.*, 5(4):600–612, 2015.
- [14] L. N. Mishra, S. K. Tiwari, V. N. Mishra, and I. A Khan. Unique fixed point theorems for generalized contractive mappings in partial metric spaces. *Journal of Function Spaces*, 2015, 2015.
- [15] V. N. Mishra. *Some problems on approximations of functions in Banach spaces*. PhD thesis, Ph. D. thesis, 2007.
- [16] B. Patir, N. Goswami, and L. N. Mishra. Fixed point theorems in fuzzy metric spaces for mappings with some contractive type conditions. *Korean journal of mathematics*, 26(2):307–326, 2018.
- [17] B. Patir, N. Goswami, and V. N. Mishra. Some results on fixed point theory for a class of generalized nonexpansive mappings. *Fixed point theory and applications*, 2018(1):1–18, 2018.
- [18] M. G. Sangago. Weak convergence of iterations for nonexpansive mappings. *International Journal of Mathematics & Computation*, 6(M10):69–76, 2010.
- [19] T. Suzuki. Strong convergence theorems for infinite families of nonexpansive mappings in general banach spaces. *Fixed Point Theory and Applications*, 2005(1):1–21, 2005.
- [20] T. Suzuki. Fixed point theorems and convergence theorems for some generalized nonexpansive mappings. *Journal of mathematical analysis and applications*, 340(2):1088–1095, 2008.
- [21] D Thakur, B. S. Thakur, and M. Postolache. Convergence theorems for generalized nonexpansive mappings in uniformly convex banach spaces. *Fixed Point Theory and Applications*, 2015(1):1–12, 2015.