

(Un)boundedness of directional maximal operators through a notion of “Perron capacity” and an application

Emma D’Aniello, Anthony Gauvan and Laurent Moonens

June 15, 2022

Abstract

We introduce the notion of *Perron capacity* of a set of slopes $\Omega \subset \mathbb{R}$. Precisely, we prove that if the Perron capacity of Ω is finite then the directional maximal operator associated M_Ω is not bounded on $L^p(\mathbb{R}^2)$ for any $1 < p < \infty$. This allows us to prove that the set

$$\Omega_e = \left\{ \frac{\cos n}{n} : n \in \mathbb{N}^* \right\}$$

is not finitely lacunary which answers a question raised by A. Stokolos.

1 Introduction

It is well known, since the original work of Jensen, Marcinkiewicz and Zygmund [16], that Lebesgue’s differentiation formula, *i.e.* the equality for a.e. $x \in \mathbb{R}^n$:

$$f(x) = \lim_{\substack{R \ni x \\ \text{diam } R \rightarrow 0}} \frac{1}{|R|} \int_R f, \quad (1)$$

holds along n -dimensional *rectangles* R parallel to the coordinate hyperplanes (called standard rectangles), whenever one has $f \in L \log^{n-1} L(\mathbb{R}^n)$ — the latter Orlicz space being, moreover, the largest for which the above formula holds if one does not assume any extra hypothesis on the rectangles R ; the beautiful survey by A. Stokolos [21] provides a quite exhaustive state of the art on the topic — see also [5] for sufficient conditions (of geometric nature) on \mathcal{R} ensuring that $L \log^{n-1} L(\mathbb{R}^n)$ is the largest space for which (1) holds whenever R is restricted to belong to the class \mathcal{R} .

In dimension $n = 2$, the nature of the problem changes when one allows rectangles R to rotate around one of their vertices, requiring that the slope of their longest side

⁰2020 *Mathematics Subject Classification*: Primary: 42B25, 26B05; Secondary: 42B35.

Keywords: Lebesgue’s Differentiation Theorem, Directional Maximal Functions, Lacunary Sets of Finite Order.

belongs to a fixed set Ω (call \mathcal{R}_Ω the family of such rectangles in the plane). It has been shown *e.g.* by Cordoba and Fefferman [4] that when $\Omega = \{\omega_k : k \in \mathbb{N}\}$ is a lacunary sequence (see below for a definition), then formula (1) holds for any $f \in L^p(\mathbb{R}^2)$ if one has $p \in [2, +\infty[$. This was later generalized for arbitrary $1 < p < \infty$ by Nagel, Stein and Wainger [19]. Given a set of slopes Ω , the question of finding Orlicz spaces (the largest possible) beyond L^p spaces, for all elements of which formula (1) holds is often delicate; issues in these directions have been studied in works involving some of the current authors, see [7] and [6] for example.

Concerning the possibility of (1) holding for all $f \in L^p(\mathbb{R}^2)$, $1 < p \leq \infty$, it has been shown *e.g.* by Katz [17], Bateman and Katz [2], Hare [14] and more recently by Bateman [1] that Cantor sets (and even uncountable sets) never give rise to formula (1) for all $f \in L^p(\mathbb{R}^2)$, $1 < p \leq \infty$ — Bateman’s work [1] even gives a complete characterization, even though not always simple to implement *in practice* (see below), of sets Ω for which it does hold in any $L^p(\mathbb{R}^2)$, $1 < p \leq \infty$: those are sets one will call *finitely lacunary* in the sequel.

Before to state more precisely Bateman’s result, let us mention an important tool in its proof: the possibility, when a set of slopes Ω is too large, to find finite families of rectangles in \mathcal{R}_Ω such that the union of their (centered) homothetic expansions occupies a much larger area in the plane than their union does; we call this a *Keakeya blow*.

When no restriction is made on directions of rectangles in play, the possibility of exhibiting a Keakeya blow is a standard result in measure theory (see Busemann and Feller’s work [3] for a first construction of this type):

Theorem 1.1 (Keakeya Blow). *For any large $A \gg 1$, there exists a finite family of rectangles $\{R_i\}_{i \leq N}$ such that we have*

$$A \left| \bigcup_{i \leq N} R_i \right| \leq \left| \bigcup_{i \leq N} 4R_i \right|.$$

Here, $4R$ stands for the 4-fold dilation of R by its center.

Keakeya blows have deep implications in harmonic analysis: for example, a concrete construction proving the above theorem allowed Fefferman to disprove the Disc multiplier conjecture in [10].

A natural question, given a set of slopes Ω , then becomes the following: can we still construct Keakeya blows using *only* rectangles in \mathcal{R}_Ω , namely rectangles the longest side of which makes an angle $\omega \in \Omega$ with the horizontal axis (see also Stokolos [20] and Hagelstein and Stokolos [13] concerning the links between directional maximal operators and multipliers)? In his study of the so-called *directional maximal operator* M_Ω defined for a locally integrable $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $x \in \mathbb{R}^2$ as

$$M_\Omega f(x) := \sup_{R \in \mathcal{R}_\Omega} \frac{1}{|R|} \int_R |f|,$$

Bateman [1] introduced a notion we shall call *finitely lacunary set* and proved the following Theorem.

Theorem 1.2 (Bateman). *The following conditions are equivalent*

- *the set Ω is not finitely lacunary ;*
- *for any $A \gg 1$, there exists a finite family of rectangles $\{R_i\}$ such that the longest side of each makes an angle $\omega \in \Omega$ with the x -axis and also satisfying $A \left| \bigcup_{i \in I} R_i \right| \leq \left| \bigcup_{i \in I} 4R_i \right|$;*
- *the operator M_Ω is not bounded on $L^p(\mathbb{R}^2)$ for any $1 < p < \infty$.*

Moreover, the following conditions are also equivalent

- *the set Ω is finitely lacunary*
- *the operator M_Ω is bounded on $L^p(\mathbb{R}^2)$ for any $1 < p < \infty$.*

In the latter theorem, it needs to be explained what is meant by “finitely lacunary”. We recall it here, following the presentation by Kroc and Pramanik [18]. We start by defining the notion of *lacunary sequence* and then the notion of *lacunary set of finite order*.

Definition 1.3 (Lacunary sequence). We say that a sequence of real numbers $L = (L_k)$ is a lacunary sequence converging to $\ell \in \mathbb{R}$ when there holds

$$|\ell - \ell_{k+1}| \leq \frac{1}{2} |\ell - \ell_k|$$

for any k .

For example the sequences $L_1 := \left(\frac{1}{2^k}\right)_{k \geq 2}$ and $L_2 := \left(\frac{1}{k!}\right)_{k \geq 4}$ are lacunary.

Definition 1.4 (Lacunary set of finite order). A lacunary set of order 0 in \mathbb{R} is a set which is either empty or a singleton. Recursively, for $N \in \mathbb{N}^*$, we say that a set Ω included in \mathbb{R} is a lacunary set of order at most $N + 1$ — and write $\Omega \in \Lambda(N + 1)$ — if there exists a lacunary sequence L with the following properties : for any $a, b \in L$ such that $a < b$ and $(a, b) \cap L = \emptyset$, the set $\Omega \cap (a, b)$ is a lacunary set of order at most N *i.e.* $\Omega \cap (a, b) \in \Lambda(N)$.

For example the set

$$\Omega := \left\{ \frac{1}{2^k} + \frac{1}{4^l} : k, l \in \mathbb{N}, l \leq k \right\}$$

is a lacunary set of order 2. In this case, observe that the set Ω cannot be re-written as $\Omega = \{\omega_k : k \in \mathbb{N}^*\}$ where (ω_k) is a monotone sequence, since it has several points of accumulation. We can finally give a definition of a finitely lacunary set.

Definition 1.5 (Finitely lacunary set). A set Ω in \mathbb{R} is said to be *finitely lacunary* if there exists a finite number of set $\Omega_1, \dots, \Omega_M$ which are lacunary of finite order such that

$$\Omega \subset \bigcup_{k \leq M} \Omega_k.$$




Figure 1: A representation of a lacunary sequence and a lacunary set of order 2.

Even if Theorem 1.2 gives a full characterization of the behavior of directional maximal operators, it is usually hard *in practice* to decide whether a set of slopes Ω is finitely lacunary. For example, it was known that the set of slopes Ω_0 defined by

$$\Omega_0 := \left\{ \frac{1}{n} : n \in \mathbb{N}^* \right\}$$

generates a directional maximal operator M_{Ω_0} which is not bounded on $L^p(\mathbb{R}^2)$ for any $1 < p < \infty$ (see *e.g.* [9, Theorem 2.1]); using Theorem 1.2 one can hence assert that the set of directions Ω_0 is not finitely lacunary. In the case where the set of directions Ω is a discrete set which can be ordered in \mathbb{R}_+^* , more precisely in the case where $\Omega = \{\frac{1}{u} : u \in U\}$ and we can write $U = \{u_k : k \in \mathbb{N}^*\} \subset \mathbb{R}_+^*$ for an increasing sequence $(u_k)_{k \in \mathbb{N}^*}$, Hare and Rönning [15] provided a *quantitative* sufficient condition on U which ensures that *the maximal operator associated M_U is not bounded on $L^p(\mathbb{R}^2)$* . Precisely, they proved the following Theorem.

Theorem 1.6 (Hare-Rönning [15]). *Assume that $U = (u_k)_{k \in \mathbb{N}^*} \subset (0, \pi/2)$ is an increasing sequence of directions and that one has*

$$G_U := \sup_{\substack{k \geq 1 \\ l \leq k}} \left(\frac{u_{k+2l} - u_{k+l}}{u_{k+l} - u_k} + \frac{u_{k+l} - u_k}{u_{k+2l} - u_{k+l}} \right) < \infty.$$

Then for any $p \in (1, \infty)$, one has $\|M_U\|_p = \infty$.

In particular, applying Bateman's Theorem, it follows that if we have $G_U < \infty$ then U is not finitely lacunary — other uses of Perron trees have also been made in works involving the current authors, see *e.g.* [8], [11] and [12]. Hence, usually one does not prove *directly* that a set of slopes Ω is not finitely lacunary (or that it is): in general, it is a consequence of the (un)-boundedness of the associated maximal operator M_Ω — obtained by other methods — and Bateman's Theorem.

Our results are motivated by the following question raised by A. Stokolos : what can be said (for example) about the set of directions Ω_e defined as

$$\Omega_e := \left\{ \frac{\cos n}{n} : n \in \mathbb{N}^* \right\} ?$$

Can one prove that it is finitely lacunary or not? In this text, we prove it is *not* finitely lacunary.

2 Results

We denote by \mathcal{R} the set containing all rectangles of the plane and for $R \in \mathcal{R}$, we denote by $\omega_R \in [0, \pi)$ the angle that the longest side of the rectangle R makes with the x -axis. For any set $\Omega \subset \mathbb{R}$ (which should be thought as a set of *slopes*) we define the family of rectangle \mathcal{R}_Ω as

$$\mathcal{R}_\Omega := \{R \in \mathcal{R} : \tan(\omega_R) \in \Omega\}.$$

We then define the *directional maximal operator* M_Ω as

$$M_\Omega f(x) := \sup_{x \in R \in \mathcal{R}_\Omega} \frac{1}{|R|} \int_R |f|.$$

We finally define the *Perron capacity* of a set of slopes Ω included in \mathbb{R} as follow.

Definition 2.1 (Perron capacity of a set of slopes). For any set of slopes $\Omega \subset \mathbb{R}$, we define its *Perron capacity* as

$$P_\Omega := \lim_{N \rightarrow \infty} \inf_{\substack{U \subset \Omega \\ \#U = 2^N}} G_U \in [2, \infty]$$

where $G_U = \sup_{\substack{k, l \geq 1 \\ k+2l \leq 2^N}} \left(\frac{u_{k+2l} - u_{k+l}}{u_{k+l} - u_k} + \frac{u_{k+l} - u_k}{u_{k+2l} - u_{k+l}} \right)$ if $U = \{u_1, \dots, u_{2^N}\}$ with $u_1 < u_2 < \dots < u_{2^N}$.

Our first result is the following: it gives a *sufficient* and *quantitative* condition on an arbitrary set Ω which ensures that the associated maximal operator is unbounded on $L^p(\mathbb{R}^2)$ for any $1 < p < \infty$. In contrast with Hare and Ronning Theorem, we do not assume that our set of slopes is ordered.

Theorem 2.2. *Fix any set of slopes $\Omega \subset \mathbb{R}$ and suppose that we have*

$$P_\Omega < \infty.$$

Then for any $p \in (1, \infty)$, one has $\|M_\Omega\|_p = \infty$.

Loosely speaking, the fact that $P_\Omega < \infty$ indicates whether the set Ω contains arbitrary large subsets which are (more or less) uniformly distributed in \mathbb{R} . Our second result is an application of Theorem 2.2 and deals with the set of slopes Ω_e defined as

$$\Omega_e := \left\{ \frac{\cos n}{n} : n \in \mathbb{N}^* \right\}.$$

Namely, we prove the following Theorem.

Theorem 2.3. *The Perron capacity of the set Ω_e is finite i.e. we have $P(\Omega_e) < \infty$.*

Hence, as a consequence of Theorems 2.3 and 2.2 we know that — for any $1 < p < \infty$ — we have

$$\|M_{\Omega_e}\|_p = \infty.$$

As an application of Bateman's Theorem, we can hence say that the set Ω_e is not finitely lacunary.

3 Proof of Theorem 2.2

Recall that for any set $U := \{u_1, \dots, u_{2^N}\} \subset \mathbb{R}_+$ with $u_1 < u_2 < \dots < u_{2^N}$ (so that U has 2^N elements), one define its *Perron factor* as

$$G_U := \sup_{\substack{k, l \geq 1 \\ k+2l \leq 2^N}} \left(\frac{u_{k+2l} - u_{k+l}}{u_{k+l} - u_k} + \frac{u_{k+l} - u_k}{u_{k+2l} - u_{k+l}} \right) \in (0, \infty).$$

The following proposition is a careful reading of Hare and Rönning's work [15].

Proposition 3.1. *There exists $\varepsilon_0 \in (0, 1)$ such that for any $\alpha \in [1 - \varepsilon_0, 1)$ there exists a set $X \subseteq \mathbb{R}^2$ for which one has:*

$$|X| \leq [\alpha^{2^N} + G_U(1 - \alpha)^2] \left| \left\{ M_U \mathbf{1}_X > \frac{1}{4} \right\} \right|.$$

Hence for any $0 < \alpha < 1$ close enough to 1, this gives us the following lower bound for any $p \in (1, \infty)$:

$$\|M_U\|_p^p \gtrsim_p \frac{1}{\alpha^{2^N} + G_U(1 - \alpha)^2},$$

where \gtrsim_p means that the inequality holds up to a multiplicative constant that does only depend on p .

Indeed, we have

$$\|M_U \mathbf{1}_X\|_p^p \geq \frac{1}{4^p} \left| \left\{ M_U \mathbf{1}_X > \frac{1}{4} \right\} \right| \gtrsim_p \frac{|X|}{\alpha^{2^N} + G_U(1 - \alpha)^2}.$$

This allows us now to easily conclude the proof of Theorem 2.2. Indeed, fix an arbitrary set $\Omega \subset \mathbb{R}$ and suppose that its Perron capacity is finite *i.e.* assume that one has:

$$P_\Omega < \infty.$$

By definition of P_Ω , there thus exists a strictly increasing sequence of integers $\{N_k : k \in \mathbb{N}^*\}$ such that for any k there is a set $U_k \subset \Omega$ such that $G(U_k) < 2P_\Omega$ and $\#U_k = 2^{N_k}$. Since there holds $U_k \subset \Omega$, we obtain, for any $0 < \alpha < 1$ sufficiently close to 1 and any $k \geq 1$:

$$\|M_\Omega\|_p^p \geq \|M_{U_k}\|_p^p \gtrsim \frac{1}{\alpha^{2^{N_k}} + 2P_\Omega(1 - \alpha)^2}.$$

Since this holds for any $k \geq 1$ and any α close to 1, this implies that we have $\|M_\Omega\|_p = \infty$.

4 Homogeneous sets

We fix an arbitrary set U in \mathbb{R}_+^* whose cardinal is 2^N for some $N \in \mathbb{N}^*$. The following proposition shows that it is equivalent for U to be uniformly distributed and its Perron factor to equal 2.

Proposition 4.1. *We have $G_U = 2$ if and only if the elements of U are in arithmetic progression i.e. U is of the form*

$$U = \{a + k\delta : 1 \leq k \leq 2^N\}$$

for some $a \in \mathbb{R}$ and $\delta > 0$.

Proof. If U is in arithmetic progression we have easily $G_U = 2$. On the other hand if we have $G_U = 2$ since $x + \frac{1}{x} = 2$ if and only if $x = 1$ we have for any k (taking $\ell = 1$)

$$u_{k+2} - u_{k+1} = u_{k+1} - u_k$$

which concludes the proof. □

We will be particularly interested by *homogeneous sets* that is to say sets H of the form

$$H := H_{a,N} = \{ka : 1 \leq k \leq 2^N\}$$

for some integers $a \in \mathbb{N}^*$ and $N \in \mathbb{N}$. In particular, we wish to perturb a little an homogeneous set $H_{a,N}$ into a set H' such that the Perron's constant of H' is still bounded. More precisely, fix any $a, N \in \mathbb{N}^*$ and let ε be an arbitrary function

$$\varepsilon : H_{a,N} \rightarrow (0, 1).$$

Define then the set $H_{a,N}(\varepsilon)$ as

$$H_{a,N}(\varepsilon) := \{(1 + \varepsilon(x))x : x \in H_{a,N}\}.$$




Figure 2: A representation of a uniformly distributed set and its perturbation.

Proposition 4.2. *Assume that $a \in \mathbb{R}_+^*$, $N \in \mathbb{N}^*$ and $\varepsilon : H_{a,N} \rightarrow (0, 1)$ are given. If one has*

$$2^N \|\varepsilon\|_\infty \leq \frac{1}{2},$$

then we have $G_{H_{a,N}} \leq 6$.

Proof. To simplify the notations, let for $1 \leq k \leq 2^N$, $h_k := [1 + \varepsilon(ka)]ka$. First, observe that for $1 \leq k \leq 2^N - 1$, one has:

$$\begin{aligned} h_{k+1} - h_k &= \{1 + \varepsilon[(k+1)a]\}(k+1)a - [1 + \varepsilon(ka)]ka \\ &\geq a - ak\varepsilon(ka) = a[1 - k\varepsilon(ka)] \geq a[1 - (2^N - 1)\|\varepsilon\|_\infty] > 0, \end{aligned}$$

so that there holds $h_1 < h_2 < \dots < h_{2^N}$.

Now compute, for $1 \leq l \leq k$ such that $k + 2l \leq 2^N$ (implying in particular that one has $\frac{k}{l} \leq 2^N - 2$):

$$\frac{h_{k+2l} - h_{k+l}}{h_{k+l} - h_k} \leq \frac{la + 2^N a \|\varepsilon\|_\infty}{la - \|\varepsilon\|_\infty ka} = \frac{1 + \frac{2^N}{l} \|\varepsilon\|_\infty}{1 - \frac{k}{l} \|\varepsilon\|_\infty} \leq \frac{1 + 2^N \|\varepsilon\|_\infty}{1 - (2^N - 2) \|\varepsilon\|_\infty} \leq 3.$$

One obtains a similar inequality for the symmetric ratio. □

5 Proof of Theorem 2.3

Recall that we have defined

$$\Omega_e := \left\{ \frac{n}{\cos n} : n \in \mathbb{N}^* \right\}$$

and that we wish to prove that the Perron capacity of Ω_e is finite *i.e.* that we have $P(\Omega_e) < \infty$. To do so, we are simply going to prove that the set Ω_e contains “small perturbations” of *arbitrarily long homogeneous sets*. Specifically, for any $N \in \mathbb{N}$, we consider the set

$$E(N) := \{n \in \mathbb{N}^* : \exists m \in \mathbb{Z}, |n + 2\pi m| < 2^{-N}\}.$$

To begin with, we claim the following.

Claim 1. For any $N \in \mathbb{N}$ we have $\#E(N) = \infty$.

Proof of the claim. The claim follows from the fact that $G' = \{n + 2\pi m : n \in \mathbb{N}, m \in \mathbb{Z}\}$ is dense in \mathbb{R} , which can be shown easily using standard techniques from basic real analysis¹. □

Claim 2. For any $n \in \mathbb{N}$ and any $n \in E(N)$, one has:

$$1 < \frac{1}{\cos n} \leq 1 + 2^{-2N}.$$

¹See *e.g.* <https://lefevre.perso.math.cnrs.fr/PagesPerso/enseignement/Archives/SSgrpesAdd.pdf> P. Lefèvre’s webpage for a proof.

Proof of the claim. Choosing $m \in \mathbb{Z}$ such that one has $|n + 2\pi m| < 2^{-N}$ (which exists by definition since one has $n \in E(N)$), one obtains, using the inequality $1 - \cos x \leq \frac{1}{2}x^2$ valid for any $x \in \mathbb{R}$:

$$1 - \cos n = 1 - \cos(n + 2\pi m) \leq \frac{1}{2}(n + 2\pi m)^2 \leq \frac{1}{2}2^{-2N}.$$

Note that, in particular, this yields $0 < \cos n < 1$.

Using the fact that one has $\frac{1}{1-x} \leq 1 + 2x$ for any $0 \leq x \leq \frac{1}{2}$, one finally computes:

$$\frac{1}{\cos n} = \frac{1}{1 - (1 - \cos n)} \leq 1 + 2(1 - \cos n) \leq 1 + 2^{-2N},$$

as was announced. \square

Fix an arbitrary large $N \in \mathbb{N}$ and any integer $a \in E(2N)$. We claim the following.

Claim 3. For any $a \in E(2N)$, we have

$$H_{a,N} \subset E(N).$$

Proof. Fix $a \in E(2N)$; by definition we have $m \in \mathbb{Z}$ such that

$$|a + 2\pi m| < 2^{-2N}.$$

Hence for any $1 \leq k \leq 2^N$ we have

$$|ka + 2\pi km| < k2^{-2N} \leq 2^{-N}$$

i.e. for any $1 \leq k \leq 2^N$ we have $ka \in E(N)$. \square

Fix now $N \in \mathbb{N}^*$ and $a \in E(2N)$ and define $\varepsilon : H_{a,N} \rightarrow (0, 1)$ by the formula $1 + \varepsilon(n) = \frac{1}{\cos n}$ for $n \in H_{a,N}$ — we hence see the set:

$$\left\{ \frac{n}{\cos n} : n \in H_{a,N} \right\} = \{[1 + \varepsilon(n)]n : n \in H_{a,N}\},$$

as a perturbation of the above type of the set $H_{a,N}$ — . We compute, using Claim 2:

$$2^N \|\varepsilon\|_\infty \leq 2^{-N} \leq \frac{1}{2},$$

so that Proposition 4.2 yields $G_{H_{a,N}(\varepsilon)} \leq 6$.

Finally, observe that we have by construction the inclusion

$$H_{a,N}(\varepsilon) \subset \Omega.$$

Since this holds for any $N \in \mathbb{N}^*$, it follows that we have $P_\Omega \leq 6$ as desired. \square

Remark 5.1. The same conclusion holds if in Theorem 2.3 we replace $\Omega_e := \left\{ \frac{\cos n}{n} : n \in \mathbb{N}^* \right\}$ with

$$\Omega_s := \left\{ \frac{\sin n}{n} : n \in \mathbb{N}^* \right\}.$$

As an application of Bateman's Theorem, we can hence say that also the set Ω_s is not finitely lacunary. The argument is the same but, in this case, we approximate $\frac{\pi}{2}$ by elements of G' and take $1 + \varepsilon(x) = \frac{1}{\sin x}$.

Acknowledgements.

This research has been supported by the *École Normale Supérieure*, Paris, by the “Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni dell’Istituto Nazionale di Alta Matematica F. Severi” and by the project *Vain-Hopes* within the program *Valere* of *Università degli Studi della Campania “Luigi Vanvitelli”*. It has also been partially accomplished within the *UMI Group TAA “Approximation Theory and Applications”*.

E. D’Aniello would like to thank the *École Normale Supérieure*, Paris, for its warm hospitality in March-April, 2022; L. Moonens would like to thank the *Dipartimento di Matematica e Fisica* of the *Università degli Studi della Campania “Luigi Vanvitelli”* for its warm hospitality in April-May, 2022.

References

- [1] Michael Bateman. *Keakeya sets and directional maximal operators in the plane.* *Duke Math. J.*, 147(1):55–77, 2009.
- [2] Michael Bateman and Nets Hawk Katz. *Keakeya sets in Cantor directions.* *Math. Res. Lett.*, 15(1):73–81, 2008.
- [3] Herbert Busemann and William Feller. *Zur differentiation der lebesgueschen integrale.* *Fundamenta Mathematicae*, 22(1):226–256, 1934.
- [4] Antonio Córdoba and Robert Fefferman. *On differentiation of integrals.* *Proc. Nat. Acad. Sci. U.S.A.*, 74(6):2211–2213, 1977.
- [5] Emma D’Aniello and Laurent Moonens. *Averaging on n -dimensional rectangles.* *Ann. Acad. Sci. Fenn. Math.*, 42(1):119–133, 2017.
- [6] Emma D’Aniello and Laurent Moonens. *Differentiating Orlicz spaces with rectangles having fixed shapes in a set of directions.* *Z. Anal. Anwend.*, 39(4):461–473, 2020.
- [7] Emma D’Aniello, Laurent Moonens, and Joseph M. Rosenblatt. *Differentiating Orlicz spaces with rare bases of rectangles.* *Ann. Acad. Sci. Fenn. Math.*, 45(1):411–427, 2020.
- [8] D’Aniello, Emma and Gauvan, Anthony and Moonens, Laurent and Rosenblatt, Joseph M. *Almost Everywhere Convergence for Lebesgue Differentiation Processes along Rectangles.* *Preprint*, 2022.
- [9] Miguel de Guzmán. *Differentiation of integrals in R^n .* *Lecture Notes in Mathematics*, Vol. 481. Springer-Verlag, Berlin-New York, 1975. With appendices by Antonio Córdoba, and Robert Fefferman, and two by Roberto Moriyón.

- [10] Charles Fefferman. The multiplier problem for the ball. *Ann. of Math. (2)*, 94:330–336, 1971.
- [11] Anthony Gauvan. Application of perron trees to geometric maximal operators, <https://arxiv.org/abs/2204.02640>, 2022.
- [12] Anthony Gauvan. Kakeya-type sets for geometric maximal operators, <https://arxiv.org/abs/2204.00253>, 2022.
- [13] Paul Hagelstein and Alexander M. Stokolos. An extension of the Córdoba-Fefferman theorem on the equivalence between the boundedness of certain classes of maximal and multiplier operators. *C. R. Math. Acad. Sci. Paris*, 346(19-20):1063–1065, 2008.
- [14] Kathryn E. Hare. Maximal operators and Cantor sets. *Canad. Math. Bull.*, 43(3):330–342, 2000.
- [15] Kathryn E. Hare and Jan-Olav Rønning. Applications of generalized Perron trees to maximal functions and density bases. *J. Fourier Anal. Appl.*, 4(2):215–227, 1998.
- [16] Børge Jessen, Józef Marcinkiewicz, and Antoni Zygmund. Note on the differentiability of multiple integrals. *Fundam. Math.*, 25:217–234, 1935.
- [17] Nets Hawk Katz. A counterexample for maximal operators over a Cantor set of directions. *Math. Res. Lett.*, 3(4):527–536, 1996.
- [18] Edward Kroc and Malabika Pramanik. Kakeya-type sets over Cantor sets of directions in \mathbb{R}^{d+1} . *J. Fourier Anal. Appl.*, 22(3):623–674, 2016.
- [19] Alexander Nagel, Elias M. Stein, and Stephen Wainger. Differentiation in lacunary directions. *Proc. Nat. Acad. Sci. U.S.A.*, 75(3):1060–1062, 1978.
- [20] Alexander M. Stokolos. On a problem of M. de Guzmán on Fourier multipliers of polygonal domains. *Sibirsk. Mat. Zh.*, 36(6):1392–1398, iii, 1995.
- [21] Alexander M. Stokolos. Zygmund’s program: some partial solutions. *Ann. Inst. Fourier (Grenoble)*, 55(5):1439–1453, 2005.

EMMA D’ANIELLO

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI DELLA CAMPANIA “LUIGI VANVITELLI”, VIALE LINCOLN N. 5, 81100 CASERTA, ITALIA
emma.daniello@unicampania.it.

ANTHONY GAUVAN

LABORATOIRE DE MATHÉMATIQUES D’ORSAY, UNIVERSITÉ PARIS-SUD, CNRS UMR8628, UNIVERSITÉ PARIS-SACLAY, BÂTIMENT 307, F-91405 ORSAY CEDEX, FRANCE
anthony.gauvan@universite-paris-saclay.fr.

LAURENT MOONENS

LABORATOIRE DE MATHÉMATIQUES D’ORSAY, UNIVERSITÉ PARIS-SACLAY, CNRS UMR

8628, BÂTIMENT 307, F-91405 ORSAY CEDEX, FRANCE

laurent.moonens@universite-paris-saclay.fr.

ÉCOLE NORMALE SUPÉRIEURE-PSL UNIVERSITY, CNRS UMR 8553, 45, RUE D'ULM, F-75230 PARIS CEDEX 3, FRANCE

laurent.moonens@ens.fr.