

SANDWICH CELLULARITY AND A VERSION OF CELL THEORY

DANIEL TUBBENHAUER

ABSTRACT. We explain how the theory of sandwich cellular algebras can be seen as a version of cell theory for algebras. We apply this theory to many examples such as Hecke algebras, and various monoid and diagram algebras.

CONTENTS

1. Introduction	1
2. Cell theory for algebras	2
3. Kazhdan–Lusztig bases and sandwich cellularity	10
4. Diagram algebras and sandwich cellularity	15
References	29

1. INTRODUCTION

One of the main tools to study the representation theory of algebras is the notion of a *cellular algebra* due to Graham–Lehrer [GL96], and cellular algebras are nowadays ubiquitous in representation theory. *Sandwich cellular algebras* can be seen as a common and strict generalization of Graham–Lehrer’s cellular algebras and affine cellular algebras as in [KX12], as well as partial generalizations of Kazhdan–Lusztig bases, monoid and diagram algebras. All of these, in a precise sense, fit under the umbrella of sandwich cellularity.

These sandwich cellular algebras are certain algebras equipped with a sandwich cell datum which in turn gives rise to the notion of *cells* for these algebras. Cells partition sandwich cellular algebras in the same way as *Green’s relations* [Gre51] partition monoids, or more generally semigroups (we stay with monoids in this paper for simplicity), and *Kazhdan–Lusztig cells* [KL79] partition Hecke algebras. (The latter is the reason why we use the name cells.) Sandwich cellular structures were also successfully applied very early on, albeit in disguise, for example in the study of Brauer algebras as in [Bro55] and [FG95].

The analogy between sandwich cellular algebras and monoids respectively Hecke algebras goes even further. Part of a sandwich cell datum are *sandwich cellular bases* and *sandwiched algebras*. For cellular algebras all sandwiched algebras are trivial, and the sandwiched algebras are the main new ingredient in the theory. These sandwiched algebras play the role of Green’s H -groups [Gre51] and asymptotic Hecke algebras associated to intersections of left and right cells as e.g. in [Lus87]. The sandwich cellular bases are versions of Kazhdan–Lusztig bases in the theory of sandwich cellularity.

The most important theorem regarding sandwich cellular algebras is *H -reduction*, see [Theorem 2A.17](#). H -reduction classifies simple modules of sandwich cellular algebras by using their cells and the sandwiched algebras. In the theory of monoids H -reduction is known as the celebrated *Clifford–Munn–Ponizovskiĭ theorem* that classifies simple modules of monoids by Green’s J -classes and the simple modules of the H -groups, see [GMS09] or [Ste16] for modern expositions. In the theory of Kazhdan–Lusztig cells the corresponding theorem does not have a name (as far as we know), and is weaker than the Clifford–Munn–Ponizovskiĭ theorem, see [Section 3](#) for details. A variant of the H -reduction for sandwich cellular algebras is the theorem with the same name in categorification, see e.g. [MMMZ20] and [MMM⁺21], where we got its name from. Note however that this categorical H -reduction is similar in spirit but different in nature, which becomes evident when comparing [MMM⁺19] with [Section 3](#).

Mathematics Subject Classification 2020. Primary: 16G10, 16G99; Secondary: 18M30, 20C08, 20M30.

Keywords. Cellular algebras, representations of monoids and semigroups, Kazhdan–Lusztig theory and Hecke algebras, diagram algebras.

It turns out that all algebras are sandwich cellular, and the main point is to find a useful sandwich cell structure. Examples of sandwich cellular algebras with such cell structures include many algebras from various diverse fields of mathematics: all cellular algebras with useful cell datum, some Hecke algebras and their (p) -Kazhdan–Lusztig, many monoid algebras such as transformation monoids, many diagram algebras such as Brauer algebras, (weighted) KLRW algebras seem to have natural sandwich cellular structures, see e.g. [Bow22], [MT21] and [MT22], diagram algebras that appear in the study of higher genus knot invariants [RT21], [TV21] have useful sandwich cell structures. Note that some, but not all, of the named algebras are cellular but the sandwich cell structure is often much easier as we will see.

Sandwich cellular algebras are around for a long time, however, in various disguises and often implicit in the literature. They originate from at least four perspectives as mentioned above. In historical order, they appeared via the Clifford–Munn–Ponizovskii theorem in monoid theory, in the study of the Brauer algebra, they are related to Kazhdan–Lusztig cells, and are generalizations of cellular algebras. In this paper we will draw connections between these different fields by taking all of these perspectives at once. In this paper we:

- Give a concise summary and advance the theory of sandwich cellular algebras at the same time, see Section 2. In Section 2 we give a reformulation of the original definition that is useful in practice and explains our choice of using cell theory in the title. We also make the connection to cellular algebras and monoid theory precise.
- In Section 3 we apply the theory of sandwich cellular algebras to Hecke algebras of finite Coxeter type and their Kazhdan–Lusztig and p -Kazhdan–Lusztig bases.
- In Section 4 we study various diagram algebras from the viewpoint of sandwich cellular algebras. This includes diagram algebras without antiinvolution such as transformation and planar transformation monoids, as well as diagram algebras with an antiinvolution such as Brauer and Temperley–Lieb algebras.
- In all the examples we study, we classify simple modules using H -reduction and also compute some dimensions of simple modules, which is often doable with the general theory of sandwich cellular algebras at hand.

Quite a few, but not all, results in this paper have been obtain before, sometimes a long time ago. However, our point is that all of them fit under the umbrella of sandwich cellularity. At the end of each section we, for convenience, systematically collected references and explain how the results in this paper compare to known theorems in the literature.

Acknowledgments. We like to thank the organizers of QUACKS (Quantum groups, Categorification, Knot invariants, and Soergel bimodules) 2022 for the opportunity to present the theory of cells during that event. This made us think more carefully about the various incarnations of cells and ultimately led to this paper. Special thanks to James East for pointers to the monoid/semigroup literature and valuable help with GAP.

We are supported, in part, by the Australian Research Council as well as by the aphorism “Publish or perish”.

2. CELL THEORY FOR ALGEBRAS

We now recall the notion of a sandwich cellular algebra, and discuss a modification of sandwich cellularity which is useful in practice (in particular, for all algebras in this paper).

2A. Sandwich cellular algebras. We start with notation:

Notation 2A.1. We use the following conventions.

- (a) We fix a commutative unital ring \mathbb{K} . This is our ground ring throughout, and e.g. ranks $\text{rk}_{\mathbb{K}}$ are with respect to \mathbb{K} if not specified otherwise. We sometimes we need \mathbb{K} to be a field.
- (b) Throughout this section let \mathcal{A} denote an associative unital \mathbb{K} -algebra. Algebras in this paper are always assumed to be associative and unital.
- (c) Whenever we define an order, say $<_{\mathcal{P}}$, then we will also use $>_{\mathcal{P}}$, $\leq_{\mathcal{P}}$ or $\geq_{\mathcal{P}}$, having the evident meanings. We also write e.g. $\mathcal{A}^{>_{\mathcal{P}}\lambda}$ which stands for the \mathbb{K} -submodule of \mathcal{A} spanned by $\{c_{U,n,V}^{\mu} \mid \mu >_{\mathcal{P}} \lambda, U \in \mathcal{T}(\mu), V \in \mathcal{B}(\mu), n \in B_{\mu}\}$.

- (d) Unless otherwise specified, modules will always be left modules. In diagrammatic terms these are given by acting from the top, and we use

$$ab = \begin{array}{|c|} \hline a \\ \hline b \\ \hline \end{array}.$$

We sometimes use right modules and bimodules, and we will stress whenever that is the case.

Remark 2A.2. As in [Notation 2A.1](#), we will use colors in this paper. The colors are a visual aid only and the paper is readable in black-and-white without restrictions.

The following definition is a modification of [[MT22](#), Definition 2A.2], see also [[TV21](#), Section 2] for the same definition using a slightly different formulation.

Definition 2A.3. A *sandwich cell datum* for \mathcal{A} is a quadruple $(\mathcal{P}, (\mathcal{T}, \mathcal{B}), (\mathcal{H}_\lambda, B_\lambda), C)$, where:

- $\mathcal{P} = (\mathcal{P}, <_{\mathcal{P}})$ is a poset (the *middle poset* with *sandwich order* $<_{\mathcal{P}}$),
- $\mathcal{T} = \bigcup_{\lambda \in \mathcal{P}} \mathcal{T}(\lambda)$ and $\mathcal{B} = \bigcup_{\lambda \in \mathcal{P}} \mathcal{B}(\lambda)$ are collections of finite sets (the *top/bottom sets*),
- For $\lambda \in \mathcal{P}$ we have algebras \mathcal{H}_λ (the *sandwiched algebras*) and bases B_λ of \mathcal{H}_λ ,
- $C: \prod_{\lambda \in \mathcal{P}} \mathcal{T}(\lambda) \times B_\lambda \times \mathcal{B}(\lambda) \rightarrow \mathcal{A}; (T, m, B) \mapsto c_{T,m,B}^\lambda$ is an injective map,

such that:

(AC₁) The set $B_{\mathcal{A}} = \{c_{T,m,B}^\lambda \mid \lambda \in \mathcal{P}, T \in \mathcal{T}(\lambda), B \in \mathcal{B}(\lambda), m \in B_\lambda\}$ is a basis of \mathcal{A} . (We call $B_{\mathcal{A}}$ a *sandwich cellular basis*.)

(AC₂) For all $x \in \mathcal{A}$ there exist scalars $r_{TU}^x \in \mathbb{K}$ that do not depend on B or on m , such that

$$(2A.4) \quad xc_{T,m,B}^\lambda \equiv \sum_{U \in \mathcal{T}(\lambda)} r_{TU}^x c_{U,m,B}^\lambda \pmod{\mathcal{A}^{>_{\mathcal{P}}\lambda}}.$$

(AC₃) There exists a free \mathcal{A} - \mathcal{H}_λ -bimodule $\Delta(\lambda)$, a free \mathcal{H}_λ - \mathcal{A} -bimodule $\nabla(\lambda)$, and an \mathcal{A} -bimodule isomorphism

$$(2A.5) \quad \mathcal{A}_\lambda = \mathcal{A}^{\geq_{\mathcal{P}}\lambda} / \mathcal{A}^{>_{\mathcal{P}}\lambda} \cong \Delta(\lambda) \otimes_{\mathcal{H}_\lambda} \nabla(\lambda).$$

We call \mathcal{A}_λ the *cell algebra*, and $\Delta(\lambda)$ and $\nabla(\lambda)$ left and right *cell modules*.

The algebra \mathcal{A} is a *sandwich cellular algebra* if it has a sandwich cell datum.

Definition 2A.6. In the setup of [Definition 2A.3](#) assume that $\mathcal{T}(\lambda) = \mathcal{B}(\lambda)$ for all $\lambda \in \mathcal{P}$, and that there is a antiinvolution $(-)^*: \mathcal{A} \rightarrow \mathcal{A}$ such that:

(AC₄) We have $(c_{T,m,B}^\lambda)^* \equiv c_{B,m,T}^\lambda \pmod{\mathcal{A}^{>_{\mathcal{P}}\lambda}}$.

In this case we call the sandwich cell datum *involutive* and write $(\mathcal{P}, \mathcal{T}, (\mathcal{H}_\lambda, B_\lambda), C, (-)^*)$ for it.

Notation 2A.7. We will also say, for example, that \mathcal{A} itself is sandwich cellular, although this is a bit misleading since an algebra can be sandwich cellular with different sandwich cell data.

Example 2A.8. Two crucial special cases of involutive sandwich cellular algebras are:

- If all sandwiched algebras are isomorphic to \mathbb{K} , then $(\mathcal{P}, \mathcal{T}, (\mathcal{H}_\lambda \cong \mathbb{K}, B_\lambda = \{1\}), C, (-)^*)$ is a cell datum and \mathcal{A} is a *cellular algebra* in the sense of [[GL96](#)]. (Strictly speaking \mathcal{A} is cellular in the sense of [[GG11](#)] due to the weakened condition (AC₄) on the antiinvolution. Throughout, we will always use this weaker version of cellularity.)
- If all sandwiched algebras are commutative, then $(\mathcal{P}, \mathcal{T}, (\mathcal{H}_\lambda, B_\lambda), C, (-)^*)$ is an affine cell datum and \mathcal{A} is an *affine cellular algebra* in the sense of [[KX12](#)].

The mnemonic (in particular for readers familiar with diagram algebras) is:

$$(2A.9) \quad \begin{array}{l} \text{cellular: } c_{T,1,B}^\lambda \rightsquigarrow \begin{array}{|c|} \hline T \\ \hline B \\ \hline \end{array} \leftarrow \mathcal{H}_\lambda \cong \mathbb{K}, \quad \text{affine cellular: } c_{T,m,B}^\lambda \rightsquigarrow \begin{array}{|c|} \hline T \\ \hline m \\ \hline B \\ \hline \end{array} \leftarrow \text{commutative } \mathcal{H}_\lambda, \\ \\ \text{sandwich cellular: } c_{T,m,B}^\lambda \rightsquigarrow \begin{array}{|c|} \hline T \\ \hline m \\ \hline B \\ \hline \end{array} \leftarrow \text{general } \mathcal{H}_\lambda, \end{array}$$

where we assume the existence of an antiinvolution for the top two pictures. ◇

The comparison of sandwich cellular and cellular is as follows.

Proposition 2A.10. *An involutive $(\mathcal{P}, \mathcal{T}, (\mathcal{H}_\lambda, B_\lambda), C, (-)^*)$ sandwich cell datum is a cell datum if and only if $\mathcal{H}_\lambda \cong \mathbb{K}$ for all $\lambda \in \mathcal{P}$.*

Moreover, an involutive sandwich cellular algebra \mathcal{A} such that all \mathcal{H}_λ are cellular (with the same antiinvolution) is cellular with a refined sandwich cell datum. Conversely, if at least one \mathcal{H}_λ is non-cellular, then \mathcal{A} is non-cellular.

Proof. The first claim is immediate, for the second see [TV21, Proposition 2.9]. \square

Example 2A.11. Let $\mathcal{A} = \mathbb{K}[G]$ for a group G . Then the group element basis is a sandwich cellular basis which is not a cellular or affine cellular basis in general.

In Proposition 2D.1 we give a condition (that can likely be weakened) on a monoid that ensures that the monoid element basis is a sandwich cellular basis. \diamond

We will discuss more examples, and also return to Example 2A.11, later on.

Proposition 2A.12. *Any algebra has the structure of a sandwich cellular algebra. Moreover, if \mathbb{K} is an algebraically closed field, then there exists a sandwich cell datum with $\mathcal{H}_\lambda \cong \mathbb{K}$ for all $\lambda \in \mathcal{P}$.*

Proof. For the first statement take all ingredients to be trivial, e.g. $\mathcal{P} = \{0\}$, $\mathcal{T}(0) = \mathcal{B}(0) = \{0\}$, $\mathcal{H}_0 = \mathcal{A}$ etc. After comparison of definitions, the second claim is [CZ19, Corollary B]. \square

Remark 2A.13. By Proposition 2A.12, the main point is not to prove that an algebra is sandwich cellular, but rather to find a useful sandwich cell datum. Note also that (the proof of) Proposition 2A.10 and Proposition 2A.12 imply that any algebra is cellular over an algebraically closed field if one not assumes the existence of an antiinvolution.

Notation 2A.14. The \mathcal{A} -bimodule \mathcal{A}_λ has an induced multiplication, but might not have a unit in this multiplication. If it does not have a unit, then we always adjoin a unit whenever we want to see it as an algebra.

An *apex* of a \mathcal{A} -module M , if it exists, is a maximal $\lambda \in \mathcal{P}$ such that $\mathbb{K}\mathcal{J}$ does not annihilate M . The same notion is used for \mathcal{A}_λ .

Example 2A.15. The cell algebra \mathcal{A}_λ has at most two apexes, one for the unit and λ . \diamond

Lemma 2A.16. *Every simple \mathcal{A} -module has an apex $\lambda \in \mathcal{P}$, and similarly for simple \mathcal{A}_λ -modules.*

Proof. See [TV21, Lemma 2.15]. \square

The main theory about sandwich cellular algebras is:

Theorem 2A.17. (*H-reduction a.k.a. Clifford–Munn–Ponizovskii theorem*) *Let \mathbb{K} be a field.*

(a) *If $\lambda \in \mathcal{P}$ is an apex and \mathcal{H}_λ is Artinian, then we have bijections*

$$\begin{array}{c} \text{Start: } \{ \text{simple } \mathcal{A}\text{-modules with apex } \lambda \} / \cong \\ \leftarrow \begin{array}{c} 1:1 \\ \leftarrow \rightarrow \end{array} \\ \text{J-reduction: } \{ \text{simple } \mathcal{A}_\lambda\text{-modules with apex } \lambda \} / \cong \\ \leftarrow \begin{array}{c} 1:1 \\ \leftarrow \rightarrow \end{array} \\ \text{H-reduction: } \quad \{ \text{simple } \mathcal{H}_\lambda\text{-modules} \} / \cong . \end{array}$$

(b) *All \mathcal{A} -modules with apex λ have composition factors of apex μ with $\mu \leq_{\mathcal{P}} \lambda$.*

Proof. See [TV21, Theorem 2.16]. \square

Notation 2A.18. Whenever Theorem 2A.17 applies, we will write $\mathcal{P}^{ap} \subset \mathcal{P}$ for the set of apexes, and $L(\lambda, K)$ for the simple \mathcal{A} -modules associated to $\lambda \in \mathcal{P}^{ap}$ and a simple \mathcal{H}_λ -module K . If $\mathcal{H}_\lambda \cong \mathbb{K}$, then we write $L(\lambda)$ for the unique simple \mathcal{A} -module of apex λ .

Remark 2A.19. The bijections in Theorem 2A.17 can be made explicit and $L(\lambda, K)$ is the head of the induction of K to \mathcal{A} , see [TV21, Theorem 2.16] for details.

Notation 2A.20. As already indicated by the Artinian condition in Theorem 2A.17, there are some technicalities when working with infinite dimensional algebras, see [TV21, Section 2] for a more detailed treatment. To simplify the exposition, we will from now on assume that our algebras are finite dimensional.

Remark 2A.21. Sandwich cellular algebras are inspired by the theory of cellular algebras on the one hand, see e.g. [GL96], [HM10], [KX12], [AST18], [ET21], [GW15] or [TV21], and ideas coming from monoid representation theory on the other hand, see e.g. [Gre51], [GMS09] or [KST22]. There was also a huge influence from Kazhdan–Lusztig (KL) theory and based algebras [KL79], [Lus87], [KM16]. Sandwich cellular algebras have appeared in disguise in, for example, [Bro55], [FG95], [KX99] and [KX01]. We learned the idea underlying sandwich cellular algebras from [GW15].

2B. Green’s theory of cells in algebras. We now discuss a special case of sandwich cellular algebras for which we can reformulate [Definition 2A.3](#) to be closer to Green’s classical theory of cells a.k.a. Green’s relations, and KL theory.

Remark 2B.1. The main slogan of cell theory is the following. In contrast to groups, the multiplication in an algebra often destroys information. For example, if $b = ca$, then b can be obtained from a by left multiplication, and we can say that b is left bigger than a . In a group we can go back by $c^{-1}b = a$ so a is also left bigger than b , but this is not always possible in an algebra. Cells can then be thought of as keeping track of the information loss during multiplication. The reader is encouraged to keep this analogy in mind while reading the text below.

Recall that \mathcal{A} denotes an algebra as in [Notation 2A.1](#). Fix a basis $B_{\mathcal{A}}$ of \mathcal{A} . Everything below depends on the choice of this basis, and we will use the *(based) pair* $(\mathcal{A}, B_{\mathcal{A}})$.

For $a, b, c \in B_{\mathcal{A}}$ we write $b \in ca$ if, when expanded in terms of $B_{\mathcal{A}}$, b appear with nonzero coefficient in ca . We define preorders on $B_{\mathcal{A}}$ by

$$(a \leq_l b) \Leftrightarrow \exists c : b \in ca, \quad (a \leq_r b) \Leftrightarrow \exists c : b \in ac, \quad (a \leq_{lr} b) \Leftrightarrow \exists c, d : b \in cad.$$

We call these left, right and two-sided *cell orders*.

Definition 2B.2. We define equivalence relations by

$$(a \sim_l b) \Leftrightarrow (a \leq_l b \text{ and } b \leq_l a), \quad (a \sim_r b) \Leftrightarrow (a \leq_r b \text{ and } b \leq_r a), \quad (a \sim_{lr} b) \Leftrightarrow (a \leq_{lr} b \text{ and } b \leq_{lr} a).$$

The respective equivalence classes are called left, right respectively two-sided *cells*.

We also say *J-cells* instead of two-sided cells, following the notation in [Gre51].

Definition 2B.3. An *H-cell* $\mathcal{H} = \mathcal{H}(\mathcal{L}, \mathcal{R}) = \mathcal{L} \cap \mathcal{R}$ is an intersection of a left \mathcal{L} and a right cell \mathcal{R} .

The following is easy:

Lemma 2B.4. *The left, right and J-cell orders induce preorders on the left, right and J-cells, respectively. Similarly, we can also compare elements and cells via the cell orders.* \square

We will use the same symbols for the various preorders.

Example 2B.5. Let $\mathcal{A} = \mathbb{K}S$ for a finite monoid S (all monoids we use are finite, and we drop that adjective), and fix $B_{\mathcal{A}} = S$, the monoid basis.

(a) The above recovers Green’s relations. That is, the preorders simplify to


$$(a \leq_l b) \Leftrightarrow \exists c : b = ca, \quad (a \leq_r b) \Leftrightarrow \exists c : b = ac, \quad (a \leq_{lr} b) \Leftrightarrow \exists c, d : b = cad.$$

In monoid theory the corresponding cells are called \mathcal{L} , \mathcal{R} , \mathcal{J} and \mathcal{H} -*classes*.

(b) Let $G \subset S$ be the group of invertible elements. We have $G \leq_l a$, $G \leq_r a$ and $G \leq_{lr} a$ for all $a \in S$. This can be seen by $a = ag^{-1}g$ and similar calculations for the right and *J*-orders. In particular, if $G = S$ (thus, S is a group), then there is only one cell, which is a left, right, *J* and *H*-cell at the same time.

If not specified otherwise, when referring to monoids or groups, then we will always fix $B_{\mathcal{A}}$ to be the monoid basis. \diamond

Remark 2B.6. We think of cells as a matrix decomposition of $B_{\mathcal{A}}$:



In the left picture we use matrix notation for the twelve H -cells in \mathcal{J} so that \mathcal{H}_{ij} is the intersection of the i th right with the j th left cell of \mathcal{J} . The J -cell \mathcal{J} is then a matrix with entries from H -cells. Moreover, the left cells are column vectors and the right cells are row vectors of \mathcal{J} .

The right picture is a GAP output using the package Semigroups, see <https://www.gap-system.org/Packages/semigroups.html> (link last checked June 15, 2022). The illustration shows the cell structure of a given, not further specified, monoid in the matrix notation. The J -order is illustrated as well. The shaded H -cells are strictly idempotent in the sense of Definition 2B.12 below.

We denote the indexing set for the J -cells by \mathcal{P} , and denote the J -cells by \mathcal{J}_λ for $\lambda \in \mathcal{P}$.

Notation 2B.8. We use a subscript λ to indicate that we are working in a fixed J -cell \mathcal{J}_λ .

The next lemma is immediate and will be used throughout.

Lemma 2B.9. $\mathcal{A}^{>lr^\lambda} = \mathbb{K}\{\bigcup_{\mu \in \mathcal{P}, \mu > lr^\lambda} \mathcal{J}_\mu\}$ is a two-sided ideal in \mathcal{A} . \square

Example 2B.10. If \mathcal{J}_λ is the J -cell of size 3-3 in (2B.7), then $\mathcal{A}^{>lr^\lambda}$ is the two-sided ideal supported on the size 9-9 and the top J -cells. Note that we can ignore the J -cell of size 2-2 as the multiplication will never get us from \mathcal{J}_λ into it. \diamond

Notation 2B.11. An equation etc. holds *up to higher order terms* if it holds modulo $\mathcal{A}^{>lr^\lambda}$.

Definition 2B.12. If the \mathbb{K} -linear span of a J -cell \mathcal{J} contains a pseudo-idempotent up to higher order terms, that is, $e \in \mathbb{K}\mathcal{J}$ with $e^2 = s(e) \cdot e \pmod{\mathcal{A}^{>lr^\lambda}}$ for $s(e) \in \mathbb{K} \setminus \{0\}$, then we call \mathcal{J} *idempotent*. We say \mathcal{J} is *strictly idempotent* if \mathcal{J} itself contains a pseudo-idempotent up to higher order terms. We use the same terminology for H -cells.

Note that pseudo-idempotents can be rescaled to idempotents if $s(e)$ is invertible, which is always true, for example, if \mathbb{K} is a field. We write $\mathcal{J}_\lambda(e)$ and $\mathcal{H}_\lambda(e) \subset \mathcal{J}_\lambda(e)$ for idempotent cells.

Example 2B.13. All idempotent H and J -cells are strictly idempotent for monoids. The notion then corresponds to having an idempotent. \diamond

A *bottom J -cell* is a J -cell that is minimal in the J -order. Similarly, a *top J -cell* is a J -cell that is maximal in the J -order.

Proposition 2B.14. For the pair $(\mathcal{A}, B_{\mathcal{A}})$ we have:

- (a) Every H -cell is contained in some J -cell, and every J -cell is a disjoint union of H -cells.
 - (b) If \mathbb{K} is a field, then $\mathcal{H}_\lambda = \mathbb{K}\mathcal{H}_\lambda(e) / \mathcal{A}^{>lr^\lambda}$ is an algebra with identity $\frac{1}{s(e)}e$. This algebra is a subalgebra of $\mathcal{A}^{\leq lr^\lambda} = \mathcal{A} / \mathcal{A}^{>lr^\lambda}$ and $\mathcal{A}^\lambda = \mathcal{A}^{\leq lr^\lambda} \cap \mathbb{K}\mathcal{J}_\lambda$.
 - (c) Assume that $1 \in B_{\mathcal{A}}$. The pair $(\mathcal{A}, B_{\mathcal{A}})$ has a unique bottom J -cell, and this J -cell is strictly idempotent.
 - (d) The pair $(\mathcal{A}, B_{\mathcal{A}})$ has a unique top J -cell, which is an H -cell at the same time.
- (a) works analogously for left and right cells in a fixed J cell.

If they exists, then we write \mathcal{J}_b and \mathcal{J}_t for the bottom and top J -cell, respectively.

These are obtained from the previous ones by putting idempotents (above in denoted as a box) on the bottom. \square

Definition 2C.4. Under the assumptions from Lemma 2C.3, the *sandwich matrix* $S^{\lambda, K}$ (associated to $\lambda \in \mathcal{P}^{ap}$ and a simple \mathcal{H}_λ -module K) is the $(|\mathcal{H}_\lambda| \cdot \#\mathcal{R}_\lambda)$ - $(\#\mathcal{L}_\lambda \cdot |\mathcal{H}_\lambda|)$ -matrix with values in \mathcal{H}_λ given by Lemma 2C.3.

For a sandwich pair $(\mathcal{A}, B_{\mathcal{A}})$, the *Gram matrix* G^λ (associated to $\lambda \in \mathcal{P}^{ap}$) is the $(\#\mathcal{R}_\lambda)$ - $(\#\mathcal{L}_\lambda)$ -matrix with values in \mathbb{K} defined by

$$(G^\lambda)_{i,j} = \begin{cases} s(e) & \text{if } \mathcal{H}_{ij} \text{ is idempotent with eigenvalue } s(e), \\ 0 & \text{else.} \end{cases}$$

In other words, G^λ is the matrix with entries being the eigenvalues of pseudo-idempotents.

Example 2C.5. For admissible monoids the sandwich matrix is a variation of the matrix with the same name in monoid theory, see e.g. [Ste16, Section 5.4]. Precisely, the matrix given in [Ste16, Section 5.4] would be the analog of the matrix for ϕ^λ .

The Gram matrices for the monoid with cell structure as in (2B.7) are identity matrices of sizes 1-1 (twice), 3-3, 2-2 and 9-9. \diamond

Proposition 2C.6. Let \mathbb{K} be a field and $(\mathcal{A}, B_{\mathcal{A}})$ be a sandwich pair.

(a) Under the assumptions from Lemma 2C.3, for $\lambda \in \mathcal{P}^{ap}$ and a simple \mathcal{H}_λ -modules K , we have

$$\dim_{\mathbb{K}}(L(\lambda, K)) = \text{rank}_{\mathbb{K}}(S^{\lambda, K}).$$

(b) For $\mathcal{H}_\lambda \cong \mathbb{K}$ and all $\lambda \in \mathcal{P}^{ap}$ we have

$$\dim_{\mathbb{K}}(L(\lambda)) = \text{rank}_{\mathbb{K}}(G^\lambda).$$

Proof. Statement (a). This follows from [TV21, Lemma 2.13] and the proof of Lemma 2C.3.

Statement (b). For $\mathcal{H}_\lambda \cong \mathbb{K}$, the Gram matrix is the sandwich matrix, so (a) applies. \square

As we will see in Theorem 4A.17 is in general not enough to get the simple dimensions and one needs to know the (in general much bigger) sandwich matrices.

Remark 2C.7. As we wrote above, the idea of using pairing and matrices to determine information about modules is everywhere in quantum algebra and related fields. Its a bit hard to track down, but see [CP61, Section 5.2] for an early reference. The above is the version in the theory of sandwich cellular algebras.

2D. Monoids and sandwich cellularity. We call a monoid S *admissible* if all J -cells are either idempotent (with H -cells of arbitrary size), or have H -cells of size one.

Proposition 2D.1. For any admissible monoid S (with an antiinvolution) we have a(n involutive) sandwich pair $(\mathbb{K}S, S)$.

Proof. The crucial fact about monoids we need is that left, right and H -cells in one J -cells are always of the same size, see [Gre51, Theorem 1]. In particular, we can let \mathcal{P} be the poset coming from Green cells, we can let \mathcal{T} and \mathcal{B} be indexed by right and left cells, and $B_\lambda = \mathcal{H}$ to be any H -cell in \mathcal{J}_λ . This choice satisfies (AC₁) and (AC₂), by construction.

We need to work a bit more for (AC₃). As free \mathbb{K} -modules we can take $\Delta(\lambda) = \mathbb{K}\mathcal{L}$, $\nabla(\lambda) = \mathbb{K}\mathcal{R}$ and $\mathcal{H}_\lambda = \mathbb{K}\mathcal{H}$ for any choice of cells. As a free \mathbb{K} -module we thus get $\mathcal{A}_\lambda \cong \Delta(\lambda) \otimes_{\mathbb{K}} \mathcal{H}_\lambda \otimes_{\mathbb{K}} \nabla(\lambda)$. If the H -cells in \mathcal{J} are of size one, then this construction satisfies (AC₃).

The final case is when $\mathcal{J}_\lambda(e)$ is idempotent, but has arbitrary sized H -cells. In this case $\mathcal{J}_\lambda(e)$ is strictly idempotent by Example 2B.13. It then follows that $\mathcal{H}_\lambda(e)$ is a subgroup by [Gre51, Theorem 7]. We can take $\mathcal{H}_\lambda = \mathbb{K}\mathcal{H}_\lambda(e)$ and $\Delta(\lambda) = \mathbb{K}\mathcal{L}$, $\nabla(\lambda) = \mathbb{K}\mathcal{R}$ for the left and right cells defining $\mathcal{H}_\lambda(e)$ via their intersection. It is then easy to see (again with reference to [Gre51]) that (AC₃) holds for these choices since $\mathcal{H}_\lambda(e)$ acts freely on its defining left and right cell.

Finally, having an antiinvolution on S clearly implies (AC₄). \square

Example 2D.2. For admissible monoids the cell modules $\Delta(\lambda)$ are the classical Schützenberger modules with their origin in [Sch58]. \diamond

Proposition 2D.3. *For any admissible monoid S with an antiinvolution the sandwich pair $(\mathbb{K}S, S)$ can be refined into a cellular pair if and only if all $\mathbb{K}\mathcal{H}(e)$ are cellular algebras with a compatible antiinvolution.*

Proof. Directly from [Proposition 2A.10](#) and the proof of [Proposition 2D.1](#). \square

Remark 2D.4. The discussion in this section is new, but the question whether monoid algebras are cellular is well-studied, see e.g. [\[Eas06\]](#) which is very similar to [Proposition 2D.3](#), but our point is that this follows from the general theory of sandwich cellular algebras.

3. KAZHDAN–LUSZTIG BASES AND SANDWICH CELLULARITY

The KL bases of Hecke algebras partially motivated the definition of sandwich cellularity. In this section we discuss the relation between these two notions.

3A. The classical story. Throughout this section, let $W = (W, S)$ be a (connected finite) Coxeter system. We identify W with its Coxeter diagram.

Definition 3A.1. For a parameter v let $H_W = H_{(W,S)}$ denote the $\mathbb{Z}[v, v^{-1}]$ -algebra generated by $\{H_s \mid s \in S\}$ subject to the braid relations and $H_s^2 = 1 + (v^{-1} - v)H_s$.

Notation 3A.2. Throughout this section \mathbb{K} will denote a specialization of $\mathbb{Z}[v, v^{-1}]$. Everything below is defined over \mathbb{K} because the main ingredients are defined over $\mathbb{Z}[v, v^{-1}]$.

By [\[KL79, Theorem 1.1\]](#) there exists a distinguished basis of H_W that we call the **KL basis** and denote by $B_W = B_{(W,S)} = \{b_w \mid w \in W\}$. We will consider the pair (H_W, B_W) .

Remark 3A.3. The precise conventions for H_W and its distinguished basis will only be of importance when we do explicit calculations in dihedral type. To not distract the reader from the main points, we specify our conventions in [Section 3C](#) after the main statements.

Example 3A.4. For the pair (H_W, B_W) the cells are the KL cells from [\[KL79\]](#). Explicitly, take $\mathbb{K} = \mathbb{C}(v)$ and consider the dihedral type Coxeter system $I_2(n)$ determined by \bullet^n with the left node called 1 and the right node called 2. Then we have the cell structure

$$\begin{array}{ccc}
 \mathcal{J}_t & \begin{array}{|c|} \hline b_{1212} \\ \hline \end{array} & \mathcal{J}_t & \begin{array}{|c|} \hline b_{12121} \\ \hline \end{array} \\
 n = 4 \text{ (type } B_2\text{): } \mathcal{J}_m & \begin{array}{|c|c|} \hline b_1, b_{121} & b_{12} \\ \hline b_{21} & b_2, b_{212} \\ \hline \end{array} & , & n = 5 \text{ (type } H_2\text{): } \mathcal{J}_m & \begin{array}{|c|c|} \hline b_1, b_{121} & b_{12}, b_{1212} \\ \hline b_{21}, b_{2121} & b_2, b_{212} \\ \hline \end{array} \\
 \mathcal{J}_b & \begin{array}{|c|} \hline b_\emptyset \\ \hline \end{array} & & \mathcal{J}_b & \begin{array}{|c|} \hline b_\emptyset \\ \hline \end{array}
 \end{array}$$

The pattern for general even and n odd is the same: in the even case the diagonal H -cells have one more element than the off-diagonal H -cells, while they have the same size for n odd.

As before, the marked H -cells indicate strictly idempotent (diagonals) and idempotent but not strictly idempotent (off-diagonals), respectively, in the sense of [Definition 2B.12](#). This is where we use that $\mathbb{K} = \mathbb{C}(v)$ in this example, the rest works over any \mathbb{K} . \diamond

Theorem 3A.5. *For a Coxeter system W we have:*

- (a) *The pair (H_W, B_W) is a sandwich pair if and only if W is of type A or type $I_2(n)$ for n odd.*
- (b) *If W is of type A or type $I_2(n)$ for n odd, then the sandwich pair (H_W, B_W) is involutive with the usual bar involution.*

Proof. Statement (a). If the Coxeter graph is not of type A or type $I_2(n)$ for n odd, then one will always find H -cells of different sizes within one J -cell, see e.g. [\[Lus84\]](#) for Weyl types and [\[MMM⁺19, Section 8\]](#) for the other types. So these types can not give a sandwich pair (H_W, B_W) by [Theorem 2B.18](#). Conversely, in type A the KL basis is a cellular basis, as follows from [\[KL79\]](#), and for type $I_2(n)$ with n odd sandwich cellularity can be easily verified by hand using the calculations in [Section 3C](#).

Statements (b). This can be proven using the results in [\[KL79\]](#). \square

The following example shows that the assumption of H -cells being of the same size within one J -cell is crucial for [Theorem 2A.17](#) to work.

Example 3A.6. In Example 3A.4 consider the non-quantum case over $\mathbb{K} = \mathbb{C}$, that is the group algebras of dihedral groups equipped with the KL basis B_{I_n} (we write I_n for $I_2(n)$).

We consider $n = 4$ and $n = 5$, the dihedral groups D_4 and D_5 of orders 8 and 10. The character tables are (the conjugacy classes index the columns, the simple characters the rows and ϕ is the golden ratio):

$$(3A.7) \quad D_4: \begin{array}{c|ccccc} & \chi^1 & \chi^2 & \chi^3 & \chi^4 & \chi^5 \\ \hline \chi_t & 1 & 1 & 1 & 1 & 1 \\ \chi_b & 1 & 1 & -1 & 1 & -1 \\ \chi_{m1} & 1 & 1 & 1 & -1 & -1 \\ \chi_{m2} & 1 & 1 & -1 & -1 & 1 \\ \chi_{m3} & 2 & -2 & 0 & 0 & 0 \end{array}, \quad D_5: \begin{array}{c|cccc} & \chi^1 & \chi^2 & \chi^3 & \chi^4 \\ \hline \chi_t & 1 & 1 & 1 & 1 \\ \chi_b & 1 & -1 & 1 & 1 \\ \chi_{m1} & 2 & 0 & \phi & \phi^2 \\ \chi_{m2} & 2 & 0 & \phi^2 & \phi \end{array}.$$

Now, $(\mathbb{C}[D_5], B_{I_5})$ is a sandwich pair, with sandwiched algebras $\mathcal{H}_b \cong \mathbb{C}$, $\mathcal{H}_m \cong \mathbb{C}[\mathbb{Z}/2\mathbb{Z}]$ and $\mathcal{H}_t \cong \mathbb{C}$. Thus, H -reduction Theorem 2A.17 gives the expected classification of simple D_5 -modules as in (3A.7). In contrast, $(\mathbb{C}[D_4], B_{I_4})$ is not a sandwich pair since the algebras one gets are $\mathcal{H}_b \cong \mathbb{C}$, $\mathcal{H}_m \cong \mathbb{C}[\mathbb{Z}/2\mathbb{Z}]$ or $\mathcal{H}_m \cong \mathbb{C}$ and $\mathcal{H}_t \cong \mathbb{C}$. The H -reduction for the middle cell fails: neither $\mathbb{C}[\mathbb{Z}/2\mathbb{Z}]$ nor \mathbb{C} give the expected count of three simple D_4 -modules. \diamond

For \mathbb{K} let $p \in \mathbb{Z}_{\geq 0}$ be minimal such that $p \cdot 1 = 0 \in \mathbb{K}$, and let $p = \infty$ if no such p exists. That is, $p = \text{char}(\mathbb{K})$ but we consider $\text{char}(\mathbb{K}) = 0$ as $p = \infty$.

We use the English convention for Young diagrams, i.e. we draw left-justified rows of boxes successively down the page. Let $P(\lambda|p)$ the set of p -restricted Young diagrams with λ boxes. That is, each element of $P(\lambda|p)$ is a Young diagram with λ boxes such that the difference between the length of two consecutive columns is $< p$ (empty columns are of length zero). Note that the set $P(\lambda|p)$ for $p > \lambda$ is the set of all Young diagrams with λ boxes. The set $P(\lambda|p) = (P(\lambda|p), <_d)$ is a poset when equipped with the dominance order $<_d$ (with conventions specified by Example 3A.8).

Example 3A.8. For example,

$$p > 3: \left\{ \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} <_d \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} <_d \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right\}, \quad p \in \{2, 3\}: \left\{ \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} <_d \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right\},$$

are the sets of p -restricted partitions of 3 for all possible p . \diamond

For $a \in \mathbb{Z}_{\geq 0}$, let $[a] = \frac{v^a - v^{-a}}{v - v^{-1}} \in \mathbb{Z}[v, v^{-1}]$ denote the usual quantum numbers. Below we will need the condition that $\mathbb{K} = \mathbb{F}(q)$ for \mathbb{F} an algebraically closed field and \mathbb{K} is such that quantum numbers in q are invertible. We call such fields **admissible**.

Example 3A.9. The fields $\mathbb{K} = \mathbb{K}(v)$ or $\mathbb{K} = \mathbb{C}$, where $q = v$ and $q = 1$ respectively, are examples of admissible fields. \diamond

Theorem 3A.10. Let (H_W, B_W) be an involutive sandwich pair.

(a) Let W be of type A .

(i) The J -cells of H_A are given by $\mathcal{P} = (P(\lambda|\infty), <_d)$.

(ii) The left cells and right cells of H_A are determined by the Robinson–Schensted correspondence.

(iii) A J -cell of H_A is strictly idempotent if and only if its corresponding partition is p -restricted. For strictly idempotent J -cells and \mathbb{K} a field we have

$$\mathcal{H}_\lambda \cong \mathbb{K}.$$

(iv) Let \mathbb{K} be a field. The set of apexes for simple H_A -modules is $\mathcal{P}^{ap} = (P(\lambda|p), <_d)$, and there is precisely one simple H_A -module $L(\lambda)$ for $\lambda \in \mathcal{P}^{ap}$.

(b) Let W be odd dihedral type $I_2(n)$.

(i) The J -cells of H_{I_n} are given by $\mathcal{P} = \{b <_{lr} m <_{lr} t\}$.

(ii) \mathcal{J}_b and \mathcal{J}_t are left and right cells, while the left cells in \mathcal{J}_m are given by reduced expressions starting with either 1 or 2, and the right cells by reduced expressions that end with either 1 or 2.

- (iii) \mathcal{J}_b is always strictly idempotent, \mathcal{J}_m is strictly idempotent if $[2]$ is invertible in \mathbb{K} , and \mathcal{J}_m is strictly idempotent if $[2][n]$ is invertible in \mathbb{K} . Assume \mathbb{K} is an admissible field. Then we have

$$\mathcal{H}_b \cong \mathbb{K}, \quad \mathcal{H}_m \cong \mathbb{K}[\mathbb{Z}/(\frac{n-1}{2})\mathbb{Z}], \quad \mathcal{H}_t \cong \mathbb{K}.$$

- (iv) Let \mathbb{K} be an admissible field. The set of apexes for simple H_{In} -modules is $\mathcal{P}^{ap} = \mathcal{P}$, and there is precisely one simple H_{In} -module for $\lambda \in \{b, t\} \subset \mathcal{P}^{ap}$, and there are precisely $\frac{n-1}{2}$ simple H_{In} -modules for $m \in \mathcal{P}^{ap}$. The dimensions of the simple H_{In} -modules are 1, 1 and 2, the latter $\frac{n-1}{2}$ times.

- (c) Let \mathbb{K} be an admissible field. We have that H_A and H_{In} for n odd are semisimple.

Proof. Statement (a). Proving how the cell structure looks like is not trivial, but still well-known and follows from [KL79]. Thus, it remains to verify part (iv). That $\mathcal{P}^{ap} = (P(\lambda|p), <_d)$ follows from the calculation of eigenvalues of distinguished involutions in the Hecke algebra, see [Lus03, Conjectures 14.2 and Chapter 15] (we mean the arXiv version of the book). The rest is then Theorem 3A.5 and H -reduction Theorem 2A.17.

Statement (b). By Section 3C below. That is, the hardest parts are proven in Lemma 3C.3 and Lemma 3C.4. The rest is easy to prove using the explicit description of the multiplication of KL basis elements given in Section 3C.

Statement (c). By the above and Proposition 2B.23, it remains to show that the $\Delta(\lambda)$ are simple. However, we already know the dimensions of the simples by (a).(iv) and (b).(iv), and the proof completes. \square

Example 3A.11. Back to Example 3A.6 for $n = 5$. The sandwich matrices for the bottom and top J -cell are $S^b = (b_\emptyset)$ and $S^t = (10 \cdot b_{12121})$, respectively. We compute the sandwich matrices $S^{m,1}$ and $S^{m,2}$ in the proof of Lemma 3C.4. These matrices are of rank 2. Thus, the bottom and top J -cells give one dimensional simple \mathcal{A} -modules, while the middle J -cell gives two simple \mathcal{A} -modules of dimension 2. \diamond

Proposition 3A.12. *The algebra H_{In} is cellular when n is odd.*

Proof. By Theorem 3A.5 and Proposition 2A.10 (and some care with the antiinvolution). \square

Remark 3A.13. Theorem 3A.5 also shows that the KL basis is cellular if and only if W is of type A. That the KL basis of type A is cellular follows from [KL79], as we wrote above. The converse is known, but hard to find in the literature. The case of general sandwich cellularity is new.

We point out that all finite type Hecke algebras are sandwich cellular, even cellular by [Gec07], and the statement in Theorem 3A.5 is that the KL basis is not a sandwich cellular basis.

Finally, our sandwich approach to classify simple H_{In} -modules from Theorem 3A.10 is new, but the results are certainly not new: In the semisimple case the classification follows directly from Tits' deformation theorem (explicitly, [GP00, Theorem 8.1.7]) and classical theory. We also stress that the approach taken in Theorem 3A.10 with a bit more work, see Remark 3C.6 below, also classifies simple H_{In} -modules over arbitrary fields. This classification is again well-known, see e.g. [GP00, Chapter 8] for a concise treatment and bibliographical remarks, but our point is that it also follows as part of the general theory of sandwich cellular algebras.

3B. p -Kazhdan–Lusztig bases and sandwich cellularity. For a fixed prime p e.g. [JW17] defines another distinguished basis $B_W^p = \{b_w^p \mid w \in W\}$ of H_W that we call the p -KL basis. Not much is known about the (sandwich) cellularity of the p -KL basis, and all we can say is:

Theorem 3B.1. *We have the following.*

- (a) *If W is of type A, then (H_W, B_W^p) is a sandwich pair for all primes, and the cell structure is the same as for (H_W, B_W) .*
- (b) *If W is of type $B_2, C_2, G_2, B_3, C_3, B_4, C_4$ or D_4 , then (H_W, B_W^p) is never a sandwich pair.*

Note that Theorem 3A.5.(c) and Theorem 3B.1.(a) give the same classification of simple H_W -modules via H -reduction Theorem 2A.17.

Proof. Statement (a). This is [Jen20b, Theorem 5.14].

Statement (b). We will show that (H_W, B_W^p) can not be a sandwich pair by contradicting [Theorem 2B.18.\(d\)](#). For $p > 2$ in types B_2 and C_2 and for $p > 3$ in type G_2 the p -KL basis is the usual KL basis, so [Theorem 3A.5](#) applies. For the remaining cases we simply list the p -KL cells, which have H -cells of different sizes:

$$\begin{array}{c}
 \text{type } B_2: \quad \begin{array}{c} b_{1212}^2 \\ \hline \begin{array}{|c|c|} \hline b_{121}^2 & b_{12}^2 \\ \hline b_{21}^2 & b_2^2, b_{212}^2 \\ \hline \end{array} \\ \hline b_1^2 \\ \hline b_\emptyset^2 \end{array}, \quad \text{type } C_2: \quad \begin{array}{c} b_{1212}^2 \\ \hline \begin{array}{|c|c|} \hline b_1^2, b_{121}^2 & b_{12}^2 \\ \hline b_{21}^2 & b_{212}^2 \\ \hline \end{array} \\ \hline b_2^2 \\ \hline b_\emptyset^2 \end{array}, \\
 \\
 \text{type } G_2: \quad \begin{array}{c} b_{1212}^2 \\ \hline \begin{array}{|c|c|} \hline b_{121}^2, b_{12121}^2 & b_{1212}^2 \\ \hline b_{2121}^2 & b_{212}^2, b_{21212}^2 \\ \hline \end{array} \\ \hline \begin{array}{|c|c|} \hline b_1^2 & b_{12}^2 \\ \hline b_{21}^2 & b_2^2 \\ \hline \end{array} \\ \hline b_\emptyset^2 \end{array}, \quad \begin{array}{c} b_{121212}^3 \\ \hline \begin{array}{|c|c|} \hline b_{121}^3, b_{12121}^3 & b_{12}^3, b_{1212}^3 \\ \hline b_{21}^3, b_{2121}^3 & b_2^3, b_{212}^3, b_{21212}^3 \\ \hline \end{array} \\ \hline b_1^3 \\ \hline b_\emptyset^3 \end{array}.
 \end{array}$$

Here we used \leftrightarrow , \leftrightarrow and \leftrightarrow with the same labeling of vertices as in the dihedral case. Similar calculations verify the remaining cases. Note hereby that for p big enough we can use [Theorem 3A.5](#) since the p -KL basis and the KL agree for big enough primes, and there are only finitely many cases left to check which we verified with computer help (one can verify this without computer, but we did it by computer). \square

Remark 3B.2. For finite Coxeter systems of Dynkin type calculations suggest that (H_W, B_W^p) is a sandwich pair if and only if W is of type A .

Remark 3B.3. The statements in [Theorem 3B.1](#) are reformulations of the vast literature on p -KL cells, see e.g. [\[JW17\]](#), [\[Jen20a\]](#) or [\[Jen20b\]](#), into the theory of sandwich cellular algebras.

3C. Kazhdan–Lusztig structure constants in dihedral type. Consider ∞ with nodes labeled 1 and 2. The Coxeter group associated to this Coxeter diagram is the infinite dihedral group $D_\infty = \langle 1, 2 \mid 1^2 = 2^2 = \emptyset \rangle$. Every element of D_∞ has a unique reduced expression and we write $k21$ and $k12$ for the reduced expressions $\dots 21$ and $\dots 12$ in k symbols.

The associated Hecke algebra H_{D_∞} has a KL basis $\{b_w \mid w \in D_\infty\}$ (whose precise definition does not matter) with identity b_\emptyset . Set $b_{0ab} = 0$. The nonidentity multiplication rules are given by the *(scaled) Clebsch–Gordan formula* where the steps size is two:

$$b_{k12}b_{j12} = \begin{cases} [2]b_{(|k-j|+1)12} + \dots + [2]b_{(|k+j|-1)12} & j12=2\dots 12, \\ b_{|k-j|12} + 2b_{(|k-j|+2)12} + \dots + 2b_{(|k+j|-2)12} + b_{|k+j|12} & j12=1\dots 12. \end{cases}$$

There are also similar formulas with b_{j21} on the right.

Let $D_n = \langle 1, 2 \mid 1^2 = 2^2 = (12)^n = \emptyset \rangle$ be the dihedral group of the n gon. The longest element is $w_0 = n, 1, 2 = n, 2, 1$. Let $[2]_i = (v^i + v^{-i})$. With respect to the KL basis and its multiplication rules, the only change compare to D_∞ is that expressions of the form (here $d > 0$)

$$b_{(n-d)12} + b_{(n+d)12} \mapsto [2]_d b_{w_0}, \quad b_{(n-d)21} + b_{(n+d)21} \mapsto [2]_d b_{w_0},$$

are replaced as indicated. This is the *(scaled) truncated Clebsch–Gordan formula*.

Example 3C.1. For the infinite dihedral group we get

$$\begin{aligned}
 b_{1212}b_{21212} &= [2]b_{12} + [2]b_{1212} + [2]b_{121212} + [2]b_{12121212}, \\
 b_{1212}b_{121212} &= b_{12} + 2b_{1212} + 2b_{121212} + 2b_{12121212} + b_{1212121212},
 \end{aligned}$$

from the Clebsch–Gordan formula. Moreover,

$$b_{1212}b_{21212} = [2]b_{12} + [2]b_{1212} + [2]b_{121212} + [2]b_{12121212} = [2]b_{12} + ([2]_3 + 2[2])b_{121212},$$

$$b_{1212}b_{121212} = b_{12} + 2b_{1212} + 2b_{121212} + 2b_{12121212} + b_{1212121212} = ([2]_4 + 2[2]_2 + 2[2]_0)b_{121212},$$

are calculations for $n = 6$. \diamond

The next lemma can be easily proven using the truncated Clebsch–Gordan formulas.

Lemma 3C.2. *For the pair (H_W, B_W) the cell structure for H_{I_n} is as in [Example 3A.4](#). \square*

Lemma 3C.3. *Let \mathbb{K} be an admissible field. For the involutive sandwich pair (H_W, B_W) in odd dihedral type $I_2(n)$ we have $\mathcal{H}_m \cong \mathbb{K}[\mathbb{Z}/(\frac{n-1}{2}\mathbb{Z})]$.*

Proof. We focus on the H -cell containing b_1 . For $n = 3$ we have $b_1^2 = [2]b_1$. Hence, $\mathcal{H}_m \cong \mathbb{K}$ and we will assume that $n > 3$.

For $n > 3$, the multiplication by the element $b_{121} \in \mathcal{H}_m$ satisfies $P_{(n-1)/2}(b_{121}) = 0$ where the polynomial $P_n(X)$ is given recursively by $P_0(X) = 1$, $P_1(X) = X$ and $P_{k+1}(X) = \frac{X-[2]}{[2]}P_k(X) - P_{k-1}(X)$ for $k > 1$. To see this one either uses the explicit formulas above. Or, alternatively, one observes that the Clebsch–Gordan formulas show that, up to scaling, \mathcal{H}_m is the Grothendieck algebra of $\mathrm{SO}_{2n}(3)$, the semisimplification of quantum $\mathfrak{so}_3(\mathbb{C})$ tilting modules for a primitive complex $2n$ th root of unity, with b_{121} corresponding to the generating object of $\mathrm{SO}_{2n}(3)$. (In other words, b_{121} corresponds to multiplication by [3].)

Hence, we have $\mathcal{H}_m \cong \mathbb{K}[X]/(P_{(n-1)/2}(X))$. When interpreting the quantum parameter as a generic complex number or as 1, the polynomial $P_k(X)$ has k distinct roots. Thus, $\mathcal{H}_m \cong \mathbb{K}[X]/(P_{(n-1)/2}(X))$ and the Chinese remainder theorem imply the lemma. \square

Lemma 3C.4. *Let \mathbb{K} be an admissible field. For the involutive sandwich pair (H_W, B_W) the ranks of the sandwich matrices are 1 for the bottom, 2 for all middle ones, and 1 for the top.*

The involved calculations are a bit ugly and, for the sake of brevity, we only sketch the proof:

Sketch of a proof of [Lemma 3C.4](#). The claim is immediate for the bottom and top J -cells. For \mathcal{J}_m we first compute the general pairing matrix. We scale such that we do not need to worry about [2] and write $c_w = \frac{1}{[2]}b_w$. Take $C = (C_1, \dots) = (c_1, c_{121}, \dots, c_{12}, c_{1212}, \dots)$ and $R = (R_1, \dots) = (c_1, c_{121}, \dots, c_{21}, c_{2121}, \dots)$ as index sets for columns and rows. Next, we write down a matrix M with i - j entry being $R_i C_j \pmod{\mathbb{K}\{b_{w_0}\}}$. For $n = 5$ we for example get

$$M = \begin{array}{c|c|c|c|c} & c_1 & c_{121} & c_{12} & c_{1212} \\ \hline c_1 & c_1 & c_{121} & c_{12} & c_{1212} \\ \hline c_{121} & c_{121} & c_1 + c_{121} & c_{12} + c_{1212} & c_{12} \\ \hline c_{21} & c_{21} & c_{21} + c_{2121} & c_2 + c_{212} & c_{212} \\ \hline c_{2121} & c_{2121} & c_{21} & c_{212} & c_2 \end{array}.$$

The isomorphism constructed in the proof of [Lemma 3C.3](#) implies that the matrix M , up to scaling, can be replaced by a matrix N with four blocks corresponding to $\mathbb{K}[\mathbb{Z}/(\frac{n-1}{2}\mathbb{Z})]$. For example, for $n = 5$ one gets

$$N = \begin{array}{c|c|c|c|c} & d_1 & d_{121} & d_{12} & d_{1212} \\ \hline d_1 & d_1 & d_{121} & d_{12} & d_{1212} \\ \hline d_{121} & d_{121} & d_1 & d_{1212} & d_{12} \\ \hline d_{21} & d_{21} & d_{2121} & d_2 & d_{212} \\ \hline d_{2121} & d_{2121} & d_{21} & d_{212} & d_2 \end{array}.$$

In this example, the units in the four blocks corresponding to $\mathbb{K}[\mathbb{Z}/2\mathbb{Z}]$ are d_1 , d_{1212} , d_{2121} and d_2 . In general the units are d_1 , $d_{(n-1)12}$, $d_{(n-1)21}$ and d_2 .

The idempotents for the simple $\mathbb{K}[\mathbb{Z}/(\frac{n-1}{2}\mathbb{Z})]$ -modules are, of course, easy to write down, and it is then also easy to show that the ranks of the associated sandwich matrices are $\mathrm{rank}_{\mathbb{K}}(S^{m,i}) = 2$. For example, for $n = 5$ the idempotents for the northwest corner are $e_1 = d_1 + d_{121}$ and $e_2 = d_1 - d_{121}$. Let $e_3 = d_{1212} + d_{12}$ be the idempotent for the southwest corner. For e_1 we get the matrix

$$Ne_1 = \begin{pmatrix} e_1 & e_1 & e_1 & e_1 \\ e_1 & e_1 & e_1 & e_1 \\ e_3 & e_3 & e_3 & e_3 \\ e_3 & e_3 & e_3 & e_3 \end{pmatrix}.$$

The matrix is clearly of rank two, which shows that the sandwich matrix for e_1 is also of rank two. The rank of the sandwich matrix for e_2 can be computed similarly. \square

Example 3C.5. With appropriate care one can use H -reduction [Theorem 2A.17](#) for arbitrary fields to classify simple D_n -modules for n odd as well as their quantum counterparts.

For example, take $n = 5$ and $v = 1$ and let us exclude $\text{char}(\mathbb{K}) = 2$. The polynomial from the proof of [Lemma 3C.3](#) defining the quotient is $P_2(X) = \frac{1}{2}X^2 - X - 1$. Substituting $X \mapsto \sqrt{2Y^2 - 2Y + 1} + 1$ gives $P_2(Y) = Y^2 - Y - 1$ whose roots are the golden ratio Φ , interpreted in \mathbb{K} , appearing in [\(3A.7\)](#) and its conjugate. (For the general case note that Φ is [2] specialized at $\exp(\pi i/n)$ with $n = 5$.) Thus, unless $\text{char}(\mathbb{K}) = 5$, we can apply H -reduction for the middle J -cell and obtain two simple D_5 -modules of that apex. For $\text{char}(\mathbb{K}) = 5$ one needs to distinguish the cases when $\mathbb{K} = \mathbb{F}_{25}$ and $\mathbb{K} = \mathbb{F}_5$, but one still gets the correct number of simple D_5 -modules with the middle apex.

In general, the recursion for the polynomial giving the character table of D_n spot on is $P'_0(X) = 1$, $P'_1(X) = X$ and $P'_{k+1}(X) = (X - 1)P'_k(X) - P'_{k-1}(X)$ for $k > 1$. \diamond

Remark 3C.6. As [Example 3C.5](#), it is not hard to discuss the case for general \mathbb{K} , but there is some annoying rescaling involved so we decide not to include this into this paper.

Remark 3C.7. The multiplication rules for the KL basis of the dihedral group are well-known. The above, in particular, is a reformulation of [[dC06](#), Section 4]. The application to dihedral representation theory via sandwich cellularity, e.g. [Lemma 3C.2](#), [Lemma 3C.3](#) and [Lemma 3C.4](#), is new.

4. DIAGRAM ALGEBRAS AND SANDWICH CELLULARITY

We now discuss several examples of diagram algebras, all of which fit into the theory of sandwich cellular algebras.

4A. Some non-involutive diagram algebras. Let $S_n = \text{Aut}(\{1, \dots, n\})$ be the *symmetric group* on the set $\{1, \dots, n\}$, and in this paper the *planar symmetric group* is $S_n^p = 1$, independent of n . We now discuss two examples, which in some sense are analogs of the (planar) symmetric group in the theory of monoids.

Definition 4A.1. The *transformation monoid* T_n on the set $\{1, \dots, n\}$ is $\text{End}(\{1, \dots, n\})$.

Note that the subgroup of invertible elements $G \subset T_n$ is isomorphic to S_n and we will identify S_n as a subgroup of T_n in this way.

The elements of T_n can be written in one-line notation with $(ijk\dots)$ denoting the map $1 \mapsto i$, $2 \mapsto j$, $3 \mapsto k$ etc. Alternatively we can model T_n using string diagrams as follows.

We consider isotopy classes of diagrams of $2n$ points in the rectangle $[0, 1] \times [0, 1]$, with n equally spaced points at the bottom and top. For $(ijk\dots)$, connect the first bottom point with the i th on the top, the second with the j th, the third with the k th and so on. Two diagrams represent the same element if and only if they represent the same map in T_n . For example:

$$(4A.2) \quad \begin{array}{l} (24138567) \rightsquigarrow \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array}, \quad (24637158) \rightsquigarrow \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}, \\ (23135555) \rightsquigarrow \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array}, \quad (11335577) \rightsquigarrow \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array}. \end{array}$$


The dots on strings used in the pictures are just reminders that there are top point not in the image of the displayed maps. The first two diagrams represent elements of $S_8 \subset T_8$.

Notation 4A.3. We have *crossings*, *merges* and *end dots*, which are:

$$\text{crossings: } \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}, \quad \text{merges: } \begin{array}{c} \diagup \diagdown \\ | \end{array}, \begin{array}{c} \diagdown \diagup \\ | \end{array}, \dots, \quad \text{end dots: } \downarrow.$$

We will use this terminology throughout this section.

The multiplication \circ in the transformation monoid is then given by vertical gluing (and rescaling), using the convention from [Notation 2A.1](#). For example,



Definition 4A.4. The *planar transformation monoid* T_n^p on the set $\{1, \dots, n\}$ is the submonoid of T_n for which we can draw planar string diagrams without leaving the defining rectangle $[0, 1] \times [0, 1]$.

Example 4A.5. From the four elements displayed in [\(4A.2\)](#) only the southeast is in T_8^p . In particular, $T_8^p \cap S_8 = \{id\}$, where id is the unit in T_8 . \diamond


Remark 4A.6. In the semigroup literature T_n is also called *full transformation monoid* and T_n^p is called *order-preserving transformation monoid*.

Using the diagrammatic description we find:

Lemma 4A.7. For $a \in T_n$ there is a unique factorization of the form $a = \tau \circ \sigma_\lambda \circ \beta$ such that β has a minimal number of crossings, τ has no crossings, β contains no end dots, τ contains no merges and $\sigma_\lambda \in S_\lambda$ for minimal λ .

Similarly for $a \in T_n^p$ but with $\sigma_\lambda \in S_n^p = 1$.

Proof. The rearrangement



generalizes immediately. \square

With respect to the factorization in [Lemma 4A.7](#), we call λ the number of *through strands*, β the *bottom*, τ the *top* and σ_λ the *middle* of a . [Lemma 4A.7](#) also motivates:

Definition 4A.8. We define the *symmetric* $\mathbb{K}T_n = \mathbb{K}[T_n]$ and the *planar diagram algebra* $\mathbb{K}T_n^p = \mathbb{K}[T_n^p]$ associated the (planar) transformation monoid.

Notation 4A.9. We write D_n for either T_n or T_n^p .

Let $\left\{ \begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right\}$ denote the (n, λ) th Stirling number (of the second kind), see also [Remark 4A.11](#) below.

Proposition 4A.10. We have the following for the pair $(\mathbb{K}D_n, D_n)$.

- (a) The J -cells of $\mathbb{K}D_n$ are given by diagrams with a fixed number of through strands λ . The \leq_{lr} -order is a total order and increases as the number of through strands decreases. That is,

$$\mathcal{P} = \{n <_{lr} n-1 <_{lr} \dots <_{lr} 1\}.$$

- (b) The left cells of $\mathbb{K}D_n$ are given by diagrams where one fixes the bottom of the diagram, and similarly right cells are given by diagrams where one fixes the top of the diagram. The \leq_l and the \leq_r -order increases as the number of through strands decreases. Within \mathcal{J}_λ we have

$$\mathbb{K}T_n: \#\mathcal{L}_\lambda = \left\{ \begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right\}, \quad \mathbb{K}T_n^p: \#\mathcal{L}_\lambda = \binom{n-1}{\lambda-1}, \quad \mathbb{K}D_n: \#\mathcal{R}_\lambda = \binom{n}{\lambda}.$$

- (c) Each J -cell of $\mathbb{K}D_n$ is strictly idempotent, and


$$\mathbb{K}T_n: \mathcal{H}_\lambda \cong \mathbb{K}[S_\lambda], \quad \mathbb{K}T_n^p: \mathcal{H}_\lambda \cong \mathbb{K}[S_\lambda^p] \cong \mathbb{K}.$$

- (d) The pair $(\mathbb{K}D_n, D_n)$ is a sandwich pair, that comes neither from a cellular nor an affine cellular algebra.

- (e) Let \mathbb{K} be a field with $\text{char}(\mathbb{K}) = 0$. For $\lambda \neq 1$ we have $\text{rank}_{\mathbb{K}}(G^\lambda) = \binom{n}{\lambda}$ for $\mathbb{K}T_n$ and $\text{rank}_{\mathbb{K}}(G^\lambda) = \binom{n-1}{\lambda-1}$ for $\mathbb{K}T_n^p$. We also have $\text{rank}_{\mathbb{K}}(G^1) = 1$ for both.

Note that the numerical data given in [Proposition 4A.10](#) is sufficient to determine the numbers and sizes of left, right, J and H -cells by [Lemma 2B.21](#).

Remark 4A.11. Very similar to binomials, the **Stirling numbers** $\left\{ \begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right\}$ count certain combinatorial set partitions, and thus appear very often for diagram algebras. Explicitly, $\left\{ \begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right\}$ counts the number of ways to partition a set of n labeled objects into λ nonempty unlabeled subsets. Not surprisingly, $\left\{ \begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right\}$ also admit a triangle description:



The recursion is $\left\{ \begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right\} = \lambda \cdot \left\{ \begin{smallmatrix} n-1 \\ \lambda \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n-1 \\ \lambda-1 \end{smallmatrix} \right\}$ with the starting conditions $\left\{ \begin{smallmatrix} 0 \\ 0 \end{smallmatrix} \right\} = 1$ and $\left\{ \begin{smallmatrix} j \\ 0 \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} 0 \\ j \end{smallmatrix} \right\} = 0$ for $j > 0$. For example, $90 = 3 \cdot 25 + 15$.

Proof of Proposition 4A.10. Statements (a), (b) and (c). Except the counts for the left and the right cells, this follows from [Lemma 4A.7](#) and the construction of an idempotent for all J -cells. The construction is: let e_λ be the idempotent that sends the first λ entries to 1 and fixes the remaining ones. For example, if $n = 5$ and $\lambda = 3$, then

$$e_\lambda = \begin{array}{c} \diagdown \quad | \quad | \\ \diagup \quad | \quad | \end{array}$$

Clearly, $e_\lambda \in \mathcal{J}_\lambda$ and $e_\lambda e_\lambda = e_\lambda$.

To count the number of right cells, observe that they are determined by the number of end dots placements at the top of the diagram, and thus, we get $\binom{n}{n-\lambda} = \binom{n}{\lambda}$ of these. Analogously, for left cells we need to count bottom diagrams. For $\mathbb{K}T_n$ this count is given by the Stirling numbers, by construction of the latter, see [Remark 4A.11](#). In the planar case, the bottom diagrams are counted by compositions of n into λ parts, and the number of such compositions is known to be $\binom{n-1}{\lambda-1}$. The claim follows.

Statement (d). From (a), (b) and (c), [Proposition 2D.1](#) and [Lemma 2B.17](#).

Statement (e). Recall from the above that each H -cell of $\mathbb{K}D_n$ is determined by fixing a bottom and top diagram. The idempotent H -cells in $\mathbb{K}D_n$ are easy to describe:

Claim A. An H -cell of $\mathbb{K}D_n$ is idempotent if and only if each bottom vertex that gets merged has an end dot at the top.

Proof of Claim A. This is just



and the claim follows.

Now, if $\lambda = 1$, then Statement (e) is immediate from the cell structure of $\mathbb{K}D_n$, so let $\lambda > 1$.



For $\mathbb{K}T_n^p$ we calculated the rank of a submatrix of G^λ to be $\binom{n-1}{\lambda-1}$. To this end, we use induction on $n + \lambda$. For small values the claim is easily verified, see also [Example 4A.14](#) below. Assume inductively that the claim holds. Decompose the diagrams with n strands and λ through strands by fixing the leftmost strand to be a through strand:

$$(4A.12) \quad \begin{array}{c} \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \boxed{\begin{smallmatrix} n-2 \\ \lambda \end{smallmatrix}} \end{array}, \quad \begin{array}{c} | \quad | \quad | \quad | \\ \boxed{\begin{smallmatrix} n-2 \\ \lambda-1 \end{smallmatrix}} \end{array}.$$


(We pull end dots into the boxes if necessary.) The diagrams in these boxes have $n - 1$ strands with λ and $\lambda - 1$ through strands, respectively. Claim A shows that if we put these together in a submatrix the ranks add. Thus, we get that a submatrix of rank $\binom{n-1}{\lambda-1} = \binom{n-2}{\lambda-1} + \binom{n-2}{\lambda-2}$ by induction.

For $\mathbb{K}T_n$ the argument is the same, using induction, diagrams as in (4A.12), Claim A and the formula $\binom{n}{\lambda} = \binom{n-1}{\lambda} + \binom{n-1}{\lambda-1}$. \square

Example 4A.13. For $n = 3$ we get




which display the cells of \mathbb{KT}_3 in diagrammatic notation and in one-line notation. For \mathbb{KT}_3^p one gets



(We have not illustrate the cell structure in one-line notation.) ◇

Note that there is redundant information in the cell pictures in [Example 4A.13](#). For example, the H -cells of \mathbb{KT}_n within \mathcal{J}_λ are always of size $\lambda!$, so we actually only need to remember which ones are idempotent. GAP's package Semigroups does exactly that:

Example 4A.14. The cell structures of \mathbb{KT}_n and \mathbb{KT}_n^p are of the form



To prove that $\Delta(\lambda)e_K$ is a simple $\mathbb{K}\mathbb{T}_n$ -module, we use induction on $n - \lambda$. If $n - \lambda = 0$, then we are in the bottom cell and the claim is clear. So let $n - \lambda = k$ and assume that we have proven the claim for $n - \lambda < k$. Define idempotents

$$e_{i,j} = \left| \begin{array}{c} i \\ \text{||||} \\ i \quad j \end{array} \right|, \quad 1 \leq i < j \leq n,$$

that merge the i th and j th strands and are the identity everywhere else. Note that the top of the diagrams for the $e_{i,j}$ have $n - 1$ strands.

For $x \in \Delta(\lambda)e_K$ suppose that $e_{i,j}x \neq 0$ for some $e_{i,j}$. Using the permutation $\sigma_{i,j}$ of the i th and j th strand, this implies that $y = e_{1,n}\sigma_{i,j}x \in e_{1,n}\Delta(\lambda)e_K$ is nonzero. Since $\mathbb{K}\mathbb{T}_{n-1} \cong e_{1,n}\mathbb{K}\mathbb{T}_ne_{1,n}$, we know by induction that $e_{1,n}\Delta(\lambda)e_K$ is simple as a $\mathbb{K}\mathbb{T}_{n-1}$ -module and we get that $\Delta(\lambda)e_K \subset \mathbb{K}\mathbb{T}_ne_{1,n}\Delta(\lambda)e_K \subset \mathbb{K}\mathbb{T}_nx$, which shows that $\Delta(\lambda)e_K$ is a simple $\mathbb{K}\mathbb{T}_n$ -module.

From (4A.18) it follows that at least one $e_{i,j}$ does not annihilate $\Delta(\lambda)e_K$: in the case where $K = K(\mu)$ is not the sign \mathcal{H}_λ -module the idempotent e_K has at least one nontrivial symmetrizer box at the top and hitting it will produce a nonzero diagram.

Case 2: K is the sign \mathcal{H}_λ -module. The case where K is the sign \mathcal{H}_λ -module is special and better to be analyzed by different means. Let e_λ be the idempotent as in the proof of Proposition 4A.10. One easily sees that $e_\lambda\mathbb{T}_ne_\lambda \cong \mathbb{S}_\lambda$. Let $M = \{(x_1, \dots, x_n) \in \mathbb{K}^n \mid x_1 + \dots + x_n = 0\}$ denote the $\mathbb{K}\mathbb{T}_n$ -module with action induced from the natural action of $\mathbb{K}\mathbb{T}_n$ on \mathbb{K}^n . The space M is a $e_\lambda\mathbb{T}_ne_\lambda \cong \mathbb{S}_\lambda$ -module by restriction and hence, we get a $e_\lambda\mathbb{K}\mathbb{T}_ne_\lambda \cong \mathbb{K}[\mathbb{S}_\lambda]$ -module $\bigwedge^{\lambda-1} M$. It is known that $\bigwedge^{\lambda-1} M$ is a simple $\mathbb{K}[\mathbb{S}_\lambda]$ -module, and thus, is also a simple $\mathbb{K}\mathbb{T}_n$ -module. Its dimension is $\binom{n-1}{\lambda-1}$ and $\bigwedge^{\lambda-1} M$ has apex λ , by construction. Since we already know all the other simple $\mathbb{K}\mathbb{T}_n$ -module with apex λ , and they are of bigger dimensions, we conclude that $L(\lambda, K) \cong \bigwedge^{\lambda-1} M$.

Statement (b). Directly from Proposition 2C.6 and Proposition 4A.10 \square

Remark 4A.19. Note that the dimensions of the simple \mathbb{S}_λ -modules that appear in Theorem 4A.17 can be computed using well-known formulas since we assume that $\text{char}(\mathbb{K}) = 0$. Thus, Theorem 4A.17 gives an explicit description of the dimensions of the simple $\mathbb{K}\mathbb{D}_n$ -modules.

In the proof of Theorem 4A.17 we could have alternatively computed the sandwich matrices as the following example shows.

Example 4A.20. Continuing Example 4A.16, and let \mathbb{K} be such that $\text{char}(\mathbb{K}) \nmid 3!$. Then the dimensions of simple $\mathbb{K}\mathbb{T}_3$ -modules are given by 1, 2, 1 for the bottom apex, 3, 2 for the middle apex and 1 for the top apex. The two sandwich matrices of ranks 3 and 2 for the middle apex are

$$S^{m, \text{triv}} = \begin{pmatrix} e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} & 0 & 0 \\ e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} & 0 & 0 \\ e_{\text{triv}} & e_{\text{triv}} & 0 & 0 & e_{\text{triv}} & e_{\text{triv}} \\ e_{\text{triv}} & e_{\text{triv}} & 0 & 0 & e_{\text{triv}} & e_{\text{triv}} \\ 0 & 0 & e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} \\ 0 & 0 & e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} & e_{\text{triv}} \end{pmatrix}, \quad S^{m, \text{sign}} = \begin{pmatrix} e_{\text{sign}} & -e_{\text{sign}} & e_{\text{sign}} & -e_{\text{sign}} & 0 & 0 \\ -e_{\text{sign}} & e_{\text{sign}} & -e_{\text{sign}} & e_{\text{sign}} & 0 & 0 \\ e_{\text{sign}} & -e_{\text{sign}} & 0 & 0 & -e_{\text{sign}} & e_{\text{sign}} \\ -e_{\text{sign}} & e_{\text{sign}} & 0 & 0 & e_{\text{sign}} & -e_{\text{sign}} \\ 0 & 0 & -e_{\text{sign}} & e_{\text{sign}} & -e_{\text{sign}} & e_{\text{sign}} \\ 0 & 0 & e_{\text{sign}} & -e_{\text{sign}} & e_{\text{sign}} & -e_{\text{sign}} \end{pmatrix},$$

where e_{triv} and e_{sign} are the idempotents for the trivial and the sign \mathcal{H}_2 -module, respectively.

For $\mathbb{K}\mathbb{T}_3^p$ there is actually no restriction on the field since the Gram matrices are

$$G^b = (1), \quad G^m = \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad G^t = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

and we get 1, 2, 1 as the simple dimensions. \diamond

Remark 4A.21. The quantum version of the above discussion simply replaces crossings by **over** and **undercrossings**:

$$\text{X} \rightsquigarrow \text{Y} \text{ or } \text{Z}.$$

In this case the sandwiched algebras are the Hecke algebras of type A as in Section 3. Otherwise there is no difference.

Remark 4A.22. The transformation monoid and its planar counterpart are around for Donkey's years, and so are their diagrammatic description. The diagrammatic incarnations appear, for example, in the formulation of categories of (planar) transformation, see e.g. [ER20, Figure 8].

The representation theory of $\mathbb{K}T_n$ was studied since the early days of monoid representation theory. Often relying on versions of [Theorem 2A.17](#), but in a less general setting with the focus on monoids. For example, the classification of simple $\mathbb{K}T_n$ -modules is given in [Ste16, Section 5.3], but in a different language and not using sandwich cellularity. That section also lists a few original references, starting with [HZ57].

We do not know any reference for the representation theory of the planar transformation monoid. This discussion appears to be new.

4B. Some involutive diagram algebras.

In the previous section we found the cells and parameterized the simple $\mathbb{K}D_n$ -modules, for D_n being T_n or T_n^p , using their diagrammatic incarnation. The same strategy works, by its very construction, for a wide range of diagram monoids and algebras.

We list now a few such algebras (the reader unfamiliar with these is referred to e.g. [HR05] for more details). These algebras have in common that their multiplication is defined via an underlying monoid. As before, *planar* means the submonoids having only planar diagrams of the same type, while the other listed diagrammatic descriptions are *symmetric*.

- The *partition monoid* Pa_n of all diagrams of partitions of a $2n$ -element set. The *planar partition monoid* Pa_n^p is the respective planar submonoid of Pa_n .

$$\begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \in \text{Pa}_n, \quad \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \in \text{Pa}_n^p.$$

- The *rook-Brauer monoid* RoBr_n consisting of all diagrams with components of size 1 or 2. The planar rook-Brauer monoid $\text{RoBr}_n^p = \text{Mo}_n$ is also called *Motzkin monoid*.

$$\begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \in \text{RoBr}_n, \quad \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \in \text{Mo}_n.$$

- The *Brauer monoid* Br_n consisting of all diagrams with components of size 2. The planar Brauer monoid $\text{Br}_n^p = \text{TL}_n$ is known as the *Temperley-Lieb monoid*.

$$\begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \in \text{Br}_n, \quad \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \in \text{TL}_n.$$

- The *rook monoid* or *symmetric inverse semigroup* Ro_n consisting of all diagrams with components of size 1 or 2, and all partitions have at most one component at the bottom and at most one at the top. The *planar rook monoid* Ro_n^p is the corresponding submonoid.

$$\begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \in \text{Ro}_n, \quad \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \in \text{Ro}_n^p.$$

- The *symmetric group* S_n consisting of all matchings with components of size 1. The *planar symmetric group* is trivial $S_n^p \cong 1$.

$$\begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \in S_n, \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \in S_n^p.$$

The monoids S_n and S_n^p are, of course, the same groups as in [Section 4A](#). S_n is isomorphic to a subgroup of any of the symmetric monoids, and S_n^p is isomorphic to a subgroup of any of the planar monoids, both in the evident way. We will use this below.

Notation 4B.1. Below we write D_n for any of the monoids listed above.

Notation 4B.2. Additionally to [Notation 4A.3](#) we now also have


$$\text{caps: } \begin{array}{c} \text{---} \\ \text{---} \end{array}, \quad \text{cups: } \begin{array}{c} \text{---} \\ \text{---} \end{array}, \quad \text{splits: } \begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array}, \dots, \quad \text{start dots: } \begin{array}{c} \text{---} \\ \text{---} \end{array},$$

which we give the names *caps*, *cups*, *splits* and *start dots*.

Definition 4B.3. Fix $\delta \in \mathbb{K}$. For D_n we define its associated (symmetric or planar) *diagram algebra* $D_n(\delta)$ to be the algebra with basis D_n and multiplication of basis elements given by the monoid multiplication except that all closed components are evaluated to δ .

Note that $D_n(\delta)$ is \mathbb{K} -linear, but we decided to use $D_n(\delta)$ as the notation instead of e.g. $\mathbb{K}[D_n](\delta)$.

Example 4B.4. Typical examples how the multiplication in diagram algebras works is



which illustrate the multiplication of $Mo_6(\delta)$. ◇

Remark 4B.5. For some of the D_n one can define associated multiparameter diagram algebras. For example, for Mo_n one could evaluate circles to δ_1 and intervals to δ_2 . Our discussion below works mutatis mutandis for these multiparameter diagram algebras as well.

A main difference between D_n and T_n , T_n^p is that D_n is involutive using the **diagrammatic antiinvolution** $-^*$, e.g.:

$$\left(\begin{array}{c} \text{Diagram with arcs and dots} \end{array} \right)^* = \begin{array}{c} \text{Diagram with arcs and dots} \end{array}.$$

The \mathbb{K} -linear extensions of the diagrammatic antiinvolution endows $D_n(\delta)$ with the structure of an involutive sandwich cellular algebra, as we will see in [Proposition 4B.9](#) below. (We will always use the diagrammatic antiinvolution in this section.) Hence, by [Lemma 2B.17](#) the J -cells are squares and we can focus on describing left cells and right cells come for free.

Example 4B.6. Let us give some examples how the cells structure of $TL_n(\delta)$ looks like.

	\mathcal{J}_1		$\mathcal{H}_1 \cong 1$,
	\mathcal{J}_3		$\mathcal{H}_3 \cong 1$	
(4B.7)	\mathcal{J}_0		$\mathcal{H}_0 \cong 1$	
	\mathcal{J}_2		$\mathcal{H}_2 \cong 1$.
	\mathcal{J}_4		$\mathcal{H}_4 \cong 1$	

These are the cells of $TL_3(\delta)$ and $TL_4(\delta)$ for invertible δ . If δ is not invertible, then the picture is the same but with uncolored diagonal H -cells for \mathcal{J}_1 and \mathcal{J}_2 , and \mathcal{J}_0 having no idempotent at all. ◇

The following analog of [Lemma 4A.7](#) is easy to verify.

Lemma 4B.8. *Let D_n be symmetric. For $a \in D_n$ there is a unique factorization of the form $a = \tau \circ \sigma_\lambda \circ \beta$ such that β and τ have a minimal number of crossings, β contains no cups, splits or end dots, τ contains no caps, merges or start dots and $\sigma_\lambda \in S_\lambda$ for minimal λ .*

Similarly for $a \in D_n$ when D_n is planar, but with $\sigma_\lambda \in S_n^p = 1$. □

As before, we get through strands etc.

Proposition 4B.9. *We have the following for the pair $(D_n(\delta), D_n)$.*

- (a) *The J -cells of $D_n(\delta)$ are given by diagrams with a fixed number of through strands λ . The \leq_l -order is a total order and increases as the number of through strands decreases. See [\(4B.10\)](#) for a summary.*
- (b) *The left cells of $D_n(\delta)$ are given by diagrams where one fixes the bottom of the diagram, and similarly right cells are given by diagrams where one fixes the top of the diagram. The \leq_l and the \leq_r -order increases as the number of through strands decreases. For $\#\mathcal{L}_\lambda$ see [\(4B.10\)](#).*

(c) The idempotency of the J -cells are as follows. Assume δ is invertible in \mathbb{K} . Then all J -cells of $D_n(\delta)$ are strictly idempotent. Second, assume δ is not invertible in \mathbb{K} . Then the following J -cells for $D_n(\delta)$ are strictly idempotent, while all other J -cells are not idempotent (we list the various D_n):

(i) All J -cells except the top for Pa_n , Pa_n^p , RoBr_n and Mo_n , and Br_n , TL_n for even n .

(ii) Only the bottom J -cell for Ro_n and Ro_n^p .

(iii) All J -cells for S_n and S_n^p , and Br_n , TL_n for n odd.

See also (4B.10). The sandwiched algebras are given in (4B.10).


(d) The pair $(D_n(\delta), D_n)$ is an involutive sandwich pair, that for D_n symmetric comes neither from a cellular nor an affine cellular algebra, but can be refined into a cellular pair.

The following table summarizes the cell structure of these diagram monoids where $\lambda \in \mathcal{P}$ (the column for \mathcal{H}_λ only applies if the parent J -cell is strictly idempotent):

Monoid	$\mathcal{P} = \mathcal{P}^{ap}$ for $\delta^{-1} \in \mathbb{K}$	\mathcal{P}^{ap} for δ not inv.	$\#\mathcal{L}_\lambda$	\mathcal{H}_λ
Pa_n	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 0\}$	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 1\}$	$\sum_{t=0}^n \binom{n}{t} \binom{t}{\lambda}$	$\cong S_\lambda$
Pa_n^p	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 0\}$	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 1\}$	$\frac{4\lambda+2}{2n+2\lambda+2} \binom{2n}{(2n-2\lambda)/2}$	$\cong 1$
RoBr_n	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 0\}$	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 1\}$	$\sum_{t=0}^n \binom{n}{\lambda} \binom{n-\lambda}{2t} (2t-1)!!$	$\cong S_\lambda$
Mo_n	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 0\}$	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 1\}$	$\sum_{t=0}^n \frac{\lambda+1}{\lambda+t+1} \binom{n}{\lambda+2t} \binom{\lambda+2t}{t}$	$\cong 1$
Br_n	$\{n <_{lr} n-2 <_{lr} \dots <_{lr} 0, 1\}$	$\{n <_{lr} n-2 <_{lr} \dots <_{lr} 2, 1\}$	$\binom{n}{\lambda} (n-\lambda-1)!!$	$\cong S_\lambda$
TL_n	$\{n <_{lr} n-2 <_{lr} \dots <_{lr} 0, 1\}$	$\{n <_{lr} n-2 <_{lr} \dots <_{lr} 2, 1\}$	$\frac{2\lambda+2}{n+\lambda+2} \binom{n}{(n-\lambda)/2}$	$\cong 1$
Ro_n	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 0\}$	$\{n\}$	$\binom{n}{\lambda}$	$\cong S_\lambda$
Ro_n^p	$\{n <_{lr} n-1 <_{lr} \dots <_{lr} 0\}$	$\{n\}$	$\binom{n}{\lambda}$	$\cong 1$
S_n	$\{n\}$	$\{n\}$	1	$\cong S_\lambda$
S_n^p	$\{n\}$	$\{n\}$	1	$\cong 1$

For TL_n and Br_n the last entry of \mathcal{P} is either 0 or 1, depending on the parity of n . Similarly, the last entry of \mathcal{P}^{ap} is either 2 or 1, again depending on the parity of n .

Example 4B.11. GAP produces the following outputs for the underlying monoids:



The monoids S_n and S_n^p are not illustrated as they just have one cell. ◇

Proof of Proposition 4B.9. Statements (a) and (b). Only the counts for $\#\mathcal{L}_\lambda$ are not immediate. Counting them is a combinatorial exercise that has been solved several times in the literature. The

respective triangles, cf. [Remark 4A.11](#), are A049020, A008313, A096713, A064189, A111062, and A007318 on OEIS.

Statement (c). Easy to see and omitted. The main trick is to use variations of

$$\begin{array}{c} \text{S} \\ \text{=} \\ | \end{array}$$

to ensure that certain cells stay strictly idempotent even when δ is not invertible.

Statement (d). Since D_n is an admissible monoid by the above, [Proposition 2D.1](#) applies for $(\mathbb{K}[D_n], D_n)$. This result can be pulled over to $(D_n(\delta), D_n)$. The second claim about cellularity follows then from [Lemma 2B.17](#) and [Proposition 2A.10](#) (and some care with the antiinvolution). \square

From now on we use that $(D_n(\delta), D_n)$ is an involutive sandwich pair.


Theorem 4B.12. *Assume that \mathbb{K} is a field, and consider the involutive sandwich pair $(D_n(\delta), D_n)$.*

- (a) *Assume that $\delta \neq 0$. The set of apexes for simple $D_n(\delta)$ -modules is \mathcal{P}^{ap} as in [\(4B.10\)](#), and there are precisely $|\mathcal{P}(\lambda|p)|$ (symmetric) or one (planar) simple $D_n(\delta)$ -modules for $\lambda \in \mathcal{P}^{ap}$.*
- (b) *Assume that $\delta = 0$. Then the same statement holds for the restricted set of apexes as detailed in [\(4B.10\)](#).*


Proof. Clear by [Proposition 4B.9](#) and H -reduction [Theorem 2A.17](#). \square

Below we compute a few Gram matrices. To get started, here is an example:

Example 4B.13. For $TL_5(1)$ we have



$$G^1 = \begin{pmatrix} \delta^2 & \delta & 1 & \delta & 1 \\ \delta & \delta^2 & \delta & 1 & \delta \\ 1 & \delta & \delta^2 & \delta & 1 \\ \delta & 1 & \delta & \delta^2 & \delta \\ 1 & \delta & 1 & \delta & \delta^2 \end{pmatrix}, \quad \det(G^1) = (\delta - 1)^4(\delta + 1)^4(\delta^2 - 2),$$



$$G^3 = \begin{pmatrix} \delta & 1 & 0 & 0 \\ 1 & \delta & 1 & 0 \\ 0 & 1 & \delta & 1 \\ 0 & 0 & 1 & \delta \end{pmatrix}, \quad \det(G^3) = (\delta^2 + \delta - 1)(\delta^2 - \delta - 1),$$



$$G^5 = (1), \quad \text{rank}_{\mathbb{K}}(G^5) = 1,$$

as cells, Gram matrices and their determinants. The ranks depend on δ with the exception of the bottom cell. \diamond

The following proposition computes ranks of Gram matrices for $D_n \in \{S_n, S_n^p, Ro_n, Ro_n^p, TL_n, Pa_n^p\}$. We do not know nice formulas for the ranks of the remaining monoids, but we give some partial results by computing the ranks of the cells close to the bottom.

Proposition 4B.14. *Assume that \mathbb{K} is a field, and consider the involutive sandwich pair $(D_n(\delta), D_n)$. The ranks of (some of the) Gram matrices are as follows.*

- (a) *For $D_n \in \{S_n, S_n^p\}$ we have $\text{rank}_{\mathbb{K}}(G^n) = 1$ for $n \in \mathcal{P}^{ap}$.*
- (b) *For $D_n \in \{Ro_n, Ro_n^p\}$ we have $\text{rank}_{\mathbb{K}}(G^\lambda) = \binom{n}{\lambda}$ for $\lambda \in \mathcal{P}^{ap}$.*
- (c) *For $D_n = TL_n$ we have $\text{rank}_{\mathbb{K}}(G^\lambda) = tl_n^\lambda$ for $\lambda \in \mathcal{P}^{ap}$, with tl_n^λ explicitly given in [Remark 4B.15](#) below. For $D_n = Pa_n^p$ we have $\text{rank}_{\mathbb{K}}(G^\lambda) = tl_{2n}^{2\lambda}$ for $\lambda \in \mathcal{P}^{ap}$.*
- (d) *For $D_n \in \{Br_n, RoBr_n, Mo_n, Pa_n\}$ we have $\text{rank}_{\mathbb{K}}(G^n) = 1$ for $n \in \mathcal{P}^{ap}$.*
- (e) *For $D_n = Br_n$, $\text{char}(\mathbb{K}) = 0$ and $n - 2 \in \mathcal{P}^{ap}$ we have*

$$\text{rank}_{\mathbb{K}}(G^{n-2}) = \begin{cases} \frac{1}{2}n(n-1) & \text{if } \delta \notin \{2, -n+4, -2n+4\}, \\ n & \text{if } \delta = 2, \\ \frac{1}{2}n(n-3) + 1 & \text{if } \delta = -n+4, \\ \frac{1}{2}(n+1)(n-2) & \text{if } \delta = -2n+4. \end{cases}$$

isomorphism $\text{Pa}_n^p \cong \text{TL}_{2n}$ from [HR05, Section 1] can then be used to prove the statement for the planar partition algebra from the Temperley–Lieb algebra.

Case $\text{D}_n \in \{\text{Br}_n, \text{Mo}_n, \text{RoBr}_n, \text{Pa}_n\}$, bottom cell. Clear.

Case Br_n , $\lambda = n - 2$. In this case there is one cap and one cup. Thus, it suffices to remember the endpoints of the middle. We do this by using

$$(4B.17) \quad \begin{array}{c} \text{---} \\ | \\ | \\ | \\ \text{---} \\ i \qquad j \end{array} \leftrightarrow b[i, j], 1 \leq i < j \leq n, \quad \left| \begin{array}{c} k \qquad l \\ \text{---} \\ | \\ | \\ | \\ \text{---} \end{array} \right. \leftrightarrow t[k, l], 1 \leq k < l \leq n.$$

The numbers in these pictures are the endpoints of the strings, read from left to right. The Gram matrix becomes symbolically $\mathbf{G}^{n-2} = (b[i, j]t[k, l])_{i, j, k, l}$ for $1 \leq i < j \leq n$ and $1 \leq k < l \leq n$. The entries of the Gram matrix are then determined by

$$(4B.18) \quad \begin{cases} b[i, j]t[k, l] = \delta & \text{if both endpoints match,} \\ b[i, j]t[k, l] = 1 & \text{if one endpoint matches,} \\ b[i, j]t[k, l] = 0 & \text{else,} \end{cases} \quad \begin{array}{c} 2 \qquad 4 \\ \text{---} \\ | \\ | \\ | \\ \text{---} \\ 1 \qquad 4 \end{array} \circ \begin{array}{c} 2 \qquad 4 \\ \text{---} \\ | \\ | \\ | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \\ | \\ | \\ \text{---} \end{array} \rightsquigarrow 1.$$

This can be seen as indicated above. The matrices are then easy to write down, for example

$$n = 4: \mathbf{G}^{n-2} = \begin{pmatrix} \delta & 1 & 1 & 1 & 1 & 0 \\ 1 & \delta & 1 & 1 & 0 & 1 \\ 1 & 1 & \delta & 0 & 1 & 1 \\ 1 & 1 & 0 & \delta & 1 & 1 \\ 1 & 0 & 1 & 1 & \delta & 1 \\ 0 & 1 & 1 & 1 & 1 & \delta \end{pmatrix}.$$

Induction verifies that the determinant is

$$\det(\mathbf{G}^{n-2}) = (\delta - 2)^{\frac{1}{2}n(n-3)}(\delta + n - 4)^{n-1}(\delta + 2n - 4).$$

Thus, unless $\delta \in \{2, -n + 4, -2n + 4\}$, we get $\text{rk}_{\mathbb{K}}(\mathbf{G}^{n-2}) = \frac{1}{2}n(n-1)$. For the remaining cases we get $\text{rk}_{\mathbb{K}}(\mathbf{G}^{n-2}) = n$ if $\delta = 2$, $\text{rk}_{\mathbb{K}}(\mathbf{G}^{n-2}) = \frac{1}{2}n(n-3) + 1$ if $\delta = -n + 4$ and $\text{rk}_{\mathbb{K}}(\mathbf{G}^{n-2}) = \frac{1}{2}(n+1)(n-2)$ if $\delta = -2n + 4$.

Case RoBr_n and Mo_n , $\lambda = n - 1$. The only way to reduce the number of through strands by one is to have a start and an end dot. Thus, this cell is exactly as in (4B.16), up to permutations, and we get the same Gram matrix as for Ro_n and Ro_n^p , namely the δ times identity.

Case Mo_n , $\lambda = n - 2$. In order to get $n - 2$ through strands one either needs to have one cap or two start dots at the bottom, and one cup or two end dots at the top of the diagram. Having one cap-cup pair is the Temperley–Lieb case, having two start and two end dots is the planar rook monoid case, and then there are the mixed cases. In the following illustration we again mark the endpoints of strings by their positions read from left to right. As for Br_n above, that is (4B.17), we use a translation from diagrams to symbols:

$$\begin{array}{c} \text{---} \\ \text{---} \\ i \quad i+1 \end{array} \leftrightarrow b[i, i+1], 1 \leq i \leq n, \quad \begin{array}{c} \bullet \quad \bullet \\ | \quad | \\ i \quad i+1 \end{array} \leftrightarrow c[i, i+1], 1 \leq i \leq n, \quad \begin{array}{c} \bullet \quad | \quad \bullet \\ | \quad | \\ j \quad k \end{array} \leftrightarrow d[j, k], 1 \leq j < k \leq n.$$

We use the notation $t[i, i+1]$, $u[i, i+1]$ and $v[j, k]$ for right cells. Additionally to (4B.18) we have:

$$\begin{cases} b[i, i+1]u[j, j+1] = \delta & \text{if } i = j, \\ b[i, i+1]u[j, j+1] = 0 & \text{else,} \\ b[i, i+1]v[j, k] = 0 & \text{always,} \\ u[i, i+1]u[j, j+1] = \delta^2 & \text{if } i = j, \end{cases} \quad \begin{cases} u[i, i+1]u[j, j+1] = 0 & \text{else,} \\ u[i, i+1]v[j, k] = 0 & \text{always,} \\ v[i, j]v[k, l] = \delta^2 & \text{if both endpoints match,} \\ v[i, j]v[k, l] = 0 & \text{else.} \end{cases}$$

Using these formulas and the corresponding symbolic Gram matrix, an analysis as for Br_n shows that the Gram matrix of square size $(n-1) + (n-1) + (\frac{1}{2}(n-1)(n-2))$ is

$$(4B.19) \quad \mathbf{G}^{n-2} = \left(\begin{array}{c|c|c} A_{11} & A_{12} & 0 \\ \hline A_{12} & A_{22} & 0 \\ \hline 0 & 0 & A_{33} \end{array} \right),$$

$$A_{11} = \begin{pmatrix} \delta & 1 & 0 & 0 & 0 & \dots & 0 \\ 1 & \delta & 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & \delta & 1 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \end{pmatrix}, \quad A_{12} = \delta id, \quad A_{22} = \delta^2 id, \quad A_{33} = \delta^2 id.$$

The northwest corner is the same as for TL_n . An example is

$$n = 4: G^{n-2} = \begin{pmatrix} \delta & 1 & 0 & \delta & \delta & 0 & 0 & 0 & 0 & 0 \\ 1 & \delta & 1 & 0 & \delta & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & \delta & 0 & 0 & \delta & 0 & 0 & 0 & 0 \\ \delta & 0 & 0 & \delta^2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \delta & 0 & 0 & \delta^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \delta & 0 & 0 & \delta^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \delta^2 \end{pmatrix}.$$

Let $U_k(X)$ again be the (normalized) Chebyshev polynomial of the second kind which is defined by the recursion $U_{k+1}(X) = XU_k(X) - U_{k-1}(X)$ together with $U_0(X) = 1$ and $U_1(X) = X$, see [Remark 4B.15](#). It is not hard to see (and well-known) that $\det(A_{11}) = U_{n-1}(\delta)$. Using this determinant formula and [\(4B.19\)](#) we get

$$\det(G^{n-2}) = \delta^{n(n-1)} U_{n-1}(\delta - 1).$$

The Chebyshev polynomial $U_{n-1}(X - 1)$ is a polynomial of degree $n - 1$ and has distinct roots (given by explicit formulas) and we denote the set of these roots by Roots_{n-1} . Thus, for $\delta \notin \{0\} \cup \text{Roots}_{n-1}$, we get $\text{rk}_{\mathbb{K}}(G^{n-2}) = \frac{1}{2}(n^2 + n - 2)$. If we have $\delta \neq 0$ and $\delta \in \text{Roots}_{n-1}$, then we get $\text{rk}_{\mathbb{K}}(G^{n-2}) = \frac{1}{2}(n^2 + n - 4)$. In the final case $\delta = 0$ we recall that $0 \in \text{Roots}_{n-1}$ holds if and only if $n \equiv 0 \pmod{2}$. Thus, we get $\text{rk}_{\mathbb{K}}(G^{n-2}) = n - 1$ if $n \not\equiv 0 \pmod{2}$, and $\text{rk}_{\mathbb{K}}(G^{n-2}) = n - 2$ if $n \equiv 0 \pmod{2}$.

Case RoBr_n , $\lambda = n - 2$. This case is very similar to Br_n and Mo_n for $\lambda = n - 2$ with some small differences. That is, we still use the shorthand notation from [\(4B.17\)](#) but now also

$$(4B.20) \quad \begin{array}{c} \uparrow \\ i \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{c} \uparrow \\ j \end{array} \longleftrightarrow c[i, j], \quad \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{c} \uparrow \\ k \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{c} \uparrow \\ l \end{array} \longleftrightarrow u[k, l].$$

As one easily checks by drawing the relevant pictures, we then get

$$\begin{cases} b[i, j]t[k, l] = \delta & \text{if both endpoints match,} \\ b[i, j]t[k, l] = 1 & \text{if one endpoint matches,} \\ b[i, j]t[k, l] = 0 & \text{else,} \\ c[i, j]u[k, l] = \delta^2 & \text{if both endpoints match,} \\ c[i, j]u[k, l] = 0 & \text{else,} \end{cases} \quad \begin{cases} b[i, j]u[k, l] = \delta & \text{if both endpoints match,} \\ b[i, j]u[k, l] = 0 & \text{else,} \\ c[i, j]t[k, l] = \delta & \text{if both endpoints match,} \\ c[i, j]t[k, l] = 0 & \text{else,} \end{cases}$$

as entries of the Gram matrix. Thus, we have the same block decomposition as in [\(4B.19\)](#) but with northwest corner corresponding to Br_n and A_{33} being empty. For example,

$$n = 4: G^{n-2} = \begin{pmatrix} \delta & 1 & 1 & 1 & 0 & \delta & \delta & 0 & 0 & 0 & 0 & 0 \\ 1 & \delta & 1 & 1 & 0 & 1 & 0 & \delta & 0 & 0 & 0 & 0 \\ 1 & 1 & \delta & 0 & 1 & 1 & 0 & 0 & \delta & 0 & 0 & 0 \\ 1 & 1 & 0 & \delta & 1 & 1 & 0 & 0 & 0 & \delta & 0 & 0 \\ 1 & 0 & 1 & 1 & \delta & 1 & 0 & 0 & 0 & 0 & \delta & 0 \\ 0 & 1 & 1 & 1 & 1 & \delta & 0 & 0 & 0 & 0 & 0 & \delta \\ \delta & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & \delta & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & \delta & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & \delta & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & \delta & 0 & 0 & 0 & 0 & 0 & \delta^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & \delta & 0 & 0 & 0 & 0 & 0 & \delta^2 \end{pmatrix}.$$

From the previous formulas one easily gets that

$$\det(G^{n-2}) = \delta^{\frac{1}{2}n(n-1)} (\delta - 3)^{\frac{1}{2}n(n-3)} (\delta + n - 5)^{n-1} (\delta + 2n - 5).$$

Analyzing the rank using this determinant formula gives the claimed result. That is, unless $\delta \in \{0, 3, 5 - n, 5 - 2n\}$ we have $\text{rk}_{\mathbb{K}}(G^{n-2}) = n(n - 1)$. Otherwise, reading $\{0, 3, 5 - n, 5 - 2n\}$ left to right, we get $\text{rk}_{\mathbb{K}}(G^{n-2}) = \frac{1}{2}n(n - 1)$, $\text{rk}_{\mathbb{K}}(G^{n-2}) = \frac{1}{2}n(n + 1)$, $\text{rk}_{\mathbb{K}}(G^{n-2}) = (n - 1)^2$ and $\text{rk}_{\mathbb{K}}(G^{n-2}) = n(n - 1) - 1$. \square

We now compute the dimensions of the simple D_n -modules for $D_n \in \{S_n, S_n^p, \text{Ro}_n, \text{Ro}_n^p, \text{TL}_n, \text{Pa}_n^p\}$, and give some partial results for the remaining ones.

Theorem 4B.21. *Assume that \mathbb{K} is a field with $\text{char}(\mathbb{K}) = 0$ for the symmetric monoids, and consider the involutive sandwich pair $(D_n(\delta), D_n)$.*

- (a) *For $D_n \in \{S_n, S_n^p, \text{Ro}_n, \text{Ro}_n^p, \text{TL}_n, \text{Pa}_n^p\}$ the dimension of the simple $D_n(\delta)$ -modules for $\lambda \in \mathcal{P}^{ap}$, and additionally a simple \mathcal{H}_λ -module K for S_n and Ro_n , are*

$$\dim_{\mathbb{K}}(L(\lambda)) = \text{rank}_{\mathbb{K}}(G^\lambda), \quad \dim_{\mathbb{K}}(L(\lambda, K)) = \text{rank}_{\mathbb{K}}(G^\lambda) \cdot \dim_{\mathbb{K}}(K).$$

(The ranks of the Gram matrices are computed in Proposition 4B.14.)

- (b) *For $D_n \in \{S_n, S_n^p, \text{Ro}_n, \text{Ro}_n^p, \text{TL}_n, \text{Pa}_n^p\}$ the algebra $D_n(\delta)$ is semisimple if and only if we are in the following cases:*

(i) *All cases for $D_n \in \{S_n, S_n^p\}$.*

(ii) *The parameter δ is nonzero for $D_n \in \{\text{Ro}_n, \text{Ro}_n^p\}$.*

(iii) *The number l from Remark 4B.15 is $l = \infty$ for $D_n \in \{\text{TL}_n, \text{Pa}_n^p\}$.*

- (c) *For $\lambda \in \mathcal{P}^{ap}$ we have the following lower bounds, where we always want to bound the left side by the right side:*

$$\dim_{\mathbb{K}}(L(K_{triv})_{\text{Pa}_n}) \geq tl_{2n}^{2\lambda}, \quad \dim_{\mathbb{K}}(L(K_{triv})_{\text{RoBr}_n}) \geq \dim_{\mathbb{K}}(L(\lambda)_{\text{Mo}_n}) \geq tl_n^\lambda, \quad \dim_{\mathbb{K}}(L(K_{triv})_{\text{Br}_n}) \geq tl_n^\lambda.$$

Here K_{triv} always denotes the trivial \mathcal{H}_λ -module and we use subscripts to indicate what kind of simple $D_n(\delta)$ -modules we consider.

Proof. Statement (a). For the planar diagram algebras this is just Proposition 2C.6, and for $D_n = S_n$ the statement is clear. For $\text{Ro}_n(\delta)$ and $\lambda \in \mathcal{P}^{ap}$ we define a pseudo-idempotent e_λ that have a start-end dot pair for all positions 1 to $n - \lambda$ and is the identity otherwise. For example, for $n = 5$ and $\lambda = 3$ we get

$$e_3 = \begin{array}{c} \bullet \bullet \bullet \\ \bullet \bullet \bullet \\ \bullet \bullet \bullet \\ | \\ | \end{array}.$$

Hence, we get idempotents $e'_\lambda = \frac{1}{\delta^{n-\lambda}} e_\lambda$ in \mathcal{J}_λ . Note also that $\Delta(\lambda) \cong \text{Ro}_n(\delta)e_\lambda$ as $\text{Ro}_n(\delta)$ -modules. Let $e_{\sigma(\lambda)}$ denote the projection to the component of $\mathcal{H}_\lambda^{\oplus k}$ determined by the top through strand points of σe_λ , where we write $k = \binom{n}{\lambda}$. The \mathbb{K} -linear map $f_\lambda: \Delta(\lambda) \rightarrow \mathcal{H}_\lambda^{\oplus k}$ given by $f_\lambda(\sigma e'_\lambda) = \sigma e_{\sigma(\lambda)}$ is an isomorphism of \mathcal{H}_λ -modules. Thus, $\Delta(\lambda, K) \cong K^{\oplus k}$ for all simple \mathcal{H}_λ -modules K . Using now the permutation action of \mathcal{H}_n , which connects the various direct summands, one sees that $\Delta(\lambda, K)$ is a simple $\text{Ro}_n(\delta)$ -module.

Statement (b). By (a) and Proposition 2B.23.

Statement (c). This follows because we have algebra embeddings $\text{Pa}_n^p(\delta) \hookrightarrow \text{Pa}_n(\delta)$, $\text{TL}_n(\delta) \hookrightarrow \text{Mo}_n(\delta) \hookrightarrow \text{RoBr}_n(\delta)$ and $\text{TL}_n(\delta) \hookrightarrow \text{Br}_n(\delta)$ given by sending diagrams to themselves interpreted in the bigger monoids. Thus, using an appropriate order, the sandwich matrices for the small monoids are submatrices of the sandwich matrices for the bigger monoids and Proposition 4B.14 completes the proof. \square

Remark 4B.22. Similarly as in Remark 4A.21, the quantum versions of the diagram algebras $D_n(\delta)$ (e.g. BMW algebras) can be studied verbatim, and the Hecke algebras of type A appear again as the sandwiched algebras.

Remark 4B.23. The study of diagram algebras has a long history, which we can not cover here in any satisfying way. Nevertheless, let us give a few references for some of the results above which have appeared in the literature, but not under the umbrella of sandwich cellularity, which is new in this paper.

The representation theory of S_n and S_n^p is, of course, well-studied, and we do not comment on these any further.

Statements such as Proposition 4B.9 and Proposition 4B.14 appear throughout the semigroup literature, see e.g. [HR05], [DEE⁺19], [HJ20], sometimes as disguised lemmas such as [EMRT18, Remark 9.21 and Lemma 9.19], and many more. They have very similar appearances for diagram algebras with the catch that the parameter δ is not necessarily 1 as for the monoids.

Computations of dimensions of simple modules of diagram algebras such as in Theorem 4B.21 have also appeared in many works (although some of the results above appear to be new). For example, for Ro_n see [Mun57] and [Sol02], and for Ro_n^p see [FHH09]. The formulas from Remark 4B.15 for TL_n go back to [RSA14], [And19] and [Spe20], and the ones for Pa_n^p follows then from the isomorphism

to TL_{2n} from [HR05]. We do not know any reference for the dimensions of simple modules for any of the remaining symmetric monoids. The bounds however have been studied in [KST22]. Let us also note that a lot of these diagram algebras and similar diagram algebra can be studied using Schur–Weyl–Brauer duality, see e.g. [AST17, Section 3] for a collection of these dualities.

The quantum versions, which also fit into the theory, cf. Remark 4B.22, have also been studied extensively, see e.g. [Eny04] for the BMW case. Note also that Proposition 4B.9.(d) shows that all the diagram algebras in this section are cellular. This is known, see e.g. [GL96], [Xi99] and [Eny04]. But again our point is that we get all of this by using the theory of sandwich cellular algebras.

REFERENCES

- [AM98] A.K. Aiston and H.R. Morton. Idempotents of Hecke algebras of type A . *J. Knot Theory Ramifications*, 7(4):463–487, 1998. URL: <https://arxiv.org/abs/q-alg/9702017>, doi:10.1142/S0218216598000243.
- [And19] H.H. Andersen. Simple modules for Temperley–Lieb algebras and related algebras. *J. Algebra*, 520:276–308, 2019. URL: <https://arxiv.org/abs/1709.00248>, doi:10.1016/j.jalgebra.2018.10.035.
- [AST17] H.H. Andersen, C. Stroppel, and D. Tubbenhauer. Semisimplicity of Hecke and (walled) Brauer algebras. *J. Aust. Math. Soc.*, 103(1):1–44, 2017. URL: <https://arxiv.org/abs/1507.07676>, doi:10.1017/S1446788716000392.
- [AST18] H.H. Andersen, C. Stroppel, and D. Tubbenhauer. Cellular structures using U_q -tilting modules. *Pacific J. Math.*, 292(1):21–59, 2018. URL: <https://arxiv.org/abs/1503.00224>, doi:10.2140/pjm.2018.292.21.
- [Bow22] C. Bowman. The many integral graded cellular bases of Hecke algebras of complex reflection groups. *Amer. J. Math.*, 144(2):437–504, 2022. URL: <https://arxiv.org/abs/1702.06579>, doi:10.1353/ajm.2022.0008.
- [Bro55] W.P. Brown. Generalized matrix algebras. *Canadian J. Math.*, 7:188–190, 1955. doi:10.4153/CJM-1955-023-2.
- [CP61] A.H. Clifford and G.B. Preston. *The algebraic theory of semigroups. Vol. I*. Mathematical Surveys, No. 7. American Mathematical Society, Providence, R.I., 1961.
- [CZ19] K. Coulembier and R. Zhang. Borelic pairs for stratified algebras. *Adv. Math.*, 345:53–115, 2019. URL: <https://arxiv.org/abs/1607.01867>, doi:10.1016/j.aim.2019.01.002.
- [dC06] F. du Cloux. Positivity results for the Hecke algebras of noncrystallographic finite Coxeter groups. *J. Algebra*, 303(2):731–741, 2006. URL: <https://arxiv.org/abs/math/0506448>, doi:10.1016/j.jalgebra.2005.10.004.
- [DEE⁺19] I. Dolinka, J. East, A. Evangelou, D. FitzGerald, N. Ham, J. Hyde, N. Loughlin, and J.D. Mitchell. Enumeration of idempotents in planar diagram monoids. *J. Algebra*, 522:351–385, 2019. URL: <https://arxiv.org/abs/1507.04838>, doi:10.1016/j.jalgebra.2018.11.014.
- [Eas06] J. East. Cellular algebras and inverse semigroups. *J. Algebra*, 296(2):505–519, 2006. doi:10.1016/j.jalgebra.2005.04.027.
- [EMRT18] J. East, J.D. Mitchell, N. Ruškuc, and M. Torpey. Congruence lattices of finite diagram monoids. *Adv. Math.*, 333:931–1003, 2018. URL: <https://arxiv.org/abs/1709.00142>, doi:10.1016/j.aim.2018.05.016.
- [Eny04] J. Enyang. Cellular bases for the Brauer and Birman–Murakami–Wenzl algebras. *J. Algebra*, 281(2):413–449, 2004. doi:10.1016/j.jalgebra.2003.03.002.
- [ER20] J. East and N. Ruškuc. Congruence lattices of ideals in categories and (partial) semigroups. 2020. URL: <https://arxiv.org/abs/2001.01909>.
- [ET21] M. Ehrig and D. Tubbenhauer. Relative cellular algebras. *Transform. Groups*, 26(1):229–277, 2021. URL: <https://arxiv.org/abs/1710.02851>, doi:10.1007/S00031-019-09544-5.
- [FG95] S. Fishel and I. Grojnowski. Canonical bases for the Brauer centralizer algebra. *Math. Res. Lett.*, 2(1):15–26, 1995. doi:10.4310/MRL.1995.v2.n1.a3.

- [FHH09] D. Flath, T. Halverson, and K. Herbig. The planar rook algebra and Pascal’s triangle. *Enseign. Math. (2)*, 55(1-2):77–92, 2009. URL: <https://arxiv.org/abs/0806.3960>, doi:10.4171/lem/55-1-3.
- [Gec07] M. Geck. Hecke algebras of finite type are cellular. *Invent. Math.*, 169(3):501–517, 2007. URL: <https://arxiv.org/abs/math/0611941>, doi:10.1007/s00222-007-0053-2.
- [GG11] F.M. Goodman and J. Graber. Cellularity and the Jones basic construction. *Adv. in Appl. Math.*, 46(1-4):312–362, 2011. URL: <https://arxiv.org/abs/0906.1496>, doi:10.1016/j.aam.2010.10.003.
- [GL96] J.J. Graham and G. Lehrer. Cellular algebras. *Invent. Math.*, 123(1):1–34, 1996. doi:10.1007/BF01232365.
- [GMS09] O. Ganyushkin, V. Mazorchuk, and B. Steinberg. On the irreducible representations of a finite semigroup. *Proc. Amer. Math. Soc.*, 137(11):3585–3592, 2009. URL: <https://arxiv.org/abs/0712.2076>, doi:10.1090/S0002-9939-09-09857-8.
- [GP00] M. Geck and G. Pfeiffer. *Characters of finite Coxeter groups and Iwahori–Hecke algebras*, volume 21 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 2000.
- [Gre51] J.A. Green. On the structure of semigroups. *Ann. of Math. (2)*, 54:163–172, 1951. doi:10.2307/1969317.
- [GW15] N. Guay and S. Wilcox. Almost cellular algebras. *J. Pure Appl. Algebra*, 219(9):4105–4116, 2015. doi:10.1016/j.jpaa.2015.02.010.
- [HJ20] T. Halverson and T.N. Jacobson. Set-partition tableaux and representations of diagram algebras. *Algebr. Comb.*, 3(2):509–538, 2020. URL: <https://arxiv.org/abs/1808.08118>, doi:10.5802/alco.102.
- [HM10] J. Hu and A. Mathas. Graded cellular bases for the cyclotomic Khovanov–Lauda–Rouquier algebras of type A. *Adv. Math.*, 225(2):598–642, 2010. URL: <http://arxiv.org/abs/0907.2985>, doi:10.1016/j.aim.2010.03.002.
- [HR05] T. Halverson and A. Ram. Partition algebras. *European J. Combin.*, 26(6):869–921, 2005. URL: <https://arxiv.org/abs/math/0401314>, doi:10.1016/j.ejc.2004.06.005.
- [HZ57] E. Hewitt and H.S. Zuckerman. The irreducible representations of a semi-group related to the symmetric group. *Illinois J. Math.*, 1:188–213, 1957.
- [Jen20a] L.T. Jensen. The ABC of p -cells. *Selecta Math. (N.S.)*, 26(2):Paper No. 28, 46, 2020. URL: <https://arxiv.org/abs/1901.02323>, doi:10.1007/s00029-020-0552-1.
- [Jen20b] L.T. Jensen. Cellularity of the p -canonical basis for symmetric groups. 2020. URL: <https://arxiv.org/abs/2009.11715>.
- [JW17] L.T. Jensen and G. Williamson. The p -canonical basis for Hecke algebras. In *Categorification and higher representation theory*, volume 683 of *Contemp. Math.*, pages 333–361. Amer. Math. Soc., Providence, RI, 2017. URL: <https://arxiv.org/abs/1510.01556>, doi:10.1090/conm/683.
- [KL79] D. Kazhdan and G. Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.*, 53(2):165–184, 1979. doi:10.1007/BF01390031.
- [KM16] T. Kildetoft and V. Mazorchuk. Special modules over positively based algebras. *Doc. Math.*, 21:1171–1192, 2016. URL: <https://arxiv.org/abs/1601.06975>.
- [KST22] M. Khovanov, M. Sitaraman, and D. Tubbenhauer. Monoidal categories, representation gap and cryptography. 2022. URL: <https://arxiv.org/abs/2201.01805>.
- [KX99] S. König and C. Xi. Cellular algebras: inflations and Morita equivalences. *J. London Math. Soc. (2)*, 60(3):700–722, 1999. doi:10.1112/S0024610799008212.
- [KX01] S. König and C. Xi. A characteristic free approach to Brauer algebras. *Trans. Amer. Math. Soc.*, 353(4):1489–1505, 2001. doi:10.1090/S0002-9947-00-02724-0.
- [KX12] S. König and C. Xi. Affine cellular algebras. *Adv. Math.*, 229(1):139–182, 2012. doi:10.1016/j.aim.2011.08.010.
- [Lus84] G. Lusztig. *Characters of reductive groups over a finite field*, volume 107 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1984. doi:10.1515/9781400881772.
- [Lus87] G. Lusztig. Leading coefficients of character values of Hecke algebras. In *The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986)*, volume 47 of *Proc. Sympos. Pure Math.*, pages 235–262. Amer. Math. Soc., Providence, RI, 1987.

- [Lus03] G. Lusztig. *Hecke algebras with unequal parameters*, volume 18 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 2003. URL: <https://arxiv.org/abs/math/0208154>.
- [Mat99] A. Mathas. *Iwahori-Hecke algebras and Schur algebras of the symmetric group*, volume 15 of *University Lecture Series*. American Mathematical Society, Providence, RI, 1999. doi:10.1090/ulect/015.
- [MMM⁺19] M. Mackaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer, and X. Zhang. Simple transitive 2-representations of Soergel bimodules for finite Coxeter types. 2019. URL: <https://arxiv.org/abs/1906.11468>.
- [MMM⁺21] M. Mackaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer, and X. Zhang. Finitary birepresentations of finitary bicategories. *Forum Math.*, 33(5):1261–1320, 2021. URL: <https://arxiv.org/abs/2008.01658>, doi:10.1515/forum-2021-0021.
- [MMMZ20] M. Mackaay, Volodymyr M., V. Miemietz, and X. Zhang. Analogues of centralizer subalgebras for fiat 2-categories and their 2-representations. *J. Inst. Math. Jussieu*, 19(6):1793–1829, 2020. URL: <https://arxiv.org/abs/1802.02078>, doi:10.1017/s1474748018000555.
- [MT21] A. Mathas and D. Tubbenhauer. Subdivision and cellularity for weighted KLRW algebras. 2021. URL: <https://arxiv.org/abs/2111.12949>.
- [MT22] A. Mathas and D. Tubbenhauer. Cellularity for weighted KLRW algebras of types B, $A^{(2)}$, $D^{(2)}$. 2022. URL: <https://arxiv.org/abs/2201.01998>.
- [Mun57] W.D. Munn. The characters of the symmetric inverse semigroup. *Proc. Cambridge Philos. Soc.*, 53:13–18, 1957. doi:10.1017/s0305004100031947.
- [RSA14] D. Ridout and Y. Saint-Aubin. Standard modules, induction and the structure of the Temperley–Lieb algebra. *Adv. Theor. Math. Phys.*, 18(5):957–1041, 2014. URL: <https://arxiv.org/abs/1204.4505>.
- [RT21] D.E.V. Rose and D. Tubbenhauer. HOMFLYPT homology for links in handlebodies via type A Soergel bimodules. *Quantum Topol.*, 12(2):373–410, 2021. URL: <https://arxiv.org/abs/1908.06878>, doi:10.4171/qt/152.
- [Sch58] M.-P. Schützenberger. Sur la représentation monomiale des demi-groupes. *C. R. Acad. Sci. Paris*, 246:865–867, 1958.
- [Sol02] L. Solomon. Representations of the rook monoid. *J. Algebra*, 256(2):309–342, 2002. doi:10.1016/S0021-8693(02)00004-2.
- [Spe20] R.A. Spencer. The modular Temperley–Lieb algebra. 2020. URL: <https://arxiv.org/abs/2011.01328>.
- [Ste16] B. Steinberg. *Representation theory of finite monoids*. Universitext. Springer, Cham, 2016. doi:10.1007/978-3-319-43932-7.
- [TV21] D. Tubbenhauer and P. Vaz. Handlebody diagram algebras. 2021. To appear in *Rev. Mat. Iberoam.* URL: <https://arxiv.org/abs/2105.07049>.
- [TVW17] D. Tubbenhauer, P. Vaz, and P. Wedrich. Super q -Howe duality and web categories. *Algebr. Geom. Topol.*, 17(6):3703–3749, 2017. URL: <https://arxiv.org/abs/1504.05069>, doi:10.2140/agt.2017.17.3703.
- [Xi99] C. Xi. Partition algebras are cellular. *Compositio Math.*, 119(1):99–109, 1999. doi:10.1023/A:1001776125173.

D.T.: THE UNIVERSITY OF SYDNEY, SCHOOL OF MATHEMATICS AND STATISTICS F07, OFFICE CARSLAW 827, NSW 2006, AUSTRALIA, WWW.DTUBBENHAUER.COM
 Email address: daniel.tubbenhauer@sydney.edu.au