

A Mixed Graph Achieving a Moore-like Bound

Geoffrey Exoo

Department of Mathematics and Computer Science
Indiana State University
Terre Haute, IN 47802
ge@cs.indstate.edu

Thursday June 16, 2022

Abstract

Mixed graphs have both directed and undirected edges. A mixed cage is a regular mixed graph of given girth with minimum possible order. In this paper we construct a mixed cage of order 30 that achieves the mixed graph analogue of the Moore bound for degree 3, out-degree 1, and girth 6.

1 Notation and Terminology

A mixed graph is a graph with both directed and undirected edges. We refer to directed edges as *arcs* and undirected edges as *edges*. The *degree* of a vertex v in a mixed graph G is the number of edges incident with v , whereas the *in-degree* and *out-degree* are the numbers of arcs incident to and from v . A G is *regular* if all three degrees are constant as v ranges over $V(G)$.

A *cycle* in a mixed graph is a sequence of vertices v_0, v_1, \dots, v_k such that there are no repeated vertices except that $v_0 = v_k$, each pair of consecutive vertices (v_i, v_{i+1}) is either an edge or an arc, and there are no repeated edges or arcs. The *girth* of a mixed graph is the length of a shortest cycle. Note that this definition considers the possibility of 1-cycles (loops) and 2-cycles.

Mixed graphs have been studied in the context of the degree/diameter problem. We follow the notation used in that literature and denote the degree and outdegree of a regular mixed graph by r and z , respectively.

An (r, z, g) -graph is a regular mixed graph with degree r , out-degree z , and girth g . An (r, z, g) -cage is an (r, z, g) -graph of minimum possible order. We denote this minimum order by $f(r, z, g)$.

Recall [2] that the Moore bound for an r -regular graph of diameter d is given by:

$$n(r, d) \geq \frac{r(r-1)^d - 2}{r-2} \quad (1)$$

In [1] Araujo-Pardo, Hernández-Cruz, and Montellano-Ballesteros consider the problem of finding mixed cages. They focus on the case $z = 1$ and determine a lower bound for $f(r, 1, g)$ based on

the Moore bound. Their idea is to attach undirected Moore trees to each vertex of a directed path of length $g-1$, choosing trees whose depth is as large as possible while still guaranteeing that all tree vertices are distinct.

So let v_0, v_1, \dots, v_{g-1} be the vertices of a directed path of length $g-1$. Using edges, attach a Moore tree of depth i to both v_i and v_{g-1-i} for $0 \leq i \leq \lfloor g/2 \rfloor$. Note: if $i = g-i-1$ we attach only one tree. The base path contains arcs and the Moore trees contain edges. This gives the following bound.

Theorem 1 (The AHM Bound [1])

$$f(r, 1, g) \geq \sum_{i=0}^{g-1} n(r, \min(i, g-i-1))$$

Theorem 2

$$f(3, 1, 6) = 30$$

Proof. The AHM bound for $r = 3$, $z = 1$, and $g = 6$ is 30. The graph G shown in the figure has order 30 and has the require parameters.

The graph can be describe algebraically as follows. Let

$$V(G) = \{v(i, j) \mid 0 \leq i < 3, 0 \leq j < 10\}$$

be the vertex set. For $0 \leq i < 3$, let

$$V_i = \{v(i, j) \mid 0 \leq j < 10\}.$$

Each V_i induces a directed 10-cycle. These three cycles are shown in figure. Let V_0 be the set of vertices in the lower partion of the figure and let V_1 and V_2 be the sets of vertices on the outer and inner (resp.) cycles in the upper part of the figure. Imagine the vertices labeled such that vertex $v_{i,0}$ is the rightmost vertex on each of the three directed 10-cycles and that the other vertices are labeled in counter-clockwise order. The arcs and edges are given as follows, where second indices are computed module 10:

- (a) arc($v(0, j), v(0, j+1)$)
- (b) edge($v(0, j), v(1, j)$)
- (c) edge($v(0, j), v(2, j+5)$)
- (d) edge($v(1, j), v(2, j+2)$)
- (e) edge($v(1, j), v(2, j-2)$)

In the figure, edges of type (b) and (c) are not drawn but indicated by color: vertices in V_0 are adjacent to those vertices in V_1 and V_2 that have the matching color.




Figure 1: The unique smallest $(3, 1, 6)$ -graph of order 30. Vertices in the lower figure are adjacent to vertices in the upper figure that have the matching color. The automorphism group has two generators: a rotation of $\pi/5$ (in both figures) and an involution that transposes the inner and outer cycles in the upper figure. These generators commute, so the group is $Z_2 \times Z_{10}$.

References

- [1] G. Araujo-Pardo, C. Hernández-Cruz, and J. J. Montellano-Ballesteros, Mixed Cages. *Graphs and Combinatorics*, 35 (2019), 989–999.
- [2] M. Miller and J. Sirán, Moore Graphs and Beyond: A Survey of the Degree/Diameter Problem, *Electronic Journal of Combinatorics* DS14, May 16, 2013.