

# RECURRENCES FOR THE DERIVATIVE POLYNOMIALS FOR TANGENT AND SECANT

GUO-NIU HAN AND SHI-MEI MA

**ABSTRACT.** In this paper, we choose the derivative polynomials for tangent and secant as basis sets of polynomial space. From this viewpoint, we first give an expansion of the derivative polynomials for tangent in terms of the derivative polynomials for secant, and we then present a result in the reverse direction. We also discuss the relationships between alternating derivative polynomials and Eulerian polynomials. As applications, we give certain expansions of the alternating derivative polynomials, which indicate that the alternating derivative polynomials share more properties with the Chebyshev polynomials.

**Keywords:** Derivative polynomials; Alternating derivative polynomials; Eulerian polynomials;

## 1. INTRODUCTION

Let  $\mathfrak{S}_n$  be the symmetric group of all permutations of  $[n]$ , where  $[n] = \{1, 2, \dots, n\}$ . Let  $\pi = \pi(1)\pi(2)\cdots\pi(n) \in \mathfrak{S}_n$ . We say that  $\pi$  is *alternating* if

$$\pi(1) > \pi(2) < \pi(3) > \cdots \pi(n).$$

In other words,  $\pi(i) < \pi(i+1)$  if  $i$  is even and  $\pi(i) > \pi(i+1)$  if  $i$  is odd. Let  $E_n$  denote the number of alternating permutations in  $\mathfrak{S}_n$ . For instance,  $E_4 = \#\{2143, 3142, 3241, 4132, 4231\}$ . A famous result of André [2] says that

$$\sum_{n=0}^{\infty} E_n \frac{z^n}{n!} = \tan z + \sec z = 1 + z + \frac{z^2}{2!} + 2\frac{z^3}{3!} + 5\frac{z^4}{4!} + 16\frac{z^5}{5!} + \cdots. \quad (1)$$

Since Euler used (1) as the definition of  $E_n$ , the numbers  $E_n$  are called *Euler numbers*. Note that

$$\sum_{n=0}^{\infty} E_{2n+1} \frac{z^{2n+1}}{(2n+1)!} = \tan z, \quad \sum_{n=0}^{\infty} E_{2n} \frac{z^{2n}}{(2n)!} = \sec z.$$

For this reason the numbers  $E_{2n+1}$  are sometimes called *tangent numbers* and the numbers  $E_{2n}$  are called *secant numbers*.

It is clear that

$$\begin{cases} \frac{d}{d\theta} \tan \theta = 1 + \tan^2 \theta, \\ \frac{d}{d\theta} \sec \theta = \tan \theta \sec \theta. \end{cases} \quad (2)$$

Besides (2), there are some differential systems can be also generated by trigonometric functions. As illustrations, we list four examples.

**Example 1.** Setting  $y(\theta) = \sec \theta$  and  $z(\theta) = \tan \theta$ , one has

$$\begin{cases} \frac{d}{d\theta}y(\theta) = y(\theta)z(\theta), \\ \frac{d}{d\theta}z(\theta) = y^2(\theta), \end{cases}$$

which can be used to generate peak and left peak polynomials (see [8, 20] for details).

**Example 2.** Let  $x(\theta) = (\tan \theta + \sec \theta)^q$ ,  $y(\theta) = \sec \theta$  and  $z(\theta) = \tan \theta$ , where  $q$  is a given parameter. One has

$$\begin{cases} \frac{d}{d\theta}x(\theta) = qx(\theta)y(\theta), \\ \frac{d}{d\theta}y(\theta) = y(\theta)z(\theta), \\ \frac{d}{d\theta}z(\theta) = y^2(\theta), \end{cases}$$

which can be used to generate run polynomials of permutations (see [23, Section 3]).

**Example 3.** Setting  $x(\theta) = \sec \theta (\tan \theta + \sec \theta)$  and  $y(\theta) = \tan \theta + \sec \theta$ , one has

$$\begin{cases} \frac{d}{d\theta}x(\theta) = x(\theta)y(\theta), \\ \frac{d}{d\theta}y(\theta) = x(\theta), \end{cases}$$

which can be used to generate André polynomials of 0-1-2 increasing trees (see [8, 11]).

**Example 4.** Let  $x(\theta) = \sec(2\theta)$  and  $y(\theta) = 2 \tan(2\theta)$ . One has

$$\begin{cases} \frac{d}{d\theta}x(\theta) = x(\theta)y(\theta), \\ \frac{d}{d\theta}y(\theta) = 4x^2(\theta), \end{cases}$$

which can be used to generate  $\gamma$ -coefficient polynomials of the types A and B Eulerian and Narayana polynomials (see [22] for details).

In this paper, we focus on the derivative polynomials generated by (2). Note that

$$\frac{d}{d\theta}(\tan^n \theta) = n \tan^{n-1} \theta (1 + \tan^2 \theta),$$

$$\frac{d}{d\theta}(\sec \theta \tan^n \theta) = \sec \theta (n \tan^{n-1} \theta + (n+1) \tan^{n+1} \theta).$$

The derivative polynomials for tangent and secant are respectively defined as follows:

$$\frac{d^n}{d\theta^n} \tan \theta = P_n(\tan \theta), \quad \frac{d^n}{d\theta^n} \sec \theta = \sec \theta Q_n(\tan \theta).$$

The study of these polynomials was initiated by Knuth and Buckholtz [19]. Recently, there has been much work on various properties and applications of derivative polynomials (see [4, 10, 17, 18, 21]). As discussed by Adamchik [1], Boyadzhiev [4], Hoffman [16, 17] and Qi [26], the derivative polynomials can be used to express some improper integrals and infinite series, including Hurwitz zeta functions, Dirichlet  $L$ -series and series of powers of reciprocals of integers.

In [19], Knuth and Buckholtz noted that

$$P_{2n+1}(0) = E_{2n+1}, \quad Q_{2n}(0) = E_{2n},$$

and by the chain rule, they deduced that

$$P_{n+1}(x) = (1+x^2) \frac{d}{dx} P_n(x), \quad Q_{n+1}(x) = (1+x^2) \frac{d}{dx} Q_n(x) + x Q_n(x), \quad (3)$$

with  $P_0(x) = x$  and  $Q_0(x) = 1$ . Carlitz and Scoville [7] derived that

$$P(x; z) = \sum_{n=0}^{\infty} P_n(x) \frac{z^n}{n!} = \frac{x + \tan z}{1 - x \tan z}, \quad Q(x; z) = \sum_{n=0}^{\infty} Q_n(x) \frac{z^n}{n!} = \frac{\sec z}{1 - x \tan z}. \quad (4)$$

In [16], Hoffman noted that

$$P_n(1) = 2^n(P_n(0) + Q_n(0)) = \begin{cases} 2^n Q_n(0) = 2^{2k} E_{2k}, & \text{if } n = 2k \text{ is even;} \\ 2^n P_n(0) = 2^{2k+1} E_{2k+1}, & \text{if } n = 2k + 1 \text{ is odd.} \end{cases} \quad (5)$$

Subsequently, Hoffman [17] discussed the other particular values of derivative polynomials. For example, Hoffman [17] noted that the numbers  $Q_n(1)$  are the Springer numbers of root systems of type  $B_n$ , which also count snakes of type  $B_n$ . A *snake* of type  $B_n$  is a signed permutation  $\pi(1)\pi(2)\cdots\pi(n)$  of  $B_n$  such that  $0 < \pi(1) > \pi(2) < \cdots < \pi(n)$ . Setting  $s_n = Q_n(1)$ , one has

$$\sum_{n=0}^{\infty} s_n \frac{z^n}{n!} = \frac{1}{\cos z - \sin z}.$$

One of the main results of Hoffman is the following expression of  $Q_n(1)$  (see [17, Theorem 3.1]):

$$Q_n(1) = -\sin \frac{n\pi}{2} + \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} (-1)^k P_{n-2k}(1). \quad (6)$$

This paper is motivated by the following problem.

**Problem 5.** *Are there some generalizations or variations of (6)?*

The Chebyshev polynomials  $T_n(x)$  of the first kind are defined by

$$T_n(x) = \cos(n\theta), \text{ when } x = \cos(\theta),$$

while the second kind Chebyshev polynomials  $U_n(x)$  are defined by

$$U_n(x) = \frac{\sin((n+1)\theta)}{\sin(\theta)}, \text{ when } x = \cos(\theta).$$

It is well known that the polynomials  $T_n(x)$  and  $U_n(x)$  are both alternately even and odd, and there are some close connections between the Chebyshev polynomials of the first and second kinds, see [24] for details. For example, the trigonometric identity

$$\sin((n+1)\theta) - \sin((n-1)\theta) = 2 \sin(\theta) \cos(n\theta)$$

leads to an identity:  $U_n(x) - U_{n-2}(x) = 2T_n(x)$ . It is well known that the Chebyshev polynomials form an orthogonal basis over the finite interval  $[-1, 1]$ .

In [15], Heteyi showed that derivative polynomials share some properties with Chebyshev polynomials, and derivative polynomials are closely related to the face enumerating polynomials of the Chebyshev transforms of the Boolean algebras. As an illustration, we give an example.

**Example 6** ([15, Corollary 8.7]). *The zeros of  $P_n(x)$  and  $Q_n(x)$  are pure imaginary, have multiplicity 1, belong to the line segment  $[-i, i]$  and are interlaced.*

Note that  $P_n(-x) = (-1)^{n+1}P_n(x)$  and  $Q_n(x) = (-1)^nQ_n(-x)$ . Then the polynomials  $P_n(x)$  and  $Q_n(x)$  are both alternately even and odd. Below are  $P_n(x)$  and  $Q_n(x)$  for  $n \leq 5$ :

$$\begin{aligned} P_0(x) &= x, & P_1(x) &= 1 + x^2, & P_2(x) &= 2x + 2x^3, & P_3(x) &= 2 + 8x^2 + 6x^4, \\ P_4(x) &= 16x + 40x^3 + 24x^5, & P_5(x) &= 16 + 136x^2 + 240x^4 + 120x^6, \\ Q_0(x) &= 1, & Q_1(x) &= x, & Q_2(x) &= 1 + 2x^2, & Q_3(x) &= 5x + 6x^3, \\ Q_4(x) &= 5 + 28x^2 + 24x^4, & Q_5(x) &= 61x + 180x^3 + 120x^5. \end{aligned}$$

Motivated by Proposition 6, we shall explore the following problem.

**Problem 7.** *If we choose the derivative polynomials for tangent or secant as basis sets of polynomial space, whether there are any interesting findings?*

In the next section, we present various results concerning Problems 5 and 7. In Section 3, we first introduce the definition of alternating derivative polynomials, and we then present certain combinatorial expansions of these polynomials.

## 2. RECURRENCES FOR DERIVATIVE POLYNOMIALS AND EULER NUMBERS

Let  $P(x; z)$  and  $Q(x; z)$  be given by (4). Hoffman [16, Eq. (5), Theorem 3.2] found that

$$\begin{aligned} \frac{d}{dz}P(x; z) &= 1 + P^2(x; z), \\ \frac{d}{dz}P(x; z) &= (1 + x^2)Q^2(x; z), \\ \frac{d}{dz}Q(x; z) &= P(x; z)Q(x; z), \end{aligned}$$

which yield three convolution formulas:

$$\begin{aligned} P_{n+1}(x) &= \delta_{0n} + \sum_{i=0}^n \binom{n}{i} P_i(x)P_{n-i}(x), \\ P_{n+1}(x) &= (1 + x^2) \sum_{i=0}^n \binom{n}{i} Q_i(x)Q_{n-i}(x), \\ Q_{n+1}(x) &= \sum_{i=0}^n \binom{n}{i} P_i(x)Q_{n-i}(x). \end{aligned}$$

Comparing the last two convolution formula with (3), we get the following result.

**Proposition 8.** *We have*

$$\begin{aligned} \frac{d}{dx}P_n(x) &= \sum_{i=0}^n \binom{n}{i} Q_i(x)Q_{n-i}(x) \text{ for } n \geq 0, \\ \frac{d}{dx}Q_n(x) &= \frac{1}{1 + x^2} \sum_{i=1}^n \binom{n}{i} P_i(x)Q_{n-i}(x) \text{ for } n \geq 1. \end{aligned}$$

Putting  $P_{-1}(x) = 1$ , then the polynomials  $P_{-1}(x)$ ,  $P_0(x)$ ,  $P_1(x)$ ,  $P_2(x)$ ,  $\dots$ ,  $P_n(x)$  form a basis for polynomials with degree less than or equal to  $n + 1$ , since they have different degrees. We can now present the following result.

**Theorem 9.** For any  $n \geq 0$ , one has

$$Q_{2n}(x) = (-1)^n + \sum_{k=0}^{n-1} \binom{2n}{2k+1} (-1)^k P_{2n-2k-1}(x), \quad (7)$$

$$Q_{2n+1}(x) = \sum_{k=0}^n \binom{2n+1}{2k+1} (-1)^k P_{2n-2k}(x). \quad (8)$$

*Proof.* By using (4), we obtain

$$\begin{aligned} Q(x; z) &= \frac{\sin^2(z) + \cos^2(z)}{\cos(z)} \frac{1}{1 - x \tan(z)} \\ &= \left( \cos(z) - x \sin(z) + x \sin(z) + \frac{\sin^2(z)}{\cos(z)} \right) \frac{1}{1 - x \tan(z)} \\ &= \frac{\cos(z) - x \sin(z)}{1 - x \tan(z)} + \sin(z) \frac{x + \tan(z)}{1 - x \tan(z)}. \end{aligned}$$

Thus

$$Q(x; z) = \cos(z) + \sin(z)P(x; z). \quad (9)$$

So we get

$$\begin{aligned} &\sum_{n=0}^{\infty} Q_{2n}(x) \frac{z^{2n}}{(2n)!} + \sum_{n=0}^{\infty} Q_{2n+1}(x) \frac{z^{2n+1}}{(2n+1)!} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n}}{(2n)!} + \sum_{k=0}^{\infty} \frac{(-1)^k z^{2k+1}}{(2k+1)!} \times \left( \sum_{n=1}^{\infty} \frac{P_{2n-1}(x)}{(2n-1)!} z^{2n-1} + \sum_{n=0}^{\infty} \frac{P_{2n}(x)}{(2n)!} z^{2n} \right) \end{aligned}$$

Selecting the coefficients of  $z^{2n}$  and  $z^{2n+1}$ , we get

$$\begin{aligned} \frac{Q_{2n}(x)}{(2n)!} &= \frac{(-1)^n}{(2n)!} + \sum_{k=0}^{n-1} \frac{(-1)^k}{(2k+1)!} \frac{P_{2n-2k-1}(x)}{(2n-2k-1)!}, \\ \frac{Q_{2n+1}(x)}{(2n+1)!} &= \sum_{k=0}^n \frac{(-1)^k}{(2k+1)!} \frac{P_{2n-2k}(x)}{(2n-2k)!}, \end{aligned}$$

and so the proof is complete.  $\square$

**Example 10.** Now specialize to the case  $n = 2$  in Theorem 9, we have

$$Q_4(x) = 5 + 28x^2 + 24x^4 = 4P_3(x) - 4P_1(x) + 1,$$

$$Q_5(x) = 61x + 180x^3 + 120x^5 = 5P_4(x) - 10P_2(x) + P_0(x).$$

**Theorem 11.** For any  $m \geq 0$ , we have

$$\sum_{k=0}^m \binom{2m}{2k} (-1)^k P_{2m-2k}(x) - x \sum_{k=0}^{m-1} \binom{2m}{2k+1} (-1)^k P_{2m-2k-1}(x) = (-1)^m x, \quad (10)$$

$$x \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k P_{2m-2k}(x) - \sum_{k=0}^m \binom{2m+1}{2k} (-1)^k P_{2m-2k+1}(x) = (-1)^{m+1}. \quad (11)$$

*Proof.* (i) Combining (3) and (7), we obtain

$$\begin{aligned}
Q_{2m+1}(x) &= (1+x^2) \frac{d}{dx} Q_{2m}(x) + xQ_{2m}(x) \\
&= \sum_{k=0}^{m-1} \binom{2m}{2k+1} (-1)^k (1+x^2) \frac{d}{dx} P_{2m-2k-1}(x) + xQ_{2m}(x) \\
&= \sum_{k=0}^{m-1} \binom{2m}{2k+1} (-1)^k P_{2m-2k}(x) + xQ_{2m}(x) \\
&= \sum_{k=0}^{m-1} \left( \binom{2m+1}{2k+1} - \binom{2m}{2k} \right) (-1)^k P_{2m-2k}(x) + xQ_{2m}(x) \\
&= \sum_{k=0}^{m-1} \binom{2m+1}{2k+1} (-1)^k P_{2m-2k}(x) - \sum_{k=0}^{m-1} \binom{2m}{2k} (-1)^k P_{2m-2k}(x) + xQ_{2m}(x) \\
&= Q_{2m+1}(x) - (-1)^m P_0(x) - \sum_{k=0}^{m-1} \binom{2m}{2k} (-1)^k P_{2m-2k}(x) + xQ_{2m}(x) \\
&= Q_{2m+1}(x) - \sum_{k=0}^m \binom{2m}{2k} (-1)^k P_{2m-2k}(x) + xQ_{2m}(x).
\end{aligned}$$

Hence

$$xQ_{2m}(x) - \sum_{k=0}^m \binom{2m}{2k} (-1)^k P_{2m-2k}(x) = 0.$$

It follows from (7) that

$$(-1)^m x + x \sum_{k=0}^{m-1} \binom{2m}{2k+1} (-1)^k P_{2m-2k-1}(x) - \sum_{k=0}^m \binom{2m}{2k} (-1)^k P_{2m-2k}(x) = 0,$$

and so we obtain (10).

(ii) Combining (3) and (8), we get

$$\begin{aligned}
Q_{2m+2}(x) &= (1+x^2) \frac{d}{dx} Q_{2m+1}(x) + xQ_{2m+1}(x) \\
&= \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k \left( (1+x^2) \frac{d}{dx} P_{2m-2k}(x) + xP_{2m-2k}(x) \right) \\
&= \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k (P_{2m-2k+1}(x) + xP_{2m-2k}(x)).
\end{aligned}$$

Since

$$\binom{2m+2}{2k+1} = \binom{2m+1}{2k+1} + \binom{2m+1}{2k},$$

we have

$$\begin{aligned}
&Q_{2m+2}(x) \\
&= \sum_{k=0}^m \left( \binom{2m+2}{2k+1} - \binom{2m+1}{2k} \right) (-1)^k P_{2m-2k+1}(x) + x \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k P_{2m-2k}(x) \\
&= \sum_{k=0}^m \binom{2m+2}{2k+1} (-1)^k P_{2m-2k+1}(x) + S_m(x),
\end{aligned}$$

where

$$S_m(x) = x \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k P_{2m-2k}(x) - \sum_{k=0}^m \binom{2m+1}{2k} (-1)^k P_{2m-2k+1}(x).$$

In view of (7), we find  $S_m(x) = (-1)^{m+1}$ . This completes the proof  $\square$

Recall that  $P_{2m-2k+1}(0) = E_{2m-2k+1}$  for any  $m \geq 0$ . Now specialize to the case  $x = 0$  in (11), one can get the following result.

**Corollary 12.** *For any  $m \geq 0$ , we have*

$$\sum_{k=0}^m \binom{2m+1}{2k} (-1)^k E_{2m-2k+1} = (-1)^m.$$

Combining (5) and Theorem 11, one can derive the following result.

**Corollary 13.** *For any  $m \geq 0$ , we have*

$$\begin{aligned} \sum_{k=0}^m \binom{2m}{2k} (-1)^k 2^{2m-2k} E_{2m-2k} - \sum_{k=0}^{m-1} \binom{2m}{2k+1} (-1)^k 2^{2m-2k-1} E_{2m-2k-1} &= (-1)^m, \\ \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k 2^{2m-2k} E_{2m-2k} - \sum_{k=0}^m \binom{2m+1}{2k} (-1)^k 2^{2m-2k+1} E_{2m-2k+1} &= (-1)^{m+1}. \end{aligned}$$

The *Bernoulli numbers*  $B_n$  can be defined by the exponential generating function

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} B_n \frac{z^n}{n!} = 1 - \frac{z}{2} + \frac{1}{6} \frac{z^2}{2!} - \frac{1}{30} \frac{z^4}{4!} + \frac{1}{42} \frac{z^6}{6!} - \frac{1}{30} \frac{z^8}{8!} + \dots$$

In particular,  $B_{2n+1} = 0$  for  $n \geq 1$ , since  $\frac{z}{2} \coth\left(\frac{z}{2}\right)$  is an even function and

$$\frac{z}{2} \coth\left(\frac{z}{2}\right) = \frac{z e^{z/2} + e^{-z/2}}{2 e^{z/2} - e^{-z/2}} = \frac{z}{e^z - 1} + \frac{z}{2}.$$

It is well known that the Bernoulli number has an explicit formula:

$$B_n = \sum_{k=0}^n \frac{(-1)^k k!}{k+1} \left\{ \begin{matrix} n \\ k \end{matrix} \right\},$$

where  $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$  is the Stirling number of the second kind, which counts the number of partitions of  $[n] = \{1, 2, \dots, n\}$  into  $k$  nonempty blocks. The Bernoulli numbers appear often in the coefficients of trigonometric functions (see [14, Chapter 6] for details). For example,

$$z \csc(z) = 1 + \sum_{n=1}^{\infty} (-1)^{n+1} (4^n - 2) B_{2n} \frac{z^{2n}}{(2n)!}, \quad z \cot(z) = \sum_{n=0}^{\infty} (-4)^n B_{2n} \frac{z^{2n}}{(2n)!} \quad (0 < |z| < \pi).$$

We can now present a result in the reverse direction of Theorem 9.

**Theorem 14.** *For any  $n \geq 0$ , one has*

$$\begin{aligned} (2n+1)P_{2n}(x) &= Q_{2n+1}(x) + \sum_{i=0}^{n-1} \binom{2n+1}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} Q_{2n-2i-1}(x), \\ (2n+2)P_{2n+1}(x) &= Q_{2n+2}(x) + \sum_{i=0}^n \binom{2n+2}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} Q_{2n-2i}(x) - (-4)^{n+1} B_{2n+2}. \end{aligned}$$

*Proof.* From (9), we see that  $P(x; z) = \csc(z)Q(x; z) - \cot(z)$ . Thus

$$zP(x; z) = z \csc(z)Q(x; z) - z \cot(z).$$

Then we have

$$\begin{aligned} & \sum_{n=0}^{\infty} P_{2n}(x) \frac{z^{2n+1}}{(2n)!} + \sum_{n=0}^{\infty} P_{2n+1}(x) \frac{z^{2n+2}}{(2n+1)!} \\ &= \left( 1 + \sum_{i=0}^{\infty} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} \frac{z^{2i+2}}{(2i+2)!} \right) \left( \sum_{n=0}^{\infty} Q_{2n}(x) \frac{z^{2n}}{(2n)!} + \sum_{n=0}^{\infty} Q_{2n+1}(x) \frac{z^{2n+1}}{(2n+1)!} \right) - \\ & \sum_{n=0}^{\infty} (-4)^n B_{2n} \frac{z^{2n}}{(2n)!}. \end{aligned}$$

Equating the coefficients of  $z^{2n+1}$  and  $z^{2n+2}$ , we get

$$\begin{aligned} \frac{P_{2n}(x)}{(2n)!} &= \frac{Q_{2n+1}(x)}{(2n+1)!} + \sum_{i=0}^{n-1} \frac{(-1)^{i+2} (4^{i+1} - 2) B_{2i+2}}{(2i+2)!} \frac{Q_{2n-2i-1}(x)}{(2n-2i-1)!}, \\ \frac{P_{2n+1}(x)}{(2n+1)!} &= \frac{Q_{2n+2}(x)}{(2n+2)!} + \sum_{i=0}^n \frac{(-1)^{i+2} (4^{i+1} - 2) B_{2i+2}}{(2i+2)!} \frac{Q_{2n-2i}(x)}{(2n-2i)!} - \frac{(-4)^{n+1} B_{2n+2}}{(2n+2)!}, \end{aligned}$$

and so the proof is complete.  $\square$

**Example 15.** Now specialize to the case  $n = 2$  in Theorem 14, we have

$$\begin{aligned} & Q_5(x) + \sum_{i=0}^1 \binom{5}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} Q_{3-2i}(x) \\ &= Q_5(x) + 10(4-2)B_2Q_3(x) - 5(4^2-2)B_4Q_1(x) \\ &= 61x + 180x^3 + 120x^5 + \frac{20}{6}(5x + 6x^3) + \frac{7}{3}x \\ &= 5P_4(x), \\ & Q_6(x) + \sum_{i=0}^2 \binom{6}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} Q_{4-2i}(x) - (-4)^3 B_6 \\ &= Q_6(x) + 30B_2Q_4(x) - 210B_4Q_2(x) + 62B_6 + 64B_6 \\ &= 61 + 662x^2 + 1320x^4 + 720x^6 + 5(5 + 28x^2 + 24x^4) + 7(1 + 2x^2) + 3 \\ &= 6P_5(x). \end{aligned}$$

Taking  $x = 1$  in Theorem 14 leads to the following result, which gives connections among tangent numbers, secant numbers, Springer numbers and Bernoulli numbers.

**Corollary 16.** For any  $n \geq 0$ , one has

$$\begin{aligned} (2n+1)2^{2n}E_{2n} &= s_{2n+1} + \sum_{i=0}^{n-1} \binom{2n+1}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} s_{2n-2i-1}, \\ (2n+2)2^{2n+1}E_{2n+1} &= s_{2n+2} + \sum_{i=0}^n \binom{2n+2}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} s_{2n-2i} - (-4)^{n+1} B_{2n+2}. \end{aligned}$$

Putting  $x = 0$  in Theorem 14 leads to the following result.



**Corollary 17.** *For any  $n \geq 0$ , one has*

$$E_{2n+2} = (2n+2)E_{2n+1} - \sum_{i=0}^n \binom{2n+2}{2i+2} (-1)^{i+2} (4^{i+1} - 2) B_{2i+2} E_{2n-2i} + (-4)^{n+1} B_{2n+2}.$$

### 3. EULERIAN POLYNOMIALS AND ALTERNATING DERIVATIVE POLYNOMIALS

Let  $\mathfrak{S}_n$  denote the symmetric group of all permutations of  $[n]$ . A *descent* of  $\pi \in \mathfrak{S}_n$  is an index  $i \in [n-1]$  such that  $\pi(i) > \pi(i+1)$ . Denote by  $\text{des}(\pi)$  the number of descents of  $\pi$ . The *Eulerian polynomials of type A* are defined by

$$A_n(x) = \sum_{\pi \in \mathfrak{S}_n} x^{\text{des}(\pi)+1}.$$

Let  $\pm[n] = [n] \cup \{\bar{1}, \dots, \bar{n}\}$ , where  $\bar{i} = -i$ . Denote by  $\mathcal{B}_n$  the hyperoctahedral group of rank  $n$ . Elements of  $\mathcal{B}_n$  are signed permutations  $\sigma$  of the set  $\pm[n]$  such that  $\sigma(-i) = -\sigma(i)$  for all  $i$ . Let  $\sigma = \sigma(1)\sigma(2) \cdots \sigma(n) \in \mathcal{B}_n$ . The *type B Eulerian polynomials* are defined by

$$B_n(x) = \sum_{\sigma \in \mathcal{B}_n} x^{\text{des}_B(\sigma)},$$

where  $\text{des}_B(\sigma) = \#\{i \in [0, n-1] : \sigma(i) > \sigma(i+1), \sigma(0) = 0\}$ . For  $n \geq 1$ , they satisfy the following recurrence relations (see [5, 9, 25] for instance):

$$\begin{cases} A_n(x) = nx A_{n-1}(x) + x(1-x) \frac{d}{dx} A_{n-1}(x), & A_0(x) = 1 \\ B_n(x) = (2nx+1-x) B_{n-1}(x) + 2x(1-x) \frac{d}{dx} B_{n-1}(x), & B_0(x) = 1. \end{cases} \quad (12)$$

The derivative polynomials for hyperbolic tangent and secant are defined by

$$\frac{d^n}{d\theta^n} \tanh(\theta) = \tilde{P}_n(\tanh(\theta)) \quad \text{and} \quad \frac{d^n}{d\theta^n} \text{sech}(\theta) = \text{sech}(\theta) \tilde{Q}_n(\tanh(\theta)).$$

Let  $i = \sqrt{-1}$ . Since  $\tanh(\theta) = -i \tan(i\theta)$  and  $\text{sech}(\theta) = \sec(i\theta)$ , we have

$$\tilde{P}_n(x) = i^{n-1} P_n(ix) \quad \text{and} \quad \tilde{Q}_n(x) = i^n Q_n(ix).$$

Using (3), we obtain that

$$\begin{cases} \tilde{P}_{n+1}(x) = (1-x^2) \frac{d}{dx} \tilde{P}_n(x), & \tilde{P}_0(x) = x; \\ \tilde{Q}_{n+1}(x) = (1-x^2) \frac{d}{dx} \tilde{Q}_n(x) - x \tilde{Q}_n(x), & \tilde{Q}_0(x) = 1. \end{cases} \quad (13)$$

**Definition 18.** *The alternatingly derivative polynomials for tangent and secant are respectively defined as follows:*

$$\begin{aligned} p_n(x) &= (-1)^n \tilde{P}_n(x) = (-1)^n i^{n-1} P_n(ix) \quad \text{for } n \geq 1, \quad p_0(x) = x, \\ q_n(x) &= (-1)^n 2^n \tilde{Q}_n(x) = (-1)^n 2^n i^n Q_n(ix) \quad \text{for } n \geq 0, \quad q_0(x) = 1. \end{aligned}$$

Let

$$P_n(x) = \sum_{k=0}^{\lfloor (n+1)/2 \rfloor} p(n, n-2k+1) x^{n-2k+1}, \quad Q_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} q(n, n-2k) x^{n-2k}.$$

It is easy to verify that

$$p_n(x) = \sum_{k=0}^{\lfloor (n+1)/2 \rfloor} p(n, n-2k+1) (-1)^k x^{n-2k+1}, \quad q_n(x) = 2^n \sum_{k=0}^{\lfloor n/2 \rfloor} q(n, n-2k) (-1)^k x^{n-2k}.$$

Using (13), we get

$$\begin{cases} p_{n+1}(x) = (x^2 - 1) \frac{d}{dx} p_n(x), & p_0(x) = x; \\ q_{n+1}(x) = 2(x^2 - 1) \frac{d}{dx} q_n(x) + 2xq_n(x), & q_0(x) = 1. \end{cases} \quad (14)$$

It should be noted that the distribution of zeros of  $p_n(x)$  and  $q_n(x)$  has been given by Heteyi [15, Theorem 8.6]. Comparing (12) and (14), it is routine to verify the following result, which is equivalent to [13, Theorem 5, Theorem 6]. For the sake of brevity we omit the proof.

**Theorem 19.** *For  $n \geq 1$ , we have*

$$p_n(x) = (x+1)^{n+1} A_n \left( \frac{x-1}{x+1} \right), \quad q_n(x) = (x+1)^n B_n \left( \frac{x-1}{x+1} \right).$$

In the past decades, Eulerian polynomials have been extensively studied, see [25] and references therein. Using Theorem 19, one can derive several properties of alternating derivative polynomials, including generating functions, particular values, explicit formulas and combinatorial expansions. In the sequel we shall give two applications of Theorem 19.

The most famous expansion of Eulerian polynomials is the *Frobenius formula*:

$$A_n(x) = x \sum_{k=0}^n k! \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x-1)^{n-k}. \quad (15)$$

Brenti [5, Theorem 3.14] found a generalization of (15):

$$B_n(x) = \sum_{k=0}^n k! \sum_{i=k}^n \binom{n}{i} 2^i \left\{ \begin{matrix} i \\ k \end{matrix} \right\} (x-1)^{n-k}. \quad (16)$$

Therefore, the following result follows from (15), (16) and Theorem 19.

**Corollary 20.** *For  $n \geq 1$ , we have*

$$\begin{aligned} p_n(x) &= (x-1) \sum_{k=0}^n (-2)^{n-k} k! \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x+1)^k, \\ q_n(x) &= \sum_{k=0}^n (-2)^{n-k} k! \sum_{i=k}^n \binom{n}{i} 2^i \left\{ \begin{matrix} i \\ k \end{matrix} \right\} (x+1)^k. \end{aligned}$$

Let  $f(x) = \sum_{i=0}^n f_i x^i$  be a symmetric polynomial, i.e.,  $f_i = f_{n-i}$  for any  $0 \leq i \leq n$ . Then  $f(x)$  can be expanded uniquely as  $f(x) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_k x^k (1+x)^{n-2k}$ , and it is said to be  $\gamma$ -positive if  $\gamma_k \geq 0$  for  $0 \leq k \leq \lfloor \frac{n}{2} \rfloor$ . The  $\gamma$ -positivity provides a natural approach to study symmetric and unimodal polynomials, see [3] and references therein.

Let  $\pi \in \mathfrak{S}_n$ . An index  $i \in [n]$  is a *peak* (resp. *double descent*) of  $\pi$  if  $\pi(i-1) < \pi(i) > \pi(i+1)$  (resp.  $\pi(i-1) > \pi(i) > \pi(i+1)$ ), where  $\pi(0) = \pi(n+1) = 0$ . Let  $a(n, k)$  be the number of permutations in  $\mathfrak{S}_n$  with  $k$  peaks and without double descents. Foata and Schützenberger [12] discovered that

$$A_n(x) = \sum_{k=1}^{\lfloor (n+1)/2 \rfloor} a(n, k) x^k (1+x)^{n+1-2k}. \quad (17)$$

Following [9, 25], one has

$$B_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} 4^k b(n, k) x^k (1+x)^{n-2k}, \quad (18)$$

where  $b(n, k) = \#\{\pi \in \mathfrak{S}_n : i \in [n-1], \pi(i-1) < \pi(i) > \pi(i+1), \pi(0) = 0\}$ . Substituting (17) and (18) into Theorem 19, we deduce the following.

**Corollary 21.** *For  $n \geq 1$ , we have*

$$p_n(x) = \sum_{k=1}^{\lfloor (n+1)/2 \rfloor} a(n, k) (x^2 - 1)^k (2x)^{n+1-2k},$$

$$q_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} 4^k b(n, k) (x^2 - 1)^k (2x)^{n-2k}.$$

The Chebyshev polynomials can be defined in terms of the sums (see [24] for instance):

$$T_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{2k} (x^2 - 1)^k x^{n-2k}, \quad U_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n+1}{2k+1} (x^2 - 1)^k x^{n-2k}. \quad (19)$$

Therefore, Corollary 21 can be seen as a dual of (19), which indicate that the alternating derivative polynomials deserve more work.

#### ACKNOWLEDGEMENTS.

This work is supported by NSFC (12071063).

#### REFERENCES

- [1] V.S. ADAMCHIK, *On the Hurwitz function for rational arguments*, Appl. Math. Comp., 187 (2007), 3–12.
- [2] D. ANDRÉ, *Développement de sec x et tan x*, C. R. Math. Acad. Sci. Paris., 88 (1879), 965–979.
- [3] C.A. ATHANASIADIS, *Gamma-positivity in combinatorics and geometry*, Sémin. Lothar. Combin., 77 (2018), Article B77i.
- [4] K.N. BOYADZHEV, *Derivative polynomials for tanh, tan, sech and sec in explicit form*, Fibonacci Quart., 45 (2007), 291–303.
- [5] F. BRENTI, *q-Eulerian polynomials arising from Coxeter groups*, European J. Combin., 15 (1994), 417–441.
- [6] C.-O. CHOW, *On certain combinatorial expansions of the Eulerian polynomials*, Adv. in Appl. Math., 41 (2008), 133–157.
- [7] L. CARLITZ, R. SCOVILLE, *Tangent numbers and operators*, Duke Math. J., 39 (1972), 413–429.
- [8] W.Y.C. CHEN, A.M. FU, *Context-free grammars for permutations and increasing trees*, Adv. in Appl. Math., 82 (2017), 58–82.
- [9] C.-O. CHOW, *On certain combinatorial expansions of the Eulerian polynomials*, Adv. in Appl. Math., 41 (2008), 133–157.
- [10] D. CVIJOVIĆ, *Derivative polynomials and closed-form higher derivative formulae*, Appl. Math. Comput., 185 (2009), 3002–3006.
- [11] D. DUMONT, *Grammaires de William Chen et dérivations dans les arbres et arborescences*, Sémin. Lothar. Combin., 37, Art. B37a (1996), 1–21.
- [12] D. FOATA, M. P. SCHÜTZENBERGER, *Théorie géométrique des polynômes eulériens*, Lecture Notes in Math., vol. 138, Springer, Berlin, 1970.

- [13] G.R. FRANSSENS, FUNCTIONS WITH DERIVATIVES GIVEN BY POLYNOMIALS IN THE FUNCTION ITSELF OR A RELATED FUNCTION, *Anal. Math.* 33 (2007), 17–36.
- [14] R.L. GRAHAM, D.E. KNUTH, O. PATASHNIK, *Concrete Mathematics*, Addison-Wesley, Reading, MA, 2nd. ed., 1994.
- [15] G. HETYEI, *Chebyshev triangulations of stable simplicial complexes*, *J. Combin. Theory Ser. A*, 115 (2008), 569–592.
- [16] M.E. HOFFMAN, *Derivative polynomials for tangent and secant*, *Amer. Math. Monthly*, 102 (1995), 23–30.
- [17] M.E. HOFFMAN, *Derivative polynomials, Euler polynomials, and associated integer sequences*, *Electron. J. Combin.*, 6 (1999), #R21.
- [18] M. Josuat-Vergès, Enumeration of snakes and cycle-alternating permutations, *Aust. J. Combin.*, 60(3) (2014), 279–305.
- [19] D.E. KNUTH, T.J. BUCKHOLTZ, *Computation of tangent, Euler, and Bernoulli numbers*, *Math. Comp.*, 21 (1967), 663–688.
- [20] S.-M. MA, *Derivative polynomials and enumeration of permutations by number of interior and left peaks*, *Discrete Math.*, 312 (2012), 405–412.
- [21] S.-M. MA, *An explicit formula for the number of permutations with a given number of alternating runs*, *J. Combin. Theory Ser. A*, 119 (2012), 1660–1664.
- [22] S.-M. MA, *On  $\gamma$ -vectors and the derivatives of the tangent and secant functions*, *Bull. Aust. Math. Soc.* 90 (2) (2014), 177–185.
- [23] S.-M. MA, J. MA, Y.-N. YEH, *David-Barton type identities and the alternating run polynomials*, *Adv. in Appl. Math.*, 114 (2020), 101978.
- [24] J.C. MASON, D.C. HANDSCOMB, *Chebyshev Polynomials*, Chapman and Hall/CRC, 2003.
- [25] T.K. PETERSEN, *Eulerian Numbers*, Birkhäuser/Springer, New York, 2015.
- [26] F. QI, *Derivatives of tangent function and tangent numbers*, *Appl. Math. Comput.*, 268 (2015), 844–858.

I.R.M.A., UMR 7501, UNIVERSITÉ DE STRASBOURG ET CNRS, 7 RUE RENÉ DESCARTES, F-67084 STRASBOURG, FRANCE

*Email address:* guoniu.han@unistra.fr (G.-N. Han)

SCHOOL OF MATHEMATICS AND STATISTICS, NORTHEASTERN UNIVERSITY AT QINHUANGDAO, HEBEI 066000, P.R. CHINA

*Email address:* shimeimapapers@163.com (S.-M. Ma)