

On the application of the Adomian-based methods

Francisco M. Fernández*
 INIFTA, DQT, Sucursal 4, C.C. 16,
 1900 La Plata, Argentina

Abstract

We analyse a recent application of two Adomian-based approximate methods to the Chen-Lee-Liu equation. We prove that the outcome of these approaches is merely the Taylor expansion of the solution about $t = 0$ and, consequently, they are valid only for sufficiently small values of the time variable t .

In a recent paper Mohammed and Bakodah (MB) [1] compared the performance of the Adomian decomposition method (ADM) and improved Adomian decomposition method (IADM) on the nonlinear partial differential equation

$$iu_t + au_{xx} + ib|u|^2u_x = 0, \quad (1)$$

and concluded that the later approximate method is slightly more accurate. In order to determine the accuracy of these algorithms the authors resorted to the exact solution [2]

$$u(x, t) = \sqrt{\gamma \pm \eta \operatorname{sech}[\lambda(x - \nu t)]} e^{i[\omega t - kx + \theta(x - \nu t)]}, \quad (2)$$

where γ , η and λ depend on the model parameters a and b as well as on some arbitrary constants k , B , ν and ω [1]. MB did not comment on the, apparently necessary, extra phase function $\theta(s)$ of the travelling variable $s = x - \nu t$ [2].

*fernande@quimica.unlp.edu.ar

MB showed tables with absolute errors and figures for some real function that they failed to indicate (probably $|u(x, t)|$ or $|u(x, t)|^2$). They chose $a = 5$ and $b = 10$ for their calculations but omitted the selected values of k , B , ν and ω . Therefore, it is not possible to reproduce their results obtained for $t = 0.3$ and $t = 0.5$. This fact does not affect the aim of the present Comment that is to show that they did not choose larger values of t because neither the ADM or the IADM are expected to yield acceptable results for sufficiently large values of the time variable.

For simplicity, we consider the ADM in what follows. MB did not show the application of the ADM to equation (1) but to the closely related one

$$u_t + iau_{xx} - b|u|^2u_x = 0. \quad (3)$$

Curiously, they did not comment on the connection between the solutions to these two nonlinear partial differential equations. The approach is based on an expansion of the solution as

$$u(x, t) = \sum_{j=0}^{\infty} u_j(x, t), \quad u_0(x, t) = u(x, 0), \quad (4)$$

from which it follows that $A = |u|^2u_x$ can be expanded as

$$A(x, t) = \sum_{j=0}^{\infty} A_j(x, t). \quad (5)$$

MB also forgot to say how many terms of the expansion (4) were considered in their calculations and, most importantly, to test the rate of convergence of the algorithms. From MB's equation (3.6) one easily shows that

$$A_n = \sum_{j=0}^n u_{n-j} \sum_{k=0}^j u_k \bar{u}_{j-k}, \quad (6)$$

where \bar{u}_j is the complex conjugate of u_j . Curiously, this simple and efficient closed-form expression has not been taken into account neither in this paper or in earlier ones where the authors have been doing this same calculation again and again [3–5]. One can easily verify that equation (6) already yields the expressions given in one of those papers for $n = 1, 2, 3, 4$ [4].

The ADM corrections are given by the recurrence relation [1]

$$u_{k+1}(x, t) = ai \int_0^t u_{k,x}(x, t') dt' - b \int_0^t A_k(x, t') dt', \quad k \geq 0. \quad (7)$$

Since $u_{0,x}(x, t)$ and $A_0(x, t) = |u(x, 0)|^2$ are independent of t , then $u_1(x, t)$ is of the form $u_1(x, t) = v_1(x)t$, where $v_1(x) = iau_{0,x}(x, 0) - b|u(x, 0)|^2$. In the next step we have $A_1(x, t) = u_1(x, t)\bar{u}_0(x, t)u_{0,x}(x, t) + u_0(x, t)\bar{u}_1(x, t)u_{0,x}(x, t) + u_0(x, t)\bar{u}_0(x, t)u_{1,x}(x, t) = B_1(x)t$ so that $u_2(x, t) = v_2(x)t^2$ where $v_2(x)$ is straightforwardly given by equation (7). Repeating this argument we conclude that $u_j(x, t) = v_j(x)t^j$ and $A_j(x, t) = B_j(x)t^j$, $j = 0, 1, \dots$. Therefore, the expansion (4) reads

$$u(x, t) = \sum_{j=0}^{\infty} v_j(x)t^j, \quad (8)$$

and shows that the ADM simply produces the Taylor expansion of $u(x, t)$ about $t = 0$. Consequently, the ADM approximation to the solution $u(x, t)$ will be valid only for sufficiently small values of t . MB's choices $t = 0.3$ and $t = 0.5$ do not reveal the shortcomings of the method. In earlier applications of similar approaches to the same type of equations the authors also chose values of the time variable in the range $0.1 \leq t \leq 0.5$ [3–5]. It is worth mentioning that the functions $v_j(x)$ can be obtained recursively from

$$v_{k+1}(x) = \frac{1}{k+1} [aiv_{k,x}(x) - bB_k(x)], \quad (9)$$

that appears to be somewhat simpler than equation (7).

The IAM consists of writing $u(x, t) = u_1(x, t) + iu_2(x, t)$, u_1 and u_2 real, and applying the ADM to the real and imaginary parts of equation (3) separately. A simple analysis, similar to the one above, shows that in this case one obtains the t -power series for $u_1(x, t)$ and $u_2(x, t)$ and, consequently, for $u(x, t)$. Therefore, the IADM exhibits the same drawback as the ADM.

Other authors have unknowingly derived Taylor series by means of unsuitable implementations of approximate methods [6–10].

References

- [1] Mohammed A S H F and Bakodah H O 2021 *Phys. Scr.* **96** 035206.
- [2] Triki H, Zhou Q, Moshokoa S P, Ullah M Z, Biswas A, and Belic M 2018 *Optik* **155** 208.
- [3] Mohammed A S H F, Bakodah H O, Banaja M A, Alshaery A A, Zhou Q, Biswas A, Moshokoa S P, and Belic M R 2019 *Optik* **181** 964.
- [4] Mohammed A S H F and Bakodah H O 2020 *Am. J. Comput. Math.* **10** 398.
- [5] Mohammed A S H F and Bakodah H O 2020 *Nonlin. Anal. Diff. Eq.* **8** 67.
- [6] Fernández F M 2009 *Phys. Scr.* **79** 055003.
- [7] Fernández F M 2009 *Appl. Math. Comput.* **215** 168.
- [8] Fernández F M 2010 *Phys. Scr.* **81** 037002.
- [9] Fernández F M 2014 *Appl. Math. Comput.* **230** 208.
- [10] Fernández F M 2020 *J. Phys. A* **53** 388001.