Geometry, conformal Killing-Yano tensors and conserved "currents"

Ulf Lindström a,b1 and Özgür Sarıoğlu a2

^aDepartment of Physics, Faculty of Arts and Sciences, Middle East Technical University, 06800, Ankara, Turkey

^bDepartment of Physics and Astronomy, Theoretical Physics, Uppsala University SE-751 20 Uppsala, Sweden

Abstract

In this brief letter we derive some useful identities relating conformal Killing-Yano tensors (CKYTs) and geometric quantities. We then use these identities to construct covariantly conserved "currents". We conclude that rank-n currents linear in rank-n CKYTs k and second order in derivatives must have a simple form in terms of dk. Using the Plebański-Demiański and the Kerr-Newman metrics, we show how these currents can be used to define charges. By construction, these currents are covariant under a general conformal rescaling of the metric.

 1 email: ulf.lindstrom@physics.uu.se

²email: sarioglu@metu.edu.tr

Contents

1	Introduction	2
2	rank-n CKYT identities	3
3	Currents of rank- n	4
4	Some special cases	5
	4.1 $n=1$	5
	4.2 $n=2$	6
	4.3 Plebański-Demiański metric	7
	4.4 Kerr-Newman metric	8
5	Discussion	8
A	CKYTs under Weyl scalings	g
В	Killing-Yano Cotton current	10
\mathbf{C}	Lie derivative of curvature tensors	11

1 Introduction

In recent papers [1–3], we have investigated Killing tensors (KTs) and Killing-Yano tensors (KYTs), identified new geometric identities constraining the geometries that carry such tensors and shown the existence of covariantly conserved antisymmetric tensors (that we call "currents") and worked out their related charges and asymptotic charges. Our inspiration originally came from analyzing the current construction in [4]. So far, we have only considered conformal Killing-Yano Tensors (CKYTs) in connection with the Cotton currents in [1], but a natural question to ask is which of the KYT currents can be generalized to include CKYTs. In the present paper, we address this question for rank-n currents constructed using rank-n CKYTs.

We follow the previous pattern of construction using the curvature tensor and its contractions and derive new identities between these and the CKYTs needed to evaluate the divergences of the currents. Using our identities, we find the unexpected result that the divergences of all currents can be expressed in a simple "obvious" form. In the language of differential forms, they only involve dk of the n-form CKYT k.

Section 2 contains a derivation of identities of general interest, some of which are needed for the derivation of the rank-n conserved currents in section 3. Some interesting special cases are collected in section 4. Our conclusions and a discussion are contained in section 5. In appendix A, we derive the transformation properties of a CKYT under a rescaling of the metric¹ and use

For $n \neq 2$ our formulae are at variance with some of the literature.

these to show that our "obvious" currents are covariant under such a rescaling. In appendix B, we review the Cotton current and prove the important property of covariance under rescalings for it as well. In appendix C, finally, we discuss Lie derivatives along conformal Killing vectors (CKVs). In the references we include a list of publications containing useful general background information about (C)KYTs and their applications [5,6].

2 rank-n CKYT identities

In this section we define a CKYT and derive identities involving CKYTs and geometric tensors. The geometric setting is always curved (pseudo-)Riemannian geometry. A CKYT k generalizes a KYT in the way a conformal Killing vector generalizes a Killing vector. The definition of a rank-n CKYT k is

$$\nabla_{a_1} k_{a_2...a_{n+1}} = \nabla_{[a_1} k_{a_2...a_{n+1}]} + n g_{a_1[a_2} K_{a_3...a_{n+1}]},$$

$$K_{a_1...a_{n-1}} := \frac{1}{(D-n+1)} \nabla_c k^c_{a_1...a_{n-1}} \text{ and } \nabla_{a_1} K^{a_1...a_{n-1}} = 0.$$
(2.1)

The definition reduces to that of a rank-n KYT f by setting $K^{a_1...a_{n-1}} = 0$. An important identity for KYTs [4] may be generalized to CKYTs as follows

$$\nabla_a \nabla_b k_{c_1...c_n} = (-1)^{n+1} \frac{(n+1)}{2} R^d_{a[bc_1} k_{c_2...c_n]d} - (n+1) g_{a[b} \nabla_{c_1} K_{c_2...c_n]} + n \nabla_a \left(g_{b[c_1} K_{c_2...c_n]} \right) . \tag{2.2}$$

Using (2.1) and (2.2) in the commutator $[\nabla_a, \nabla_b] \nabla_c k_{c_1...c_n}$, and contracting the index pairs (a, c_n) and (b, c) on both sides of the resultant expression, we get

$$(n-1)\left(\nabla^{b}R^{a}_{[c_{1}}\right)k_{c_{2}...c_{n-1}]ab} + \frac{1}{2}(\nabla^{a}R)k_{ac_{1}...c_{n-1}} + RK_{c_{1}...c_{n-1}} + (D-n)\square K_{c_{1}...c_{n-1}} - \frac{1}{3}(n-1)(n-2)(D-n-1)K^{ab}_{[c_{1}...c_{n-3}}\left(R_{c_{n-2}|a|c_{n-1}]b} + R_{c_{n-2}c_{n-1}]ab}\right) + (n-1)(D-n-2)K_{[c_{1}...c_{n-2}}{}^{a}R_{c_{n-1}]a} - (n-1)(D-n)\nabla_{a}\nabla_{[c_{1}}K^{a}_{c_{2}...c_{n-1}]} = 0.$$
 (2.3)

Since

$$2K^{ab}_{[c_1...c_{n-3}}R_{c_{n-2}|a|c_{n-1}]b} = K^{ab}_{[c_1...c_{n-3}}R_{c_{n-2}c_{n-1}]ab},$$
(2.4)

we have

$$-\frac{1}{3}K^{ab}_{[c_1...c_{n-3}]}(R_{c_{n-2}|a|c_{n-1}]b} + R_{c_{n-2}c_{n-1}]ab}) = -\frac{1}{2}K^{ab}_{[c_1...c_{n-3}]}R_{c_{n-2}c_{n-1}]ab}.$$
(2.5)

Moreover

$$\nabla_{a}\nabla_{[c_{1}}K^{a}{}_{c_{2}...c_{n-1}]} = [\nabla_{a}, \nabla_{[c_{1}]}K^{a}{}_{c_{2}...c_{n-1}]} = R_{a[c_{1}}K^{a}{}_{c_{2}...c_{n-1}]} - (n-2)R_{b[c_{1}c_{2}|a|}K^{ab}{}_{c_{3}...c_{n-1}]}$$

$$\Longrightarrow \nabla_{a}\nabla_{[c_{1}}K^{a}{}_{c_{2}...c_{n-1}]} = K_{[c_{1}...c_{n-2}}{}^{a}R_{c_{n-1}]a} - (n-2)R_{b[c_{n-2}c_{n-1}|a|}K^{ab}{}_{c_{1}...c_{n-3}]}. \tag{2.6}$$

These can be used for simplifying the second and third lines of (2.3) to arrive at

$$(n-1)\left(\nabla^{b}R^{a}_{[c_{1}}\right)k_{c_{2}...c_{n-1}]ab} + \frac{1}{2}(\nabla^{a}R)k_{ac_{1}...c_{n-1}} + (D-n)\Box K_{c_{1}...c_{n-1}} + RK_{c_{1}...c_{n-1}} - 2(n-1)K_{[c_{1}...c_{n-2}}{}^{a}R_{c_{n-1}]a} + \frac{1}{2}(n-1)(n-2)K^{ab}_{[c_{1}...c_{n-3}}R_{c_{n-2}c_{n-1}]ab} = 0.$$

$$(2.7)$$

On the other hand, by contracting the (a, b) indices in (2.2), we also have

$$\Box k^{c_1...c_n} = \frac{(n-1)}{2} k^{ab[c_1...c_{n-2}} R^{c_{n-1}c_n]}{}_{ab} - R^{a[c_1} k_a{}^{c_2...c_n]} + (2n-D) \nabla^{[c_1} K^{c_2...c_n]} . \quad (2.8)$$

These may be combined to show that the following current is covariantly conserved²

$$F^{c_1...c_n} = \Box k^{c_1...c_n} - n\nabla^{[c_1}K^{c_2...c_n]} = \nabla_a \left(\nabla^{[a}k^{c_1...c_n]}\right) \quad \text{and} \quad \nabla_{c_1}F^{c_1...c_n} = 0.$$
 (2.9)

3 Currents of rank-n

In this section, we use some of the formulae derived in section 2 to investigate a rank-n current constructed from a rank-n CKYT. More specifically, we look for a covariantly conserved rank-n tensor J linear in the CKYT and its contraction, formed from geometric tensors and covariant derivatives, such that J is second-order in covariant derivatives. This specification is in line with previous currents constructed using KYTs [2,4].

We start by listing a number of relations between terms of the right kind:

$$R^{[a_1|cd|a_2}k^{a_3...a_n]}{}_{cd} = -\frac{1}{2}R_{cd}^{[a_1a_2}k^{a_3...a_n]cd}, \qquad (3.1)$$

$$\nabla_c \nabla^{[a_1} k^{a_2...a_n]c} = \nabla_c \nabla^{[a_1} k^{a_2...a_nc]} + (-1)^{n+1} \nabla^{[a_1} K^{a_2...a_n]}, \tag{3.2}$$

$$\nabla^{[a_1}\nabla_c k^{a_2...a_n]c} = (-1)^{n+1}(D+1-n)\nabla^{[a_1}K^{a_2...a_n]},$$
(3.3)

$$[\nabla_c, \nabla^{[a_1]}] k^{a_2...a_n]c} = (-1)^n \frac{(n-1)}{2} R_{cd}^{[a_1 a_2} k^{a_3...a_n]cd} + R_c^{[a_1} k^{a_2...a_n]c},$$
(3.4)

$$\Box k^{a_1...a_n} = \nabla_c \nabla^{[c} k^{a_1...a_n]} + n \nabla^{[a_1} K^{a_2...a_n]}.$$
 (3.5)

Notice that the difference of (3.2) and (3.3) gives (3.4), and thus

$$F^{a_1...a_n} + (D-n)\nabla^{[a_1}K^{a_2...a_n]} = \frac{(n-1)}{2}R_{cd}^{[a_1a_2}k^{a_3...a_n]cd} + (-1)^nR_c^{[a_1}k^{a_2...a_n]c}.$$
 (3.6)

These relations let us choose the following rank-n, skew-symmetric, independent combinations

$$R_{cd}^{[a_1a_2} k^{a_3...a_n]cd}, R_c^{[a_1} k^{a_2...a_n]c}, R_c^{[a_1...a_n]}, \nabla^{[a_1} K^{a_2...a_n]},$$
 (3.7)

This kind of CKYT current, which is obtained by replacing $f \to k$ in the simplest of the KYT currents $\nabla_a \nabla^{[a} f^{b_1...b_n]}$ will be called *obvious*.

as the building blocks of a possible current. For computational ease, without loss of generality, we replace the first two terms in the list (3.7) by

$$\check{\mathcal{K}}^{a_1...a_n} := -\frac{(n-1)}{4} R^{[a_1 a_2} {}_{bc} k^{a_3...a_n]bc} + \frac{1}{2n} R k^{a_1...a_n},$$
(3.8)

and

$$\hat{\mathcal{K}}^{a_1...a_n} := R_c^{[a_1} k^{a_2...a_n]c} + \frac{(-1)^n}{n} R k^{a_1...a_n}, \qquad (3.9)$$

modeled after two conserved currents in [2].

With these preambles, we make an ansatz for the conserved current:

$$J^{a_1...a_n} = 2\check{\mathcal{K}}^{a_1...a_n} + \alpha\hat{\mathcal{K}}^{a_1...a_n} + \beta Rk^{a_1...a_n} + \gamma \nabla^{[a_1} K^{a_2...a_n]}. \tag{3.10}$$

To this, we may add any amount of the conserved current $F^{a_1...a_n}$ in (2.9), its coefficient will remain arbitrary.

To proceed, we rewrite the identity (2.8) or (3.6) in terms of the tensors (3.8) and (3.9):

$$F^{a_1...a_n} = (-1)^n \hat{\mathcal{K}}^{a_1...a_n} - 2 \check{\mathcal{K}}^{a_1...a_n} + (n-D) \nabla^{[a_1} K^{a_2...a_n]}. \tag{3.11}$$

Using this identity in (3.10) yields for the divergence

$$\nabla_{a_1} J^{a_1 \dots a_n} = (\alpha + (-1)^n) \nabla_{a_1} \hat{\mathcal{K}}^{a_1 \dots a_n} + (\gamma + n - D) \nabla_{a_1} \nabla^{[a_1} K^{a_2 \dots a_n]} + \beta \nabla_{a_1} (Rk^{a_1 \dots a_n}) + \nabla_{a_1} F^{a_1 \dots a_n}.$$
(3.12)

The last term is zero, of course, and the coefficients of the others give $\alpha = (-1)^{n+1}$, $\beta = 0$ and $\gamma = D - n$. Plugging these values into (3.10) and using (3.11), the rank-n current is thus shown to be obvious, i.e. proportional to the current $F^{a_1...a_n}$ in (2.9).

4 Some special cases

Since we are considering rank-n forms, we are restricted by the dimension D of the underlying manifold. Interesting cases are therefore $n = 1 \dots D/2$ in even and $n = 1 \dots (D-1)/2$ in odd dimensions (modulo their Hodge duals). Below we comment on some of these.

4.1 n=1

This is the special case when the CKYT is a conformal Killing vector (CKV). Consider thus the following contravariant vector

$$J^a = R^{ab}k_b + \alpha Rk^a + \beta \nabla^a K . (4.1)$$

Then

$$\nabla_a J^a = k^a \nabla_a R \left(\frac{1}{2} + \alpha + \frac{\beta}{2(1-D)} \right) + KR \left(1 + \alpha D + \frac{\beta}{(1-D)} \right). \tag{4.2}$$

Demanding $\nabla_a J^a = 0$ implies that either "D = 2, $\beta = 1 + 2\alpha$ with α left arbitrary" or " $D \neq 2$, $\alpha = 0$, $\beta = D - 1$ ". So one has a conserved vector J^a for

$$D = 2: \quad J^{a} = R^{ab}k_{b} + \alpha Rk^{a} + (1 + 2\alpha)\nabla^{a}K,$$

$$D \neq 2: \quad J^{a} = R^{ab}k_{b} + (D - 1)\nabla^{a}K = -\Box k^{a} + \nabla^{a}K = -F^{a}.$$
(4.3)

This follows the pattern described in Sec.3 for $D \neq 2$.

Since we are dealing with a CKV, a natural question to ask is on the transformation properties of J^a under Weyl scaling of the metric $\tilde{g}_{ab} = \Omega^2 g_{ab}$. A CKV transforms as

$$\tilde{k}^a = k^a \quad \text{and} \quad \tilde{K} = K + \Omega^{-1}(\nabla_a \Omega)k^a$$
. (4.4)

The transformations of the Ricci and scalar curvature is given by

$$\tilde{R}_{ab} = R_{ab} - \Omega^{-1} \left((D - 2) \nabla_a \nabla_b \Omega + g_{ab} \Box \Omega \right)$$

$$+ \Omega^{-2} \left(2(D - 2) (\nabla_a \Omega) \nabla_b \Omega + (3 - D) g_{ab} (\nabla_c \Omega) \nabla^c \Omega \right),$$

$$\tilde{R} = \Omega^{-2} \left(R + 2(1 - D) \Omega^{-1} \Box \Omega + (D - 1) (4 - D) \Omega^{-2} (\nabla_c \Omega) \nabla^c \Omega \right).$$

$$(4.5)$$

For D=2, these give

$$\tilde{J}^{a} = \Omega^{-2} J^{a} - (1 + 2\alpha) \Omega^{-2} \left(k^{a} \Box \ln \Omega + \nabla^{a} (k^{c} \nabla_{c} \ln \Omega) \right), \tag{4.6}$$

$$\tilde{\nabla}_{a} \tilde{J}^{a} = -(1 + 2\alpha) \Omega^{-2} \left(2K \Box \ln \Omega + k^{a} \nabla_{a} \Box \ln \Omega + \Box (k^{a} \nabla_{a} \ln \Omega) \right).$$

So one has $\tilde{\nabla}_a \tilde{J}^a = 0$ only when $\alpha = -1/2$, for which $J^a = G^{ab}k_b$, with G^{ab} the Einstein tensor. The $D \neq 2$ case can be obtained from the discussion on the Weyl scalings of a generic rank-n current F given in appendix A, by setting n = 1, of course.

4.2 n=2

For n=2, following the general case we find the conserved (obvious) current

$$J^{ab} = -\frac{1}{2}\nabla_c \nabla^{[a} k^{bc]}. \tag{4.7}$$

As may be seen from the following relations, proven in [2]

$$(D-3)G^{ac}K_c = \nabla_c \left(G^{b[c}k_b{}^{a]} + (D-2)\nabla^{[a}K^{c]} \right)$$
(4.8)

and

$$\nabla_a \check{\mathcal{K}}^{ab} = \nabla_a \hat{\mathcal{K}}^{ab} + \frac{1}{2} (D - 3) G^b{}_c K^c \,, \tag{4.9}$$

the case D=3 requires special attention.

When D=3, the Riemann tensor can be expressed in terms of the Ricci tensor and the curvature scalar only, and one finds $\check{\mathcal{K}}^{ab}=\hat{\mathcal{K}}^{ab}$ identically. This implies

$$\frac{1}{2}R^{abcd}k_{cd} + R_c^{[a}k^{b]c} = -\hat{\mathcal{K}}^{ab} = -\nabla^{[a}K^{b]} + \Box k^{ab}, \qquad (4.10)$$

where the last equality follows from (2.8) evaluated for n = 2. On the other hand, when D = 3 one has

$$\nabla_c \left(\hat{\mathcal{K}}^{ac} + \nabla^{[a} K^{c]} \right) = 0 \tag{4.11}$$

from (4.8). The discrepancy between (4.10) and (4.11) is accounted for by the obvious current (2.9) for n = 2 also in D = 3.

4.3 Plebański-Demiański metric

The Plebański-Demiański metric [7,8] is

$$ds^{2} = \Omega^{2} \left(-\frac{Q(r)(d\tau - p^{2}d\sigma)^{2}}{r^{2} + p^{2}} + \frac{P(p)(d\tau + r^{2}d\sigma)^{2}}{r^{2} + p^{2}} + \frac{r^{2} + p^{2}}{P(p)}dp^{2} + \frac{r^{2} + p^{2}}{Q(r)}dr^{2} \right), \quad (4.12)$$

with the metric functions Q(r) and P(p)

$$Q(r) = k + e^{2} + g^{2} - 2mr + \epsilon r^{2} - 2nr^{3} - (k + \Lambda/3) r^{4},$$

$$P(p) = k + 2np - \epsilon p^{2} + 2mp^{3} - (k + e^{2} + g^{2} + \Lambda/3) p^{4},$$
(4.13)

and the conformal factor

$$\Omega(p,r) = (1 - pr)^{-1}. \tag{4.14}$$

The Plebański-Demiański metric (4.12) (independent of the special choices (4.13), (4.14) and the vector potential, not displayed here) admits two rank-2 CKYTs

$$k = \Omega^{3} \left(p dr \wedge \left(d\tau - p^{2} d\sigma \right) + r dp \wedge \left(d\tau + r^{2} d\sigma \right) \right), \tag{4.15}$$

$$h = \Omega^{3} \left(r dr \wedge \left(p^{2} d\sigma - d\tau \right) + p dp \wedge \left(r^{2} d\sigma + d\tau \right) \right). \tag{4.16}$$

(See e.g. [1] for further details.)

For the obvious current F^{ac} (2.9), one may, in analogy to the discussion in subsection 4.4 (see also further details given in Sec. 3 of [1]), define the charge

$$Q^c := \int_{\Sigma_{\tau}} d^3x \sqrt{\gamma} \, n_a \, F^{ac} \,. \tag{4.17}$$

Identifying $\mathcal{P}^c := \sqrt{\gamma} n_a F^{ac}$ for convenience, we find that

$$\mathcal{P}^{p}(k) = -\Omega^{2}P(p)\frac{(p^{2}+r^{2})Q'(r)+2rP(p)-2rQ(r)}{2(p^{2}+r^{2})(P(p)-Q(r))} ,$$

$$\mathcal{P}^{r}(k) = -\Omega^{2}Q(r)\frac{(p^{2}+r^{2})P'(p)-2pP(p)+2pQ(r)}{2(p^{2}+r^{2})(P(p)-Q(r))} ,$$

$$\mathcal{P}^{p}(h) = -\Omega^{2}P(p)\frac{p^{2}(p^{2}+r^{2})Q'(r)+2\Omega\left(p^{5}+2p^{3}r^{2}+r^{3}\right)Q(r)-2r^{3}P(p)}{2(p^{2}+r^{2})(P(p)-Q(r))} ,$$

$$\mathcal{P}^{r}(h) = -\Omega^{2}Q(r)\frac{2p^{3}Q(r)-r^{2}(p^{2}+r^{2})P'(p)-2\Omega P(p)\left(p^{3}+2p^{2}r^{3}+r^{5}\right)}{2(p^{2}+r^{2})(P(p)-Q(r))} .$$

Here we have presented the derivation, but to evaluate the integrals (4.17) we need to know the ranges of the coordinates on the hypersurface Σ_{τ} . This is tantamount to finding the roots of the polynomials P(p) and Q(r), and choosing ranges which make P(p) > 0 and Q(r) > 0 and the integrals convergent. We plan to return to these questions in a forthcoming publication.

4.4 Kerr-Newman metric

Note from (4.8) that $\mathcal{J}^a := G^{ac}K_c$, where K_c is the contracted part of a possibly closed CKYT, is itself covariantly conserved with a manifest potential, thanks to

$$R^{ac}\nabla_a K_c = R^{ac}\nabla_{(a}K_{c)} = 0, \qquad (4.18)$$

shown in [2]. This is an example of a non-obvious current and can be used for defining a charge in the usual way, e.g. as described in Sec. 3 of [1]:

$$Q \equiv \int_{\Sigma_t} d^3 x \sqrt{\gamma} \, n_a \, \mathcal{J}^a \,. \tag{4.19}$$

Let us see how things go on the example of the Kerr-Newman metric

$$ds^{2} = -\frac{Q}{\rho^{2}} \left(dt - a \sin^{2}\theta \, d\varphi \right)^{2} + \frac{\rho^{2}}{Q} dr^{2} + \rho^{2} d\theta^{2} + \frac{\sin^{2}\theta}{\rho^{2}} \left(adt - (r^{2} + a^{2}) \, d\varphi \right)^{2}, \tag{4.20}$$

where

$$Q(r) = (r - r_{+})(r - r_{-}), \text{ with } r_{\pm} \equiv m \pm \sqrt{m^{2} - a^{2} - e^{2} - g^{2}},$$

 $\rho^{2} = r^{2} + a^{2} \cos^{2} \theta,$

with the rank-2 CKYT

$$k = r dr \wedge \left[dt - a \sin^2 \theta d\varphi \right] + a \cos \theta \sin \theta d\theta \wedge \left[a dt - \left(r^2 + a^2 \right) d\varphi \right], \tag{4.21}$$

that remarkably has $K^a = -(\partial_t)^a$. These give

$$Q = 2\pi \left(e^2 + g^2\right) \int_{r_{-}}^{\infty} dr \int_{0}^{\pi} \frac{d\theta \sin\theta \left(a^2 + 2(r - r_{-})(r - r_{+}) - a^2 \cos 2\theta\right)}{\left(a^2 + 2r^2 + a^2 \cos 2\theta\right) \left((r - r_{-})(r - r_{+}) - a^2 \sin^2\theta\right)}.$$
 (4.22)

This integral is equivalent to the integral found in Sec. 5.3 of [1]. It is convergent and finite, but as discussed in [1], even though the θ integral can be taken exactly, we were unable to evaluate the r-integral. See Sec. 5.3 of [1] for details.

The CKYT (4.21) is a closed CKYT, i.e. $\nabla_{[a}k_{bc]}=0$ identically, so it is not suitable for defining a conserved charge using the obvious current (2.9) for n=2.

5 Discussion

We have discussed the construction of conserved "currents" i.e. covariantly divergence-free tensors of rank-n constructed out of geometric tensors and linear in CKYTs also of rank-n. As seen

from our ansatz (3.10), we also require at most two covariant derivatives in the construction. For all $n \ge 2$, (in fact also for n = 1, see the second equation of (4.4)), the resulting conserved current reduces to

$$F^{c_1...c_n} = \Box k^{c_1...c_n} - n\nabla^{[c_1}K^{c_2...c_n]} = \nabla_a \left(\nabla^{[a}k^{c_1...c_n]}\right). \tag{5.1}$$

This motivates the following Conjecture/Theorem:

The only covariantly-conserved rank-n current in $D \neq 2$ dimensions for a rank-n CKYT that is local, geometric, linear in k and second-order in ∇s is uniquely given by "obvious" currents as in (5.1).

If we drop the requirement that the current has the same rank as the CKYT or that the current is second rank in derivatives, there are more possibilities. For n=2 we have discussed the Einstein current in terms of the contracted second rank CKYT in subsection 4.4. It is *third* order in ∇ s and defined for a rank-2 CKYT k_{ab} while the current is rank-1. Similarly, the Cotton current [1]

$$J^a = C^{abc} k_{bc} \,, \tag{5.2}$$

is again third order in ∇s and defined for a rank-2 CKYT k_{ab} while the current is rank-1.

In this context it is interesting to note that since the dual of a CKYT is again a CKYT, we may use the Cotton current with C^{abc} the Cotton tensor, defined for a rank-2 CKYT k_{bc} , to find a current for its rank-(D-2) dual

$$\mathcal{J}^{a} = C^{abc} \epsilon_{bcd_{1}...d_{D-2}} k^{d_{1}...d_{D-2}}.$$
(5.3)

In three dimensions this relates the Cotton current for a rank-2 CKYT to a current for its rank-1 dual defined with the dual of the Cotton tensor, the York tensor.

In [1] the Cotton current was lifted to supergravity in three dimensions. An interesting open problem is to find supergravity versions also of the obvious currents.

Acknowledgments

Ö.S. would like to thank D.O. Devecioğlu for help with xAct. The research of U.L. is supported in part by the 2236 Co-Funded Scheme2 (CoCirculation2) of TÜBİTAK (Project No:120C067)³.

Appendices

A CKYTs under Weyl scalings

In this appendix we show that the covariant-conservation of the obvious current (2.9) is left-invariant under Weyl scalings of the metric.

 $^{^{3}}$ However the entire responsibility for the publication is ours. The financial support received from TÜBİTAK does not mean that the content of the publication is approved in a scientific sense by TÜBİTAK.

Under conformal transformations $\tilde{g}_{ab} = \Omega^2 g_{ab}$, a generic rank-n CKYT k and its rank-(n-1) companion K (2.1) transform as

$$\tilde{k}^{a_1...a_n} = \Omega^{1-n} k^{a_1...a_n} \quad \text{and} \quad \tilde{K}_{a_1...a_{n-1}} = \Omega^{n-1} K_{a_1...a_{n-1}} + \Omega^{n-2} (\nabla_c \Omega) k^c_{a_1...a_{n-1}} \,. \tag{A.1}$$

Note that for all $n \neq 2$, the second part on \tilde{K} differs significantly from the formulae given in appendix A of [6].

With these, we find

$$\tilde{\nabla}^{[c}\tilde{k}^{a_1...a_n]} = \Omega^{-(n+1)}\nabla^{[c}k^{a_1...a_n]} + (n+1)\Omega^{-(n+2)}(\nabla^{[c}\Omega)k^{a_1...a_n]}$$
(A.2)

leading to the transformation rule

$$\tilde{F}^{a_1...a_n} = \Omega^{-(n+1)} F^{a_1...a_n} + (D-n-1) \Omega^{-(n+2)} (\nabla_c \Omega) \nabla^{[c} k^{a_1...a_n]}$$

$$+ (n+1) \Omega^{-(n+2)} \left(\nabla_c \left((\nabla^{[c} \Omega) k^{a_1...a_n]} \right) + (D-n-2) \Omega^{-1} (\nabla_c \Omega) (\nabla^{[c} \Omega) k^{a_1...a_n]} \right)$$
(A.3)

for a generic rank-n current F. Since for any skew-symmetric rank-(n+1) tensor \tilde{T} , one has

$$[\tilde{\nabla}_{a_1}, \tilde{\nabla}_c]\tilde{T}^{ca_1...a_n} = 2\tilde{\nabla}_{a_1}\tilde{\nabla}_c\tilde{T}^{ca_1...a_n} = 0, \qquad (A.4)$$

which can be shown by the expansion of the commutator, the symmetries of the curvature tensors and the first Bianchi identity, it follows straightforwardly that $\tilde{\nabla}_{a_1}\tilde{F}^{a_1...a_n}=0$ by the identifications $\tilde{T}^{ca_1...a_n}=\tilde{\nabla}^{[c}\tilde{k}^{a_1...a_n]}$ and $\tilde{F}^{a_1...a_n}=\tilde{\nabla}_c\tilde{T}^{ca_1...a_n}$. So the covariant-conservation of the obvious current F is left invariant under conformal transformations.

B Killing-Yano Cotton current

In [1], we showed that the Killing-Yano Cotton current $J^a \equiv C^{abc}k_{bc}$, constructed out of a rank-2 CKYT k and the Cotton tensor C, is covariantly conserved. It is known that the Cotton tensor transforms as

$$\tilde{C}_{abc} = C_{abc} - (D - 2)(\nabla_d \ln \Omega) W^d_{abc}$$
(B.1)

under conformal transformations $\tilde{g}_{ab} = \Omega^2 g_{ab}$. Meanwhile, one has⁴

$$\tilde{k}_{ab} = \Omega^3 k_{ab}$$
 and $\tilde{K}_c = \Omega K_c + (\nabla_a \Omega) k^a_c$. (B.2)

These give

$$\tilde{J}^a = \Omega^{-3} \left(J^a - (D-2)(\nabla_d \ln \Omega) W^{da}{}_{bc} k^{bc} \right). \tag{B.3}$$

Using these and the property $(D-3)C_{bdc} = (D-2)\nabla_a W^a{}_{bcd}$, it is straightforward to show that

$$\tilde{\nabla}_a \tilde{J}^a = \nabla_a \tilde{J}^a + D(\nabla_a \ln \Omega) \tilde{J}^a = 0,$$

independent of the dimension D.

⁴Here we correct an error in [1], where a factor of (D-1) was inadvertently forgotten in the second term on the right-hand-side of the second equation in eqn. (2.12) in that paper, with no effect on the ensuing discussion.

C Lie derivative of curvature tensors

While not used in this article, we still feel that this section fills a much needed gap.

Let us study the consequences of (2.2) for n=1 from a geometrical perspective. First note that one has

$$[\nabla_a, \nabla_b] \nabla_c k_d = R_{abc}{}^i \nabla_i k_d + R_{abd}{}^i \nabla_c k_i \tag{C.1}$$

from the definition of the commutator. Meanwhile, the left hand side of this expression can also be calculated by employing (2.2)

$$[\nabla_a, \nabla_b] \nabla_c k_d = \nabla_a \left(R^i{}_{bcd} k_i - 2g_{b[c} \nabla_{d]} K + g_{cd} \nabla_b K \right) - \nabla_b \left(R^i{}_{acd} k_i - 2g_{a[c} \nabla_{d]} K + g_{cd} \nabla_a K \right)$$
$$= k^i \nabla_i R_{abcd} + R^i{}_{bcd} \nabla_a k_i + R_a{}^i{}_{cd} \nabla_b k_i - 2g_{b[c} \nabla_{d]} \nabla_a K + 2g_{a[c} \nabla_{d]} \nabla_b K , \quad (C.2)$$

using the second Bianchi identity and $\nabla_{[a}\nabla_{b]}K = 0$. Equating the right hand sides of (C.1) and (C.2), using $\nabla_{(a}k_{b)} = g_{ab}K$ from (2.1), and recalling the properties of the Lie derivative

$$\mathcal{L}_k R_{abcd} = k^i \nabla_i R_{abcd} + R_{ibcd} \nabla_a k^i + R_{aicd} \nabla_b k^i + R_{abid} \nabla_c k^i + R_{abci} \nabla_d k^i ,$$

one finds

$$\mathcal{L}_k R_{abcd} = 2K R_{abcd} + 2g_{b[c} \nabla_{d]} \nabla_a K - 2g_{a[c} \nabla_{d]} \nabla_b K. \tag{C.3}$$

Since $\mathcal{L}_k g_{ab} = 2Kg_{ab}$, and hence $\mathcal{L}_k g^{ab} = -2Kg^{ab}$, it easily follows that

$$\mathcal{L}_k R_{ac} = g^{bd} \mathcal{L}_k R_{abcd} - 2K R_{ac} = (2 - D) \nabla_a \nabla_c K - g_{ac} \Box K, \qquad (C.4)$$

$$\mathcal{L}_k R = g^{ac} \mathcal{L}_k R_{ac} - 2KR = 2(1-D)\Box K - 2KR. \tag{C.5}$$

Finally, from

$$S_{ab} \equiv \frac{1}{(D-2)} \left(R_{ab} - \frac{1}{2(D-1)} R g_{ab} \right) ,$$

$$R_{ab} = (D-2) S_{ab} + S g_{ab} \quad \text{with} \quad S \equiv g^{ab} S_{ab} \text{ so that } R = 2(D-1) S ,$$

$$R^{ab}_{cd} = W^{ab}_{cd} + 4 \delta^{[a}_{[c} S^{b]}_{d]} \quad \text{and} \quad G_{ab} = (D-2) \left(S_{ab} - S g_{ab} \right) ,$$
(C.6)

it follows that

$$\mathcal{L}_k S_{ac} = -\nabla_a \nabla_c K, \qquad \mathcal{L}_k W_{abcd} = 2K W_{abcd}, \qquad \mathcal{L}_k W_{abc}{}^d = 0.$$
 (C.7)

Since one has $\tilde{W}_{abc}{}^d = W_{abc}{}^d$ under a conformal transformation $\tilde{g}_{ab} = \Omega^2 g_{ab}$, the last inequality in (C.7) is on a par with the latter invariance.

References

[1] U. Lindström and Ö. Sarıoğlu, "Killing-Yano Cotton Currents," *JHEP* **03** (2022) 029; arXiv:2110.03470 [hep-th].

- [2] U. Lindström and Ö. Sarıoğlu, "New currents with Killing-Yano tensors," Class. Quant. Grav. 38 no.19, 195011 (2021); arXiv:2104.12451 [hep-th].
- [3] U. Lindström and Ö. Sarıoğlu, "Tensionless strings and Killing(-Yano) tensors," *Phys. Lett. B* **829** (2022), 137088; arXiv:2202.06542 [hep-th].
- [4] D. Kastor and J. Traschen, "Conserved gravitational charges from Yano tensors," JHEP 08 (2004) 045; arXiv:hep-th/0406052 [hep-th].
- [5] U. Lindström and Ö. Sarıoğlu, "Uses of Killing and Killing-Yano Tensors," arXiv:2202.07226 [hep-th].
 - P. S. Howe and U. Lindström, "Some remarks on (super)-conformal Killing-Yano tensors," *JHEP* 11 (2018) 049; arXiv:1808.00583 [hep-th].
 - V. Frolov, P. Krtous and D. Kubiznak, "Black holes, hidden symmetries, and complete integrability," *Living Rev. Rel.* **20**, no.1, 6 (2017); arXiv:1705.05482 [gr-qc].
 - C. Batista, "Integrability Conditions for Killing-Yano Tensors and Conformal Killing-Yano Tensors," *Phys. Rev.* D91, 024013 (2015); arXiv:1406.3069 [gr-qc].
 - M. Cariglia, "Hidden Symmetries of the Dirac Equation in Curved Space-Time," Springer Proc. Phys. 157 (2014) 25; arXiv:1209.6406 [gr-qc].
 - O. P. Santillan, "Hidden symmetries and supergravity solutions," J. Math. Phys. 53 (2012) 043509; arXiv:1108.0149 [hep-th].
 - G. W. Gibbons, R. H. Rietdijk and J. W. van Holten, "Susy in the sky," *Nucl. Phys. B* **404** (1993) 42-64; arXiv:hep-th/9303112 [hep-th].
 - S. Tachibana and T. Kashiwada, "On the integrability of Killing-Yano's equation," *J. Math. Soc. Japan* **21** (1969) No.2, 259-265.
 - S. Tachibana, "On conformal Killing tensor in a Riemannian space", *Tôhoku Math. Journ.* **21** (1969), 56-64.:
- [6] Y. Chervonyi and O. Lunin, "Killing(-Yano) Tensors in String Theory," JHEP 09 (2015) 182; arXiv:1505.06154 [hep-th].
- [7] J. F. Plebanski and M. Demianski, "Rotating, charged, and uniformly accelerating mass in general relativity," *Annals Phys.* **98**, 98-127 (1976).
- [8] D. Kubiznak and P. Krtous, "On conformal Killing-Yano tensors for Plebanski-Demianski family of solutions," *Phys. Rev. D* **76**, 084036 (2007); arXiv:0707.0409 [gr-qc].