

# COHERENT ORIENTATIONS IN SYMPLECTIC FIELD THEORY REVISITED

ERKAO BAO

ABSTRACT. In symplectic field theory (SFT), the moduli spaces of  $J$ -holomorphic curves can be oriented coherently (compatible with gluing). In this note, we correct the signs involved in the generating function  $\mathbf{H}$  in SFT so that the master equation  $\mathbf{H} \circ \mathbf{H} = 0$  holds assuming transversality. The orientation convention that we use is consistent with that of Hutchings-Taubes from [HT09].

## 1. INTRODUCTION

Symplectic Field Theory (SFT) was introduced by Eliashberg, Hofer and Givental in [EGH00], and it is a generalization of the Gromov-Witten invariants in the spirit of topological field theory. SFT packs into a potential function  $\mathbf{H}$  a signed count of elements of moduli spaces (with virtual dimension zero) of  $J$ -holomorphic curves in the symplectization of a contact manifold. The potential function  $\mathbf{H}$  satisfies the master equation  $\mathbf{H} \circ \mathbf{H} = 0$ . There are different choices to orient the moduli spaces of  $J$ -holomorphic curves, such as those of Bourgeois-Mohnke [BM04] and Hutchings-Taubes [HT09]. However, the algebraic setup of  $\mathbf{H}$  in [EGH00] is not compatible with either of the two orientation conventions. We correct the signs of  $\mathbf{H}$  so that they match with the orientation convention of [HT09]<sup>1</sup>. We reformulate and generalize the coherent orientation in [HT09] so that the proof of  $\mathbf{H} \circ \mathbf{H} = 0$  becomes obvious assuming transversality.

In Section 2 we recall the coherent orientation for Cauchy-Riemann tuples, and we reformulate the coherent orientation used in [HT09]. In Section 3 we orient moduli spaces of  $J$ -holomorphic curves. In Section 4 we correct the definition of  $\mathbf{H}$  and prove  $\mathbf{H} \circ \mathbf{H} = 0$ .

## 2. COHERENT ORIENTATIONS OF CAUCHY-RIEMANN TUPLES

### 2.1. Cauchy-Riemann tuples.

**Definition 2.1.** A decorated Riemann surface with  $(k_+, k_-)$  marked points is a tuple  $(\Sigma, j, \mathbf{p}, \mathbf{r})$  such that

- (1)  $(\Sigma, j)$  is a possibly *disconnected* Riemann surface,

---

2010 *Mathematics Subject Classification.* Primary 53D10; Secondary 53D40.

*Key words and phrases.* coherent orientation, contact structure, contact homology, symplectic field theory.

<sup>1</sup>It is likely that a different correction of  $\mathbf{H}$  can match with the orientation convention of [Bou02] resulting in an equivalent SFT.

- (2)  $\mathbf{p} = (\mathbf{p}^+, \mathbf{p}^-)$  and  $\mathbf{p}^\pm = (p_1^\pm, \dots, p_{k_\pm}^\pm)$  is a (ordered) tuple of points on  $\Sigma$ ,
- (3)  $\mathbf{r} = (\mathbf{r}^+, \mathbf{r}^-)$  and  $\mathbf{r}^\pm = (r_1^\pm, \dots, r_{k_\pm}^\pm)$  is a tuple of rays on  $\Sigma$  at  $\mathbf{p}$ , i.e.,  $r_i^\pm \in T_{p_i^\pm} \Sigma - 0$ .

**Definition 2.2** ([Sch95]). A smooth loop of symmetric matrices

$$S \in C^\infty(S^1, \text{End}(\mathbb{R}^{2n-2}))$$

is called *admissible* if the ordinary differential equation

$$\begin{aligned} \dot{x}(t) &= J_0 S(t) x(t), \\ x : S^1 &\rightarrow \mathbb{R}^{2n-2} \end{aligned}$$

has only the zero solution, where  $J_0$  is the standard complex structure on  $\mathbb{R}^{2n-2}$ .

**Definition 2.3.** A *Cauchy-Riemann tuple* (CR tuple for short) is a tuple

$$\mathcal{T} = (\Sigma, j, \mathbf{p}, \mathbf{r}, E, J, \psi = \{\psi_i^\pm\}_i, \mathcal{S} = \{S_i^\pm\}_i)$$

consists of the following data:

- (1) a decorated Riemann surface  $(\Sigma, j, \mathbf{p}, \mathbf{r})$  with  $(k_+, k_-)$  marked points,
  - (2) a complex vector bundle  $(E, J)$  over  $\dot{\Sigma} = \Sigma - \mathbf{p}$ , such that for each  $p_i^\pm$ , there are
    - (a) a neighborhood  $\mathcal{U}_{p_i^\pm}$  of  $p_i^\pm$  in  $\Sigma$  and a trivialization  $\psi_i^\pm$  of
- (2.1.1) 
$$(E, J)|_{\dot{\mathcal{U}}_{p_i^\pm}} \simeq (\mathbb{R}^{2n} \times \dot{\mathcal{U}}_{p_i^\pm}, J_0),$$
- where  $\dot{\mathcal{U}}_{p_i^\pm} = \mathcal{U}_{p_i^\pm} - p_i^\pm$ , and  $J_0$  is the standard complex structure on  $\mathbb{R}^{2n}$ , and
- (b) an admissible  $S_i \in C^\infty(S^1, \text{End}(\mathbb{R}^{2n-2}))$ .

Given an admissible loop  $S$ , we obtain a path of symplectic matrices  $B(t) \in \text{Symp}(2n-2)$  that solves  $\dot{B}(t) = J_0 S(t) B(t)$  and  $B(0) = \text{Id}$ . We define the Conley-Zehnder index of  $S$  by  $\mu_{\text{CZ}}(S) = \mu(\{B(t)\}_t)$ , where  $\mu$  is the Maslov-index, and we grade  $S$  over  $\mathbb{Z}_2$  by  $|S| = \mu_{\text{CZ}}(S) + (n-1) \pmod{2}$ .

**Lemma 2.4** ([BM04]). *Given an admissible  $S \in C^\infty(S^1, \text{End}(\mathbb{R}^{2n-2}))$ , the associated operator  $A = J_0 \frac{\partial}{\partial t} + S : W^{1,p}(S^1, \mathbb{R}^{2n-2}) \rightarrow L^p(S^1, \mathbb{R}^{2n-2})$  has discrete spectrum  $\sigma(A) \subset \mathbb{R} - \{0\}$ .*

For an admissible loop  $S$ , we define  $\lambda_S = \min\{-\lambda_{-1}, \lambda_1\} > 0$  where  $\lambda_1$  is the smallest positive eigenvalue of  $J_0 \frac{\partial}{\partial t} + S$  and  $\lambda_{-1}$  is the largest negative eigenvalue of  $J_0 \frac{\partial}{\partial t} + S$ .

Fix  $k \in \mathbb{Z}^{\geq 0}$ , and  $p > 1$  such that  $kp > 2$ .

**Definition 2.5** (Cauchy-Riemann operators). For a CR tuple  $\mathcal{T}$ , we define  $\mathfrak{D}(\mathcal{T})$  to be the topological space of linear operators, called *Cauchy-Riemann operators* (CR operators for short)

$$(2.1.2) \quad L : W_\delta^{k,p}(\dot{\Sigma}, E) \oplus \mathcal{V} \rightarrow W_\delta^{k-1,p}(\dot{\Sigma}, \wedge^{0,1} T^* \dot{\Sigma} \otimes_{\mathbb{C}} E),$$

such that

- (1)  $0 < \delta < \min_i \lambda_{S_i}$ , and  $W_\delta^{k,p}(\cdot)$  and  $W_\delta^{k-1,p}(\cdot)$  are the weighted Sobolev spaces;
- (2)  $\mathcal{V} = (\oplus_{i=1}^{k_-} \mathcal{V}_i^-) \oplus (\oplus_{i=1}^{k_+} \mathcal{V}_i^+)$ , where  $\mathcal{V}_i^\pm = \text{span}\{\beta_i^\pm \cdot e_1, \beta_i^\pm \cdot e_2\}$ ,  $\{e_1, \dots, e_n\}$  is the standard basis of  $\mathbb{R}^2 \oplus \mathbb{R}^{2n-2}$ ,  $0 \leq \beta_i^\pm \leq 1$  is a bump function that is supported in  $\mathcal{U}_{p_i^\pm}$  and satisfies  $\beta_i^\pm(p_i^\pm) = 1$ , and hence  $\beta_i^\pm \cdot e_1$  and  $\beta_i^\pm \cdot e_2$  are sections of  $E$ ;
- (3)  $L$  is a real Cauchy-Riemann operator (Appendix C.1 in [MS12]), i.e.,  $L = L_0 + L_1$ , where  $L_1$  is a complex Cauchy-Riemann operator, that is,  $L_1(f\eta) = f(L_1\eta) + (\bar{\partial}f)\eta$  for any  $f \in C_c^\infty(\dot{\Sigma})$  and  $\eta \in W_\delta^{k,p}(\dot{\Sigma}, E) \oplus \mathcal{V}$  and  $L_0 \in W^{k-1,p}(\dot{\Sigma}, \wedge^{0,1}T^*\dot{\Sigma} \otimes_{\mathbb{R}} \text{End}_{\mathbb{R}}(E))$ ;
- (4) with respect to the coordinate  $h_i^\pm$  and the trivialization  $\psi_i^\pm$ , we require that for any  $\eta \in W_\delta^{k,p}(\dot{\Sigma}, E) \oplus \mathcal{V}$  with support in  $\dot{\mathcal{U}}_{p_i^\pm}$ ,

$$L\eta(s, t) = \left( \frac{\partial \eta}{\partial s} + \tilde{J}_i^\pm(s, t) \frac{\partial \eta}{\partial t} + \tilde{S}_i^\pm(s, t) \eta \right) \otimes (ds - \sqrt{-1}dt),$$

where  $\tilde{J}_i^\pm(s, t)$  is a complex structure on  $\mathbb{R}^{2n}$ ,  $\tilde{S}_i^\pm(s, t) \in \text{End}(\mathbb{R}^{2n})$ , and there exists some constants  $C_i^\pm > 0$  such that for all  $\beta = (\beta_1, \beta_2) \in \mathbb{Z}^{\geq 0} \times \mathbb{Z}^{\geq 0}$  with  $\beta_1 + \beta_2 \leq k$  one has

$$|\partial^\beta(\tilde{J}_i^\pm(s, t) - \hat{J}_0)| \leq C_i^\pm e^{\mp \frac{1}{2}\lambda_{S_i^\pm} s},$$

$$|\partial^\beta(\tilde{S}_i^\pm(s, t) - \hat{S}_i^\pm(t))| \leq C_i^\pm e^{\mp \frac{1}{2}\lambda_{S_i^\pm} s},$$

where  $\hat{J}_0$  is the standard complex structure on  $\mathbb{R}^{2n}$ ,

$$\hat{S}_i^\pm = \begin{pmatrix} 0 & & \\ & 0 & \\ & & S_i^\pm \end{pmatrix} \quad \text{and} \quad \partial^\beta = \partial_s^{\beta_1} \partial_t^{\beta_2}.$$

**Proposition 2.6** ([Bou02; BM04]). *Any  $L \in \mathcal{D}(\mathcal{T})$  is Fredholm, and its Fredholm index is*

$$\text{ind } L = \sum_{i=1}^{k_+} \mu_{\text{CZ}}(S_i^+) - \sum_{i=1}^{k_-} \mu_{\text{CZ}}(S_i^-) - (n-1)(k_- + k_+) + 2c_1(E) + n(2-2g).$$

We define the mod 2 indexes

$$\text{ind}^\pm \mathcal{T} := \sum_{i=1}^{k_\pm} |S_i^\pm| \pmod{2},$$

and

$$\text{ind } \mathcal{T} := \text{ind}^+ \mathcal{T} + \text{ind}^- \mathcal{T} = \text{ind } L \pmod{2}.$$

For any CR operator  $L$  we define its determinant line by

$$\det L = \wedge^{\text{top}} \ker L \otimes \wedge^{\text{top}} (\text{coker } L)^*.$$

An orientation  $o(L)$  of  $L$  is a choice of non-zero vector in  $\det L$  up to a positive scalar multiplication.

**Example 2.7.** We call a CR tuple  $\mathcal{T}$  *trivial*, if  $k_+ = k_- = 1$  and  $S_1^+ = S_1^-$ . For a trivial  $\mathcal{T}$ ,  $L \in \mathfrak{D}(\mathcal{T})$  is said to be *trivial*, if  $J_1^+ = J_1^-$ , and  $\tilde{J}_1^\pm$  and  $\tilde{S}_1^\pm$  are independent of  $s$ . In this case,  $\text{ind } L = 2$ , and  $\text{coker } L = \{0\}$ , and  $\ker L$  is spanned by translation and rotation.

**Example 2.8.** In the case when  $k_- = 0 = k_+$ , the operator  $L$  is homotopic to a complex CR operator, whose  $\ker$  and  $\text{coker}$  are complex vector spaces. We have the canonical orientation  $o_{\text{can}}(L)$  of  $\det L$  coming from the complex structure.

It is convenient to have the following lemma:

**Lemma 2.9** ([FOOO09] p. 676). *Let  $V, W$  be Banach spaces, and  $\phi : V \rightarrow W$  a linear Fredholm operator. Let  $F \subset W$  be a finite dimensional subspace of  $W$  such that  $W = \text{im}(\phi) + F$ . Then there is an isomorphism*

$$\det \phi \simeq \wedge^{\text{top}}(\phi^{-1}(F)) \otimes \wedge^{\text{top}} F^*,$$

which is natural up to a positive constant. More precisely, suppose  $\phi^{-1}(F) = \ker \phi \oplus H \subseteq V$ ,  $\{e_1, \dots, e_n\}$  is a basis of  $\ker \phi$ ,  $\{h_1, \dots, h_m\}$  is a basis of  $H$ , and  $\{v_1, \dots, v_\ell, \phi(h_1), \dots, \phi(h_m)\}$  is a basis of  $F$ . Then the isomorphism is given by

$$\begin{aligned} e_1 \wedge \dots \wedge e_n \otimes v_\ell^* \wedge \dots \wedge v_1^* \mapsto & e_1 \wedge \dots \wedge e_n \wedge h_1 \wedge \dots \wedge h_m \\ & \otimes \phi(h_m)^* \wedge \dots \wedge \phi(h_1)^* \wedge v_\ell^* \wedge \dots \wedge v_1^*. \end{aligned}$$

Given a continuous family of Fredholm operators  $\{\phi_\tau\}_{\tau \in [0,1]}$ , we can find a finite dimensional space  $F \subset W$  and a subspace  $U \subset W$  such that  $W = F \oplus U$  and  $W = \text{im}(\phi_\tau) + F$ . Let  $\pi_U : W \rightarrow U$  be the projection map. Then the map  $\pi_U \circ \phi_\tau : V \rightarrow U$  is surjective, and  $\ker(\pi_U \circ \phi_\tau) = \phi_\tau^{-1}(F)$ . By Lemma 2.9, we have  $\det \phi_\tau \simeq \wedge^{\text{top}}(\phi_\tau^{-1}(F)) \otimes \wedge^{\text{top}} F^* \simeq \wedge^{\text{top}} \ker(\pi_U \circ \phi_\tau) \otimes \wedge^{\text{top}} F^*$ . Since both  $\ker(\pi_U \circ \phi_\tau)$  and  $F^*$  form vector bundles over  $[0, 1]$ ,  $\det \phi_\tau$  forms a line bundle over  $[0, 1]$ . More generally, a homotopy of CR tuples  $\{L_\tau\}_{0 \leq \tau \leq 1}$  induces an isomorphism  $\det L_0 \simeq \det L_1$ .

Since the space  $\mathfrak{D}(\mathcal{T})$  is contractible,  $\det L$  and  $\det L'$  are canonically isomorphic for any  $L, L' \in \mathfrak{D}(\mathcal{T})$ . For this reason, we also write  $\det \mathcal{T}$ ,  $o(\mathcal{T})$  and  $o_{\text{can}}(\mathcal{T})$ .

**2.2. Disjoint union.** The disjoint union  $\mathcal{T} \amalg \mathcal{T}'$  of two Cauchy Riemann tuples  $\mathcal{T}$  and  $\mathcal{T}'$  are defined in the obvious way. The punctures of  $\dot{\Sigma} \amalg \dot{\Sigma}'$  are ordered in the way that the punctures of  $\dot{\Sigma}$  comes before those of  $\dot{\Sigma}'$ , and the relative orders of the punctures of  $\dot{\Sigma}$  and  $\dot{\Sigma}'$  are preserved respectively. For any  $L \in \mathfrak{D}(\mathcal{T})$  and  $L' \in \mathfrak{D}(\mathcal{T}')$ , the disjoint union map induces an isomorphism

$$(2.2.1) \quad \det L \otimes \det L' \rightarrow \det(L \amalg L'),$$

where  $L \amalg L' \in \mathfrak{D}(\mathcal{T} \amalg \mathcal{T}')$  is defined as follows: Let  $\{e_1, \dots, e_n\}$  be a basis of  $\ker L$ ,  $\{f_1, \dots, f_m\}$  be a basis of  $\text{coker } L$ ,  $\{e'_1, \dots, e'_{n'}\}$  be a basis of  $\ker L'$ ,

$\{f'_1, \dots, f'_{m'}\}$  be a basis of  $\text{coker } L'$ . Then the isomorphism is given by

$$\begin{aligned} e_1 \wedge \dots \wedge e_n \otimes f_m^* \wedge \dots \wedge f_1^* \\ \otimes e'_1 \wedge \dots \wedge e'_{n'} \otimes f'_{m'} \wedge \dots \wedge f'_1 \mapsto e_1 \wedge \dots \wedge e_n \wedge \otimes e'_1 \wedge \dots \wedge e'_{n'} \\ \otimes f'_{m'} \wedge \dots \wedge f'_1 \otimes f_m^* \wedge \dots \wedge f_1^* \end{aligned}$$

In the case when both  $L$  and  $L'$  are surjective,  $L \amalg L'$  is also surjective, and the isomorphism simplifies to

$$e_1 \wedge \dots \wedge e_n \otimes e'_1 \wedge \dots \wedge e'_{n'} \mapsto e_1 \wedge \dots \wedge e_n \wedge e'_1 \wedge \dots \wedge e'_{n'}.$$

The isomorphism 2.2.1 is continuous with respect to the homotopy of CR operators  $L$  and  $L'$ , hence induces a map on CR tuples  $\mathcal{T}$  and  $\mathcal{T}'$ :

$$\det \mathcal{T} \otimes \det \mathcal{T}' \rightarrow \det(\mathcal{T} \amalg \mathcal{T}')$$

and we denote the image of  $v \otimes v'$  by  $v \amalg v'$ .

Since  $\det \mathcal{T}$  by definition is independent of the ordering of  $\mathcal{T}$ ,  $\det(\mathcal{T} \amalg \mathcal{T}') = \det(\mathcal{T}' \amalg \mathcal{T})$ . The following lemma is clear from the above definition.

**Lemma 2.10.**

$$v \amalg v' = (-1)^{\text{ind } \mathcal{T} + \text{ind } \mathcal{T}'} v' \amalg v,$$

for any  $v \in \det \mathcal{T}$  and  $v' \in \det \mathcal{T}'$ .

**2.3. Gluing.** Given two CR tuples  $\mathcal{T}$  and  $\mathcal{T}'$ , and  $L \in \mathfrak{D}(\mathcal{T})$  and  $L' \in \mathfrak{D}(\mathcal{T}')$ , a positive integer  $\tau \leq \min(k_-, k'_+)$ , and a sufficiently large gluing parameter  $R > 0$ , if the first  $\tau$  negative ends of  $\dot{\Sigma}$  matches the last  $\tau$  positive ends of  $\dot{\Sigma}'$ , i.e., for  $1 \leq i \leq \tau$ ,

$$S_i^- = S'_{k'_+ - \tau + i},$$

then we can glue  $\mathcal{T}$  and  $\mathcal{T}'$  to obtain a new CR tuple  $\mathcal{T}''$ , and glue  $L$  and  $L'$  to obtain  $L'' \in \mathfrak{D}(\mathcal{T}'')$ . The gluing construction is straightforward: for each  $1 \leq i \leq \tau$  we ‘‘chop off’’ the end  $(-\infty, -2R] \times S^1$  from  $\dot{\Sigma}$  around  $p_i^-$ , and the end  $[2R, \infty) \times S^1$  from  $\dot{\Sigma}'$ , identify the regions  $[-2R, -R] \times S^1 \subset \dot{\Sigma}$  and  $[R, 2R] \times S^1 \subset \dot{\Sigma}'$ , and the bundles  $E$  and  $E'$  above these regions, and interpolate  $L$  and  $L'$  over the identified regions. We now explain how  $\mathbf{p}''$  is ordered. The positive (negative) marked points of  $\mathbf{p}'$  are ordered before the positive (negative) marked points of  $\mathbf{p}$ , more precisely,  $\mathbf{p}''_+ = (p'_+{}^1, \dots, p'_+{}^{k'_+ - \tau}, p_+{}^1, \dots, p_+{}^{k_+})$  and  $\mathbf{p}''_- = (p_-{}^1, \dots, p_-{}^{k_-}, p'_-{}^{\tau+1}, \dots, p'_-{}^{k'_-})$ . We denote  $\mathcal{T}'' = \mathcal{T} \sharp_{\tau, R} \mathcal{T}'$  and  $L'' = L \sharp_{\tau, R} L'$ .

We follow the *complete gluing* convention in [HT09] by restricting the gluing to the case when  $k_- = k'_+ = \tau$ , i.e., all the negative punctures of  $\dot{\Sigma}$  match with all the positive punctures of  $\dot{\Sigma}'$  and we glue along all of them. For complete gluing we write  $\mathcal{T}'' = \mathcal{T} \sharp_R \mathcal{T}'$  and  $L'' = L \sharp_R L'$ . To get a non-complete gluing from complete gluing, one can add a few trivial CR tuples before gluing in the obvious way. See Figure (1).

*From now on, all gluings are complete gluings unless specified.*




FIGURE 1. Taking disjoint union with trivial cylinders before gluing.

**Lemma 2.11** (Corollary 7 in [BM04], Lemma 9.6 in [HT09]). *The gluing map induces an isomorphism*

$$\det L \otimes \det L' \rightarrow \det(L \sharp_R L'),$$

*which is continuous with respect to the homotopies of CR operators  $L$  and  $L'$  and the gluing parameter  $R$ .*

For this reason we write  $\sharp$  instead of  $\sharp_R$  meaning gluing with some unspecified gluing parameter. The gluing map induces an isomorphism

$$\det \mathcal{T} \otimes \det \mathcal{T}' \rightarrow \det(\mathcal{T} \sharp \mathcal{T}'),$$

and we denote the image of  $v \otimes v'$  under the isomorphism by  $v \sharp v'$ .

**Lemma 2.12** (Lemma 9.7 in [HT09]). *The gluing operation of CR tuples is associative up to homotopy, and so is the induced operation on determinants:*

$$(v \sharp v') \sharp v'' = v \sharp (v' \sharp v'')$$

for any  $v \in \det \mathcal{T}$ ,  $v' \in \det \mathcal{T}'$ , and  $v'' \in \det \mathcal{T}''$ .

**2.4. Choices.** To assign an orientation to all CR tuples, we follow the approach in [BM04]:

- (1) for each loop of admissible symmetric matrices  $S$ ,
  - (a) we pick a CR tuple  $\mathcal{T}_S^+$  with one positive puncture and zero negative punctures and the associated admissible loop being  $S$ ,
  - (b) we choose an orientation  $o(\mathcal{T}_S^+)$ ,
  - (c) we pick a CR tuple  $\mathcal{T}_S^-$  with zero positive puncture and one negative puncture and the associated admissible loop being  $S$ ,
  - (d) we determine the orientation  $o(\mathcal{T}_S^-)$  by gluing  $\mathcal{T}_S^+$  and  $\mathcal{T}_S^-$ . More precisely, the equation

$$o(\mathcal{T}_S^+) \sharp o(\mathcal{T}_S^-) = o_{\text{can}}(\mathcal{T}_S^+ \sharp \mathcal{T}_S^-)$$

uniquely determines the  $o(\mathcal{T}_S^-)$ .

We call  $\mathcal{T}_{\mathcal{S}^\pm}$  the *positive (negative) capping CR tuple*, and  $o(\mathcal{T}_{\mathcal{S}^\pm})$  the *positive (negative) capping orientation*.

- (2) for an arbitrary CR tuple  $\mathcal{T}$ , let  $\mathcal{T}_{\mathcal{S}^\pm}$  be the CR tuple that is obtained by taking iterated disjoint union of  $\mathcal{T}_{\mathcal{S}_i^\pm}$  in the  $i$  increasing order, where  $\mathcal{S}_i^\pm$  is the admissible loop of symmetric matrices at the  $i$ -th positive (negative) puncture. Then  $o(\mathcal{T}_{\mathcal{S}^\pm})$  is determined by

$$(2.4.1) \quad o(\mathcal{T}_{\mathcal{S}^\pm}) = \epsilon^{\mathbb{I}} \cdot o(\mathcal{T}_{\mathcal{S}_1^\pm}) \prod \cdots \prod o(\mathcal{T}_{\mathcal{S}_{k^\pm}^\pm}),$$

where  $\epsilon^{\mathbb{I}} \in \{\pm 1\}$  to be chosen.

- (3) we glue  $\mathcal{T}_{\mathcal{S}^+}$ ,  $\mathcal{T}$ , and  $\mathcal{T}_{\mathcal{S}^-}$  to get a CR tuple with no punctures, and  $o(\mathcal{T})$  is determined by

$$(2.4.2) \quad o(\mathcal{T}_{\mathcal{S}^+}) \# o(\mathcal{T}) \# o(\mathcal{T}_{\mathcal{S}^-}) = \epsilon^\# o_{\text{can}}(\mathcal{T}_{\mathcal{S}^+} \# \mathcal{T} \# \mathcal{T}_{\mathcal{S}^-}),$$

where  $\epsilon^\# \in \{\pm 1\}$  to be chosen.

The choices of  $\epsilon^{\mathbb{I}}$  and  $\epsilon^\#$  are different in [BM04] and  $o_{ht}$  in [HT09], and different choices result in different properties of the coherent orientations. We list the properties and the choices of  $\epsilon^{\mathbb{I}}$  and  $\epsilon^\#$  of coherent orientations  $o_{bm}$  in [BM04] and  $o_{ht}$  in [HT09] below.

**Theorem 2.13** ([BM04]). *There exists a choice of orientation  $o_{bm}$  for all CR tuples such that*

- (1) *the gluing map preserves the orientation with a sign correction:*

$$o_{bm}(\mathcal{T}) \# o_{bm}(\mathcal{T}') = \epsilon_{bm}^\# \cdot o_{bm}(\mathcal{T} \# \mathcal{T}'),$$

where

$$\epsilon_{bm}^\# = (-1)^{\prod_{1 \leq a < b \leq k_-} |S_a^-| \cdot |S_b^-|}.$$

*The sign that comes from reverting the ordering of negative ends of  $\mathcal{T}$ ;*

- (2) *disjoint union map preserves the orientation with a sign correction:*

$$o_{bm}(\mathcal{T}) \prod o_{bm}(\mathcal{T}') = \epsilon_{bm}^{\mathbb{I}} \cdot o_{bm}(\mathcal{T} \prod \mathcal{T}'),$$

where

$$\epsilon_{bm}^{\mathbb{I}} = (-1)^{\text{ind}^- \mathcal{T} \cdot \text{ind}^+ \mathcal{T}'};$$

- (3) *when  $k_- + k_+ = 0$ ,  $o_{bm}(\mathcal{T}) = o_{\text{can}}(\mathcal{T})$ ; and*  
(4) *an isomorphism between CR tuples preserves orientation.*

The gluing convention in this paper is different from that of [BM04]<sup>2</sup>, which is source of the extra sign  $\epsilon_{bm}^\#$ .

**Theorem 2.14** ([HT09]). *There exists a choice of orientation  $o_{ht}$  for all CR tuples such that*

- (1) *the gluing map preserves the orientation, i.e.,*

$$o_{ht}(\mathcal{T}) \# o_{ht}(\mathcal{T}') = o_{ht}(\mathcal{T} \# \mathcal{T}');$$

<sup>2</sup>The matching ordering between the negative punctures of  $\dot{\Sigma}$  and the positive punctures of  $\dot{\Sigma}'$  is reversed in [BM04].

(2) disjoint union map preserves the orientation with a sign correction:

$$o_{ht}(\mathcal{T}) \amalg o_{ht}(\mathcal{T}') = \epsilon_{ht}^{\amalg} \cdot o_{ht}(\mathcal{T} \amalg \mathcal{T}'),$$

where

$$\epsilon_{ht}^{\amalg} = (-1)^{\text{ind}^- \mathcal{T} \cdot \text{ind} \mathcal{T}'};$$

(3) when  $k_- + k_+ = 0$ ,  $o_{ht}(\mathcal{T}) = o_{\text{can}}(\mathcal{T}')$ ; and

(4) an isomorphism between CR tuples preserves orientation.

Let  $\mathcal{T}'$  be the same CR tuple as  $\mathcal{T}$  except that the  $i$ -th and the  $(i+1)$ -th positive (or negative) punctures swapped. Note that  $\det \mathcal{T}$  does not depend on the ordering of the punctures of  $\mathcal{T}$ , so  $\det \mathcal{T} = \det \mathcal{T}'$ . On the other hand, the coherent orientation depends on the ordering of the punctures. Indeed,

**Corollary 2.15.** *Let  $o = o_{bm}$  or  $o_{ht}$ . Then*

$$o(\mathcal{T}) = (-1)^{|S_i^\pm| \cdot |S_{i+1}^\pm|} o(\mathcal{T}').$$

*Proof.* The case  $o = o_{bm}$  is covered in [BM04]. The proof for  $o = o_{ht}$  can be proved similarly. We provide a proof for the case when  $\mathcal{T}$  has two positive punctures and zero negative punctures. Let  $\mathcal{T}_{S_i^+}^+$  for  $i = 1, 2$  be the positive capping CR tuple. Then the two gluings  $(\mathcal{T}_{S_1^+}^+ \amalg \mathcal{T}_{S_2^+}^+) \# \mathcal{T}$  and  $(\mathcal{T}_{S_2^+}^+ \amalg \mathcal{T}_{S_1^+}^+) \# \mathcal{T}'$  give the same CR tuple  $\mathcal{T}''$  with no punctures. Hence, we have

$$\begin{aligned} & \left( (-1)^{|S_1^+| \cdot |S_2^+|} o(\mathcal{T}_{S_1^+}^+) \amalg o(\mathcal{T}_{S_2^+}^+) \right) \# o(\mathcal{T}) \\ &= o(\mathcal{T}'') = \left( (-1)^{|S_1^+| \cdot |S_2^+|} o(\mathcal{T}_{S_2^+}^+) \amalg o(\mathcal{T}_{S_1^+}^+) \right) \# o(\mathcal{T}'). \end{aligned}$$

By Lemma 2.10

$$o(\mathcal{T}_{S_1^+}^+) \amalg o(\mathcal{T}_{S_2^+}^+) = (-1)^{|S_1^+| \cdot |S_2^+|} o(\mathcal{T}_{S_2^+}^+) \amalg o(\mathcal{T}_{S_1^+}^+),$$

which implies the statement.  $\square$

**2.5. Reformulation of  $o_{ht}$ .** We reformulate  $o_{ht}$  so that the calculation becomes simple. For each CR tuple  $\mathcal{T}$ , we pick a nonzero element  $o(\mathcal{T}) \in \det \mathcal{T}$  such that when  $\mathcal{T}$  has no punctures,  $o(\mathcal{T}) = o_{\text{can}}(\mathcal{T})$ . For any  $\mathcal{T}$  with punctures, as before we cap them to reduce it to the no puncture case. In this process, what matters are the asymptotic data at the punctures, the number and the ordering of the punctures. This can be described by the multiplicative group  $\mathcal{O}$  with elements of the form

$$\epsilon o(\mathcal{T}_1) \cdots o(\mathcal{T}_n)$$

for some  $n \geq 0$ , where  $\epsilon \in \{\pm 1\}$ ,  $\mathcal{T}_i$  is a CR tuple for  $i \in \{1, \dots, n\}$ , such that the following conditions are satisfied:

- (1)  $o(\mathcal{T}) = 1$  for any CR tuple  $\mathcal{T}$  without punctures;
- (2)  $o(\mathcal{T})o(\mathcal{T}') = (-1)^{\text{ind} \mathcal{T} \cdot \text{ind} \mathcal{T}'} o(\mathcal{T}')o(\mathcal{T})$ ;
- (3)  $o(\mathcal{T})o(\mathcal{T}') = \epsilon_{ht}^{\amalg} o(\mathcal{T} \amalg \mathcal{T}')$ , for some  $\epsilon_{ht}^{\amalg} \in \{\pm 1\}$  depending on  $\mathcal{T}$  and  $\mathcal{T}'$ ;



- (4) when all the negative punctures of  $\mathcal{T}$  match all the positive punctures of  $\mathcal{T}'$ ,

$$o(\mathcal{T})o(\mathcal{T}') = \epsilon^\sharp o(\mathcal{T} \sharp \mathcal{T}'),$$

for some  $\epsilon^\sharp \in \{\pm 1\}$  depending on  $\mathcal{T}$  and  $\mathcal{T}'$ , and in particular  $\epsilon^\sharp = 1$  when  $\mathcal{T}$  is a positive capping CR tuple and  $\mathcal{T}'$  is the corresponding negative capping CR tuple; and

- (5)  $o(\mathcal{T})$  satisfies

$$(2.5.1) \quad o(\mathcal{T})o_1^- \cdots o_{k_-}^- o_{k_+}^+ \cdots o_1^+ = 1,$$

where  $o_i^\pm$  is the positive (negative) capping orientation of the  $i$ th positive (negative) puncture of  $\mathcal{T}$ .

It is easy to see that the group  $\mathcal{O}$  is generated by  $o(\mathcal{T}_S^+)$  for each admissible  $S$  subject to Condition (2), and

$$o(\mathcal{T}) = o(\mathcal{T}_{S_1^+}^+) \cdots o(\mathcal{T}_{S_{k_+}^+}^+) o(\mathcal{T}_{S_{k_-}^-}^-)^{-1} \cdots o(\mathcal{T}_{S_1^-}^-)^{-1},$$

for any CR tuple  $\mathcal{T}$ . The coherent orientation  $o(\mathcal{T})$  is determined by Equation (2.4.1) and (2.4.2), and the constants  $\epsilon^{\amalg}$  and  $\epsilon^\sharp$  can be determined by Equation (2.5.3) in the group  $\mathcal{O}$ .

*We do not distinguish notation of  $o(\mathcal{T})$  from the notation of equivalence class of  $o(\mathcal{T})$  in  $\mathcal{O}$ , when there is no confusion.*

**Lemma 2.16.** *The orientation  $o$  defined above is  $o_{ht}$ , and in particular,  $\epsilon^{\amalg} = \epsilon_{ht}^{\amalg}$  and  $\epsilon^\sharp = \epsilon_{ht}^\sharp$ .*

*Proof.* It is easy to check that  $o = o_{ht}$  satisfies Equation (2.5.1).  $\square$

Although logically unnecessary, we provide a proof of property (1) and (2) in Theorem 2.14 using our reformulation. It explains how the constants  $\epsilon_{ht}^{\amalg}$  and  $\epsilon_{ht}^\sharp$  appear.

*Proof of (2) in Theorem 2.14.* By definition, we have

$$(2.5.2) \quad o(\mathcal{T})o^- o^+ = 1,$$

$$(2.5.3) \quad o(\mathcal{T}')o'^- o'^+ = 1,$$

and

$$(2.5.4) \quad o(\mathcal{T} \amalg \mathcal{T}')o^- o'^- o'^+ o^+ = 1,$$

where

- $o^- = o_1^- \cdots o_{k_-}^-$ ,
- $o^+ = o_{k_+}^+ \cdots o_1^+$ ,
- $o'^- = o'_1^- \cdots o'_{k_-}^-$ , and
- $o'^+ = o'_{k_+}^+ \cdots o'_1^+$ .

From Equation (2.5.2) and (2.5.3) we get

$$\begin{aligned} o(\mathcal{T})o^-(o(\mathcal{T}')o'^-o'^+)o^+ &= 1 \\ (-1)^{\text{ind } o^- \text{ ind } \mathcal{T}'} o(\mathcal{T})o(\mathcal{T}')o^-o'^-o'^+o^+ &= 1. \end{aligned}$$

Compare it with Equation (2.5.4) one gets

$$\epsilon \amalg = (-1)^{\text{ind } o^- \text{ ind } \mathcal{T}'}.$$

□

*Proof of (3) in Theorem 2.14.* Suppose all the negative ends of  $\mathcal{T}$  match all the positive end of  $\mathcal{T}'$ . By definition, one has

$$(2.5.5) \quad o(\mathcal{T} \# \mathcal{T}')o'^-o^+ = 1.$$

From Equation (2.5.2) and (2.5.3) we get

$$\begin{aligned} o(\mathcal{T})(o(\mathcal{T}')o'^-o'^+)o^-o^+ &= 1 \\ o(\mathcal{T})o(\mathcal{T}')o'^-(o'^+o^-)o^+ &= 1, \end{aligned}$$

but

$$o'^+o^- = o'_{k_+}^+ \cdots o'_1^+ o_1^- \cdots o_{k_-}^- = 1,$$

so

$$\epsilon^\# = 1.$$

□

### 3. COHERENT ORIENTATION OF THE MODULI SPACES

Let  $(M, \xi)$  be a contact manifold of dimension  $2n - 1$ , and  $\alpha$  is a contact 1-form such that  $\xi = \ker \alpha$ . Denote by  $R_\alpha$  the Reeb vector field of  $\alpha$ . We say a Reeb orbit is *non-degenerate* if the Poincaré return map restricted to  $\xi$  along the Reeb orbit does not have 1 as an eigenvalue. We assume that all Reeb orbits are non-degenerate.

For each simple (i.e., not multiply covered) Reeb orbit  $\gamma$ , we choose a trivialization  $\tau$  of the symplectic vector bundle  $(\xi, d\alpha|_\xi)$  restricted to  $\gamma$ . Then the linearized flow of  $R_\alpha$  along  $\gamma$  gives a path of symplectic matrices, whose Maslov index is called the *Conley-Zehnder index* of  $\gamma$  and is denoted by  $\mu_{\text{CZ}, \tau}(\gamma)$ . We assign  $\gamma$  the  $\mathbb{Z}_2$ -grading  $|\gamma| = \mu_{\text{CZ}, \tau}(\gamma) + n - 1 \pmod{2}$ , which is independent of the choice of  $\tau$ . On each simple Reeb orbit  $\gamma$  we choose a fixed point  $x_\gamma$ , which we call an asymptotic marker.

Let  $J$  be an ( $\mathbb{R}$ -invariant)  $\alpha$ -tame (Definition 3.1.1 in [BH18]) almost complex structure on  $W := \mathbb{R} \times M$ . We recall the definition of moduli spaces of  $J$ -holomorphic curves.

**Definition 3.1** (moduli space of  $J$ -holomorphic curves). For integers  $k_+ \geq 1$  and  $k_- \geq 0$ , let  $\gamma_\pm = (\gamma_1^\pm, \dots, \gamma_{k_\pm}^\pm)$  be a tuple of Reeb orbits. For any  $g \in \mathbb{Z}^{\geq 0}$ ,  $A \in H_2(M; \mathbb{Z})$ , we define  $\mathcal{M}^g(\gamma_+; \gamma_-; A)$  the moduli space of  $J$ -holomorphic curves to be equivalent classes  $[\Sigma, j, \mathbf{p}, \mathbf{r}, u]$  of tuples such that

- (1)  $(\Sigma, j, \mathbf{p}, \mathbf{r})$  is a *connected*, decorated Riemann surface of genus  $g$  with  $(k_+, k_-)$  marked points.

Let  $\phi_i^\pm : D \subset \mathbb{C} \rightarrow \mathcal{U}_{p_i^\pm} \subset \Sigma$  be a biholomorphic map from the unit disc  $D$  to a neighborhood  $\mathcal{U}_{p_i^\pm}$  of  $p_i^\pm$  such that  $\phi_i^\pm(o) = p_i^\pm$  and  $d\phi_i^\pm(\frac{\partial}{\partial x}) = r_i^\pm$ , where  $\frac{\partial}{\partial x} \in T_o D$ . Let  $h_i^\pm : \mathbb{R}^{\geq 0(\leq 0)} \times S^1 \rightarrow \dot{\mathcal{U}}_{p_i^\pm}$  be the biholomorphic map defined by  $h_i^\pm(s, t) = (\phi_i^\pm)^{-1}(e^{\mp s \mp \sqrt{-1}t})$ .

- (2)  $u : \Sigma - \mathbf{p} \rightarrow W$  is a proper map satisfying
- (a)  $\bar{\partial}_J u = \frac{1}{2}(du + J \circ du \circ j) = 0$ ,
  - (b)  $\lim_{s \rightarrow \pm\infty} u \circ (\phi_i^\pm)^{-1}(e^{\mp s \mp \sqrt{-1}t}) = \gamma_i^\pm(Tt)$ , where  $T > 0$  is the period of  $\gamma_i^\pm$ , and  $\gamma_i^\pm$  is parametrized so that  $\gamma_i^\pm(0) = r_i^\pm$
- (3) the homology class obtained by ‘‘capping off’’ the punctures of  $u$  is  $A$ . See Section 9.1.2 in [BH15] or [BM04] for the ‘‘capping off’’ construction.
- (4)  $(\Sigma, j, \mathbf{p}, \mathbf{r}, u)$  is equivalent to  $(\Sigma', j', \mathbf{p}', \mathbf{r}', u')$  if there exist a biholomorphic map  $f : (\Sigma, j) \rightarrow (\Sigma', j')$  and a constant  $a \in \mathbb{R}$  such that  $f$  also maps  $(\mathbf{p}, \mathbf{r})$  to  $(\mathbf{p}', \mathbf{r}')$  and  $u' = \text{Tr}_a \circ u \circ f$ , where  $\text{Tr}_a : W \rightarrow W$  is the translation by  $a$  in the  $\mathbb{R}$ -direction.

It is convenient to have the following lemma.

**Lemma 3.2.** *Let  $U, W$  be Banach spaces, and  $\phi : U \rightarrow W$  a linear Fredholm operator. Let  $V$  be a finite dimensional vector space, and  $\psi : V \rightarrow W$  be a linear map. Suppose that the map  $\phi \oplus \psi : U \oplus V \rightarrow W$  defined as  $\phi \oplus \psi(u, v) = \phi(u) + \psi(v)$  is surjective. Then there is an isomorphism*

$$\det \phi \simeq \det(\phi \oplus \psi) \otimes \wedge^{\text{top}} V^*,$$

which is natural up to a positive constant.

*Proof.* See Exercise A.23 in [MS12] for the case when  $\psi$  injective. Let  $I = \phi(U) \cap \psi(V) \subset W$ . Suppose  $\phi^{-1}(I) = H \oplus \ker \phi$  and let  $\{u_1, \dots, u_k\}$  be a basis of  $H$ , and  $\{u_{k+1}, \dots, u_{k+\ell}\}$  be a basis of  $\ker \phi$ . Then  $\{\phi(u_1), \dots, \phi(u_k)\}$  forms a basis of  $I$ . Suppose  $V = G \oplus \psi^{-1}(I)$  and  $\psi^{-1}(I) = F \oplus \ker \psi$ . Let

$$\{v_1, \dots, v_m, v_{m+1}, \dots, v_{m+k}, v_{m+k+1}, \dots, v_{m+k+n}\}$$

be a basis of  $V$  such that

- (1)  $\{v_1, \dots, v_m\}$  is a basis of  $G$ ,
- (2)  $\{v_{m+1}, \dots, v_{m+k}\}$  is a basis of  $F$  and  $\psi(v_{m+i}) = \phi(u_i)$  for  $i = 1, \dots, k$ ,  
and
- (3)  $\{v_{k+1}, \dots, v_m\}$  is a basis of  $\ker \psi$ .

The isomorphism is given by

$$\begin{aligned} & u_{k+1} \wedge \dots \wedge u_{k+\ell} \otimes \psi(v_m)^* \wedge \dots \wedge \psi(v_1)^* \\ \mapsto & (u_{k+1}, 0) \wedge \dots \wedge (u_{k+\ell}, 0) \wedge (u_1, -v_{m+1}) \wedge \dots \wedge (u_k, -v_{m+k}) \\ & \wedge (0, v_{m+k+1}) \wedge \dots \wedge (0, v_{m+k+n}) \otimes v_{m+k+n}^* \wedge \dots \wedge v_1^*. \end{aligned}$$

□

For any

$$[\Sigma, j, \mathbf{p}, \mathbf{r}, u] \in \mathcal{M}^g(\gamma_+; \gamma_-; A),$$

consider the complex vector bundle  $E := u^*TW$  over  $\dot{\Sigma}$  where the complex structure is given by  $u^*J$ . Around each Reeb orbit  $\gamma_i^\pm \in \gamma_\pm$ , a trivialization  $\tau$  of  $(\xi, d\alpha|_\xi, J)$  restricted to  $\gamma_i^\pm$  (induced from the trivialization over the underlying simple Reeb orbit) extends to a trivialization of  $(E, J)|_{U_{p_i^\pm}}$  as in Formula (2.1.1).

This gives a CR tuple denoted by  $\mathcal{T}_u$ . A different representative of  $[\Sigma, j, \mathbf{p}, \mathbf{r}, u]$  gives an isomorphic CR tuple. Let

$$D_u : C^\infty(\dot{\Sigma}, E) \rightarrow C^\infty(\dot{\Sigma}, \wedge^{0,1}\dot{\Sigma} \otimes_{\mathbb{C}} E)$$

be the linearized  $\bar{\partial}_J$  operator, it extends to a Fredholm operator as in Formula (2.1.2). Hence, we get a CR operator denoted by  $D_u \in \mathfrak{D}(\mathcal{T}_u)$ .

Fix any  $[u] \in \mathcal{M}^g(\gamma_+; \gamma_-; A)$ . If  $6g - 6 + 2(k_+ + k_-) > 0$ , we have the full linearized  $\bar{\partial}_J$  operator at  $[u]$

$$\mathcal{D} : \mathit{Teich} \oplus C^\infty(\dot{\Sigma}, E) \rightarrow C^\infty(\dot{\Sigma}, \wedge^{0,1}\dot{\Sigma} \otimes_{\mathbb{C}} E),$$

where  $\mathit{Teich}$  is the tangent space of complex structures of  $\dot{\Sigma}$  at  $j$ , which has dimension equals  $6g - 6 + 2(k_+ + k_-)$  and  $\mathcal{D}|_{C^\infty(\dot{\Sigma}, E)} = D_u$ . Suppose that  $[u]$  is transversely cut out, i.e.,  $\text{coker } \mathcal{D} = \{0\}$ . Then

$$\wedge^{\text{top}} (\mathbb{R}\langle \partial_s \rangle \oplus T_{[u]}\mathcal{M}^g(\gamma_+; \gamma_-; A)) \simeq \wedge^{\text{top}} \ker \mathcal{D} \simeq \det D_u \otimes \wedge^{\text{top}} \mathit{Teich}^*,$$

where  $\partial_s$  is induced by the translation  $\text{Tr}_a : W \rightarrow W$  in the  $\mathbb{R}$ -direction, the last isomorphism is given by Lemma 3.2.

If  $6g - 6 + 2(k_+ + k_-) < 0$  and suppose  $[u]$  is transversely cut out, i.e.,  $\text{coker } D_u = \{0\}$ , then

$$\ker D_u = \mathbb{R}\langle \partial_s \rangle \oplus T_{[u]}\mathcal{M}^g(\gamma_+; \gamma_-; A) \oplus \mathit{Aut},$$

where  $\mathit{Aut}$  is the tangent space of the biholomorphism of  $(\dot{\Sigma}, j)$  which has dimension  $-(6g - 6 + 2(k_+ + k_-))$ .

In either case, the virtual dimension of the moduli space is given by

$$\begin{aligned} & \text{virdim } \mathcal{M}^g(\gamma_+; \gamma_-; A) \\ &= \dim_{\mathbb{R}} \ker D_u + 6g - 6 + 2(k_+ + k_-) \\ &= \sum_{i=1}^{k_+} \mu_{\text{CZ}}(S_i^+) - \sum_{i=1}^{k_-} \mu_{\text{CZ}}(S_i^-) + 2c_1(E) + (n-3)(2-2g-k_-+k_+), \end{aligned}$$

and moreover, we have the canonical isomorphism

$$(3.0.1) \quad \det \mathcal{T}_u \simeq \wedge^{\text{top}} (\mathbb{R}\langle \partial_s \rangle \oplus T_{[u]}\mathcal{M}^g(\gamma_+; \gamma_-; A))$$

as the portions  $\mathit{Teich}$  and  $\mathit{Aut}$  are complex and hence canonically oriented. Let  $o = o_{ht}$  or  $o_{bm}$ . We define  $\tilde{o}(u)$  the orientation of  $T_{[u]}\mathcal{M}^g(\gamma_+; \gamma_-; A)$  by the equation

$$(3.0.2) \quad o(\mathcal{T}_u) \simeq \partial_s \wedge \tilde{o}(u),$$

where the isomorphism is given by Formula (3.0.1). In the case when

$$\text{vir} \dim \mathcal{M}^g(\gamma_+; \gamma_-; A) = 1,$$

$\tilde{o}(u) \in \{\pm 1\}$ .

For any  $[u] \in \mathcal{M}_1 = \mathcal{M}^g(\gamma_+; \gamma_-; A)$  and  $[v] \in \mathcal{M}_2 = \mathcal{M}^{g'}(\gamma'_+; \gamma'_-; A')$  that are transversely cut out and the first  $\tau$  Reeb orbits of  $\gamma_-$  match the last  $\tau$  Reeb orbits in  $\gamma'_+$ , we can glue  $u$  and  $v$  with some fixed large gluing parameter along the  $\tau$  punctures, and get  $[w] \in \mathcal{M}_3 = \mathcal{M}^{g+g'}(\gamma_+; \gamma'_-; A + A')$ . The construction of the gluing map on the moduli spaces is standard (see for example Section 10 in [MS12] or Section 6 in [BH15]). The gluing of the CR tuples in Section 2.3 is the linearized version of this, and indeed, we have the commutative diagram

(3.0.3)

$$\begin{array}{ccc} \det \mathcal{T}_u \otimes \det \mathcal{T}_v & \xrightarrow{\#_\tau} & \det \mathcal{T}_w \\ \downarrow \simeq & & \downarrow \simeq \\ \wedge^{\text{top}}(\mathbb{R}\langle \partial_s^1 \rangle \oplus T_{[u]}\mathcal{M}_1) \otimes \wedge^{\text{top}}(\mathbb{R}\langle \partial_s^2 \rangle \oplus T_{[v]}\mathcal{M}_2) & \longrightarrow & \wedge^{\text{top}}(\mathbb{R}\langle \partial_s^3 \rangle \oplus T_{[w]}\mathcal{M}_3), \end{array}$$

where the lower horizontal map is induced by gluing. The curve  $[w]$  is transversely cut out and can be viewed as in a codimension one boundary component of  $\mathcal{M}_3$ , and hence we have the boundary orientation  $\tilde{o}^b(w)$  defined by

$$(3.0.4) \quad \tilde{o}(w) = n \wedge \tilde{o}^b(w),$$

where  $n$  is the outward pointing normal (pointing to the gluing parameter increasing direction). When  $\text{vir} \dim \mathcal{M}_3 = 1$ , we obtain  $\tilde{o}^b(w) \in \{\pm 1\}$ . Denote  $o(\mathcal{T}_u) \simeq \partial_s^u \wedge \tilde{o}(u)$ ,  $o(\mathcal{T}_v) \simeq \partial_s^v \wedge \tilde{o}(v)$ , and  $o(\mathcal{T}_w) \simeq \partial_s^w \wedge \tilde{o}(w)$ . Then we have

$$o(\mathcal{T}_u) \#_\tau o(\mathcal{T}_v) = \epsilon^\# \cdot \epsilon^\Pi \cdot o(\mathcal{T}_u \#_\tau \mathcal{T}_v) = \epsilon^\# \cdot \epsilon^\Pi \cdot o(\mathcal{T}_w),$$

where the formula for partial gluing follows from taking disjoint unions with trivial tuples before gluing, and more precisely

- (1) if  $o = o_{ht}$ ,
  - (a)  $\epsilon^\# = 1$ , and
  - (b)  $\epsilon^\Pi = (-1)^{(\sum_{i=1}^{k_+ - \tau} |S'_{+,i}|) \cdot \text{ind } \mathcal{T}_u}$ ,
- (2) if  $o = o_{bm}$ ,
  - (a)  $\epsilon^\# = \epsilon_{bm}^\#$ , and
  - (b)  $\epsilon^\Pi = (-1)^C$  with

$$\begin{aligned} C = & \sum_{1 \leq i < j \leq \tau} |S'_{+,i}| \cdot |S'_{+,j}| + \left( \sum_{i=1}^{\tau} |S'_{+,i}| \right) \left( \sum_{i=1}^{k_+} |S_{+,i}| \right) \\ & + \sum_{\tau+1 \leq i < j \leq k_-} |S_{-,i}| \cdot |S_{-,j}| + \left( \sum_{i=1}^{k'_-} |S'_{-,i}| \right) \left( \sum_{i=\tau+1}^{k_-} |S_{-,i}| \right). \end{aligned}$$

Tracking the images of  $o(\mathcal{T}_u) \otimes o(\mathcal{T}_v) \in \det \mathcal{T}_u \otimes \det \mathcal{T}_v$  in the commutative diagram (3.0.3) we have

$$\begin{aligned} \epsilon^\sharp \cdot \epsilon^\Pi \cdot \partial_s^3 \wedge n \wedge \tilde{\delta}^b(w) &= \widehat{\partial}_s^1 \wedge \widehat{\tilde{o}}(u) \wedge \widehat{\partial}_s^2 \wedge \widehat{\tilde{o}}(v) \\ &= (-1)^{\text{ind } \mathcal{T}_u - 1} \widehat{\partial}_s^1 \wedge \widehat{\partial}_s^2 \wedge \widehat{\tilde{o}}(u) \wedge \widehat{\tilde{o}}(v) \\ &= (-1)^{\text{ind } \mathcal{T}_u} \partial_s^3 \wedge n \wedge \widehat{\tilde{o}}(u) \wedge \widehat{\tilde{o}}(v), \end{aligned}$$

where the  $\widehat{\eta}$  is defined by multiplying  $\eta$  with a cut off function supported away from the punctures and projecting to the kernel of the linearized  $\bar{\partial}$  operator at  $[w]$  (see for example, Section 9.12 in [BH18] for details in the case when the domains are cylinders.); and the last equality follows from the fact that up to multiplication by a positive number,  $\partial_s^3$  is approximately  $\widehat{\partial}_s^1 + \widehat{\partial}_s^2$  and  $n$  is approximately  $\widehat{\partial}_s^1 - \widehat{\partial}_s^2$ , and hence  $\partial_s^3 \wedge n$  is approximately  $-\widehat{\partial}_s^1 \wedge \widehat{\partial}_s^2$ . In summary, we have the following lemma.

**Lemma 3.3.** *The gluing map  $\mathcal{M}_1 \times \mathcal{M}_2 \rightarrow \partial \mathcal{M}_3$  changes the orientation by the sign  $\epsilon^\sharp \epsilon^\Pi \cdot (-1)^{\text{vir dim } \mathcal{M}_1}$ .*

In particular, if we use the orientation  $o_{ht}$ , in the proof of  $\mathbf{H} \circ \mathbf{H} = 0$  (See Section 4), where  $\text{vir dim } \mathcal{M}_1 = 1$ , we have the gluing map reverses the boundary orientation.

#### 4. SIGNS IN SYMPLECTIC FIELD THEORY

For each Reeb orbit  $\gamma$ , we assign two formal variables  $p_\gamma, q_\gamma$  graded over  $\mathbb{Z}_2$  by  $|\gamma|$ . For any  $A \in H_2(M; \mathbb{Z})$ , we write it multiplicatively as  $e^A$  and grade it by 0. Let  $\hbar$  be a formal variable graded by 0 to keep track of the genus  $g$ . Consider

$$\mathfrak{W} = \mathbb{Q}[\{q_\gamma\}_\gamma, \hbar, \{e^A\}_{A \in H_2(M; \mathbb{Z})}] \llbracket \{p_\gamma\}_\gamma \rrbracket$$

the space of all formal power series in  $\{p_\gamma\}_\gamma$  over the polynomial ring

$$\mathbb{Q}[\{q_\gamma\}_\gamma, \{e^A\}_{A \in H_2(M; \mathbb{Z})}, \hbar].$$

We require that all formal variables are graded commutative except

$$(4.0.1) \quad [p_\gamma, q_\gamma] = p_\gamma q_\gamma - (-1)^{|\gamma|} q_\gamma p_\gamma = \hbar \cdot m(\gamma),$$

where  $m(\gamma)$  is the multiplicity of  $\gamma$  over the underlying simple Reeb orbit. In [EGH00], a potential function  $\mathbf{H}$  is constructed by counting  $J$ -holomorphic curves. We recall and modify the definition as follows:

$$(4.0.2) \quad \mathbf{H} = \sum_{g \geq 0} \sum_{[\gamma_+], [\gamma_-]} \sum_{A \in H_2(M)} |\overline{\mathcal{M}}^g(\gamma_+; \gamma_-; A)| q_{\gamma_-} p_{\gamma_+} e^A \hbar^{g-1},$$

where

- (1) the second summation is over pairs of unordered tuple  $[\gamma_\pm]$  of Reeb orbits,
- (2)  $\gamma_\pm = (\gamma_{\pm, i_1}, \dots, \gamma_{\pm, i_{k_\pm}})$  is an ordered tuple of Reeb orbits that represent the isomorphism class  $[\gamma_\pm]$ ,

- (3)  $\gamma_+^\dagger = (\gamma_{+,i_{k_+}}, \dots, \gamma_{+,i_1})$  is the ordered tuple obtained from  $\gamma_+$  by reversing the ordering,
- (4)  $p_{\gamma_+^\dagger}$  and  $q_{\gamma_-}$  are the monomials of  $p_\gamma, q_\gamma$  that are associated to the ordered tuples  $\gamma_+^\dagger$  and  $\gamma_-$ , respectively, and
- (5)  $|\overline{\mathcal{M}}^g(\gamma_+; \gamma_-; A)|$  is the (virtual) number of elements in the compactified moduli space based on the coherent orientation  $o_{ht}$  and is set to be 0 when the virtual dimension is not 0.

Note that Formula (4.0.2) does not depend on the choice of representatives  $\gamma_\pm$  of  $[\gamma_\pm]$  by Corollary 2.15.

**Remark 4.1.** The sign correction of  $\mathbf{H}$  as mentioned in the abstract is the usage of  $p_{\gamma_+^\dagger}$  over  $p_{\gamma_+}$ .

To make sure that  $\mathbf{H}$  is well-defined, one needs to perturb the moduli space because the multiple covered curves are not transversely cut out in general. There are several versions of perturbation theories that fit or can be generalized to fit this setting. They include but not limited to [Par19; BH15; Ish18; HWZ07]. Transversality is far beyond the scope of this paper.

**Theorem 4.2.** *Suppose that the moduli spaces are transversely cut out after some perturbation. We have*

$$\mathbf{H} \circ \mathbf{H} = 0.$$

One defines the differential  $D : \mathfrak{W} \rightarrow \mathfrak{W}$  by  $Df = [\mathbf{H}, f]$  for all  $f \in \mathfrak{W}$ , and the homology algebra  $H_*(\mathfrak{W}, D)$  to be the homology of  $(\mathfrak{W}, D)$ . Before proving the theorem, we first re-visit the example in Figure 4 of [EGH00] with some modification.

**Example 4.3.** Suppose

$$\mathbf{H} = \mathbf{H}_0 = aq_1q_2p_4\hbar^{-1} + bp_3p_2p_1\hbar^{-1} + \text{other terms},$$

where

- $a = |\mathcal{M}^0(\gamma_4; \gamma_1\gamma_2)|$  with  $|\gamma_4| + |\gamma_1| + |\gamma_2| = 1 \pmod 2$ ,
- $b = |\mathcal{M}^0(\gamma_1\gamma_2\gamma_3; \emptyset)|$  with  $|\gamma_1| + |\gamma_2| + |\gamma_3| = 1 \pmod 2$ ,
- we write  $q_i$  and  $p_i$  for  $q_{\gamma_i}$  and  $p_{\gamma_i}$  respectively,
- we drop the variable  $e^A$  for  $A \in H_2(M; \mathbb{Z})$ , and
- other terms are terms that come from moduli spaces other than  $\mathcal{M}^0(\gamma_4; \gamma_1\gamma_2)$  and  $\mathcal{M}^0(\gamma_1\gamma_2\gamma_3; \emptyset)$ .

An explicit calculation yields

$$\begin{aligned} \mathbf{H} \circ \mathbf{H} &= (-1)^{d_2+d_2d_3+d_2d_4+d_3d_4} abq_2p_4p_3p_2\hbar^{-1} \\ &\quad + (-1)^{d_1+d_1d_2+d_1d_4+d_3d_4} abq_1p_4p_3p_1\hbar^{-1} \\ &\quad + (-1)^{d_3d_4} abp_4p_3 + \text{other terms}, \end{aligned}$$

where  $d_i = |\gamma_i|$ , and two monomials that are multiples of  $q_1q_2p_4p_3p_2p_1$  cancel out. The three terms that appear in  $\mathbf{H} \circ \mathbf{H}$  matches the three gluing of the moduli spaces




FIGURE 2. There are three gluings that correspond to three terms in  $\mathbf{H} \circ \mathbf{H}$ : gluing along  $\gamma_1$ , gluing along  $\gamma_2$ , and simultaneously gluing along  $\gamma_1$  and  $\gamma_2$ .

$\mathcal{M}_I = \mathcal{M}^0(\gamma_4; \gamma_1\gamma_2)$  and  $\mathcal{M}_{II} = \mathcal{M}^0(\gamma_1\gamma_2\gamma_3; \emptyset)$  with signs (See Figure (2)). Consider the moduli space  $\mathcal{M}_{III} = \mathcal{M}^0(\gamma_2\gamma_3\gamma_4; \gamma_2)$ . We check that the number of elements in  $\partial_{I,II}\mathcal{M}_{III}$ , the boundary components of  $\mathcal{M}_{III}$  that correspond to the gluing of the moduli spaces  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$  along  $\gamma_1$ , equals  $(-1)^{d_2+d_2d_3+d_2d_4+d_3d_4+1}ab$ , which is  $(-1)$  times the first term in the coefficient of  $q_2p_4p_3p_2\hbar^{-1}$ .

$$\begin{aligned}
|\partial_{I,II}\mathcal{M}_{III}| &= \sum_{[w] \in \partial_{I,II}\mathcal{M}_{III}} \tilde{\delta}^b(w) \\
&= \sum_{[w] \in \partial_{I,II}\mathcal{M}(423;2)} (-1)^{d_4(d_2+d_3)} \tilde{\delta}^b(w) \\
&= \sum_{[u] \in \mathcal{M}(4;21)} \sum_{[v] \in \mathcal{M}(123;\emptyset)} (-1)^{d_4(d_2+d_3)+d_2(d_1+d_2+d_3)} \tilde{\delta}(u)\tilde{\delta}(v) \\
&= \sum_{[u] \in \mathcal{M}(4;12)} \sum_{[v] \in \mathcal{M}(123;\emptyset)} (-1)^{d_4(d_2+d_3)+d_2(d_1+d_2+d_3)+d_1d_2} \tilde{\delta}(u)\tilde{\delta}(v) \\
&= (-1)^{d_2+d_2d_3+d_2d_4+d_3d_4+1} \left( \sum_{[u] \in \mathcal{M}(4;12)} \tilde{\delta}(u) \right) \left( \sum_{[v] \in \mathcal{M}(123;\emptyset)} \tilde{\delta}(v) \right) \\
&= (-1)^{d_2+d_2d_3+d_2d_4+d_3d_4+1} ab,
\end{aligned}$$

where in the notation of the moduli spaces, we drop  $\gamma$  to make it clearer. We leave the computation of the other two terms for readers.

*Proof of Theorem 4.2.* Note that

$$\mathbf{H} \circ \mathbf{H} = \sum \tilde{\delta}(u)\tilde{\delta}(u')q_{\gamma_-}p_{\gamma_+^\dagger}q_{\gamma'_-}p_{\gamma'_+^\dagger}e^{A+A'}\hbar^{g+g'-2},$$

where  $\tilde{\delta}(u), \tilde{\delta}(u') \in \{\pm 1\}$ , and the summation is over  $g, g' \geq 0$ , unordered tuples of good Reeb orbits  $[\gamma_+], [\gamma_-], [\gamma'_+], [\gamma'_-]$ , homology classes  $A, A' \in H_2(M)$ , and  $[u] \in \mathcal{M}^g(\gamma_+, \gamma_-; A)$  and  $[u'] \in \mathcal{M}^{g'}(\gamma'_+, \gamma'_-; A')$ . We move the variables



in  $q_{\gamma'_-}$  across  $p_{\gamma'_+}$  to the left one by one according to the graded commutativity and Formula (4.0.1). To better keep track of the total sign introduced of the above moves, we include *totally ordered* CR tuples, which are CR tuples with the extra data: an ordering of all marked points that is consistent with the ordering of negative marked points and the reversed ordering of positive marked points. The reformulation in Section 2.5 simply generalizes to totally ordered CR tuples with Formula (2.5.2) replaced by

$$(4.0.3) \quad o(\Gamma)o_1 \cdots o_n = 1,$$

where

- $n$  is the total number of marked points; and
- if the  $i$ -th marked point is positive (negative) and is associated with the loop of symmetric matrices  $S_i$ ,  $o_i$  is the positive (negative) capping orientation of  $S_i$ .

Given a CR tuple  $\mathcal{T}$ , we can associate a compatible totally ordered CR tuple  $\Gamma(\mathcal{T})$  by ordering the negative marked points before the positive marked points, which matches the left to right ordering of  $q_{\gamma_-}p_{\gamma'_+}$ . In this case,  $o(\Gamma(\mathcal{T})) = o(\mathcal{T})$  by Formula (2.5.1). For each term  $\tilde{o}(u)\tilde{o}(u')q_{\gamma_-}p_{\gamma'_+}q_{\gamma'_-}p_{\gamma_+}$  we associate a totally ordered CR tuple  $\Gamma_{u,u'}$  by taking the disjoint union of  $\mathcal{T}_u$  and  $\mathcal{T}_{u'}$  and the ordering of the marked points is given by the ordering of  $\Gamma(\mathcal{T}_u)$  followed by that of  $\Gamma(\mathcal{T}_{u'})$ , which matches the left to right ordering of  $q_{\gamma_-}p_{\gamma'_+}q_{\gamma'_-}p_{\gamma_+}$ . It is easy to see that

$$o(\Gamma_{u,u'}) = o(\mathcal{T}_u) \wedge o(\mathcal{T}_{u'}).$$

Now it is clear that every step of re-arranging the variables  $q_{\gamma_-}p_{\gamma'_+}q_{\gamma'_-}p_{\gamma_+}$  corresponds to switching the ordering of two adjacent punctures in  $\Gamma_{u,u'}$ , when the two variables do not correspond to the same Reeb orbit. When they correspond to the same Reeb orbit, we get two totally ordered CR tuples: one by switching ordering of punctures, and one by gluing along the two punctures, whose definition is an obvious generalization of the gluing in Section 2.3 (the relative ordering of the remaining punctures stay the same). In particular, let  $\Gamma$  be a totally ordered CR tuple. Suppose that the  $i$ -th puncture is a positive puncture,  $(i+1)$ -th puncture is a negative puncture and they correspond to the same loop of admissible matrices. Let  $\Gamma'$  be the totally ordered CR tuple obtained from gluing the  $i$ -th puncture and  $(i+1)$ -th puncture. Suppose that  $o(\Gamma)$  satisfies Equation (4.0.3). Then  $o(\Gamma')$  satisfies

$$o(\Gamma')o_1 \cdots o_{i-1}o_{i+1} \cdots o_{2n} = 1,$$

as  $o_i o_{i+1} = 1$ . Hence,  $o(\Gamma) = o(\Gamma')$  viewed as elements in the group  $\mathcal{O}$ . In other words, gluing two adjacent puncture preserves orientation. Now it is easy to see that in the end of the switching operation, each algebraic term that we obtained corresponds exactly to one CR tuple. Hence, for each  $[\gamma_+], [\gamma_-], A, g \geq -1$ , the coefficient in front of the monomial  $q_{\gamma_-}p_{\gamma'_+}e^A \hbar^g$  in  $\mathbf{H} \circ \mathbf{H}$  is given by the term

$$\sum \epsilon \tilde{o}(u)\tilde{o}(u'),$$

where the sum is taken over all triples  $([u], [u'], \vartheta)$  of equivalence classes of  $J$ -holomorphic curves  $[u]$  and  $[u']$ , and a matching  $\vartheta$  of certain positive ends of  $u$  with certain negative ends of  $u'$  such that if we glue  $u$  and  $u'$  along  $\vartheta$ , we obtain a curve  $w = u \#_{\vartheta} u'$  which satisfies  $[w] \in \mathcal{M}^g(\gamma_+, \gamma_-)$ , and  $\epsilon \in \{\pm 1\}$  is the sign that comes from the algebraic operation of moving  $q$ 's across  $p$ 's. On the other hand, the sign  $\epsilon$  also appears in

$$o(\mathcal{T}_w) = \epsilon o(\mathcal{T}_{u'}) \#_{\vartheta} o(\mathcal{T}_u).$$

But  $o(\mathcal{T}_w) \simeq \partial_s \wedge n \wedge \tilde{o}^b(w)$ , where  $n$  is the outward pointing normal of the boundary of  $\mathcal{M}^g(\gamma_+; \gamma_-)$ , and  $o(\mathcal{T}_{u'}) \simeq \partial_s^{u'} \wedge \tilde{o}(u')$  and  $o(\mathcal{T}_u) \simeq \partial_s^u \wedge \tilde{o}(u)$ , as in Formula (3.0.2) (3.0.4). A similar calculation as in the proof of Lemma 3.3 with  $\#_{\tau}$  replaced by  $\#_{\vartheta}$  gives  $\tilde{o}^b(w) = -\epsilon \tilde{o}(u') \tilde{o}(u)$ , where we use the fact that  $\text{ind } \mathcal{T}_u = 1 = \text{ind } \mathcal{T}_{u'}$ . Therefore

$$\sum \epsilon \tilde{o}(u) \tilde{o}(u') = - \sum \tilde{o}_w^b = 0.$$

□

**Corollary 4.4** (Contact homology [Par19; BH15; Ish18]). *Let  $\mathfrak{A}$  be the differential graded commutative algebra generated freely by all good Reeb orbits over  $\mathbb{Q}[H_2(M)]$ . Let  $\partial : \mathfrak{A} \rightarrow \mathfrak{A}$  be the differential defined over generators by*

$$(4.0.4) \quad \partial \gamma_+ = \sum_{[\gamma_-]} \sum_{A \in H_2(M; \mathbb{Z})} \frac{1}{m(\gamma_-)} |\mathcal{M}^0(\gamma_+; \gamma_-; A)| \cdot e^A \gamma_{-,1} \dots \gamma_{-,k},$$

where  $\gamma_- = \gamma_{-,1} \dots \gamma_{-,k}$ , and  $m(\gamma_-) = m(\gamma_{-,1}) \dots m(\gamma_{-,k})$ . Then  $\partial^2 = 0$ .

*Proof.* This follows from Theorem 4.2 by restricting to the term  $g = 0$  and linear  $p_{\gamma_+}^{\dagger}$ . □

**Remark 4.5.** If one uses  $o_{bm}$ , then  $\partial$  should be defined as

$$(4.0.5) \quad \partial \gamma_+ = \sum_{[\gamma_-]} \sum_{A \in H_2(M; \mathbb{Z})} \frac{1}{m(\gamma_-)} |\mathcal{M}^0(\gamma_+; \gamma_-; A)| \cdot e^A \gamma_{-,k} \dots \gamma_{-,1}$$

as in [BH15].

Lastly, we mention the follow result:

**Proposition 4.6.** *Different choices of capping CR tuples  $\mathcal{T}_S^{\pm}$  and the capping orientations  $o(\mathcal{T}_S^{\pm})$  produce isomorphic SFT.*

*Proof.* Let  $\mathcal{T}_S^{\pm}$  and  $o'(\mathcal{T}_S^{\pm})$  be a different choices of capping CR tuples and their capping orientation. Let  $\epsilon_S \in \{\pm 1\}$  be defined by  $\epsilon_S o(\mathcal{T}_S^+) \# o'(\mathcal{T}_S^-) = o_{\text{can}}(\mathcal{T}_S^+ \# \mathcal{T}_S'^-)$ . Then the isomorphism from  $(\mathfrak{W}, D) \rightarrow (\mathfrak{W}, D')$  is defined by sending generators  $p_{\gamma}, q_{\gamma} \mapsto \epsilon_S p_{\gamma}, \epsilon_S q_{\gamma}$  for all  $\gamma$ , where  $S$  is the loop of symmetric matrices associated to  $\gamma$ . □

#### ACKNOWLEDGEMENT

We thank Ko Honda, Russell Avdek, and Fan Zheng for helpful discussions.

## REFERENCES

- [BH15] E. Bao and K. Honda. Semi-global Kuranishi charts and the definition of contact homology. *ArXiv preprint arXiv:1512.00580*, 2015.
- [BH18] E. Bao and K. Honda. Definition of cylindrical contact homology in dimension three. *Journal of Topology* 11.4, 2018, pp. 1002–1053.
- [Bou02] F. Bourgeois. A Morse-Bott approach to contact homology. PhD thesis. stanford university, 2002.
- [BM04] F. Bourgeois and K. Mohnke. Coherent orientations in symplectic field theory. *Mathematische Zeitschrift* 248.1, 2004, pp. 123–146.
- [EGH00] Y. Eliashberg, A. Givental, and H. Hofer. Introduction to symplectic field theory. *Geom. Funct. Anal.* Special Volume, Part II, 2000. GAFA 2000 (Tel Aviv, 1999), pp. 560–673.
- [FOOO09] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono. *Lagrangian Intersection Floer Theory: Anomaly and Obstruction. Parts I and II*. Vol. 46.1 and 46.2. AMS/IP Studies in Advanced Mathematics. American Mathematical Society, 2009.
- [HWZ07] H. W. Hofer, K. Wysocki, and E. Zehnder. A general Fredholm theory I: A splicing-based differential geometry. *Journal of the European Mathematical Society* 9.4, 2007, pp. 841–876.
- [HT09] M. Hutchings and C. H. Taubes. Gluing pseudoholomorphic curves along branched covered cylinders II. *Journal of Symplectic Geometry* 7.1, 2009, pp. 29–133.
- [Ish18] S. Ishikawa. Construction of general symplectic field theory. *ArXiv preprint arXiv:1807.09455*, 2018.
- [MS12] D. McDuff and D. Salamon. *J-holomorphic curves and symplectic topology*. Vol. 52. American Mathematical Soc., 2012.
- [Par19] J. Pardon. Contact homology and virtual fundamental cycles. *Journal of the American Mathematical Society* 32.3, 2019, pp. 825–919.
- [Sch95] M. Schwarz. Cohomology operations from  $S^1$ -cobordisms in Floer homology. PhD thesis. ETH Zurich, 1995.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455

*Email address:* bao@umn.edu

*URL:* <https://erkaobao.github.io/math/>