

The Wavefront Sets of Unipotent Supercuspidal Representations

Dan Ciubotaru, Lucas Mason-Brown, and Emile Okada

July 11, 2022

Abstract

Let $\mathbf{G}(\mathbf{k})$ be a semisimple p -adic group, inner to split. In this article, we compute the algebraic and canonical unramified wavefront sets of the irreducible supercuspidal representations of $\mathbf{G}(\mathbf{k})$ in Lusztig's category of unipotent representations.

Contents

1	Introduction	1
2	Preliminaries	2
2.1	The Bruhat-Tits Building	3
2.2	Nilpotent orbits	4
2.3	Wavefront sets	7
2.4	Isogenies	8
2.5	Depth-0 representations	8
2.6	Unipotent supercuspidal representations	10
2.7	Langlands classification of unipotent supercuspidal representations	10
3	Main result	11
4	Unipotent cuspidal representations of finite reductive groups	11
4.1	Classical groups	12
4.2	Exceptional groups	13
5	Langlands parameters for unipotent supercuspidal representations	14
5.1	Classical groups	14
5.2	Exceptional groups	16
6	Proof of Proposition 3.0.1	16
6.1	Classical groups	16
6.2	Exceptional groups	18

1 Introduction

Let \mathbf{G} be a connected semisimple algebraic group defined over a p -adic field \mathbf{k} with residue field \mathbb{F}_q and let $\mathbf{G}(\mathbf{k})$ be the group of \mathbf{k} -rational points. Let X be an irreducible admissible representation of $\mathbf{G}^\omega(\mathbf{k})$ with distribution character Θ_X . For each nilpotent orbit \mathcal{O} in the Lie algebra $\mathfrak{g}(\mathbf{k})$ of $\mathbf{G}(\mathbf{k})$, let $\hat{\mu}_{\mathcal{O}}$ denote the Fourier transform of the associated orbit integral. In [Har99], Harish-Chandra proved that there are complex numbers $c_{\mathcal{O}}(X) \in \mathbb{C}$ such that

$$\Theta_X(\exp(\xi)) = \sum_{\mathcal{O}} c_{\mathcal{O}}(X) \hat{\mu}_{\mathcal{O}}(\xi) \quad (1.0.1)$$

for $\xi \in \mathfrak{g}(\mathbf{k})$ a regular element in a small neighborhood of 0. The formula (2.3.1) is called the *local character expansion* of X .

One of the most fundamental invariants which can be extracted from the local character expansion is the so-called *wavefront set*. This is the set of nilpotent orbits

$$\mathrm{WF}(X) = \max\{\mathbb{O} \subset \mathfrak{g}(\mathfrak{k}) \mid c_{\mathbb{O}}(X) \neq 0\}$$

It is common in the literature to consider a slightly coarser invariant called the *geometric wavefront set*. This is the set of nilpotent orbits over an *algebraic closure* $\bar{\mathfrak{k}}$ of \mathfrak{k} which meet some orbit in $\mathrm{WF}(X)$. This set is denoted by ${}^{\bar{\mathfrak{k}}}\mathrm{WF}(X)$. A longstanding conjecture of Mœglin and Waldspurger [MW87, Page 429] asserts that ${}^{\bar{\mathfrak{k}}}\mathrm{WF}(X)$ is a singleton, for all X . Another closely related invariant is the *canonical unramified wavefront set*, which was recently introduced by the third-named author in [Oka21, Section 2.2.3]. This final invariant is a refinement of ${}^{\bar{\mathfrak{k}}}\mathrm{WF}(X)$.

Lusztig in [Lus95, Section 0.3] defined the notion of a *unipotent representation* of $\mathbf{G}(\mathfrak{k})$ (see Definition 2.6.1 below) and completed a Langlands classification for this class of representations, under the assumption that \mathbf{G} is simple and adjoint. These restrictions on \mathbf{G} have since been removed as a result of various works, culminating with [Sol18].

Let G^\vee denote the complex Langlands dual group. The (enhanced) Langlands parameter of an irreducible unipotent representation X is the G^\vee -orbit of a triple (τ, n, ρ) , where τ is a semisimple element in G^\vee , n is a nilpotent element in \mathfrak{g}^\vee (the Lie algebra of G^\vee) such that $\mathrm{Ad}(\tau)n = \rho n$, and ρ is an irreducible representation of a certain finite group A_φ^1 , see section 2.7. When \mathbf{G} is adjoint, A_φ^1 is the component group of the centralizer of τ and n in G^\vee . Let \mathbb{O}_X^\vee denote the nilpotent G^\vee -orbit of n .

It is natural to ask if and how the local character expansion is related to the Langlands parameter (τ, n, ρ) of X . At one extreme, we have the coefficient $c_{\mathbb{O}}(X)$. It has long been known that when X is tempered, $c_{\mathbb{O}}(X) \neq 0$ if and only if X is square integrable, and in this case, $c_{\mathbb{O}}(X)$ equals, up to a sign, the ratio between the formal degrees of X and the Steinberg representation. An interpretation of the formal degree in terms of the Langlands parameters was conjectured first by Reeder [Ree00]. This interpretation was verified in the case of split exceptional groups by Reeder ([Ree00]) and in the remaining cases by Opdam [Opd16].

At the other extreme, we have the wavefront set of X . In this article, we compute the geometric and canonical unramified wavefront sets of all *supercuspidal* unipotent representations of $\mathbf{G}(\mathfrak{k})$ when \mathbf{G} is an inner twist of a split group. In all such cases, we find that ${}^{\bar{\mathfrak{k}}}\mathrm{WF}(X)$ is a singleton, and is uniquely determined by the nilpotent part n of the Langlands parameter. More precisely, we prove in Theorem 3.0.2

$${}^{\bar{\mathfrak{k}}}\mathrm{WF}(X) = d(\mathbb{O}_X^\vee), \quad {}^K\mathrm{WF}(X) = d_A(\mathbb{O}_X^\vee, 1). \quad (1.0.2)$$

Here d and d_A are the duality maps defined by Spaltenstein, also Lusztig and Barbasch-Vogan, and Achar, respectively, see section 2.2. In particular, the geometric wavefront set ${}^{\bar{\mathfrak{k}}}\mathrm{WF}(X)$ determines the nilpotent orbit \mathbb{O}_X^\vee via the duality map d . We emphasize that the simplicity of the formulas (1.0.2) is due to the fact that X is supercuspidal and therefore equal to its Aubert-Zelevinsky dual ([Aub95]). In general, one expects that the wavefront set of X is obtained by duality from the nilpotent parameter associated to the *AZ dual* of X , see [CMO21]. This expression for the wavefront set is closely related to Lusztig's formula for the Kawanaka wavefront set of an irreducible unipotent representation of a finite reductive group [Lus92, Theorem 11.2]. In fact, the finite reductive group results from *loc.cit.* play an important role in the construction and analysis of test functions in the local character expansion [BM97; Oka21].

We remark that the methods in this paper show that the canonical unramified wavefront set (and so the geometric wavefront set as well) is in fact a singleton for all depth-zero supercuspidal representations, see Proposition 2.5.4. The irreducibility of the geometric wavefront set for depth-zero supercuspidal representations was established independently in [AGS22], but also as a consequence of the results in [BM97; Oka21].

2 Preliminaries

Let \mathfrak{k} be a nonarchimedean local field of characteristic 0 with residue field \mathbb{F}_q of sufficiently large characteristic, ring of integers $\mathfrak{o} \subset \mathfrak{k}$, and valuation $\mathrm{val}_{\mathfrak{k}}$. Fix an algebraic closure $\bar{\mathfrak{k}}$ of \mathfrak{k} with Galois group Γ , and let $K \subset \bar{\mathfrak{k}}$ be the maximal unramified extension of \mathfrak{k} in $\bar{\mathfrak{k}}$. Let \mathfrak{D} be the ring of integers of K . Let Frob be the geometric Frobenius element of $\mathrm{Gal}(K/\mathfrak{k})$, the topological generator which induces the inverse of the automorphism $x \rightarrow x^q$ of \mathbb{F}_q .

Let \mathbf{G} be a connected semisimple algebraic group defined over \mathbb{Z} , and let $\mathbf{T} \subset \mathbf{G}$ be a maximal torus. For any field F , we write $\mathbf{G}(F)$, $\mathbf{T}(F)$, etc. for the groups of F -rational points. The \mathbb{C} -points are denoted by G , T , etc. Let $\mathbf{G}_{\text{ad}} = \mathbf{G}/Z(\mathbf{G})$ denote the adjoint group of \mathbf{G} .

Write $X^*(\mathbf{T}, \bar{k})$ (resp. $X_*(\mathbf{T}, \bar{k})$) for the lattice of algebraic characters (resp. co-characters) of $\mathbf{T}(\bar{k})$, and write $\Phi(\mathbf{T}, \bar{k})$ (resp. $\Phi^\vee(\mathbf{T}, \bar{k})$) for the set of roots (resp. co-roots). Let

$$\mathcal{R} = (X^*(\mathbf{T}, \bar{k}), \Phi(\mathbf{T}, \bar{k}), X_*(\mathbf{T}, \bar{k}), \Phi^\vee(\mathbf{T}, \bar{k}), \langle, \rangle)$$

be the root datum corresponding to \mathbf{G} , and let W the associated (finite) Weyl group. Let \mathbf{G}^\vee be the Langlands dual group of \mathbf{G} , i.e. the connected reductive algebraic group corresponding to the dual root datum

$$\mathcal{R}^\vee = (X_*(\mathbf{T}, \bar{k}), \Phi^\vee(\mathbf{T}, \bar{k}), X^*(\mathbf{T}, \bar{k}), \Phi(\mathbf{T}, \bar{k}), \langle, \rangle).$$

Set $\Omega = X_*(\mathbf{T}, \bar{k})/\mathbb{Z}\Phi^\vee(\mathbf{T}, \bar{k})$. The center $Z(\mathbf{G}^\vee)$ can be naturally identified with the irreducible characters $\text{Irr}\Omega$, and dually, $\Omega \cong X^*(Z(\mathbf{G}^\vee))$. For $\omega \in \Omega$, let ζ_ω denote the corresponding irreducible character of $Z(\mathbf{G}^\vee)$.

For details regarding the parametrization of inner twists of a group $\mathbf{G}(k)$, see [Vog93, §2], [Kot84], [Kal16, §2], or [ABPS17, §1.3] and [FOS19, §1]. We only record here that the set of equivalence classes of inner twists of the split form of \mathbf{G} are parametrized by the Galois cohomology group

$$H^1(\Gamma, \mathbf{G}_{\text{ad}}) \cong H^1(F, \mathbf{G}_{\text{ad}}(K)) \cong \Omega_{\text{ad}} \cong \text{Irr} Z(\mathbf{G}_{\text{sc}}^\vee),$$

where $\mathbf{G}_{\text{sc}}^\vee$ is the Langlands dual group of \mathbf{G}_{ad} , i.e., the simply connected cover of \mathbf{G}^\vee , and F denotes the action of Frobenius on $\mathbf{G}(K)$. We identify Ω_{ad} with the fundamental group of \mathbf{G}_{ad} . The isomorphism above is determined as follows: for a cohomology class h in $H^1(F, \mathbf{G}_{\text{ad}}(K))$, let z be a representative cocycle. Let $u \in \mathbf{G}_{\text{ad}}(K)$ be the image of F under z , and let ω denote the image of u in Ω_{ad} . Set $F_\omega = \text{Ad}(u) \circ F$. The corresponding rational structure of \mathbf{G} is given by F_ω . Let \mathbf{G}^ω be the connected semisimple group defined over k such that $\mathbf{G}(K)^{F_\omega} = \mathbf{G}^\omega(k)$. Note that $\mathbf{G}^1 = \mathbf{G}$ (where we view \mathbf{G} as an algebraic group over k for this equality).

If H is a complex reductive group and x is an element of H or \mathfrak{h} , we write $H(x)$ for the centralizer of x in H , and $A_H(x)$ for the group of connected components of $H(x)$. If S is a subset of H or \mathfrak{h} (or indeed, of $H \cup \mathfrak{h}$), we can similarly define $H(S)$ and $A_H(S)$. We will sometimes write $A(x)$, $A(S)$ when the group H is implicit. The subgroups of H of the form $H(x)$ where x is a semisimple element of H are called *pseudo-Levi* subgroups of H .

Let $\mathcal{C}(\mathbf{G}(k))$ be the category of smooth complex $\mathbf{G}(k)$ -representations and let $\Pi(\mathbf{G}(k)) \subset \mathcal{C}(\mathbf{G}(k))$ be the set of irreducible objects. Let $R(\mathbf{G}(k))$ denote the Grothendieck group of $\mathcal{C}(\mathbf{G}(k))$.

2.1 The Bruhat-Tits Building

In this section we will recall some standard facts about the Bruhat-Tits building (all of which can be found in [Tit79]).

Fix a $\omega \in \Omega$ and let \mathbf{G}^ω be the inner twist of \mathbf{G} corresponding ω as defined in the previous section. Let $\mathcal{B}(\mathbf{G}^\omega, k)$ denote the (enlarged) Bruhat-Tits building for $\mathbf{G}^\omega(k)$. Let $\mathcal{B}(\mathbf{G}, K)$ denote the (enlarged) Bruhat-Tits for $\mathbf{G}(K)$. For an apartment \mathcal{A} of $\mathcal{B}(\mathbf{G}, K)$ and $\Omega \subseteq \mathcal{A}$ we write $\mathcal{A}(\Omega, \mathcal{A})$ for the smallest affine subspace of \mathcal{A} containing Ω . The inner twist \mathbf{G}^ω of \mathbf{G} gives rise to an action of the Galois group $\text{Gal}(K/k)$ on $\mathcal{B}(\mathbf{G}, K)$ and we can (and will) identify $\mathcal{B}(\mathbf{G}^\omega, k)$ with the fixed points of this action. We use the notation $c \subseteq \mathcal{B}(\mathbf{G}^\omega, k)$ to indicate that c is a face of $\mathcal{B}(\mathbf{G}^\omega, k)$. Given a maximal k -split torus \mathbf{T} of \mathbf{G}^ω , write $\mathcal{A}(\mathbf{T}, k)$ for the corresponding apartment in $\mathcal{B}(\mathbf{G}^\omega, k)$. Write $\Phi(\mathbf{T}, k)$ (resp. $\Psi(\mathbf{T}, k)$) for the set of roots of $\mathbf{G}(k)$ (resp. affine roots) of $\mathbf{T}(k)$ on $\mathbf{G}^\omega(k)$. For $\psi \in \Psi(\mathbf{T}, k)$ write $\psi \in \Phi(\mathbf{T}, k)$ for the gradient of ψ , and $W = W(\mathbf{T}, k)$ for the Weyl group of $\mathbf{G}(k)$ with respect to $\mathbf{T}(k)$. For a face $c \subseteq \mathcal{B}(\mathbf{G}^\omega, k)$ there is a group \mathbf{P}_c^\dagger defined over \mathfrak{o} such that $\mathbf{P}_c^\dagger(\mathfrak{o})$ identifies with the stabiliser of c in $\mathbf{G}(k)$. There is an exact sequence

$$1 \rightarrow \mathbf{U}_c(\mathfrak{o}) \rightarrow \mathbf{P}_c^\dagger(\mathfrak{o}) \rightarrow \mathbf{L}_c^\dagger(\mathbb{F}_q) \rightarrow 1, \quad (2.1.1)$$

where $\mathbf{U}_c(\mathfrak{o})$ is the pro-unipotent radical of $\mathbf{P}_c^\dagger(\mathfrak{o})$ and \mathbf{L}_c^\dagger is the reductive quotient of the special fibre of \mathbf{P}_c^\dagger . Let \mathbf{L}_c denote the identity component of \mathbf{L}_c^\dagger , and let \mathbf{P}_c be the subgroup of \mathbf{P}_c^\dagger defined over \mathfrak{o} such that $\mathbf{P}_c(\mathfrak{o})$ is the inverse image of $\mathbf{L}_c(\overline{\mathbb{F}}_q)$ in $\mathbf{P}_c^\dagger(\mathfrak{o})$. We also write \mathbf{T} for the well defined split torus scheme over \mathfrak{o} with generic fibre \mathbf{T} . This scheme \mathbf{T} defined over \mathfrak{o} is a subgroup of \mathbf{P}_c and the special fibre of \mathbf{T} , denoted $\overline{\mathbf{T}}$, is a maximal torus of \mathbf{L}_c . For c viewed as a face of $\mathcal{B}(\mathbf{G}, K)$, the stabiliser of c in $\mathbf{G}(K)$ identifies with $\mathbf{P}_c^\dagger(\mathfrak{O})$. It has pro-unipotent radical $\mathbf{U}_c(\mathfrak{O})$ and $\mathbf{P}_c^\dagger(\mathfrak{O})/\mathbf{U}_c(\mathfrak{O}) = \mathbf{L}_c^\dagger(\overline{\mathbb{F}}_q)$. For c a face lying in $\mathcal{B}(\mathbf{G}^\omega, k) \subseteq \mathcal{B}(\mathbf{G}, K)$, F_ω stabilises $\mathbf{P}_c(\mathfrak{O})$ and induces a Frobenius on $\mathbf{L}_c(\overline{\mathbb{F}}_q)$. The group $\mathbf{L}_c(\overline{\mathbb{F}}_q)$ consists of the fixed points of this Frobenius. The groups $\mathbf{P}_c(\mathfrak{o})$ obtained in this manner are called (*k*-)parahoric subgroups of \mathbf{G}^ω . When c is a chamber, then we call $\mathbf{P}_c(\mathfrak{o})$ an *Iwahori subgroup* of \mathbf{G} .

For this paper it will be convenient to fix a maximal *k*-split torus \mathbf{T} of \mathbf{G}^ω and a maximal *K*-split torus \mathbf{T}_1 of \mathbf{G}^ω containing \mathbf{T} and defined over *k*. We have that $\mathcal{A}(\mathbf{T}, k) = \mathcal{A}(\mathbf{T}_1, K)^{\text{Gal}(K/k)}$. We will also fix a $\text{Gal}(K/k)$ -stable chamber c_0 of $\mathcal{A}(\mathbf{T}_1, K)$ and a special point $x_0 \in c_0$. Let $\widetilde{W} = W \rtimes X_*(\mathbf{T}_1, K)$ be the (extended) affine Weyl group. The choice of special point x_0 of $\mathcal{B}(\mathbf{G}, K)$ fixes an inclusion $\Phi(\mathbf{T}_1, K) \rightarrow \Psi(\mathbf{T}_1, K)$ and an isomorphism between \widetilde{W} and $N_{\mathbf{G}(K)}(\mathbf{T}_1(K))/\mathbf{T}_1(\mathfrak{O}^\times)$. Write

$$\widetilde{W} \rightarrow W, \quad w \mapsto \dot{w} \tag{2.1.2}$$

for the natural projection map. For a face $c \subseteq \mathcal{A}$ let W_c be the subgroup of \widetilde{W} generated by reflections in the hyperplanes through c . The special fibre of \mathbf{T}_1 (as a scheme over \mathfrak{O}) which we denote by $\overline{\mathbf{T}}_1$, is a split maximal torus of $\mathbf{L}_c(\overline{\mathbb{F}}_q)$. Write $\Phi_c(\overline{\mathbf{T}}_1, \overline{\mathbb{F}}_q)$ for the root system of \mathbf{L}_c with respect to $\overline{\mathbf{T}}_1$. Then $\Phi_c(\overline{\mathbf{T}}_1, \overline{\mathbb{F}}_q)$ naturally identifies with the set of $\psi \in \Psi(\mathbf{T}_1, K)$ that vanish on c , and the Weyl group of $\overline{\mathbf{T}}_1$ in \mathbf{L}_c is isomorphic to W_c .

Recall that a choice of x_0 fixes an embedding $\Phi(\mathbf{T}_1, K) \rightarrow \Psi(\mathbf{T}_1, L)$. If we fix a set of simple roots $\Delta \subset \Phi(\mathbf{T}_1, K)$, this embedding determines a set of extended simple roots $\tilde{\Delta} \subseteq \Psi(\mathbf{T}_1, K)$. When $\Phi(\mathbf{T}_1, K)$ is irreducible, $\tilde{\Delta}$ is just the set $\Delta \cup \{1 - \alpha_0\}$ where α_0 is the highest root of $\Phi(\mathbf{T}_1, K)$ with respect to Δ . When $\Phi(\mathbf{T}_1, K)$ is reducible, say $\Phi(\mathbf{T}_1, K) = \cup_i \Phi_i$ where each Φ_i is irreducible, then $\tilde{\Delta} = \cup_i \tilde{\Delta}_i$ where $\tilde{\Delta}_i = \Phi_i \cap \tilde{\Delta}$. Fix Δ so that the chamber cut out by $\tilde{\Delta}$ is c_0 . Let

$$\mathbf{P}(\tilde{\Delta}) := \{J \subsetneq \tilde{\Delta} : J \cap \tilde{\Delta}_i \subsetneq \tilde{\Delta}_i, \forall i\}.$$

Each $J \in \mathbf{P}(\tilde{\Delta})$ cuts out a face of c_0 which we denote by $c(J)$. In particular $c(\Delta) = x_0$. Note that since $\Omega \simeq \widetilde{W}/W \rtimes \mathbb{Z}\Phi(\mathbf{T}_1, K)$ (recall \mathbf{G} is semisimple), and $W \rtimes \mathbb{Z}\Phi(\mathbf{T}_1, K)$ acts simply transitively on the chambers of $\mathcal{A}(\mathbf{T}_1, K)$, the action of \widetilde{W} on $\mathcal{A}(\mathbf{T}_1, K)$ induces an action of Ω on the faces of c_0 and hence on $\tilde{\Delta}$ (and $\mathbf{P}(\tilde{\Delta})$). For $\omega \in \Omega$ let σ_ω denote the corresponding permutation of $\tilde{\Delta}$. Let

$$\mathbf{P}^\omega(\tilde{\Delta}) := \{J \in \mathbf{P}(\tilde{\Delta}) \mid \sigma_\omega(J) = J\}$$

and let c_0^ω be the chamber of $\mathcal{B}(\mathbf{G}^\omega, k)$ lying in c_0 . The set $\mathbf{P}^\omega(\tilde{\Delta})$ is an indexing set for the faces of c_0^ω . For $J \in \mathbf{P}^\omega(\tilde{\Delta})$ write $c^\omega(J)$ for the face of c_0^ω corresponding to J . The face $c^\omega(J)$ lies in $c(J)$. Moreover for $J, J' \in \mathbf{P}^\omega(\tilde{\Delta})$ (resp. $\mathbf{P}(\tilde{\Delta})$) we have $J \subseteq J'$ if and only if $c^\omega(J) \supseteq c^\omega(J')$ (resp. $c(J) \supseteq c(J')$).

2.2 Nilpotent orbits

Let \mathcal{N} be the functor which takes a field F to the set of nilpotent elements in $\mathfrak{g}(F)$, and let \mathcal{N}_o be the functor which takes F to the set of adjoint $\mathbf{G}(F)$ -orbits on $\mathcal{N}(F)$. When F is *k* or *K*, we view $\mathcal{N}_o(F)$ as a partially ordered set with respect to the closure ordering in the topology induced by the topology on F . When F is algebraically closed, we view $\mathcal{N}_o(F)$ as a partially ordered set with respect to the closure ordering in the Zariski topology. For brevity we will write $\mathcal{N}(F'/F)$ (resp. $\mathcal{N}_o(F'/F)$) for $\mathcal{N}(F \rightarrow F')$ (resp. $\mathcal{N}_o(F \rightarrow F')$) where $F \rightarrow F'$ is a morphism of fields. For $(F, F') = (k, K)$ (resp. $(k, \bar{k}), (K, \bar{k})$), the map $\mathcal{N}_o(F'/F)$ is strictly increasing (resp. strictly increasing, non-decreasing). We will simply write \mathcal{N} for $\mathcal{N}(\mathbb{C})$ and \mathcal{N}_o for $\mathcal{N}_o(\mathbb{C})$. In this case we also define $\mathcal{N}_{o,c}$ (resp. $\mathcal{N}_{o,\bar{c}}$) to be the set of all pairs (\mathbb{O}, C) such that $\mathbb{O} \in \mathcal{N}_o$ and C is a conjugacy class in the fundamental group $A(\mathbb{O})$ of \mathbb{O} (resp. Lusztig's canonical quotient $\bar{A}(\mathbb{O})$ of $A(\mathbb{O})$, see [Som01, Section 5]). There is a natural map

$$\mathfrak{Q} : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_{o,\bar{c}}, \quad (\mathbb{O}, C) \mapsto (\mathbb{O}, \bar{C}) \tag{2.2.1}$$

where \bar{C} is the image of C in $\bar{A}(\mathbb{O})$ under the natural homomorphism $A(\mathbb{O}) \rightarrow \bar{A}(\mathbb{O})$. There are also projection maps $\text{pr}_1 : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_o$, $\text{pr}_1 : \mathcal{N}_{o,\bar{c}} \rightarrow \mathcal{N}_o$. We will typically write \mathcal{N}^\vee , \mathcal{N}_o^\vee , $\mathcal{N}_{o,c}^\vee$, and $\mathcal{N}_{o,\bar{c}}^\vee$ for the sets \mathcal{N} , \mathcal{N}_o , $\mathcal{N}_{o,c}$, and $\mathcal{N}_{o,\bar{c}}$ associated to the Langlands dual group G^\vee . When we wish to emphasise the group we are working with we include it as a superscript e.g. $\mathcal{N}_o^{\mathbf{G}^\omega}$. Note that since \mathbf{G}^ω splits over K we have that $\mathcal{N}_o^{\mathbf{G}}(F) = \mathcal{N}_o^{\mathbf{G}^\omega}(F)$ for field extensions F of K .

2.2.1 Classical results and constructions Recall the following classical results and constructions related to nilpotent orbits.

Lemma 2.2.1 (Corollary 3.5, [Pom77] and Theorem 1.5, [Pom80]). *Let F be algebraically closed with good characteristic for \mathbf{G} . Then there is canonical isomorphism of partially ordered sets $\Lambda^F : \mathcal{N}_o^{\mathbf{G}}(F) \xrightarrow{\sim} \mathcal{N}_o$.*

Write

$$d : \mathcal{N}_o \rightarrow \mathcal{N}_o^\vee, \quad d : \mathcal{N}_o^\vee \rightarrow \mathcal{N}_o. \quad (2.2.2)$$

duality maps defined by Spaltenstein ([Spa82, Proposition 10.3]), Lusztig ([Lus84, §13.3], and Barbasch-Vogan ([BV85, Appendix A]). Write

$$d_S : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_o^\vee, \quad d_S : \mathcal{N}_{o,c}^\vee \rightarrow \mathcal{N}_o \quad (2.2.3)$$

for the duality maps defined by Sommers in [Som01, Section 6] and

$$d_A : \mathcal{N}_{o,\bar{c}} \rightarrow \mathcal{N}_{o,\bar{c}}^\vee, \quad d_A : \mathcal{N}_{o,\bar{c}}^\vee \rightarrow \mathcal{N}_{o,\bar{c}} \quad (2.2.4)$$

for the duality maps defined by Achar in ([Ach03, Section 1]). These duality maps are compatible in the following sense. For $\mathbb{O} \in \mathcal{N}_o$

$$d_S(\mathbb{O}, 1) = d(\mathbb{O})$$

and for $(\mathbb{O}, C) \in \mathcal{N}_{o,c}$

$$d_A(\mathfrak{Q}(\mathbb{O}, C)) = (d_S(\mathbb{O}, C), \bar{C}')$$

for some \bar{C}' .

There is a natural pre-order \leq_A on $\mathcal{N}_{o,c}$ defined by Achar in [Ach03, Introduction] by

$$(\mathbb{O}, C) \leq_A (\mathbb{O}', C') \iff \mathbb{O} \leq \mathbb{O}' \text{ and } d_S(\mathbb{O}, C) \geq d_S(\mathbb{O}', C').$$

Write \sim_A for the equivalence relation on $\mathcal{N}_{o,c}$ induced by this pre-order, i.e.

$$(\mathbb{O}_1, C_1) \sim_A (\mathbb{O}_2, C_2) \iff (\mathbb{O}_1, C_1) \leq_A (\mathbb{O}_2, C_2) \text{ and } (\mathbb{O}_2, C_2) \leq_A (\mathbb{O}_1, C_1)$$

Write $[(\mathbb{O}, C)]$ for the equivalence class of $(\mathbb{O}, C) \in \mathcal{N}_{o,c}$. The \sim_A -equivalence classes in $\mathcal{N}_{o,c}$ coincide with the fibres of the projection map $\mathfrak{Q} : \mathcal{N}_{o,c} \rightarrow \mathcal{N}_{o,\bar{c}}$ [Ach03, Theorem 1]. So \leq_A descends to a partial order on $\mathcal{N}_{o,\bar{c}}$, also denoted by \leq_A . The maps d, d_S, d_A are all order reversing with respect to the relevant pre/partial orders.

2.2.2 Structure of $\mathcal{N}_o^{\mathbf{G}}(K)$ In [Oka21, Section 2] the third-named author establishes a number of results about the structure of $\mathcal{N}_o^{\mathbf{G}}(K) = \mathcal{N}_o^{\mathbf{G}^\omega}(K)$ which we now briefly summarize.

Let \mathbf{T} be a maximal \mathfrak{k} -split torus of \mathbf{G}^ω , \mathbf{T}_1 be a maximal K -split torus of \mathbf{G}^ω defined over \mathfrak{k} and containing \mathbf{T} , and x_0 be a special point in $\mathcal{A}(\mathbf{T}_1, K)$. In [Oka21, Section 2.1.5] the third-named author constructs a bijection

$$\theta_{x_0, \mathbf{T}_1} : \mathcal{N}_o^{\mathbf{G}^\omega}(K) \xrightarrow{\sim} \mathcal{N}_{o,c}$$

Theorem 2.2.2. [Oka21, Theorem 2.20, Theorem 2.27, Proposition 2.29] *The bijection*

$$\theta_{x_0, \mathbf{T}_1} : \mathcal{N}_o^{\mathbf{G}^\omega}(K) \xrightarrow{\sim} \mathcal{N}_{o,c}$$

is natural in \mathbf{T}_1 , equivariant in x_0 , and makes the following diagram commute:

$$\begin{array}{ccc} \mathcal{N}_o^{\mathbf{G}^\omega}(K) & \xrightarrow{\theta_{x_0, \mathbf{T}_1}} & \mathcal{N}_{o,c} \\ \mathcal{N}_o(\bar{\mathfrak{k}}/K) \downarrow & & \downarrow \text{pr}_1 \\ \mathcal{N}_o(\bar{\mathfrak{k}}) & \xrightarrow{\Lambda^{\bar{\mathfrak{k}}}} & \mathcal{N}_o. \end{array} \quad (2.2.5)$$

The composition

$$d_{S, \mathbf{T}_1} := d_S \circ \theta_{x_0, \mathbf{T}_1}$$

is independent of the choice of x_0 and natural in \mathbf{T}_1 [Oka21, Proposition 2.32].

For $\mathbb{O}_1, \mathbb{O}_2 \in \mathcal{N}_o(K)$ define $\mathbb{O}_1 \leq_A \mathbb{O}_2$ by

$$\mathbb{O}_1 \leq_A \mathbb{O}_2 \iff \mathcal{N}_o(\bar{k}/K)(\mathbb{O}_1) \leq \mathcal{N}_o(\bar{k}/K)(\mathbb{O}_2), \text{ and } d_{S, \mathbf{T}_1}(\mathbb{O}_1) \geq d_{S, \mathbf{T}_2}(\mathbb{O}_2)$$

and let \sim_A denote the equivalence classes of this pre-order. This pre-order is independent of the choice of \mathbf{T}_1 and the map

$$\theta_{x_0, \mathbf{T}_1} : (\mathcal{N}_o(K), \leq_A) \rightarrow (\mathcal{N}_{o, c}, \leq_A)$$

is an isomorphism of pre-orders.

Theorem 2.2.3. *The composition $\Omega \circ \theta_{x_0, \mathbf{T}_1} : \mathcal{N}_o(K) \rightarrow \mathcal{N}_{o, \bar{c}}$ descends to a (natural in \mathbf{T}_1) bijection*

$$\bar{\theta}_{\mathbf{T}_1} : \mathcal{N}_o(K) / \sim_A \rightarrow \mathcal{N}_{o, \bar{c}}$$

which does not depend on x_0 .

2.2.3 Lifting nilpotent orbits Define

$$\mathcal{I}_o = \{(c, \mathbb{O}) \mid c \subseteq \mathcal{B}(\mathbf{G}), \mathbb{O} \in \mathcal{N}_o^{\mathbf{L}_c}(\bar{\mathbb{F}}_q)\}. \quad (2.2.6)$$

There is a partial order on \mathcal{I}_o , defined by

$$(c_1, \mathbb{O}_1) \leq (c_2, \mathbb{O}_2) \iff c_1 = c_2 \text{ and } \mathbb{O}_1 \leq \mathbb{O}_2$$

In [Oka21, Section 1.1.2] the third author defines a strictly increasing surjective map

$$\mathcal{L} : (\mathcal{I}_o, \leq) \rightarrow (\mathcal{N}_o(K), \leq). \quad (2.2.7)$$

The composition

$$[\bullet] \circ \mathcal{L} : (\mathcal{I}_o, \leq) \rightarrow (\mathcal{N}_o(K) / \sim_A, \leq_A)$$

is also strictly increasing [Oka21, Corollary 4.7, Lemma 5.3].

Recall the groups $G = \mathbf{G}(\mathbb{C}), T = \mathbf{T}(\mathbb{C})$ from section 2. Call a pseudo-Levi subgroup L of G *standard* if it contains T and write Z_L for its center. Let $\mathcal{A} = \mathcal{A}(\mathbf{T}_1, K)$.

Lemma 2.2.4. [Oka21, Section 2.14, Corollary 2.19] *There is a W -equivariant map*

$$\mathfrak{L}_{x_0} : \{\text{faces of } \mathcal{A}\} \rightarrow \{(L, tZ_L^\circ) \mid L \text{ a standard pseudo-Levi, } Z_G^\circ(tZ_L^\circ) = L\} \quad (2.2.8)$$

where c_1, c_2 lie in the same fibre iff

$$\mathcal{A}(c_1, \mathcal{A}) + X_*(\mathbf{T}_1, K) = \mathcal{A}(c_2, \mathcal{A}) + X_*(\mathbf{T}_1, K).$$

Moreover, if $\mathfrak{L}_{x_0}(c) = (L, tZ_L^\circ)$ then L is the complex reductive group with the same root datum as $\mathbf{L}_c(\bar{\mathbb{F}}_q)$ and thus there is an isomorphism $\Lambda_c^{\bar{\mathbb{F}}_q} : \mathcal{N}_o^{\mathbf{L}_c}(\bar{\mathbb{F}}_q) \xrightarrow{\sim} \mathcal{N}_o^L$.

Recall from the end of section 2.1 the definitions of $\Delta, \tilde{\Delta}, c_0, \mathbf{P}(\tilde{\Delta})$ and $c(J)$. Note that the definitions of c_0 and $c(J)$ depend on a choice of x_0 and Δ . Let L_J denote the pseudo-Levi subgroup of G generated by T and the root groups corresponding to $\dot{\alpha}$ for $\alpha \in J$. Then $\text{pr}_1 \circ \mathfrak{L}_{x_0}(c(J)) = L_J$ (indeed by [Oka21, Lemma 2.21], this group should not depend on x_0). Define

$$\mathcal{I}_{x_0, \tilde{\Delta}} = \{(J, \mathbb{O}) \mid J \in \mathbf{P}(\tilde{\Delta}), \mathbb{O} \in \mathcal{N}_o^{\mathbf{L}_{c(J)}}(\bar{\mathbb{F}}_q)\}, \quad (2.2.9)$$

$$\mathcal{K}_{\tilde{\Delta}} = \{(J, \mathbb{O}) \mid J \in \mathbf{P}(\tilde{\Delta}), \mathbb{O} \in \mathcal{N}_o^{L_J}(\mathbb{C})\}. \quad (2.2.10)$$

The map

$$\iota_{x_0} : \mathcal{I}_{x_0, \bar{\Delta}} \rightarrow \mathcal{K}_{\bar{\Delta}}, \quad (J, \mathbb{O}) \mapsto (J, \Lambda_{c(J)}^{\overline{\mathbb{F}}_q}(\mathbb{O}))$$

is an isomorphism. Let

$$\mathbb{L} : \mathcal{K}_{\bar{\Delta}} \rightarrow \mathcal{N}_{c, o} \tag{2.2.11}$$

be the map that sends (J, \mathbb{O}) to $(Gx, tZ_G^\circ(x))$ where $x \in \mathbb{O}$ and $(L, tZ_L^\circ) = \mathfrak{L}(c(J))$ where \mathfrak{L} is the map from [Oka21, Corollary 2.19]. By [CMO21, Theorem 2.1.7] the diagram

$$\begin{array}{ccc} \mathcal{I}_{x_0, \bar{\Delta}} & \xrightarrow{\sim} & \mathcal{K}_{\bar{\Delta}} \\ \downarrow \mathcal{L} & & \downarrow \mathbb{L} \\ \mathcal{N}_o(K) & \xrightarrow{\sim} & \mathcal{N}_{o, c} \end{array} \tag{2.2.12}$$

commutes. Define

$$\overline{\mathbb{L}} = \mathfrak{Q} \circ \mathbb{L}.$$

This map can be computed using Achar's algorithms in [Ach03, Section 3.4].

2.3 Wavefront sets

Let X be an admissible smooth representation of $\mathbf{G}^\omega(\mathfrak{k})$ and let Θ_X be the character of X . Recall that for each nilpotent orbit $\mathbb{O} \in \mathcal{N}_o^{\mathbf{G}^\omega}(\mathfrak{k})$ there is an associated distribution $\mu_{\mathbb{O}}$ on $C_c^\infty(\mathfrak{g}^\omega(\mathfrak{k}))$ called the *nilpotent orbital integral* of \mathbb{O} [Ran72]. Write $\hat{\mu}_{\mathbb{O}}$ for the Fourier transform of this distribution. Generalizing a result of Howe ([How74]), Harish-Chandra in [Har99] showed that there are complex numbers $c_{\mathbb{O}}(X) \in \mathbb{C}$ such that

$$\Theta_X(\exp(\xi)) = \sum_{\mathbb{O}} c_{\mathbb{O}}(X) \hat{\mu}_{\mathbb{O}}(\xi) \tag{2.3.1}$$

for $\xi \in \mathfrak{g}^\omega(\mathfrak{k})$ a regular element in a small neighborhood of 0. The formula (2.3.1) is called the *local character expansion* of π . The (*p-adic*) *wavefront set* of X is

$$\text{WF}(X) := \max\{\mathbb{O} \mid c_{\mathbb{O}}(X) \neq 0\} \subseteq \mathcal{N}_o(\mathfrak{k}).$$

The *geometric wavefront set* of X is

$$\bar{\mathfrak{k}}\text{WF}(X) := \max\{\mathcal{N}_o(\bar{\mathfrak{k}}/\mathfrak{k})(\mathbb{O}) \mid c_{\mathbb{O}}(X) \neq 0\} \subseteq \mathcal{N}_o(\bar{\mathfrak{k}}),$$

see [Wal18, p. 1108] (warning: in [Wal18], the invariant $\bar{\mathfrak{k}}\text{WF}(X)$ is called simply the ‘wavefront set’ of X).

In [Oka21, Section 2.2.3] the third author has introduced a third type of wavefront set for depth-0 representations, called the *canonical unramified wavefront set*. This invariant is a natural refinement of $\bar{\mathfrak{k}}\text{WF}(X)$. We will now define ${}^K\text{WF}(X)$ and explain how to compute it.

Recall from Equation 2.2.7 the lifting map \mathcal{L} . For every face $c \subseteq \mathcal{B}(\mathbf{G})$, the space of invariants $X^{\mathbf{U}_c(o)}$ is a (finite-dimensional) $\mathbf{L}_c(\mathbb{F}_q)$ -representation. Let $\text{WF}(X^{\mathbf{U}_c(o)}) \subseteq \mathcal{N}_o^{\mathbf{L}_c}(\overline{\mathbb{F}}_q)$ denote the Kawanaka wavefront set [Kaw87] and let

$${}^K\text{WF}_c(X) := \{[\mathcal{L}(c, \mathbb{O})] \mid \mathbb{O} \in \text{WF}(X^{\mathbf{U}_c(o)})\} \subseteq \mathcal{N}_o(K). \tag{2.3.2}$$

Definition 2.3.1. *Let X be a depth-0 representation of $\mathbf{G}^\omega(\mathfrak{k})$. The canonical unramified wavefront set of X is*

$${}^K\text{WF}(X) := \max\{{}^K\text{WF}_c(X) \mid c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})\} \subseteq \mathcal{N}_o(K) / \sim_A. \tag{2.3.3}$$

Fix $\mathbf{T}, \mathbf{T}_1, c_0, x_0, \Delta$ as at the end of section 2.1. By [Oka21, Lemma 2.36] we have that

$${}^K\text{WF}(X) = \max\{{}^K\text{WF}_{c(J)}(X) \mid J \in \mathbf{P}^\omega(\bar{\Delta})\}. \tag{2.3.4}$$

We will often want to view ${}^K\text{WF}(X)$ and ${}^K\text{WF}_c(X)$ as subsets of $\mathcal{N}_{o, \bar{\varepsilon}}$ using the identification $\bar{\theta}_{\mathbf{T}_1}$ from Theorem 2.2.3. We will write

$${}^K\text{WF}(X, \mathbb{C}) := \bar{\theta}_{\mathbf{T}_1}({}^K\text{WF}(X)), \quad {}^K\text{WF}_c(X, \mathbb{C}) := \bar{\theta}_{\mathbf{T}_1}({}^K\text{WF}_c(X)).$$

We will also want to view $\bar{k}\text{WF}(X)$, which is naturally contained in $\mathcal{N}_o(\bar{k})$, as a subset of \mathcal{N}_o and so will write

$$\bar{k}\text{WF}(X, \mathbb{C}) := \Lambda^{\bar{k}}(\bar{k}\text{WF}(X)).$$

By [Oka21, Theorem 2.37], if ${}^K\text{WF}(X)$ is a singleton, then $\bar{k}\text{WF}(X)$ is also a singleton and

$${}^K\text{WF}(X, \mathbb{C}) = (\bar{k}\text{WF}(X, \mathbb{C}), \bar{C}). \quad (2.3.5)$$

for some conjugacy class \bar{C} in $\bar{A}(\bar{k}\text{WF}(X, \mathbb{C}))$.

2.4 Isogenies

Let $f : \mathbf{H}' \rightarrow \mathbf{H}$ be an isogeny of connected reductive groups defined over k . Let $f_k : \mathbf{H}'(k) \rightarrow \mathbf{H}(k)$ denote the corresponding homomorphism of k -points. We note that $\mathcal{N}_{o, \bar{c}}^{\mathbf{H}} \simeq \mathcal{N}_{o, \bar{c}}^{\mathbf{H}'}$ and so we can compare the canonical unramified wavefront sets of representations of the two groups. In an upcoming paper [Oka22] the third author proves the following result about the behaviour of the canonical unramified wavefront set under isogeny.

Lemma 2.4.1. *Let X be an irreducible admissible depth-0 representation of $\mathbf{H}(k)$ and write X' for the representation of $\mathbf{H}'(k)$ obtained by pulling back along f_k . Then X' decomposes as a finite sum of irreducible admissible representations $X' = \bigoplus_i X'_i$ and ${}^K\text{WF}(X) = {}^K\text{WF}(X'_i)$ for all i .*

2.5 Depth-0 representations

Let $\omega \in \Omega$. Recall that a smooth irreducible representation X of $\mathbf{G}^\omega(k)$ has depth-0 if $X^{\mathbf{U}_c(\mathfrak{o})} \neq 0$ for some $c \subseteq \mathcal{B}(\mathbf{G}^\omega, k)$. Write $\Pi^0(\mathbf{G}^\omega(k))$ for the subset of $\Pi(\mathbf{G}^\omega(k))$ consisting of depth-0 representations. Let

$$S^\omega := \{(c, \sigma) : c \subseteq \mathcal{B}(\mathbf{G}^\omega, k), \sigma \text{ a cuspidal representation of } \mathbf{L}_c(\mathbb{F}_q)\}$$

and for $(c_1, \sigma_1), (c_2, \sigma_2) \in S^\omega$ write $(c_1, \sigma_1) \sim (c_2, \sigma_2)$ if they are associate in the sense of [MP96, Section 5]. Suppose $(c_1, \sigma_1), (c_2, \sigma_2) \in S^\omega$ are such that σ_i is a subrepresentation of $X^{\mathbf{U}_{c_i}(\mathfrak{o})}$ for $i \in \{1, 2\}$. Then by [MP96, Theorem 5.2], we have that $(c_1, \sigma_1) \sim (c_2, \sigma_2)$. Thus there is a well defined map

$$\text{supp} : \Pi^0(\mathbf{G}^\omega(k)) \rightarrow S^\omega / \sim$$

which attaches to X the well-defined association class of $(c, \sigma) \in S^\omega$ where (c, σ) is such that σ appears as a subrepresentation of $X^{\mathbf{U}_c(\mathfrak{o})}$. This association class called the unrefined minimal K-type of X , but we write $\text{supp}(X)$ for brevity.

Note that for $(c_1, \sigma_1), (c_2, \sigma_2) \in S^\omega$, if $(c_1, \sigma_1) \sim (c_2, \sigma_2)$ then $(c_1, \text{WF}(\sigma_1)) \sim_K (c_2, \text{WF}(\sigma_2))$ where WF denotes the Kawanaka wavefront set and \sim_K is the equivalence relation defined in [Oka21, Section 1.1.1]. In particular, since \mathcal{L} is constant on \sim_K -classes we have that

$$\mathcal{L}(c_1, \text{WF}(\sigma_1)) = \mathcal{L}(c_2, \text{WF}(\sigma_2)). \quad (2.5.1)$$

For $X \in \Pi^0(\mathbf{G}^\omega(k))$ define ${}^K\text{WF}_{\text{supp}}(X)$ to be \sim_A -class of the well-defined orbit in Equation 2.5.1 applied to the association class of $\text{supp}(X)$. We will write

$${}^K\text{WF}_{\text{supp}}(X, \mathbb{C}) := \bar{\theta}_{\mathbf{T}_1}({}^K\text{WF}_{\text{supp}}(X)).$$

Lemma 2.5.1. *For $X \in \Pi^0(\mathbf{G}^\omega(k))$ we have that*

$${}^K\text{WF}_{\text{supp}}(X) \leq_A \mathbb{O}$$

for some $\mathbb{O} \in {}^K\text{WF}(X)$. In particular, when ${}^K\text{WF}(X)$ is a singleton we have

$${}^K\text{WF}_{\text{supp}}(X) \leq_A {}^K\text{WF}(X).$$

Proof. Let $\text{supp}(X) = [(c, \sigma)]$. Then σ is a subrepresentation of $X^{\mathbf{U}_c(\mathfrak{o})}$ and so $\text{WF}(\sigma) \leq \mathbb{O}$ for some $\mathbb{O} \in \text{WF}(X^{\mathbf{U}_c(\mathfrak{o})})$. Thus

$$[\mathcal{L}(c, \text{WF}(\sigma))] \leq_A [\mathcal{L}(c, \mathbb{O})] \in {}^K\text{WF}_c(X).$$

The result then follows from the fact that

$${}^K\text{WF}(X) = \max\{{}^K\text{WF}_c(X) : c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})\}.$$

□

Remark 2.5.2. *In fact the analogous result for the unramified wavefront set (as defined in [Oka21, Section 1.0.5]), holds as well.*

For an Iwahori-spherical representation X this inequality says nothing because $\text{supp}(X) = [(c_0, \text{triv})]$ and so ${}^K\text{WF}_{\text{supp}}(X, \mathbb{C}) = (\{0\}, 1)$. For supercuspidal representations however, the inequality is in fact an equality. We now proceed to prove this.

Lemma 2.5.3. *Let X be a depth-0 supercuspidal representation. Let c' be a face of $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ with $X^{\mathbf{U}_{c'}(\mathfrak{o})} \neq 0$ and suppose that τ is an irreducible constituent of $X^{\mathbf{U}_{c'}(\mathfrak{o})}$. Then τ is a cuspidal representation of $\mathbf{L}_{c'}(\mathbb{F}_q)$ and in particular $[(c', \tau)] = \text{supp}(X)$.*

Proof. Let $\text{supp}(X) = [(c, \sigma)]$ and c', τ be as in the statement of the lemma. Let $[(M, \tau')]$ be the cuspidal data for τ (i.e. a conjugacy class of Levi of $\mathbf{L}_{c'}(\mathbb{F}_q)$ and cuspidal representation of said Levi). In particular, if M is included into any parabolic P so that P has Levi decomposition $P = MU$, then τ' is a subrepresentation of τ^U . Now, all of the parabolics of $\mathbf{L}_{c'}(\mathfrak{o})$ are conjugate to a parabolic of the form $\mathbf{P}_{c''}(\mathfrak{o})/\mathbf{U}_{c'}(\mathfrak{o})$ where $c' \subseteq \overline{c''}$. Thus (conjugating M appropriately) we can find a c'' such that M is a Levi factor of $\mathbf{P}_{c''}(\mathfrak{o})/\mathbf{U}_{c'}(\mathfrak{o})$ and so $\mathbf{L}_{c''}(\mathbb{F}_q) \simeq M$. We thus have that

$$\tau' \subseteq \tau^U \subseteq (X^{\mathbf{U}_{c'}(\mathfrak{o})})^{\mathbf{U}_{c''}(\mathfrak{o})/\mathbf{U}_{c'}(\mathfrak{o})} = X^{\mathbf{U}_{c''}(\mathfrak{o})}.$$

In particular (c'', τ') is an unrefined minimal K -type for X . Thus by [MP96, Theorem 5.2], we have that $(c'', \tau') \sim (c, \sigma)$. In particular c'' is also a minimal face and so $c'' = c'$ and $\tau = \tau'$. Thus τ is a cuspidal representation of $\mathbf{L}_{c'}(\mathbb{F}_q)$ and $[(c, \sigma)] = [(c', \tau)]$. □

Proposition 2.5.4. *Let X be a depth-0 supercuspidal representation of $\mathbf{G}^\omega(\mathfrak{k})$. Then*

$${}^K\text{WF}_{\text{supp}}(X) = {}^K\text{WF}(X).$$

Proof. Recall that

$${}^K\text{WF}(X) = \max\{{}^K\text{WF}_c(X) : c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})\}.$$

Suppose $c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ is such that $X^{\mathbf{U}_c(\mathfrak{o})} \neq 0$. Then for any τ an irreducible constituent of $X^{\mathbf{U}_c(\mathfrak{o})}$ we have by Lemma 2.5.3 that $[(c, \tau)] = \text{supp}(X)$. Thus we must have that ${}^K\text{WF}_c(X) = {}^K\text{WF}_{\text{supp}}(X)$. If $c \subseteq \mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ is such that $X^{\mathbf{U}_c(\mathfrak{o})} = 0$ then ${}^K\text{WF}_c(X, \mathbb{C}) = (\{0\}, 1)$. Thus

$${}^K\text{WF}(X) = {}^K\text{WF}_{\text{supp}}(X).$$

□

Remark 2.5.5. *Although we have only stated the results for inner twists of the split group, the proposition in fact holds for all connected reductive groups (with identical proof).*

Note that since ${}^K\text{WF}_{\text{supp}}$ is a singleton by construction, this establishes Moeglin and Waldspurger's conjecture for all depth-0 supercuspidal representations for connected reductive groups.

2.6 Unipotent supercuspidal representations

Let

$$S_{unip}^\omega := \{(c, \sigma) \in S^\omega : \sigma \text{ is unipotent}\}.$$

Definition 2.6.1. *Let X be an irreducible $\mathbf{G}^\omega(\mathfrak{k})$ -representation. We say that X has unipotent cuspidal support if $\text{supp}(X) \in S_{unip}^\omega$. Write $\Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathfrak{k}))$ for the subset of $\Pi(\mathbf{G}(\mathfrak{k}))$ consisting of all such representations.*

Call a supercuspidal representation unipotent if it has cuspidal unipotent support. By [Mor96], the irreducible unipotent supercuspidal representations are all obtained by compact induction $\text{ind}_{\mathbf{P}_c^\dagger(\mathfrak{o})}^{\mathbf{G}^\omega(\mathfrak{k})}(\sigma^\dagger)$, where $\mathbf{P}_c(\mathfrak{o})$ is a maximal parahoric subgroup and σ^\dagger contains an irreducible Deligne-Lusztig cuspidal unipotent representation σ of $\mathbf{L}_c(\mathbb{F}_q)$ upon restriction to $\mathbf{L}_c(\mathbb{F}_q)$. In particular $X \in \Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathfrak{k}))$ is supercuspidal if and only if $\text{supp}(X) = [(c, \sigma)]$ where c is a minimal face of $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$.

2.7 Langlands classification of unipotent supercuspidal representations

Let $W_{\mathfrak{k}}$ be the Weil group of \mathfrak{k} with inertia subgroup $I_{\mathfrak{k}}$ and set $W'_{\mathfrak{k}} = W_{\mathfrak{k}} \times SL(2, \mathbb{C})$. We will think of a *Langlands parameter* for \mathbf{G} as a continuous morphism $\varphi : W'_{\mathfrak{k}} \rightarrow G^\vee$ such that $\varphi(w)$ is semisimple for each $w \in W_{\mathfrak{k}}$ and the restriction of φ to $SL(2, \mathbb{C})$ is algebraic. A Langlands parameter φ is called *unramified* if $\varphi(I_{\mathfrak{k}}) = \{1\}$. Let $G^\vee(\varphi)$ denote the centralizer of $\varphi(W'_{\mathfrak{k}})$ in G^\vee . Define

$$Z_{G_{sc}^\vee}^1(\varphi) = \text{preimage of } G^\vee(\varphi)/Z(G^\vee) \text{ under the projection } G_{sc}^\vee \rightarrow G_{ad}^\vee,$$

and let A_φ^1 denote the component group of $Z_{G_{sc}^\vee}^1(\varphi)$. An *enhanced Langlands parameter* is pair (φ, ρ) , where $\rho \in \text{Irr}(A_\varphi^1)$. A parameter (φ, ρ) is called *\mathbf{G}^ω -relevant* (recall that \mathbf{G}^ω is an inner twist of the split form, $\omega \in \Omega_{ad}$) if ρ acts on $Z(G_{sc}^\vee)$ by a multiple of the character ζ_ω .

Define the elements

$$s_\varphi = \varphi(\text{Frob}, 1), \quad u_\varphi = \varphi\left(1, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right).$$

Following [AMS18], consider the possibly disconnected reductive group

$$\mathcal{G}_\varphi = Z_{G_{sc}^\vee}^1(\varphi(W_{\mathfrak{k}})),$$

which is defined analogously to $Z_{G_{sc}^\vee}^1(\varphi)$. Then $u_\varphi \in \mathcal{G}_\varphi^\circ$ and by [AMS18, (92)]

$$A_\varphi^1 \cong \mathcal{G}_\varphi(u_\varphi)/\mathcal{G}_\varphi(u_\varphi)^\circ.$$

An enhanced Langlands parameter (φ, ρ) is called *discrete* if $G^\vee(\varphi)$ does not contain a nontrivial torus (this notion is independent of ρ). A discrete parameter is called *cuspidal* if (u_φ, ρ) is a cuspidal pair. This means that every ρ° which occurs in the restriction of ρ to $A_{\mathcal{G}_\varphi^\circ}(u_\varphi)$ defines a $\mathcal{G}_\varphi^\circ$ -equivariant local system on the $\mathcal{G}_\varphi^\circ$ -conjugacy class of u_φ which is cuspidal in the sense of Lusztig.

A Langlands correspondence for unipotent supercuspidal representations has been obtained by [Mor96] when \mathbf{G} is simple and adjoint, see also [Lus95]. For arbitrary reductive K -split groups, this correspondence is available by [FOS19; FO20]. Let $\text{Irr}(\mathbf{G}^\omega(\mathfrak{k}))_{\text{cusp, unip}}$ denote the set of equivalence classes of irreducible unipotent supercuspidal $\mathbf{G}^\omega(\mathfrak{k})$ -representations. Let $\Phi(G^\vee)_{\text{cusp, nr}}^\omega$ denote the set of G^\vee -equivalence classes of unramified cuspidal enhanced Langlands parameters (φ, ρ) which are \mathbf{G}^ω -relevant.

Theorem 2.7.1. *For every $\omega \in \Omega_{ad}$, there is a bijection*

$$\Phi(G^\vee)_{\text{cusp, nr}}^\omega \longleftrightarrow \text{Irr}(\mathbf{G}^\omega(\mathfrak{k}))_{\text{cusp, unip}}.$$

This bijection satisfies several natural desiderata (including formal degrees, equivariance with respect to tensoring by weakly unramified characters), see [FOS19, Theorem 2].

For X a unipotent supercuspidal representation of $\mathbf{G}^\omega(k)$ let φ denote the corresponding Langlands parameter. We will write $\mathbb{O}_X^\vee \in \mathcal{N}_o^\vee$ for the G^\vee -orbit of

$$n_\varphi = d\varphi(0, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}).$$

Lemma 2.7.2. *Let $X, X' \in \text{Irr}(\mathbf{G}^\omega(k))_{\text{cusp,unip}}$. If $\text{supp}(X) = \text{supp}(X')$ then $\mathbb{O}_X^\vee = \mathbb{O}_{X'}^\vee$.*

Proof. This follows by inspecting the explicit classification in [FOS19]. □

For $[(c, \sigma)] \in S^\omega$ with c a minimal face we write $\mathbb{O}^\vee(c, \sigma)$ for the common nilpotent parameter of all $X \in \text{Irr}(\mathbf{G}^\omega(k))_{\text{cusp,unip}}$ with $\text{supp}(X) = [(c, \sigma)]$.

We will recall the explicit classification in the section 5.

3 Main result

Proposition 3.0.1. *Suppose \mathbf{G} is simple and adjoint and let $[(c, \sigma)] \in S^\omega$ be such that c is a minimal face. Then*

$$\overline{\mathbb{L}}(c, \text{WF}(\sigma)) = d_A(\mathbb{O}^\vee(c, \sigma), 1).$$

This proposition will be proved in Section 6.

Theorem 3.0.2. *Let \mathbf{G} be a split semisimple group defined over k . Let $\omega \in \Omega$ and let \mathbf{G}^ω denote the corresponding inner twist of \mathbf{G} . Let X be an irreducible supercuspidal $\mathbf{G}^\omega(k)$ -representation with unipotent cuspidal support. Then ${}^K\text{WF}(X), {}^{\bar{k}}\text{WF}(X)$ are singletons, and*

$$\begin{aligned} {}^K\text{WF}(X, \mathbb{C}) &= d_A(\mathbb{O}_X^\vee, 1) \\ {}^{\bar{k}}\text{WF}(X, \mathbb{C}) &= d(\mathbb{O}_X^\vee). \end{aligned}$$

Proof. Suppose first that \mathbf{G} is simple and adjoint.

Let X be a unipotent supercuspidal representation of $\mathbf{G}^\omega(k)$. By Proposition 2.5.4 we have that

$${}^K\text{WF}(X) = {}^K\text{WF}_{\text{supp}}(X).$$

Write $\text{supp}(X) = [(c, \sigma)]$. By definition

$${}^K\text{WF}_{\text{supp}}(X, \mathbb{C}) = \overline{\mathbb{L}}(c, \text{WF}(\sigma)).$$

By Proposition 3.0.1 we have that

$$\overline{\mathbb{L}}(c, \text{WF}(\sigma)) = d_A(\mathbb{O}^\vee(c, \sigma), 1).$$

Since $\mathbb{O}_X^\vee = \mathbb{O}^\vee(c, \sigma)$ we get that

$${}^K\text{WF}(X; \mathbb{C}) = d_A(\mathbb{O}_X^\vee, 1)$$

as required.

Applying Lemma 2.4.1 we get that the theorem holds for all simply-connected simple groups. Since wavefront sets behave as expected with respect to products, the theorem holds for all simply-connected semisimple groups. Finally, applying Lemma 2.4.1 again we get that the theorem holds for all split semisimple groups \mathbf{G} . □

4 Unipotent cuspidal representations of finite reductive groups

For the explicit results about the parametrization of unipotent representations of finite reductive groups, we refer to [Lus84, §4, §8.1] and [Car93, §13.8, §13.9]. The relevant results for the Kawanaka wavefront sets and unipotent support are in [Lus92, §10, §11]. The classification of unipotent representations is independent of the isogeny, so in this section, we may assume without loss of generality that the group G is simple and adjoint.

4.1 Classical groups

4.1.1 $A_{n-1}(q)$. The group $G = PGL(n)$ does not have unipotent cuspidal representations.

4.1.2 ${}^2A_n(q^2)$. The group $G = PU(n+1)$ has unipotent representations if and only if $n = \frac{r(r+1)}{2} - 1$, for some integer $r \geq 2$. The unipotent ${}^2A_n(q^2)$ -representations are in one-to-one correspondence with partitions of $n+1$, and so are the geometric nilpotent orbits of G . When $n = \frac{r(r+1)}{2} - 1$, the cuspidal unipotent representation σ is unique and it is parametrized by the partition

$$(1, 2, 3, \dots, r).$$

The Kawanaka wavefront set is $\text{WF}(\sigma) = (1, 2, 3, \dots, r)$.

4.1.3 $B_n(q), C_n(q)$. Suppose G is $SO(2n+1)$ or $PSp(2n)$ over \mathbb{F}_q . The group $G(\mathbb{F}_q)$ has a unipotent cuspidal representation (and in this case the cuspidal representation is unique) if and only if $n = r^2 + r$ for a positive integer r . The unipotent representations of $G(\mathbb{F}_q)$ are parametrized by symbols

$$\begin{pmatrix} \lambda_1 & \lambda_2 & \dots & \lambda_a \\ & \mu_1 & \dots & \mu_b \end{pmatrix},$$

$0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_a$, $0 \leq \mu_1 < \mu_2 < \dots < \mu_b$, $a - b$ odd and positive, and λ_1, μ_1 are not both zero, such that

$$n = \sum \lambda_i + \sum \mu_j - \left(\frac{a+b-1}{2} \right)^2.$$

Let $d = a - b$ be the defect of the symbol. Two unipotent representations belong to the same family if their symbols have the same entries with the same multiplicities. For the unipotent cuspidal representation σ , the corresponding symbol has defect $d = 2r + 1$ and it is

$$\begin{pmatrix} 0 & 1 & 2 & \dots & 2r \\ & & - & & \end{pmatrix}.$$

The geometric nilpotent orbits of $SO(2n+1)$ (resp., $PSp(2n)$) are parametrized by partitions of $2n+1$ (resp., $2n$), where the even (resp., odd) parts occur with even multiplicity. The Kawanaka wavefront set of the unipotent cuspidal representation σ is

$$\text{WF}(\sigma) = \begin{cases} (1, 1, 3, 3, \dots, 2r-1, 2r-1, 2r+1), & G = SO(2n+1) \\ (2, 2, 4, 4, \dots, 2r, 2r), & G = PSp(2n). \end{cases} \quad (4.1.1)$$

4.1.4 $D_n(q)$. Suppose G is the split orthogonal group $PSO(2n)$ over \mathbb{F}_q . There exists a unipotent cuspidal representation (and in this case it is unique) if and only if $n = r^2$ for a positive even integer r . The type D_n -symbols are

$$\begin{pmatrix} \lambda_1 & \lambda_2 & \dots & \lambda_a \\ & \mu_1 & \dots & \mu_b \end{pmatrix},$$

$0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_a$, $0 \leq \mu_1 < \mu_2 < \dots < \mu_b$, $a - b$ is divisible by 4, and λ_1, μ_1 are not both zero, such that

$$n = \sum \lambda_i + \sum \mu_j - \frac{(a+b)(a+b-2)}{4}.$$

One symbol and the symbol if the row swapped are regarded the same. The irreducible unipotent $G(\mathbb{F}_q)$ -representations are in one-to-one correspondence with the type D_n -symbols, except if the symbol has identical rows, then there are two nonisomorphic irreducible unipotent representations attached to it. The defect $d = a - b$ is even.

For the unipotent cuspidal representation σ , the corresponding symbol has defect $d = 2r$ and it is

$$\begin{pmatrix} 0 & 1 & 2 & \dots & 2r-1 \\ & & - & & \end{pmatrix}.$$

The geometric nilpotent orbits of $PSO(2n)$ are parametrized by partitions of $2n$ with the even parts occurring with even multiplicity. The Kawanaka wavefront set of the unipotent cuspidal representation is

$$\text{WF}(\sigma) = (1, 1, 3, 3, \dots, 2r-1, 2r-1). \quad (4.1.2)$$

4.1.5 ${}^2D_n(q^2)$. The group ${}^2D_n(q^2)$ admits unipotent cuspidal representations if and only if $n = r^2$, for some odd positive integer r , and in this case the unipotent cuspidal representation is unique. The type 2D_n -symbols are

$$\left(\begin{array}{cccc} \lambda_1 & \lambda_2 & \dots & \lambda_a \\ & \mu_1 & & \mu_b \end{array} \right),$$

$0 \leq \lambda_1 < \lambda_2 < \dots < \lambda_a$, $0 \leq \mu_1 < \mu_2 < \dots < \mu_b$, $a - b \equiv 2 \pmod{4}$, and λ_1, μ_1 are not both zero, such that

$$n = \sum \lambda_i + \sum \mu_j - \frac{(a+b)(a+b-2)}{4}.$$

One symbol and the symbol if the row swapped are regarded the same. The irreducible unipotent ${}^2D_n(q^2)$ -representations are in one-to-one correspondence with the type 2D_n -symbols.

For the unipotent cuspidal representation σ , the corresponding symbol and Kawanaka wavefront set are the same as in the split case $D_n(q)$ (except r is now odd).

4.1.6 ${}^3D_4(q^3)$ The group ${}^3D_4(q^3)$ has eight unipotent representations: six are in the principal series, in one-to-one correspondence with the irreducible representations of the Weyl group of type G_2 , and two unipotent cuspidal representations, denoted ${}^3D_4[1]$ and ${}^3D_4[-1]$.

The geometric nilpotent orbits of ${}^3D_4(q^3)$ are parametrized by partitions of 8 with even parts occurring with even multiplicity. The unipotent cuspidal representations have Kawanaka wavefront set

$$\text{WF}({}^3D_4[1]) = \text{WF}({}^3D_4[-1]) = (1, 1, 3, 3). \quad (4.1.3)$$

4.2 Exceptional groups

Suppose $G(\mathbb{F}_q)$ is \mathbb{F}_q -split. In the table below, we will list all unipotent cuspidal $G(\mathbb{F}_q)$ -representations. The irreducible unipotent representations of $G(\mathbb{F}_q)$ are partitioned into families, each family being in one-to-one correspondence with the set

$$M(\Gamma) = \Gamma\text{-orbits in } \{(x, \tau) \mid x \in \Gamma, \tau \in \widehat{Z_\Gamma(x)}\},$$

for a finite group Γ . Each group Γ is uniquely attached to a special nilpotent orbit \mathbb{O}^\vee in the dual Lie algebra, such that $\Gamma = \bar{A}(\mathbb{O}^\vee)$, where $\bar{A}(\mathbb{O}^\vee)$ is Lusztig's canonical quotient.

In Table 1, for each unipotent cuspidal representation σ , we will record the corresponding Kawanaka wavefront set, the nilpotent orbit \mathbb{O}^\vee corresponding to σ and its canonical quotient $\bar{A}(\mathbb{O}^\vee)$, the pair $(x, \tau) \in M(\bar{A}(\mathbb{O}^\vee))$ that parametrizes σ . The geometric nilpotent orbits are given in the Bala-Carter notation.

$G(\mathbb{F}_q)$	Cuspidal σ	$\text{WF}(\sigma)$	\mathbb{O}_σ^\vee	$\bar{A}(\mathbb{O}^\vee)$	(x, τ)
G_2	$G_2[1]$ $G_2[-1]$ $G_2[\theta^l], l = 1, 2$	$G_2(a_1)$	$G_2(a_1)$	S_3	$(1, \epsilon)$ (g_2, ϵ) (g_3, θ^l)
F_4	$F_4^{III}[1]$ $F_4[-1]$ $F_4^I[1]$ $F_4[\theta^l], l = 1, 2$ $F_4[\pm i]$	$F_4(a_3)$	$F_4(a_3)$	S_4	$(1, \lambda^3)$ (g_2, ϵ) (g'_2, ϵ) (g_3, θ^l) $(g_4, \pm i)$
E_6	$E_6[\theta^l], l = 1, 2$	$D_4(a_1)$	$D_4(a_1)$	S_3	(g_3, θ^l)
E_7	$E_7[\zeta]$ $E_7[\zeta]$	$A_4 + A_1$	$A_4 + A_1$	$\mathbb{Z}/2$	$(g_2, 1)$ (g_2, ϵ)

E_8	$E_8^{II}[1]$ $E_8[-1]$ $E_8^I[1]$ $E_8[\theta^l], l = 1, 2$ $E_8[-\theta^l], l = 1, 2$ $E_8[\pm i]$ $E_8[\zeta^j], 1 \leq j \leq 4$	$E_8(a_7)$	$E_8(a_7)$	S_5	$(1, \lambda^4)$ $(g_2, -\epsilon)$ (g'_2, ϵ) $(g_3, \epsilon\theta^l)$ $(g_6, -\theta^l)$ $(g_4, \pm i)$ (g_5, ζ^j)
-------	--	------------	------------	-------	---

Table 1: Unipotent cuspidal representations of exceptional groups $G(\mathbb{F}_q)$

Finally, for the twisted group ${}^2E_6(q^2)$, there are three unipotent cuspidal representations, denoted ${}^2E_6[1]$, ${}^2E_6[\theta]$, ${}^2E_6[\theta^2]$. All three of them have Kawanaka wavefront set $D_4(a_1)$ in E_6 .

5 Langlands parameters for unipotent supercuspidal representations

Recall that $X \in \Pi^{\text{Lus}}(\mathbf{G}^\omega(\mathfrak{k}))$ is supercuspidal if and only if $\text{supp}(X) = [(c, \sigma)]$ where c is a minimal face of $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$. Since every association class of faces of $\mathcal{B}(\mathbf{G}^\omega, \mathfrak{k})$ contains a face of c_0^ω we may assume that c is of the form $c^\omega(J)$ for some $J \in \mathbf{P}^\omega(\tilde{\Delta})$. Moreover $c^\omega(J)$ is a minimal face if and only if J is maximal in $\mathbf{P}^\omega(\tilde{\Delta})$. In this section we list the set of possible pairs (J, σ) (up to \sim) along with $\mathbb{O}^\vee(c^\omega(J), \sigma)$, where $J \in \mathbf{P}^\omega(\tilde{\Delta})$ is maximal and σ is a unipotent cuspidal representation of $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$, for \mathbf{G} split, simple and adjoint, and $\omega \in \Omega$. We use the conventions of [Lus95, Section 6.10] to specify the set $J \subseteq \tilde{\Delta}$. Note that when \mathbf{G} is of classical type, the group $\mathbf{L}_{c^\omega(J)}(\mathbb{F}_q)$ is also of classical type and so if it admits a unipotent cuspidal representation, then it has *exactly one* unipotent cuspidal representation. Thus for the classical types we will only record the J and $\mathbb{O}^\vee(c^\omega(J), \sigma)$. The explicit parameters can be found in [Lus95], [Ree00], [FO20, §4.7], and [FOS19].

5.1 Classical groups

5.1.1 $PGL(n)$ If $\mathbf{G} = PGL(n)$, then $G^\vee = SL(n, \mathbb{C})$ and $Z(G^\vee) = \mathbb{Z}/n\mathbb{Z}$. Hence $\Omega = \text{Irr}(Z(G^\vee))$ can be identified with C_n . For $\omega \in \Omega$, the inner form \mathbf{G}^ω admits unipotent supercuspidal representations if and only if ω has order n and $J = \emptyset$. In this case $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is the principal nilpotent orbit.

5.1.2 $SO(2n+1)$ If $\mathbf{G} = SO(2n+1)$, $G^\vee = Sp(2n, \mathbb{C})$ and $Z(G^\vee) = \mathbb{Z}/2\mathbb{Z}$. The inner forms are parametrized by $\widehat{Z}(G^\vee) \cong C_2 = \{1, -1\}$.

1. If $\omega = 1$, then J is of the form $D_\ell \times B_t$, where $\ell + t = n$, $\ell = a^2$, $t = b(b+1)$, a, b nonnegative integers, a even. Let

$$\delta = \begin{cases} b - a & \text{if } b \geq a \\ a - b - 1 & \text{if } a > b, \end{cases} \quad (5.1.1)$$

and $\Sigma = a + b$. The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is parameterized by the partition

$$\lambda = (2, 4, \dots, 2\delta) \cup (2, 4, \dots, 2\Sigma). \quad (5.1.2)$$

2. If $\omega = -1$, then J is of the form $D_\ell \times B_t$, where $\ell + t = n$, $\ell = a^2$, $t = b(b+1)$, a, b nonnegative integers, where a is now odd. The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is defined analogously to the $\omega = 1$ case.

5.1.3 $PSp(2n)$ If $\mathbf{G} = PSp(2n)$, then $G^\vee = Spin(2n+1, \mathbb{C})$, and $Z(G^\vee) = \mathbb{Z}/2\mathbb{Z}$. The inner forms are parametrized by $\widehat{Z}(G^\vee) \cong C_2 = \{1, -1\}$.

1. If $\omega = 1$, then J is of the form $C_\ell \times C_t$, where $\ell + t = n$, $\ell = a(a+1)$, $t = b(b+1)$, a, b nonnegative integers and $a \geq b$. Let $\delta = a - b$ and $\Sigma = a + b$. The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is parameterized by the partition

$$\lambda = (1, 3, \dots, 2\delta - 1) \cup (1, 3, \dots, 2\Sigma + 1) \quad (5.1.3)$$

where \cup means union of partitions.

2. If $\omega = -1$, then J is of the form $J = C_\ell {}^2A_t C_\ell$, where $2\ell + t = n - 1$ and $t = \frac{a(a+1)}{2} - 1$, $\ell = b(b+1)$, a, b are nonnegative integers. If $a = 0, 1$, we interpret J as being $J = C_\ell \times C_\ell$. Let a' be such that $a = 2a'$ if a is even and $a = 2a' + 1$ if a is odd. Let $\Sigma = b + a'$ and

$$\delta = \begin{cases} b - a' & \text{if } 2b \geq a \\ a' - b & \text{if } 2b < a. \end{cases}$$

The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is parameterized by the partition

$$\lambda = \begin{cases} (1, 5, \dots, 4\Sigma + 1) \cup (3, 7, \dots, 4\delta - 1) & \text{if } a \text{ is even and } 2b \geq a \\ (1, 5, \dots, 4\Sigma + 1) \cup (1, 5, \dots, 4\delta - 3) & \text{if } a \text{ is even and } 2b < a \\ (3, 7, \dots, 4\Sigma + 3) \cup (1, 5, \dots, 4\delta - 3) & \text{if } a \text{ is odd and } 2b \geq a \\ (3, 7, \dots, 4\Sigma + 3) \cup (3, 7, \dots, 4\delta - 1) & \text{if } a \text{ is odd and } 2b < a. \end{cases}$$

5.1.4 *PSO(2n)* If $\mathbf{G} = PSO(2n)$, then $G^\vee = Spin(2n, \mathbb{C})$, and

$$Z(G^\vee) = \begin{cases} (\mathbb{Z}/2\mathbb{Z})^2, & n \text{ even,} \\ \mathbb{Z}/4\mathbb{Z}, & n \text{ odd.} \end{cases}$$

Let τ be the standard diagram automorphism of type D_n . Let $\{1, -1\}$ be the kernel of the isogeny $Spin(2n, \mathbb{C}) \rightarrow SO(2n, \mathbb{C})$. Write the four characters of $Z(G^\vee)$ as $\Omega = \{1, \eta, \rho, \eta\rho\}$, where $\tau(\eta) = \eta$ and $\eta(-1) = 1$.

1. If $\omega = 1$, then J is of the form $D_\ell \times D_t$, where $\ell + t = n$, $\ell = a^2$, $t = b^2$, a, b even non-negative integers, $a \geq b$. Let $\delta = a - b$, $\Sigma = a + b$. The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is parameterized by the partition

$$\lambda = (1, 3, \dots, 2\delta - 1) \cup (1, 3, \dots, 2\Sigma - 1). \quad (5.1.4)$$

2. If $\omega = \eta$, then J is of the form ${}^2D_\ell \times {}^2D_t$, where $\ell + t = n$, $\ell = a^2$, $t = b^2$, and a, b are odd positive integers, $a \geq b$. The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is defined analogously to the $\omega = 1$ case.

3. If $\omega = \rho, \eta\rho$, then J can take one of the following two forms:

- (i) J is of the form 2A_t , where $t = n - 1$ is even, $t = \frac{a(a+1)}{2} - 1$, a is a non-negative integer. This means that $a \equiv 0, 3 \pmod{4}$. There are four ways to embed J into the affine Dynkin diagram \tilde{D}_n , two of them are ρ -stable, and the other two $\eta\rho$ -stable. In all cases the nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is parameterized by the partition

$$\lambda = \begin{cases} (3, 3, 7, 7, \dots, 2a - 1, 2a - 1), & a \equiv 0 \pmod{4}, \\ (1, 1, 5, 5, \dots, 2a - 1, 2a - 1), & a \equiv 3 \pmod{4}. \end{cases} \quad (5.1.5)$$

- (ii) J is of the form $D_\ell {}^2A_t D_\ell$, where $2\ell + t = n - 1$, $t = \frac{a(a+1)}{2} - 1$ and $\ell = b^2$, a, b are non-negative integers. Let a' be such that $a = 2a'$ if a is even and $a = 2a' + 1$ if a is odd. Let $\Sigma = b + a'$ and

$$\delta = \begin{cases} b - a' & \text{if } 2b > a \\ a' - b & \text{if } 2b \leq a. \end{cases}$$

The nilpotent orbit $\mathbb{O}^\vee(c^\omega(J), \sigma)$ is parameterized by the partition

$$\lambda = \begin{cases} (3, 7, \dots, 4\Sigma - 1) \cup (1, 5, \dots, 4\delta - 3) & \text{if } a \text{ is even and } 2b > a \\ (3, 7, \dots, 4\Sigma - 1) \cup (3, 7, \dots, 4\delta - 1) & \text{if } a \text{ is even and } 2b \leq a \\ (1, 5, \dots, 4\Sigma + 1) \cup (3, 7, \dots, 4\delta - 5) & \text{if } a \text{ is odd and } 2b > a \\ (1, 5, \dots, 4\Sigma + 1) \cup (1, 5, \dots, 4\delta + 1) & \text{if } a \text{ is odd and } 2b \leq a. \end{cases}$$

5.2 Exceptional groups

5.2.1 G_2 If $\mathbf{G} = G_2$, then $G^\vee = G_2(\mathbb{C})$, and $Z(G^\vee) = \{1\}$. If $\omega = 1$ then J is of the form G_2 and there are 4 choices for σ as enumerated in Table 1. In all cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = G_2(a_1).$$

5.2.2 F_4 If $\mathbf{G} = F_4$, then $G^\vee = F_4(\mathbb{C})$, and $Z(G^\vee) = \{1\}$. If $\omega = 1$ then J is of the form F_4 and there are 7 choices for σ as enumerated in Table 1. In all cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = F_4(a_3).$$

5.2.3 E_6 If $\mathbf{G} = E_6$, then $G^\vee = E_6(\mathbb{C})$, and $Z(G^\vee) = \{1, \zeta, \zeta^2\}$.

1. If $\omega = 1$ then J is of the form E_6 and there are 2 choices for σ as enumerated in Table 1. In both cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = D_4(a_1).$$

2. If $\omega \in \{\zeta, \zeta^2\}$ then J is of the form 3D_4 and $\sigma = D_4[1]$ or $D_4[-1]$. In both cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = E_6(a_3).$$

5.2.4 E_7 If $\mathbf{G} = E_7$, then $G^\vee = E_7(\mathbb{C})$, and $Z(G^\vee) = \{1, -1\}$.

1. If $\omega = 1$ then J is of the form E_7 and there are 2 choices for σ as enumerated in Table 1. In both cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = A_4 + A_1.$$

2. If $\omega = -1$, then J is of the form 2E_6 . There are three cuspidal unipotent representations afforded by J : ${}^2E_6[1]$, ${}^2E_6[\theta]$, ${}^2E_6[\theta^2]$. In all cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = E_7(a_5).$$

5.2.5 E_8 If $\mathbf{G} = E_8$, then $G^\vee = E_8(\mathbb{C})$, and $Z(G^\vee) = \{1\}$. If $\omega = 1$ then J is of the form E_8 and there are 13 choices for σ as enumerated in Table 1. In all cases

$$\mathbb{O}^\vee(c^\omega(J), \sigma) = E_8(a_7).$$

6 Proof of Proposition 3.0.1

In this section, we will prove Proposition 3.0.1.

6.1 Classical groups

In each case we show that

$$\overline{\mathbb{L}}(J, \text{WF}(\sigma)) = d_A(\lambda, 1)$$

where λ is the partition parameterizing $\mathbb{O}^\vee(c^\omega(J), \sigma)$. We will use the machinery of [Ach03, Section 3.4] to prove this equality.

6.1.1 $PGL(n)$ Let $\omega \in \Omega \simeq C_n$ be of order n . Let $J = \emptyset$. Then $\sigma = \text{triv}$, $\text{WF}(\sigma) = \{0\}$, and $\mathbb{O}^\vee(c^\omega(\emptyset), \sigma)$ is the principal orbit $\mathbb{O}_{\text{prin}}^\vee$. We need to show that

$$\overline{\mathbb{L}}(\emptyset, \{0\}) = d_A(\mathbb{O}_{\text{prin}}^\vee, 1).$$

But both sides are equal to $(\{0\}, 1)$ and so we have equality.

6.1.2 $SO(2n+1)$ Consider the cases $\omega = 1, -1$ simultaneously. Fix integers a, b as in section 5.1.2 to fix J and hence σ . By section 4

$$\text{WF}(\sigma) = (1, 1, 3, 3, \dots, 2a-1, 2a-1) \times (1, 1, 3, 3, \dots, 2b-1, 2b-1, 2b+1).$$

Let δ, Σ, λ be as in section 5.1.2. We have that

$$\begin{aligned} \lambda^t &= (\delta, \delta, \delta-1, \delta-1, \dots, 1, 1) \vee (\Sigma, \Sigma, \Sigma-1, \Sigma-1, \dots, 1, 1) \\ &= \begin{cases} (2b, 2b, 2b-2, 2b-2, \dots, 2a, 2a, 2a-1, 2a-1, \dots, 1, 1) & \text{if } b \geq a \\ (2a-1, 2a-1, 2a-3, 2a-3, \dots, 2b+1, 2b+1, 2b, 2b, \dots, 1, 1) & \text{if } a > b \end{cases} \end{aligned}$$

so $\pi(\lambda) = \emptyset$. We also have

$$d(\lambda) = (2b+1, 2b-1, 2b-1, \dots, 1, 1) \cup (2a-1, 2a-1, \dots, 1, 1).$$

Since $(1, 1, 3, 3, \dots, 2a-1, 2a-1)$ only has parts with even multiplicity,

$$\overline{\mathbb{L}}(J, \text{WF}(\sigma)) = \langle (1, 1, 3, 3, \dots, 2a-1, 2a-1) \rangle d(\lambda) = \langle \emptyset \rangle d(\lambda) = \langle \pi(\lambda) \rangle d(\lambda) = d_A(\lambda, 1)$$

where $\pi(\lambda)$ is the subpartition of λ^t defined by Achar in ([Ach03], Equation 8).

6.1.3 $PSp(2n)$

1. Let $\omega = 1$. Fix integers a, b as in section 5.1.3 (1) to fix J and hence σ . By section 4

$$\text{WF}(\sigma) = (2, 2, 4, 4, \dots, 2a, 2a) \times (2, 2, 4, 4, \dots, 2b, 2b).$$

Let δ, Σ, λ be as in section 5.1.3 (1). We have that

$$\begin{aligned} \lambda^t &= (\delta, \delta-1, \delta-1, \dots, 1, 1) \vee (\Sigma+1, \Sigma, \Sigma, \dots, 1, 1) \\ &= (2a+1, 2a-1, 2a-1, \dots, 2b+1, 2b+1, 2b, 2b, \dots, 1, 1) \end{aligned}$$

so $\pi(\lambda) = \emptyset$. We also have

$$d(\lambda) = (2a, 2a, \dots, 2b+2, 2b+2, 2b, 2b, 2b, 2b, \dots, 2, 2, 2, 2).$$

Since $(2, 2, 4, 4, \dots, 2a, 2a)$ only has parts with even multiplicity,

$$\overline{\mathbb{L}}(J, \text{WF}(\sigma)) = \langle (2, 2, 4, 4, \dots, 2a, 2a) \rangle d(\lambda) = \langle \emptyset \rangle d(\lambda) = \langle \pi(\lambda) \rangle d(\lambda) = d_A(\lambda, 1).$$

2. Let $\omega = -1$. Fix integers a, b as in section 5.1.3 (2) to fix J and hence σ . Then

$$\text{WF}(\sigma) = (2, 2, 4, 4, \dots, 2b, 2b) \times (1, 2, \dots, a) \times (2, 2, 4, 4, \dots, 2b, 2b).$$

Let δ, Σ, λ be as in section 5.1.3 (2). We have that

$$\lambda^t = \begin{cases} (\Sigma+1, \Sigma^4, \dots, 1^4) \vee (\delta^3, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b \geq a \\ (\Sigma+1, \Sigma^4, \dots, 1^4) \vee (\delta, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b < a \\ ((\Sigma+1)^3, \Sigma^4, \dots, 1^4) \vee (\delta, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b \geq a \\ ((\Sigma+1)^3, \Sigma^4, \dots, 1^4) \vee (\delta^3, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b < a \end{cases} \quad (6.1.1)$$

$$= \begin{cases} (2b+1, (2b)^2, \dots, (a+1)^2, a^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b \geq a \\ (a+1, (a-1)^4, \dots, (2b+1)^4, (2b)^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b < a \\ (2b+1, (2b)^2, \dots, (a+1)^2, a^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b \geq a \\ (a^3, (a-2)^4, \dots, (2b+1)^4, (2b)^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b < a. \end{cases} \quad (6.1.2)$$

Thus $\pi(\lambda) = \emptyset$ since all even parts of λ^t have even multiplicity. Moreover

$$d(\lambda) = (2, 2, 4, 4, \dots, 2b, 2b) \cup (1, 1, 2, 2, \dots, a, a) \cup (2, 2, 4, 4, \dots, 2b, 2b)$$

in all cases. Thus

$$\begin{aligned} \overline{\mathbb{L}}(J, \text{WF}(\sigma)) &= \overline{\mathbb{L}}(\tilde{J}, (2, 2, \dots, 2b, 2b) \times (1, 1, \dots, a, a) \cup (2, 2, \dots, 2b, 2b)) \\ &= \langle (2, 2, 4, 4, \dots, 2b, 2b) \rangle d(\lambda) = \langle \emptyset \rangle d(\lambda) = \langle \pi(\lambda) \rangle d(\lambda) = d_A(\lambda, 1) \end{aligned}$$

where $\tilde{J} = C_l \times C_{t+1+l}$.

6.1.4 $PSO(2n)$

1. Let $\omega \in \{1, \eta\}$. Fix integers a, b as in section 5.1.4 (1), (2) to fix J and hence σ . By section 4

$$\text{WF}(\sigma) = (1, 1, 3, 3, \dots, 2a-1, 2a-1) \times (1, 1, 3, 3, \dots, 2b-1, 2b-1).$$

Let δ, Σ, λ be as in section 5.1.4 (1). We have that

$$\begin{aligned} \lambda^t &= (\delta, \delta-1, \delta-1, \dots, 1, 1) \vee (\Sigma, \Sigma-1, \Sigma-1, \dots, 1, 1) \\ &= (2a, 2a-2, 2a-2, \dots, 2b, 2b, 2b-1, 2b-1, \dots, 1, 1) \end{aligned}$$

so $\pi(\lambda) = \emptyset$ since all odd parts have even multiplicity. We also have

$$d(\lambda) = (2a-1, 2a-1, \dots, 2b+1, 2b+1, 2b-1, 2b-1, 2b-1, 2b-1, \dots, 1, 1, 1, 1).$$

Since $(1, 1, 3, 3, \dots, 2a-1, 2a-1)$ only has parts with even multiplicity,

$$\overline{\mathbb{L}}(J, \text{WF}(\sigma)) = \langle (1, 1, 3, 3, \dots, 2a-1, 2a-1) \rangle d(\lambda) = \langle \emptyset \rangle d(\lambda) = \pi(\lambda) d(\lambda) = d_A(\lambda, 1).$$

2. Let $\omega \in \{\rho, \eta\rho\}$. We will treat the cases (i) and (ii) simultaneously. Fix integers a, b as in section 5.1.4 (3) (ii) to fix J and hence σ (we treat (i) as the case with $b = 0$). By section 4

$$\text{WF}(\sigma) = (1, 1, 3, 3, \dots, 2b-1, 2b-1) \times (1, 2, \dots, a) \times (1, 1, 3, 3, \dots, 2b-1, 2b-1).$$

Let δ, Σ, λ be as in section 5.1.4 (3) (ii). We have that

$$\lambda^t = \begin{cases} (\Sigma^3, (\Sigma-1)^4, \dots, 1^4) \vee (\delta, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b \leq a \\ (\Sigma^3, (\Sigma-1)^4, \dots, 1^4) \vee (\delta^3, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b < a \\ (\Sigma+1, \Sigma^4, \dots, 1^4) \vee (\delta, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b < a \\ ((\Sigma+1)^3, \Sigma^4, \dots, 1^4) \vee (\delta^3, (\delta-1)^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b < a \end{cases} \quad (6.1.3)$$

$$= \begin{cases} (2b+1, (2b)^2, \dots, (a+1)^2, a^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b \geq a \\ (a+1, (a-1)^4, \dots, (2b+1)^4, (2b)^4, \dots, 1^4) & \text{if } a \text{ is even and } 2b < a \\ (2b+1, (2b)^2, \dots, (a+1)^2, a^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b \geq a \\ (a^3, (a-2)^4, \dots, (2b+1)^4, (2b)^4, \dots, 1^4) & \text{if } a \text{ is odd and } 2b < a. \end{cases} \quad (6.1.4)$$

Thus $\pi(\lambda) = \emptyset$ since all even parts of λ^t have even multiplicity. Moreover

$$d(\lambda) = (2, 2, 4, 4, \dots, 2b, 2b) \cup (1, 1, 2, 2, \dots, a, a) \cup (2, 2, 4, 4, \dots, 2b, 2b)$$

in all cases. Thus

$$\begin{aligned} \overline{\mathbb{L}}(J, \text{WF}(\sigma)) &= \overline{\mathbb{L}}(\tilde{J}, (1, 1, \dots, 2b-1, 2b-1) \times (1, 1, \dots, a, a) \cup (1, 1, \dots, 2b-1, 2b-1)) \\ &= \langle (2, 2, 4, 4, \dots, 2b, 2b) \rangle d(\lambda) = \langle \emptyset \rangle d(\lambda) = \langle \pi(\lambda) \rangle d(\lambda) = d_A(\lambda, 1) \end{aligned}$$

where $\tilde{J} = D_l \times D_{t+1+l}$.

6.2 Exceptional groups

6.2.1 Split forms Suppose that \mathbf{G} is split, of exceptional type, and that $\omega = 1$. As can be seen in Section 5.2, J is always equal to Δ . Thus,

$$\overline{\mathbb{L}}(J, \text{WF}(\sigma)) = (\text{WF}(\sigma), 1).$$

On the other hand, the nilpotent orbit $\mathbb{O}^\vee := \mathbb{O}^\vee(c^\omega(J), \sigma)$ is always special. Thus,

$$d_A(\mathbb{O}^\vee, 1) = (d(\mathbb{O}^\vee), 1)$$

by the general properties of d_A , see [Ach03, Section 3]. So for Proposition 3.0.1 it suffices to show that

$$\text{WF}(\sigma) = d(\mathbb{O}^\vee)$$

for all σ . This follows by comparing the orbits in Table 1 and in section 5.2.

6.2.2 Non-split forms of E_6 Suppose \mathbf{G} is of type E_6 and $\omega \in \{\zeta, \zeta^2\}$. Then J is of the form 3D_4 , and $\text{WF}(\sigma) = (1, 1, 3, 3)$ for both $\sigma = D_4[1]$ and $\sigma = D_4[-1]$. The orbit $(1, 1, 3, 3)$ is the orbit A_2 in Bala-Carter notation. Thus we need to show that

$$\overline{\mathbb{L}}(J, (1, 1, 3, 3)) = d_A(E_6(a_3), 1).$$

We note that $E_6(a_3)$ is special and $d(E_6(a_3)) = A_2$ so we must show that

$$\overline{\mathbb{L}}(J, A_2) = (A_2, 1).$$

Since $J \subseteq \Delta$ this follows from [Oka21, Proposition 2.30].

6.2.3 Non-split forms of E_7 Suppose \mathbf{G} is of type E_7 and $\omega = -1$. Then J is of the form 2E_6 , and $\text{WF}(\sigma) = D_4(a_1)$ for all possible σ . Thus we need to show that

$$\overline{\mathbb{L}}(J, D_4(a_1)) = d_A(E_7(a_5), 1).$$

We note that $E_7(a_5)$ is special and $d(E_7(a_5)) = D_4(a_1)$ so we must show that

$$\overline{\mathbb{L}}(J, D_4(a_1)) = (D_4(a_1), 1).$$

Since $J \subseteq \Delta$ this follows from [Oka21, Proposition 2.30].

References

- [Ach03] P. Achar. “An order-reversing duality map for conjugacy classes in Lusztig’s canonical quotient”. In: *Transform. Groups* 8 (2003), pp. 107–145.
- [AGS22] A. Aizenbud, D. Gourevitch, and E. Sayag. *Irreducibility of wave front sets for depth zero cuspidal representations*. 2022.
- [Aub95] A. M. Aubert. “Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d’un groupe réductif p-adique”. In: *Trans. Amer. Math. Soc.* 347.6 (1995), pp. 2179–2189.
- [ABPS17] A.-M. Aubert, P. Baum, R. Plymen, and M. Solleveld. “Conjectures about p -adic groups and their non commutative geometry”. In: *Around Langlands correspondences*. Vol. 691. Contemp. Math. Amer. Math. Soc., Providence, RI, 2017, pp. 5–21.
- [AMS18] A.-M. Aubert, A. Moussaoui, and M. Solleveld. “Generalizations of the Springer correspondence and cuspidal Langlands parameters”. In: *Manuscripta Math.* 157 (1-2 2018), pp. 121–192.
- [BV85] D. Barbasch and D. Vogan. “Unipotent representations of complex semisimple groups”. In: *Ann. of Math. (2)* 121.1 (1985), pp. 41–110.
- [BM97] Dan Barbasch and Allen Moy. “Local character expansions”. In: *Ann. Sci. École Norm. Sup. (4)* 30.5 (1997), pp. 553–567.
- [Car93] R. Carter. *Finite groups of Lie type*. Wiley Classics Library. John Wiley & Sons, Ltd., 1993.
- [CMO21] D. Ciubotaru, L. Mason-Brown, and E. Okada. *Some Unipotent Arthur Packets for Reductive p -adic Groups I*. 2021.
- [FO20] Y. Feng and E. Opdam. “On a uniqueness property of supercuspidal unipotent representations”. In: *Adv. Math.* 375 (2020), 107406, 62 pp.
- [FOS19] Y. Feng, E. Opdam, and M. Solleveld. *On formal degrees of unipotent representations*. 2019.
- [Har99] Harish-Chandra. “Harish-Chandra Admissible invariant distributions on reductive p -adic groups. With a preface and notes by Stephen DeBacker and Paul J. Sally, Jr.” In: vol. 16. University Lecture Series. American Mathematical Society, Providence, RI, 1999, xiv+97 pp.
- [How74] R. Howe. “The Fourier Transform and Germs of Characters (Case of $Gl(n)$ over a p -Adic Field).” In: *Mathematische Annalen* 208 (1974), pp. 305–322.

- [Kal16] T. Kaletha. “The local Langlands conjecture for non-quasi-split groups”. In: *Families of automorphic forms and the trace formula*. Simons Symp. Springer, 2016, pp. 217–257.
- [Kaw87] Noriaki Kawanaka. “Shintani lifting and Gel’fand-Graev representations”. In: *The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986)*. Vol. 47. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 1987, pp. 147–163.
- [Kot84] R. Kottwitz. “Stable trace formula: cuspidal tempered terms”. In: *Duke Math. J.* 51 (3 1984), pp. 611–650.
- [Lus95] G. Lusztig. “Classification of unipotent representations of simple p -adic groups”. In: *IMRN* 11 (1995), pp. 517–589.
- [Lus84] George Lusztig. *Characters of Reductive Groups over a Finite Field. (AM-107)*. Princeton University Press, 1984.
- [Lus92] George Lusztig. “A unipotent support for irreducible representations”. In: *Adv. Math.* 94.2 (1992), pp. 139–179.
- [MW87] C. Mœglin and J.-L. Waldspurger. “Modèles de Whittaker dégénérés pour des groupes p -adiques”. In: *Math. Z.* 196.3 (1987), pp. 427–452. ISSN: 0025-5874. DOI: 10.1007/BF01200363. URL: <https://ezproxy-prd.bodleian.ox.ac.uk:2102/10.1007/BF01200363>.
- [Mor96] L. Morris. “Tamely ramified supercuspidal representations”. In: *Ann. Sci, École Norm. Sup.* 29 (5 1996), pp. 639–667.
- [MP96] Allen Moy and Gopal Prasad. “Jacquet functors and unrefined minimal K -types”. In: *Comment. Math. Helv.* 71.1 (1996), pp. 98–121. ISSN: 0010-2571. DOI: 10.1007/BF02566411. URL: <https://ezproxy-prd.bodleian.ox.ac.uk:2102/10.1007/BF02566411>.
- [Oka21] Emile Takahiro Okada. *The wavefront set over a maximal unramified field extension*. 2021. arXiv: 2107.10591 [math.RT].
- [Oka22] Emile Takahiro Okada. *Wavefront sets and isogenies [In preparation]*. 2022.
- [Opd16] E. Opdam. “Spectral transfer morphisms for unipotent affine Hecke algebras”. In: *Selecta Math. (N.S.)* 22 (4 2016), pp. 2143–2207.
- [Pom77] K. Pommerening. “Über die unipotenten Klassen reductiver Gruppen”. In: *J. Algebra* (1977), pp. 373–398.
- [Pom80] K. Pommerening. “Über die unipotenten Klassen reductiver Gruppen II”. In: *J. Algebra* 65 (1980), pp. 525–536.
- [Ran72] R. Ranga Rao. “Orbital integrals in reductive groups”. In: *Ann. of Math. (2)* 96 (1972), pp. 505–510. ISSN: 0003-486X. DOI: 10.2307/1970822. URL: <https://ezproxy-prd.bodleian.ox.ac.uk:2102/10.2307/1970822>.
- [Ree00] M. Reeder. “Formal degrees and L -packets of unipotent discrete series representations of exceptional p -adic groups. With an appendix by Frank Lübeck”. In: *J. Reine Angew. Math.* 520 (2000), pp. 37–93.
- [Sol18] M. Solleveld. *A local Langlands correspondence for unipotent representations*. 2018. eprint: 1806.11357 (math.RT).
- [Som01] E. Sommers. “Lusztig’s canonical quotient and generalized duality”. In: *J. Algebra* 243 (2001), pp. 790–812.
- [Spa82] N. Spaltenstein. *Classes Unipotentes et Sous-groupes de Borel*. Springer Berlin Heidelberg, 1982.
- [Tit79] J. Tits. “Reductive groups over local fields”. In: *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*. Proc. Sympos. Pure Math., XXXIII. Amer. Math. Soc., Providence, R.I., 1979, pp. 29–69.
- [Vog93] D.A. Vogan. “The local Langlands conjecture”. In: *Representation theory of groups and algebras*. Vol. 145. Contemp. Math. Amer. Math. Soc., Providence, RI, 1993, pp. 305–379.
- [Wal18] J.-L. Waldspurger. “Représentations de réduction unipotente pour $SO(2n + 1)$, III: exemples de fronts d’onde”. In: *Algebra Number Theory* 12.5 (2018), pp. 1107–1171.