

# Orthonormal Expansions for Translation-Invariant Kernels

Filip Tronarp<sup>1</sup> — Toni Karvonen<sup>2</sup>

<sup>1</sup>Department of Computer Science, University of Tübingen, Germany

<sup>2</sup>Department of Mathematics and Statistics, University of Helsinki, Finland

June 20, 2022

## Abstract

We present a general Fourier analytic technique for constructing orthonormal basis expansions of translation-invariant kernels from orthonormal bases of  $\mathcal{L}_2(\mathbb{R})$ . This allows us to derive explicit expansions on the real line for (i) Matérn kernels of all half-integer orders in terms of associated Laguerre functions, (ii) the Cauchy kernel in terms of rational functions, and (iii) the Gaussian kernel in terms of Hermite functions.

**Keywords:** positive-definite kernels, radial basis functions, orthonormal expansions, orthogonal polynomials

**MSC2020:** 65D12, 46E22, 33C45, 60G10

## 1 Introduction

Let  $r: \Omega \times \Omega \rightarrow \mathbb{R}$  be a symmetric positive-semidefinite kernel on some domain  $\Omega$  and  $\mathcal{H}_r(\Omega)$  its reproducing kernel Hilbert space (RKHS), which is equipped with an inner product  $\langle \cdot, \cdot \rangle_r$  and the associated norm  $\|\cdot\|_r$ . Practically every commonly used kernel induces an infinite-dimensional RKHS that is separable (see [17] for a short review on separability of RKHSs), which means that  $\mathcal{H}_r(\Omega)$  has an orthonormal basis  $\{\psi_m\}_{m \in I}$  for some countably infinite index set  $I$  (e.g.,  $I = \mathbb{N}$ ) in terms of which the kernel has the pointwise convergent orthonormal expansion

$$r(t, u) = \sum_{m \in I} \psi_m^*(t) \psi_m(u) \quad \text{for all } t, u \in \Omega. \quad (1.1)$$

If  $\Omega$  is a compact subset of  $\mathbb{R}^d$  and  $r$  is continuous, the expansion converges uniformly [18, Section 11.3]. One can approach kernels and their orthonormal expansions in two different ways. The *space-centric* approach is to start with a Hilbert space or its orthonormal basis, show that this space is an RKHS, and construct its reproducing kernel via (1.1); under fortuitous circumstances the kernel is available in closed form. A prime example of this approach is how Korobov spaces and their kernels, which can be expressed in terms of Bernoulli polynomials, are used in the quasi-Monte Carlo literature [e.g., 5, Section 5.8]. Other examples include Hardy spaces [18, Section 1.4.2], power series kernels [29], and Hermite spaces [12]. In the *kernel-centric* approach one instead starts with a kernel that has, in some sense, desirable or intuitive properties and, if necessary for whatever purpose, finds its RKHS or orthonormal expansion. This approach is dominant in scattered data approximation [25] and Gaussian process modelling as used in, for example, spatial statistics [21] and machine learning [19]. Given only the algebraic form of the kernel, it is usually a non-trivial task to find an orthonormal expansion. However, access to an expansion of the form in (1.1) is often needed to develop reduced rank methods of sub-cubic

computational complexity [20], to improve numerical stability [9], or to illustrate and validate theoretical results [13, Section 5].

Translation-invariant kernels, which take the form  $r(t, u) = \Phi(t - u)$  for some function  $\Phi: \Omega \rightarrow \mathbb{R}$ , are a mainstay of scattered data approximation and Gaussian process modelling. Nevertheless, few orthonormal expansions appear to have been constructed for these kernels. To the best of our knowledge, the Matérn- $\frac{1}{2}$  kernel  $r(t, u) = \exp(-\lambda |t - u|)$  and the Gaussian kernel  $r(t, u) = \exp(-\frac{1}{2}\lambda^2(t - u)^2)$  on subsets of  $\mathbb{R}$  are the only commonly used translation-invariant kernels for which orthonormal expansions have been found. For various expansions of the Matérn- $\frac{1}{2}$  kernel, see Section 4 in [10], Section 3.4.1 in [24], Example 4.1 in [27], and Example 2.5 and Appendix A.2 in [8]. For the Gaussian kernel both a simple non-Mercer expansion based on a Taylor expansion of the exponential function [e.g., 16] and a class of Mercer expansions [8, Section 12.2.1], which appear to have originated in [28, Section 4], are available. A large collection of expansions for kernels which are not translation-invariant can be found in [8, Appendix A]. In this article we describe a general and conceptually simple technique for constructing orthonormal bases for translation-invariant kernels on  $\mathbb{R}$  out of orthonormal bases of  $\mathcal{L}_2(\mathbb{R})$ .

## 1.1 Construction of orthonormal bases

Let  $z^*$  denote the complex conjugate and  $|z|$  the modulus of  $z \in \mathbb{C}$ . The spaces  $\mathcal{L}_2(\mathbb{R})$  and  $\mathcal{L}_2(\mathbb{R}, 1/2\pi)$  consist of all square-integrable functions  $f: \mathbb{R} \rightarrow \mathbb{C}$  and are equipped with the inner products

$$\langle f, g \rangle_{\mathcal{L}_2(\mathbb{R})} = \int_{-\infty}^{\infty} f^*(t)g(t) dt \quad \text{and} \quad \langle f, g \rangle_{\mathcal{L}_2(\mathbb{R}, 1/2\pi)} = \frac{1}{2\pi} \int_{-\infty}^{\infty} f^*(t)g(t) dt.$$

The Fourier transform and the corresponding inverse transform for any integrable or square-integrable function  $f$  are defined as

$$\hat{f}(\omega) = \int_{-\infty}^{\infty} f(t)e^{-i\omega t} dt \quad \text{and} \quad f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}(\omega)e^{i\omega t} d\omega.$$

The Fourier transform defines an isometry from  $\mathcal{L}_2(\mathbb{R})$  to  $\mathcal{L}_2(\mathbb{R}, 1/2\pi)$  via the Plancherel theorem

$$\int_{-\infty}^{\infty} f^*(t)g(t) dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}^*(\omega)\hat{g}(\omega) d\omega.$$

The functions  $f$  and  $\hat{f}$  are referred to as time domain and Fourier domain representations, respectively. Our  $\mathcal{H}_r(\mathbb{R})$ -orthonormal expansions are derived from the following rather straightforward theorem. Let  $I$  be a countably infinite index set, typically either  $\mathbb{N}$  or  $\mathbb{Z}$ .

**Theorem 1.1** (Construction of orthonormal bases). *Let  $r(t, u) = \Phi(t - u)$  be a translation-invariant symmetric positive-definite kernel defined by  $\Phi \in C(\mathbb{R}) \cap \mathcal{L}_1(\mathbb{R})$  and  $\{\varphi_m\}_{m \in I}$  an orthonormal basis of  $\mathcal{L}_2(\mathbb{R})$ . Let  $h$  be a function such that  $|\hat{h}(\omega)| = \hat{\Phi}(\omega)^{1/2}$ . Then the functions*

$$\psi_m(t) = \int_{-\infty}^{\infty} h(t - \tau)\varphi_m(\tau) d\tau \quad \text{with Fourier transforms} \quad \hat{\psi}_m(\omega) = \hat{h}(\omega)\hat{\varphi}_m(\omega)$$

for  $m \in I$  form an orthonormal basis of  $\mathcal{H}_r(\mathbb{R})$  and the kernel  $r$  has the pointwise convergent expansion

$$r(t, u) = \sum_{m \in I} \psi_m^*(t)\psi_m(u) \quad \text{for all} \quad t, u \in \mathbb{R}.$$

*Proof.* That  $r$  is symmetric positive-definite implies that  $\hat{\Phi}$  is real-valued and positive [25, Theorem 6.11]. For a non-vanishing function  $h$  such that  $|\hat{h}(\omega)| = \hat{\Phi}(\omega)^{1/2}$  we define a convolution operator  $\mathcal{H}: \mathcal{L}_2(\mathbb{R}) \rightarrow \mathcal{L}_2(\mathbb{R})$  via

$$(\mathcal{H}f)(t) = \int_{-\infty}^{\infty} h(t - \tau)f(\tau) d\tau \quad \text{for all } t \in \mathbb{R}.$$

Note that the convolution theorem yields  $\widehat{\mathcal{H}f}(\omega) = \hat{h}(\omega)\hat{f}(\omega)$ . By the standard characterisation (see [14] or [25, Theorem 10.12]) of the RKHS of a translation-invariant kernel,

$$\langle f, g \rangle_r = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\hat{f}^*(\omega)\hat{g}(\omega)}{\hat{\Phi}(\omega)} d\omega \quad \text{for any } f, g \in \mathcal{H}_r(\mathbb{R}). \quad (1.2)$$

For any  $f, g \in \mathcal{L}_2(\mathbb{R})$  the convolution theorem and Plancherel theorem thus give

$$\langle \mathcal{H}f, \mathcal{H}g \rangle_r = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{|\hat{h}(\omega)|^2 \hat{f}^*(\omega)\hat{g}(\omega)}{\hat{\Phi}(\omega)} d\omega = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{f}^*(\omega)\hat{g}(\omega) d\omega = \langle f, g \rangle_{\mathcal{L}_2(\mathbb{R})},$$

which shows that  $\mathcal{H}$  is an isometry from  $\mathcal{L}_2(\mathbb{R})$  to  $\mathcal{H}_r(\mathbb{R})$ . It follows from (1.2) that the inverse Fourier transform

$$(\mathcal{H}^{-1}f)(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\hat{f}(\omega)}{\hat{h}(\omega)} e^{i\omega t} d\omega \quad \text{for all } t \in \mathbb{R}$$

defines the inverse of  $\mathcal{H}$ . Therefore  $\mathcal{H}$  is an isometric isomorphism and thus maps every orthonormal basis of  $\mathcal{L}_2(\mathbb{R})$  to an orthonormal basis of  $\mathcal{H}_r(\mathbb{R})$  [11, Section 2.6].  $\square$

To obtain the basis functions  $\psi_m$  in time domain using Theorem 1.1 one has to either compute the convolution  $\int_{-\infty}^{\infty} h(t - \tau)\phi_m(\tau) d\tau$  or the inverse Fourier transform of  $\hat{h}(\omega)\hat{\phi}_m(\omega)$ . It is therefore necessary to select a basis of  $\mathcal{L}_2(\mathbb{R})$  for which either of these operations can be done in closed form. We use Theorem 1.1 to derive orthonormal expansions for (i) Matérn kernels for all half-integer orders, (ii) the Cauchy kernel (i.e., rational quadratic kernel [19, Equation (4.19)] with  $\alpha = 1$ ), and (iii) the Gaussian kernel. The expansions are summarised in Section 2.

## 1.2 On Mercer expansions

Let  $\Omega$  be a subset of  $\mathbb{R}^d$  and  $w: \Omega \rightarrow [0, \infty)$  a weight function. The Hilbert space  $\mathcal{L}_2(\Omega, w)$  is equipped with the inner product

$$\langle f, g \rangle_{\mathcal{L}_2(\Omega, w)} = \int_{\Omega} f^*(t)g(t)w(t) dt$$

and consists of all functions  $f: \mathbb{R} \rightarrow \mathbb{C}$  for which the corresponding norm is finite. Suppose that the kernel  $r$  is continuous and define the integral operator

$$\mathcal{T}_{r,w}f = \int_{-\infty}^{\infty} r(\cdot, u)f(u)w(u) du. \quad (1.3)$$

Under certain assumptions, Mercer's theorem [22] states that (i)  $\mathcal{T}_{r,w}$  has continuous eigenfunctions  $\{\phi_m\}_{m=0}^{\infty}$  and corresponding positive non-increasing eigenvalues  $\{\mu_m\}_{m=0}^{\infty}$  which tend to zero, (ii)  $\{\phi_m\}_{m=0}^{\infty}$  are an orthonormal basis of  $\mathcal{L}_2(\Omega, w)$ , and (iii)  $\{\sqrt{\mu_m}\phi_m\}_{m=0}^{\infty}$  is an orthonormal basis of  $\mathcal{H}_r(\Omega)$ . Consequently, the kernel has the pointwise convergent *Mercer expansion*

$$r(t, u) = \sum_{m=0}^{\infty} \mu_m \phi_m^*(t)\phi_m(u) \quad \text{for all } t, u \in \Omega. \quad (1.4)$$

While Mercer’s theorem and the eigenvalues of  $\mathcal{T}_{r,w}$  constitute a powerful tool for understanding, for example, optimal approximation in  $\mathcal{L}_2(\Omega, w)$ -norm (e.g., [7, Section 2.4]) and improved approximation orders in subsets of  $\mathcal{H}_r(\Omega)$  [25, Section 11.5], there is often (both in theoretical research and practical applications) no reason to prefer a Mercer expansion (1.4) over a generic RKHS-orthonormal expansion (1.1). For example, the Karhunen–Loève theorem is merely a special case of a more general result that a Gaussian process with covariance kernel  $r$  can be expanded in terms of any orthonormal basis of  $\mathcal{H}_r(\Omega)$  [1, Chapter III]. It therefore appears unjustified that Mercer expansions should be generally preferred over generic orthonormal expansions.

Constructing a Mercer expansion by first identifying a convenient weight and then finding the eigendecomposition of the integral operator (1.3) can be rather involved, which is illustrated by the construction in [8, Example 2.5] for the Matérn- $\frac{1}{2}$  kernel. What makes Theorem 1.1 convenient is therefore that it does *not* require that the expansion be Mercer for some weight. However, identifying a weight  $w$  for which the basis function  $\psi_m$  constructed via Theorem 1.1 are  $\mathcal{L}_2(\mathbb{R}, w)$ -orthogonal shows that the expansion is Mercer because the  $\mathcal{L}_2(\mathbb{R}, w)$ -normalised versions of  $\psi_m$  are the eigenfunctions of  $\mathcal{T}_{r,w}$ . It turns out that our expansion for the Gaussian kernel is Mercer and the ones for Matérn kernels are “almost” Mercer, in that all but finitely many basis functions are orthogonal in  $\mathcal{L}_2(\mathbb{R}, w)$  for a certain weight.

## 2 Summary of expansions

This section summarises the expansions that we derive using Theorem 1.1. Each expansion converges pointwise for all  $t, u \in \mathbb{R}$ . The parameter  $\lambda$  is a positive scale parameter.

### 2.1 Matérn kernels

Expansions for Matérn kernels are derived in Section 3. A Matérn kernel of order  $\alpha > 0$  is

$$r_{\alpha,\lambda}(t, u) = \frac{2^{1-\alpha}}{\Gamma(\alpha)} (\lambda|t-u|)^\alpha K_\alpha(\lambda|t-u|), \quad (2.1)$$

where  $\Gamma$  is the Gamma function and  $K_\alpha$  the modified Bessel function of the second kind of order  $\alpha$ . Let  $L_m^{(\eta)}$  denote the  $m$ th associated Laguerre polynomial of index  $\eta$ , defined in (3.13), and let  $\{\varphi_{m,\lambda}\}_{m \in \mathbb{Z}}$  be the Laguerre functions

$$\varphi_{m,\lambda}(t) = L_m(2\lambda t)e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t) \quad \text{and} \quad \varphi_{-m-1,\lambda}(t) = -L_m(-2\lambda t)e^{\lambda t} \mathbf{1}_{(-\infty,0)}$$

for  $m \in \mathbb{N}_0$ , where  $L_m = L_m^{(0)}$  and  $\mathbf{1}_A$  denotes the indicator function of a set  $A$ . The Matérn-Laguerre functions

$$\begin{aligned} \psi_{m,\lambda}^{+,\nu}(t) &= \frac{\nu!}{(2\nu)!} \frac{m!}{(m+\nu+1)!} (2\lambda t)^{\nu+1} L_m^{(\nu+1)}(2\lambda t) e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t) & \text{for } m \in \mathbb{N}_0 \\ \psi_{m,\lambda}^{-,\nu}(t) &= (-1)^\nu \psi_{m,\lambda}^{+,\nu}(-t) & \text{for } m \in \mathbb{N}_0, \\ \psi_{m,\lambda}^{0,\nu}(t) &= \frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \varphi_{m+k-\nu-1,\lambda}(t) & \text{for } m = 0, \dots, \nu \end{aligned}$$

form an orthonormal basis of the RKHS when the order is  $\alpha = \nu + 1/2$  for  $\nu \in \mathbb{N}_0$ . Therefore

$$r_{\nu+1/2,\lambda}(t, u) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,\nu}(t) \psi_{m,\lambda}^{0,\nu}(u) + \sum_{m=0}^{\infty} \psi_{m,\lambda}^{-,\nu}(t) \psi_{m,\lambda}^{-,\nu}(u) + \sum_{m=0}^{\infty} \psi_{m,\lambda}^{+,\nu}(t) \psi_{m,\lambda}^{+,\nu}(u)$$

for all  $t, u \in \mathbb{R}$ . The basis functions  $\psi_{m,\lambda}^{-,\nu}$  and  $\psi_{m,\lambda}^{+,\nu}$  are orthogonal in  $\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})$  for the weight function  $w_{\nu,\lambda}(t) = 2\lambda/|2\lambda t|^{\nu+1}$ .

## 2.2 Cauchy kernel

Expansions for the Cauchy kernel are derived in Section 4. The Cauchy kernel is

$$r_\lambda(t, u) = \frac{1}{1 + \lambda^2(t - u)^2}.$$

The complex-valued Cauchy–Laguerre functions

$$\psi_{m,1/\lambda}(t) = -\frac{1}{\sqrt{2}} \frac{(i\lambda t)^m}{(i\lambda t - 1)^{m+1}} \quad \text{and} \quad \psi_{-m-1,1/\lambda}(t) = -\frac{1}{\sqrt{2}} \frac{(i\lambda t)^m}{(i\lambda t + 1)^{m+1}}$$

for  $m \in \mathbb{N}_0$  and the real-valued trigonometric Cauchy–Laguerre functions

$$\alpha_{m,1/\lambda}(t) = \frac{1}{\sqrt{2}} (\psi_{m,1/\lambda}(t) + \psi_{m,1/\lambda}^*(t)) \quad \text{and} \quad \beta_{m,1/\lambda}(t) = \frac{1}{\sqrt{2}} (\psi_{m,1/\lambda}(t) - \psi_{m,1/\lambda}^*(t))$$

for  $m \in \mathbb{N}_0$  form orthonormal bases of the RKHS. Therefore, the Cauchy kernel has the expansions

$$r_\lambda(t, u) = \sum_{m=-\infty}^{\infty} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(u) = \sum_{m=0}^{\infty} \alpha_{m,1/\lambda}(t) \alpha_{m,1/\lambda}(u) + \sum_{m=0}^{\infty} \beta_{m,1/\lambda}(t) \beta_{m,1/\lambda}(u)$$

for all  $t, u \in \mathbb{R}$ . Expressions of  $\alpha_{m,1/\lambda}$  and  $\beta_{m,1/\lambda}$  in terms of real parameters are given in (4.5).

## 2.3 Gaussian kernel

Expansions for the Gaussian kernel are derived in Section 5. The Gaussian kernel is

$$r_\lambda(t, u) = \exp\left(-\frac{\lambda^2}{2}(t - u)^2\right).$$

The functions

$$\psi_{m,\lambda}(t) = \left(\frac{2\sqrt{2}}{3}\right)^{1/2} \sqrt{\frac{1}{6^m m!}} e^{-\lambda^2 t^2/3} \mathbf{H}_m\left(\frac{2\lambda t}{\sqrt{3}}\right) \quad \text{for } m \in \mathbb{N}_0 \quad (2.2)$$

form an orthonormal basis of the RKHS and the kernel has the expansion

$$r_\lambda(t, u) = \sum_{m=0}^{\infty} \psi_{m,\lambda}(t) \psi_{m,\lambda}(u)$$

for all  $t, u \in \mathbb{R}$ . This expansion is a special case of the well-known Mercer expansion of the Gaussian kernel [8, Section 12.2.1]. The basis functions (2.2) are orthogonal in  $\mathcal{L}_2(\mathbb{R}, w_\alpha)$  for the weight function  $w_\alpha(t) = \alpha \pi^{-1/2} e^{-\alpha^2 t^2}$  with  $\alpha = \sqrt{2/3} \lambda$ .

## 3 Matérn kernels

The Matérn kernel of order  $\alpha > 0$  in (2.1) can be written as

$$r_{\alpha,\lambda}(t, u) = \frac{2^{1-2\alpha}}{\Gamma(\alpha)} (2\lambda|t - u|)^\alpha \mathbf{K}_\alpha(\lambda|t - u|),$$

and its Fourier transform is

$$\hat{\Phi}_{\alpha,\lambda}(\omega) = 2^{1-2\alpha} \sqrt{\pi} \frac{\Gamma(\alpha + 1/2)}{\Gamma(\alpha)} \frac{(2\lambda)^{2\alpha}}{(\omega^2 + \lambda^2)^{\alpha+1/2}}.$$

From now on we assume that the kernel is of half-integer order:  $\alpha = \nu + 1/2$  for  $\nu \in \mathbb{N}_0$ . Then the Fourier transform simplifies to

$$\hat{\Phi}_{\nu+1/2,\lambda}(\omega) = \frac{(\nu!)^2}{(2\nu)!} \frac{(2\lambda)^{2\nu+1}}{(\omega^2 + \lambda^2)^{\nu+1}},$$

and a non-symmetric square-root, in the sense that  $|\hat{h}_{\nu+1/2,\lambda}(\omega)|^2 = \hat{\Phi}_{\nu+1/2,\lambda}(\omega)$ , is given by

$$\hat{h}_{\nu+1/2,\lambda}(\omega) = \frac{\nu!}{\sqrt{(2\nu)!}} \frac{(2\lambda)^{\nu+1/2}}{(i\omega + \lambda)^{\nu+1}}. \quad (3.1)$$

The corresponding time domain function is [26, Section 1.03]

$$h_{\nu+1/2,\lambda}(t) = (2\lambda)^{\nu+1/2} \frac{\nu!}{\sqrt{(2\nu)!}} \frac{t^\nu}{\nu!} e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t). \quad (3.2)$$

This function vanishes on the negative real line.

### 3.1 Laguerre functions

The following material is mostly based on Section 2.6.4 in [11] and Section 1.03 in [26]. To derive an orthonormal expansion for the Matérn kernel we use the so-called *Laguerre functions*  $\varphi_{m,\lambda}$  whose Fourier transforms are given by

$$\hat{\varphi}_{m,\lambda}(\omega) = \sqrt{2\lambda} \frac{(i\omega - \lambda)^m}{(i\omega + \lambda)^{m+1}} \quad \text{for } m \in \mathbb{Z}. \quad (3.3)$$

The functions  $\hat{\varphi}_{m,\lambda}$  form an orthonormal basis of  $\mathcal{L}_2(\mathbb{R}, 1/2\pi)$ . Because the Fourier transform is an isometry, the Laguerre functions themselves, defined by the inverse Fourier transform

$$\varphi_{m,\lambda}(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{\varphi}_{m,\lambda}(\omega) e^{i\omega t} d\omega \quad \text{for } m \in \mathbb{Z}, \quad (3.4)$$

are an orthonormal basis of  $\mathcal{L}_2(\mathbb{R})$ . Let  $L_m$  for  $m \in \mathbb{N}_0$  be the  $m$ th Laguerre polynomial

$$L_m(t) = \sum_{k=0}^m \binom{m}{k} \frac{(-1)^k}{k!} t^k. \quad (3.5)$$

For non-negative indices  $m \in \mathbb{N}_0$  the inverse Fourier transform (3.4) is given by

$$\varphi_{m,\lambda}(t) = \sqrt{2\lambda} L_m(2\lambda t) e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t). \quad (3.6)$$

The conjugate symmetry  $\hat{\varphi}_{-m-1,\lambda}^*(\omega) = -\hat{\varphi}_{m,\lambda}(\omega)$  gives the following expression for negative indices:

$$\varphi_{-m-1,\lambda}(t) = -\varphi_m(-t) = -\sqrt{2\lambda} L_m(-2\lambda t) e^{\lambda t} \mathbf{1}_{(-\infty,0)}(t) \quad \text{for } m \in \mathbb{N}_0.$$

The Laguerre functions and their Fourier transforms satisfy the scaling laws

$$\varphi_{m,\lambda}(t) = \sqrt{2\lambda} \varphi_{m,1/2}(2\lambda t) \quad \text{and} \quad \hat{\varphi}_{m,\lambda}(\omega) = \frac{1}{\sqrt{2\lambda}} \hat{\varphi}_{m,1/2}(\omega/2\lambda).$$

Furthermore, they satisfy the following useful identities:

$$\begin{aligned}
\hat{\varphi}_{-m-1,\lambda}^*(\omega) &= -\hat{\varphi}_{m,\lambda}(\omega), && \text{(conjugate symmetry)} \\
\hat{\varphi}_{m+k,\lambda}(\omega) &= \left(\frac{i\omega - \lambda}{i\omega + \lambda}\right)^k \hat{\varphi}_{m,\lambda}(\omega), && \text{(shift property)} \\
\hat{\varphi}_{m,\lambda}(\omega)\hat{\varphi}_{k,\lambda}(\omega) &= \frac{1}{\sqrt{2\lambda}}(\hat{\varphi}_{m+k,\lambda}(\omega) - \hat{\varphi}_{m+k+1,\lambda}(\omega)), && \text{(multiplication property)} \\
\frac{(2\lambda)^{\nu+1/2}}{(i\omega + \lambda)^{\nu+1}} &= \sum_{k=0}^{\nu} \binom{\nu}{k} (-1)^k \hat{\varphi}_{k,\lambda}(\omega). && \text{(binomial identity)}
\end{aligned}$$

### 3.2 Matérn–Laguerre functions

In view of Theorem 1.1, an orthonormal basis for the RKHS of the Matérn kernel  $r_{\nu+1/2,\lambda}$  is obtained from (3.1) and (3.3) in Fourier domain as

$$\hat{\psi}_{m,\lambda}^{(\nu)}(\omega) = \hat{h}_{\nu+1/2,\lambda}(\omega)\hat{\varphi}_{m,\lambda}(\omega) = (2\lambda)^{\nu+1} \frac{\nu!}{\sqrt{(2\nu)!}} \frac{(i\omega - \lambda)^m}{(i\omega + \lambda)^{m+1+\nu+1}} \quad (3.7)$$

for  $m \in \mathbb{Z}$ . We call the resulting functions the *Matérn–Laguerre functions*. They satisfy the scaling laws

$$\psi_{m,\lambda}^{(\nu)}(t) = \psi_{m,1/2}^{(\nu)}(2\lambda t) \quad \text{and} \quad \hat{\psi}_{m,\lambda}^{(\nu)}(\omega) = \frac{1}{2\lambda} \hat{\psi}_{m,1/2}^{(\nu)}(\omega/2\lambda). \quad (3.8)$$

Like the Laguerre functions, the Matérn–Laguerre functions also satisfy a certain conjugate symmetry property in the sense that

$$\hat{\psi}_{-\nu-1-m-1,\lambda}^{(\nu)}(\omega) = (-1)^\nu \hat{\psi}_{m,\lambda}^{(\nu)*}(\omega) \quad \text{for} \quad m \in \mathbb{N}_0. \quad (3.9)$$

Furthermore, by the binomial identity and the shift property of Laguerre functions, the Matérn–Laguerre functions and their Fourier transforms are

$$\psi_{m,\lambda}^{(\nu)}(t) = \frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \varphi_{m+k,\lambda}(t) \quad (3.10)$$

and

$$\hat{\psi}_{m,\lambda}^{(\nu)}(\omega) = \frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \hat{\varphi}_{m+k,\lambda}(\omega) \quad (3.11)$$

for  $m \in \mathbb{Z}$ . The Matérn kernel of order  $\nu + 1/2$  can therefore be expanded as

$$r_{\nu+1/2,\lambda}(t, u) = \sum_{m=-\infty}^{\infty} \psi_{m,\lambda}^{(\nu)}(t)\psi_{m,\lambda}^{(\nu)}(u). \quad (3.12)$$

The following proposition provides a uniform upper bound on the Matérn–Laguerre functions.

**Proposition 3.1** (Matérn–Laguerre upper bound). *For all  $t \in \mathbb{R}$  and  $m \in \mathbb{Z}$ , the Matérn–Laguerre functions satisfy*

$$|\psi_{m,\lambda}^{(\nu)}(t)| \leq \frac{2^\nu \nu!}{\sqrt{(2\nu)!}} \sim (\pi\nu)^{1/4} \quad \text{as} \quad \nu \rightarrow \infty.$$

*Proof.* By (3.11) and the binomial identity for Laguerre functions,

$$\psi_m^{(\nu)}(t) = \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu} \binom{\nu}{k} (-1)^k \frac{1}{2\pi} \int_{-\infty}^{\infty} \hat{\varphi}_{m,\lambda}(\omega) \hat{\varphi}_{k,\lambda}(\omega) e^{i\omega t} d\omega.$$

Apply triangle inequality, the Cauchy–Schwartz inequality, and the orthonormality in  $\mathcal{L}_2(\mathbb{R}, 1/2\pi)$  of  $\hat{\varphi}_{m,\lambda}$  to arrive at

$$\begin{aligned} |\psi_{m,\lambda}^{(\nu)}(t)| &\leq \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu} \binom{\nu}{k} \frac{1}{2\pi} \left| \int_{-\infty}^{\infty} e^{i\omega t} \hat{\varphi}_{m,\lambda}(\omega) \hat{\varphi}_{k,\lambda}(\omega) d\omega \right| \\ &\leq \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu} \binom{\nu}{k} \frac{1}{2\pi} \left( \int_{-\infty}^{\infty} |\hat{\varphi}_{m,\lambda}(\omega)|^2 d\omega \right)^{1/2} \left( \int_{-\infty}^{\infty} |\hat{\varphi}_{k,\lambda}(\omega)|^2 d\omega \right)^{1/2} \\ &= \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu} \binom{\nu}{k} \\ &= \frac{2^\nu \nu!}{\sqrt{(2\nu)!}}. \end{aligned}$$

The asymptotic equivalence as  $\nu \rightarrow \infty$  follows from Stirling’s formula.  $\square$

It appears difficult to improve upon the bound in Proposition 3.1. Consequently, uniform convergence of Matérn–Laguerre expansions on  $\mathbb{R}$  is likely unattainable.

### 3.3 Classification of Matérn–Laguerre functions

For  $m \in \mathbb{N}_0$ , a more compact and convenient expression of the Matérn–Laguerre functions (3.10) may be obtained by using the convolution formula in Theorem 1.1. For  $\eta \in \mathbb{N}_0$ , the associated Laguerre polynomial  $L_m^{(\eta)}$  is defined as

$$L_m^{(\eta)}(t) = \sum_{k=0}^m \binom{m+\eta}{m-k} \frac{(-1)^k}{k!} t^k. \quad (3.13)$$

The associated Laguerre polynomial  $L_m^{(0)}$  equals the Laguerre polynomial  $L_m$  in (3.5). For  $t > 0$ , we get from (3.2) and (3.6) that

$$\begin{aligned} \psi_{m,1/2}^{(\nu)}(t) &= \int_{-\infty}^{\infty} h_{\nu+1/2,1/2}(t-\tau) \varphi_{m,1/2}(\tau) d\tau \\ &= \frac{\nu!}{\sqrt{(2\nu)!}} \int_0^t e^{-(t-\tau)/2} \frac{(t-\tau)^\nu}{\nu!} e^{-\tau/2} L_m(\tau) d\tau \\ &= \frac{\nu!}{\sqrt{(2\nu)!}} e^{-t/2} \int_0^t \frac{(t-\tau)^\nu}{\nu!} L_m(\tau) d\tau \\ &= \frac{\nu!}{\sqrt{(2\nu)!}} \frac{m!}{(m+\nu+1)!} t^{\nu+1} L_m^{(\nu+1)}(t) e^{-t/2}, \end{aligned}$$

where the last equality follows from a convolution identity for Laguerre polynomials [3, Chapter 6, Problem (3)]. For  $t < 0$ , the Laguerre functions  $\varphi_{m,1/2}(t)$  vanish and the convolution evaluates to zero. Consequently, the scaling law (3.8) gives

$$\psi_{m,\lambda}^{(\nu)}(t) = \frac{\nu!}{\sqrt{(2\nu)!}} \frac{m!}{(m+\nu+1)!} (2\lambda t)^{\nu+1} L_m^{(\nu+1)}(2\lambda t) e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t) \quad \text{for } m \in \mathbb{N}_0.$$






Figure 1: The Matérn-Laguerre functions  $\psi_{m,\lambda}^{+,\nu}$  in (3.15) for  $\lambda = 4/5$  and  $m = 0, \dots, 6$ . Observe that the functions vanish on the negative real line.

For negative indices  $m \leq -\nu - 2$  a similar expression is obtained from the conjugate symmetry (3.9):

$$\begin{aligned} \psi_{-\nu-1-m-1,\lambda}^{(\nu)}(t) &= (-1)^\nu \psi_{m,\lambda}^{(\nu)}(-t) \\ &= -\frac{\nu!}{\sqrt{(2\nu)!}} \frac{m!}{(m+\nu+1)!} (2\lambda t)^{\nu+1} \mathbb{L}_m^{(\nu+1)}(2\lambda|t|) e^{-\lambda|t|} \mathbf{1}_{(-\infty,0)}(t) \end{aligned} \quad (3.14)$$

for  $m \in \mathbb{N}_0$ . This motivates the following notation for the three classes of Matérn-Laguerre functions that comprise an orthonormal basis:

$$\psi_{m,\lambda}^{+,\nu}(t) = \psi_{m,\lambda}^{(\nu)}(t) \quad \text{for } m \in \mathbb{N}_0, \quad (3.15)$$

$$\psi_{m,\lambda}^{-,\nu}(t) = (-1)^\nu \psi_{m,\lambda}^{(\nu)}(-t) \quad \text{for } m \in \mathbb{N}_0, \quad (3.16)$$

$$\psi_{m,\lambda}^{0,\nu}(t) = \psi_{-\nu-1+m,\lambda}^{(\nu)}(t) \quad \text{for } m = 0, 1, \dots, \nu. \quad (3.17)$$

For convenience, define the corresponding sets

$$\mathcal{M}_{\nu,\lambda}^+ = \{\psi_{m,\lambda}^{+,\nu}\}_{m \in \mathbb{N}_0}, \quad \mathcal{M}_{\nu,\lambda}^- = \{\psi_{m,\lambda}^{-,\nu}\}_{m \in \mathbb{N}_0}, \quad \mathcal{M}_{\nu,\lambda}^0 = \{\psi_{m,\lambda}^{0,\nu}\}_{m=0}^\nu,$$

the union  $\mathcal{M}_{\nu,\lambda} = \mathcal{M}_{\nu,\lambda}^- \cup \mathcal{M}_{\nu,\lambda}^+$ , and the kernels

$$\rho_{\nu+1/2,\lambda}^-(t,u) = \sum_{m=0}^{\infty} \psi_{m,\lambda}^{-,\nu}(t) \psi_{m,\lambda}^{-,\nu}(u) \quad \text{and} \quad \rho_{\nu+1/2,\lambda}^+(t,u) = \sum_{m=0}^{\infty} \psi_{m,\lambda}^{+,\nu}(t) \psi_{m,\lambda}^{+,\nu}(u). \quad (3.18)$$

We call the set  $\mathcal{M}_{\nu,\lambda}^0$  the *null-space* and study it in more detail in Section 3.5. For now, note that the null-space functions are supported on  $\mathbb{R}$  because from (3.10) one can see that for  $m = 0, \dots, \nu$  the sum that defines  $\psi_{-\nu-1+m,\lambda}^{(\nu)}$  contains Laguerre functions with both negative and non-negative index. Some of the basis functions are shown in Figures 1 and 4.

The Matérn expansion (3.12) can now be written in terms of these functions and kernels as

$$r_{\nu+1/2,\lambda}(t,u) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,\nu}(t) \psi_{m,\lambda}^{0,\nu}(u) + \rho_{\nu+1/2,\lambda}^-(t,u) + \rho_{\nu+1/2,\lambda}^+(t,u).$$

It is clear that the functions in  $\mathcal{M}_{\nu,\lambda}^-$  are supported on the negative real line and the functions in

$\mathcal{M}_{\nu,\lambda}^+$  on the positive real line. This observation yields the following simplifications:

$$r_{\nu+1/2,\lambda}(t, u) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,(\nu)}(t)\psi_{m,\lambda}^{0,(\nu)}(u) + \rho_{\nu+1/2,\lambda}^+(t, u) \quad \text{if } t \geq 0 \text{ or } u \geq 0, \quad (3.19)$$

$$r_{\nu+1/2,\lambda}(t, u) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,(\nu)}(t)\psi_{m,\lambda}^{0,(\nu)}(u) + \rho_{\nu+1/2,\lambda}^-(t, u) \quad \text{if } t \leq 0 \text{ or } u \leq 0, \quad (3.20)$$

$$r_{\nu+1/2,\lambda}(t, u) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,(\nu)}(t)\psi_{m,\lambda}^{0,(\nu)}(u) \quad \text{if } \text{sign } t \neq \text{sign } u. \quad (3.21)$$

We next show that  $\mathcal{M}_{\nu,\lambda}$ ,  $\mathcal{M}_{\nu,\lambda}^-$ , and  $\mathcal{M}_{\nu,\lambda}^+$  form orthogonal bases with respect to the weight function

$$w_{\nu,\lambda}(t) = 2\lambda/|2\lambda t|^{\nu+1}.$$

This justifies saying that the expansions we have derived for Matérn kernels are ‘‘almost’’ Mercer.

**Proposition 3.2** (Matérn–Laguerre orthogonality). *The sets  $\mathcal{M}_{\nu,\lambda}$ ,  $\mathcal{M}_{\nu,\lambda}^+$ , and  $\mathcal{M}_{\nu,\lambda}^-$  form orthogonal bases in  $\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})$ ,  $\mathcal{L}_2(\mathbb{R}_+, w_{\nu,\lambda})$ , and  $\mathcal{L}_2(\mathbb{R}_-, w_{\nu,\lambda})$ , respectively. Furthermore,*

$$\|\psi_{m,\lambda}^{+,( \nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})}^2 = \|\psi_{m,\lambda}^{-,( \nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})}^2 = \frac{(\nu!)^2}{(2\nu)!} \frac{m!}{(m + \nu + 1)!} \quad \text{for every } m \in \mathbb{N}_0.$$

*Proof.* That  $\mathcal{M}_{\nu,\lambda}^+$  forms an orthogonal basis in  $\mathcal{L}_2(\mathbb{R}_+, w_{\nu,\lambda})$  follows from the fact that the functions

$$t^{\nu/2+1/2} L_m^{(\nu+1)}(t) e^{-t/2} \quad \text{for } m \in \mathbb{N}_0 \quad (3.22)$$

form an orthonormal basis in  $\mathcal{L}_2(\mathbb{R}_+)$  [23, Theorem 5.7.1]. Furthermore, the norms of the functions in  $\mathcal{M}_{\nu,\lambda}^+$  are readily computed from the norms of the corresponding Laguerre polynomials:

$$\begin{aligned} \|\psi_{m,\lambda}^{+,( \nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})}^2 &= \frac{(\nu!)^2}{(2\nu)!} \left( \frac{m!}{(m + \nu + 1)!} \right)^2 \int_0^\infty [L_m^{(\nu+1)}(t)]^2 t^{\nu+1} e^{-t} dt \\ &= \frac{(\nu!)^2}{(2\nu)!} \frac{m!}{(m + \nu + 1)!}. \end{aligned}$$

The statement pertaining to  $\mathcal{M}_{\nu,\lambda}^-$  follows from the symmetry (3.14) and the statement pertaining to  $\mathcal{M}_{\nu,\lambda}$  from the fact that  $\mathcal{L}_2(\mathbb{R}) = \mathcal{L}_2(\mathbb{R}_-) \oplus \mathcal{L}_2(\mathbb{R}_+)$ .  $\square$

### 3.4 Truncation error

Define the kernel

$$\rho_{\nu+1/2,\lambda}(t, u) = \rho_{\nu+1/2,\lambda}^-(t, u) + \rho_{\nu+1/2,\lambda}^+(t, u) \quad (3.23)$$

in terms of the kernels in (3.18). A few translates of this kernel are displayed in Figure 2. The full Matérn kernel is therefore

$$r_{\nu+1/2,\lambda}(t, u) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,(\nu)}(t)\psi_{m,\lambda}^{0,(\nu)}(u) + \rho_{\nu+1/2,\lambda}(t, u).$$

From Proposition 3.2 we see that the kernel  $\rho_{\nu+1/2,\lambda}$  is an element of  $\mathcal{L}_2(\mathbb{R} \times \mathbb{R}, w_{\nu,\lambda} \otimes w_{\nu,\lambda})$  and that its squared norm is given by

$$\begin{aligned} &\int_{-\infty}^\infty \int_{-\infty}^\infty \rho_{\nu+1/2,\lambda}^2(t, u) w_{\nu,\lambda}(t) w_{\nu,\lambda}(u) dt du \\ &= \sum_{m=0}^\infty (\|\psi_{m,\lambda}^{-,( \nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})}^4 + \|\psi_{m,\lambda}^{+,( \nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu,\lambda})}^4). \end{aligned}$$




Figure 2: Translates  $\rho_{\nu+1/2, \lambda}(\cdot, u)$  of the kernel in (3.23) for  $u \in \{-2, -1.2, -0.4, 0.4, 1.2, 2\}$  and  $\lambda = 4/5$ . Observe that each translate is supported on the axis that  $u$  lies on.

This implies that  $\rho_{\nu+1/2, \lambda}$  defines a Hilbert–Schmidt operator on  $\mathcal{L}_2(\mathbb{R}, w_{\nu, \lambda})$  via (1.3) and that the above norm is precisely the squared Hilbert–Schmidt norm of this operator [15, Chapter 1, §1]. Next the approximation errors for appropriately truncated approximations of the Matérn kernel are examined in terms of the Hilbert–Schmidt norm. Let  $n \geq 1$  and define the truncated kernels

$$\rho_{\nu+1/2, \lambda, n}(t, u) = \sum_{m=0}^{n-1} \psi_{m, \lambda}^{-, (\nu)}(t) \psi_{m, \lambda}^{-, (\nu)}(u) + \sum_{m=0}^{n-1} \psi_{m, \lambda}^{+, (\nu)}(t) \psi_{m, \lambda}^{+, (\nu)}(u), \quad (3.24)$$

$$r_{\nu+1/2, \lambda, n}(t, u) = \sum_{m=0}^{\nu} \psi_{m, \lambda}^{0, (\nu)}(t) \psi_{m, \lambda}^{0, (\nu)}(u) + \rho_{\nu+1/2, \lambda, n}(t, u). \quad (3.25)$$

Observe that  $r_{\nu+1/2, \lambda, n}$  is a finite expansion of  $\nu + 1 + 2n$  terms. Some truncations of Matérn kernels are displayed in Figure 3.

**Proposition 3.3** (Matérn truncation). *For every  $n \in \mathbb{N}$  it holds that*

$$\left( \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (r_{\nu+1/2, \lambda}(t, u) - r_{\nu+1/2, \lambda, n}(t, u))^2 w_{\nu, \lambda}(t) w_{\nu, \lambda}(u) dt du \right)^{1/2} \leq \frac{c_{\nu}}{n^{\nu+1/2}},$$

where

$$c_{\nu} = \frac{(\nu!)^2}{(2\nu)!} \sqrt{\frac{2(2\nu+2)}{2\nu+1}} \sim \frac{1}{2^{2\nu}} \sqrt{\frac{2\pi(2\nu+2)\nu}{2\nu+1}} \quad \text{as } \nu \rightarrow \infty.$$

*Proof.* Firstly, the truncation error is

$$\begin{aligned} & r_{\nu+1/2, \lambda}(t, u) - r_{\nu+1/2, \lambda, n}(t, u) \\ &= \rho_{\nu+1/2, \lambda}(t, u) - \rho_{\nu+1/2, \lambda, n}(t, u) \\ &= \sum_{m=n}^{\infty} [\psi_{m, \lambda}^{-, (\nu)}(t, u) \psi_{m, \lambda}^{-, (\nu)}(t, u) + \psi_{m, \lambda}^{+, (\nu)}(t, u) \psi_{m, \lambda}^{+, (\nu)}(t, u)]. \end{aligned}$$




Figure 3: The truncation in (3.25) for two Matérn kernels with  $\lambda = 4/5$ . Because the second kernel argument has been fixed to a positive value, the truncations are exact on the negative real line by (3.21).

Using Proposition 3.2, the squared norm of the truncation error is straight-forwardly computed as

$$\begin{aligned}
& \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (r_{\nu+1/2, \lambda}(t, u) - r_{\nu+1/2, \lambda, n}(t, u))^2 w_{\nu, \lambda}(t) w_{\nu, \lambda}(u) dt du \\
&= \sum_{m=n}^{\infty} (\|\psi_{m, \lambda}^{-, (\nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu, \lambda})}^4 + \|\psi_{m, \lambda}^{+, (\nu)}\|_{\mathcal{L}_2(\mathbb{R}, w_{\nu, \lambda})}^4) \\
&= 2 \left( \frac{(\nu!)^2}{(2\nu)!} \right)^2 \sum_{m=n}^{\infty} \left( \frac{m!}{(m + \nu + 1)!} \right)^2 \\
&\leq 2 \left( \frac{(\nu!)^2}{(2\nu)!} \right)^2 \sum_{m=n}^{\infty} \frac{1}{m^{2\nu+2}}.
\end{aligned}$$

The sum may be estimated with an integral as

$$\sum_{m=n}^{\infty} \frac{1}{m^{2\nu+2}} \leq \frac{1}{n^{2\nu+2}} + \int_n^{\infty} \frac{1}{t^{2\nu+2}} dt = \frac{1}{n^{2\nu+2}} + \frac{1}{2\nu+1} \frac{1}{n^{2\nu+1}} \leq \frac{2\nu+2}{2\nu+1} \frac{1}{n^{2\nu+1}},$$

where  $n \geq 1$  was used in the last inequality. This yields the desired upper bound. The asymptotic equivalence for  $c_\nu$  as  $\nu \rightarrow \infty$  follows from Stirling's formula.  $\square$

### 3.5 The null-space $\mathcal{M}_{\nu, \lambda}^0$

In view of Proposition 3.2,  $\mathcal{M}_{\nu, \lambda}^0$  is left as the odd set out. From (3.1) and (3.7) we compute that

$$\hat{\psi}_{m, \lambda}^{0, (\nu)}(\omega) = (-1)^{\nu+1} \hat{h}_{\nu+1/2, \lambda}^*(\omega) \hat{\varphi}_{m, \lambda}(\omega).$$

Furthermore, the functions

$$(i\omega + \lambda)^{\nu+1} \hat{\psi}_{m, \lambda}^{0, (\nu)}(\omega)$$

viewed as functions of  $i\omega$ , have no poles in the left half-plane. Therefore  $\mathcal{M}_{\nu, \lambda}^0$  are annihilated on the positive real line by the differential operator  $(D + \lambda)^{\nu+1}$ . That is,

$$(D + \lambda)^{\nu+1} \hat{\psi}_{m, \lambda}^{0, (\nu)}(t) = 0 \quad \text{for every } t > 0.$$




Figure 4: The null-space Matérn–Laguerre functions  $\psi_{m,\lambda}^{0,(\nu)}$  in (3.17) for  $\lambda = 4/5$ .

For this reason we refer to these functions as the null-space functions. The null space functions have a symmetry property similar to that of the functions  $\mathcal{M}_{\nu,\lambda}$  given by (3.15) and (3.16).

**Proposition 3.4** (Null-space symmetry). *The null-space functions satisfy*

$$\psi_{\nu-m,\lambda}^{0,(\nu)}(t) = (-1)^\nu \psi_m^{0,(\nu)}(-t) \quad \text{and} \quad \hat{\psi}_{\nu-m,\lambda}^{0,(\nu)}(\omega) = (-1)^\nu \hat{\psi}_m^{0,(\nu)*}(\omega).$$

for  $m = 0, 1, \dots, \nu$ .

*Proof.* Starting from (3.11), using the conjugate symmetry of Laguerre functions, and then changing the order of summation gives

$$\begin{aligned} \hat{\psi}_{m,\lambda}^{0,(\nu)}(\omega) &= \frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \hat{\varphi}_{-\nu-1+m+k,\lambda}(\omega) \\ &= \frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \hat{\varphi}_{-(\nu-m-k)-1,\lambda}(\omega) \\ &= -\frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \hat{\varphi}_{\nu-m-k,\lambda}^*(\omega) \\ &= -\frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{\nu+1-k} (-1)^{\nu+1-k} \hat{\varphi}_{\nu-m-(\nu+1-k),\lambda}^*(\omega) \\ &= (-1)^\nu \frac{1}{\sqrt{2\lambda}} \frac{\nu!}{\sqrt{(2\nu)!}} \sum_{k=0}^{\nu+1} \binom{\nu+1}{k} (-1)^k \hat{\varphi}_{-\nu-1+\nu-m+k,\lambda}^*(\omega) \\ &= (-1)^\nu \hat{\psi}_{\nu-m,\lambda}^{0,(\nu)*}(\omega), \end{aligned}$$

which is the Fourier domain symmetry. The time domain symmetry is then obtained from Fourier inversion.  $\square$

**Example 3.5** (Null-space functions). *The set  $\mathcal{M}_{0,\lambda}^0$  (i.e.,  $\nu = 0$ ) consists of the function*

$$\psi_{0,\lambda}^{0,(0)}(t) = -e^{-\lambda|t|}.$$

The set  $\mathcal{M}_{1,\lambda}^0$  (i.e.,  $\nu = 1$ ) consists of the functions

$$\begin{aligned}\psi_{0,\lambda}^{0,(1)}(t) &= \frac{1}{\sqrt{2}}(2\lambda t e^{\lambda t} \mathbf{1}_{(-\infty,0)}(t) + e^{-\lambda|t|}), \\ \psi_{1,\lambda}^{0,(1)}(t) &= -\frac{1}{\sqrt{2}}(2\lambda t e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t) + e^{-\lambda|t|}).\end{aligned}$$

The set  $\mathcal{M}_{2,\lambda}^0$  (i.e.,  $\nu = 2$ ) consists of the functions

$$\begin{aligned}\psi_{0,\lambda}^{0,(2)}(t) &= \frac{2}{\sqrt{4!}}(2(-\lambda^2 t^2 + \lambda t)e^{\lambda t} \mathbf{1}_{(-\infty,0)}(t) - e^{-\lambda|t|}), \\ \psi_{1,\lambda}^{0,(2)}(t) &= \frac{4}{\sqrt{4!}}(\lambda|t| + 1)e^{-\lambda|t|}, \\ \psi_{2,\lambda}^{0,(2)}(t) &= \frac{2}{\sqrt{4!}}(-2(\lambda^2 t^2 + \lambda t)e^{-\lambda t} \mathbf{1}_{[0,\infty)}(t) - e^{-\lambda|t|}).\end{aligned}$$

Some null space functions are depicted in Figure 4. Unlike the basis functions  $\mathcal{M}_{\nu,\lambda}^+$  depicted in Figure 1, the null space functions are supported on the entire real line. For  $d = |t - u|$ , a Matérn kernel can be written as

$$r_{\nu+1/2,\lambda}(t, u) = r_{\nu+1/2,\lambda}(0, d) = \sum_{m=0}^{\nu} \psi_{m,\lambda}^{0,(\nu)}(0) \psi_{m,\lambda}^{0,(\nu)}(d),$$

where we have used (3.19) and the fact that the kernel  $\rho_{\nu+1/2,\lambda}^+(t, u)$  vanishes if  $t = 0$  or  $u = 0$ . Upon substitution of the expressions in Example 3.5 we obtain the well-known explicit forms of Matérn kernels in terms of  $d$ , such as

$$r_{3/2,\lambda}(t, u) = (1 + \lambda d)e^{-\lambda d} \quad \text{and} \quad r_{5/2,\lambda}(t, u) = \left(1 + \lambda d + \frac{\lambda^2 d^2}{3}\right)e^{-\lambda d}.$$

## 4 Cauchy kernel

The Cauchy kernel and its Fourier transform are

$$r_{\lambda}(t, u) = \frac{1}{1 + \lambda^2(t - u)^2} \quad \text{and} \quad \hat{\Phi}_{\lambda}(\omega) = \frac{\pi}{\lambda} e^{-|\omega|/\lambda}. \quad (4.1)$$

The Cauchy kernel is thus a Fourier dual to the Matérn kernel of smoothness index  $\alpha = 1/2$  (i.e.,  $\nu = 0$ ). In what follows this will inform the construction of an RKHS basis. To emphasize this relationship, we use the reparametrisation  $\lambda = 1/\gamma$ . A square-root of  $\hat{\Phi}_{1/\gamma}(\omega)$  is then given by

$$\hat{h}_{1/\gamma}(\omega) = \hat{\Phi}_{1/\gamma}(\omega)^{1/2} = \sqrt{\pi\gamma} e^{-\gamma|\omega|/2}. \quad (4.2)$$

### 4.1 Expansion in complex-valued Cauchy–Laguerre functions

In view of the Fourier dualism with the Matérn- $\frac{1}{2}$  kernel and the fact that the Fourier transform is an isometry from  $\mathcal{L}_2(\mathbb{R})$  to  $\mathcal{L}_2(\mathbb{R}, 1/2\pi)$ , a straight-forward way to construct a suitable basis in  $\mathcal{L}_2(\mathbb{R}, 1/2\pi)$  is to pick the Laguerre functions  $\varphi_{m,\gamma}(\omega)$  from Section 3.1 and scale them by  $\sqrt{2\pi}$ . These  $\mathcal{L}_2(\mathbb{R})$  basis functions and (4.2) yield the RKHS basis functions

$$\hat{\psi}_{m,\gamma}(\omega) = \sqrt{\pi\gamma} e^{-\gamma|\omega|/2} \sqrt{2\pi} \varphi_{m,\gamma/2}(\omega)$$

in the Fourier domain. Because their inverse Fourier transforms are complex-valued, we call these functions the *complex-valued Cauchy–Laguerre functions*. For  $m \in \mathbb{N}_0$ , Fourier inversion gives

$$\begin{aligned}\psi_{m,\gamma}(t) &= \frac{\sqrt{2\gamma}}{2} \int_{-\infty}^{\infty} e^{-\gamma|w|/2} \varphi_{m,\gamma/2}(\omega) e^{i\omega t} d\omega = \frac{\sqrt{\gamma}}{\sqrt{2}} \int_0^{\infty} \varphi_{m,\gamma/2}(\omega) e^{-i\omega(-i\gamma/2-t)} d\omega \\ &= \frac{\sqrt{\gamma}}{\sqrt{2}} \hat{\varphi}_{m,\gamma/2}(-i\gamma/2-t) \\ &= \frac{\gamma}{\sqrt{2}} \frac{(\gamma/2 - it - \gamma/2)^m}{(\gamma/2 - it + \gamma/2)^{m+1}} \\ &= -\frac{\gamma}{\sqrt{2}} \frac{(it)^m}{(it - \gamma)^{m+1}}.\end{aligned}$$

Similarly, for negative indices we get

$$\begin{aligned}\psi_{-m,\gamma}(t) &= \frac{\sqrt{2\gamma}}{2} \int_{-\infty}^{\infty} e^{-\gamma|w|/2} \varphi_{-m,\gamma/2}(\omega) e^{i\omega t} d\omega \\ &= -\frac{\sqrt{\gamma}}{\sqrt{2}} \int_{-\infty}^0 \varphi_{m-1,\gamma/2}(-\omega) e^{\omega(\gamma/2+it)} d\omega \\ &= -\frac{\sqrt{\gamma}}{\sqrt{2}} \int_0^{\infty} \varphi_{m-1,\gamma/2}(\omega) e^{-i\omega(-i\gamma/2+t)} d\omega \\ &= -\frac{\sqrt{\gamma}}{\sqrt{2}} \hat{\varphi}_{m-1,\gamma/2}(-i\gamma/2+t) = -\frac{\gamma}{\sqrt{2}} \frac{(\gamma/2 + it - \gamma/2)^{m-1}}{(\gamma/2 + it + \gamma/2)^m} = -\frac{\gamma}{\sqrt{2}} \frac{(it)^{m-1}}{(it + \gamma)^m}.\end{aligned}$$

To summarise, the complex valued Cauchy–Laguerre functions are

$$\psi_{m,\gamma}(t) = -\frac{\gamma}{\sqrt{2}} \frac{(it)^m}{(it - \gamma)^{m+1}} \quad \text{for } m \in \mathbb{N}_0, \quad (4.3a)$$

$$\psi_{-m-1,\gamma}(t) = -\frac{\gamma}{\sqrt{2}} \frac{(it)^m}{(it + \gamma)^{m+1}} \quad \text{for } m \in \mathbb{N}_0. \quad (4.3b)$$

They have the conjugate symmetry property

$$\psi_{m,\gamma}^*(t) = -\psi_{-m-1,\gamma}(t) = \psi_{m,\gamma}(-t) \quad \text{for } m \in \mathbb{Z}.$$

An expansion of the Cauchy kernel (4.1) in terms of complex-valued Cauchy–Laguerre functions is thus given by

$$r_\lambda(t, u) = \sum_{m=-\infty}^{\infty} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(u).$$

This expansion is remarkably easy to verify since geometric summation and conjugate symmetry yield

$$\sum_{m=0}^{\infty} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(u) = \frac{1}{2\lambda^2} \frac{1}{(it - 1/\lambda)(-iu - 1/\lambda) - tu}$$

and

$$\sum_{m=-\infty}^{-1} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(u) = \left( \sum_{m=0}^{\infty} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(u) \right)^*.$$

Hence

$$\sum_{m=-\infty}^{\infty} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(u) = \frac{1}{2} \frac{1}{1 - i\lambda(t - u)} + \frac{1}{2} \frac{1}{1 + i\lambda(t - u)} = \frac{1}{1 + \lambda^2(t - u)^2},$$




Figure 5: The trigonometric Cauchy–Laguerre functions  $\alpha_{m,\gamma}$  and  $\beta_{m,\gamma}$  in (4.4) for  $\lambda = \gamma = 1$ .

which indeed is the Cauchy kernel. An appropriate  $\mathcal{L}_2(\mathbb{R}, w)$  space in which the complex-valued Cauchy–Laguerre functions form a complete orthogonal set remains elusive to us. However, just as with the Matérn–Laguerre expansions in Section 3, the present expansion is very good at origin since all but two terms vanish:

$$r_\lambda(t, 0) = \sum_{m=-\infty}^{\infty} \psi_{m,1/\lambda}^*(t) \psi_{m,1/\lambda}(0) = -\frac{1}{\lambda} (\psi_{-1,1/\lambda}(t) + \psi_{0,1/\lambda}(t)).$$

## 4.2 Expansion in trigonometric Cauchy–Laguerre functions

It would be desirable to obtain a real-valued basis for the Cauchy RKHS. This can be done by scaling the real and imaginary parts of  $\hat{\varphi}_{m,\lambda}$  [4]. The resulting *trigonometric Laguerre functions*, which form an orthonormal basis of  $\mathcal{L}_2(\mathbb{R})$ , are given by

$$\begin{aligned} \hat{C}_{m,\gamma}(\omega) &= \frac{1}{\sqrt{2}} (\hat{\varphi}_{m,\gamma}(\omega) + \hat{\varphi}_{m,\gamma}^*(\omega)) \quad \text{for } m \in \mathbb{N}_0, \\ \hat{S}_{m,\gamma}(\omega) &= \frac{1}{i\sqrt{2}} (\hat{\varphi}_{m,\gamma}(\omega) - \hat{\varphi}_{m,\gamma}^*(\omega)) \quad \text{for } m \in \mathbb{N}_0 \end{aligned}$$

in the Fourier domain. These functions are so named because they resemble the trigonometric functions by being even and odd, respectively. Explicit expressions are [4, Section 3]

$$\begin{aligned} \hat{C}_{m,\gamma}(\omega) &= 2 \frac{\sqrt{\gamma} (-1)^{m+2}}{(\omega^2 + \gamma^2)^{m+1}} \sum_{n=0}^m \binom{2m+1}{2n} (-1)^n \omega^{2n} (2\gamma)^{2m+1-2n}, \\ \hat{S}_{m,\gamma}(\omega) &= 2 \frac{\sqrt{\gamma} (-1)^{m+1}}{(\omega^2 + \gamma^2)^{m+1}} \sum_{n=0}^m \binom{2m+1}{2n+1} (-1)^n \omega^{2n+1} (2\gamma)^{2m-2n}. \end{aligned}$$

In time domain, the functions are

$$C_{m,\gamma}(t) = \frac{1}{\sqrt{2}} (\varphi_{m,\gamma}(t) + \varphi_{m,\gamma}(-t)) \quad \text{and} \quad S_{m,\gamma}(t) = \frac{1}{i\sqrt{2}} (\varphi_{m,\gamma}(t) - \varphi_{m,\gamma}(-t)),$$

which have the more compact expressions

$$C_{m,\gamma}(t) = \frac{1}{\sqrt{2}} \varphi_{m,\gamma}(|t|) \quad \text{and} \quad S_{m,\gamma}(t) = \frac{1}{i\sqrt{2}} \text{sign}(t) \varphi_{m,\gamma}(|t|).$$



The results in Section 4.1 give the RKHS basis functions

$$\alpha_{m,\gamma}(t) = \frac{1}{\sqrt{2}}(\psi_{m,\gamma}(t) + \psi_{m,\gamma}^*(t)) \quad \text{and} \quad \beta_{m,\gamma}(t) = \frac{1}{i\sqrt{2}}(\psi_{m,\gamma}(t) - \psi_{m,\gamma}^*(t)) \quad (4.4)$$

for  $m \in \mathbb{N}_0$ , where  $\psi_{m,\gamma}$  are the complex-valued Cauchy–Laguerre functions in (4.3). We call the functions  $\alpha_{m,\gamma}$  and  $\beta_{m,\gamma}$  the *trigonometric Cauchy–Laguerre functions*. These functions can be computed by taking the real and imaginary parts of  $\sqrt{2}\psi_{m,\gamma}(t)$ . The binomial theorem yields the explicit expressions

$$\alpha_{m,\gamma}(t) = \frac{\gamma}{2} \frac{(-1)^m (it)^m}{(t^2 + \gamma^2)^{m+1}} \sum_{k=0}^{m+1} \binom{m+1}{k} (it)^k \gamma^{m+1-k} (1 - (-1)^{m+1-k}),$$

$$\beta_{m,\gamma}(t) = \frac{\gamma}{i2} \frac{(-1)^m (it)^m}{(t^2 + \gamma^2)^{m+1}} \sum_{k=0}^{m+1} \binom{m+1}{k} (it)^k \gamma^{m+1-k} (1 + (-1)^{m+1-k}),$$

which can be transformed into expressions of only real parameters by considering even and odd  $m$  separately. This yields

$$\alpha_{2m,\gamma}(t) = \gamma \frac{(-1)^m t^{2m}}{(t^2 + \gamma^2)^{2m+1}} \sum_{k=0}^m \binom{2m+1}{2k} (-1)^k t^{2k} \gamma^{2m+1-2k}, \quad (4.5a)$$

$$\alpha_{2m+1,\gamma}(t) = \gamma \frac{(-1)^m t^{2m+1}}{(t^2 + \gamma^2)^{2m+2}} \sum_{k=0}^m \binom{2m+2}{2k+1} (-1)^k t^{2k+1} \gamma^{2m+1-2k}, \quad (4.5b)$$

$$\beta_{2m,\gamma}(t) = \gamma \frac{(-1)^m t^{2m}}{(t^2 + \gamma^2)^{2m+1}} \sum_{k=0}^m \binom{2m+1}{2k+1} (-1)^k t^{2k+1} \gamma^{2m-2k}, \quad (4.5c)$$

$$\beta_{2m+1,\gamma}(t) = \gamma \frac{(-1)^{m+1} t^{2m+1}}{(t^2 + \gamma^2)^{2m+2}} \sum_{k=0}^{m+1} \binom{2m+2}{2k} (-1)^k t^{2k} \gamma^{2m+2-2k}. \quad (4.5d)$$

An expansion of the Cauchy kernel (4.1) in terms of real functions is thus given by

$$r_\lambda(t, u) = \sum_{m=0}^{\infty} \alpha_{m,1/\lambda}(t) \alpha_{m,1/\lambda}(u) + \sum_{m=0}^{\infty} \beta_{m,1/\lambda}(t) \beta_{m,1/\lambda}(u). \quad (4.6)$$

At the origin, this reduces to the finite term expansion

$$r_\lambda(t, 0) = \alpha_{0,1/\lambda}(t) \alpha_{0,1/\lambda}(0).$$

The basis functions  $\alpha_{m,\gamma}$  and  $\beta_{m,\gamma}$  and truncations of the expansion (4.6) are displayed in Figures 5 and 6.

## 5 Gaussian kernel

The Gaussian kernel and its Fourier transform are

$$r_\lambda(t, u) = \exp\left(-\frac{\lambda^2}{2}(t-u)^2\right) \quad \text{and} \quad \hat{\Phi}_\lambda(\omega) = \sqrt{\frac{2\pi}{\lambda^2}} e^{-\omega^2/2\lambda^2}. \quad (5.1)$$

A square-root is

$$\hat{h}_\lambda(\omega) = \hat{\Phi}_\lambda(\omega)^{1/2} = \left(\frac{2\pi}{\lambda^2}\right)^{1/4} e^{-\omega^2/4\lambda^2},$$

so that taking the inverse Fourier transform gives the function  $h$  in Theorem 1.1 as

$$h_\lambda(t) = 2^{1/4} \pi^{-1/4} \sqrt{\lambda} e^{-\lambda^2 t^2}. \quad (5.2)$$




Figure 6: Truncations  $\sum_{m=0}^{n-1} \alpha_{m,1/\lambda}(t)\alpha_{m,1/\lambda}(u) + \sum_{m=0}^{n-1} \beta_{m,1/\lambda}(t)\beta_{m,1/\lambda}(u)$  of the Cauchy expansion (4.6) for  $\lambda = \gamma = 1$ .

## 5.1 Expansion for the Gaussian kernel

As an orthonormal basis of  $\mathcal{L}_2(\mathbb{R})$  we use the *Hermite functions* (for them being an orthonormal basis, see [23, Theorem 5.7.1])

$$\varphi_{m,\lambda}(t) = \sqrt{\frac{\lambda}{2^m m! \sqrt{\pi}}} e^{-\lambda^2 t^2 / 2} \mathbf{H}_m(\lambda t) \quad \text{for } m \in \mathbb{N}_0. \quad (5.3)$$

Here  $\mathbf{H}_m$  is the  $m$ th physicist's Hermite polynomial given by

$$\mathbf{H}_m(t) = m! \sum_{k=0}^{\lfloor m/2 \rfloor} \frac{(-1)^k}{k!(m-2k)!} (2t)^{m-2k}. \quad (5.4)$$

By Theorem 1.1, the functions

$$\begin{aligned} \psi_{m,\lambda}(t) &= \int_{-\infty}^{\infty} h_\lambda(t-\tau) \varphi_{m,\lambda}(\tau) d\tau \\ &= \left( \frac{\sqrt{2}\lambda^2}{\pi} \right)^{1/2} \sqrt{\frac{1}{2^m m!}} \int_{-\infty}^{\infty} e^{-\lambda^2(t-\tau)^2} e^{-\lambda^2 \tau^2 / 2} \mathbf{H}_m(\lambda \tau) d\tau \end{aligned}$$

form an orthonormal basis of the RKHS of the Gaussian kernel (5.1). Equation (17) in Section 16.5 of [6] states that

$$\int_{-\infty}^{\infty} e^{-(s-\tau)^2} \mathbf{H}_m(a\tau) d\tau = \sqrt{\pi} (1-a^2)^{m/2} \mathbf{H}_m\left(\frac{as}{\sqrt{1-a^2}}\right)$$

for any reals  $s$  and  $a$ . Completing the square, doing a change of variables, and using this equation yields

$$\begin{aligned} \int_{-\infty}^{\infty} e^{-\lambda^2(t-\tau)^2} e^{-\lambda^2 \tau^2 / 2} \mathbf{H}_m(\lambda \tau) d\tau &= \sqrt{\frac{2}{3\lambda^2}} e^{-\lambda^2 t^2 / 3} \int_{-\infty}^{\infty} e^{-(\sqrt{2/3}\lambda t - \tau)^2} \mathbf{H}_m(\sqrt{2/3}\tau) d\tau \\ &= \sqrt{\frac{2\pi}{3\lambda^2}} 3^{-m/2} e^{-\lambda^2 t^2 / 3} \mathbf{H}_m\left(\frac{2\lambda t}{\sqrt{3}}\right). \end{aligned}$$

We thus obtain the basis functions

$$\psi_{m,\lambda}(t) = \left( \frac{2\sqrt{2}}{3} \right)^{1/2} \sqrt{\frac{1}{6^m m!}} e^{-\lambda^2 t^2 / 3} \mathbf{H}_m\left(\frac{2\lambda t}{\sqrt{3}}\right) \quad \text{for } m \in \mathbb{N}_0 \quad (5.5)$$




Figure 7: The first six basis functions  $\psi_{m,\lambda}$  in (5.5) of the Gaussian kernel (5.1) with  $\lambda = 4/5$ .

and the resulting expansion

$$r_\lambda(t, u) = \sum_{m=0}^{\infty} \psi_{m,\lambda}(t)\psi_{m,\lambda}(u) \quad (5.6)$$

of the Gaussian kernel in (5.1). Figure 7 displays some of the basis functions.

Note that the basis functions can be written in terms of the Hermite functions (5.3) by using the multiplication theorem

$$H_m(bt) = \sum_{k=0}^{\lfloor m/2 \rfloor} b^{m-2k} (b^2 - 1)^k \binom{m}{2k} \frac{(2k)!}{k!} H_{m-2k}(t)$$

for Hermite polynomials. Setting  $b = \sqrt{2}$  gives

$$H_m\left(\frac{2\lambda t}{\sqrt{3}}\right) = 2^{m/2} \sum_{k=0}^{\lfloor m/2 \rfloor} 2^{-k} \binom{m}{2k} \frac{(2k)!}{k!} H_{m-2k}\left(\frac{\sqrt{2}\lambda t}{\sqrt{3}}\right),$$

so that

$$\psi_{m,\lambda}(t) = \left(\frac{2\sqrt{\pi}}{\sqrt{3}\lambda}\right)^{1/2} \sqrt{\frac{2^m m!}{3^m}} \sum_{k=0}^{\lfloor m/2 \rfloor} \frac{1}{4^k k! \sqrt{(m-2k)!}} \varphi_{m-2k,\lambda}\left(\frac{\sqrt{2}t}{\sqrt{3}}\right).$$

It would be interesting to be able to connect  $\psi_{m,\lambda}$  to the associated Hermite polynomials [2] like the Matérn–Laguerre functions are connected to associated Laguerre functions in Section 3.3.

## 5.2 Mercer basis and Mehler's formula

The expansion (5.6) that we derived for the Gaussian kernel by the use of the basis functions in (5.5) can also be derived by setting

$$\rho = \frac{1}{3}, \quad x = \frac{2\lambda t}{\sqrt{3}}, \quad \text{and} \quad y = \frac{2\lambda u}{\sqrt{3}}$$

in Mehler's formula

$$\sum_{m=0}^{\infty} \frac{(\rho/2)^m}{m!} H_m(x)H_m(y)e^{-(x^2+y^2)/2} = \sqrt{\frac{1}{1-\rho^2}} \exp\left(\frac{4xy\rho - (1+\rho^2)(x^2+y^2)}{2(1-\rho^2)}\right)$$

and subsequently multiplying both sides by  $e^{-\lambda^2(t^2+u^2)/3}$ . This suggests that the expansion derived in the preceding section is a special case of the relatively well known Mercer expansion

of the Gaussian kernel, which too can be derived from Mehler's formula [8, Section 12.2.1]. Let  $\alpha > 0$  and define the constants

$$\beta = \left(1 + \frac{2\lambda^2}{\alpha^2}\right)^{1/4} \quad \text{and} \quad \delta^2 = \frac{\alpha^2}{2}(\beta^2 - 1).$$

The Mercer expansion of the Gaussian kernel with respect to the weight function

$$w_\alpha(t) = \frac{\alpha}{\sqrt{\pi}} e^{-\alpha^2 t^2}$$

on the real line is

$$r_\lambda(t, u) = \sum_{m=0}^{\infty} \mu_{m,\lambda,\alpha} \phi_{m,\lambda,\alpha}(t) \phi_{m,\lambda,\alpha}(u), \quad (5.7)$$

where

$$\mu_{m,\lambda,\alpha} = \sqrt{\frac{\alpha^2}{\alpha^2 + \delta^2 + \lambda^2/2}} \left( \frac{\lambda^2/2}{\alpha^2 + \delta^2 + \lambda^2/2} \right)^m$$

are the eigenvalues and

$$\phi_{m,\lambda,\alpha}(t) = \sqrt{\frac{\beta}{2^m m!}} e^{-\delta^2 t^2} \mathbb{H}_m(\alpha \beta t) \quad (5.8)$$

the  $\mathcal{L}_2(\mathbb{R}, w_\alpha)$ -orthonormal eigenfunctions of the integral operator in (1.3). By requiring that  $\alpha\beta = 2\lambda/\sqrt{3}$ , so that the Hermite polynomials appearing in (5.5) and (5.8) have the same scaling, it is straight-forward to solve that

$$\psi_{m,\lambda} = \sqrt{\mu_{m,\lambda,\alpha}} \phi_{m,\lambda,\alpha} = \sqrt{\frac{2}{3^{m+1}}} \phi_{m,\lambda,\alpha} \quad \text{when} \quad \alpha = \sqrt{\frac{2}{3}} \lambda,$$

which shows that the basis (5.5) is a special case of the Mercer basis. Results of some of the above computations are collected in the following proposition.

**Proposition 5.1** (Orthogonality of the Gaussian basis). *Let  $\alpha = \sqrt{2/3} \lambda$ . The functions*

$$\sqrt{\frac{3^{m+1}}{2}} \psi_{m,\lambda} = 2^{1/4} \sqrt{\frac{1}{2^m m!}} e^{-\lambda^2 t^2/3} \mathbb{H}_m\left(\frac{2\lambda t}{\sqrt{3}}\right) \quad \text{for} \quad m \in \mathbb{N}_0$$

*form an orthonormal basis of  $\mathcal{L}_2(\mathbb{R}, w_\alpha)$ .*

Although the Mercer expansion (5.7) has been known for some time, apparently originating in [28, Section 4], all its derivations in the literature that we are aware of are based on Mehler's formula and integral identities for Hermite polynomials (the only detailed derivation that we know of is given in [8, Section 12.2.1]). The expansion (5.6) is therefore the first Mercer expansion for the Gaussian kernel that has been derived from some general principle, which in this case is Theorem 1.1, instead of utilising *ad hoc* calculations. The relative simplicity of the basis functions (5.5) and the fact that the Hermite functions (5.3) have the same exponential decay as the kernel suggest that the choice  $\alpha = \sqrt{2/3} \lambda$  for the standard deviation of the Gaussian weight  $w_\alpha$  may be in some sense the most natural one. More discussion on the selection of  $\alpha$  may be found in [9, Section 5.3].

### 5.3 Truncation error

Define the truncated kernel

$$r_{\lambda,n}(t, u) = \sum_{m=0}^{n-1} \psi_{m,\lambda}(t) \psi_{m,\lambda}(u) \quad (5.9)$$




Figure 8: The Gaussian kernel (5.1) with  $s = 0$  and  $\lambda = 4/5$  and its truncated expansions (5.9). For  $n = 3$  and  $n = 11$  the truncated kernels become negative.

for any  $n \in \mathbb{N}$ . A few truncations are shown in Figure 8. The truncated kernel converges to the full Gaussian kernel  $r_\lambda$  pointwise on  $\mathbb{R} \times \mathbb{R}$ . The following proposition shows that the convergence of (5.9) to  $r_\lambda$  is exponential in  $\mathcal{L}_2(\mathbb{R} \times \mathbb{R}, w_\alpha \otimes w_\alpha)$ .

**Proposition 5.2** (Gaussian truncation). *Let  $\alpha = \sqrt{2/3} \lambda$ . For every  $n \in \mathbb{N}$  it holds that*

$$\left( \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (r_\lambda(t, u) - r_{\lambda, n}(t, u))^2 w_\alpha(t) w_\alpha(u) dt du \right)^{1/2} = \frac{1}{\sqrt{2}} \frac{1}{3^n}.$$

*Proof.* As in the proof of Proposition 3.3, we get

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (r_\lambda(t, u) - r_{\lambda, n}(t, u))^2 w_\alpha(t) w_\alpha(u) dt du = \sum_{m=n}^{\infty} \|\psi_{m, \lambda}\|_{\mathcal{L}_2(\mathbb{R}, w_\alpha)}^4.$$

By Proposition 5.1,

$$\sum_{m=n}^{\infty} \|\psi_{m, \lambda}\|_{\mathcal{L}_2(\mathbb{R}, w_\alpha)}^4 = \sum_{m=n}^{\infty} \frac{4}{9^{m+1}} = \frac{1}{2} \frac{1}{9^n}.$$

This completes the proof.  $\square$

## 6 Conclusion

In this article, we have demonstrated that Theorem 1.1 is a simple and powerful tool for constructing orthonormal expansions of translation-invariant kernels. In particular, using the Cholesky factor of the Fourier transform of the kernel together with the Laguerre functions led to an interesting decomposition of the RKHS of the Matérn kernel for half-integer smoothness parameters in terms of a finite dimensional space and a Hilbert space of functions vanishing at origin. This might be deemed unsatisfying, and a possible avenue to obtaining basis functions for Matérns in a common space would be to investigate constructions based on the symmetric square-root. The expansion for the Cauchy kernel was derived from the Fourier duality with the Matérn kernel of smoothness  $\alpha = 1/2$ . It remains an open problem to find a weighted  $\mathcal{L}_2$  space in which the Cauchy basis functions are orthogonal. For the Gaussian kernel, our construction is a means to reproduce certain Mercer expansions that are typically derived from Mehler's formula.

Observe that in each case we have used a basis of  $\mathcal{L}_2(\mathbb{R})$  that is “compatible” with the kernel, in being parametrised by  $\lambda$  in a suitable way that mirrors how this parameter appears in the kernel or its Fourier transform. For example, compare (3.1) and (3.3) or (5.1) and (5.3). It may be possible to replace  $\lambda$  with an arbitrary positive  $\kappa$  in the basis of  $\mathcal{L}_2(\mathbb{R})$  and base constructions

on the functions  $\varphi_{m,\kappa}$ . We do not know how to proceed with Matérns or the Cauchy kernel, but for the Gaussian kernel one can proceed exactly as in Section 5.1. Doing so yields the basis functions

$$\psi_{m,\lambda,\kappa}(t) = \left(\frac{\sqrt{2\kappa\lambda}}{a^2}\right)^{1/2} \sqrt{\frac{1}{2^m m!}} \left(1 - \frac{\kappa^2}{a^2}\right)^{m/2} e^{-\lambda^2(1-\kappa^2/a^2)t^2} H_m\left(\frac{\kappa\lambda^2 t}{a^2\sqrt{1-\kappa^2/a^2}}\right),$$

where  $a = \sqrt{\lambda^2 + \kappa^2/2}$ , for any  $\kappa \in (0, \sqrt{2}\lambda)$ .

## Acknowledgements

FT gratefully acknowledges financial support by the German Federal Ministry of Education and Research (BMBF) through Project ADIMEM (FKZ 01IS18052B), and financial support by the European Research Council through ERC StG Action 757275 / PANAMA; the DFG Cluster of Excellence “Machine Learning - New Perspectives for Science”, EXC 2064/1, project number 390727645; the German Federal Ministry of Education and Research (BMBF) through the Tübingen AI Center (FKZ: 01IS18039A); and funds from the Ministry of Science, Research and Arts of the State of Baden-Württemberg. TK was supported by the Academy of Finland postdoctoral researcher grant 338567 “Scalable, adaptive and reliable probabilistic integration”. Most of this article was written while TK was visiting the University of Tübingen in May 2022.

## References

- [1] ADLER, R. J. *An Introduction to Continuity, Extrema, and Related Topics for General Gaussian Processes*. No. 12 in Lecture Notes–Monograph Series. Institute of Mathematical Statistics, 1990.
- [2] ASKEY, R., AND WIMP, J. Associated Laguerre and Hermite polynomials. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics* 96, 1–2 (1984), 15–37.
- [3] BELL, W. W. *Special Functions for Scientists and Engineers*. Courier Corporation, 2004.
- [4] CHRISTOV, C. A complete orthonormal system of functions in  $L^2(-\infty, \infty)$  space. *SIAM Journal on Applied Mathematics* 42, 6 (1982), 1337–1344.
- [5] DICK, J., KUO, F. Y., AND SLOAN, I. H. High-dimensional integration: The quasi-Monte Carlo way. *Acta Numerica* 22 (2013), 133–288.
- [6] ERDÉLYI, A. *Tables of Integral Transforms. Volume II*. McGraw-Hill, 1954.
- [7] FASSHAUER, G., HICKERNELL, F., AND WOŹNIAKOWSKI, H. On dimension-independent rates of convergence for function approximation with Gaussian kernels. *SIAM Journal on Numerical Analysis* 50, 1 (2012), 247–271.
- [8] FASSHAUER, G., AND MCCOURT, M. *Kernel-based Approximation Methods using MATLAB*. No. 19 in Interdisciplinary Mathematical Sciences. World Scientific Publishing, 2015.
- [9] FASSHAUER, G. E., AND MCCOURT, M. J. Stable evaluation of Gaussian radial basis function interpolants. *SIAM Journal on Scientific Computing* 34, 2 (2012), A737–A762.
- [10] HAWKINS, D. L. Some practical problems in implementing a certain sieve estimator of the Gaussian mean function. *Communications in Statistics - Simulation and Computation* 18, 2 (1989), 481–500.
- [11] HIGGINS, J. R. *Completeness and Basis Properties of Sets of Special Functions*. No. 72 in Cambridge Tracts in Mathematics. Cambridge University Press, 1977.
- [12] IRRGEHER, C., AND LEOBACHER, G. High-dimensional integration on  $\mathbb{R}^d$ , weighted Hermite spaces, and orthogonal transforms. *Journal of Complexity* 31, 2 (2015), 174–205.
- [13] KARVONEN, T. Small sample spaces for Gaussian processes. *Bernoulli* (2022). To appear.

- [14] KIMELDORF, G. S., AND WAHBA, G. A correspondence between Bayesian estimation on stochastic processes and smoothing by splines. *The Annals of Mathematical Statistics* 41, 2 (1970), 495–502.
- [15] KUO, H.-H. *Gaussian Measures in Banach Spaces*. No. 463 in Lecture Notes in Mathematics. Springer, 1975.
- [16] MINH, H. Q. Some properties of Gaussian reproducing kernel Hilbert spaces and their implications for function approximation and learning theory. *Constructive Approximation* 32, 2 (2010), 307–338.
- [17] OWHADI, H., AND SCOVEL, C. Separability of reproducing kernel spaces. *Proceedings of the American Mathematical Society* 145, 5 (2017), 2131–2138.
- [18] PAULSEN, V. I., AND RAGHUPATHI, M. *An Introduction to the Theory of Reproducing Kernel Hilbert Spaces*. No. 152 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2016.
- [19] RASMUSSEN, C. E., AND WILLIAMS, C. K. I. *Gaussian Processes for Machine Learning*. Adaptive Computation and Machine Learning. MIT Press, 2006.
- [20] SOLIN, A., AND SÄRKKÄ, S. Hilbert space methods for reduced-rank Gaussian process regression. *Statistics and Computing* 30, 2 (2020), 419–446.
- [21] STEIN, M. L. *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Series in Statistics. Springer, 1999.
- [22] STEINWART, I., AND SCOVEL, C. Mercer’s theorem on general domains: On the interaction between measures, kernels, and RKHSs. *Constructive Approximation* 35 (2012), 363–417.
- [23] SZEGŐ, G. *Orthogonal Polynomials*. No. 23 in Colloquium Publications. American Mathematical Society, 1939.
- [24] VAN TREES, H. L. *Detection Estimation and Modulation Theory: Part I*. Wiley-Interscience, 2001.
- [25] WENDLAND, H. *Scattered Data Approximation*. No. 17 in Cambridge Monographs on Applied and Computational Mathematics. Cambridge University Press, 2005.
- [26] WIENER, N. *Extrapolation, Interpolation, and Smoothing of Stationary Time Series: With Engineering Applications*. MIT Press, 1949.
- [27] XIU, D. *Numerical Methods for Stochastic Computations*. Princeton University Press, 2010.
- [28] ZHU, H., WILLIAMS, C. K. I., ROHWER, R., AND MORCINIEC, M. Gaussian regression and optimal finite dimensional linear models. In *Neural Networks and Machine Learning*, C. M. Bishop, Ed., vol. 168 of *NATO ASI Series. Series F: Computer and Systems Science*. Springer, 1998, pp. 167–184.
- [29] ZWICKNAGL, B. Power series kernels. *Constructive Approximation* 29, 1 (2009), 61–84.